

A Review of Option Pricing Models and Simulations

Shi Guo

June 15, 2020

Abstract In this article, we review the strategies and a variety of methods for pricing options. We first review the theory of the Black-Scholes-Merton model, the Heston model with stochastic volatility and the Bates model that contains diffusion with jumps. Then we discuss the Monte Carlo simulation algorithms of each model and how to treat the stopping time for American style options. Finally, we present the simulation result.

1 European Option Pricing Models

In this section we firstly present two approaches for pricing European options: One is to solve Black-Scholes-Merton partial equation and the other one is using Martingale property. We are going to show these two ways are equivalent as they are connected through Feynman-Kac formula. Then we present the analytical solutions and discussions.

1.1 Black Scholes Merton Equation: Introduction

Here is basic idea for pricing options. We consider a portfolio composed of certain shares of stock and a saving account with an interest rate then use the portfolio to replicate the payoff of the option. If we are able to replicate the payoff successfully, we use the value of the portfolio to price the option. We briefly go through the steps to derive the Black Scholes Merton equation. We assume the stock prices following a geometric Brownian motion

1) Stock price:

$$dS(t) = \alpha S(t)dt + \sigma S(t)dW(t)$$

2) We have a portfolio $X(t)$ which consists of $\Delta(t)$ share of stock $\Delta(t)S(t)$, and $(X(t) - \Delta(t)S(t))$ money market account with interest rate r .

$$dX(t) = \Delta(t)dS(t) + r(X(t) - \Delta(t)S(t))dt$$

3) Change of the portfolio with respect to time

$$\begin{aligned} dX(t) &= \Delta(t)dS(t) + r(X(t) - \Delta(t)S(t))dt \\ &= rX(t)dt + \Delta(t)(\alpha - r)S(t) + \Delta(t)\sigma S(t)dW(t) \end{aligned}$$

4) Change of the present value of the stock with respect to time

$$d(e^{-rt}S(t)) = (\alpha - r)e^{-rt}S(t)dt + \sigma e^{-rt}S(t)dW(t)$$

5) With a few steps, we get change of the present value of the portfolio with respect to time

$$d(e^{-rt}X(t)) = \Delta(t)(\alpha - r)e^{-rt}S(t)dt + \Delta(t)\sigma e^{-rt}S(t)dW(t) \quad (1)$$

6) Assume the option value is $c(t, S(t))$ and we apply Ito's formula

$$\begin{aligned} &d(e^{-rt}c(t, S(t))) \\ &= e^{-rt}[-rc(t, S(t)) + c_t(t, S(t)) + \alpha S(t)\frac{\partial c(t, S(t))}{\partial S(t)} + \frac{1}{2}\sigma^2 S^2(t)\frac{\partial^2 c(t, S(t))}{\partial S^2(t)}]dt \\ &\quad + e^{-rt}\sigma S(t)\frac{\partial c(t, S(t))}{\partial S(t)}dW(t) \end{aligned}$$

7) Now equate Equation in 5) and 6), we get $dW(t)$ term:

$$\Delta(t) = \frac{\partial c(t, S(t))}{\partial S(t)}$$

dt term:

$$rc(t, S) = c_t(t, S(t)) + rS(t)\frac{\partial c(t, S(t))}{\partial S(t)} + \frac{1}{2}\sigma^2 S(t)\frac{\partial^2 c(t, S(t))}{\partial S^2(t)} \quad (2)$$

which is known as Black-Scholes-Merton partial differential equation. The terminal condition the equation satisfies for call option is

$$c(T, S) = (S(T) - K)^+$$

Similarly, for put option

$$p(T, S) = (K - S(T))^+$$

1.2 Martingale Approach

In risk-neutral measure, we write the stock price as

$$dS(t) = rS(t)dt + \sigma S(t)d\tilde{W}(t)$$

Where $\tilde{W}(t)$ is a standard Brownian motion under risk-neutral measure. So by substituting $\alpha = r$ in Eq. 1, we have

$$d(e^{-rt}X(t)) = \Delta(t)\sigma e^{-rt}S(t)dW(t)$$

As the expectation of $dW(t)$ is 0, the expectation of discounted value of the portfolio does not change with time, which implies the discounted value of the portfolio is a martingale. Using this martingale property, we can get the value of the portfolio at time t

$$X(t) = \tilde{E}[e^{-r(T-t)}X(T)|F(t)]$$

where $X(T) = V(T)$. We set the option price $c(t, S(t))$ to be $X(t)$

$$c(t, S(t)) = \tilde{E}[e^{-r(T-t)}V(T)|F(t)]$$

1.3 Connection to Feynman-Kac formula

From above, we know

$$c(t, S(t)) = \tilde{E}[e^{-r(T-t)}V(T)|F(t)] = \tilde{E}[e^{-r(T-t)}h(S(T))|F(t)] \quad (3)$$

Based on Feynman-Kac formula, there is a function $c(t, S(t))$ which must satisfy discounted partial differential equation

$$c_t(t, S) + rSc(t, S) + \frac{1}{2}\sigma^2 S^2 c_{SS}(t, S) = rc(t, S)$$

Where $c(T, S(T)) = V(S(T))$. Now we have seen two ways of showing the Black-Scholes-Merton(BSM) equation. One way is to reproduce the payoff of the option using a portfolio that consists of a saving account. Another way is based on the risk-neutral pricing formula and Feynman-Kac formula. These two ways are equivalent. Because under risk-neutral measure, the payoff of a derivative is the same as a saving account, which imply we are able to reproduce the payoff using portfolio that consisting of a saving account.

1.4 Black-Scholes-Merton Model: Analytic Solution for European Option

1. European call option

For European call option with payoff to be $V(T) = S(T) - K$, with K as strike price, let us assume constant volatility σ , and constant interest rate r . Then we can obtain the solution to the BSM equation with martingale property without bothering solving the complex partial differential equation. The call option value satisfies

$$c(t, S(t)) = \tilde{E}[e^{-r(T-t)}(S(T) - K)^+ | F(t)]$$

We write

$$\begin{aligned} S(T) &= S(t) \exp\{\sigma(\tilde{W}(T) - \tilde{W}(t)) + (r - \frac{1}{2}\sigma^2)\tau\} \\ &= S(t) \exp\{-\sigma\sqrt{\tau}Y + (r - \frac{1}{2}\sigma^2)\tau\} \end{aligned}$$

Where Y is the standard normal random variable and $\tau = T - t$ is the time to expiration.

$$Y = -\frac{\tilde{W}(T) - \tilde{W}(t)}{\sqrt{T-t}}$$

So we see that $S(T)$ is the product of the $F(t)$ measurable random variable $S(t)$ and random variable

$$\exp\{-\sigma\sqrt{\tau}Y + (r - \frac{1}{2}\sigma^2)\tau\}$$

Which is independent of $F(t)$. Therefore based on risk-neutral pricing formula[3]

$$\begin{aligned} c(t, x) &= \tilde{E}[e^{-r\tau}(x \exp\{-\sigma\sqrt{\tau}Y + (r - \frac{1}{2}\sigma^2)\tau\} - K)^+] \\ &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-\tau r} (x \exp\{-\sigma\sqrt{\tau}y + (r - \frac{1}{2}\sigma^2)\tau\} - K)^+ e^{-\frac{1}{2}y^2} dy \end{aligned}$$

After a little bit of math with integration, we have the solution to the Black-Scholes-Merton model for European call option

$$c(\tau, x; K, r, \sigma) = xN(d_1(\tau, x)) - e^{-r\tau}KN(d_2(\tau, x))$$

Where

$$\begin{aligned} d_1 &= \frac{1}{\sigma\sqrt{\tau}}[\ln(\frac{S_t}{K}) + (r + \frac{\sigma^2}{2})\tau] \\ d_2 &= d_1 - \sigma\sqrt{\tau} \end{aligned}$$

$N(\cdot)$ is the cumulative distribution function of the standard normal distribution

2. European put option

The payoff for the European put option is $V(T) = K - S(T)$, we can follow a similar derivation and get the formula for put option

$$p(\tau, x; K, r, \sigma) = Ke^{-r\tau}N(-d_2) - xN(-d_1)$$

3. Options based on dividend-paying stock

The value of dividend-paying stock consists of two parts. One is the value of the stock, the other one is the value of dividend. Suppose the dividend rate is a , then the change of value of dividend-paying stock over time dt is $dS(t) + aS(t)dt$. In order to make the discounted value a martingale, we require

$$\frac{dS(t) + aS(t)dt}{S(t)} = rdt + \sigma d\tilde{W}(t)$$

so

$$dS(t) = (r - a)S(t)dt + \sigma S(t)d\tilde{W}(t)$$

$$S(T) = S(t) \exp\{\sigma(\tilde{W}(T) - \tilde{W}(t)) + (r - a - \frac{1}{2}\sigma^2)\tau\}$$

Following similar derivation, for call option we can get

$$c(\tau, x; K, r, \sigma) = xe^{-a\tau} N(d_1(\tau, x)) - e^{-r\tau} K N(d_2(\tau, x))$$

$$p(\tau, x; K, r, \sigma) = Ke^{-r\tau} N(-d_2) - xe^{-a\tau} N(-d_1)$$

4. Boundary conditons

Using the solution $c(t, x)$ and $p(t, x)$, we can easily check the boundary conditions when time t approaches to expiration time T .

As we know

$$d_1 = \frac{1}{\sigma\sqrt{\tau}} \ln\left(\frac{S}{K}\right) + \frac{1}{\sigma}\left(r + \frac{\sigma^2}{2}\sqrt{\tau}\right)$$

When $\tau \rightarrow 0$, the second term decays much faster, so it vanishes. When $S > K$, d_1 goes to infinity, when $S < K$, d_1 goes to negative infinity. Therefore, when $S > K$


$$c(t, x) = S * N(+\infty) - K * N(+\infty) = S - K$$


Therefore, when $S < K$

$$c(t, x) = S * N(-\infty) - K * N(-\infty) = 0 - 0 = 0$$

5. Examples

The following graphs show the change of option price with respect to different





6. Alternative formulation

If we introduce $F = e^{(r-a)\tau}S$, which is the forward price of the asset S . Then the equation pricing equation becomes

$$\begin{aligned} C(F, \tau) &= D[N(d_+)F - N(d_-)K] \\ P(F, \tau) &= D[N(-d_-)K - N(-d_+)F] \\ d_{+/-} &= \frac{1}{\sigma\sqrt{\tau}}[\ln(\frac{F}{K}) + / - \frac{1}{2}\sigma^2\tau] \end{aligned}$$

The variables are:

$\tau = T - t$ is the time to expiry

$D = e^{-r\tau}$ is the discount factor

1.5 Heston Stochastic Volatility Model

The Black-Scholes equation assumes the volatility is constant, which is the ideal case and not practical in the real market. The Heston model[1] assumes the volatility to follow a stochastic process. Suppose a stock price under risk-neutral measure is governed by

$$dS(t) = rS(t)dt + \sqrt{V(t)}S(t)\tilde{d}W_1(t) \quad (4)$$

and the volatility itself is governed by the equation

$$dV(t) = (a - bV(t))dt + \sigma\sqrt{V(t)}\tilde{d}W_2(t) \quad (5)$$

$$= b(\frac{a}{b} - V(t))dt + \sigma\sqrt{V(t)}\tilde{d}W_2(t) \quad (6)$$

$$(7)$$

Where

$$\tilde{d}W_1(t)\tilde{d}W_2(t) = \rho dt$$

The parameter $\frac{a}{b}$ can be viewed as a long-term average variance. And the instantaneous variance is pulled elastically towards the long-term variance by a speed controlled by parameter b .

Fourier Transform Solution to Heston Model

For Heston model, the same martingale property still holds. And the option price is

$$c(t, S(t)) = \tilde{E}[e^{-r(T-t)}V(T)|F(t)] = \tilde{E}[e^{-r(T-t)}(S_T - K)^+|F(t)]$$

Let $x = \ln(S_T)$, and $\tau = T - t$, then, and pdf of $\ln(x_T)$ is $p(x)$

$$\begin{aligned} C(t, S(t)) &= e^{-r\tau} \int_{\ln K}^{\infty} (e^x - K)p(x)dx \\ &= e^{-r\tau} \int_{\ln K}^{\infty} e^x p(x)dx - e^{-r\tau} K \int_{\ln K}^{\infty} p(x)dx \\ &= S_0 f - e^{-r\tau} K g \end{aligned}$$

where

$$f = \frac{1}{S_0} e^{-r\tau} \int_{\ln K}^{\infty} e^x p(x)dx$$

$$g = \int_{\ln K}^{\infty} p(x)dx$$

To evaluate f and g is difficult as the analytical form of $p(x)$ does not exist. However, literature has proposed that the characteristic function of pdf of x exist. Heston [1] proposed a solution using characteristic function. We list the result here and we will elaborate it in the Appendix. Suppose $\phi(u)$ is the characteristic function of $p(x)$, then using inverse Fourier transform

$$p(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-iux} \phi(u) du$$

with $\phi(u)$,

$$g = \int_{\ln K}^{\infty} p(x)dx = \int_{\ln K}^{\infty} \left(\frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-iux} \phi(u) du \right) dx$$

Changing the order the integration yields

$$\begin{aligned} g &= \frac{1}{2\pi} \int_{-\infty}^{\infty} \phi(u) \left(\int_{\ln K}^{\infty} e^{-iux} dx \right) du \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} \phi(u) \left(\int_{-\infty}^{\infty} \mathbf{1}_{x > \ln K} e^{-iux} dx \right) du \end{aligned}$$

The inner integral is the inverse Fourier transform of a step function and the result is well-known.

$$\begin{aligned}
g &= \frac{1}{2\pi} \int_{-\infty}^{\infty} \phi(u) \left(e^{-iulnK} \frac{1}{iu} + \pi \delta(u) \right) du \\
&= \frac{1}{2\pi} \int_{-\infty}^{\infty} \phi(u) \left(e^{-iulnK} \frac{1}{iu} \right) du + \frac{1}{2} \\
&= \frac{1}{\pi} \int_0^{\infty} \phi(u) e^{-iulnK} \frac{1}{iu} du + \frac{1}{2}
\end{aligned}$$

f is a little tricky,

$$f = \frac{1}{S_0} e^{-r\tau} \int_{lnK}^{\infty} e^x p(x) dx = \frac{1}{S_0} e^{-r\tau} \frac{\int_{lnK}^{\infty} e^x p(x) dx}{\int_{-\infty}^{\infty} e^x p(x) dx} \int_{-\infty}^{\infty} e^x p df(x) dx$$

Based on martingale property $\int_{-\infty}^{\infty} e^x p(x) dx = e^{rT} S_0$, so

$$f = \int_{lnK}^{\infty} \frac{e^x p(x) dx}{\int_{-\infty}^{\infty} e^x p(x) dx} = \int_{lnK}^{\infty} p' dx$$

The Fourier transform of function p' is

$$\begin{aligned}
&\frac{\int_{-\infty}^{\infty} e^{iux+x} p(x) dx}{\int_{-\infty}^{\infty} e^{i(-i)x} p(x) dx} \\
&= \frac{\phi(u-i)}{\phi(-i)}
\end{aligned}$$

Therefore we can write f as

$$f = \frac{1}{\pi} \int_0^{\infty} \frac{e^{-iuln(K)} \phi(u-i)}{iu\phi(-i)} du$$

1.6 Jump Diffusion Model

The BSM model and the Heston model assumes the asset prices follows the stochastic process driven by Brownian motion. Under this assumption, the asset price is continuous in time. The jump diffusion model extends the BSM model and Heston model by assuming asset price has discrete jump in time. Therefore in jump diffusion model, the stochastic process contains a continuous Brownian motion process and a discrete jump process.

1. Number of jumps during per unit time: Poission distribution

The number of jump should be either zero or a positive integer. Let N be the number of jump per unit of time and k be a nonnegative integer. Then the probability of $N = k$ should satisfies

$$\sum_{k=0}^{\infty} P(N = k) = 1$$

In order to find the distribution, we borrow the idea of Taylor's expansion. We consider a function e^λ , its Taylor's expansion is

$$e^\lambda = \sum_{k=0}^{\infty} \frac{\lambda^k}{k!}$$

Dividing e^λ on both sides, we can

$$1 = \sum_{k=0}^{\infty} \frac{\lambda^k}{k!} e^{-\lambda}$$

So we can assign the probability

$$P(N = k) = \frac{\lambda^k}{k!} e^{-\lambda}$$

This is the well-known Poisson distribution. It is easy to show Poisson distribution has mean and variance equal to λ . So λ can be interpreted as the average number of jumps per unit of time.

2. Number of jumps during time interval Δt : Poisson process

Now we consider the number of jumps in a given time interval Δt , and we call it ΔN . The average jump during Δt is $\lambda \Delta t$. So we can get the probability distribution by substitute $\lambda \Delta t$ to λ

$$P(\Delta N = k) = \frac{(\lambda \Delta t)^k}{k!} e^{-\lambda \Delta t}$$

The mean of ΔN is $\lambda \Delta t$.

Let Δt be really small time interval dt , and from above we see

$$\begin{aligned} P(dN = 0) &= e^{-\lambda dt} \approx 1 - \lambda dt \\ P(dN = 1) &= \lambda dt e^{-\lambda dt} \approx \lambda dt (1 - \lambda dt) \approx \lambda dt \\ P(dN = 2) &= (-\lambda dt)^2 e^{-\lambda dt} \approx 0 \end{aligned}$$

This makes perfectly sense, as during an infinitesimal time interval, the jump can either not happen or happen only once.

We define S_k is the time when the k th jump occurs and we assume $S_0 = 0$. Then the Poisson process is

$$N(t) = k \text{ if } S_k \leq t < S_{k+1}$$

With $E[N(t)] = \lambda t$. Note the Poisson process defined above is right-continuous.

3. Compensated Poisson process

The Poisson process has a mean that is a function of time. As time evolves, the process has a non-zero drift so it is not a martingale. We define a compensated Poisson process $M(t)$ which obeys the martingale property

$$\begin{aligned} M(t) &= N(t) - \lambda t \\ E[M(t)|F(s)] &= M(s) \end{aligned}$$

4. Compound Poisson process

In the above definition, we assume the jump size in the Poisson process is 1. Now we allow the jump size to be random. Let $N(t)$ be a Poisson process with mean λ and Y_i be a sequence of random variable with mean $E[Y] = m$. We define compound Poisson process

$$Q(t) = \sum_{i=1}^{N(t)} Y_i$$

The number of jumps follows Poisson distribution, and at each jump, the size of the jump is Y_i . In other words, the first jump size is Y_1 and the second jump size is Y_2 , etc.

The mean of the compound Poisson process is the product of the number of jumps and the size of the jumps, as these two are independent.

$$EQ(t) = m\lambda t$$

We can also define a compensated compound Poisson process

$$\tilde{M}(t) = Q(t) - m\lambda t$$

And this is a martingale.

5. Geometric Poisson process with constant jump size

In the compound Poisson process, let the jump size be a constant y . When jump occurs, we have

$$Q(t_i) = Q(t_{i-}) + y$$

The asset value in the financial market follows the geometric Poisson process. Let $S(t)$ be the asset price, let

$$\begin{aligned} S(t_i) &= S(t_{i-})y \\ \frac{S(t_i) - S(t_{i-})}{S(t_{i-})} &= y - 1 \\ \Delta S(t_i) &= (y - 1)S(t_{i-})\Delta N(t) \end{aligned}$$

Where $\Delta N(t) = 1$ when jump occurs, 0 otherwise. When $N(t) = 1$, there is one jump from time zero up to t , so $S(t) = yS(0)$. When $N(t) = 2$, $S(t) = y^2S(0)$. So for any arbitrary $N(t)$

$$S(t) = S(0)y^{N(t)}$$

In order to fulfill the martingale condition, we write

$$dS(t) = (y - 1)S(t)dN(t) - (y - 1)\lambda S(t)dt$$

Its integral form is

$$S(t) = S(0)y^{N(t)}\exp(-(y - 1)\lambda t)$$

6. Geometric Poisson process with log-normal jump size: Bates jump diffusion model

If the jump size Y is random in the geometric Poisson process, then to model the asset price, we need to modify the last equation by changing $y^{N(t)}$ to $\prod_{i=1}^{N(t)} Y_i$, and $y - 1$ to $E[Y] - 1$.

$$S(t) = S(0) \prod_{i=1}^{N(t)} Y_i \exp(-(E[Y] - 1)\lambda t)$$

We need to model the jump size Y_i . Y_i is a strictly positive random variable so a good candidate is the exponential of normal random variable which is log-normal random variable. If we have a normal distributed random variable $J \sim N(\nu, \delta^2)$, and let Y be e^J . Then Y follows log-normal distribution with mean $e^{\nu + \frac{1}{2}\delta^2}$. We can write $S(t)$ using J

$$S(t) = S(0) \exp \sum_i^{N(t)} J_i \exp(-(e^{\nu + \frac{1}{2}\delta^2} - 1)\lambda t)$$

Let $Z = \sum_i^{N(t)} J_i$, which is a sum of normal random variable. Then $Z \sim Normal(N(t)\nu, N(t)\delta^2)$. So

$$S(t) = S(0) \exp(Z - (e^{\nu + \frac{1}{2}\delta^2} - 1)\lambda t)$$

Where $Z \sim Normal(N(t)\nu, N(t)\delta^2)$ given $N(t) \sim Poisson(\lambda t)$.

2 Monte Carlo Simulation Processes

In simulation, we decompose the time evolution of an asset price into three components: drift, diffusion, jump. The drift term is dt term, the diffusion term has Brownian motion component, the jump term has Poisson component. Let us look at the following examples.

1. BSM process

In BSM process, the asset price follows

$$S(t) = S(0) \exp((r - \frac{1}{2}\sigma^2)dt + \sigma dW(t))$$

The drift term is clearly $r - \frac{1}{2}\sigma^2$ for any given dt . For the diffusion term, we need to generate $W(t)$. We know the $W(t)$ is a Normal random variable with mean 0 and variance t . So suppose we generate a Gaussian random variable \hat{W} , it turns out $W = \sqrt{t}\hat{W}$ is normally distributed with variance t .

$$\begin{aligned} \mu &= r - \frac{1}{2}\sigma^2 \\ S^{(t+dt)} &= S^{(t)} \exp(\mu dt + \sigma \sqrt{dt} \hat{W}) \end{aligned}$$

2. Heston process

The Heston model states

$$\begin{aligned} dS(t) &= rS(t)dt + \sqrt{V(t)}S(t)\tilde{d}W_1(t) \\ dV(t) &= (a - bV(t))dt + \sigma\sqrt{V(t)}\tilde{d}W_2(t) \\ \tilde{d}W_1(t)\tilde{d}W_2(t) &= \rho dt \end{aligned}$$

a. Generating two Brownian motions with correlation.

To simulate Heston process needs to generate two Gaussian random variables that has correlation ρ , we do this by first generate two independent Gaussian random variables W_1, W_2 , then

$$\begin{aligned} W_1' &= \hat{W}_1 \\ W_2' &= \rho\hat{W}_1 + \sqrt{1 - \rho^2}\hat{W}_2 \end{aligned}$$

We can easily check the mean of W_2'

$$\begin{aligned} E[W_2'] &= \rho E[W_1] + \sqrt{1 - \rho^2}E[W_2] = 0 \\ Var[W_2'] &= \rho^2 Var[W_1] + (1 - \rho^2)Var[W_2] = \rho^2 + 1 - \rho^2 = 1 \end{aligned}$$

$$\begin{aligned} cor(W_1', W_2') &= \frac{E[W_1'W_2']}{\sqrt{Var[W_1']}\sqrt{Var[W_2']}} \\ &= E[W_1(\rho W_1 + \sqrt{1 - \rho^2})W_2] \\ &= \rho E[W_1^2] + \sqrt{1 - \rho^2}E[W_1W_2] \\ &= \rho \end{aligned}$$

The last step uses the fact W_1 and W_2 are independent, so $E[W_1W_2] = E[W_1]E[W_2] = 0$.

b. Coping with negative $V(t)$.

The Heston model setting provides a non-negative $V(t)$. This can be seen when $V(t)$ is positively moving close to zero, the drift term adt pulls the $V(t)$ up and avoid $V(t)$ to cross zero. However, in discretized version $V(t)$ may attain negative values. There exists several methods to deal with this issue.

(1) Absorption: Keep positive part of the previous $V(t)$ and use it to calculate next step $V(t + dt)$ and $X(t+dt)$

$$V^{(t+dt)} = V^{+(t)} + (a - bV^{+(t)})\sqrt{dt} + \sigma\sqrt{V^{+(t)}}dt(\rho\hat{W}_1 + \sqrt{1 - \rho^2}\hat{W}_2)$$

(2) Relection: Keep the absolute value of the previous $V(t)$ and use it to calculate next step $V(t + dt)$ and $X(t+dt)$

$$V^{(t+dt)} = |V^{(t)}| + (a - b|V^{(t)}|)\sqrt{dt} + \sigma\sqrt{|V^{(t)}|}dt(\rho\hat{W}_1 + \sqrt{1 - \rho^2}\hat{W}_2)$$

(3) Partial truncation: Using the positive part in diffusion term of $V(t + dt)$ and in $X(t + dt)$

$$V^{(t+dt)} = V^{(t)} + (a - bV^{(t)})\sqrt{dt} + \sigma\sqrt{V^{+(t)}}dt(\rho\hat{W}_1 + \sqrt{1 - \rho^2}\hat{W}_2)$$

(4) Full truncation[3]: Using the positive part in drift and diffusion terms of $V(t + dt)$, and in $X(t + dt)$. This scheme is proved to have the lowest bias.

$$V^{(t+dt)} = V^{(t)} + (a - bV^{+(t)})\sqrt{dt} + \sigma\sqrt{V^{+(t)}}dt(\rho\hat{W}_1 + \sqrt{1 - \rho^2}\hat{W}_2)$$

Together with formula for the drift and asset price, we write

$$\begin{aligned}\mu &= r - \frac{1}{2}(V^{+(t)}) \\ S^{(t+dt)} &= S^{(t)}\exp(\mu dt + \sqrt{V^{+(t)}}\sqrt{dt}W_1) \\ V^{(t+dt)} &= V^{(t)} + (a - bV^{+(t)})\sqrt{dt} + \sigma\sqrt{V^{+(t)}}dt(\rho\hat{W}_1 + \sqrt{1 - \rho^2}\hat{W}_2)\end{aligned}$$

3. Bates process

$$\begin{aligned}\mu &= r - \frac{1}{2}(V^{+(t)}) - \lambda(e^{\nu + \frac{1}{2}\delta^2} - 1) \\ S^{(t+dt)} &= S^{(t)}\exp(\mu dt + \sqrt{V^{+(t)}}\sqrt{dt}dW_1 + (\nu_*N + \delta^2 * \sqrt{N}) * dW_3) \\ V^{(t+dt)} &= V^{(t)} + (a - bV^{+(t)})dt + \sigma\sqrt{V^{+(t)}}dt(\rho\hat{W}_1 + \sqrt{1 - \rho^2}\hat{W}_2)\end{aligned}$$

3 Simulation for American Option

American Option can be exercised at any time. So in Monte Carlo Simulation, for each path we need to determine the stopping time, in other words, time for the option to be exercised. For each sample path at each time step point, we need to compare the current payoff value with the value to keep holding this option. If the current payoff value is great than the value to keep holding this option, we choose to exercise, otherwise we hold it. To evaluate the value to hold this option requires another Monte Carlo simulation, so the simulation for American option has a "Monte Carlo on Monte Carlo feature", and this is very computationally expensive. To see why it is expensive, consider the case of M samples and we are at time step 1. To determine the value of holding the option, we need generate another Monte Carlo simulation which has M sample paths. As we move on the time step, each step needs the number of sample path M times the previous one, therefore, the total number of Monte Carlo simulation grows exponentially as time step grows.

However, this does not mean there is no feasible way to evaluate American options. One idea is to avoid doing additional Monte Carlo simulation at each step to determine the value of holding the option and utilizing the simulation path of the European options. LongStaff[2] proposed an algorithm to determine the stopping time by using least square regression and here we briefly outline the algorithm.

Let us assume the option has a payoff function $h(S, K)$, where K is strike price. The algorithm simply has two parts. The first part is to follow the standard simulation scheme for European options. We generate M samples path from timestep 0 to timestep T . For each sample, the asset price forms a price path from $i = 0$ to T and the current asset price at a given timestep i is $S_i^{(m)}$, where $0 \leq i \leq T$. We define a discount factor $D_{i,j}$ to discount the asset price at time j to time i . For example $D_{1,3} = e^{-(3-1)r} = e^{-2r}$. Then the second part is to define a stopping time $\tau^{(m)}$ for each path.

- (1) set $\tau^{(m)}$ to T for each sample path m .
- (2) for t from $T - 1$ to 1:
for each sample:
 - (2b) calculate discount payoff value: $DV^{(m)} = D_{t,\tau}h(S_\tau^{(m)}, K)$
 - (2c) calculate the current payoff value: $h(S_t^{(m)})$
 - (2d) regress DV on S_t using least square and get estimated discount payoff value $EDV = f(X_t)$
 - (2e) if $h(S_t^{(m)}) > EDV^{(m)}(X_t^{(m)})$,
this means we can exercise the option, then upate $\tau^{(m)} = t$.
- (3) for each path calculate the present value of the payoff at stopping time τ
 $PV^{(m)} = D_{t,\tau}h(S_\tau^{(m)}, K)$
- (4) average $PV^{(m)}$ to get the price of the option

4 Result

Simulation of option value with BSM process using both analytical(ANAL) and Monte Carlo(MC)

We calculate value of option of different type and style listed in Table. 1. For MC method, we choose 100 timesteps and 100000 sample paths. For European style, we also compare the result with the analytical one. Our data matches the data presented in [2]. The result verified that for European option, there is no incentive to exercise early as the payoff function given the risk-neutral measure is a submartingale and has the trend to increase. For American put option, the NPV is larger than the NPV of European option by 1.05, so the early exercise value is clearly visible.

A Appledices

A.1 Semi-analytical Solution to Heston Model

$$dS(t) = rS(t)dt + \sqrt{V(t)}S(t)\tilde{d}W_1(t) \quad (8)$$

and the volatility itself is governed by the equation

$$dV(t) = (a - bV(t))dt + \sigma\sqrt{V(t)}\tilde{d}W_2(t) \quad (9)$$

$$= b\left(\frac{a}{b} - V(t)\right)dt + \sigma\sqrt{V(t)}\tilde{d}W_2(t) \quad (10)$$

$$(11)$$

Style	Type	S	K	r	σ	NPV(ANAL)	NPV(MC)	Ref[2]
European	Call	36.0	40	0.06	0.2	4.28	4.27(3)	
American	Call	36.0	40	0.06	0.2	N/A	4.28(3)	
European	Put	36.0	40	0.06	0.2	3.76	3.76(2)	
American	Put	36.0	40	0.06	0.2	N/A	4.81(3)	4.82(1)
European	Call	36.0	40	0.06	0.4	8.22	8.27(6)	
American	Call	36.0	40	0.06	0.4	N/A	8.23(6)	
European	Put	36.0	40	0.06	0.4	7.70	7.70(4)	
American	Put	36.0	40	0.06	0.4	N/A	8.50(3)	8.49(2)
European	Call	44.0	40	0.06	0.2	9.95	9.93(5)	
American	Call	44.0	40	0.06	0.2	N/A	9.91(5)	
European	Put	44.0	40	0.06	0.2	1.43	1.44(1)	
American	Put	44.0	40	0.06	0.2	N/A	1.67(3)	1.68(1)
European	Call	44.0	40	0.06	0.4	13.73	13.81(9)	
American	Call	44.0	40	0.06	0.4	N/A	13.75(8)	
European	Put	44.0	40	0.06	0.4	5.20	5.18(3)	
American	Put	44.0	40	0.06	0.4	N/A	5.58(3)	5.62(2)

Table 1: Simulation result of different options using BSM process. S: Underlying Asset Price. K: strike price. r: interest rate. σ : volatility. NPV(ANAL): Analytical result of net present value. NPV(MC): Monte Carlo result of net present value.

Where

$$\tilde{d}W_1(t)\tilde{d}W_2(t) = \rho dt$$

At time t , the risk-neutral price of a call expiring at time $T \geq t$ in this model is

$$c(t, S(t), V(t)) = \tilde{E}[e^{-r(T-t)}(S(T) - K)^+ | F(t)]$$

If we move the term e^{-rt} to the left hand side, we see

$$e^{-rt}c(t, S(t), V(t)) = \tilde{E}[e^{-rT}(S(T) - K)^+ | F(t)] \quad (12)$$

which satisfies the martingale property. Then we take the differentiation of

$e^{-rt}c(t, S(t), V(t))$. We get

$$\begin{aligned} & d(e^{-rt}c(t, S(t), V(t))) \\ &= \frac{\partial e^{-rt}}{\partial t}c(t, S(t), V(t)) + e^{-rt}\frac{\partial c(t, S(t), V(t))}{\partial t}dt \\ &= -re^{-rt}c(t, S(t), V(t))dt \quad (1) \end{aligned}$$

$$+e^{-rt}\frac{\partial c}{\partial t}dt \quad (2)$$

$$+e^{-rt}\frac{\partial c}{\partial S}dS \quad (3)$$

$$+e^{-rt}\frac{\partial^2 c}{\partial S^2}dSdS \quad (4)$$

$$+e^{-rt}\frac{\partial c}{\partial V}dV \quad (5)$$

$$+e^{-rt}\frac{\partial^2 c}{\partial V^2}dVdV \quad (6)$$

$$+e^{-rt}\frac{\partial^2 c}{\partial V\partial S}dVdS \quad (7)$$

As we are interested in only the dt terms, we find out the dt terms from (1) to (7) the dt term in (1) is

$$-rc(t, S(t), V(t))e^{-rt}dt$$

the dt term in (2) is

$$\frac{\partial c}{\partial t}e^{-rt}dt$$

the dt term in (3) is

$$\frac{\partial c}{\partial S}rSe^{-rt}dt$$

the dt term in (4) is

$$\frac{1}{2}\frac{\partial^2 c}{\partial S^2}VS^2e^{-rt}dt$$

the dt term in (5) is

$$\frac{\partial c}{\partial V}(a - bV(t))e^{-rt}dt$$

the dt term in (6) is

$$\frac{1}{2}\frac{\partial^2 c}{\partial V^2}V\sigma^2e^{-rt}dt$$

the dt term in (7) is

$$\frac{\partial^2 c}{\partial V\partial S}VS\sigma e^{-rt}dt$$

Collect all the dt terms and let those terms equal to zero, we get

$$c_t + rsc_s + (a - bv)c_v + \frac{1}{2}s^2vc_{ss} + \rho\sigma svc_{sv} + \frac{1}{2}\sigma^2vc_{vv} = rc \quad (13)$$

The function $c(t, s, v)$ satisfies boundary condition

$$\begin{aligned} c(T, s, v) &= (s - K)^+ \\ c(t, 0, v) &= 0 \\ c(t, s, 0) &= (s - e^{-r(T-t)}K)^+ \\ \lim_{s \rightarrow \infty} \frac{c(t, s, v)}{s - K} &= 1 \\ \lim_{v \rightarrow \infty} c(t, s, v) &= s \end{aligned}$$

Based on the solution to the BSM equation, we can guess that the solution has the following form

$$c(t, s, v) = sf(t, \log s, v) - e^{-r(T-t)}Kg(t, \log s, v) \quad (14)$$

Where f and g can be interpreted as a cumulative distribution function. Then since $c(t, s, v)$ satisfies the partial differential equation 13, we can show that f and g satisfy the following

$$f_t + (r + \frac{1}{2}v)f_x + (a - bv + \rho\sigma v)f_v + \frac{1}{2}vf_{xx} + \rho\sigma vf_{xv} + \frac{1}{2}\sigma^2vf_{vv} = 0 \quad (15)$$

$$(16)$$

$$g_t + (r - \frac{1}{2}v)g_x + (a - bv)g_v + \frac{1}{2}vg_{xx} + \rho\sigma vg_{xv} + \frac{1}{2}\sigma^2vg_{vv} = 0 \quad (17)$$

$$(18)$$

The derivation is straightforward but one needs to keep in mind here we treat x and v as two independent variables. The above PDE for f and g satisfy boundary condition

$$\begin{aligned} f(T, x, v) &= 1_{x \geq \log K} \\ g(T, x, v) &= 1_{x \geq \log K} \end{aligned}$$

This implies that f and g can be interpreted as "Probabilities". We can define

$$f(t, x, v) = E^{t, x, v} 1_{x \geq \log K}$$

We suppose a pair of stochastic process $X(t)$, $V(t)$ given by the following expression

$$dX(t) = (r + \frac{1}{2}V(t))dt + \sqrt{V(t)}dW_1(t) dV(t) = (a - bV(t) + \rho\sigma V(t))dt + \sigma\sqrt{V(t)}dW_2(t)$$

By F-K formula, we can show that f satisfies the PDE above. Similarly, we have

$$g(t, x, v) = E^{t,x,v} 1_{x \geq \log K}$$

and the stochastic process of $X(t)$ and $V(t)$ are

$$\begin{aligned} dX(t) &= (r - \frac{1}{2}V(t))dt + \sqrt{V(t)}dW_1(t) \\ dV(t) &= (a - bV(t))dt + \sigma\sqrt{V(t)}dW_2(t) \end{aligned}$$

To find the analytical solution of $f(t, x, v)$ and $g(t, x, v)$ is not an easy task. Instead, we do Fourier transform of f and g . First we work out the function $f(t, x, v)$. Let $\tau = T - t$

$$\tilde{f}(k, v, \tau) = \int_{-\infty}^{\infty} dx e^{-ikx} f(x, v, \tau)$$

The inverse Fourier transform is

$$f(x, v, \tau) = \int_{-\infty}^{\infty} \frac{dk}{2\pi} e^{ikx} \tilde{f}(k, v, \tau)$$

Substitute this into equation 15, then

$$-\frac{\partial \tilde{f}}{\partial \tau} + (r + \frac{1}{2})ik\tilde{f} + (a - bv + \rho\sigma v)\frac{\partial \tilde{f}}{\partial v} - \frac{1}{2}vk^2\tilde{f} + \rho\sigma vik\frac{\partial \tilde{f}}{\partial v} + \frac{1}{2}\sigma^2 v\frac{\partial^2 \tilde{f}}{\partial v^2} = 0 \quad (19)$$

$$(20)$$

Now the problem is to solve for \tilde{f} . We note when $\tau = 0$,

$$\tilde{f}(k, v, 0) = \int_{-\infty}^{\infty} dx e^{-ikx} f(x, v, \tau = 0) = \int_{-\infty}^{\infty} dx e^{-ikx} 1_{x \geq \log K} = \int_0^{\infty} dx e^{-ikx} = \pi\delta(k) + \frac{1}{ik}$$

when $\tau \neq 0$, we guess a general solution which has the following form

$$\tilde{f}(k, v, \tau) = \exp(C\tau + D\tau v)\tilde{f}(k, v, 0)$$

From above we easily see it match the terminal condition at $\tau \rightarrow 0$. With inverse Fourier transform

$$f(x, v, \tau, x) = \int_{-\infty}^{\infty} \frac{dk}{2\pi} e^{ikx} \tilde{f}(k, v, \tau) \quad (21)$$

$$= \int_0^{\infty} \frac{dk}{\pi} e^{ikx} \exp(C\tau + D\tau v)\tilde{f}(k, v, 0) \quad (22)$$

$$= \int_0^{\infty} \frac{dk}{\pi} e^{ikx} \exp(C\tau + D\tau v)(\pi\delta(k) + \frac{1}{ik}) \quad (23)$$

$$= \frac{1}{2} + \int_0^{\infty} \frac{dk}{\pi} e^{ikx} \exp(\frac{C\tau + D\tau v}{ik}) \quad (24)$$

Now the only remaining task is to find C and D . If we substitute 21 into 19, we can get the expression C and D .

References

- [1] SL Heston, *The Review of Financial Studies*, "A Closed-Form Solution for Options with Stochastic Volatility with Applications to Bond and Currency Options" Vol. 6, Issue 2, April 1993, Pages 327-343
- [2] Francis Longstaff, Eduardo Schwartz, "Valuing American Options by Simulation: A Simple Least-Squares Approach" *The Review of Financial Studies*, Vol. 14, Issue 1, January 2001, pp. 113-147.
- [3] Roger Lord, Remmert Koekkoek, Dick Van Dijk, "A comparison of biased simulation schemes for stochastic volatility models", *Quantitative Finance*, Vol. 10, pp. 177-194, 2010.