

In order to describe the conductivity, physicists have developed several models, starting from classical model, semiclassical model to quantum model. These model can be summarized as:

- 1) In classical model, the electrons are treated classically, and the movement is governed by the Newton's law. This model is good enough to explain the Ohm's law.
- 2) In semiclassical model, we consider the behavior of electron as a wave, and treat the movement of electrons as the propagation of the wavepacket, therefore the velocity of the electron is group velocity of the wave. This model utilize the particle-wave duality, and is capable of explaining the conduction in metal.

1 Current Operator

Derivation of velocity operator

In quantum mechanics, the velocity operator is defined as

$$v = \frac{\partial H}{\partial p} = \frac{\partial H}{\hbar \partial k}$$

There exist several ways to understand this. First, we can recall the equation of motion in analytical mechanics. Given a Hamiltonian, the equation of motion is

$$v = \dot{q} = \frac{\partial H}{\partial p}$$

Wavefunction subject to adiabatic evolution

$$|\Psi(t)\rangle = \exp\left(-\frac{i}{\hbar} \int_{t_0}^t dt' E_n(t')\right) (|n\rangle - i\hbar \sum_{n' \neq n} |n'\rangle \frac{\langle n' | \frac{\partial}{\partial t} | n \rangle}{E_n - E_{n'}})$$

The expectation value of the velocity

$$\bar{v}(k, t) = \frac{\partial E_n(k)}{\hbar \partial k} - i \sum_{n' \neq n} (\langle n | \frac{\partial H}{\partial k} | n' \rangle \frac{\langle n' | \frac{\partial}{\partial t} | n \rangle}{E_n - E_{n'}} - c.c.)$$

$$\bar{v}(k, t) = \frac{\partial E_n(k)}{\hbar \partial k} - i (\langle \frac{\partial n}{\partial k} | \frac{\partial n}{\partial t} \rangle - \langle \frac{\partial n}{\partial t} | \frac{\partial n}{\partial k} \rangle)$$

Where the second term is the Berry phase. The current operator is

$$j = -2e \frac{dk}{2\pi} f(k) v(k)$$