

Bridging Gravity and Quantum Mechanics: A Study of Gravitationally Induced Entanglement

Prateek Gupta

18th August 2023



MSc in Theoretical Physics
The University of Edinburgh
2023

Abstract

Understanding gravity in the framework of quantum mechanics is one of the most significant challenges in modern physics. The challenge comes from the clash between general relativity and quantum mechanics. General Relativity has proven to be extraordinarily incompatible with the framework of Quantum Field Theory. So much so that some even argue whether gravity could be quantized in the first place. Several attempts have been made to probe this question of the nature of gravity. This paper details one of the alternative directions to probe this question: by studying the entanglement that could be produced between two bodies if gravity was a quantum entity. We consider the example of two harmonic oscillators and consider two cases. One when they are static and one when they are non-static. By relying on quantum perturbation theory, we compute entanglement entropy and concurrence for each case for a quantum gravitational interaction. We show that these two quantities are positive for all cases and that should gravity be a quantum entity, these values can be tested experimentally. We finally conclude by considering what kind of future work can be undertaken in this new direction.

Declaration

I declare that this dissertation was composed entirely by myself.

Chapters 1 and 2 provide an introduction to the subject area and a description of previous work on this topic. They do not contain original research.

Chapter 3 describes work that was carried out entirely by me. The results of this chapter were previously carried out by Professor Suraj N. Gupta of the University of Manchester in 1952, but the results are modified to correspond with the metric adopted in this dissertation. Few of the other results were previously obtained by Professor Marios Christodoulou of the Austrian Academy of Sciences and the University of Vienna.

Chapter 4 and 5 describes work that was carried out entirely by me. Some of the results of this chapter have been obtained previously by Professor Sougato Bose of University College London, but our results vary slightly.

Chapters 4 also contains my original work. This original work in Chapter 4 was carried out in collaboration with Dr Suddhasattwa Brahma.

All calculations were done by hand. Jupyter, NumPy, SciPy and matplotlib packages of Python were used to visualize some of the results.

Personal Statement

This project began with an introduction to entanglement and gravity, with my main sources being S. Bose’s “Spin Entanglement Witness for Quantum Gravity”, “Mechanism for the Quantum Natured Gravitons to Entangle Masses”, and M. Christodoulou’s “Locally Mediated Entanglement in Linearised Quantum Gravity”. I read the first paper, and acquainted myself with the experimental setup and the thought experiments.

Once I was comfortable with that, I moved to S. Bose’s second paper, where I spent some time understanding how gravity was quantised, as the paper presented a mechanism for entanglement in a perturbative setting.

The next thing I worked on was reproducing the results from the second paper, before moving to original work. The goal was to understand how they calculated Concurrence, a measure of entanglement, and how the formalism worked.

The third stage of our project was to calculate Entanglement Entropy, another measure for Entanglement, and corroborate my results with those found using Concurrence. The second and third steps took the most time as I ran into some discrepancies that I had to sort out.

The fourth and final stage of my project was to reproduce the results of M. Christodoulou’s “Locally Mediated Entanglement in Linearised Quantum Gravity,” and try and understand how we could obtain our interaction Hamiltonian from first principles, and deepen my understanding of how first principles can be used to derive the Newtonian potential between N point masses covariantly.

During the course of the project, I met with my supervisor roughly every week, in order to discuss my progress and the direction I would head next.

I started writing this dissertation in around the end of July, and I spent the first three weeks of August working on it full-time.

Overall, I feel that the project was a success, and I found it to be extremely enjoyable throughout.

Acknowledgements

I'd like to thank my supervisor Dr Suddhasattwa Brahma for making this project possible, and for his immense patience and his detailed functional explanations of the thought experiments and mathematical expertise. Thanks also to my fellow MSc student Han Wang, also working under the supervision of Dr. Brahma on a similar topic, for being a sounding board for some of my ideas.

I would also like to thank my fellow MSc students Maegan Anderson and Joe Milarvie for their various inputs throughout the dissertation process.

The document was typeset using L^AT_EX and the template for this document was provided by the School of Physics for MSc Theoretical Physics.

Contents

1	Introduction	1
2	Background Theory	4
2.1	Quantum Entanglement and the BMV Experiment	4
2.2	Entanglement Due to Quantum Interactions	6
3	Quantization of Gravity and Quantum Gravitational Interactions	9
3.1	Linear Approximation For The Gravitational Field	9
3.2	Canonical Quantization of Gravity	11
3.3	Quantum Gravitational Interaction	13
3.4	Covariant Action for Gravitational Field of Moving Particles	14
3.4.1	Slow-motion Approximation	16
4	Entanglement via the Graviton	17
4.1	Entanglement in the Static Limit	17
4.2	Entanglement in the Non-Static Limit	21
4.2.1	Correction 1	24
4.2.2	Correction 2	25
4.3	Concurrence	25
4.3.1	Static Limit	26
4.3.2	Non-Static Limit	26
4.4	Entanglement Entropy	27
4.4.1	Static Limit	28
4.4.2	Non-Static Limit	28
5	Analysis and Discussion	29
6	Conclusions	34

List of Figures

2.1	Schematic Diagrams for the BMV Experiment.	5
5.1	Comparison of Coupling for Different Cases: (a) Static Case; (b) Non-static Correction 1 (c) Non Static Correction 2	30
5.2	Comparison of Concurrence for Different Cases: (a) Static Case; (b) Non- static Correction 1 (c) Non Static Correction 2	31
5.3	Comparison of the Perturbative Entanglement Entropy and its Approximation.	32
5.4	Comparison of Entanglement Entropies for Different Cases: (a) Static Case; (b) Non-static Correction 1 (c) Non Static Correction 2	33

Chapter 1

Introduction

To talk about quantum gravity, I will start with the story about the clash between General Relativity and Quantum Mechanics and how it gave birth to one of the biggest crises known to Physics.

Ten years after Einstein shook the world with the *Annus Mirabilis* papers, which gave the world the Theory of Special Relativity and the world-famous equation for mass-energy equivalence, $E = mc^2$, he shocked the world of physics once again with his General Theory of Relativity, in a series of papers published in 1915. Furthermore, by the end of 1915, he published the Field Equations of Gravity, providing a description of gravity as a geometric property of space and time and how the curvature of the four-dimensional spacetime was directly related to the energy and momentum of matter and radiation [1]. And thus, we had the first classical field theory of gravity. This also forms the basis of the Standard Model of Cosmology we use today.

Around the same time, a group of physicists which included the likes of Max Born, Werner Heisenberg and Wolfgang Pauli, made great strides in the field of quantum mechanics, which govern the mechanics of sub-atomic particles [2, 3]. However, by the mid-1920s, they hit a seemingly insurmountable problem: quantising the electromagnetic field. They found that to quantise electromagnetism correctly, and they had to quantise the classical field theory of electromagnetism [4]. Furthermore, since the field equations could be rewritten using Einstein's Relativity theory, they had to incorporate special relativity into their work. Thus, the framework of Quantum Field Theory was born, and by 1927, Paul Dirac published the first theory of Quantum Electrodynamics or QED [5].

Nevertheless, as in any developmental story of a theory, they found more problems. One such problem was the problem of infinities. In the early 1930s, Robert Oppenheimer showed that higher-order perturbative calculations of QED always resulted in infinite values, such as the electron self-energy, which is the energy of the electron had as a result of changes that it causes in its environment, or the vacuum zero point energy of photon and electrons [6]. Thus, in the 1950s, the concept of renormalisation was introduced by Richard Feynman, Shinichiro Tomonaga [7, 8], Julian Schwinger [9], and Freeman Dyson, for which the first three of them even jointly won the Nobel Prize in Physics [10]. This framework allowed them to eliminate the problem of infinities in perturbative quantum field theory, and we finally had a more complete theory of Quantum Electrodynamics.

Thus, the field of Quantum Field Theory boomed. Physicists formulated a quantum field theory of the strong force, which they named Quantum Chromodynamics, unified the Weak and Electromagnetic forces, and theorised the Higgs Mechanism. Furthermore, by the 1970s, they formulated a Standard Model of Particle Physics incorporating all of these theories, and we now had a model where three of the four fundamental forces were governed by the laws of Quantum Field Theory.

Surely, physicists thought at the time, if Quantum Field Theory could describe three of the four fundamental forces, the final known fundamental force, gravity, would be no different. They thought that finally, finally, they could grasp a Unified Theory which could explain everything. Thus, Einstein's field theory equations, published in 1915, met the new and improved framework of Quantum Field Theory. However, as always, luck was not on the side of the physicists. It turned out that Einstein's Field Theory for Gravity was extraordinarily incompatible with the framework of Quantum Field Theory. Not only did classical General Relativity have singularities, like singularities in Black Holes and the Big Bang singularity, but it also had UV behaviour near Planck scales.

However, the biggest of these problems was that the field equations had a coupling with units, G_N , which made the field equations utterly non-renormalisable, and without renormalizability, a quantum field theory could not be generated. Several theories have been proposed to circumvent or counter this. Examples include String Theory, a framework in which the point-like particles are replaced by one-dimensional objects called strings [11]; Loop Quantum Gravity; Supergravity, which employs the framework of Superstring theory; or Double Copy Theory, which hypothesises a perturbative duality between gauge theory and gravity [12]. Moreover, many of these theories are highly untestable by experiments or have huge inconsistencies which are yet to be resolved.

Even then, one could say that General Relativity contains the seeds of its own destruction [11]. We can say that the singularities, the UV behaviour, and the non-renormalizability are all *known* unknowns or known problems that need to be solved for a theory of quantum gravity. However, the theory could host several *unknown* unknowns, problems we did not even know existed. One such problem, as some physicists have pointed out, mainly because of the lack of experiments, is the question about the nature of gravity itself. Is gravity even quantisable in the first place? Does it even have a quantum nature? Because if it is not quantisable and does not have a quantum nature, we must look into different theories to resolve the problems of singularities and divergent behaviour of General Relativity.

To this end, many physicists in the 21st Century have developed various gedankenexperiments [13, 14], or thought experiments, and tabletop experiments. I will briefly introduce one such experiment called the Bose-Marletto-Vedral (BMV) [15, 16] experiment in the next chapter, but the main feature of many of these experiments, including the BMV experiment, include entanglement due to gravitational interactions. This is because all quantum interactions induce entanglement, as I will also show in the next chapter. And if gravity can induce entanglement, then indeed, one could say with some certainty that gravity has a quantum nature.

In this dissertation, I will first explain the conceptual underpinnings of the BMV experiment, and how it uses a few central principles of Quantum Information Theory, how quantum interactions can induce entanglement within a system of two quantum harmonic

oscillators, and how to calculate two separate measures of entanglement called Concurrence and Entanglement Entropy [17]. I will then briefly describe how one can linearise Einstein's Field equations and then quantise them using canonical quantisation [18]. I will also describe how we could use this quantisation as a driving force for the interaction between the quantum oscillators.

I will also show how we can derive the same interaction Hamiltonian from first principles and how the slow-motion approximation of the action for an N point masses would yield an action with potential energy similar to that of a Newtonian gravitational potential energy [19].

I will then go back and describe a proposed mechanism for gravity to entangle the two quantum harmonic oscillators and show that if we were to calculate a shift in the energy of the quantised graviton from the earlier canonical quantisation is also the same as the Newtonian gravitational potential energy between two bodies. I will then derive the concurrence [17] and entanglement entropy, the two measures of entanglement, for these interactions. Then, I will analyse the results in more depth and study the analytical and numerical behaviour.

Finally, I will conclude with the implications and summary of the results and potential experiments that one can conduct to see evidence of this mechanism and the consequences of these results for the story of quantum gravity.

Chapter 2

Background Theory

2.1 Quantum Entanglement and the BMV Experiment

Before getting into the theory behind the BMV Experiment, let us first define Quantum Entanglement. Quantum Entanglement is defined as the phenomenon where a pair of particles interact in a way such that the state of each particle cannot be described independently of the state of the other, even if they are at large distances [20]. Erwin Schrödinger defined this as the *characteristic* trait of a quantum interaction [20].

As stated before, due to the lack of evidence for quantum aspects, there is a great debate about the nature of gravity [15, 21]. Some even use the breakdown of quantum mechanics itself at massive scales capable of producing gravitational effects to justify that gravity need not be a quantised field [22]. A few of the proposed tests aimed at probing this question of gravity’s quantum nature have traditionally focused on phenomenology and cosmological observations [15, 23]. However, Bose *et al.* [15] and Marletto and Vedral [16] opened up a different avenue to study the same question: probing the quantum coherent aspect of gravity, which seems to offer a much more prominent and unambiguous witness to quantum gravity [15].

In the same spirit as a Bell inequality certifies the “non-local” character of quantum mechanics [24], the BMV thought experiment shows that the growth of entanglement between two mesoscopic masses held in a superposition of spatially localised states $|L\rangle$ and $|R\rangle$ (as shown in Fig. 2.1(a)[15]), say in adjacent wave-matter interferometers as shown in Fig 2.1(b)[15], can be used to certify the quantum nature of gravitational interactions.

The proposal relies on the assumption that the gravitational interaction between the two masses is mediated by a gravitational field, and there are no other forces, like the short range Casimir-Polder force, in play. Upon making this assumption, the proposal borrows and uses a central principle of Quantum Information Theory, that entanglement between two systems *cannot* be created by Local Operations and Classical Communications (LOCC) [25]. Here, “Local Operations” means that the two subsystems are each allowed to do anything allowed by quantum mechanics to their subsystem but nothing that requires an interaction between them. “Classical Communication” means the subsystems “communicate” through a classical channel [15].




Figure 2.1: Schematic Diagrams for the BMV Experiment.

It can be proved that in the absence of closed timelike loops [26] and under the assumption of validity of the chronology protection conjecture [27], as long as the notion of classicity itself is not extended [28], LOCC keeps any initially unentangled state separable [15]. If gravitational interactions entangle the masses, the entanglement must have occurred through local operations and “quantum” communication (LOQC), where “quantum” is essentially anything that does not fall under the aforementioned notion of classicity. That is, if mutual gravitational interaction between two masses entangles them, then the mediating gravitational field is *necessarily* quantum mechanical in nature [15].

Consider now, the schematic version presented in Fig 2.1(a). Two neutral test masses m_1 and m_2 are held steadily in a superposition of two spatially separated states $|L\rangle$ and $|R\rangle$, whose centres are separated by a distance Δx and each state is a localised Gaussian wave packet with small enough width to assume $\langle L|R\rangle = 0$. There is a separation d between the centres of the superpositions so that the short range Casimir-Polder force is negligible even at the closest approach of the masses. As the masses are separated by different distances, distinct components of the superposition have distinct gravitational interaction energies and thus have different rates of phase evolution. Therefore the time evolution of the joint state of the two masses is purely due to gravitational interactions between them [15].

Fig 2.1(b) shows the Stern-Gerlach (SG) interferometry which includes three steps: (1) A spin dependent spatial splitting of the centre of mass state of a test mass in an inhomogenous magnetic field, (2) “Holding” the coherent superposition created above for a time τ , where the magnetic field of the SG is effectively turned off and finally (3) bring back the superposition through unitary transformations. If the state is proven to be entangled, then the mediator, the gravitational field, is a quantum entity. A more detailed scheme is presented in Bose *et al.* [15].

I have now explained the conceptual theory behind the BMV experiment. Detailed information and a nice review about this experiment and its workings, as well as the

demanding nature of this experiment, can be found in a separate work done by Han Wang [29] for his Master's dissertation (submitted 2023), and in Christodoulou and Rovelli [30]. The main focus of this dissertation is the review and analysis of the concepts behind gravitationally induced entanglement using these concepts.

2.2 Entanglement Due to Quantum Interactions

Let us now study quantum interactions and how they induce entanglement. Consider a setup of two Quantum Harmonic Oscillators (QHOs) in the presence of a gravitational field, denoted by A and B, located at $\pm d/2$. Assume that the harmonic oscillators are well-localised, such that,

$$\hat{x}_A = -\frac{d}{2} + \delta\hat{x}_A, \quad \hat{x}_B = \frac{d}{2} + \delta\hat{x}_B, \quad (2.1)$$

where \hat{x}_A , \hat{x}_B are the positions, and $\delta\hat{x}_A$, $\delta\hat{x}_B$ denote small displacements from the equilibrium. Thus the Hamiltonian for the two QHOs is given by [17, 31, 32],

$$\hat{H}_{\text{matter}} = \frac{\hat{p}_A^2}{2m} + \frac{\hat{p}_B^2}{2m} + \frac{m\omega_m^2}{2}\delta\hat{x}_A^2 + \frac{m\omega_m^2}{2}\delta\hat{x}_B^2, \quad (2.2)$$

where \hat{p}_A and \hat{p}_B are the conjugate momenta, and ω_m is the harmonic frequency of the QHOs. We will assume, for simplicity, that the aforementioned frequency is equal for both QHOs. Then, the mode operators can be introduced using,

$$\delta\hat{x}_A = \sqrt{\frac{\hbar}{2m\omega_m}}(\hat{a} + \hat{a}^\dagger), \quad \delta\hat{x}_B = \sqrt{\frac{\hbar}{2m\omega_m}}(\hat{b} + \hat{b}^\dagger), \quad (2.3)$$

$$\hat{p}_A = i\sqrt{\frac{\hbar m\omega_m}{2}}(\hat{a} - \hat{a}^\dagger), \quad \hat{p}_B = i\sqrt{\frac{\hbar m\omega_m}{2}}(\hat{b} - \hat{b}^\dagger), \quad (2.4)$$

where the mode operators follow the usual commutation relations $[\hat{a}, \hat{a}^\dagger] = 1$ and $[\hat{b}, \hat{b}^\dagger] = 1$. These results and derivations can be found in any introductory quantum mechanics textbook such as Griffiths and Schroeter [31] and Cohen-Tannoudji *et al.* [32]. Thus, the Hamiltonian can be written as,

$$\hat{H}_{\text{matter}} = \hat{H}_A + \hat{H}_B, \quad (2.5)$$

where $\hat{H}_A = \hbar\omega_m\hat{a}^\dagger\hat{a}$, and $\hat{H}_B = \hbar\omega_m\hat{b}^\dagger\hat{b}$.

Let us now investigate the initial state of the matter system when perturbed by an interaction Hamiltonian H_{AB} .

Let us assume that the initial state of the matter system is given by,

$$|\psi_i\rangle = |0\rangle_A |0\rangle_B, \quad (2.6)$$

where $|0\rangle_A$ and $|0\rangle_B$ denote the ground state of the first and second QHOs respectively. In future equations, we will use this same order, and omit the subscripts for ease of notation.

Suppose a perturbation is introduced using an interaction potential λH_{AB} , where λ is a small parameter, between the two systems. Then the perturbed state is given by,

$$|\psi_f\rangle = \frac{1}{\sqrt{\mathcal{N}}} \sum_{n,N} C_{nN} |n\rangle |N\rangle \quad (2.7)$$

where $|n\rangle, |N\rangle$ denote number states of A and B respectively, while \mathcal{N} is the normalisation given by $\mathcal{N} = \sum_{n,N} |C_{nN}|^2$. We know from Eq. 2.6 that the coefficient of the unperturbed state is $C_{00} = 1$. The remaining coefficients are given by [17, 33],

$$C_{nN} = \lambda \frac{\langle n | \langle N | \hat{H}_{AB} | 0 \rangle | 0 \rangle}{2E_0 - E_n - E_N}, \quad (2.8)$$

where E_0 is the ground state (assumed to be the same for both systems for simplicity), and E_n and E_N are the excited state energies. Note that the role of \hat{H}_{AB} is that of a quantum operator. Were it to be a classical operator, then in Eq. 2.8, it would be a complex number which would come out of the inner product thus giving us $\langle n, N | 0, 0 \rangle$, which would give us zero by the orthogonality of the states for $n, N > 0$. Thus the coefficients would all be zero for $n, N > 0$ and the quantum state would *remain unperturbed* by the classical interaction potential. Alternatively, we can say that now the difference between LOQC and LOCC is encoded in \hat{H}_{AB} .

If we set $A_n \equiv C_{n0}$ and $B_N \equiv C_{0N}$, then we can write the perturbed state as [33],

$$\begin{aligned} |\psi_f\rangle &= \frac{1}{\sqrt{\mathcal{N}}} \left(|0\rangle |0\rangle + \sum_{n>0} A_n |n\rangle |0\rangle + \sum_{N>0} B_N |0\rangle |N\rangle + \sum_{n,N>0} C_{nN} |n\rangle |N\rangle \right) \\ &= \frac{1}{\sqrt{\mathcal{N}}} \left(\left(|0\rangle + \sum_{n>0} A_n |n\rangle \right) |0\rangle + \left(|0\rangle + \sum_{n>0} A_n |n\rangle \right) \sum_{N>0} B_N |N\rangle \right. \\ &\quad \left. + \sum_{n,N>0} (C_{nN} - A_n B_N) |n\rangle |N\rangle \right) \\ &= \frac{1}{\sqrt{\mathcal{N}}} \left(\left(|0\rangle + \sum_{n>0} A_n |n\rangle \right) \left(|0\rangle + \sum_{N>0} B_N |N\rangle \right) + \sum_{n,N>0} (C_{nN} - A_n B_N) |n\rangle |N\rangle \right) \end{aligned} \quad (2.9)$$

Here, the first term would yield a separable state and will not be responsible for the entanglement of the system, as interactions acting as operators on only one of the two quantum systems cannot entangle the two systems [17]. Thus the A_n and B_N terms will not contribute to the entanglement in first order perturbation theory. The second term in the equation is responsible for the entanglement between the two matter systems. This shows the stark difference between LOCC and LOQC as explained in the previous chapter. The non-trivial part of an LOQC mechanism encoded in the Hamiltonian produces the second term in the last line of the above equation, and LOCC can only produce the first term, not the second [17].

To compute the degree of entanglement, one can calculate two different quantities, the *Concurrence* [34, 35], and the *von Neumann Entanglement Entropy* [33, 36, 37].

The Concurrence is defined by [34, 35],

$$C \equiv \sqrt{2(1 - \text{Tr}[\hat{\rho}_A^2])}, \quad (2.10)$$

where $\hat{\rho}_A$ is the density operator, and can be computed by tracing away the B state as,

$$\begin{aligned} \hat{\rho}_A &= \sum_N \langle N | \psi_f \rangle \langle \psi_f | N \rangle \\ &= \frac{1}{\mathcal{N}} \sum_{n, n', N} C_{nN} |n\rangle \langle N | N \rangle C_{n'N}^* \langle n' | \langle N | N \rangle \\ &= \frac{1}{\mathcal{N}} \sum_{n, n', N} C_{nN} C_{n'N}^* |n\rangle \langle n'| \end{aligned} \quad (2.11)$$

Thus the trace of $\hat{\rho}_A^2$ is given by,

$$\begin{aligned} \text{Tr}[\hat{\rho}_A^2] &= \text{Tr} \left[\frac{1}{\mathcal{N}} \sum_{n, n', N} C_{nN} C_{n'N}^* |n\rangle \langle n'| \frac{1}{\mathcal{N}} \sum_{n'', n''', N'} C_{n''N'} C_{n'''N'}^* |n''\rangle \langle n'''| \right] \\ &= \frac{1}{\mathcal{N}^2} \sum_{n, n', N, N'} C_{nN} C_{n'N}^* C_{n'N'} C_{nN'}^* \end{aligned} \quad (2.12)$$

Therefore, the concurrence is given by,

$$C = \sqrt{2 \left(1 - \frac{1}{\mathcal{N}^2} \sum_{n, n', N, N'} C_{nN} C_{n'N}^* C_{n'N'} C_{nN'}^* \right)} \quad (2.13)$$

As it can be seen in Eq. 2.13, for a system with an infinite dimensional Hilbert space¹, the range for the Concurrence is $C \in [0, \sqrt{2}]$. It is 0 for a separable state and $\sqrt{2}$ for a maximally entangled state. Similarly, for a separable state, the trace of the density operator squared will be 1 and 0 for a maximally entangled system.

The other measurement of the degree of entanglement is the von Neumann Entanglement Entropy [33, 36, 37]. That is given by,

$$\mathcal{S}(\hat{\rho}_A) = -\text{Tr}(\hat{\rho}_A \log \hat{\rho}_A) \quad (2.14)$$

This can be calculated relatively easily by calculating the logarithm of the density operator, and remembering that for a diagonalizable matrix M , $\log M = P \log D P^{-1}$, and for a diagonal matrix D , $\log D = \text{diag}(\log d_1, \dots, \log d_n)$. If, in Eq. 2.8, λ is small enough, then to $\mathcal{O}(\lambda^3)$, we can calculate the von Neumann Entanglement Entropy to be [33],

$$\mathcal{S}(\hat{\rho}_A) = -\lambda^2 \log(\lambda^2) \sum_{n, N > 0} \frac{|\langle n | \langle N | \hat{H}_{AB} | 0 \rangle | 0 \rangle|^2}{(2E_0 - E_n - E_N)^2} + \mathcal{O}(\lambda^3). \quad (2.15)$$

Both the Concurrence and Entanglement Entropy grow as entanglement grows. In Chapter 4, I will show how these quantities can be calculated in a perturbative fashion for entanglement due to gravitational interactions.

¹Technically, the maximum of C as in Eq. 2.10 and Eq. 2.13 depends on the minimum value of $\text{Tr}[\hat{\rho}_A^2]$, also known as “Purity” of the system in Quantum Information Theory, which is dependent on the number of dimensions of the Hilbert space of the system. More on this in Chapter 5.

Chapter 3

Quantization of Gravity and Quantum Gravitational Interactions

In the previous chapter, we saw how quantum interactions induce entanglement. Now, we want to use gravity as the source of the quantum interaction, through a gravitational field. And as we want to consider the above setup in a gravitational field, we must first understand how the gravitational field is quantised, since we want to work in the regime of small perturbations about the Minkowski background. This section will give a brief overview of the linearization and quantization process of the gravitational field, following Gupta [18] closely. The metric is given by $g_{\mu\nu}$, where $\mu, \nu = 0, 1, 2, 3$ and we will use the $(-, +, +, +)$ signature throughout.¹

3.1 Linear Approximation For The Gravitational Field

The fundamental equation for the Gravitational Field is given by Einstein's equation [18, 38],

$$R^{\mu\nu} - \frac{1}{2}Rg^{\mu\nu} = -\frac{1}{2}\kappa^2 T^{\mu\nu}, \quad (3.1)$$

where $R^{\mu\nu}$ and R are the Ricci Tensor and Scalar respectively, κ is a constant such that $\kappa^2 = \frac{8\pi G}{c^4}$, and $T^{\mu\nu}$ is the Stress-Energy tensor of the matter field.

Since the covariant divergence of the Stress-Energy tensor is 0, one can obtain an expression for the Stress-Energy tensor *density* $\mathfrak{T}^{\mu\nu} = \sqrt{-g}T^{\mu\nu}$ (where g is the determinant of the metric) [18],

$$\frac{\partial \mathfrak{T}_\mu^\nu}{\partial x^\nu} - \frac{1}{2}\mathfrak{T}^{\alpha\beta} \frac{\partial g_{\alpha\beta}}{\partial x^\mu} = 0. \quad (3.2)$$

We can thus express the conservation of energy and momentum as,

$$\frac{\partial}{\partial x^\nu}(\mathfrak{T}_\mu^\nu + t_\mu^\nu) = 0, \quad (3.3)$$

¹Gupta [18] uses a slightly different signature, such that their spacetime coordinates are (it, x_1, x_2, x_3) . We will modify their work slightly to correspond to our chosen signature.

where $t^{\mu\nu}$ is the Stress-Energy *pseudo-tensor density* [18] for the gravitational field, thus satisfying the equation,

$$\frac{\partial t_{\mu}^{\nu}}{\partial x^{\nu}} = -\frac{1}{2}\mathfrak{T}_{\alpha\beta}\frac{\partial g_{\alpha\beta}}{\partial x^{\mu}}. \quad (3.4)$$

Following Gupta [18] and Einstein [38] we can obtain a linear approximation for the field by considering small perturbations $h_{\mu\nu}$, where $|h_{\mu\nu}| \ll 1$, around the Minkowski background. Thus, the metric becomes,

$$g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}, \quad (3.5)$$

where $\eta_{\mu\nu} = \text{diag}(-1, +1, +1, +1)$. Then using the flat spacetime tensor notation, we can rewrite Eq. 3.1 and Eq. 3.4 as [18],

$$\square^2 h_{\mu\nu} - \left(\frac{\partial^2 h_{\mu\lambda}}{\partial x_{\nu}\partial x_{\lambda}} + \frac{\partial^2 h_{\nu\lambda}}{\partial x_{\mu}\partial x_{\lambda}} \right) + \frac{\partial^2 h_{\lambda\lambda}}{\partial x_{\mu}\partial x_{\nu}} + \eta_{\mu\nu} \left(\frac{\partial^2 h_{\lambda\rho}}{\partial x_{\lambda}\partial x_{\rho}} - \square^2 h_{\lambda\lambda} \right) = \kappa^2 T_{\mu\nu}, \quad (3.6)$$

$$\frac{\partial t_{\mu\nu}}{\partial x_{\nu}} = \frac{\kappa^2}{2} \frac{\partial h_{\nu\lambda}}{\partial x_{\mu}} T_{\nu\lambda} \quad (3.7)$$

where $\square^2 = \partial^{\mu}\partial_{\mu} = \eta^{\mu\nu}\partial_{\nu}\partial_{\mu}$. According to Einstein [38], we can also choose coordinate/supplementary conditions given by,

$$\frac{\partial h_{\mu\nu}}{\partial x_{\nu}} - \frac{1}{2} \frac{\partial h_{\lambda\lambda}}{\partial x_{\mu}} = 0. \quad (3.8)$$

When this condition is applied, Eq. 3.6 reduces to,

$$\square^2 h_{\mu\nu} - \frac{1}{2} \eta_{\mu\nu} \square^2 h_{\lambda\lambda} = \kappa^2 T_{\mu\nu}. \quad (3.9)$$

It is now convenient to decompose $h_{\mu\nu}$ as,

$$h_{\mu\nu} = \gamma_{\mu\nu} - \frac{1}{2} \eta_{\mu\nu} \gamma_{\lambda\lambda}. \quad (3.10)$$

Thus, Eq. 3.7, Eq. 3.8 and Eq. 3.9 reduce to,

$$\frac{\partial t_{\mu\nu}}{\partial x_{\nu}} = \frac{\kappa^2}{2} \frac{\partial \gamma_{\nu\lambda}}{\partial x_{\mu}} T_{\nu\lambda} - \frac{\kappa^2}{4} \frac{\partial \gamma_{\lambda\lambda}}{\partial x_{\mu}} T_{\nu\nu}, \quad (3.11)$$

$$\frac{\partial \gamma_{\mu\nu}}{\partial x_{\nu}} = 0, \quad (3.12)$$

$$\square^2 \gamma_{\mu\nu} = \kappa^2 T_{\mu\nu}. \quad (3.13)$$

Thus, equations 3.13, 3.12 and 3.11 describe the linearised gravitational field [18].

It should be noted that the field equation Eq. 3.13 and the supplementary condition Eq. 3.12 are only compatible in an approximate sense, as they respectively give,

$$\square^2 \frac{\partial \gamma_{\mu\nu}}{\partial x_{\nu}} = \kappa^2 \frac{\partial T_{\mu\nu}}{\partial x_{\nu}} \quad \text{and} \quad \square^2 \frac{\partial \gamma_{\mu\nu}}{\partial x_{\nu}} = 0, \quad (3.14)$$

which only agree in the first approximation [18].

Thus solving for $t_{\mu\nu}$, one gets [18],

$$t_{\mu\nu} = \frac{1}{2} \left[\frac{\partial\gamma_{\lambda\rho}}{\partial x_\mu} \frac{\partial\gamma_{\lambda\rho}}{\partial x_\nu} - \frac{1}{2} \frac{\partial\gamma_{\lambda\lambda}}{\partial x_\mu} \frac{\partial\gamma_{\rho\rho}}{\partial x_\nu} - \frac{1}{2} \eta_{\mu\nu} \left(\frac{\partial\gamma_{\lambda\rho}}{\partial x_\sigma} \frac{\partial\gamma_{\lambda\rho}}{\partial x_\sigma} - \frac{1}{2} \frac{\partial\gamma_{\lambda\lambda}}{\partial x_\sigma} \frac{\partial\gamma_{\rho\rho}}{\partial x_\sigma} \right) \right] \quad (3.15)$$

Thus, we can get the Hamiltonian Density as,

$$H = t_{00} = \frac{1}{2} \left[\frac{\partial\gamma_{\lambda\rho}}{\partial t} \frac{\partial\gamma_{\lambda\rho}}{\partial t} - \frac{1}{2} \frac{\partial\gamma_{\lambda\lambda}}{\partial t} \frac{\partial\gamma_{\rho\rho}}{\partial t} + \frac{1}{2} \left(\frac{\partial\gamma_{\lambda\rho}}{\partial x_\sigma} \frac{\partial\gamma_{\lambda\rho}}{\partial x_\sigma} - \frac{1}{2} \frac{\partial\gamma_{\lambda\lambda}}{\partial x_\sigma} \frac{\partial\gamma_{\rho\rho}}{\partial x_\sigma} \right) \right] \quad (3.16)$$

3.2 Canonical Quantization of Gravity

We can now write the Lagrangian Density for the linear gravitational field as,

$$L = -\frac{1}{4} \left(\frac{\partial\gamma_{\mu\nu}}{\partial x_\lambda} \frac{\partial\gamma_{\mu\nu}}{\partial x_\lambda} - \frac{1}{2} \frac{\partial\gamma}{\partial x_\lambda} \frac{\partial\gamma}{\partial x_\lambda} \right), \quad (3.17)$$

where $\gamma \equiv \eta_{\mu\nu}\gamma^{\mu\nu}$ and $h_{\mu\nu} = \gamma_{\mu\nu} - \frac{1}{2}\eta_{\mu\nu}\gamma$. γ is Spin-0 and $\gamma_{\mu\nu}$ is Spin-2. The field equations and the Hamiltonian Density are,

$$\square^2\gamma_{\mu\nu} = 0, \quad \square^2\gamma = 0 \quad (3.18)$$

$$H = \frac{1}{2} \left[\frac{\partial\gamma_{\mu\nu}}{\partial t} \frac{\partial\gamma_{\mu\nu}}{\partial t} - \frac{1}{2} \frac{\partial\gamma}{\partial t} \frac{\partial\gamma}{\partial t} + \frac{1}{2} \left(\frac{\partial\gamma_{\mu\nu}}{\partial x_\lambda} \frac{\partial\gamma_{\mu\nu}}{\partial x_\lambda} - \frac{1}{2} \frac{\partial\gamma}{\partial x_\lambda} \frac{\partial\gamma}{\partial x_\lambda} \right) \right]. \quad (3.19)$$

Now, for the commutation relations for $\gamma_{\mu\nu}$ and γ , first let us consider their canonical conjugates and the commutation relations between them and the canonical conjugates. Let us remember that similar to the case of the Klein-Gordon Field as a Harmonic Oscillator and in Space-Time [39], the equal time commutation relations between the γ -fields and their conjugates are $[\gamma_{\mu\nu}(x), \pi_{\mu\nu}(x')] = i\delta(\mathbf{x} - \mathbf{x}')$ and $[\gamma(x), \pi(x')] = i\delta(\mathbf{x} - \mathbf{x}')$, where \mathbf{x}, \mathbf{x}' are three-vectors and $\pi_{\mu\nu}(x)$ and $\pi(x)$ are the canonical conjugates of $\gamma_{\mu\nu}(x)$ and $\gamma(x)$ respectively. First let us consider the diagonal elements of $\gamma_{\mu\nu}$. The canonical conjugate of γ_{00} is,

$$\pi_{00} = \frac{\partial L}{\partial(\partial\gamma_{00}/\partial t)} = \frac{1}{2} \frac{\partial\gamma_{00}}{\partial t}. \quad (3.20)$$

Thus the commutation relation between γ_{00} and π_{00} gives us,

$$\begin{aligned} [\gamma_{00}(x), \pi_{00}(x')] &= i\delta(\mathbf{x} - \mathbf{x}') \\ \implies [\gamma_{00}(x), (\partial\gamma_{00}/\partial t)(x')] &= 2i\delta(\mathbf{x} - \mathbf{x}') \end{aligned} \quad (3.21)$$

Thus, the ETCR is,

$$[\gamma_{00}(x), \dot{\gamma}_{00}(x')] = 2i\delta(\mathbf{x} - \mathbf{x}'), \quad (3.22)$$

Similarly,

$$[\gamma_{11}(x), \dot{\gamma}_{11}(x')] = [\gamma_{22}(x), \dot{\gamma}_{22}(x')] = [\gamma_{33}(x), \dot{\gamma}_{33}(x')] = 2i\delta(\mathbf{x} - \mathbf{x}'). \quad (3.23)$$

Now consider off-diagonal elements of $\gamma_{\mu\nu}$. By our decomposition, $\gamma_{\mu\nu}$ is symmetric. That is, $\gamma_{\mu\nu} = \gamma_{\nu\mu}$. Now consider the case of an off-diagonal element, γ_{12} . Since $\gamma_{12} = \gamma_{21}$, for the canonical conjugate, we get,

$$\pi_{12} = \frac{\partial L}{\partial(\partial\gamma_{12}/\partial t)} = \frac{\partial\gamma_{12}}{\partial t} = \frac{\partial\gamma_{21}}{\partial t} = \pi_{21}. \quad (3.24)$$

Thus, the commutation relation we get is,

$$[\gamma_{12}(x), \dot{\gamma}_{12}(x')] = i\delta(\mathbf{x} - \mathbf{x}'). \quad (3.25)$$

This is the same for all off-diagonal elements. Thus we can write that the commutation relation using the above relations is given by,

$$[\gamma_{\mu\nu}(x), \dot{\gamma}_{\lambda\rho}(x')] = i(\eta_{\mu\lambda}\eta_{\nu\rho} + \eta_{\mu\rho}\eta_{\nu\lambda})\delta(\mathbf{x} - \mathbf{x}'). \quad (3.26)$$

Using Schwinger Notation [9, 18], we can write this as,

$$[\gamma_{\mu\nu}(x), \gamma_{\lambda\rho}(x')] = i(\eta_{\mu\lambda}\eta_{\nu\rho} + \eta_{\mu\rho}\eta_{\nu\lambda})D(x - x'), \quad (3.27)$$

where $D(x - y)$ is a Green's function similar to the Feynman Propagator for a K-G field (see Peskin and Schroeder [39, §2.4, Eq. 2.59] for more details). Similarly, for the canonical conjugate of γ ,

$$\pi = \frac{\partial L}{\partial(\partial\gamma/\partial t)} = -\frac{1}{4} \frac{\partial\gamma}{\partial t}, \quad (3.28)$$

yielding the commutation relation as,

$$[\gamma(x), \gamma(x')] = -4iD(x - x'). \quad (3.29)$$

Now let us promote the fluctuations around the Minkowski background into quantum operators, giving the mode expansion as [17],

$$\hat{h}_{\mu\nu} = \mathcal{A} \int d\mathbf{k} \sqrt{\frac{\hbar}{2\omega_k(2\pi)^3}} (\hat{P}_{\mu\nu}^\dagger(\mathbf{k})e^{-i\mathbf{k}\cdot\mathbf{r}} + \hat{P}_{\mu\nu}(\mathbf{k})e^{i\mathbf{k}\cdot\mathbf{r}}), \quad (3.30)$$

where $\mathcal{A} = \sqrt{16\pi G/c^2}$, \mathbf{k} is a three vector, and $\hat{P}_{\mu\nu}^\dagger$ & $\hat{P}_{\mu\nu}$ are the creation and annihilation operators respectively.

Then, using the decomposition of $\hat{h}_{\mu\nu}$, we get [17],

$$\hat{\gamma}_{\mu\nu} = \mathcal{A} \int d\mathbf{k} \sqrt{\frac{\hbar}{2\omega_k(2\pi)^3}} (\hat{P}_{\mu\nu}^\dagger(\mathbf{k})e^{-i\mathbf{k}\cdot\mathbf{r}} + \hat{P}_{\mu\nu}(\mathbf{k})e^{i\mathbf{k}\cdot\mathbf{r}}), \quad (3.31)$$

$$\hat{\gamma} = 2\mathcal{A} \int d\mathbf{k} \sqrt{\frac{\hbar}{2\omega_k(2\pi)^3}} (\hat{P}^\dagger(\mathbf{k})e^{-i\mathbf{k}\cdot\mathbf{r}} + \hat{P}(\mathbf{k})e^{i\mathbf{k}\cdot\mathbf{r}}). \quad (3.32)$$

Using the above commutation relations, and methods similar to those for finding the commutation relations of the creation and annihilation operators for a K-G field (see Peskin and Schroeder [39, chap. 2]), we get [17],

$$[\hat{P}_{\mu\nu}(\mathbf{k}), \hat{P}_{\lambda\rho}^\dagger(\mathbf{k}')] = (\eta_{\mu\lambda}\eta_{\nu\rho} + \eta_{\mu\rho}\eta_{\nu\lambda})\delta(\mathbf{k} - \mathbf{k}'), \quad (3.33)$$

$$[\hat{P}(\mathbf{k}), \hat{P}^\dagger(\mathbf{k}')] = -\delta(\mathbf{k} - \mathbf{k}'). \quad (3.34)$$

Substituting the mode expansions of $\gamma_{\mu\nu}$ and γ in the Hamiltonian Density (Eq. 3.19), and using the above commutation relations for the mode operators, we get that the Hamiltonian for the graviton is given by [17, 18],

$$\hat{H}_g = \int d\mathbf{k} \hbar\omega_k \left(\frac{1}{2} \hat{P}_{\mu\nu}^\dagger(\mathbf{k}) \hat{P}^{\mu\nu}(\mathbf{k}) - \hat{P}^\dagger(\mathbf{k}) \hat{P}(\mathbf{k}) \right). \quad (3.35)$$

This completes the canonical quantization of linear gravity.

3.3 Quantum Gravitational Interaction

For our purpose of quantifying the entanglement, we are more interested in the *shift* of the energy of the graviton vacuum, arising from interactions with matter. According to Weinberg [40], the operator valued interaction Hamiltonian is given by,

$$\hat{H}_{\text{int}} = -\frac{1}{2} \int d\mathbf{r} \hat{h}^{\mu\nu}(\mathbf{r}) \hat{T}_{\mu\nu}(\mathbf{r}), \quad (3.36)$$

where \mathbf{r} is a three-vector.

Consider the first order contribution to the shift in energy, given by $\langle 0 | \hat{H}_{\text{int}} | 0 \rangle$.

$$\begin{aligned} \langle 0 | \hat{H}_{\text{int}} | 0 \rangle &= \langle 0 | \left(-\frac{1}{2} \int d\mathbf{r} (\hat{\gamma}_{\mu\nu}(\mathbf{r}) - \frac{1}{2} \eta_{\mu\nu} \hat{\gamma}(\mathbf{r})) \hat{T}^{\mu\nu}(\mathbf{r}) \right) | 0 \rangle \\ &= -\frac{1}{2} \int d\mathbf{r} \langle 0 | (\hat{\gamma}_{\mu\nu}(\mathbf{r}) - \frac{1}{2} \eta_{\mu\nu} \hat{\gamma}(\mathbf{r})) | 0 \rangle \hat{T}^{\mu\nu}(\mathbf{r}) \end{aligned} \quad (3.37)$$

Now consider, $\langle 0 | (\hat{\gamma}_{\mu\nu}(\mathbf{r}) - \frac{1}{2} \eta_{\mu\nu} \hat{\gamma}(\mathbf{r})) | 0 \rangle$.

$$\begin{aligned} \langle 0 | (\hat{\gamma}_{\mu\nu}(\mathbf{r}) - \frac{1}{2} \eta_{\mu\nu} \hat{\gamma}(\mathbf{r})) | 0 \rangle &= \mathcal{A} \int d\mathbf{k} \sqrt{\frac{\hbar}{2\omega_k (2\pi)^3}} \left(\langle 0 | (\hat{P}_{\mu\nu}^\dagger(\mathbf{k}) e^{-i\mathbf{k}\cdot\mathbf{r}} + \hat{P}_{\mu\nu}(\mathbf{k}) e^{i\mathbf{k}\cdot\mathbf{r}}) | 0 \rangle \right. \\ &\quad \left. - \eta_{\mu\nu} \langle 0 | (\hat{P}^\dagger(\mathbf{k}) e^{-i\mathbf{k}\cdot\mathbf{r}} + \hat{P}(\mathbf{k}) e^{i\mathbf{k}\cdot\mathbf{r}}) | 0 \rangle \right) \end{aligned} \quad (3.38)$$

But, by the definition of creation and annihilation operators, $\langle 0 | \hat{P}_{\mu\nu}^\dagger = \hat{P}_{\mu\nu} | 0 \rangle = 0$ and $\langle 0 | \hat{P}^\dagger = \hat{P} | 0 \rangle = 0$. Thus,

$$\begin{aligned} \langle 0 | (\hat{\gamma}_{\mu\nu}(\mathbf{r}) - \frac{1}{2} \eta_{\mu\nu} \hat{\gamma}(\mathbf{r})) | 0 \rangle &= 0 \\ \implies \langle 0 | \hat{H}_{\text{int}} | 0 \rangle &= 0. \end{aligned} \quad (3.39)$$

This implies that the first order term vanishes, and the main contributor to the shift in energy is the second order term, given by [17],

$$\Delta \hat{H}_g \equiv \sum \int d\mathbf{k} \frac{\langle 0 | \hat{H}_{\text{int}} | \mathbf{k} \rangle \langle \mathbf{k} | \hat{H}_{\text{int}} | 0 \rangle}{E_0 - E_{\mathbf{k}}}, \quad (3.40)$$

where the sum indicates summation over all one particle projectors $|\mathbf{k}\rangle\langle\mathbf{k}|$, and $E_{\mathbf{k}} = E_0 + \hbar\omega_k$. We will soon see in Chapter 4 that for the non-relativistic limit (or the static limit), this shift is the same as the Newtonian gravitational potential energy.

3.4 Covariant Action for Gravitational Field of Moving Particles

We can also derive results similar to the previous section directly from the action for linearised gravity. We will first find the interaction term of the action from the action for the linearised gravity, following Christodoulou *et al.* [19][Supplementary Material] closely. The metric is once again given by $g_{\mu\nu}$, where $\mu, \nu = 0, 1, 2, 3$ and we will use the $(-, +, +, +)$ signature throughout. Keeping with Christodoulou *et al.* [19], let us also define the notation for the trace reversed tensor to be $\bar{A}_{\mu\nu} = A_{\mu\nu} - \frac{1}{2}\eta_{\mu\nu}A$, where $A_{\mu\nu}$ is an arbitrary tensor and $A = \eta_{\mu\nu}A^{\mu\nu}$.

Then, the action for linearised gravity coupled to matter is given by [41, 42],

$$S_h = \frac{c^4}{64\pi G} \int d^4x \left(-\partial_\rho h_{\mu\nu} \partial^\rho h^{\mu\nu} + 2\partial_\rho h_{\mu\nu} \partial^\nu h^{\mu\rho} - 2\partial_\nu h^{\mu\nu} \partial_\mu h + \partial^\mu h \partial_\mu h \right) + \frac{1}{2} \int d^4x h_{\mu\nu} T^{\mu\nu}, \quad (3.41)$$

where once again, $h_{\mu\nu}$ is the metric perturbation and $|h_{\mu\nu}| \ll 1$. The action is invariant under an infinitesimal change of co-ordinates. Let us take advantage of the gauge invariance of S_h , and apply the Lorenz gauge, where the field $h_{\mu\nu}$ satisfies,

$$\partial^\nu \bar{h}_{\mu\nu} = 0 \implies \partial^\nu h_{\mu\nu} - \frac{1}{2} \partial_\mu h = 0. \quad (3.42)$$

Now consider the second and third terms of the action. The second term, under this gauge, simplifies to,

$$\begin{aligned} 2\partial_\rho h_{\mu\nu} \partial^\nu h^{\mu\rho} &= \frac{2}{4} \partial^\mu h \partial_\mu h \\ &= \frac{1}{2} \partial^\mu h \partial_\mu h. \end{aligned} \quad (3.43)$$

The third term, under this gauge simplifies to,

$$\begin{aligned} -2\partial_\nu h^{\mu\nu} \partial_\mu h &= -\frac{2}{2} \partial_\mu h \partial^\mu h \\ &= -\partial_\mu h \partial^\mu h. \end{aligned} \quad (3.44)$$

Thus the overall action simplifies to,

$$S_h = \frac{c^4}{64\pi G} \int d^4x \left(-\partial_\rho h_{\mu\nu} \partial^\rho h^{\mu\nu} + \frac{1}{2} \partial^\mu h \partial_\mu h \right) + \frac{1}{2} \int d^4x h_{\mu\nu} T^{\mu\nu}. \quad (3.45)$$

The Euler-Lagrange equations for the field are [40, 43, 44],

$$\square^2 h_{\mu\nu} = -\frac{16\pi G}{c^4} \bar{T}_{\mu\nu} \quad (3.46)$$

Integrating the first term of the action by parts gives us two terms of the form $h_{\mu\nu} \square^2 h^{\mu\nu}$ and then by applying the E-L equations to those terms, we get the action [19],

$$S_h = \frac{1}{4} \int d^4x h_{\mu\nu} T^{\mu\nu}. \quad (3.47)$$

For a system of N point masses, $T_{\mu\nu}$ is defined as [19],

$$T^{\mu\nu}(t, \mathbf{x}) = \sum_{a=1}^N m_a \delta^3(\mathbf{x} - \mathbf{x}_a(t)) V_a^{\mu\nu}(t), \quad (3.48)$$

where,

$$V_a^{\mu\nu}(t) = \gamma_a(t) v_a^\mu(t) v_a^\nu(t), \quad (3.49)$$

where $v_a^\mu(t) = (c, \mathbf{v}_a)$, $\mathbf{v}_a = d\mathbf{x}_a/dt$ being the velocity of particle a and $\gamma_a(t) = (1 - |\mathbf{v}_a(t)|^2/c^2)^{-1/2}$ the corresponding Lorentz factor. Now we can solve the E-L equations using a retarded Green's function to the wave equation [43, 44]. A wave equation is usually of the form

$$\square^2 f(t, \mathbf{x}) = S(t, \mathbf{x}), \quad (3.50)$$

where $f(t, \mathbf{x})$ is the radiative field, and $S(t, \mathbf{x})$ is the source. The Green's function is a field that arises from delta function sources, that is, a Green's function $G(t, \mathbf{x}; t', \mathbf{x}')$ satisfies the equation

$$\square^2 G(t, \mathbf{x}; t', \mathbf{x}') = \delta(t - t') \delta^3(\mathbf{x} - \mathbf{x}'), \quad (3.51)$$

and one finds the wave solution $f(t, \mathbf{x})$ by integrating $G(t, \mathbf{x}; t', \mathbf{x}')$ against $S(t, \mathbf{x})$ [43]. The Green's function for the wave operator, \square^2 , is well known (e.g. see Jackson [45, §6.4]) and is given by,

$$G(t, \mathbf{x}; t', \mathbf{x}') = -\frac{1}{4\pi} \frac{\delta(t - t_r)}{|\mathbf{x} - \mathbf{x}'|}, \quad (3.52)$$

where $t_r = t - |\mathbf{x}' - \mathbf{x}|/c$ is the retarded time. Applying this against our E-L equations, we get,

$$h_{\mu\nu}(t, \mathbf{x}) = \frac{4G}{c^4} \int d^3x' \frac{\bar{T}_{\mu\nu}(t_r, \mathbf{x}')}{|\mathbf{x} - \mathbf{x}'|}. \quad (3.53)$$

Plugging in the expression for $T_{\mu\nu}$ from Eq. 3.48 we get,

$$h_{\mu\nu}(t, \mathbf{x}) = \frac{4G}{c^4} \sum_{a=1}^N m_a \int d^3x' \frac{\delta^3(\mathbf{x}' - \mathbf{x}_a(t_r)) \bar{V}_a^{\mu\nu}(t_r)}{|\mathbf{x} - \mathbf{x}'|}. \quad (3.54)$$

We can remove the dependence of t_r on \mathbf{x}' by integrating under a dummy variable t' and a delta function $\delta(t' - \tilde{t})$, where $\tilde{t} = \tilde{t}(\mathbf{x}, t, t') = t - |\mathbf{x} - \mathbf{x}_a(t')|/c$. Thus, we get [19],

$$h_{\mu\nu}(t, \mathbf{x}) = \frac{4G}{c^4} \sum_a m_a \int d^3x' \int dt' \frac{\bar{V}_{\mu\nu}(t') \delta(\mathbf{x}' - \mathbf{x}_a(t')) \delta(t' - \tilde{t})}{|\mathbf{x} - \mathbf{x}'|}. \quad (3.55)$$

We can now perform the spatial integration to get [19],

$$h_{\mu\nu}(t, \mathbf{x}) = \frac{4G}{c^4} \sum_a m_a \int dt' \frac{\bar{V}_{\mu\nu}(t') \delta(t' - \tilde{t}(\mathbf{x}, t, t'))}{|\mathbf{x} - \mathbf{x}_a(t')|}. \quad (3.56)$$

To complete the integration, we will use an identity of Dirac Delta functions, which is,

$$\delta(f(x)) = \sum_i \frac{\delta(x - x_i)}{|\partial_x f(x_i)|}, \quad (3.57)$$

where x_i are zeroes of $f(x)$. Applying this to the delta function within our integral, we get,

$$\delta(t' - \tilde{t}(\mathbf{x}, t, t')) = \frac{\delta(t' - t_a)}{1 - \mathbf{d}_a \cdot \mathbf{v}_a(t_a)/(d_a c)}, \quad (3.58)$$

where t_a is the retarded time satisfying $c(t - t_a) = |\mathbf{x} - \mathbf{x}_a(t_a)|$ and d_a is the magnitude of the retarded displacement $\mathbf{d}_a = \mathbf{x} - \mathbf{x}_a(t_a)$.

Thus, the solution to the field equations is,

$$h^{\mu\nu}(t, \mathbf{x}) = \frac{4G}{c^4} \sum_a \frac{m_a \bar{V}_a^{\mu\nu}(t_a)}{d_a - \mathbf{d}_a \cdot \mathbf{v}_a(t_a)/c}. \quad (3.59)$$

This equation is the gravitational analogue of the Liénard-Weichert Potential in Electrodynamics [45] and is already known [19, 46]. Plugging this solution and the expression for $T_{\mu\nu}$ from Eq. 3.48 in Eq. 3.47, we get the action for N interacting point masses as [19],

$$S_h = \frac{G}{c^4} \sum_{a,b}^{a \neq b} \int dt \frac{m_a m_b \bar{V}_a^{\mu\nu}(t_{ab}) V_{b\mu\nu}(t)}{d_{ab} - \mathbf{d}_{ab} \cdot \mathbf{v}_a(t_{ab})/c}, \quad (3.60)$$

where t_{ab} is the retarded time satisfying $c(t - t_{ab}) = |\mathbf{x}_b - \mathbf{x}_a(t_{ab})|$, and d_{ab} is the magnitude of the retarded displacement $\mathbf{d}_{ab} = \mathbf{x}_b(t) - \mathbf{x}_a(t_{ab})$.

3.4.1 Slow-motion Approximation

We now have an action that gives a shift in the gravitational field energy similar to the one derived in the previous section, but in purely covariant manner and without quantum operators.

We now want to take a look at the slow-moving approximation of this action. When the source is ‘slow-moving’, that is the N point masses are moving at non-relativistic speeds $|\mathbf{v}_a| \ll c$, then we can approximate $\bar{V}_a^{\mu\nu}(t_{ab}) V_{b\mu\nu}(t)$ as [19],

$$\bar{V}_a^{\mu\nu}(t_{ab}) V_{b\mu\nu}(t) \approx c^4 + \mathcal{O}(c^3 |\mathbf{v}_a|), \quad (3.61)$$

and thus, we can approximate Eq. 3.60 as,

$$S_h \approx \frac{1}{2} G \sum_{a,b}^{a \neq b} \int dt \frac{m_a m_b}{d_{ab}(t)}. \quad (3.62)$$

As we can see here, this action is very similar to the Newtonian Gravitational Potential Energy. In fact, we can regard this action as the causal version of Newton’s Law of Gravitation [19].

In the next chapter, we will go back to the shift in graviton vacuum energy derived in Eq. 3.40, and derive the expression for the static limit, and check if there too we recover the Newtonian potential, and check for non-static corrections, while also investigating the entanglement produced via gravitational interactions.

Chapter 4

Entanglement via the Graviton

Now that we have the quantum gravitational interaction expressions, we want to investigate the entanglement induced by such interactions. We will consider two cases, one in which the coupling is induced by the component \hat{T}_{00} in the static limit, and by the full Stress-Energy tensor $\hat{T}_{\mu\nu}$ in the non-static case, mainly following Bose *et al.* [17].

4.1 Entanglement in the Static Limit

Consider two particles of mass m . The Stress-Energy tensor then, in the static limit is,

$$\hat{T}_{00}(\mathbf{r}) \equiv mc^2(\delta(\mathbf{r} - \hat{\mathbf{r}}_A) + \delta(\mathbf{r} - \hat{\mathbf{r}}_B)), \quad (4.1)$$

where $\hat{\mathbf{r}}_A = (\hat{x}_A, 0, 0)$ and $\hat{\mathbf{r}}_B = (\hat{x}_B, 0, 0)$ represent the positions of the matter system. We can do a Fourier Transform of Eq. 4.1, and we get,

$$\begin{aligned} \hat{T}_{00}(\mathbf{r}) &= \frac{1}{\sqrt{(2\pi)^3}} \int d\mathbf{r} e^{-i\mathbf{k}\cdot\mathbf{r}} \hat{T}_{00}(\mathbf{r}) \\ &= \frac{1}{\sqrt{(2\pi)^3}} \int d\mathbf{r} e^{-i\mathbf{k}\cdot\mathbf{r}} mc^2(\delta(\mathbf{r} - \hat{\mathbf{r}}_A) + \delta(\mathbf{r} - \hat{\mathbf{r}}_B)) \\ &= \frac{mc^2}{\sqrt{(2\pi)^3}} (e^{-i\mathbf{k}\cdot\mathbf{r}_A} + e^{-i\mathbf{k}\cdot\mathbf{r}_B}). \end{aligned} \quad (4.2)$$

Now we can calculate the interaction Hamiltonian for the static limit, given by [17, 40],

$$\begin{aligned} \hat{H}_{\text{int}} &= -\frac{1}{2} \int d\mathbf{r} \hat{h}^{\mu\nu}(\mathbf{r}) \hat{T}_{\mu\nu}(\mathbf{r}) \\ &= -\frac{1}{2} \int d\mathbf{r} \hat{h}^{00}(\mathbf{r}) \hat{T}_{00}(\mathbf{r}) \\ &= -\frac{1}{2} \int d\mathbf{r} \left(\hat{\gamma}_{00}(\mathbf{r}) + \frac{1}{2} \hat{\gamma}(\mathbf{r}) \right) \hat{T}_{00}(\mathbf{r}) \\ &= -\frac{1}{2} \int d\mathbf{r} \mathcal{A} \int d\mathbf{k}' \sqrt{\frac{\hbar}{2\omega_{k'}(2\pi)^3}} \left(\hat{P}_{00}^\dagger(\mathbf{k}') e^{-i\mathbf{k}'\cdot\mathbf{r}} + \hat{P}_{00}(\mathbf{k}') e^{i\mathbf{k}'\cdot\mathbf{r}} \right. \\ &\quad \left. + \hat{P}^\dagger(\mathbf{k}') e^{-i\mathbf{k}'\cdot\mathbf{r}} + \hat{P}(\mathbf{k}') e^{i\mathbf{k}'\cdot\mathbf{r}} \right) \hat{T}_{00}(\mathbf{r}) \end{aligned} \quad (4.3)$$

As shown above in Eq. 3.40, in the shift of the graviton vacuum energy, there are no first order contributions, and the main contributions come from the second order perturbation term. In this case the second order perturbation term is,

$$\Delta \hat{H}_g = \int d\mathbf{k} \frac{\langle 0 | \hat{H}_{\text{int}} | \mathbf{k} \rangle \langle \mathbf{k} | \hat{H}_{\text{int}} | 0 \rangle}{E_0 - E_{\mathbf{k}}}, \quad (4.4)$$

where $|\mathbf{k}\rangle = (\hat{P}_{00}^\dagger(\mathbf{k}) + \hat{P}^\dagger(\mathbf{k}))|0\rangle$ is the one particle state constructed in the unperturbed background, $E_{\mathbf{k}} = E_0 + \hbar\omega_{\mathbf{k}}$ is the energy of the one particle state and E_0 is the energy of the vacuum state. The mediated graviton is off shell, which is why there is an integration over \mathbf{k} , and hence the graviton does not obey classical equations of motion.

We now want to compute $\langle \mathbf{k} | \hat{H}_{\text{int}} | 0 \rangle$. To do this, let us first consider \hat{H}_{int} again.

$$\begin{aligned} \hat{H}_{\text{int}} &= -\frac{1}{2} \int d\mathbf{r} \mathcal{A} \int d\mathbf{k}' \sqrt{\frac{\hbar}{2\omega_{\mathbf{k}'}(2\pi)^3}} \left(\hat{P}_{00}^\dagger(\mathbf{k}') e^{-i\mathbf{k}' \cdot \mathbf{r}} + \hat{P}_{00}(\mathbf{k}') e^{i\mathbf{k}' \cdot \mathbf{r}} \right. \\ &\quad \left. + \hat{P}^\dagger(\mathbf{k}') e^{-i\mathbf{k}' \cdot \mathbf{r}} + \hat{P}(\mathbf{k}') e^{i\mathbf{k}' \cdot \mathbf{r}} \right) \hat{T}_{00}(\mathbf{r}) \\ &= -\frac{1}{2} \mathcal{A} \int d\mathbf{r} \int d\mathbf{k}' \sqrt{\frac{\hbar}{2\omega_{\mathbf{k}'}(2\pi)^3}} \left(\left(\hat{P}_{00}^\dagger(\mathbf{k}') + \hat{P}^\dagger(\mathbf{k}') \right) e^{-i\mathbf{k}' \cdot \mathbf{r}} \right. \\ &\quad \left. + \left(\hat{P}_{00}(\mathbf{k}') + \hat{P}(\mathbf{k}') \right) e^{i\mathbf{k}' \cdot \mathbf{r}} \right) \hat{T}_{00}(\mathbf{r}). \end{aligned} \quad (4.5)$$

Consider $\langle \mathbf{k} | \hat{P}_{00}(\mathbf{k}') + \hat{P}(\mathbf{k}') | 0 \rangle$.

$$\langle \mathbf{k} | \hat{P}_{00}(\mathbf{k}') + \hat{P}(\mathbf{k}') | 0 \rangle = 0, \quad (4.6)$$

as $\hat{P}_{00}|0\rangle = 0$ and $\hat{P}|0\rangle = 0$. Now consider $\langle \mathbf{k} | \hat{P}_{00}^\dagger(\mathbf{k}') + \hat{P}^\dagger(\mathbf{k}') | 0 \rangle$.

$$\begin{aligned} \langle \mathbf{k} | \hat{P}_{00}^\dagger(\mathbf{k}') + \hat{P}^\dagger(\mathbf{k}') | 0 \rangle &= \langle 0 | \hat{P}_{00}(\mathbf{k}) \hat{P}_{00}^\dagger(\mathbf{k}') + \hat{P}(\mathbf{k}) \hat{P}^\dagger(\mathbf{k}') | 0 \rangle \\ &= \langle 0 | \hat{P}_{00}(\mathbf{k}) \hat{P}_{00}^\dagger(\mathbf{k}') | 0 \rangle + \langle 0 | \hat{P}(\mathbf{k}) \hat{P}^\dagger(\mathbf{k}') | 0 \rangle. \end{aligned} \quad (4.7)$$

Since $\hat{P}_{00}|0\rangle = 0$ and $\hat{P}|0\rangle = 0$, we can write $\langle 0 | \hat{P}_{00}(\mathbf{k}) \hat{P}_{00}^\dagger(\mathbf{k}') | 0 \rangle$ and $\langle 0 | \hat{P}(\mathbf{k}) \hat{P}^\dagger(\mathbf{k}') | 0 \rangle$ in terms of the ETCRs derived in Eqs. 3.33 and 3.34 as follows,

$$\langle 0 | \hat{P}_{00}(\mathbf{k}) \hat{P}_{00}^\dagger(\mathbf{k}') | 0 \rangle = \langle 0 | [\hat{P}_{00}(\mathbf{k}), \hat{P}_{00}^\dagger(\mathbf{k}')] | 0 \rangle \quad (4.8)$$

and

$$\langle 0 | \hat{P}(\mathbf{k}) \hat{P}^\dagger(\mathbf{k}') | 0 \rangle = \langle 0 | [\hat{P}(\mathbf{k}), \hat{P}^\dagger(\mathbf{k}')] | 0 \rangle, \quad (4.9)$$

because $\langle 0 | \hat{P}_{00}^\dagger(\mathbf{k}) \hat{P}_{00}(\mathbf{k}') | 0 \rangle = 0$ and $\langle 0 | \hat{P}^\dagger(\mathbf{k}) \hat{P}(\mathbf{k}') | 0 \rangle = 0$. Thus we can write Eq. 4.7 as,

$$\begin{aligned} \langle \mathbf{k} | \hat{P}_{00}^\dagger(\mathbf{k}') + \hat{P}^\dagger(\mathbf{k}') | 0 \rangle &= \langle 0 | [\hat{P}_{00}(\mathbf{k}), \hat{P}_{00}^\dagger(\mathbf{k}')] | 0 \rangle + \langle 0 | [\hat{P}(\mathbf{k}), \hat{P}^\dagger(\mathbf{k}')] | 0 \rangle \\ &= 2\delta(\mathbf{k} - \mathbf{k}') - \delta(\mathbf{k} - \mathbf{k}') \\ &= \delta(\mathbf{k} - \mathbf{k}'). \end{aligned} \quad (4.10)$$

Thus, we can write $\langle \mathbf{k} | \hat{H}_{\text{int}} | 0 \rangle$ as,

$$\begin{aligned}
\langle \mathbf{k} | \hat{H}_{\text{int}} | 0 \rangle &= -\frac{1}{2} \mathcal{A} \int d\mathbf{r} \int d\mathbf{k}' \sqrt{\frac{\hbar}{2\omega_{\mathbf{k}'}(2\pi)^3}} \delta(\mathbf{k} - \mathbf{k}') e^{-i\mathbf{k}' \cdot \mathbf{r}} \hat{T}_{00}(\mathbf{r}) \\
&= -\frac{1}{2} \mathcal{A} \sqrt{\frac{\hbar}{2\omega_k}} \int d\mathbf{r} \sqrt{\frac{1}{(2\pi)^3}} e^{-i\mathbf{k} \cdot \mathbf{r}} \hat{T}_{00}(\mathbf{r}) \\
&= -\frac{\mathcal{A}}{2} \sqrt{\frac{\hbar}{2\omega_k}} \hat{T}_{00}(\mathbf{k}),
\end{aligned} \tag{4.11}$$

where $\hat{T}_{00}(\mathbf{k})$ is the Fourier Transform of $\hat{T}_{00}(\mathbf{r})$ as shown in Eq.4.2. We then get,

$$\langle 0 | \hat{H}_{\text{int}} | \mathbf{k} \rangle \langle \mathbf{k} | \hat{H}_{\text{int}} | 0 \rangle = \frac{\hbar \mathcal{A}^2 \hat{T}_{00}^\dagger(\mathbf{k}) \hat{T}_{00}(\mathbf{k})}{8\omega_k}. \tag{4.12}$$

Plugging in Eq. 4.2 and simplifying, we get,

$$\langle 0 | \hat{H}_{\text{int}} | \mathbf{k} \rangle \langle \mathbf{k} | \hat{H}_{\text{int}} | 0 \rangle = \frac{\hbar \mathcal{A}^2 m^2 c^4}{8\omega_k (2\pi)^3} (2 + e^{i\mathbf{k} \cdot (\mathbf{r}_B - \mathbf{r}_A)} + e^{-i\mathbf{k} \cdot (\mathbf{r}_B - \mathbf{r}_A)}). \tag{4.13}$$

Plugging this into Eq. 4.4, we get,

$$\begin{aligned}
\Delta \hat{H}_g &= \int d\mathbf{k} \frac{\langle 0 | \hat{H}_{\text{int}} | \mathbf{k} \rangle \langle \mathbf{k} | \hat{H}_{\text{int}} | 0 \rangle}{E_0 - E_{\mathbf{k}}} \\
&= \int d\mathbf{k} \frac{\hbar \mathcal{A}^2 m^2 c^4}{8\omega_k (2\pi)^3} \frac{(2 + e^{i\mathbf{k} \cdot (\mathbf{r}_B - \mathbf{r}_A)} + e^{-i\mathbf{k} \cdot (\mathbf{r}_B - \mathbf{r}_A)})}{-\hbar\omega_k} \\
&= \Delta \hat{H}_{gI} + \Delta \hat{H}_{gII} + \Delta \hat{H}_{gIII}
\end{aligned} \tag{4.14}$$

We can ignore the $\Delta \hat{H}_{gI}$ term as that corresponds to self energy of this off-shell graviton with UV divergence, and assume, without loss of generality, that \mathbf{k} points in a direction such that, in spherical coordinates, $\mathbf{k} \cdot (\mathbf{r}_B - \mathbf{r}_A) = k(r_B - r_A) \cos \theta$. And, since the graviton is off-shell, $\omega_k^2 = k^2 c^2$. Therefore, solving these integrals in spherical coordinates, we get,

$$\begin{aligned}
\Delta \hat{H}_{gII} &= - \int d\mathbf{k} \frac{\mathcal{A}^2 m^2 c^4}{8\omega_k^2 (2\pi)^3} (e^{i\mathbf{k} \cdot (\mathbf{r}_B - \mathbf{r}_A)}) \\
&= \frac{-\mathcal{A}^2 m^2 c^4}{8(2\pi)^3} 2\pi \int k^2 dk \frac{1}{k^2 c^2} \int_{-1}^1 d \cos \theta (e^{ik(r_B - r_A) \cos \theta}) \\
&= \frac{-\mathcal{A}^2 m^2 c^2}{8(2\pi)^2} \int_0^\infty dk \frac{2i}{2i} \frac{e^{ik(r_B - r_A)} - e^{-ik(r_B - r_A)}}{ik(r_B - r_A)} \\
&= \frac{-\mathcal{A}^2 m^2 c^2}{8(2\pi)^2} \int_0^\infty dk 2 \frac{\sin k(r_B - r_A)}{k(r_B - r_A)} \\
&= \frac{-\mathcal{A}^2 m^2 c^2}{8(2\pi)^2} 2 \frac{\pi}{2 |\hat{x}_a - \hat{x}_b|} = \frac{-\mathcal{A}^2 m^2 c^2}{32\pi |\hat{x}_a - \hat{x}_b|}
\end{aligned} \tag{4.15}$$

Similarly, one can show that $\Delta\hat{H}_{gIII} = \frac{-\mathcal{A}^2 m^2 c^2}{32\pi|\hat{x}_a - \hat{x}_b|}$. Thus, we get that the shift in vacuum energy of the graviton in the static case is given by,

$$\begin{aligned}\Delta\hat{H}_g &= \frac{-\mathcal{A}^2 m^2 c^2}{32\pi|\hat{x}_a - \hat{x}_b|} - \frac{\mathcal{A}^2 m^2 c^2}{32\pi|\hat{x}_a - \hat{x}_b|} \\ &= \frac{-\mathcal{A}^2 m^2 c^2}{16\pi|\hat{x}_a - \hat{x}_b|}\end{aligned}\tag{4.16}$$

Now using the value of $\mathcal{A} = \sqrt{16\pi G/c^2}$, we get,

$$\Delta\hat{H}_g = -\frac{Gm^2}{|\hat{x}_a - \hat{x}_b|}.\tag{4.17}$$

This is exactly the Newtonian potential, obtained from quantum operators. This is also similar to the slow-motion approximation obtained in Section 3.4.1. There is a more covariant way to obtain this result through quantum operators, discussed in Marshman *et al.* [36], where they compute the potential by using the full graviton propagator and the scattering amplitude between two masses via the exchange of the spin-2 and spin-0 components of the graviton (see [36, Appendix]). Here, we have calculated the shift in graviton vacuum, as per Bose *et al.* [17]. However, in the non-relativistic limit, both methods yield the same potential.

It is also important to note that $\Delta\hat{H}_g$ is an operator valued function of the two matter systems, owing to the interaction between the graviton and the particles.

If the two matter systems are not in sharply defined positions, say they are in a spatial superposition, then the change in graviton energy will not be a real number, but an operator-valued quantity, which is a bonafide quantum entity [17]. Let us now calculate the excited wave function $|\psi_f\rangle$ for the QHOs described in Section 2.2. Using Eq. 2.1 to expand Eq. 4.17, we get,

$$\Delta\hat{H}_g \approx -\frac{Gm^2}{d} + \frac{Gm^2}{d^2}(\delta\hat{x}_B - \delta\hat{x}_A) - \frac{Gm^2}{d^3}(\delta\hat{x}_B - \delta\hat{x}_A)^2.\tag{4.18}$$

The last term contains the lowest order matter-matter interaction, as,

$$-\frac{Gm^2}{d^3}(\delta\hat{x}_B - \delta\hat{x}_A)^2 = -\frac{Gm^2}{d^3}(\delta\hat{x}_B^2 + \delta\hat{x}_A^2 - 2\delta\hat{x}_A\delta\hat{x}_B).\tag{4.19}$$

Thus, we get,

$$\hat{H}_{AB} = \frac{2Gm^2}{d^3} \delta\hat{x}_A\delta\hat{x}_B\tag{4.20}$$

It is critical to realise that this term would not have arisen if we had a real-valued shift to the graviton energy. We must then conclude that entanglement from gravitational interactions is a quantum signature of the gravitational field. We can now use the modes in 2.3 to expand \hat{H}_{AB} and get,

$$\begin{aligned}\hat{H}_{AB} &= \frac{2Gm^2}{d^3} \delta\hat{x}_A\delta\hat{x}_B \\ &= \frac{2Gm^2}{d^3} \frac{\hbar}{2m\omega_m} (\hat{a} + \hat{a}^\dagger)(\hat{b} + \hat{b}^\dagger) \\ &= \hbar\mathbf{g}(\hat{a}\hat{b} + \hat{a}\hat{b}^\dagger + \hat{a}^\dagger\hat{b} + \hat{a}^\dagger\hat{b}^\dagger),\end{aligned}\tag{4.21}$$

where we define the coupling, \mathfrak{g} , as,

$$\mathfrak{g} = \frac{Gm}{d^3\omega_m}. \quad (4.22)$$

Using \hat{H}_{AB} and the definition of the coefficients for the final state from Eq. 2.8, we get, using $\lambda = 1$,

$$\begin{aligned} C_{nN} &= \frac{\langle n | \langle N | \hat{H}_{AB} | 0 \rangle | 0 \rangle}{2E_0 - E_n - E_N} \\ &= \frac{\langle n | \langle N | \hbar\mathfrak{g}(\hat{a}\hat{b} + \hat{a}\hat{b}^\dagger + \hat{a}^\dagger\hat{b} + \hat{a}^\dagger\hat{b}^\dagger) | 0 \rangle | 0 \rangle}{2E_0 - E_n - E_N} \end{aligned} \quad (4.23)$$

But $\hat{a}|0\rangle = 0$ and $\hat{b}|0\rangle = 0$. Therefore,

$$C_{nN} = \frac{\langle n | \langle N | \hbar\mathfrak{g}(\hat{a}^\dagger\hat{b}^\dagger) | 0 \rangle | 0 \rangle}{2E_0 - E_n - E_N} \quad (4.24)$$

Thus the only coefficient that is non-zero is C_{11} given by,

$$C_{11} = \frac{\hbar\mathfrak{g}}{-2\hbar\omega_m} = -\frac{\mathfrak{g}}{2\omega_m}. \quad (4.25)$$

Note that $\hat{a}^\dagger\hat{b}^\dagger$ term only generates the first excited state. We also have $C_{00} = 1$ for the unperturbed state. Therefore, the final state is given by,

$$|\psi_f\rangle \equiv \frac{1}{\sqrt{1 + (\mathfrak{g}/(2\omega_m))^2}} [|0\rangle|0\rangle - \frac{\mathfrak{g}}{2\omega_m} |1\rangle|1\rangle]. \quad (4.26)$$

This is the *entangled* state involving the ground and first excited states. I will calculate the Concurrence and Entanglement Entropy in Sections 4.3 and 4.4 respectively. All of the results in this section are also derived in Bose *et al.* [17], but they have different signs for some expressions (e.g. The \hat{H}_{int} that I have computed here differs from them by a negative sign).

4.2 Entanglement in the Non-Static Limit

Now we are interested in the entanglement of two bodies with non-zero T_{ij} . Specifically, we will consider two particles in QHOs moving along the x -axis such that the only non-zero components of $T_{\mu\nu}$ are given by T_{00} , $T_{01} = T_{10}$, and T_{11} .

Then, the corresponding components of $\hat{h}_{\mu\nu}$ are $\hat{h}_{00} = \hat{\gamma}_{00} + 1/2\hat{\gamma}$, $\hat{h}_{01} = \hat{h}_{10} = \hat{\gamma}_{01}$, and by $\hat{h}_{11} = \hat{\gamma}_{11} - (1/2)\hat{\gamma}$ [17]. Recall that, like in the static case (Eq. 4.3), \hat{H}_{int} in this case is also given by [17],

$$\begin{aligned} \hat{H}_{\text{int}} &= -\frac{1}{2} \int d\mathbf{r} [\hat{\gamma}_{00}(\mathbf{r}) + (1/2)\hat{\gamma}(\mathbf{r})] \hat{T}_{00}(\mathbf{r}) \\ &\quad - \int d\mathbf{r} \hat{\gamma}_{01}(\mathbf{r}) \hat{T}_{01}(\mathbf{r}) \\ &\quad - \frac{1}{2} \int d\mathbf{r} [\hat{\gamma}_{11}(\mathbf{r}) - (1/2)\hat{\gamma}(\mathbf{r})] \hat{T}_{11}(\mathbf{r}), \end{aligned} \quad (4.27)$$

Here, the first line is the same as the static case, and the second and third lines come from the degrees of freedom of the Gravitational Waves corresponding to the + polarisation [17, 40]. Since we are considering two QHOs that are moving along the x -axis, the stress-energy tensor is given by,

$$\hat{T}_{\mu\nu}(\mathbf{r}) \equiv \frac{\hat{p}_\mu \hat{p}_\nu}{E/c^2} (\delta(\mathbf{r} - \hat{\mathbf{r}}_A) + \delta(\mathbf{r} - \hat{\mathbf{r}}_B)), \quad (4.28)$$

where $p_\mu = (-E/c, \mathbf{p})$, $E = \sqrt{\mathbf{p}^2 c^2 + m^2 c^4}$, $\mu, \nu = 0, 1$, and $\hat{\mathbf{r}}_A = (\hat{x}_A, 0, 0)$, $\hat{\mathbf{r}}_B = (\hat{x}_B, 0, 0)$ denote the positions of the two matter systems [17]. The classical expression of the stress-energy tensor has been promoted to a quantum operator following the Weyl quantization prescription to ensure that this is a Hermitian operator [17]. However, to simplify notation, following the method followed by Bose *et al.* [17], I will write the unsymmetrized expressions, e.g. $\hat{x}\hat{p}$, implicitly assuming that all expressions need to be interpreted in their symmetrized ordering, e.g. $(\hat{x}\hat{p} + \hat{p}\hat{x})/2$.

Now, the Fourier transform of the stress-energy tensor is given by,

$$\hat{T}_{\mu\nu}(\mathbf{k}) = \frac{1}{\sqrt{(2\pi)^3}} \int d\mathbf{r} e^{-i\mathbf{k}\cdot\mathbf{r}} \hat{T}_{\mu\nu}(\mathbf{r}). \quad (4.29)$$

Using the definitions of $\hat{\gamma}_{\mu\nu}$ and $\hat{\gamma}$ from Eqs. 3.31 and 3.32, and using the Fourier Transform of $\hat{T}_{\mu\nu}$, we can rewrite the interaction Hamiltonian to be,

$$\begin{aligned} \hat{H}_{\text{int}} = & -\frac{\mathcal{A}}{2} \int d\mathbf{k} \sqrt{\frac{\hbar}{2\omega_{\mathbf{k}}}} \left[\hat{P}_{00}(\mathbf{k}) + \hat{P}(\mathbf{k}) \right] \hat{T}_{00}(\mathbf{k}) + \text{H.c.}) \\ & - \mathcal{A} \int d\mathbf{k} \sqrt{\frac{\hbar}{2\omega_{\mathbf{k}}}} (\hat{P}_{01}(\mathbf{k}) \hat{T}_{01}(\mathbf{k}) + \text{H.c.}) \\ & - \frac{\mathcal{A}}{2} \int d\mathbf{k} \sqrt{\frac{\hbar}{2\omega_{\mathbf{k}}}} \left[\hat{P}_{11}(\mathbf{k}) - \hat{P}(\mathbf{k}) \right] \hat{T}_{11}(\mathbf{k}) + \text{H.c.}, \end{aligned} \quad (4.30)$$

where ‘‘H.c.’’ represents the Hermitian conjugate in the usual sense.

We can write the Fourier transforms of the non-zero $\hat{T}_{\mu\nu}$ components as follows,

$$\hat{T}_{00}(\mathbf{k}) = \frac{1}{\sqrt{(2\pi)^3}} (\hat{E}_A e^{i\mathbf{k}\cdot\hat{\mathbf{r}}_A} + \hat{E}_B e^{i\mathbf{k}\cdot\hat{\mathbf{r}}_B}), \quad (4.31)$$

$$\hat{T}_{01}(\mathbf{k}) = -\frac{1}{\sqrt{(2\pi)^3}} (\hat{p}_A c e^{i\mathbf{k}\cdot\hat{\mathbf{r}}_A} + \hat{p}_B c e^{i\mathbf{k}\cdot\hat{\mathbf{r}}_B}), \quad (4.32)$$

$$\hat{T}_{11}(\mathbf{k}) = \frac{1}{\sqrt{(2\pi)^3}} \left(\frac{\hat{p}_A^2 c^2}{E_A} e^{i\mathbf{k}\cdot\hat{\mathbf{r}}_A} + \frac{\hat{p}_B^2 c^2}{E_B} e^{i\mathbf{k}\cdot\hat{\mathbf{r}}_B} \right). \quad (4.33)$$

As before in the static case, the energy shift in the graviton vacuum $|0\rangle$ is given by the second order perturbation theory (once again, the first order perturbation term vanishes). Thus,

$$\Delta \hat{H}_g \equiv \sum \int d\mathbf{k} \frac{\langle 0 | \hat{H}_{\text{int}} | \mathbf{k} \rangle \langle \mathbf{k} | \hat{H}_{\text{int}} | 0 \rangle}{E_0 - E_{\mathbf{k}}}. \quad (4.34)$$

Here the sum represents the summation over the one particle projectors $|\mathbf{k}\rangle\langle\mathbf{k}|$ constructed on the unperturbed vacuum. $|\mathbf{k}\rangle = \frac{1}{\sqrt{2}}\hat{P}_{00}^\dagger(\mathbf{k})|0\rangle$, $\frac{1}{\sqrt{2}}\hat{P}_{11}^\dagger(\mathbf{k})|0\rangle$, $\hat{P}_{01}^\dagger(\mathbf{k})|0\rangle$ and $\hat{P}^\dagger(\mathbf{k})|0\rangle$, where the prefactor of $\frac{1}{\sqrt{2}}$ is to ensure the correct normalisation [17]. E_0 is the energy of the vacuum state, $E_{\mathbf{k}} = E_0 + \hbar\omega_{\mathbf{k}}$ is the energy of the one particle state.

Following exactly the same procedure to the static case, we can easily evaluate the following expressions,

$$\langle 0|\hat{P}(\mathbf{k})\hat{H}_{\text{int}}|0\rangle = -\frac{\mathcal{A}}{2}\sqrt{\frac{\hbar}{2\omega_{\mathbf{k}}}}(\hat{T}_{00}(\mathbf{k}) - \hat{T}_{11}(\mathbf{k})), \quad (4.35)$$

$$\langle 0|\hat{P}_{01}(\mathbf{k})\hat{H}_{\text{int}}|0\rangle = -\mathcal{A}\sqrt{\frac{\hbar}{2\omega_{\mathbf{k}}}}\hat{T}_{01}(\mathbf{k}), \quad (4.36)$$

$$\langle 0|\hat{P}_{00}(\mathbf{k})\hat{H}_{\text{int}}|0\rangle = -\mathcal{A}\sqrt{\frac{\hbar}{2\omega_{\mathbf{k}}}}\hat{T}_{00}(\mathbf{k}), \quad (4.37)$$

$$\langle 0|\hat{P}_{11}(\mathbf{k})\hat{H}_{\text{int}}|0\rangle = -\mathcal{A}\sqrt{\frac{\hbar}{2\omega_{\mathbf{k}}}}\hat{T}_{11}(\mathbf{k}). \quad (4.38)$$

Using these equations, we can now calculate the shift in the graviton vacuum by plugging them into Eq. 4.34. Thus, we get [17],

$$\begin{aligned} \Delta\hat{H}_g = & -\mathcal{A}^2 \int d\mathbf{k} \frac{\hat{T}_{00}^\dagger(\mathbf{k})\hat{T}_{00}(\mathbf{k}) + \hat{T}_{11}^\dagger(\mathbf{k})\hat{T}_{11}(\mathbf{k})}{8c^2\mathbf{k}^2} \\ & - \mathcal{A}^2 \int d\mathbf{k} \frac{(\hat{T}_{00}^\dagger(\mathbf{k})\hat{T}_{11}(\mathbf{k}) + \text{H.c.})}{8c^2\mathbf{k}^2} \\ & + 4\mathcal{A}^2 \int d\mathbf{k} \frac{\hat{T}_{01}^\dagger(\mathbf{k})\hat{T}_{01}(\mathbf{k})}{8c^2\mathbf{k}^2}. \end{aligned} \quad (4.39)$$

Now, using the fact that the two particles are confined along the x -axis, we can set $\hat{p}_{Ay} = \hat{p}_{Az} = \hat{p}_{By} = \hat{p}_{Bz} = 0$. We can also define $\hat{p}_A \equiv \hat{p}_{Ax}$ and $\hat{p}_B \equiv \hat{p}_{Bx}$. As usual, $\hat{\mathbf{r}}_A = (x_A, 0, 0)$, $\hat{\mathbf{r}}_B = (x_B, 0, 0)$ and, $\mathbf{k} = (k_x, k_y, k_z)$. Once again, removing any self energy terms (e.g. terms that are solely $\hat{E}_A\hat{E}_A$ or $\hat{E}_B\hat{E}_B$), we can then write the shift in graviton vacuum as,

$$\begin{aligned} \Delta\hat{H}_g = & -\frac{\mathcal{A}^2}{(2\pi)^3} \int d\mathbf{k} \left(\frac{\hat{E}_A\hat{E}_B + \frac{\hat{p}_A^2 c^2}{\hat{E}_A} \frac{\hat{p}_B^2 c^2}{\hat{E}_B}}{8c^2\mathbf{k}^2} + \frac{\hat{E}_A \frac{\hat{p}_B^2 c^2}{\hat{E}_B} + \hat{E}_B \frac{\hat{p}_A^2 c^2}{\hat{E}_A}}{8c^2\mathbf{k}^2} - 4 \frac{\hat{p}_A c \hat{p}_B c}{8c^2\mathbf{k}^2} \right) \\ & \times (e^{ik_x(\hat{x}_A - \hat{x}_B)} + e^{-ik_x(\hat{x}_A - \hat{x}_B)}). \end{aligned} \quad (4.40)$$

Once again, without loss of generality, we can assume that within spherical coordinates, \mathbf{k} points in a direction w.r.t $(\mathbf{r}_A - \mathbf{r}_B)$ such that $\mathbf{k} \cdot (\mathbf{r}_B - \mathbf{r}_A) = k(r_A - r_B) \cos \theta$. Then following exactly the same steps as before for the static case, performing the integration in spherical coordinates and plugging in the value of \mathcal{A} (although this time it is slightly more tedious to do), and then following the extra step of substituting \hat{E}_A and \hat{E}_B and expanding in powers of $1/c^2$ upto $\mathcal{O}(1/c^4)$, we get the following expression for $\Delta\hat{H}_g$ [17],

$$\Delta\hat{H}_g = -\frac{Gm^2}{|\hat{x}_A - \hat{x}_B|} - \frac{G(3\hat{p}_A^2 - 8\hat{p}_A\hat{p}_B + 3\hat{p}_B^2)}{2c^2|\hat{x}_A - \hat{x}_B|} - \frac{G(5\hat{p}_A^4 - 18\hat{p}_A^2\hat{p}_B^2 + 5\hat{p}_B^4)}{8c^4m^2|\hat{x}_A - \hat{x}_B|}. \quad (4.41)$$

The first term is exactly the same as the static case, and the second and third terms are what I am terming as non-static “corrections” to the static case. I will consider these two terms separately as they are simpler to deal with that way. Note that if we returned this to the static case, i.e. set \hat{p}_A and \hat{p}_B to zero, then the last two terms vanish [17]. Again, if the shift was classical, it would have been real and not operator valued.

4.2.1 Correction 1

Now consider the first correction. Let us first expand the expression using Eq. 2.3.

$$\begin{aligned} -\frac{G(3\hat{p}_A^2 - 8\hat{p}_A\hat{p}_B + 3\hat{p}_B^2)}{2c^2|\hat{x}_A - \hat{x}_B|} &= -\frac{3G\hat{p}_A^2}{2c^2|\hat{x}_A - \hat{x}_B|} + \frac{4G\hat{p}_A\hat{p}_B}{c^2|\hat{x}_A - \hat{x}_B|} - \frac{3G\hat{p}_B^2}{2c^2|\hat{x}_A - \hat{x}_B|} \\ &\approx -\frac{3G\hat{p}_A^2}{2c^2d} + \frac{4G\hat{p}_A\hat{p}_B}{c^2d} - \frac{G(3\hat{p}_B^2)}{2c^2d} + \dots \end{aligned} \quad (4.42)$$

Therefore, the lowest order interaction term is,

$$\hat{H}_{AB} \approx \frac{4G\hat{p}_A\hat{p}_B}{c^2d} + \dots \quad (4.43)$$

At the lowest order in the expansion, $\delta\hat{x}_A$ and $\delta\hat{x}_B$ do not appear, and the interaction is dominated by the momentum operators. Using Eq. 2.4 to write this in terms of the mode expansions, we get,

$$\begin{aligned} \frac{4G\hat{p}_A\hat{p}_B}{c^2d} &= -\frac{4G}{c^2d} \frac{\hbar m \omega_m}{2} (\hat{a}^\dagger - \hat{a})(\hat{b}^\dagger - \hat{b}) \\ &= -\hbar \mathfrak{g} (\hat{a}^\dagger \hat{b}^\dagger - \hat{a}^\dagger \hat{b} - \hat{a} \hat{b}^\dagger + \hat{a} \hat{b}), \end{aligned} \quad (4.44)$$

where the coupling has been defined as,

$$\mathfrak{g} = \frac{2Gm\omega_m}{c^2d} \quad (4.45)$$

Once again, since $\hat{a}|0\rangle = \hat{b}|0\rangle = 0$, only relevant term comes from $\hat{a}^\dagger \hat{b}^\dagger$. Thus, the only coefficient that is non-zero is C_{11} given by,

$$C_{11} = \frac{-\hbar \mathfrak{g}}{-2\hbar \omega_m} = \frac{\mathfrak{g}}{2\omega_m} \quad (4.46)$$

We also have $C_{00} = 1$ for the unperturbed state. Therefore, the final state is given by,

$$|\psi_f\rangle \equiv \frac{1}{\sqrt{1 + (\mathfrak{g}/(2\omega_m))^2}} [|0\rangle|0\rangle + \frac{\mathfrak{g}}{2\omega_m} |1\rangle|1\rangle]. \quad (4.47)$$

This is the *entangled* state involving the ground and first excited states for the first correction term. Again, these results are derived in Bose *et al.* [17], but there are a few sign differences. The most major one is that our coefficient C_{11} have different signs, resulting in slightly different final states. Again, I will calculate the Concurrence and Entanglement Entropy in Sections 4.3 and 4.4 respectively.

4.2.2 Correction 2

Let us now move on to the next correction term. Again, let us first expand the expression using Eq. 2.3.

$$\begin{aligned} -\frac{G(5\hat{p}_A^4 - 18\hat{p}_A^2\hat{p}_B^2 + 5\hat{p}_B^4)}{8c^4m^2|\hat{x}_A - \hat{x}_B|} &= -\frac{5G\hat{p}_A^4}{8c^4m^2|\hat{x}_A - \hat{x}_B|} + \frac{9G\hat{p}_A^2\hat{p}_B^2}{4c^4m^2|\hat{x}_A - \hat{x}_B|} - \frac{5G\hat{p}_B^4}{8c^4m^2|\hat{x}_A - \hat{x}_B|} \\ &\approx -\frac{5G\hat{p}_A^4}{8c^4m^2d} + \frac{9G\hat{p}_A^2\hat{p}_B^2}{4c^4m^2d} - \frac{5G\hat{p}_B^4}{8c^4m^2d} + \dots \end{aligned} \quad (4.48)$$

Thus, the interaction Hamiltonian is,

$$\hat{H}_{AB} \approx \frac{9G\hat{p}_A^2\hat{p}_B^2}{4c^4m^2d} + \dots \quad (4.49)$$

Once again, at the lowest order in the expansion, $\delta\hat{x}_A$ and $\delta\hat{x}_B$ do not appear, and the interaction is now dominated by the square of the momentum operators. Using Eq. 2.4 to write this in terms of the mode expansions, we get,

$$\begin{aligned} \frac{9G\hat{p}_A^2\hat{p}_B^2}{4c^4m^2d} &= \frac{9G}{4c^4m^2d} \frac{\hbar^2 m^2 \omega_m^2}{2^2} (\hat{a}^\dagger - \hat{a})^2 (\hat{b}^\dagger - \hat{b})^2 \\ &= -\hbar\mathfrak{g}((\hat{a}^\dagger\hat{b}^\dagger)^2 + \dots) \end{aligned} \quad (4.50)$$

where the coupling has been defined as,

$$\mathfrak{g} = \frac{9G\hbar\omega_m^2}{16c^4d} \quad (4.51)$$

Here, the only term of relevance is $(\hat{a}^\dagger\hat{b}^\dagger)^2$, as the other terms will vanish. This signifies that the final state is a linear combination of the ground state and $|2\rangle|2\rangle$. The gravitons also carry twice the energy of the harmonic oscillator, and $\omega_{\mathbf{k}} = 2\omega_m$. Thus, using Eq. 2.8, we get,

$$C_{22} = \frac{\hbar\mathfrak{g}}{-2\hbar\omega_m} = -\frac{\mathfrak{g}}{2\omega_m} \quad (4.52)$$

We also have $C_{00} = 1$ for the unperturbed state. Therefore, the final state is given by,

$$|\psi_f\rangle \equiv \frac{1}{\sqrt{1 + (\mathfrak{g}/(2\omega_m))^2}} [|0\rangle|0\rangle - \frac{\mathfrak{g}}{2\omega_m} |2\rangle|2\rangle]. \quad (4.53)$$

This is the *entangled* state involving the ground and first excited states for the second correction term. Again, these results are derived in Bose *et al.* [17], but there are a few sign differences. The most major one is that our coefficient C_{22} have different signs, resulting in slightly different final states. Again, I will calculate the Concurrence and Entanglement Entropy in the next two sections.

4.3 Concurrence

We can now calculate the concurrence for the final states of each case.

4.3.1 Static Limit

Recall that for the static limit, the final state is given by Eq. 4.26. Using Eq. 2.13, we can calculate the concurrence easily. In this case, $\mathcal{N} = \sqrt{1 + (\mathfrak{g}/(2\omega_m))^2}$, and the concurrence therefore is given by,

$$C = \sqrt{2 \left(1 - \frac{1 + (\mathfrak{g}/(2\omega_m))^4}{(1 + (\mathfrak{g}/(2\omega_m))^2)^2} \right)} \approx \frac{\mathfrak{g}}{\omega_m} \quad (4.54)$$

This approximation is valid when $\mathfrak{g}/2\omega_m \ll 1$. Substituting the value of the coupling, to first order, we get that the concurrence is,

$$C = \frac{Gm}{d^3\omega_m^2}. \quad (4.55)$$

This is also derived in Bose *et al.* [17], but they have an extra factor of $\sqrt{2}$. I will analyze this result in the next chapter.

4.3.2 Non-Static Limit

Here, we will find the concurrence for each of the correction terms.

Correction 1

Again, the final state is given by Eq. 4.47. Using Eq. 2.13, we can calculate the concurrence easily. In this case, $\mathcal{N} = \sqrt{1 + (\mathfrak{g}/(2\omega_m))^2}$, and the concurrence therefore is given by,

$$C = \sqrt{2 \left(1 - \frac{1 + (\mathfrak{g}/(2\omega_m))^4}{(1 + (\mathfrak{g}/(2\omega_m))^2)^2} \right)} \approx \frac{\mathfrak{g}}{\omega_m} \quad (4.56)$$

This approximation is valid when $\mathfrak{g}/2\omega_m \ll 1$. Substituting the value of the coupling, to first order, we get that the concurrence is,

$$C = \frac{2Gm}{c^2d}. \quad (4.57)$$

This is also derived in Bose *et al.* [17], but they have an extra factor of $\sqrt{2}$. I will analyze this result in the next chapter.

Correction 2

Again, the final state is given by Eq. 4.53. Using Eq. 2.13, we can calculate the concurrence easily. In this case, $\mathcal{N} = \sqrt{1 + (\mathfrak{g}/(2\omega_m))^2}$, and the concurrence therefore is given by,

$$C = \sqrt{2 \left(1 - \frac{1 + (\mathfrak{g}/(2\omega_m))^4}{(1 + (\mathfrak{g}/(2\omega_m))^2)^2} \right)} \approx \frac{\mathfrak{g}}{\omega_m} \quad (4.58)$$

This approximation is valid when $\mathfrak{g}/2\omega_m \ll 1$. Substituting the value of the coupling, to first order, we get that the concurrence is,

$$C = \frac{9G\hbar\omega_m}{16c^4d}. \quad (4.59)$$

This is also derived in Bose *et al.* [17], but they have an extra factor of $\sqrt{2}$. I will analyze this result in the next chapter.

4.4 Entanglement Entropy

I will now calculate the entanglement entropy for each case. But before that, let us take a look at the expression for the density matrix using 2.11,

$$\hat{\rho}_A = \frac{1}{\mathcal{N}} \sum_{n,n',N} C_{nN} C_{n'N}^* |n\rangle\langle n'|, \quad (4.60)$$

And then take a look at each of the three final states in Eqs. 4.26, 4.47, and 4.53. Coincidentally, they all are very similar, and were we to define $\xi = \frac{\mathfrak{g}}{2\omega_m}$, indeed we would find a common expression for the density matrix as a function of ξ and then we can use that to derive a common expression for the entanglement entropy, which would then only require the input of the coupling to give the entanglement entropy for each case. Let us calculate the density matrix, which is given by Eq. 2.11,

$$\rho_A(\xi(\mathfrak{g})) = \frac{1}{1+\xi^2} \begin{bmatrix} 1 & 0 \\ 0 & \xi^2 \end{bmatrix}, \quad (4.61)$$

Therefore, $\log \rho_A$ is given by,

$$\log \rho_A = \begin{bmatrix} \log\left(\frac{1}{1+\xi^2}\right) & 0 \\ 0 & \log\left(\frac{\xi^2}{1+\xi^2}\right) \end{bmatrix}. \quad (4.62)$$

Thus, the entanglement entropy is given by,

$$\begin{aligned} \mathcal{S}(\rho_A(\mathfrak{g})) &= -\text{Tr}[\rho_A \log \rho_A] \\ &= -\left(\frac{1}{1+\xi^2} \log\left(\frac{1}{1+\xi^2}\right) + \frac{\xi^2}{1+\xi^2} \log\left(\frac{\xi^2}{1+\xi^2}\right) \right) \\ &= -\frac{1}{1+\xi^2} \left(\log\left(\frac{1}{1+\xi^2}\right) + \xi^2 \log\left(\frac{\xi^2}{1+\xi^2}\right) \right) \\ &= \frac{1}{1+\xi^2} \left(\log(1+\xi^2)(1+\xi^2) - 2\xi^2 \log(\xi) \right) \\ &= \log(1+\xi^2) - \frac{2\xi^2}{1+\xi^2} \log(\xi) \\ &= \log\left(1 + \left(\frac{\mathfrak{g}}{2\omega_m}\right)^2\right) - \frac{2\left(\frac{\mathfrak{g}}{2\omega_m}\right)^2}{1 + \left(\frac{\mathfrak{g}}{2\omega_m}\right)^2} \log\left(\frac{\mathfrak{g}}{2\omega_m}\right). \end{aligned} \quad (4.63)$$

Now this quantity is always positive, because for all cases $\frac{g}{2\omega_m} \ll 1 \implies \log\left(\frac{g}{2\omega_m}\right) < 0$, so the second term in the above equation is inherently positive, and so is the first term. Since $\frac{g}{2\omega_m} \ll 1$, we can approximate this equation by expanding around zero.

However, $\log\left(\frac{g}{2\omega_m}\right)$ does not have a series expansion near zero and must be kept the same throughout the expansion. The expansion of $\log(1 + \xi^2)$ around zero is,

$$\log(1 + \xi^2) = \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n} \xi^{2n} = \xi^2 - \frac{\xi^4}{2} + \mathcal{O}(\xi^6), \quad (4.64)$$

which is convergent if $|\xi| < 1$, which for us is true as $\xi \ll 1$. The Taylor expansion of the second term is,

$$\frac{2X^2}{1 + \xi^2} \log(\xi) = \left(2X^2 - 2X^4 + \mathcal{O}(\xi^6)\right) \log \xi, \quad (4.65)$$

which is convergent if $0 < |\xi| < 1$, which for us is true as $0 < \xi \ll 1$. Thus, the full expansion for Eq. 4.63 is,

$$\mathcal{S}_A \approx \xi^2 \left((1 - 2 \log(\xi)) \right) + \xi^4 \left(2 \log(\xi) - \frac{1}{2} \right) + \mathcal{O}(\xi^6), \quad (4.66)$$

which is valid near zero. I will talk about this and the following case results in more detail in the next chapter, but for comparison between the perturbative entanglement entropy and the series expansion, please see Fig. 5.3.

4.4.1 Static Limit

Now all that remains is to plug in the individual $\xi = g/2\omega_m$ for each case. For this case, $\xi = \frac{Gm}{2d^3\omega_m^2}$. Thus, the entanglement entropy is,

$$\mathcal{S}_A \approx \left(\frac{Gm}{2d^3\omega_m^2} \right)^2 \left((1 - 2 \log \left(\frac{Gm}{2d^3\omega_m^2} \right)) \right) + \left(\frac{Gm}{2d^3\omega_m^2} \right)^4 \left(2 \log \left(\frac{Gm}{2d^3\omega_m^2} \right) - \frac{1}{2} \right) \quad (4.67)$$

4.4.2 Non-Static Limit

Correction 1

For this case, $\xi = \frac{Gm}{c^2 d}$. Thus, the entanglement entropy is,

$$\mathcal{S}_A \approx \left(\frac{Gm}{c^2 d} \right)^2 \left((1 - 2 \log \left(\frac{Gm}{c^2 d} \right)) \right) + \left(\frac{Gm}{c^2 d} \right)^4 \left(2 \log \left(\frac{Gm}{c^2 d} \right) - \frac{1}{2} \right) \quad (4.68)$$

Correction 2

For this case, $\xi = \frac{9G\hbar\omega_m}{32c^4 d}$. Thus, the entanglement entropy is,

$$\mathcal{S}_A \approx \left(\frac{9G\hbar\omega_m}{32c^4 d} \right)^2 \left((1 - 2 \log \left(\frac{9G\hbar\omega_m}{32c^4 d} \right)) \right) + \left(\frac{9G\hbar\omega_m}{32c^4 d} \right)^4 \left(2 \log \left(\frac{9G\hbar\omega_m}{32c^4 d} \right) - \frac{1}{2} \right). \quad (4.69)$$

Chapter 5

Analysis and Discussion

Let us start with summarising what we have done so far. We looked at underpinning concepts of the BMV experiment [15, 16]. We then saw how quantum interactions induce entanglement between two bodies, and how it is a characteristic trait of quantum interactions. We then wanted to use gravitational interactions as a stand-in for the quantum interactions to see how they would induce entanglement. But for that, we had to first go through the process of linearising Einstein’s field equations, and then do a canonical quantization of the equations. Only then could we look at quantum gravitational interactions, and arrive at expression for the shift in the vacuum energy of the graviton.

Before we jumped into entanglement via the graviton, we first looked at how we could arrive at a similar result purely from a General Relativity point of view, and so we found the action for N interacting point masses, which was similar to the shift in vacuum energy of the graviton we already found, and which, at the the slow-motion approximation, also gave us the Newtonian potential back. We then went back to the derived quantization and quantum gravitational interaction, and used it to calculate the shift in the graviton vacuum energy for when the particles were static and when they were non-static. When they were static, we found that the shift for the vacuum energy was equal to the Newtonian potential, a result we were already expecting, thanks to the action for N interacting point masses we derived earlier.

Then we saw that for the non-static case, not only did we get the Newtonian potential, but two non-static corrections to it, owing to the movement of the particles along the x -axis. These results also match those that have been previously obtained through different methods [17, 47–50]. We then saw that each of these results were operator valued which could not be produced by an operator valued shift, and that if we considered the underlying gravitational field to be classical, then the final states would never have evolved into an entangled state. We then went ahead and calculated the coupling constants, the concurrence and entanglement entropy.

Now that we have the expressions for the coupling, concurrence and entanglement entropy for each of the three cases, we can take a closer look at their behaviour. Let us start with the coupling. The coupling for each of the cases grows as Entanglement grows, and vice versa. All of them also grow inversely with the distance between the two oscillators, but to varying levels. This can be seen in Fig. 5.1. The graph shows a plot of the coupling as

a function of the distance between the two bodies, for all three cases. Constant values of mass and ω_m were chosen as 10^{-14} kg and 10^8 Hz respectively. We can see that, with varying degrees, the coupling has an inverse relationship with the distance. Another feature presented here is that for this range of distances, the static coupling dominates over the other non-static corrections, as can be seen from the scale of each subplot.




Figure 5.1: Comparison of Coupling for Different Cases: (a) Static Case; (b) Non-static Correction 1 (c) Non Static Correction 2

Next, let us look at concurrence. First let us write down the common expression for concurrence found in each of the three cases. This expression was,

$$C = \sqrt{2 \left(1 - \frac{1 + (\mathfrak{g}/(2\omega_m))^4}{(1 + (\mathfrak{g}/(2\omega_m))^2)^2} \right)} \approx \frac{\mathfrak{g}}{\omega_m} \quad (5.1)$$

If one was to calculate the maximum of this function, they would find that the maximum value of this expression is 1 for $\mathfrak{g}/(2\omega_m) = 1$. But it was previously stated in Sec. 2.2 under Eq. 2.13 that the maximum value of concurrence for a maximally entangled state of a system with an infinite-dimensional Hilbert space is $\sqrt{2}$. And as we know, 1D Quantum Harmonic Oscillators do have infinite dimensional Hilbert spaces. So what is going on? Well, this is a feature of the perturbative nature of our calculations. For each case, we took the interaction Hamiltonian as the lowest order interaction term, which only produced a final state which was a linear combination of only two states. So effectively, we reduced the dimensionality of the Hilbert space of the system from infinite to two. And for a two dimensional Hilbert space, the concurrence has a maximum value of 1, because the minimum value of purity, $\gamma = \text{Tr}[\rho_A^2]$ becomes $1/2$. This just tells us that our perturbative calculations are only valid when $\mathfrak{g}/(2\omega_m) \ll 1$, which was also the basis of our approximation.

Now let us look at the graph of concurrence plotted against distance in Fig. 5.2. The graph shows a plot of the concurrence as a function of the distance between the two bodies, for all three cases. Again, constant values of mass and ω_m were chosen as 10^{-14} kg and 10^8 Hz respectively. Here we can see that the degree of entanglement grows inversely




Figure 5.2: Comparison of Concurrence for Different Cases: (a) Static Case; (b) Non-static Correction 1 (c) Non Static Correction 2

with distance. Another feature presented here is that for this range of distances, the static coupling dominates over the other non-static corrections, as can be seen from the scale of each subplot. In fact, if we consider the same parameters as the graph, and we consider the first non-static correction and the concurrence for the static case, the first non-static correction would only start to dominate at $d = 1\text{m}$. And if we were to take the same parameters as the graph and consider the second non-static correction, that would only dominate at $d \approx 10^{15}$ m. One important fact here is that our results are different from Bose *et al.* [17], even though we followed the same methods. The difference is of a factor of $\sqrt{2}$, and it could be that they approximated the concurrence expression differently. Here, I found the first order approximation the usual way, by taking $\mathbf{g}/(2\omega_m) \ll 1$.

Finally let us consider the entanglement entropy. Once again, let us first write down the common expression for Entanglement Entropy found in each of the three cases. This perturbative expression was,

$$\mathcal{S}_A = \log \left(1 + \left(\frac{\mathbf{g}}{2\omega_m} \right)^2 \right) - \frac{2 \left(\frac{\mathbf{g}}{2\omega_m} \right)^2}{1 + \left(\frac{\mathbf{g}}{2\omega_m} \right)^2} \log \left(\frac{\mathbf{g}}{2\omega_m} \right), \quad (5.2)$$

and we approximated this as,

$$\mathcal{S}_A \approx \left(\frac{\mathbf{g}}{2\omega_m} \right)^2 \left(1 - 2 \log \left(\frac{\mathbf{g}}{2\omega_m} \right) \right) + \left(\frac{\mathbf{g}}{2\omega_m} \right)^4 \left(2 \log \left(\frac{\mathbf{g}}{2\omega_m} \right) - \frac{1}{2} \right). \quad (5.3)$$

Let us first look at the perturbative expression. Once again, if one was to take the maxima, they would get $\log(2)$ for $\mathfrak{g}/(2\omega_m) = 1$. This again is a feature of the perturbative nature of our calculations. Usually we would not expect this value, but because of the reasons explained above, this value simply signifies that our perturbation is not valid when $\mathfrak{g}/(2\omega_m) \approx 1$, and is only valid when $\mathfrak{g}/(2\omega_m) \ll 1$. Another feature of this expression is that when you take the limit of the expression for $\mathfrak{g}/(2\omega_m) \rightarrow 0$, we get that the limit is also 0. This is despite the nasty-looking $\log(\mathfrak{g}/(2\omega_m))$ which could have acted as a pole. And the reason for this is because of the coefficient of $\log(\mathfrak{g}/(2\omega_m))$ being an algebraic expression that falls faster than the logarithm.




Figure 5.3: Comparison of the Perturbative Entanglement Entropy and its Approximation.

Let us now take a look at the graph of the Entanglement Entropy against $\mathfrak{g}/(2\omega_m)$ in Fig. 5.3. Specifically at how much agreement there is between the approximation and the perturbative expressions. The approximation is a bit curious, because it has a term of $\log(\mathfrak{g}/(2\omega_m))$, which cannot be expanded around zero, and yet our approximation has the condition of working best near zero. Thus we had to keep the logarithm term in our expansion. And we can see that this led to a good agreement between the perturbative expression and its expansion. The expressions only seem to diverge when $\mathfrak{g}/(2\omega_m)$ reaches near 0.3. But, as stated before, $\mathfrak{g}/(2\omega_m) \ll 1$, and thus our expansion works very well with that range.

Finally let us take a look at the graph in Fig. 5.4. The graph shows a plot of the entanglement entropy as a function of the distance between the two bodies, for all three cases. Again, constant values of mass and ω_m were chosen as 10^{-14} kg and 10^8 Hz respectively.

Once again, we can see that the entanglement entropy grows inversely with distance. How fast it decays depends on the case, but it decays nonetheless. Another feature presented

here is that for this range of distances, the static coupling dominates over the other non-static corrections, as can be seen from the scale of each subplot. The static case has the highest entanglement entropy. The coupling and the concurrence also decayed with distance, but the corrections did not have such difference in magnitude. Here, the entanglement entropy is dominant by more than twice the order of magnitude when it comes to the second non-static correction.

We have now computed the entanglement entropy and concurrence and showed that both are positive for a quantum gravitational field.




Figure 5.4: Comparison of Entanglement Entropies for Different Cases: (a) Static Case; (b) Non-static Correction 1 (c) Non Static Correction 2

The code for the graphs can be found in the Jupyter notebook at https://github.com/guprat/MSc_Dissertation.

Chapter 6

Conclusions

We have now obtained all the results we set out to compute, by relying on elementary perturbation theory and some aspects of general relativity. Our results reinforce the idea that a quantum interaction between gravity and matter is crucial to detect entanglement between two bodies, and if gravity is a classical entity, any experiment based on the mechanisms presented here will not yield expected values. To be clear, if gravity turns out to be a classical entity, all work done here will be moot. On the other hand, if it is quantum, then these mechanisms can be used to build experiments and test the nature of gravity.

And there are a number of similar mechanisms and thought experiments that have been proposed, but none that could be tested. If gravity is indeed quantum, then our results corroborate the importance of an experiment on gravitationally induced entanglement, which relies on the fact that two masses can entangle via the graviton. This would be crucial in not only confirming the quantum properties of gravity, but also unveil crucial properties of the graviton, which as of now, is only a hypothetical particle responsible for spacetime fluctuations.

Although these results are important, there is still a lot of room for research into this mechanism. For one, it would be very interesting to study the effect of this mechanism massive bodies. Another possible area of research is the effect of curvature on this mechanism. So far, we have only studied this mechanism on a Minkowskian background. What happens if we switch it out with a deSitter background. How would the curvature play a role in the mechanism to entangle to masses? If two particles get entangled, the results from the deSitter background are surely to differ.

It makes it all the more worthwhile to study the effect of curvature because the very early universe has been approximated to have a deSitter curvature. And if particles were entangled by this mechanism in the early universe, then this ceases to be a thought experiment anymore, for we would be able to see its evidence in the cosmic microwave background. Those results could be the first step in discovering the elusive secrets of gravity that humanity has been trying to understand for decades, and give clues to settling one of the biggest crises in Physics.

Bibliography

- [1] A. Einstein, Sitzungsberichte der Königlich Preußischen Akademie der Wissenschaften (Berlin), 844 (1915).
- [2] M. Born and P. Jordan, Zeitschrift für Physik **34**, 858 (1925).
- [3] M. Born, W. Heisenberg, and P. Jordan, Zeitschrift für Physik **35**, 557 (1926).
- [4] P. Jordan and W. Pauli, Zeitschrift für Physik **47**, 151 (1928).
- [5] P. A. M. Dirac and N. H. D. Bohr, Proceedings of the Royal Society of London. Series A, Containing Papers of a Mathematical and Physical Character **114**, 243 (1927).
- [6] J. R. Oppenheimer, Phys. Rev. **38**, 725 (1931).
- [7] S. Tomonaga and J. R. Oppenheimer, Physical Review **74**, 224 (1948).
- [8] H. Fukuda, Y. Miyamoto, and S. Tomonaga, Progress of Theoretical Physics **4**, 391 (1948).
- [9] J. Schwinger, Phys. Rev. **74**, 1439 (1948).
- [10] S. S. Schweber and S. B. Treiman, Physics Today **47**, 59 (1994).
- [11] D. Tong, Lectures on string theory (2009).
- [12] Z. Bern, J. J. M. Carrasco, and H. Johansson, Phys. Rev. Lett. **105**, 061602 (2010).
- [13] A. Belenchia, R. M. Wald, F. Giacomini, E. Castro-Ruiz, Č. Brukner, and M. Aspelmeyer, Phys. Rev. D **98**, 126009 (2018).
- [14] A. Belenchia, R. M. Wald, F. Giacomini, E. Castro-Ruiz, Č. Brukner, and M. Aspelmeyer, International Journal of Modern Physics D **28**, 1943001 (2019).
- [15] S. Bose, A. Mazumdar, G. W. Morley, H. Ulbricht, M. Toroš, M. Paternostro, A. A. Geraci, P. F. Barker, M. S. Kim, and G. Milburn, Phys. Rev. Lett. **119**, 240401 (2017).
- [16] C. Marletto and V. Vedral, Phys. Rev. Lett. **119**, 240402 (2017).
- [17] S. Bose, A. Mazumdar, M. Schut, and M. Toroš, Phys. Rev. D **105**, 106028 (2022).
- [18] S. N. Gupta, Proceedings of the Physical Society. Section A **65**, 161 (1952).

- [19] M. Christodoulou, A. Di Biagio, M. Aspelmeyer, Č. Brukner, C. Rovelli, and R. Howl, *Phys. Rev. Lett.* **130**, 100202 (2023).
- [20] E. Schrödinger, *Mathematical Proceedings of the Cambridge Philosophical Society* **31**, 555–563 (1935).
- [21] F. Dyson, *International Journal of Modern Physics A* **28**, 1330041 (2013).
- [22] R. Penrose, *General Relativity and Gravitation* **28**, 581 (1996).
- [23] S. Hossenfelder, Experimental search for quantum gravity, in *Classical and quantum gravity: Theory, Analysis, and Applications*, edited by V. R. Frignanni (Nova Science Publishers, 2012) p. 257–282.
- [24] J. S. Bell, *Physics Physique Fizika* **1**, 195 (1964).
- [25] R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki, *Rev. Mod. Phys.* **81**, 865 (2009).
- [26] S. R. Moulick and P. K. Panigrahi, *Scientific Reports* **6**, 37958 (2016).
- [27] S. Deser, R. Jackiw, and G. 't Hooft, *Phys. Rev. Lett.* **68**, 267 (1992).
- [28] M. J. W. Hall and M. Reginatto, *Journal of Physics A: Mathematical and Theoretical* **51**, 085303 (2018).
- [29] H. Wang, *A Tabletop Experiment for Perturbative Quantum Gravity: A Status Report*, Master's Dissertation, 2023 (The University of Edinburgh).
- [30] M. Christodoulou and C. Rovelli, *Frontiers in Physics* **8**, 10.3389/fphy.2020.00207 (2020).
- [31] D. J. Griffiths and D. F. Schroeter, *Introduction to Quantum Mechanics*, 3rd ed. (Cambridge University Press, 2018).
- [32] C. Cohen-Tannoudji, B. Diu, and F. Laloë, *Quantum Mechanics*, 2nd ed. (Wiley-VCH, 2019).
- [33] V. Balasubramanian, M. B. McDermott, and M. Van Raamsdonk, *Phys. Rev. D* **86**, 045014 (2012).
- [34] S. A. Hill and W. K. Wootters, *Phys. Rev. Lett.* **78**, 5022 (1997).
- [35] P. Rungta, V. Bužek, C. M. Caves, M. Hillery, and G. J. Milburn, *Phys. Rev. A* **64**, 042315 (2001).
- [36] R. J. Marshman, A. Mazumdar, and S. Bose, *Phys. Rev. A* **101**, 052110 (2020).
- [37] S. Brahma, O. Alaryani, and R. Brandenberger, *Phys. Rev. D* **102**, 043529 (2020).
- [38] A. Einstein, *Sitzungsberichte der Königlich Preußischen Akademie der Wissenschaften (Berlin)*, 154 (1918).

- [39] M. E. Peskin and D. V. Schroeder, *An Introduction To Quantum Field Theory*, Frontiers in Physics (Westview Press, Boulder, CO, 1995).
- [40] S. Weinberg, *Gravitation and Cosmology : Principles and Applications of the General Theory of Relativity* (Wiley, New York, 1972).
- [41] M. Maggiore, *Gravitational Waves: Volume 1: Theory and Experiments* (Oxford University Press, 2007).
- [42] S. M. Carroll, *Spacetime and Geometry: An Introduction to General Relativity* (Cambridge University Press, 2019).
- [43] Éanna É Flanagan and S. A. Hughes, *New Journal of Physics* **7**, 204 (2005).
- [44] E. Poisson and C. M. Will, *Gravity: Newtonian, Post-Newtonian, Relativistic* (Cambridge University Press, 2014).
- [45] J. D. Jackson, *Classical Electrodynamics*, 3rd ed. (Wiley, New York, 1999).
- [46] S. M. Kopeikin and G. Schäfer, *Phys. Rev. D* **60**, 124002 (1999).
- [47] A. Cristofoli, N. E. J. Bjerrum-Bohr, P. H. Damgaard, and P. Vanhove, *Phys. Rev. D* **100**, 084040 (2019).
- [48] Y. Iwasaki, *Progress of Theoretical Physics* **46**, 1587 (1971).
- [49] K. Hiida and H. Okamura, *Progress of Theoretical Physics* **47**, 1743 (1972).
- [50] G. Grignani, T. Harmark, M. Orselli, and A. Placidi, *Journal of High Energy Physics* **2020**, 142 (2020).