

Solution of GATE-ST 2023 Q58

SUJAL GUPTA - EE22BTECH11052

Consider the following regression model

$$y_t = \alpha_0 + \alpha_1 t + \alpha_2 t^2 + \epsilon_t, \quad t = 1, 2, \dots, n \quad (1)$$

where α_0 , α_1 and α_2 are unknown parameters and ϵ_t 's are independent and identically distributed random variables each having $\mathcal{N}(\mu, 1)$ distribution with μ unknown. Then which of the following statements is/are true?

- 1) There exists an unbiased estimator of α_1
- 2) There exists an unbiased estimator of α_2
- 3) There exists an unbiased estimator of α_0
- 4) There exists an unbiased estimator of μ

Solution: Assuming that the model is

$$y_t = \alpha_0 + \alpha_1 t + \alpha_2 t^2 + \epsilon_t \quad (2)$$

$$\begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix} = \begin{bmatrix} 1 & A_{11} & A_{12} \\ 1 & A_{21} & A_{22} \\ \vdots & \vdots & \vdots \\ 1 & A_{n1} & A_{n2} \end{bmatrix} \begin{bmatrix} \alpha_0 \\ \alpha_1 \\ \alpha_2 \end{bmatrix} + \begin{bmatrix} \epsilon_1 \\ \epsilon_2 \\ \vdots \\ \epsilon_n \end{bmatrix} \quad (3)$$

Finding mean of the ϵ ,

$$\mathbf{y} = \mathbf{A}\mathbf{x} + \boldsymbol{\epsilon} \quad (4)$$

$$E\boldsymbol{\epsilon} = \begin{bmatrix} E\epsilon_1 \\ E\epsilon_2 \\ \vdots \\ E\epsilon_n \end{bmatrix} = \begin{bmatrix} \mu \\ \mu \\ \vdots \\ \mu \end{bmatrix}$$

The covariance matrix for ϵ is:

$$\mathbf{C}_\epsilon = \mathbf{E}[(\boldsymbol{\epsilon} - \mathbf{E}\boldsymbol{\epsilon})(\boldsymbol{\epsilon} - \mathbf{E}\boldsymbol{\epsilon})^T] \quad (5)$$

$$= E \begin{bmatrix} (\epsilon_1 - E\epsilon_1)^2 & (\epsilon_1 - E\epsilon_1)(\epsilon_2 - E\epsilon_2) & \dots & (\epsilon_1 - E\epsilon_1)(\epsilon_n - E\epsilon_n) \\ (\epsilon_2 - E\epsilon_2)(\epsilon_1 - E\epsilon_1) & (\epsilon_2 - E\epsilon_2)^2 & \dots & (\epsilon_2 - E\epsilon_2)(\epsilon_n - E\epsilon_n) \\ \vdots & \vdots & \ddots & \vdots \\ (\epsilon_n - E\epsilon_n)(\epsilon_1 - E\epsilon_1) & (\epsilon_n - E\epsilon_n)(\epsilon_2 - E\epsilon_2) & \dots & (\epsilon_n - E\epsilon_n)^2 \end{bmatrix} \quad (6)$$

$$= \begin{bmatrix} \text{Var}(\epsilon_1) & \text{Cov}(\epsilon_1, \epsilon_2) & \dots & \text{Cov}(\epsilon_1, \epsilon_n) \\ \text{Cov}(\epsilon_2, \epsilon_1) & \text{Var}(\epsilon_2) & \dots & \text{Cov}(\epsilon_2, \epsilon_n) \\ \vdots & \vdots & \ddots & \vdots \\ \text{Cov}(\epsilon_n, \epsilon_1) & \text{Cov}(\epsilon_n, \epsilon_2) & \dots & \text{Var}(\epsilon_n) \end{bmatrix} \quad (7)$$

Since the ϵ 's are independent vectors,

$$\text{Cov}(\epsilon_i, \epsilon_j) = 0, \quad \forall i \neq j \quad (8)$$

$$\text{Var}(\epsilon_i) = 1, \quad \forall 1 \leq i \leq n \quad (9)$$

Hence,

$$\mathbf{C}_\epsilon = I_{n \times n} \quad (10)$$

$$E(\mathbf{y}) = E(A\mathbf{x} + \epsilon) \quad (11)$$

$$= AE(\mathbf{x}) + E(\epsilon) \quad (12)$$

$$\text{Var}(\mathbf{y}) = \text{Var}(A\mathbf{x} + \epsilon) \quad (13)$$

$$= A\text{Var}(\mathbf{x}) + \text{Var}(\epsilon) \quad (14)$$

$$= A\text{Var}(\mathbf{x}) + I \quad (15)$$