



# Indigenous African fermented dairy products: Processing technology, microbiology and health benefits

Dominic Agyei, James Owusu-Kwarteng, Fortune Akabanda & Samuel Akomea-Frempong

To cite this article: Dominic Agyei, James Owusu-Kwarteng, Fortune Akabanda & Samuel Akomea-Frempong (2019): Indigenous African fermented dairy products: Processing technology, microbiology and health benefits, Critical Reviews in Food Science and Nutrition

To link to this article: <https://doi.org/10.1080/10408398.2018.1555133>



Published online: 22 Jan 2019.



Submit your article to this journal [↗](#)



View Crossmark data [↗](#)

REVIEW



## Indigenous African fermented dairy products: Processing technology, microbiology and health benefits

Dominic Agyei<sup>a</sup> , James Owusu-Kwarteng<sup>b</sup> , Fortune Akabanda<sup>c</sup> , and Samuel Akomea-Frempong<sup>d</sup> 

<sup>a</sup>Department of Food Science, University of Otago, Dunedin, New Zealand; <sup>b</sup>Department of Food Science and Technology, University of Energy and Natural Resources, Sunyani, Ghana; <sup>c</sup>Department of Applied Biology, University for Development Studies, Navrongo, Ghana; <sup>d</sup>Department of Food Science and Technology, Kwame Nkrumah University of Science and Technology (KNUST), Kumasi, Ghana

### ABSTRACT

Africa is known for its rich, ancient tradition in fermented foods. Among these, fermented dairy products represent one category that is widely consumed, contributing to the socio-economic development and food security of the people. In Africa, traditional food fermentation lends itself as a relatively cheap food processing technology that often improves shelf life/food safety as well as nutrition and health via improvement in the levels of specific micronutrients and the action of probiotics. A range of African fermented dairy products (mainly yoghurt-like products) are produced by spontaneous fermentation, and these fermented dairy products harbor rich and valuable microbial diversity, predominated by lactic acid bacteria and yeasts. Detailed knowledge of the production processes, microbiological and biochemical aspects of traditional African dairy fermentation is critical for the development of products with enhanced quality, safety and health benefits for a sustainable food security in the region. This review therefore provides a comprehensive overview of the traditional African fermented dairy processing technology, as well as technologically relevant microorganisms and health benefits associated with fermented dairy products. Efforts aimed at harnessing the functional food potential of these fermented products could help control some food and health challenges facing many countries in the region.

### KEYWORDS

Africa; fermented dairy products; microbiology; functional foods; nutrition; probiotics

### Introduction

In many counties in Africa, milk has played a significant economic and dietary role for centuries. These areas (mostly cattle-keeping regions) include North Africa, the Sudanian Savanna regions (Senegal, Gambia, Southern Mali, Burkina Faso, and northern territories of countries such as Guinea, Cote D'Ivoire, Ghana, Togo, Benin and Nigeria) and the highlands of East Africa (FAO 1990, 2018). Indigenous and nomadic tribes such as the Fulani, Maasai, Tuareg and Borani have consumed milk since prehistoric times (Ndambi, Hemme, and Latacz-Lohmann 2007). However, due to reason of climate and low income, commercial production of milk in many developing countries in Africa has been growing at a very slow pace (FAO 2018). Also, milk constitutes the most widely consumed animal protein in Africa, but it is consumed in very small quantities (World Bank 2014). Per capita milk consumption in Africa sits at 30 liters per year; compared to a global average of 214 liters per year (FAO 2006). This low level of production and consumption of milk hinders the diversification and development of novel dairy products in Africa. Moreover, the warm climate in Africa makes it challenging to preserve highly perishable food products such as milk, particularly in resource poor communities that lack the necessary processing/preservation (chilling) systems. Fermentation as a

processing technology is therefore important in Africa, not only because it serves as a form of preservation, but also meets the need for diversifying products from milk, while helping to improve the functionality, digestibility, as well as nutritional profile of milk and its products. This review gives a comprehensive profile of several traditional fermented dairy products from Africa. An assessment of the microbiology of these products is presented, together with a discussion of the food and health application of these fermented products.

### Fermentation as a processing technology in Africa

Food processing technologies are the various set of operations that modify food ingredients and raw materials, with the intention of obtaining products with desired quality attributes that meet nutritional and functional requirements. There are primitive and contemporary food processing technologies including simple processing techniques such as drying, salting, smoking and fermentation. While food processing still has the main objective of providing a nutritious diet, other aspects such as preservation and generation of wealth for producers are added advantages. In Africa, fermentation of food is one of the oldest methods of food processing and preservation and many traditional food

processing techniques have been handed down from generation to generation for centuries (Nduko et al. 2017).

Food fermentation as a processing technology comprises of the alteration of raw materials (food ingredients) into a melange of value-added products by employing the phenomena of microbial growth and their activities on various substrates (MacDonald and Reitmeier 2017). In prehistoric times, foods were fermented with the primary purpose of extending shelf life and enhancing sensory characteristics of the fermented food products (Hui et al. 2004). However, the benefits of food fermentation has expanded to include inhibition of pathogenic microorganisms and improvement in nutritional value or digestibility (Terefe 2016). Typically, food fermentation processes practiced in Africa are mostly spontaneous. Thus, the fermentation do not rely on well-defined starter cultures, durations of fermentation and temperatures are poorly controlled, leading variations in the quality of products (Nduko et al. 2017). Notwithstanding, various fermented food products are produced in different African countries. These fermented foods include alcoholic and nonalcoholic beverages such as palm wine and *pito*, breads, porridges and fermented dairy products including yoghurts and cheeses. These fermented food products are obtained from vegetal raw materials such as starchy root crops, vegetables proteins, leaves and fruits (Marshall and Mejía-Lorío 2011) or animal sources such as fish, meats and milk.


Foods such as *poto poto* (maize) in Congo, *Ogi* (maize, millet) in Nigeria, *Kenkey* and *Banku* (maize) in Ghana and *Togwa* (sorghum) in Tanzania are value added products from fermented cereals (Abriouel et al. 2006; Greppi et al. 2013; Mugula et al. 2003). Legumes and African oil seeds are vegetables that are rich in proteins. Most of these undergo alkaline fermentation resulting in foods such as *Ugba* (African oil bean seed) in Nigeria and *Ogili* (Melon and Castor oil seeds) in East, Central and West Africa (Ogbulie, Nsofor, and Nze 2014; Barber, Achinewhu, and Ibiam 1988). Additionally, cassava is one of the major starchy root crop fermented to elongate its shelf life and reduce cyanide contents for safer consumption (Aworh 2008). *Gari*, *Kivunde* and *Lafun* are value added products of fermented cassava in West and Central Africa, Tanzania and West Africa respectively. Moreover, fish and milk are the major animal protein products that are widely fermented purposely to increase shelf life. *Koobi*, *Momone*, and *Kako* in Ghana (Abbey, Hodari-Okoe, and Osei-Yaw 1994); *Lanhoun* in Benin and Togo (Dossou-Yovo 2007) and *Dagaa* in Uganda (Koffi-Nevr et al. 2011) are all value added fermented fish products. *Wagashi* (Ghana) obtained from milk, also called *Warakasi* in Nigeria (Haaland 2007), *jben* and *Iben* (Benkerroum 2013) (northern and eastern Africa) (Aworh 2008), *Mursik* and *Amabere amaruranu* in Kenya (Nduko et al. 2017) and *Amasi* in South Africa (Chelule et al. 2010). Fermented butter such as *Kibe* and cheese (*Areesh* and *Domiat*) are other examples of fermented dairy products in Africa.

## Fermented milk products from Africa

Africa is home to a wide range of centuries-old indigenous fermented dairy products that are still consumed, playing significant role in providing key nutrient requirements and income to the people of this region (Asogwa, Okoye, and Oni 2017;

Wilson 1988). African Fermented Dairy Products (AFDP) are predominantly yoghurts or yoghurt-like foods and beverages. Depending on country or local region of production, these fermented dairy products with similar or same production methods and final product characteristics may be known by different names. Majority of AFDP are made from cow's milk, but milk from goat, buffalo, camel or ewe are also used in relatively large quantities (Jans et al. 2017; Tamang and Kailasapathy 2010). Prior to the introduction of modern refrigeration in African homes, milk and other raw food commodities were preserved by fermentation (Jespersen 2003; Mensah 1997), a process that likely improved the nutritional quality, safety and digestibility of milk. Generally, food fermentation in Africa occurs spontaneously and takes place at the household level or at a small-scale production site (Caplice and Fitzgerald 1999; Ezeji and Ojimelukwe 1993; Oyewole 1997), although a few products such as *Amasi* in South Africa and *Madila* in Botswana are currently produced on commercial scale (Ohiokpehai 2003; Schutte 2013). Spontaneous fermentation occurs due to the activities of microorganisms inherent in the raw milk, from the environment or processing equipment (Kebede et al. 2007; Oyewole 1997). During the production of fermented dairy products in Africa, milk may be mixed with cereals or other plant materials before or after fermentation to yield unique rich products of cereal and milk-based blends (Franz et al. 2014). The characteristics of African fermented dairy products are often influenced by factors such as the quality and the type of raw milk used, the production methods followed and the regional or local climatic conditions (Mensah 1997; Wouters et al. 2002). Additionally, the production processes of African fermented dairy products are largely dependent on unique recipes handed down from one generation to another. Subsequently, we provide brief descriptions of African fermented dairy products taking into account the similarities or differences that may exist in processing methods and in the final product characteristics.

In Africa, different dairy animals including cow, buffalo, camel, goat or ewe serve as sources of raw milk for the production of fermented yoghurt-like products. Irrespective of the source of milk, a number of fermented dairy products may be produced using similar or slightly different traditional processes. Thus, although several generic names are applied to the different AFDP products, the real number of distinct products may be fewer. For example, *nunu* is a generic name for spontaneously fermented yoghurt-like milk products in northern Ghana (Akabanda et al. 2013; Owusu-Kwarteng et al. 2017). However, very similar products are known by different local names in other parts of West Africa such as *nyamie* in southern Ghana, *fenè* in Mali, *nono* in Nigeria and sour milk in Côte d'Ivoire (Akabanda et al. 2013; Obodai and Dodd 2006; Owusu-Kwarteng et al. 2017; Savadogo et al. 2004; Wullschlegler et al. 2013). In other parts of Africa, similar fermented yoghurt-like dairy products include *leben/lben* in North Africa (Bensalah, Delorme, and Renault 2009; Mangia et al. 2014), *amasi* and *kefir* in South Africa (Osvik et al. 2013; Witthuhn, Schoeman, and Britz 2004), *suusac* in Kenya and Somalia (Jans et al. 2012a; Njage et al. 2011), *ergo* in Ethiopia (Gonfa et al. 1999), and *gariss* in Sudan



**Figure 1.** Pictorial representation of the country of origin for some African fermented dairy products.

(Abdelgadir et al. 2008). The names and origin of some African fermented dairy products are shown in Figure 1.


During the processing of milk into yoghurts in Africa, the freshly collected milk may be sieved or filtered to remove animal fur and other debris that may have contaminated the milk during collection and handling on the farm. Generally, African fermented dairy products are produced from raw (non-pasteurized) or heated (pasteurized) milk. In addition to destroying microbial contaminants in the milk, heating also leads to the denaturation of whey proteins. The denatured proteins together with milk caseins then precipitate at low pH, brought about by lactic acid fermentation, leading to the typical properties associated with yoghurt. For AFDP in which the raw milk is heated, milk fat may be removed prior to fermentation by churning (Figure 2). On the other hand, fat or whey removal may be practiced for non-pasteurized fermented products after the fermentation step to yield varied sets of yoghurt-like products.

An important step in the processing of fermented dairy products in Africa is the fermentation process. AFDP are generally produced by spontaneous fermentation or back-slopping, with fermentation temperatures often reflecting the ambient temperature of the local region where the fermentation takes place. Durations of fermentation varies widely among products, which may be a few hours as occurs in *nunu* in West Africa (Akabanda et al. 2013) or several months, sometimes up to a year, as occurs in the production of *chekapmkaika* (produced from extended fermentation of *Kwerionik*) in Uganda (Schutte 2013). In fact, some of these fermentations can go on for so long that the milk products tend to have very long or indefinite shelf-lives, allowing their use or

consumption during the dry seasons when fresh milk is sometimes unavailable (Ørskov 1995; Schutte 2013). A special type of continuous fermentation is reported for *gariss* in Sudan where camel milk is fermented in leather bags made of tanned goat skin, embedded in green or wet grass and then placed on the back of roaming camels leading to continuous shaking by the jerky walk of the camels. Fresh camel's milk is continuously added to the fermenting batch to make up the volume whenever part of the fermented product known as *gariss* is withdrawn for consumption, and this fermentation practice can continue for several months (Abdelgadir et al. 2008; Abdelgadir, Ahmed, and Dirar 1998; Dirar 1993).

Fermentation vessels for the production of AFDP usually include calabashes, gourds, clay pots, leathers bags or plastic containers. Another important practice during the fermentation of milk in Africa is the smoke pretreatment of fermentation gourds as occurs during the production of *amabere*, *mursik* and *suusac* in Kenya (Jans et al. 2012a; Nieminen et al. 2013; Njage et al. 2011; Nyambane et al. 2014); *pendidaam* and *kindirmou* in Cameroon (Jiwoua and Milliere 1990; Libouga, Ngang, and Halilou 2005; Mbawala et al. 2013); *ergo* in Ethiopia (Gonfa et al. 1999); and *makamo* and *kwerionik* in Uganda (Nakavuma et al. 2012; Schutte 2013). For the production of *kule-naoto* which is very popular among the Maasai community in Kenya, the fermentation gourd made from *Lagenaria siceraria* is gently rubbed with the burning end of a chopped stick of *Olea africana*, a tree locally known as *enkidogoe* (Mathara 1999; Mathara et al. 2004). On the other hand, *ititu* in Ethiopia is produced by fermenting milk in *gorfa*, a vessel traditionally woven from fibers of selected plants treated with leaves of *Ocimum basilicum* for both cleaning and imparting desirable flavors to





**Figure 2.** Generalized flow diagram for the production of various African fermented yoghurt-like dairy products (country or region of production). ‡: Fermentation is spontaneous or by back-slopping. Ψ: Product results from fermentation of milk in special smoke treated gourds. †: Product results from semi-continuous fermentation of milk in bags of goatskin hanged on camel's back.

the product, or in a wooden amuyou (Andualem and Geremew 2014; Coppock, Holden, and O'Connor 1991). These pretreatments of fermentation gourds or containers and the selection of particular plant materials as fermentation vessels impart onto the AFDP distinct natural aromas and tastes associated with these products.

African fermented dairy products (with the exception of few cheeses) can generally be considered as stirred type yoghurt. Typically, the gel formed after milk fermentation is gently stirred to obtain a smooth and viscous but still pourable product that is consumed. During stirring, the gel formed break up and lead to the formation of a highly viscous, non-Newtonian liquid product (Jaros and Rohm 2003). There are however, variations in the textural and rheological properties of African fermented yoghurts due to a number of processing factors such as quality of raw milk used, extent of boiling, fermentation time and acidity levels, and the dominant fermentation microorganisms. Similarly, pH and acidity which affect the sensory qualities also vary widely among the various products. For example, the final pH of *nunu* in Ghana can reach as low as 3.0–3.1 (Akabanda et al. 2013) while *amasi* in South Africa or Namibia ranges between 4.0 and 4.5 (Beukes, Bester, and Mostert 2001). Generally, few investigations have focused on the rheological, textural and sensory characteristics of AFDP. However, research geared towards standardizing the rheological, textural and sensory properties of African fermented milk yoghurts is required for product upgrade and consumer acceptability.

### Microbiology of African fermented dairy products (AFDP)

The microbiota of fermented dairy products significantly influences the quality, safety and general product

characteristics. Therefore, detailed knowledge of the microflora of indigenous African fermented dairy products and their technological roles in fermentation is essential in selecting and defining functional starter culture for improved product quality and safety and overall food security. Microbiological research on African fermented dairy products have been limited to the isolation and characterization of the predominant microorganisms with a few studies focusing on the technological roles of the microorganisms, aimed at developing starter cultures for controlled fermentation. Subsequently, this review will provide a brief overview of the fermentative and technologically relevant microorganisms that have so far been isolated and identified from African fermented milk products.

Generally, production of fermented milk products relies on the old-age traditional process of natural fermentation practiced throughout Africa. Raw or heated milk is allowed to ferment spontaneously, or by 'back-slopping' where a part of a previous batch of a successful fermentation is used as starter to inoculate the new batch (Holzapfel 2002; Josephsen and Jespersen 2004; Owusu-Kwarteng et al. 2017). In such fermentation practices, external factors such as regional or local climatic conditions (season, temperature), quality and composition of the raw material, and the duration of the fermentation process influence the composition and activity of the microbial community during the fermentation (Mathara et al. 2004), which in turn influence the unique organoleptic (Steinkraus 1994) and functional properties of the final products.

The microflora of African fermented dairy products is dominated by mainly lactic acid bacteria (LAB), often associated with yeasts. Table 1 provides an overview of LAB and yeast species isolated from African fermented dairy products. While LAB species are the only group of

microorganisms identified so far in some products such as *amasi* (Osvik et al. 2013), *ergo* (Gonfa et al. 1999), *fenè* (Wullschleger et al. 2013), *lben* (Ouadghiri et al. 2009), *ititu* (Kassaye et al. 1991), *kiviguto* (Karenzi et al. 2012), *mabisi* (Schoustra et al. 2013) and *zabady* (El-Baradei, Delacroix-Buchet, and Ogier 2008), both LAB and yeasts have been isolated from several other fermented dairy products such as *amabere* (Nyambane et al. 2014), *gariss* (Abdelgadir et al. 2008), *kule naoto* (Mathara et al. 2004), *lben* (Mangia et al. 2014), *nunu* (Akabanda et al. 2013), *nyarmie* (Obodai and Dodd 2006), *rob* (Abdelgadir, Ahmed, and Dirar 1998; Abdelgadir et al. 2001) and *suusac* (Jans et al. 2012a; Njage et al. 2011).

Among the LAB, *Lactococcus* (*Lc.*) *lactis* and its subspecies *lactis* and *cremoris* have been isolated from most AFDP except a few products such as *amabere* in Kenya (Nyambane et al. 2014), *gariss* in Sudan (Abdelgadir et al. 2008) and *ititu* from Ethiopia (Kassaye et al. 1991). In products where *Lc. lactis* have been identified, their counts range from about  $10^6$  to  $10^{10}$  CFU/mL in the final products where they are usually but not always the predominant species (Table 1). The predominance of *Lc. lactis* in AFDP, irrespective of the region or country of production, indicates their adaptability to the prevailing ambient temperatures under which spontaneous fermentations of milk take place in Africa, and thereby contributing to milk fermentation in the region. *Lc. lactis* are widely used as starter culture for dairy fermentations owing to their various technological properties such as high adaptation to lactose metabolism, diacetyl formation and contribution to flavor development and food preservation (Teuber 2009).

In addition to *Lactococcus lactis*, *Lactobacillus* species including *Lb. fermentum*, *Lb. delbrueckii* and *Lb. plantarum* have been isolated among the dominant species in several AFDP with detection levels of  $10^8$  to  $10^{10}$  CFU/mL (Abdelgadir et al. 2008; Beukes, Bester, and Mostert 2001; Doutoum et al. 2013; Jiwoua and Milliere 1990; Mathara et al. 2004; Nyambane et al. 2014; Obodai and Dodd 2006; Ouadghiri et al. 2009; Witthuhn, Schoeman, and Britz 2004). Other less frequently isolated lactobacilli/coccobacilli in AFDP include *Lb. casei*, *Lb. paracasei*, *Lb. bulgaricus*, *Lb. helveticus*, *W. confusa*, *Lb. acidophilus*, *Lb. rhamnosus* and *Lb. brevis* (Table 1). While *Lb. delbrueckii* and *Lb. fermentum* are known to be typical dairy microorganisms, their technological role and contribution to the safety and quality of AFDP has not been fully exploited. However, the widespread predominance of lactobacilli in AFDP across the region is an indication of their adaptation to the local climatic conditions and potential technological role during fermentation of milk in Africa. In an attempt to develop starter cultures for the fermentation of *nunu* in West Africa, Akabanda et al. (2014) evaluated *Lb. fermentum*, *Lb. plantarum*, *Lb. helveticus*, and *Leuconostoc mesenteroides* for various technological properties and further tested the strains as starter cultures in fermentation trials for *nunu* production. The selected lactobacilli strains, whether used singularly or in combinations in the fermentation trials, produced *nunu*

with highly acceptable consumer sensory characteristics and improved nutritional quality (Akabanda et al. 2014).

Another species of LAB that have recently been isolated in high numbers ( $>10^8$  cells/mL) from various AFDP is *Streptococcus infantarius* subsp. *infantarius* (*Sii*). *Sii* were not described as dairy fermentation microbiota until it was first identified as a predominant species in Sudanese spontaneously fermented camel milk, *gariss*, by Abdelgadir et al. (2008). Following this, *sii* has been isolated among the predominant LAB species in other AFDP such as *suusac* in Kenya and Somalia (Jans et al. 2012a; Njage et al. 2011), *fenè* in Mali (Wullschleger et al. 2013), and the Ivorian fermented cow milk (Jans et al. 2017). Since the novel discovery of *sii* in traditional AFDP, further investigations have provided insights into the evolutionary lineages and adaptations of *Sii* to the dairy environment (Jans et al. 2012b, 2013, 2016). As a member of the *Streptococcus bovis*/*Streptococcus equinus* complex (SBSEC), *Sii* is generally considered as commensal inhabitants of the gastrointestinal tract of animals and humans, and at the same time, opportunistic pathogens, which potentially qualifies them as pathobionts (Chow, Tang, and Mazmanian 2011; Jans et al. 2015; Schlegel et al. 2003). However, the pathogenicity of *Sii* and other novel SBSEC species cannot be very certain, as the methodologies for obtaining most epidemiological data in the past were based on less discriminative biotype classifications as opposed to current taxonomic rearrangements and DNA-based classifications (Jans et al. 2015; Schlegel et al. 2003). In addition to *Sii*, other species of *Streptococcus* including *S. thermophilus*, *Streptococcus gallolyticus* subsp. *macedonicus* (*Sgm*), *S. salivarius* and *S. agalactiae* have frequently been isolated from AFDP. Except *S. thermophilus* which is currently approved for use in dairy fermentation by European Food Safety Authority (EFSA 2016), all other species of the genus *Streptococcus* including *Sii* are neither classified by the qualified presumption of safety (QPS) nor have the status of Generally Recognized as Safe (GRAS) (EFSA 2016). However, due to the wide distribution, predominance and adaptation of *Sii* to AFDP, there is the need to further investigate their technological role in dairy fermentation and thoroughly carryout the needed safety assessments to justify their development and use as dairy starter for improved safety and quality of AFDP. A more extensive review on technologically important microorganisms in African fermented dairy products focusing on African *Streptococcus infantarius* variants has recently been published by Jans et al. (2017).

*Enterococcus* species, although described as contaminants in milk, may play some role during milk fermentation as they have frequently been isolated among dominant organisms in AFDP. *Enterococcus* spp., particularly *Enterococcus faecium* and/or *E. faecalis*, have been isolated from AFDP products such as *fenè*, *kule naoto*, *gariss*, *leben/lben*, *nunu*, *pendidam* and *suusac* (Table 1). Generally, most enterococci are part of the intestinal microbiota of mammals and birds (Ludwig, Schleifer, and Whitman 2009), and their presence in water or food is regarded as an indicator of fecal contamination from animal and human sources (Franz, Holzapfel,

**Table 1.** Lactic acid bacteria and yeasts isolated from African fermented dairy products.

Product name (country/region)	Identified microflora		Identification methodology	References
	LAB species	Yeast species		
Amabere (Kenya)	<i>Lb. plantarum</i> , <i>Leuc. mesenteroides</i> , <i>S. thermophilus</i> , <i>Lb. bulgaricus</i> , <i>Lb. helveticus</i> , <i>Lb. fermentum</i>	<i>Sac. cerevisiae</i> , <i>Trichosporon mucoides</i> , <i>C. famata</i> , <i>C. albicans</i>	API 50 CH, API 20C AUX	Nyambane et al. 2014
Amasi (South Africa)	<i>Lc. Lactis</i> , <i>E. faecalis</i> , <i>Lb. casei</i> , <i>Lb. paracasei</i> , <i>Lb. plantarum</i> , <i>Leuc. Pseudomesenteroides</i> , <i>Lc. lactis</i> subsp. <i>lactis</i> , <i>Lb. delbrueckii</i> subsp. <i>lactis</i> , <i>Lb. plantarum</i> , <i>Leuc. mesenteroides</i> subsp. <i>dextranicum</i>	ND	16S clone library and DGGE. Basic phenotypic characterization, API 50 CH	Osvik et al. 2013; Beukes, Bester, and Mostert 2001
Amasi (Zimbabwe)	<i>Lc. lactis</i> subsp. <i>lactis</i> , <i>Lb. plantarum</i> , <i>Lb. helveticus</i>	<i>Sac. cerevisiae</i> , <i>C. lusitanae</i> , <i>C. colliculosa</i> , <i>S. dairensis</i> , <i>Dekera bruxillensis</i> , <i>C. lipolytica</i> , <i>C. tropicalis</i>	Phenotypic methods, API ID32 C test strips	Feresu and Muzondo 1990; Gadaga et al. 1999
Ergo (Ethopia)	<i>Lb. mesenteroides</i> <sup>a</sup> , <i>Lc. lactis</i> subsp. <i>cremoris</i> , <i>Lc. lactis</i> subsp. <i>lactis</i> , <i>Leuc. cremoris</i> , <i>S. thermophilus</i> , <i>Lb. delbrueckii</i> , <i>Lb. homi</i> , <i>Micrococcus</i> spp.	ND	Basic phenotypic characterization	Gonfa et al. 1999
Fènè (Mali)	<i>Enterococcus</i> spp., <i>Sii</i> , <i>Lb. fermentum</i> , <i>Lb. plantarum</i> , <i>Lc. lactis</i> subsp. <i>lactis</i> , <i>P. pentosaceus</i> , <i>W. confusa</i> .	ND	Rep-PCR for clustering, species-specific PCR assay	Wullschleger et al. 2013
Gariss (Sudan)	<i>Lb. fermentum</i> , <i>Sii</i> , <i>E. faecium</i> , <i>Lb. helveticus</i> ;	<i>Kluyveromyces marxianus</i> , <i>Issatchenkia orientalis</i>	rep-PCR for clustering, 16S rRNA gene, rpoB, sodA, gtf sequencing, API 50 CHL, API ID 32C and 26S rRNA gene for yeasts	Abdelgadir et al. 2008
Ititu (Ethopia)	<i>Lb. casei</i> and <i>Lb. plantarum</i> .	ND	phenotypic methods	Kassaye et al. 1991
Kiviguto (Rwanda)	<i>Lc. lactis</i> , <i>Leuc. mesenteroides</i> subsp. <i>mesenteroides</i> <i>Leuc. pseudomesenteroides</i>	ND	phenotypic methods and sequencing of 16S rDNA and/or 16S-23S rDNA intergenic transcribed spacer (ITS) region	Karenzi et al. 2012
Kule-naoto (Kenya)	<i>E. faecium</i> , <i>Lb. fermentum</i> , <i>Lb. plantarum</i> , <i>Lc. Lactis</i> , <i>Lb. acidophilus</i> , <i>Lb. casei</i> , <i>Lb. paracasei</i> , <i>Lb. rhamnosus</i> , <i>Leuc. mesenteroides</i> ,	yeasts not identified to genus or species	Basic phenotypic characterization and API 50 CHL	Mathara et al. 2004
Kwerionik (Uganda)	<i>Lb. plantarum</i> , <i>E. faecalis</i> , <i>Lb. paracasei</i> subsp. <i>paracasei</i> , <i>Lb. casei</i> subsp. <i>casei</i> , <i>Lc. lactis</i> subsp. <i>lactis</i> , <i>E. faecium</i> and <i>Leuc. mesenteroides</i> subsp. <i>mesenteroides</i>	ND	Carbohydrate fermentation and other phenotypic tests, ITS-PCR.	Nakavuma et al. 2012; Schutte 2013
Lben (Algeria)	<i>Lc. lactis</i> subsp. <i>lactis</i> , <i>Lc. lactis</i> subsp. <i>lactis</i> biovar <i>diacetylactis</i> , <i>E. faecalis</i> , <i>Lb. brevis</i> , <i>Lc. lactis</i> subsp. <i>cremoris</i> , <i>Leuc. lactis</i> , <i>Leuc. mesenteroides</i> subsp. <i>dextranicum</i>	<i>Kluyveromyces lactis</i> , <i>Sac. cerevisiae</i>	API 50 CHL, API 20 STREP, API 20C AUX	Mangia et al. 2014
Leben (North Africa)	<i>E. faecalis</i> , <i>E. faecium</i> , <i>Lc. lactis</i> , <i>S.thermophilus</i> .	ND	Basic phenotypic characterization, 16S rRNA gene, pepN and pepO sequencing	Bensalah, Delorme, and Renault 2009
Mabisi (Zambia)	<i>Acinetobacter ursingii</i> , <i>Citrobacter freundii</i> , <i>Lc. lactis</i> , <i>S. equinus</i> , <i>S. thermophilus</i> , <i>E. durans</i> , <i>Lb. brevis</i> , <i>Lb. kefiranoferiens</i> , <i>Lb. plantarum</i> , <i>Leuc. garlicum</i> , <i>Leuc. pseudomesenteroides</i>	ND	V1–V4 region analysis of 16S rRNA gene	Schoustra et al. 2013
Madila (Botswana)	<i>Lb. plantarum</i> , <i>Lb. acidophilus</i> , <i>Lc. lactis</i> , <i>Lb. fermentum</i> , <i>Lb. brevis</i> and <i>Lb. delbrueckii</i>	ND	Basic phenotypic characterization	Ohenhen, Imarenezor, and Kihuha 2013
Mafi (South Africa)	<i>Lc. lactis</i> subsp. <i>lactis</i> , <i>Lb. delbrueckii</i> subsp. <i>lactis</i> , <i>Lb. plantarum</i> , <i>Leuc. mesenteroides</i> subsp. <i>dextranicum</i>	ND	Basic phenotypic characterization, API 50 CH	Beukes, Bester, and Mostert 2001
Masse (Mozambique)	<i>Lc. lactis</i> subsp. <i>Lactis</i> , <i>Leuc pseudo-mesenteroides</i> , <i>Leuc lactis</i> , <i>Leuc garlicum</i>	ND	PCR and DNA sequencing of the 16S ribosomal RNA (rRNA) gene	Schutte 2013
Mursik (Kenya)	<i>Lb. kefir</i> , <i>Lb. casei</i> , <i>Lb. paracasei</i> , <i>Lb. rhamnosus</i> , <i>cereus</i> , <i>Lb. brevis</i> , <i>Lb. helveticus</i> , <i>Lb. pontis</i> ,	<i>C. sphaerica</i> , <i>C. krusei</i> <i>C. kefir</i> , <i>Sac. fermentati</i>	16S and 18S rRNA gene sequencing, API 32C AUX;	Nieminen et al. 2013
Nunu (Ghana)				Akabanda et al. 2013

(continued)

Table 1. Continued.

Product name (country/region)	Identified microflora		Identification methodology	References
	LAB species	Yeast species		
	<i>Lb. fermentum</i> , <i>Lb. plantarum</i> , <i>Leuc. mesenteroides</i> ; <i>E. italicus</i> , <i>Lactococcus</i> spp., <i>Lb. helveticus</i> , <i>E. faecium</i> , <i>W. confusa</i> ;	<i>Pichia kudriavzevii</i> , <i>Sac. cerevisiae</i> , <i>C. parapsilosis</i> , <i>C. rugosa</i> , <i>C. tropicalis</i> , <i>Galactomyces geotrichum</i>	Basic phenotypic characterization followed by GTG-5 fingerprinting and 16S rRNA gene/26S rRNA gene sequencing	
Nyarmie (Ghana)	<i>Leuc. mesenteroides</i> subsp. <i>mesenteroides</i> , <i>S. thermophilus</i> , <i>Lb. delbrueckii</i> subsp. <i>bulgaricus</i> , <i>Lb. helveticus</i> , <i>Lb. delbrueckii</i> subsp. <i>Lactis</i> , <i>Lc. lactis</i> ;	<i>Sac. cerevisiae</i> , <i>Candida</i> spp., <i>Trichosporon cutaneum</i>	16S and 18S rRNA gene DGGE, API 50 CHL, API 20 STREP, API 20C AUX	Obodai and Dodd 2006
Omeshikawa (Namibia)	<i>Lb. helveticus</i> , <i>Lb. kefir</i> , <i>Lb. casei</i> , <i>Lb. rhamnosus</i> , <i>Lb. paracasei</i> , <i>Lc. lactis</i> subsp. <i>Lactis</i> , <i>Leuc. pseudomesenteroides</i> , <i>E. faecium</i> , <i>E. durans</i> ,	ND	PCR and DNA sequencing of the 16S ribosomal RNA (rRNA) gene	Schutte 2013
Pendidaam (Cameroon)	Presumptive <i>Lb. spp.</i> and <i>Streptococcus/Enterococcus</i> spp.; <i>Lb. delbrueckii</i> , <i>Lb. helveticus</i> , <i>Lb. fermentum</i> , <i>Lb. plantarum</i> , <i>E. faecalis</i> , <i>E. faecium</i> , <i>S. thermophilus</i> <sup>a</sup> , <i>Lb. casei</i> , <i>Leuc. mesenteroides</i> , <i>Leuc. paramesenteroides</i>	ND	Phenotypic characterization	Jiwoua and Milliere 1990; Mbawala et al. 2013
Rob (Sudan)	<i>Lb. fermentum</i> , <i>Lb. acidophilus</i> , <i>Lc. lactis</i> , <i>Strep. salivarius</i> .	<i>Sac. cerevisiae</i> and <i>C. kefir</i> .	ITS-PCR, API 50 CH and the ID 32 API kits	Abdelgadir, Ahmed, and Dirar 1998; Abdelgadir et al. 2001
Suusac (Kenya/Somalia)	<i>E. faecium</i> , <i>Lb. helveticus</i> , <i>Sii</i> , <i>S. salivarius/thermophilus</i> , <i>W. confusa</i> , <i>E. faecalis</i> , <i>Lb. fermentum</i> , <i>Lc. Lactis</i> subsp. <i>Lactis</i> , <i>Leuc. lactis</i> , <i>Leuc. mesenteroides</i> , <i>Lb. curvatus</i> , <i>Lb. plantarum</i> , <i>Lb. salivarius</i> , <i>Lc. raffinolactis</i> , <i>Leuc. mesenteroides</i> subsp. <i>mesenteroides</i>	<i>C. famata</i> , <i>C. inconspicua</i> , <i>C. lusitaniae</i> , <i>Cryptococcus laurentii</i> , <i>Rhodotorula mucilaginosa</i> , <i>Sac. cerevisiae</i> , <i>Trichosporon mucoides</i> , <i>Trichosporon cutaneum</i> , <i>C. krusei</i> , <i>Geotrichum penicillatum</i> .	Rep-PCR for clustering, species-specific PCR assay, 16S rRNA gene sequencing Yeasts: API 20C AUX and DNA-based. API 50 CHL and API 20C AUX	Lore, Mbugua, and Wangoh 2005; Njage et al. 2011

*Lb.*, Lactobacillus; *Lc.*, Lactococcus; *Leuc.*, Leuconostoc; *E.*, Enterococcus; *S.*, Streptococcus; *W.*, Weissella; *Sac.*, Saccharomyces; *C.*, Candida; ND, not determined.

<sup>a</sup>*Lb. mesenteroides* is not an official species designation according to DSMZ standing nomenclature.

and Stiles 1999; Godfree, Kay, and Wyer 1997). While *Enterococcus* spp. are frequently detected in dairy products, their status in terms of food safety is quite controversial due to their ability to become opportunistic pathogens (Franz, Holzapfel, and Stiles 1999). Additionally, there are concerns regarding the propensity of fecal enterococci to be resistant to antibiotics, particularly vancomycin, and to transfer such resistant traits via mobile genetic elements (Giraffa 2002; Teuber, Meile, and Schwarz 1999). Therefore, enterococci just like *Sii*, are currently not approved for general application in foods (EFSA 2016). Notwithstanding, *E. faecium* and *E. faecalis* preparations have been used as probiotics (Fuller 1989; Tournut 1989).

*Leuconostoc mesenteroides* and its subspecies *dextranicum* and subspecies *mesenteroides*, and *Leuconostoc pseudomesenteroides* have also been isolated from various AFDP (Table 1). Leuconostocs in dairy fermentation are considered secondary or associated bacteria because they do not bring about significant change in product characteristics when used alone as pure culture (Vedamuthu 2006). However, in co-culture with lactococci and leuconostocs metabolize citrate present in milk to produce diacetyl, thereby impacting on the flavor of the fermented product. Thus, lactococci and leuconostocs act synergistically in generating diacetyl from citrate found in milk. Dairy leuconostocs possess the enzyme needed to metabolize citrate but ferment milk lactose very slowly and are therefore unable to generate the optimum acidic environment necessary for citrate metabolism. On the other hand, the lactococci, which rapidly ferments lactose in

milk to facilitate the uptake of citrate by leuconostocs often lack the enzymes needed to metabolize the citrate (Vedamuthu 2006). Thus, diacetyl generation in mixed culture fermentations such as those occurring in AFDP may require the associative activity of lactococci and leuconostocs. Therefore, there is the need to consider leuconostoc-lactococci compatibility when selecting mixed starter cultures for the production of AFDP to generate the needed typical flavors associated with these products.

In addition to the predominant LAB, yeasts are frequently isolated from AFDP with counts ranging from  $10^3$  to  $10^8$  CFU/mL of fermented sample. While the fermentation of dairy products in Africa are reportedly predominated by LAB, the associated yeasts could be an essential part of the technologically relevant fermentative microflora. Generally, the role of yeasts in AFDP has been underestimated with most investigations providing partial overview of the microflora of AFDP by focusing only LAB alone or through restricted use of methods to isolate and identify the predominant microflora. Among the few AFDP in which yeasts have been studied, *Saccharomyces cerevisiae* seems to be the predominant yeasts often isolated while *Candida* spp. represent the largest genus, with many products containing more than one species of yeast. Apart from *S. cerevisiae*, there is wide variation in other yeasts species occurring in the different AFDP (Table 1). Representatives of the yeast genera *Candida*, *Cryptococcus*, *Debaryomyces*, *Geotrichum*, *Issatchenkia*, *Khuyveromyces* and *Saccharomyces cerevisiae* are frequently isolated from milk and milk products (Deak and



Beuchat 1996; Quigley et al. 2013). Due to their low pH, yoghurts and yoghurt-like milk products exhibit selective environment for the growth of yeasts (Suriyarachchi and Fleet 1981). During fermentation, yeasts may interact with the fermenting LAB to contribute positively to the fermentation by supporting the growth of the bacteria through the production of essential growth metabolites, such as amino acids and vitamins, or through lactose and galactose metabolism, proteolysis, lipolysis and enzymatic degradation, which contribute to flavor development (Fleet 1990; Jakobsen and Narvhus 1996; Quigley et al. 2013).

## Food and health applications of fermented dairy products

### Nutritional composition of milk and fermented dairy products

Generally, raw milk is a complex food that contains numerous nutrients, serving as source of lipids, proteins, amino acids, vitamins and minerals. Additionally, milk contains hormones, immunoglobulins, growth factors, cytokines, nucleotides, peptides, polyamines, enzymes and other bioactive peptides and significantly contributes to the body's requirements for calcium, magnesium, selenium, riboflavin, vitamin B12 and pantothenic acid (vitamin B5) (FAO 2013; Haug, Høstmark, and Harstad 2007). Therefore, as a rich source of macro- and micronutrients, milk and fermented dairy products made from them play significant role in human nutrition in developing countries, particularly in parts of Africa where the diets of poor people often lack diversity and consumption of animal-source foods may be inadequate. Throughout the world, milk serves as key source of dietary energy, protein and fat, and contributes on average 134 kcal of energy/capita/day, 8 g of protein/capita/day, and 7.3 g of fat/capita/day, respectively (FAOSTAT 2012). However, the contribution from milk to various nutritional components varies considerably among different regions. For example, it is estimated that milk provides only 3% of dietary energy supply in Africa compared with 8%–9% in Europe and Oceania; 6%–7% of dietary protein supply in Africa compared with 19% in Europe; and 6%–8% of dietary fat supply in Africa, compared with 11%–14% in Europe, Oceania, and the Americas (FAOSTAT 2012).

The nutrient composition of fermented dairy products is largely dependent on the composition of raw milk used, which in turn is affected by several factors including stage of lactation, breed differences, number of calvings (parity), seasonal variations, age and health of animal, feed and management effects including frequency of milking per day and herd size (Bansal et al. 2003; Jenkins and McGuire 2006; Laben 1963; Walker, Dunshea, and Doyle 2004). Due to the number of different factors that affect milk composition, there are difficulties associated with using the available data from literature to draw meaningful conclusions about the nutrient composition of milk of different species because few studies provide detailed information on these factors. Additionally, the multiplicity and variability of analytical methods and study designs can also lead to differences in




Figure 3. Nutritional composition of cow, camel and goat milk (g/100 g of milk). Source: FAO (2013).

the reports on milk composition. An overview of the proximate compositions of cow, camel and goat milks, which serve as common sources of raw the main materials for the production of various AFDPs is provided in Figure 3.

Generally, milk is considered an important source of protein in the human diet, supplying an average of about 3.3 g, 3.1 g and 3.4 g of proteins per 100 g of cow, camel and goat milk respectively (Figure 3.). Milk protein fractions, both soluble/whey protein and casein proteins, are classified as high-quality proteins in view of human amino acid requirements, digestibility, bioavailability and the essential amino acid score and protein-digestibility corrected amino acid score (Boye, Wijesinha-Bettoni, and Burlingame 2012; Schaafsma 2000). Sulieman, Ilayan, and El Faki (2006) reported the protein content of camel milk for processing *gariss* in Sudan to be about 3.3–4.7%. Similarly, total protein content of fresh cow milk for production of rob was reported to be 3.4–3.8% with an average of  $3.5 \pm 0.14\%$ , but this value increased to an average of  $4.04 \pm 0.2\%$  total protein of rob. *Omashikwa* was also reported to have total protein content of about 3.3% (Bille, Buys, and Taylor 2007), while *nunu* was reported to have protein content of 3.2 mg/mL (Nebedum and Obiakor 2007).

During milk fermentation, the proteolytic activity and metabolic activity of LAB can lead to changes in the nutritive characteristics of milk in a manner that is beneficial to the consumer. For example, proteolytic capacities of some LAB during fermentation of milk can result in the release of free amino acids (Pessione and Cirrincione 2016). Muradyan, Erzhyunyan, and Sapondzhyan (1986) reported that fermentation of milk by thermophilic lactic streptococci or acidophilic rods enriched the final products with at least 4 amino acids (cysteine, valine, proline, and arginine). Akabanda et al. (2013) also showed that *nunu* fermented with LAB starter culture can serve as a good source of essential amino acids required in human metabolism.

Milk fat is made up of many different saturated and unsaturated fatty acids, with the actual amounts and composition depending on factors such as animal origin, stage of lactation, ruminal fermentation, or feed related factors. On average, saturated fatty acids (SFAs) constitute about 70% of milk fat fraction while the remaining 30% constitute unsaturated fatty acids (Pereira 2014). Of the many different saturated fatty acids that can be found in milk, only lauric, myristic and

palmitic have the property of raising blood cholesterol. Moreover, milk and fermented dairy products contain other essential nutrients, namely calcium, linoleic acid, conjugated linoleic acid (CLA), antioxidants, and probiotic bacteria that have hypocholesterolemic and protective effects (Gurr 1992; Rogeli 2000). Generally, milk and fermented dairy products are a rich source of micronutrients particularly calcium, as well as other elements such as phosphorus, magnesium, zinc, and selenium (Gaucheron 2011). Milk vitamin fraction includes fat soluble vitamins (A, D, E and K) and also the water-soluble vitamins (B complex vitamins) and vitamin C (raw milk) (Gaucheron 2011; Haug, Høstmark, and Harstad 2007). A typical nutritional composition of low fat milk and low fat plain yoghurt (Table 2) shows that the fermentation process contributes to an increase in the variety and concentration of key nutrients, especially the vitamins, minerals and essential amino acids.

### Fermented dairy products as source of reduced/lactose-free product

Lactose, the main carbohydrate present in milk, is a disaccharide made up of glucose and galactose. Lactose in milk is

found in two isomeric forms, alpha ( $\alpha$ ) and beta ( $\beta$ ). In the small intestine mucosa membrane, lactose is hydrolyzed by the enzyme lactase (also known as  $\beta$ -galactosidase) into two simple sugars glucose and galactose which are absorbed and transported to the liver through the portal vein where galactose is converted to glucose.  $\beta$ -galactosidase activity decreases significantly after weaning in mammals and in some cases may result in deficiency or complete absence of the enzyme lactase in certain individuals. An estimated 70% of adults worldwide are lactose intolerant and develop complications and ill health such as abdominal cramps/pain, bloating, flatulence, meteorism, constipation, and diarrhea upon consuming lactose (Lomer, Parkes, and Sanderson 2008; Schaafsma 2000; Shibby and Mishra 2013). However, fermented dairy products are well tolerated by lactose-intolerant individuals because some of the lactose originally in the milk is converted into lactic acid during fermentation. Additionally, yoghurt cultures are able to release  $\beta$ -galactosidase *in vivo* during digestion in lactose intolerant individuals (Savaiano 2014). The bacterial  $\beta$ -galactosidases are able to survive the acidic conditions of the stomach, as they are physically protected within the bacterial cells and facilitated by the buffering capacity of yogurt (Marco et al. 2017).

**Table 2.** Typical nutritional composition of low fat milk and yoghurt.

Nutrient	Unit	Low fat milk	Low fat plain yogurt
<b>Proximates</b>			
Energy	kCal (kJ)	42.00 (177.00)	63.00 (265.00)
Protein	g	3.37	5.25
Ash	g	0.75	1.09
Carbohydrate, by difference	g	4.99	7.04
<b>Minerals</b>			
Calcium, Ca	mg	125.00	183.00
Iron, Fe	mg	0.03	0.08
Magnesium, Mg	mg	11.00	17.00
Phosphorus, P	mg	95.00	144.00
Potassium, K	mg	150.00	234.00
Sodium, Na	mg	44.00	70.00
Zinc, Zn	mg	0.42	0.89
Copper, Cu	mg	0.01	0.01
Manganese, Mn	mg	0.00	0.00
Selenium, Se	$\mu$ g	3.30	3.30
Fluoride, F	$\mu$ g	2.60	12.00
<b>Vitamins</b>			
Vitamin C, total ascorbic acid	mg	0.00	0.80
Thiamin	mg	0.02	0.04
Riboflavin	mg	0.19	0.21
Niacin	mg	0.09	0.11
Pantothenic acid	mg	0.36	0.59
Vitamin B-6	mg	0.04	0.05
Folate, total	$\mu$ g	5.00	11.00
Choline, total	mg	17.70	15.20
Betaine	mg	0.60	0.90
Vitamin B-12	$\mu$ g	0.47	0.56
Vitamin A, IU	IU	47.00	51.00
Vitamin E (alpha-tocopherol)	mg	0.01	0.03
Vitamin D	IU	1.00	1.00
Vitamin K (phylloquinone)	$\mu$ g	0.10	0.20
<b>Essential amino acids</b>			
Histidine	g	0.10	0.13
Isoleucine	g	0.17	0.29
Leucine	g	0.32	0.53
Lysine	g	0.28	0.47
Methionine	g	0.09	0.16
Phenylalanine	g	0.17	0.29
Threonine	g	0.14	0.22
Tryptophan	g	0.04	0.03
Valine	g	0.22	0.43

Source: USDA (2018).

### Fermented dairy products in control of antibiotic use

Fermented dairy product have an extended shelf life compared to milk samples due to the *in situ* production of organic acids, as well as bacteriocins (Silva, Silva, and Ribeiro 2018). Some of these bacteriocins (such as thuricin CD) have been shown to possess narrow spectrum of antibiotics activity (especially on *Clostridium difficile*) and is less harsh in altering the balance of mice gut microbiota, as compared with conventional antibiotics such as metronidazole (Cotter, Ross, and Hill 2013). Fermented dairy products could therefore have applications as a viable option to conventional antibiotics, or be used to complement antibiotic courses, thereby helping to reduce the challenge of antibiotic use that is currently prevalent in most African countries and is exacerbating the global problem of antibiotic resistance (Tadesse et al. 2017).

### Fermented dairy products as functional foods

Staying healthy is one of the main reasons that influence the choice of food by consumers (Shibby and Mishra 2013). Beyond meeting basic nutritional requirements, diet may influence many physiological functions and can play detrimental or helpful roles in some diseases (Granato et al. 2010). Therefore, consumers have become very particular about the kinds of food they consume. Conventionally, healthy diet has been linked with foods that are rich in a variety of essential nutrients as well as bioactive components that may have a positive effect on the well-being of the consumer. These class of foods are known as functional foods and these are foods or nutrients whose ingestion leads to important physiological changes in the body that are separate and distinct from those associated with their role as nutrients (Venugopalan, Shriner, and Wong-Beringer 2010). Foods such as conventional foods,

modified foods (fortified, enriched, or enhanced), medical foods and foods for special dietary use may all be classified as functional foods (ADA 2009). Sub-classes of fermented dairy-derived functional foods that are discussed in this review are bioactive peptides, and probiotics.

### Bioactive peptides in fermented dairy products

The traditional and contemporary role of milk as the main nutritional subsistence for infants has gone beyond that horizon (Park and Nam 2015). Milk and dairy products have been the main source of nutrients for most mammals and for the growth of children and nourishment of human adults (Park 2009). Additionally, milk possesses a broad range of biologically active compounds that protect neonates and adults from pathogens and illness (Park and Nam 2015). In infants, milk peptides increase the Lactotrope development in pituitary of suckling and Prolactin increases lymphocyte trafficking and immune development. The high levels of calcium play in milk an important role in the development, strength, and density of bones in children and in the prevention of osteoporosis in elderly people. Moreover, other milk components such as phosphopeptides, lactoferricin, casokinins and casoplatelins have are mineral binding, antimicrobial, ACE-inhibitory and antithrombotic activities respectively (Gobbetti et al. 2002; Muro Urista et al. 2011). Bioactive substances are natural constituents in foods that provide health benefits beyond the basic nutritional value of the product (Marsanasco et al. 2015). Some of these bioactive compounds include antibacterial lactoferrin, casomorphins, peptides (including phosphopeptides, immunopeptides), oligosaccharides, antimicrobial proteins, immunoglobulins and lipids, amongst others (Park 2009). The proteinogenic bioactive compounds are readily available in milk, but others are latent and need to be released by the action of proteolytic enzymes (Smacchi and Gobbetti 2000). Proteolytically-derived bioactive

peptides (BP) are have a positive influence on physiological and metabolic functions of the body and may have an ultimate beneficial effect on consumers' health (Gobbetti, Minervini, and Rizzello 2007; Kitts and Weiler 2003; Korhonen and Pihlanto 2007; Park and Nam 2015). Milk is one of the most rich sources of these peptides (Korhonen and Pihlanto 2006). Park and Nam (2015) have reported the discovery of a large number of bioactive peptides and biologically and physiologically active compounds from milk casein, whey proteins and other components of milk. These peptides have been shown to have health-promoting properties including antimicrobial, hypocholesterolemic, opioid agonist and antagonists, angiotensin-converting enzyme inhibitory, antithrombotic, immunomodulations, cytomodulation, and antioxidant activity (Hayes et al. 2007; Pessione and Cirrincione 2016; Clare, Catignani, and Swaisgood 2003; Fitzgerald and Meisel 2003; Li et al. 2004). These bioactive compounds are produced in large quantities by the lactic acid bacteria (LAB), the main fermenting microorganisms in AFDP, thereby releasing these bioactive compounds into the fermented products (Pessione and Cirrincione 2016) and enhancing their functional properties. Functional food products that contain these bioactive peptides and milk components are important for the development and improvement of systemic functions, as well as prevention of diet related chronic diseases in both healthy individuals and vulnerable consumer groups (young, old, pregnant and immune deficient individuals). Some of these bioactive peptide sequences obtained via fermentation of dairy products are given in Table 3.

### Fermented dairy products as source of probiotics

Probiotics are live microorganisms which when administered in adequate amounts confer a health benefit on the host (Fijan 2014). Probiotic bacteria have dominated most food

**Table 3.** Some bioactive peptides obtained from fermented dairy products.

Food product	Characteristics (e.g. peptide sequence)	Microorganism used for fermentation	Bioactivity and health impact/application	References
Fermented milk	Val-Pro-Pro; Ile-Pro-Pro	<i>Lactobacillus helveticus</i> LBK-16H	Inhibition of angiotensin converting enzyme (ACE)/ Antihypertensive or blood pressure lowering effects	Nakamura et al. 1995
Fermented milk	Unidentified hydrolysates (possibly Val-Pro-Pro; and Ile-Pro-Pro)	<i>Lactobacillus helveticus</i> LBK-16H	Blood pressure lowering effects	Seppo et al. 2003 <sup>a</sup>
Fermented milk (yogurt-like product)	Tyr-Pro; Lys-Val-Leu-Pro-Val-Pro-Gln	<i>Lactobacillus helveticus</i> CPN4	Antihypertensive activities	Yamamoto, Maeno, and Takano 1999
Sour milk (Calpis)	Unidentified hydrolysates	<i>Lactobacillus helveticus</i> and <i>Saccharomyces cerevisiae</i>	Blood pressure lowering effects	Hata et al. 1996 <sup>a</sup>
Fermented milk	Ala-Arg-His-Pro-His-Pro-His-Leu-Ser-Phe-Met	<i>Lactobacillus delbrueckii</i> subsp. <i>bulgaricus</i> IFO13953	Antioxidant activities	Kudoh et al. 2001
Curd ( <i>Dadhi</i> )	Unidentified hydrolysates ranging from di to octapeptides.	Mixed starters including <i>Lactobacillus bulgaricus</i> and <i>S. thermophilus</i> .	Antihypertensive activities	Dabarera, Athiththan, and Perera 2015
Yoghurts from buffalo, goat and cow milks	Unidentified hydrolysates	Commercial yoghurt samples, fermentation microorganisms were unidentified	Antioxidant and antimicrobial activities	Rahmawati and Suntornsuk 2016
Buffalo milk yoghurt	Hydrolysate containing peptides such as Phe-Val-Ala-Pro-Phe-Pro-Glu; Leu-Val-Tyr-Pro-Phe-Pro-Gly-Pro-Ile-Pro-Lys; Leu-Val-Tyr-Pro-Phe-Pro-Gly-Pro-Ile-Pro-Lys-Ser-Leu-Pro-Gln-Asn; and Leu-Tyr-Gln-Glu-Pro-Val-Leu-Gly-Pro-Val-Arg-Gly-Pro-Phe-Pro-Ile-Ile-Val	Yoghurt starters, together with <i>Lactobacillus acidophilus</i> 20552 ATCC, or <i>Lactobacillus helveticus</i> CH 5	ACE-inhibitory, antioxidant, and antibacterial activities	Taha et al. 2017

<sup>a</sup>Study involved human subjects.

products including yoghurts and fermented milks for decades due to their perceived health benefits (Mattila-Sandholm et al. 2002). Research indicate that health benefits from both commercial and household African fermented milk such as *mursik* of Kenya are attributed to probiotics used as starter culture (Digo et al. 2017). Probiotic strains such as *Lactobacillus* spp. have also been isolated from non-pasteurized Ethiopia *ergo* (Bereda et al. 2014) and Tunisian *leben* (Samet-Bali 2012).

The use of probiotic bacterial cultures helps stimulate the growth of suitable microorganism, crowds out potentially harmful bacteria, partly due to their ability to produce bacteriocins (Silva, Silva, and Ribeiro 2018), reinforces the body's natural defense mechanism, and improve gut health (Salminen et al. 1999) and also induce antihypertensive and inflammation pain relief effects (Islam 2016; Steinkraus 1994). Moreover, probiotics have been shown in several clinical trials to control radiotherapy-induced diarrhea (Liu et al. 2017), reduce significantly serum cholesterol (Wang et al. 2018), and reduce the levels of plasma ammonia in patients with hepatic encephalopathy (McGee et al. 2011). It is worth noting that most African countries are in the epidemiological transition stage involving an experience of the double burden of both chronic (life-style related non-communicable) and infectious (communicable) diseases (Agyei-Mensah and de-Graft Aikins 2010; Kuate Defo 2014). With proper research and development, probiotics from indigenous fermented dairy products could be used in dietary interventions to help mitigate some of the diseases encountered in this public health challenge.

The mechanisms by which probiotic bacteria exert their beneficial effects is still not completely understood. However, commonly suggested mechanisms underlying the health-promoting effects of probiotics may involve modification of the gut, antagonizing or exclusion of pathogens through production of antimicrobial compounds, competing for pathogen binding and receptor sites and for available nutrients and growth factors, stimulating immunomodulatory cells and producing lactase (Parvez et al. 2006; Franz et al. 2014). Similarly, Stanton et al. (2005), proposed that the mechanisms of action by probiotics are via competitive exclusion, competition for nutrients and/or stimulation of an immune response.

## Future outlook and conclusion

According to the World Bank, the consumption of animal-based foods in Africa is increasing; and consumption of milk and its products is projected to grow by 2.3% per year, reaching to 50.2 million tonnes of milk by 2050 (World Bank 2014). This projected growth is good news for the continent since it comes with implications in public health, nutrition, food security and economic growth. However, to harness the full potential of this projection, a number of challenges and unknowns (particularly in the area of production, analysis, and health impact assessment) have to be overcome and deciphered. Firstly, there is the need for investment in applied research aimed at standardizing the

production methods of the many fermented dairy products in Africa. The variable quality of most African fermented dairy products resulting from lack of standardization in raw milk quality, lack of the use of defined starter cultures for controlled fermentation processes and microbial contaminations along the processing chain (Aworh 2008; Haaland 2007) need urgent attention. Thus, techniques for assuring the traceability, safety and overall quality of AFDPs need to be developed and implemented. Secondly, analysis techniques for characterizing (i.e. fingerprinting) flavor and sensorial metabolites are needed to profile and standardize the quality characteristics of these of these fermented foods. In this regard, proteomics, metabolomics and genomic techniques as well as multivariate approach to analyze all these data, will be indispensable (El Sheikha and Hu 2018). Thirdly, there is the need for *in vivo* studies and human clinical trials in each region or country to confirm and establish the health benefits and the consequent public health impact following the consumption of the indigenous fermented dairy products. Such *in vivo* health impact assessment is important because the literature is scarce on clinical trials involving indigenous fermented dairy products from African. In conclusion, it is an established fact that fermented dairy products are a storehouse of nutrient, probiotics and bioactive compounds that have a huge impact on human health. In Africa, efforts aimed at harnessing the food and health potential of these fermented products could help control some challenges such as deficiency of essential nutrients (proteins, vitamins, minerals), food security, double burden of chronic and infectious diseases facing many countries in the region.

## Conflict of interest

The authors declare no conflict of interest.

## ORCID

Dominic Agyei  <http://orcid.org/0000-0003-2280-4096>  
James Owusu-Kwarteng  <http://orcid.org/0000-0002-2703-3393>  
Fortune Akabanda  <http://orcid.org/0000-0002-3118-7619>  
Samuel Akomea-Frempong  <http://orcid.org/0000-0001-5729-8985>

## References

- Abbey, L. D., M. Hodari-Okoe, and A. Osei-Yaw. 1994. *Studies on traditional processing and quality of fermented fish momone*, edited by FRI. Accra: Food Research Institute, Accra-Ghana.
- Abdelgadir, W. S., T. K. Ahmed, and H. A. Dirar. 1998. The traditional fermented milk products of the Sudan. *International Journal of Food Microbiology* 44 (1–2):1–13. doi: [10.1016/S0168-1605\(98\)00090-7](https://doi.org/10.1016/S0168-1605(98)00090-7).
- Abdelgadir, W. S., S. H. Hamad, P. L. Møller, and M. Jakobsen. 2001. Characterisation of the dominant microbiota of Sudanese fermented milk Rob. *International Dairy Journal* 11 (1–2):63–70. doi: [10.1016/S0958-6946\(01\)00042-5](https://doi.org/10.1016/S0958-6946(01)00042-5).
- Abdelgadir, W., D. S. Nielsen, S. Hamad, and M. Jakobsen. 2008. A traditional Sudanese fermented camel's milk product, Gariss, as a habitat of *Streptococcus infantarius* subsp. *infantarius*. *International Journal of Food Microbiology* 127 (3):215–219. doi: [10.1016/j.ijfoodmicro.2008.07.008](https://doi.org/10.1016/j.ijfoodmicro.2008.07.008).



- Abriouel, H., N. B. Omar, R. L. López, M. Martínez-Cañamero, S. Keleke, and A. Gálvez. 2006. Culture-independent analysis of the microbial composition of the African traditional fermented foods *poto poto* and *dégue* by using three different DNA extraction methods. *International Journal of Food Microbiology* 111 (3):228–233. doi: [10.1016/j.ijfoodmicro.2006.06.006](https://doi.org/10.1016/j.ijfoodmicro.2006.06.006).
- ADA. 2009. Position of the American Dietetic Association: Functional foods. *Journal of the American Dietetic Association* 109 (4):735–746. doi: [10.1016/j.jada.2009.02.023](https://doi.org/10.1016/j.jada.2009.02.023).
- Agyei-Mensah, S., and A. de-Graft Aikins. 2010. Epidemiological transition and the double burden of disease in Accra, Ghana. *Journal of Urban Health* 87 (5):879–897. doi: [10.1007/s11524-010-9492-y](https://doi.org/10.1007/s11524-010-9492-y).
- Akabanda, F., J. Owusu-Kwarteng, K. Tano-Debrah, R. L. K. Glover, D. S. Nielsen, and L. Jespersen. 2013. Taxonomic and molecular characterization of lactic acid bacteria and yeasts in *nunu*, a Ghanaian fermented milk product. *Food Microbiology* 34 (2): 277–283. doi: [10.1016/j.fm.2012.09.025](https://doi.org/10.1016/j.fm.2012.09.025).
- Akabanda, F., J. Owusu-Kwarteng, K. Tano-Debrah, C. Parkouda, and L. Jespersen. 2014. The use of lactic acid bacteria starter culture in the production of *Nunu*, a spontaneously fermented milk product in Ghana. *International Journal of Food Science* 2014:1. doi: [10.1155/2014/721067](https://doi.org/10.1155/2014/721067).
- Andualem, B., and T. Geremew. 2014. Fermented Ethiopian dairy products and their common useful microorganisms: A review. *World Journal of Agricultural Sciences* 10:121–133.
- Asogwa, I. S., J. I. Okoye, and K. Oni. 2017. Promotion of indigenous food preservation and processing knowledge and the challenge of food security in Africa. *Journal of Food Security* 5:75–87.
- Aworh, O. C. 2008. The role of traditional food processing technologies in national development: The West African experience. <https://www.iufost.org/publications/books/documents/Revd.pdf>.
- Bansal, B. K., S. S. Randhawa, K. B. Singh, and P. K. Boro. 2003. Effect of specific, nonspecific and latent mastitis on milk composition of dairy cows. *Indian Journal of Animal Sciences (India)*.
- Barber, L., S. C. Achinewhu, and E. A. Ibiama. 1988. The microbiology of *ogiri* production from castor seed (*Ricinus communis*). *Food Microbiology* 5 (4):177–183. doi: [10.1016/0740-0020\(88\)90016-0](https://doi.org/10.1016/0740-0020(88)90016-0).
- Benkerroum, N. 2013. Traditional fermented foods of North African countries: Technology and food safety challenges with regard to microbiological risks. *Comprehensive Reviews in Food Science and Food Safety* 12 (1):54–89. doi: [10.1111/j.1541-4337.2012.00215.x](https://doi.org/10.1111/j.1541-4337.2012.00215.x).
- Bensalah, F., C. Delorme, and P. Renault. 2009. Characterisation of thermotolerant cocci from indigenous flora of 'leben' in algerian arid area and DNA identification of atypical *Lactococcus lactis* strains. *Current Microbiology* 59 (2):139–146. doi: [10.1007/s00284-009-9411-1](https://doi.org/10.1007/s00284-009-9411-1).
- Bereda, A., M. Eshetu, and Z. Yilma. 2014. Microbial properties of Ethiopian dairy products: A review. *African Journal of Microbiology Research* 8 (23):264–2271. doi: [10.5897/AJMR2014.6733](https://doi.org/10.5897/AJMR2014.6733).
- Beukes, E. M., B. H. Bester, and J. F. Mostert. 2001. The microbiology of South African traditional fermented milks. *International Journal of Food Microbiology* 63 (3):189–197. doi: [10.1016/S0168-1605\(00\)00417-7](https://doi.org/10.1016/S0168-1605(00)00417-7).
- Bille, P. G., E. Buys, and J. Taylor. 2007. The technology and properties of Omashikwa, a traditional fermented buttermilk produced by small-holder milk producers in Namibia. *International Journal of Food Science & Technology* 42 (5):620–624. doi: [10.1111/j.1365-2621.2006.01381.x](https://doi.org/10.1111/j.1365-2621.2006.01381.x).
- Boye, J., R. Wijesinha-Bettoni, and B. Burlingame. 2012. Protein quality evaluation twenty years after the introduction of the protein digestibility corrected amino acid score method. *British Journal of Nutrition* 108 (S2):S183–S211. doi: [10.1017/S0007114512002309](https://doi.org/10.1017/S0007114512002309).
- Caplice, E., and G. F. Fitzgerald. 1999. Food fermentations: Role of microorganisms in food production and preservation. *International Journal of Food Microbiology* 50 (1–2):131–149. doi: [10.1016/S0168-1605\(99\)00082-3](https://doi.org/10.1016/S0168-1605(99)00082-3).
- Chelule, P. K., H. P. Mbongwa, S. Carries, and N. Gqaleni. 2010. Lactic acid fermentation improves the quality of amahewu, a traditional South African maize-based porridge. *Food Chemistry* 122 (3): 656–661. doi: [10.1016/j.foodchem.2010.03.026](https://doi.org/10.1016/j.foodchem.2010.03.026).
- Chow, J., H. Tang, and S. K. Mazmanian. 2011. Pathobionts of the gastrointestinal microbiota and inflammatory disease. *Current Opinion in Immunology* 23 (4):473–480.
- Clare, D., G. Catignani, and H. Swaisgood. 2003. Biodefense properties of milk: The role of antimicrobial proteins and peptides. *Current Pharmaceutical Design* 9 (16):1239–1255. doi: [10.2174/1381612033454874](https://doi.org/10.2174/1381612033454874).
- Coppock, D. L., S. J. Holden, and C. B. O'Connor. 1991. *Milk processing and peri-urban dairy marketing in semi-arid Ethiopia and prospects for development intervention*. Addis Ababa, Ethiopia: ILCA.
- Cotter, P. D., R. P. Ross, and C. Hill. 2013. Bacteriocins – A viable alternative to antibiotics? *Nature Reviews Microbiology* 11 (2): 95–105. doi: [10.1038/nrmicro2937](https://doi.org/10.1038/nrmicro2937).
- Dabarera, M. C., L. V. Athiththan, and R. P. Perera. 2015. Antihypertensive peptides from curd. *AYU (An International Quarterly Journal of Research in Ayurveda)* 36 (2):214–219. doi: [10.4103/0974-8520.175534](https://doi.org/10.4103/0974-8520.175534).
- Deak, T., and L. R. Beuchat. 1996. Yeasts in specific types of foods. In *Handbook of food spoilage yeasts*, 61–95. Boca Raton, FL: CRC Press.
- Digo, C. A., E. Kamau-Mbuthia, J. W. Matofari, and W. K. Ng'etich. 2017. Potential probiotics from traditional fermented milk, Mursik of Kenya. *International Journal of Nutrition and Metabolism* 10 (9): 75–81. doi: [10.5897/ijnam2016.0203](https://doi.org/10.5897/ijnam2016.0203).
- Dirar, H. A. 1993. *The indigenous fermented foods of the Sudan: A study in African food and nutrition*. Wallingford, UK: CAB international.
- Dossou-Yovo, P. 2007. Survey of the improvement of fish fermentation for *lanhouin* production in Benin. *African Journal of Food Science* 5 (17). doi: [10.5897/ajfs111.136](https://doi.org/10.5897/ajfs111.136).
- Doutoum, A. A., A. Tidjani, M. Balde, S. M. T. Tidjani, A. Abdelaziz, I. K. S. B. Sylla, M. G. Seydi, and B. S. Toguebaye. 2013. Identification of lactic bacteria of milk quail in Chad. *Journal of Applied Biology & Biotechnology* 1:7–15.
- EFSA. 2016. Update of the list of QPS-recommended biological agents intentionally added to food or feed as notified to EFSA 4: Suitability of taxonomic units notified to EFSA until March 2016. *EFSA Journal* 15 (7):4884.
- El Sheikh, A. F., and D.-M. Hu. 2018. Molecular techniques reveal more secrets of fermented foods. *Critical Reviews in Food Science and Nutrition*. doi: [10.1080/10408398.2018.1506906](https://doi.org/10.1080/10408398.2018.1506906).
- El-Baradei, G., A. Delacroix-Buchet, and J. C. Ogier. 2008. Bacterial biodiversity of traditional Zabady fermented milk. *International Journal of Food Microbiology* 121 (3):295–301. doi: [10.1016/j.ijfoodmicro.2007.11.014](https://doi.org/10.1016/j.ijfoodmicro.2007.11.014).
- Ezeji, C., and P. C. Ojimelekwu. 1993. Effect of fermentation on the nutritional quality and functional properties of infant food formulations prepared from bambarra-groundnut, fluted-pumpkin and millet seeds. *Plant Foods for Human Nutrition* 44 (3):267–276. doi: [10.1007/BF01088322](https://doi.org/10.1007/BF01088322).
- FAO. 1990. The technology of traditional milk products in developing countries. <http://www.fao.org/docrep/003/t0251e/T0251E00.htm>.
- FAO. 2006. FAO Global Perspective Studies Unit. Background data of the world agriculture: Towards 2030–2050—Interim Report. Rome.
- FAO. 2013. Milk and dairy products in human nutrition. In *Food and agriculture organization of the United Nations*, edited by A. Bennett, E. Muehlhoff, and D. McMahon. Rome: FAO.
- FAO. 2018. Gateway to dairy production and products. <http://www.fao.org/dairy-production-products/production/en/>.
- FAOSTAT. 2012. FAO statistical database. <http://faostat.fao.org/>.
- Feresu, S. B., and M. I. Muzondo. 1990. Identification of some lactic acid bacteria from two Zimbabwean fermented milk products. *World Journal of Microbiology & Biotechnology* 6 (2):178–186. doi: [10.1007/bf01200939](https://doi.org/10.1007/bf01200939).
- Fijan, S. 2014. Microorganisms with claimed probiotic properties: An overview of recent literature. *International Journal of Environmental Research and Public Health* 11 (5):4745–4767. doi: [10.3390/ijerph110504745](https://doi.org/10.3390/ijerph110504745).

- Fitzgerald, R. J., and H. Meisel. 2003. Milk protein hydrolysates and bioactive peptides. In *Advanced dairy chemistry—1 proteins*. New York, NY: Springer.
- Fleet, G. H. 1990. Yeasts in dairy products. *The Journal of Applied Bacteriology* 68 (3):199–221.
- Franz, C. M., W. H. Holzapfel, and M. E. Stiles. 1999. Enterococci at the crossroads of food safety? *International Journal of Food Microbiology* 47 (1–2):1–24.
- Franz, C. M. A. P., M. Huch, J. M. Mathara, H. Abriouel, N. Benomar, G. Reid, A. Galvez, and W. H. Holzapfel. 2014. African fermented foods and probiotics. *International Journal of Food Microbiology* 190:84–96. doi:10.1016/j.ijfoodmicro.2014.08.033.
- Fuller, R. 1989. Probiotics in man and animals. *The Journal of Applied Bacteriology* 66 (5):365–378.
- Gadaga, T. H., A. N. Mutukumira, J. A. Narvhus, and S. B. Feresu. 1999. A review of traditional fermented foods and beverages of Zimbabwe. *International Journal of Food Microbiology* 53 (1):1–11. doi:10.1016/S0168-1605(99)00154-3.
- Gaucheron, F. 2011. Milk and dairy products: A unique micronutrient combination. *Journal of the American College of Nutrition* 30 (sup5): 400S–409S. doi:10.1080/07315724.2011.10719983.
- Giraffa, G. 2002. Enterococci from foods. *FEMS Microbiology Reviews* 26(2):163–171.
- Gobbetti, M., F. Minervini, and C. G. Rizzello. 2007. Bioactive peptides in dairy products. In *Handbook of food products manufacturing*. Hoboken, NJ: John Wiley & Sons, Inc.
- Gobbetti, M., L. Stepaniak, M. De Angelis, A. Corsetti, and R. Di Cagno. 2002. Latent bioactive peptides in milk proteins: Proteolytic activation and significance in dairy processing. *Critical Reviews in Food Science and Nutrition* 42 (3):223–239. doi:10.1080/10408690290825538.
- Godfree, A. F., D. Kay, and M. D. Wyer. 1997. Faecal streptococci as indicators of faecal contamination in water. *Journal of Applied Microbiology* 83 (S1):110–119. doi:10.1046/j.1365-2672.83.s1.12.x.
- Gonfa, A., A. Fite, K. Urga, and A. G. Berhanu. 1999. Microbiological aspects of ergo (Ititu) fermentation. *SINET: Ethiopian Journal of Science* 22 (2):283–290.
- Granato, D., G. F. Branco, A. G. Cruz, J. d. A. F. Faria, and N. P. Shah. 2010. Probiotic dairy products as functional foods. *Comprehensive Reviews in Food Science and Food Safety* 9 (5): 455–470. doi:10.1111/j.1541-4337.2010.00120.x.
- Greppi, A., K. Rantsiou, W. Padonou, J. Hounhouigan, L. Jespersen, M. Jakobsen, and L. Cocolin. 2013. Determination of yeast diversity in ogi, mawè, gowé and tchoukoutou by using culture-dependent and -independent methods. *International Journal of Food Microbiology* 165 (2):84–88. doi:10.1016/j.ijfoodmicro.2013.05.005.
- Gurr, M. I. 1992. Milk products: Contribution to nutrition and health. *Journal of the Society of Dairy Technology* 45 (3):61–67.
- Haaland, R. 2007. Porridge and pot, bread and oven: Food ways and symbolism in Africa and the near East from the Neolithic to the present. *Cambridge Archaeological Journal* 17 (2):165. doi:10.1017/s0959774307000236.
- Hata, Y., M. Yamamoto, M. Ohni, K. Nakajima, Y. Nakamura, and T. Takano. 1996. A placebo-controlled study of the effect of sour milk on blood pressure in hypertensive subjects. *The American Journal of Clinical Nutrition* 64 (5):767–771. doi:10.1093/ajcn/64.5.767.
- Haug, A., A. T. Høstmark, and O. M. Harstad. 2007. Bovine milk in human nutrition – A review. *Lipids in Health and Disease* 6 (1):25.
- Hayes, M., C. Stanton, G. F. Fitzgerald, and R. P. Ross. 2007. Putting microbes to work: Dairy fermentation, cell factories and bioactive peptides. Part II: Bioactive peptide functions. *Biotechnology Journal* 2 (4):435–449. doi:10.1002/biot.200700045.
- Holzapfel, W. H. 2002. Appropriate starter culture technologies for small-scale fermentation in developing countries. *International Journal of Food Microbiology* 75 (3):197–212. doi:10.1016/S0168-1605(01)00707-3.
- Hui, Y.H., L. Meunier-Goddik, J. Josephsen, W.K. Nip, and P.S. Stanfield. 2004. *Handbook of food and beverage fermentation technology*. Boca Raton, FL: CRC Press.
- Islam, S. U. 2016. Clinical uses of probiotics. *Medicine* 95 (5):e2658. doi:10.1097/MD.0000000000002658.
- Jakobsen, M., and J. A. Narvhus. 1996. Yeasts and their possible beneficial and negative effects on the quality of dairy products. *International Dairy Journal* 6 (8–9):755–768. doi:10.1016/0958-6946(95)00071-2.
- Jans, C., J. Bugnard, P. M. K. Njage, C. Lacroix, and L. Meile. 2012b. Lactic acid bacteria diversity of African raw and fermented camel milk products reveals a highly competitive, potentially health-threatening predominant microflora. *LWT - Food Science and Technology* 47 (2):371–379. doi:10.1016/j.lwt.2012.01.034.
- Jans, C., T. deWouters, B. Bonfoh, C. Lacroix, D. W. M. Kaïndi, J. Anderegg, D. Böck, S. Vitali, T. Schmid, J. Isenring, et al. 2016. Phylogenetic, epidemiological and functional analyses of the *Streptococcus bovis*/*Streptococcus equinus* complex through an over-arching MLST scheme. *BMC Microbiology* 16:117.
- Jans, C., R. Follador, M. Hochstrasser, C. Lacroix, L. Meile, and M. J. A. Stevens. 2013. Comparative genome analysis of *Streptococcus infantarius subsp. infantarius* CJ18, an African fermented camel milk isolate with adaptations to dairy environment. *BMC Genomics* 14 (1):200. doi:10.1186/1471-2164-14-200.
- Jans, C., A. Gerber, J. Bugnard, P. M. K. Njage, C. Lacroix, and L. Meile. 2012a. Novel *Streptococcus infantarius subsp. infantarius* variants harboring lactosemetabolism genes homologous to *Streptococcus thermophilus*. *Food Microbiology* 31 (1):33–42. doi:10.1016/j.fm.2012.02.001.
- Jans, C., L. Meile, D. W. M. Kaïndi, W. Kogi-Makau, P. Lamuka, P. Renault, B. Kreikemeyer, C. Lacroix, J. Hattendorf, and J. Zinsstag. 2017. African fermented dairy products – Overview of predominant technologically important microorganisms focusing on African *Streptococcus infantarius* variants and potential future applications for enhanced food safety and security. *International Journal of Food Microbiology* 250:27–36.
- Jans, C., L. Meile, C. Lacroix, and M. J. A. Stevens. 2015. Genomics, evolution, and molecular epidemiology of the *Streptococcus bovis*/*Streptococcus equinus* complex (SBSEC). *Infection, Genetics and Evolution* 33:419–436. doi:10.1016/j.meegid.2014.09.017.
- Jaros, D., and H. Rohm. 2003. Controlling the texture of fermented dairy products: The case of yoghurt. *Dairy Processing: Maximising Quality*. doi:10.1533/9781855737075.1.155.
- Jenkins, T. C., and M. A. McGuire. 2006. Major advances in nutrition: Impact on milk composition. *Journal of Dairy Science* 89 (4): 1302–1310.
- Jespersen, L. 2003. Occurrence and taxonomic characteristics of strains of *Saccharomyces cerevisiae* predominant in African indigenous fermented foods and beverages. *FEMS Yeast Research* 3 (2):191–200.
- Jiwoua, C., and J. B. Milliere. 1990. Lactic flora and enterococci of fermented milk (Pindidam) produced in Adamaoua (Cameroon). *Lait* 70 (5–6):475–486.
- Josephsen, J., and L. Jespersen. 2004. Starter cultures and fermented products. In *Handbook of food and beverage fermentation technology*, 23–49. New York, NY: Marcel Dekker Incorporated.
- Karenzi, E., R. D. Dauphin, A. Mashaku, L. Majad, B. Munyanganizi, and P. Thonart. 2012. Fermentation of kivuguto, a Rwandese traditional milk: Selection of microbes for a starter culture. *Sciences & Technologie C* (36):9–17.
- Kassaye, T., B. K. Simpson, J. P. Smith, and C. B. O'connor. 1991. Chemical and microbiological characteristics of Ititu. *Milchwissenschaft* 46 (10):649–653.
- Kebede, A., B. C. Viljoen, T. H. Gadaga, J. A. Narvhus, and A. Lourens-Hattingh. 2007. The effect of container type on the growth of yeast and lactic acid bacteria during production of Sethemi, South African spontaneously fermented milk. *Food Research International* 40 (1):33–38. doi:10.1016/j.foodres.2006.07.012.
- Kitts, D., and K. Weiler. 2003. Bioactive proteins and peptides from food sources. Applications of bioprocesses used in isolation and recovery. *Current Pharmaceutical Design* 9 (16):1309–1323. doi:10.2174/1381612033454883. doi:10.2174/1381612033454883.
- Koffi-Nevr, R., T. S. T. Ouina, M. Koussémon, and K. Brou. 2011. Chemical composition and lactic microflora of adjuevan, a

- traditional Ivorian fermented fish condiment. *Pakistan Journal of Nutrition* 10 (4):332–337. doi:10.3923/pjn.2011.332.337.
- Korhonen, H., and A. Pihlanto. 2006. Bioactive peptides: Production and functionality. *International Dairy Journal* 16 (9):945–960. doi:10.1016/j.idairyj.2005.10.012.
- Korhonen, H., and A. Pihlanto. 2007. Bioactive peptides from food proteins. In *Handbook of food products manufacturing*. Hoboken, NJ: John Wiley & Sons, Inc.
- Kuate Defo, B. 2014. Demographic, epidemiological, and health transitions: Are they relevant to population health patterns in Africa? *Global Health Action* 7:22443. doi:10.3402/gha.v7.22443.
- Kudoh, Y., S. Matsuda, K. Igoshi, and T. Oki. 2001. Antioxidative peptide from milk fermented with *Lactobacillus delbrueckii* subsp. *bulgaricus* IFO13953. *Nippon Shokuhin Kagaku Kogaku Kaishi* 48 (1): 44–50. doi:10.3136/nskkk.48.44.
- Laben, R. C. 1963. Factors responsible for variation in milk composition. *Journal of Dairy Science* 46 (11):1293–1301. doi:10.3168/jds.S0022-0302(63)89264-4.
- Li, G.-H., G.-W. Le, Y.-H. Shi, and S. Shrestha. 2004. Angiotensin I-converting enzyme inhibitory peptides derived from food proteins and their physiological and pharmacological effects. *Nutrition Research* 24 (7):469–486. doi:10.1016/j.nutres.2003.10.014.
- Libouga, D. G., J. J. E. Ngang, and H. Halilou. 2005. Quality of some Cameroonian fermented milk. *Sciences Des Aliments* 25 (1):53–66. doi:10.3166/sda.25.53-66.
- Liu, M.-M., S.-T. Li, Y. Shu, and H.-Q. Zhan. 2017. Probiotics for prevention of radiation-induced diarrhea: A meta-analysis of randomized controlled trials. *PLoS One* 12 (6):e0178870. doi:10.1371/journal.pone.0178870.
- Lomer, M. C. E., G. C. Parkes, and J. D. Sanderson. 2008. Lactose intolerance in clinical practice—myths and realities. *Alimentary Pharmacology & Therapeutics* 27 (2):93–103. doi:10.1111/j.1365-2036.2007.03557.x.
- Lore, T. A., S. K. Mbugua, and J. Wangoh. 2005. Enumeration and identification of microflora in suusac, a Kenyan traditional fermented camel milk product. *LWT - Food Science and Technology* 38 (2):125–130. doi:10.1016/j.lwt.2004.05.008.
- Ludwig, W., K. H. Schleifer, and W. B. Whitman. 2009. Order II. Lactobacillales ord. nov. *Bergey's manual of systematic bacteriology*. vol. 3, 464. Hoboken, NJ: John Wiley & Sons, Inc.
- MacDonald, R., and C. Reitmeier. 2017. Food processing. In *Understanding food systems*, edited by Ruth MacDonald and Cheryll Reitmeier, 179–225. Cambridge, MA: Academic Press.
- Mangia, N. P., G. Garau, M. A. Murgia, A. Bennani, and P. Deiana. 2014. Influence of autochthonous lactic acid bacteria and enzymatic yeast extracts on the microbiological, biochemical and sensorial properties of Lben generic products. *Journal of Dairy Research* 81 (2):193–201. doi:10.1017/S0022029914000119.
- Marco, M. L., D. Heeney, S. Binda, C. J. Cifelli, P. D. Cotter, B. Foligné, M. Gänzle, R. Kort, G. Pasin, A. Pihlanto, et al. 2017. Health benefits of fermented foods: Microbiota and beyond. *Current Opinion in Biotechnology* 44:94–102.
- Marsanasco, M., A. L. Márquez, J. R. Wagner, N. S. Chiarmoni, and S. d. V. Alonso. 2015. Bioactive compounds as functional food ingredients: Characterization in model system and sensory evaluation in chocolate milk. *Journal of Food Engineering* 166:55–63. doi:10.1016/j.jfoodeng.2015.05.007.
- Marshall, E., and D. J. Mejía-Lorío. 2011. *Traditional fermented food and beverages for improved livelihoods*. Rural Infrastructure and Agro-Industries Division, Food and Agriculture Organization of the United Nations.
- Mathara, J. M. 1999. Studies on lactic acid producing microflora in mursik and kule naoto, traditional fermented milks from Nandi and Masai communities in Kenya. University of Nairobi.
- Mathara, J. M., U. Schillinger, P. M. Kutima, S. K. Mbugua, and W. H. Holzapfel. 2004. Isolation, identification and characterisation of the dominant microorganisms of kule naoto: The Maasai traditional fermented milk in Kenya. *International Journal of Food Microbiology* 94 (3):269–278.
- Mattila-Sandholm, T., P. Myllärinen, R. Crittenden, G. Mogensen, R. Fondén, and M. Saarela. 2002. Technological challenges for future probiotic foods. *International Dairy Journal* 12 (2–3):173–182. doi:10.1016/S0958-6946(01)00099-1.
- Mbawala, A., P. Y. Mahbou, H. T. Mouafo, and L. N. Tatsadjieu. 2013. Antibacterial activity of some lactic acid bacteria isolated from a local fermented milk product (pendidam) in ngaoundere, Cameroon. *Journal of Animal and Plant Sciences* 23 (1):157–166.
- McGee, R. G., A. Bakens, K. Wiley, S. M. Riordan, and A. C. Webster. 2011. Probiotics for patients with hepatic encephalopathy. *Cochrane Database of Systematic Reviews* (11):CD008716. doi:10.1002/14651858.CD008716.pub2.
- Mensah, P. 1997. Fermentation—The key to food safety assurance in Africa? *Food Control* 8 (5–6):271–278. doi:10.1016/S0956-7135(97)00020-0.
- Mugula, J. K., S. A. M. Nnko, J. A. Narvhus, and T. Sørhaug. 2003. Microbiological and fermentation characteristics of togwa, a Tanzanian fermented food. *International Journal of Food Microbiology* 80 (3):187–199. doi:10.1016/S0168-1605(02)00141-1.
- Muradyan, E. A., L. A. Erzhynskyan, and M. S. Saponzhyan. 1986. Composition of free amino acids in fermented milk products. *Biologicheskii Zhurnal Armenii* 29:111–112.
- Muro Urista, C., R. Álvarez Fernández, F. Riera Rodríguez, A. Arana Cuenca, and A. Téllez Jurado. 2011. Review: Production and functionality of active peptides from milk. *Food Science and Technology International* 17 (4):293–317. doi:10.1177/1082013211398801.
- Nakamura, Y., N. Yamamoto, K. Sakai, and T. Takano. 1995. Antihypertensive effect of sour milk and peptides isolated from it that are inhibitors to angiotensin I-converting enzyme. *Journal of Dairy Science* 78 (6):1253–1257. doi:10.3168/jds.S0022-0302(95)76745-5.
- Nakavuma, J. L., P. L. Møller, M. Jakobsen, P. Salimo, and G. W. Nasinyama. 2012. Processing steps and lactic acid bacteria involved in traditional cultured milk (Kwerionik) production in Uganda. *Africa Journal of Animal and Biomedical Sciences* 7 (2):82–93.
- Ndambi, O. A., T. Hemme, and U. Latacz-Lohmann. 2007. Dairying in Africa – Status and recent developments. *Livestock Research for Rural Development* 19 (8).
- Nduko, J. M., J. W. Matofari, Z. O. Nandi, and M. B. Sichangi. 2017. Spontaneously fermented Kenyan milk products: A review of the current state and future perspectives. *African Journal of Food Science* 11 (1):1–11. doi:10.5897/AJFS2016.1516.
- Nebedum, J. O., and T. Obiakor. 2007. The effects of different preservation methods on the quality of nunu, a locally fermented Nigerian dairy product. *African Journal of Biotechnology* 6 (4).
- Nieminen, M. T., L. Novak-Frazer, R. Collins, S. P. Dawsey, S. M. Dawsey, C. C. Abnet, R. E. White, N. D. Freedman, M. Mwachi, P. Bowyer, et al. 2013. Alcohol and acetaldehyde in African fermented milk mursik—A possible etiologic factor for high incidence of esophageal cancer in Western Kenya. *Cancer Epidemiology and Prevention Biomarkers* 22 (1):69–75. doi:10.1158/1055-9965.EPI-12-0908.
- Njage, P. M. K., S. Dolci, C. Jans, J. Wangoh, C. Lacroix, and L. Meile. 2011. Characterization of yeasts associated with camel milk using phenotypic and molecular identification techniques. *Research Journal of Microbiology* 6 (9):678. doi:10.3923/jm.2011.678.692.
- Nyambane, B., W. M. Thari, J. Wangoh, and P. M. K. Njage. 2014. Lactic acid bacteria and yeasts involved in the fermentation of amabere amaruranu, a Kenyan fermented milk. *Food Science & Nutrition* 2 (6):692–699. doi:10.1002/fsn3.162.
- Obodai, M., and C. E. R. Dodd. 2006. Characterization of dominant microbiota of a Ghanaian fermented milk product, nyarmie, by culture-and nonculture-based methods. *Journal of Applied Microbiology* 100 (6):1355–1363. doi:10.1111/j.1365-2672.2006.02895.x.
- Ogbulie, T. E., C. A. Nsofor, and F. C. Nze. 2014. Bacteria species associated with Ugba (Pentaclethra macrophylla) produced traditionally and in the laboratory and the effect of fermentation on product of oligosaccharide hydrolysis. *Nigerian Food Journal* 32 (2):73–80. doi:10.1016/S0189-7241(15)30120-x.



- Ohenhen, R. E., E. P. K. Imarenezor, and A. N. Kihuha. 2013. Microbiome of madila – A Southern-African fermented milk product. *International Journal of Basic and Applied Sciences* 2 (2): 170–175. doi:10.14419/ijbas.v2i2.639.
- Ohiokpehai, O. 2003. Processed food products and nutrient composition of goat milk. *Pakistan Journal of Nutrition* 2 (2):68–71. doi:10.3923/pjn.2003.68.71.
- Ørskov, E. R. 1995. A traveller's view of Outer Mongolia. *Outlook on Agriculture* 24 (2):127–129. doi:10.1177/003072709502400211.
- Osvik, R. D., S. Sperstad, E. M. Breines, J. Godfroid, Z. Zhou, P. Ren, C. Geoghegan, W. Holzapfel, and E. Ringø. 2013. Bacterial diversity of aMasi, a South African fermented milk product, determined by clone library and denaturing gradient gel electrophoresis analysis. *African Journal of Microbiology Research* 7 (32):4146–4158.
- Ouadghiri, M., M. Vancanneyt, P. Vandamme, S. Naser, D. Gevers, K. Lefebvre, J. Swings, and M. Amar. 2009. Identification of lactic acid bacteria in Moroccan raw milk and traditionally fermented skimmed milk 'Iben'. *Journal of Applied Microbiology* 106 (2):486–495. doi:10.1111/j.1365-2672.2008.04016.x.
- Owusu-Kwarteng, J., F. Akabanda, P. Johansen, L. Jespersen, and D. S. Nielsen. 2017. Chapter 15 - Nunu, A West African Fermented Yogurt-Like Milk Product. In *Yogurt in Health and Disease Prevention*, ed. N. P. Shah, 275–283. Academic Press.
- Oyewole, O. B. 1997. Lactic fermented foods in Africa and their benefits. *Food Control* 8 (5–6):289–297. doi:10.1016/S0956-7135(97)00075-3.
- Park, Y. W. 2009. Overview of bioactive components in milk and dairy products. In *Bioactive components in milk and dairy products*. Hoboken, NJ: Wiley-Blackwell.
- Park, Y. W., and M. S. Nam. 2015. Bioactive peptides in milk and dairy products: A review. *Korean Journal for Food Science of Animal Resources* 35 (6):831–840. doi:10.5851/kosfa.2015.35.6.831.
- Parvez, S., K. A. Malik, S. Ah Kang, and H. Y. Kim. 2006. Probiotics and their fermented food products are beneficial for health. *Journal of Applied Microbiology* 100 (6):1171–1185. doi:10.1111/j.1365-2672.2006.02963.x.
- Pereira, P. C. 2014. Milk nutritional composition and its role in human health. *Nutrition (Burbank, Los Angeles County, Calif.)* 30 (6): 619–627.
- Pessione, E., and S. Cirrincione. 2016. Bioactive molecules released in food by lactic acid bacteria: Encrypted peptides and biogenic amines. *Frontiers in Microbiology* 7:876. doi:10.3389/fmicb.2016.00876.
- Quigley, L., O. O'sullivan, C. Stanton, T. P. Beresford, R. P. Ross, G. F. Fitzgerald, and P. D. Cotter. 2013. The complex microbiota of raw milk. *FEMS Microbiology Reviews* 37 (5):664–698. doi:10.1111/1574-6976.12030.
- Rahmawati, I. S., and W. Suntornsuk. 2016. Effects of fermentation and storage on bioactive activities in milks and yoghurts. *Procedia Chemistry* 18:53–62. doi:10.1016/j.proche.2016.01.010.
- Rogeli, I. 2000. Milk, dairy products, nutrition and health. *Food Technology and Biotechnology* 38 (2):143–148.
- Salminen, S., A. Ouwehand, Y. Benno, and Y. K. Lee. 1999. Probiotics: How should they be defined? *Trends in Food Science & Technology* 10 (3):107–110. doi:10.1016/s0924-2244(99)00027-8.
- Samet-Bali, O. 2012. Development of fermented milk "leben made from spontaneous fermented cow's milk." *African Journal of Biotechnology* 11 (7):1829–1837. doi:10.5897/ajb11.2806.
- Savado, A., C. A. T. Ouattara, P. W. Savado, A. S. Ouattara, N. Barro, and A. S. Traore. 2004. Microorganisms involved in Fulani traditional fermented milk in Burkina Faso. *Pakistan Journal of Nutrition* 3 (2):134–139. doi:10.3923/pjn.2004.134.139.
- Savaiano, D. A. 2014. Lactose digestion from yogurt: Mechanism and relevance. *The American Journal of Clinical Nutrition* 99 (5): 1251S–1255S. doi:10.3945/ajcn.113.073023.
- Schaafsma, G. 2000. The protein digestibility–corrected amino acid score. *The Journal of Nutrition* 130 (7):1865S–1867S. doi:10.1093/jn/130.7.1865S.
- Schlegel, L., F. Grimont, E. Ageron, P. A. D. Grimont, and A. Bouvet 2003. Reappraisal of the taxonomy of the *Streptococcus bovis*/*Streptococcus equinus* complex and related species: Description of *Streptococcus gallolyticus* subsp. *gallolyticus* subsp. nov., *S. gallolyticus* subsp. *macedonicus* subsp. nov. and *S. gallolyticus* subsp. *pasteurianus* subsp. nov. *International Journal of Systematic and Evolutionary Microbiology* 53 (3):631–645. doi:10.1099/ijs.0.02361-0.
- Schoustra, S. E., C. Kasase, C. Toarta, R. Kassen, and A. J. Poulain. 2013. Microbial community structure of three traditional Zambian fermented products: Mabisi, chibwantu and munkoyo. *PLoS One* 8 (5):e63948. e63948
- Schutte, L. M. 2013. *Isolation and identification of the microbial consortium present in fermented milks from Sub-Saharan Africa*. Stellenbosch, South Africa: Stellenbosch University.
- Seppo, L., T. Jauhiainen, T. Poussa, and R. Korpela. 2003. A fermented milk high in bioactive peptides has a blood pressure-lowering effect in hypertensive subjects. *The American Journal of Clinical Nutrition* 77 (2):326–330. doi:10.1093/ajcn/77.2.326.
- Shiby, V. K., and H. N. Mishra. 2013. Fermented milks and milk products as functional foods—A review. *Critical Reviews in Food Science and Nutrition* 53 (5):482–496. doi:10.1080/10408398.2010.547398.
- Silva, C. C. G., S. P. M. Silva, and S. C. Ribeiro. 2018. Application of bacteriocins and protective cultures in dairy food preservation. *Frontiers in microbiology* 9:594. doi:10.3389/fmicb.2018.00594.
- Smacchi, E., and M. Gobbetti. 2000. Bioactive peptides in dairy products: Synthesis and interaction with proteolytic enzymes. *Food Microbiology* 17 (2):129–141. doi:10.1006/fmic.1999.0302.
- Stanton, C., R. Paul Ross, G. F. Fitzgerald, and D. V. Sinderen. 2005. Fermented functional foods based on probiotics and their biogenic metabolites. *Current Opinion in Biotechnology* 16 (2):198–203. doi:10.1016/j.copbio.2005.02.008.
- Steinkraus, K. H. 1994. Nutritional significance of fermented foods. *Food Research International* 27 (3):259–267. doi:10.1016/0963-9969(94)90094-9.
- Suliman, A. M. E., A. A. Ilayan, and A. E. El Faki. 2006. Chemical and microbiological quality of Garris, Sudanese fermented camel's milk product. *International Journal of Food Science and Technology* 41 (3):321–328. doi:10.1111/j.1365-2621.2005.01070.x.
- Suriyarachchi, V. R., and G. H. Fleet. 1981. Occurrence and growth of yeasts in yogurts. *Applied and Environmental Microbiology* 42 (4): 574–579.
- Tadesse, B. T., E. A. Ashley, S. Ongarello, J. Havumaki, M. Wijegoonewardena, J. G. Iveth, and S. Dittich. 2017. Antimicrobial resistance in Africa: A systematic review. *BMC Infectious Diseases* 17 (1):616–616. doi:10.1186/s12879-017-2713-1.
- Taha, S., M. El Abd, C. De Gobba, M. Abdel-Hamid, E. Khalil, and D. Hassan. 2017. Antioxidant and antibacterial activities of bioactive peptides in buffalo's yoghurt fermented with different starter cultures. *Food Science and Biotechnology* 26 (5):1325–1332. doi:10.1007/s10068-017-0160-9.
- Tamang, J. P., and K. Kailasapathy. 2010. *Fermented foods and beverages of the world*. Boca Raton, FL: CRC Press.
- Terefe, N. S. 2016. Food fermentation. In *Reference module in food science*. New York, NY: Elsevier.
- Teuber, M. 2009. Genus II. *Lactococcus*. *Bergey's manual of systematic bacteriology: The firmicutes*. 2nd ed. New York, NY: Springer.
- Teuber, M., L. Meile, and F. Schwarz. 1999. Acquired antibiotic resistance in lactic acid bacteria from food. In *Lactic acid bacteria: Genetics, metabolism and applications*, 115–137. New York, NY: Springer.
- Tournut, J. 1989. Applications of probiotics to animal husbandry. *Revue scientifique et technique (International Office of Epizootics)* 8 (2):551–556.
- USDA. 2018. Agricultural research service national nutrient database for standard reference. <https://ndb.nal.usda.gov/ndb/search/list>.
- Vedamuthu, E. R. 2006. Starter cultures for yogurt and fermented milks. In *Manufacturing Yogurt and fermented milks*, 89–116. Hoboken, NJ: Wiley-Blackwell.
- Venugopalan, V., K. A. Shriner, and A. Wong-Beringer. 2010. Regulatory oversight and safety of probiotic use. *Emerging Infectious Diseases* 16 (11):1661–1665. doi:10.3201/eid1611.100574.
- Walker, G. P., F. R. Dunshea, and P. T. Doyle. 2004. Effects of nutrition and management on the production and composition of milk



- fat and protein: A review. *Australian Journal of Agricultural Research* 55 (10):1009–1028. doi: [10.1071/AR03173](https://doi.org/10.1071/AR03173).
- Wang, L., M.-J. Guo, Q. Gao, J.-F. Yang, L. Yang, X.-L. Pang, and X.-J. Jiang. 2018. The effects of probiotics on total cholesterol: A meta-analysis of randomized controlled trials. *Medicine* 97 (5):e9679. doi: [10.1097/MD.00000000000009679](https://doi.org/10.1097/MD.00000000000009679).
- Wilson, R.T. 1988. *Camels*. London, UK: MacMillan Educational Press Ltd, CTA series.
- Witthuhn, R. C., T. Schoeman, and T. J. Britz. 2004. Isolation and characterization of the microbial population of different South African kefir grains. *International Journal of Dairy Technology* 57 (1):33–37. doi: [10.1111/j.1471-0307.2004.00126.x](https://doi.org/10.1111/j.1471-0307.2004.00126.x).
- World Bank. 2014. Business and livelihoods in African livestock: Investments to overcome information gaps. <http://documents.worldbank.org/curated/en/765551468209989452/pdf/860930WP0P147900Box382165B00PUBLIC0.pdf>.
- Wouters, J. T. M., E. H. E. Ayad, J. Hugenholtz, and G. Smit. 2002. Microbes from raw milk for fermented dairy products. *International Dairy Journal* 12 (2–3):91–109. doi: [10.1016/S0958-6946\(01\)00151-0](https://doi.org/10.1016/S0958-6946(01)00151-0).
- Wulschleger, S., C. Lacroix, B. Bonfoh, A. Sissoko-Thiam, S. Hugenschmidt, E. Romanens, S. Baumgartner, I. Traoré, M. Yaffee, C. Jans, and L. Meile. 2013. Analysis of lactic acid bacteria communities and their seasonal variations in a spontaneously fermented dairy product (Malian fèfè) by applying a cultivation/genotype-based binary model. *International Dairy Journal* 29 (1):28–35. doi: [10.1016/j.idairyj.2012.08.001](https://doi.org/10.1016/j.idairyj.2012.08.001).
- Yamamoto, N., M. Maeno, and T. Takano. 1999. Purification and characterization of an antihypertensive peptide from a yogurt-like product fermented by *Lactobacillus helveticus* CPN4. *Journal of Dairy Science* 82 (7):1388–1393. doi: [10.3168/jds.S0022-0302\(99\)75364-6](https://doi.org/10.3168/jds.S0022-0302(99)75364-6).