



## Critical Reviews in Food Science and Nutrition

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/bfsn20>

### Camellia Sinensis in Asymptomatic Hyperuricaemia: A Meta-analysis of Tea or Tea Extract Effects on Uric Acid Levels

Ilaria Peluso<sup>a</sup>, Alessia Teichner<sup>a</sup>, Hussein Manafikhi<sup>a</sup> & Maura Palmery<sup>a</sup>

<sup>a</sup> Department of Physiology and Pharmacology "V. Erspamer", "Sapienza" University of Rome, P.le Aldo Moro 5, 00185 Rome, Italy.

Accepted author version posted online: 07 Apr 2015.



[Click for updates](#)

To cite this article: Ilaria Peluso, Alessia Teichner, Hussein Manafikhi & Maura Palmery (2015): Camellia Sinensis in Asymptomatic Hyperuricaemia: A Meta-analysis of Tea or Tea Extract Effects on Uric Acid Levels, Critical Reviews in Food Science and Nutrition, DOI: [10.1080/10408398.2014.889653](https://doi.org/10.1080/10408398.2014.889653)

To link to this article: <http://dx.doi.org/10.1080/10408398.2014.889653>

Disclaimer: This is a version of an unedited manuscript that has been accepted for publication. As a service to authors and researchers we are providing this version of the accepted manuscript (AM). Copyediting, typesetting, and review of the resulting proof will be undertaken on this manuscript before final publication of the Version of Record (VoR). During production and pre-press, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal relate to this version also.

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

**Camellia sinensis in asymptomatic hyperuricaemia: a meta-analysis of tea or tea extract effects on uric acid levels**

Ilaria Peluso<sup>1</sup>, Alessia Teichner<sup>1</sup>, Hussein Manafikhi<sup>1</sup> and Maura Palmery<sup>1\*</sup>

<sup>1</sup>*Department of Physiology and Pharmacology "V. Erspamer", "Sapienza" University of Rome, P.le Aldo Moro 5, 00185 Rome, Italy.*

**\*Address for correspondence:**

Maura Palmery, Prof.

Department of Physiology and Pharmacology "V. Erspamer"

"Sapienza" University of Rome ,

P.le Aldo Moro 5, 00185 Rome, Italy.

[maura.palmery@uniroma1.it](mailto:maura.palmery@uniroma1.it)

Phone +39-06-49912573; Fax: +39-06-49912480

**Short title for the running head:** Camellia sinensis in asymptomatic hyperuricaemia

**Abstract**

Flavanols of *Camellia sinensis* exhibit uric acid (UA) lowering effect, through the modulation of both xanthine oxidase and urate excretion. In order to investigate the potential benefit of *Camellia Sinensis* products in asymptomatic hyperuricaemia, a meta-analysis of long-term Randomized Controlled Trials (RCT) with tea or tea extract has been conducted. From 20 human intervention studies selected only 5 RCT (13 interventions) were suitable for meta-analysis (n=472).

The current "normal" range set for hyperuricaemia fails to identify patients with potential metabolic disorders. Therefore on the basis of the literature data, we fixed cut off limits for UA baseline levels of 4.5 mg/dl for women, 6.1 mg/dl for men and 5.5 mg/dl for studies involving mixed populations.

Statistically significant effects were not found, but subgroup analysis revealed that the Pooled Estimate effect was different in subjects with baseline levels under [MD (95% CI): 0.1078 (-0.0528 to 0.2684)] and over the cut off [MD (95% CI): -0.0239 (0.3311 to 0.2833)].

However, due to the low number of RCT and to the lack of data on bioavailability, it is difficult to draw any firm conclusion and more studies are needed to establish if tea flavanols could be useful in asymptomatic hyperuricaemia treatment.

**Keywords:** long term intervention, meta-analysis, uric acid, xanthine oxidase, urate transport.

## INTRODUCTION

Black tea (BT) and green tea (GT) are very different due to the processing or harvesting times of the leaves and leaf buds of the *Camellia sinensis*, which lead to a different composition in phenolic compounds (Serafini et al., 2011a). In the case of GT, the leaves are steamed quickly after harvesting to prevent oxidation of polyphenols and the major flavonoids present are the flavanols epicatechin and its gallate derivatives (Serafini et al., 2011a). These flavonoids are present in lower amounts in BT where they are converted, during the “fermentation” period, to complex condensation products such as theaflavins and thearubigins (Serafini et al., 2011a). Despite the role of uric acid (UA) in Plasma Non-Enzymatic Antioxidant Capacity (Serafini et al., 2011b), some of the protective effects of Epigallocatechin-3-gallate (EGCG) have been ascribed to its capability to reduce excessive UA level (Yokozawa et al., 2004). Accordingly, the latter can also be a dangerous signal for immune system (Shi et al., 2003), and hyperuricaemia has been associated to oxidative stress (Lahera et al., 2006).

Hyperuricaemia is an elevated UA level in blood that is most often mild and remains asymptomatic (Kanellis et al., 2004; Mikuls et al., 2005; Zoppini et al., 2011). However, chronic asymptomatic hyperuricaemia may be complicated by gout, urolithiasis and possibly nephropathy. Gout is an inflammatory response to deposition of monosodium urate crystals in and around joints (Mandell, 2008; Tausche et al., 2009; Fink et al., 2013; Wang et al., 2012). Therapy of hyperuricaemia and gout has to depend on pathogenesis and stage of the disease. Guidelines of management for gout recommend that serum urate concentration should be maintained below 6 mg/dl to promote crystal dissolution leading to prevention of recurrent gouty

attack (Mandell, 2008; Tausche et al., 2009). In the presence of renal impairment, allopurinol, an analogue of hypoxanthine, is the treatment of choice for urate lowering therapy (Fink et al., 2013; Wang et al., 2012).

There is still no consensus on the treatment of individuals with asymptomatic hyperuricaemia (Kanellis et al., 2004; Mikuls et al., 2005; Zoppini et al., 2011). However, the latter is an important etiological factor in common morbidities such as arterial hypertension, cardiovascular disease (CVD) and type 2 diabetes mellitus (T2D) (Kanellis et al., 2004; Zoppini et al., 2011; Becker and Jolly, 2006; Strazzullo and Puig, 2007; 15. Hwu and Lin, 2010). Urate lowering drug treatment is normally not indicated in asymptomatic hyperuricaemic individuals, considering that the allopurinol hypersensitivity syndrome has been reported (Markel, 2005). For this reason Febuxostat, a more tolerable xanthine oxidase (XO) inhibitor has received attention (Tayar et al., 2012).

In this context, also theaflavins and catechins inhibit XO to produce UA and theaflavin-3,3'-digallate, the most potent inhibitor of XO among these compounds, acts as a competitive inhibitor (Lin et al., 2000; Dew et al., 2005). Moreover, EGCG (Yin et al., 2009; Palermo et al., 2005; Han et al., 2012; Park and Dong 2003; Nomura et al., 2000) and theaflavins (Park and Dong 2003; Nomura et al., 2000; Łuczaj and Skrzydlewska, 2005) inhibit the nuclear factor  $\kappa$ B (NF- $\kappa$ B) (Park and Dong 2003; Nomura et al., 2000; Łuczaj and Skrzydlewska, 2005) and the aryl hydrocarbon receptor (AhR) EGCG (Yin et al., 2009; Palermo et al., 2005; Han et al., 2012) response pathways, key regulators of XO (Martinez-Hervas et al., 2010; Sugihara et al., 2001). Flavonoids have been proposed as alternative of uricosuric agents (e.g. benzbromarone) in allopurinol-allergic patients (Yu et al., 2007). Renal urate transporter (URAT) (Yu et al., 2007;

Hu et al., 2012; Wang et al., 2010), organic anion transporters (OAT) (Hu et al., 2012; Wang et al., 2010; Sekine et al., 2006), multidrug resistance-associated proteins (MRP) (Sekine et al., 2006; El-Sheikh et al., 2008) and uromodulin (Hu et al., 2012), play important roles in renal urate excretion and are regulated by both flavonoids (Yu et al., 2007; Hu et al., 2012; Wang et al., 2010) and hyperuricaemic drugs (Yu et al., 2007; El-Sheikh et al., 2008). In particular tea catechins (Roth et al., 2011; Fuchikami et al., 2006; Vaidyanathan and Walle, 2001) interact with OAT (Roth et al., 2011; Fuchikami et al., 2006) and MRP (Vaidyanathan and Walle, 2001).

Moreover allopurinol and flavonoids reversed dys-regulations of OAT and URAT and attenuate hyperuricaemia and renal dysfunction in rats induced by fructose intake (Hu et al., 2009).

Besides, it has been suggested that the inhibitory effect of green tea on calcium oxalate urolithiasis is most likely due to antioxidative effects (Itoh et al., 2005). Contrarily to teas, other antioxidant beverages, such as orange juice and pink grapefruit juice, were shown to promote XO activity (Dew et al., 2005). Moreover, cocoa beverage significantly increase calcium and oxalic acid excretion and also leafy vegetable foods contain oxalate (Hesse et al., 1993). This could be critical in hyperuricaemia considering that increases in calcium and oxalate excretion promote kidney stones (Taylor and Curhan, 2006).

Accordingly to the urate lowering effect of *Camellia sinensis*, despite the evidence showing that reducing soft-drink consumption decreased recurrent symptomatic stone risk (Fink et al., 2013), observational studies have found that tea consumption reduces the risk of stone formation (Taylor and Curhan, 2006).

Therefore, the objective of this meta-analysis was to investigate the effect of *Camellia sinensis* products on hyperuricaemia. To this aim, in the absence of specific Randomize Controlled Trials

(RCT), a systematic review and meta-analysis of chronic intervention studies with BT, GT or extracts and UA concentrations has been conducted in this work.

## METHODS

### Data source and study selection

We performed a systematic search (Figure 1) up to October 2013. Any study that met the following criteria was selected for meta-analysis: randomized long term placebo-controlled trials where the plasma levels of UA were measured before and after placebo and treatment and that appear in an edited English journal. All human chronic studies were included if they met the above criteria regardless of the dose of BT, GT or extract and of the healthy status of subjects. First trials were identified through title or abstract. Duplicates and irrelevant references have been removed from our search and from result eligible studies were included through full text and reviewed by two independent authors (I.P. and H.M.). Discrepancies were resolved by discussion with a third reviewer (M.P.).

A total of 20 intervention studies after tea or tea extracts administration were identified as suitable and were retrieved for complete review. Figure 1 depicts the flow of studies in this review and the four-phase diagram of meta-analysis, according to the PRISMA Statement (Moher et al., 2009).

Of the reviewed studies, we excluded studies referring to bolus administration (Henning et al., 2005; Jówko et al., 2012; Leenen et al., 2000; Müller et al., 2010; Natella et al., 2002; Rabovsky et al., 2006; Van Amelsvoort et al., 2001), not placebo-controlled trials (Gomikawa et al., 2008;

Kimura et al., 2002; Sone et al., 2011) and papers reporting post-treatment but not pre-treatment values (Panza et al., 2008), not shown data (Chow et al., 2003; Maron et al., 2003; Rondanelli et al., 2009) or figures (Li et al., 2010).

The remaining 5 studies (Bahorun et al., 2010; Hsu et al., 2008; Hsu et al., 2011; Princen et al., 1998; van het Hof et al., 1997), reporting 8 intervention with BT from 3 references and 5 data from 4 RCT with GT or green tea extract (GTE), met our inclusion criteria and could provide data for the analyses (Figure 1).

### **Data extraction**

A data extraction form including quality characteristics was used and a quality score was assigned as previously described (Peluso et al., 2013). All data extraction and quality assessment were performed independently by two reviewers (I.P. and A.T.) to ensure uniformity, and all data were entered by these reviewers, who checked the extracted data and validity criteria against the original published report. Discrepancies were resolved by discussion with a third reviewer (M.P.).

Hyperuricaemia is defined as serum values of UA  $\geq 6.8$  and  $\geq 5.4$  mg/dL for men and women, respectively (Rodrigues et al., 2012). However, the current "normal" range set for hyperuricaemia often fails to identify patients with potential metabolic disorders (Zhang et al., 2013). The association of serum UA with CVD risk factor (Rodrigues et al., 2012), metabolic syndrome (Rodrigues et al., 2012; Zhang et al., 2013), T2D (Kawamoto et al., 2013) and nonalcoholic fatty liver disease (Hwang et al., 2011) is gender-specific and more pronounced in women.



Therefore, on the basis of the literature data, we fixed a cut off of 5.5 mg/dl for studies involving mixed population of male and females (Rodrigues et al., 2012; Hwang et al., 2011), and of 4.5 mg/dl and 6.1 mg/dl, respectively for women and men, when reported (Rodrigues et al., 2012; Zhang et al., 2013; Kawamoto et al., 2013) (Figure 1).

Accordingly with these cut off limits we have found 8 and 5 interventions from 3 (Bahorun et al., 2010; Princen et al., 1998; van het Hof et al., 1997) and 3 (Bahorun et al., 2010; Hsu et al., 2008; Hsu et al., 2011) studies for UA in normal range and over the limits associated with diseases risk, respectively (Figure 1).

### **Meta-analysis**

For the meta-analysis, we calculated the mean change of plasma UA concentrations from baseline to follow-up for each intervention and control group, if not reported, and the SD of delta concentration was estimated as previously described (Peluso et al., 2013). All values were converted to mg/dl, if necessary. For each study with more than one intervention group, we divided the control group evenly according to the number of intervention groups (Peluso et al., 2013). When the data were presented as SEMs, SD was obtained by multiplying SEM by the square root of the sample size.

Statistical heterogeneity, the presence of publication bias, Mean difference (MD) and confidence intervals 95% (CI 95%) were evaluated as previously described (Peluso et al., 2013).

To explore the effects of modifier factors on the primary outcomes, subgrouping was used when data from  $\geq$  three studies and  $\geq$  five interventions were included. These factors were the type of

intervention (BT or GT/GTE) and baseline levels of UA. Meta regression was not performed because less than ten studies were available (Peluso et al., 2013).

## RESULTS

### Description of included studies

Studies characteristics and the results of the assessment of risk of bias of the included trials (quality assessment) are shown in Table 1. The ingested dose ranged from 600 ml to 900 ml of tea and from 1.2 g to 3.6 g of GTE; the latter was equivalent to the consumption of 18 cups of GT per day (Princen et al., 1998).

Average length is 4-16 weeks and the number of participants in individual trials ranged from 45 to 232. Participant withdrawal rate from the trial was between 4 and 22% and 3 studies (Bahorun et al., 2010; Princen et al., 1998; van het Hof et al., 1997) involved multiple interventions in parallel design.

Age ranged between 16 and 65 years (Table 1) and one study selected only females (Hsu et al., 2008). The remaining trials included percentage of female participants from 41% to 65% and the characteristics and comorbidity status of the participants varied (Table 1). Only one study recruited healthy participants (van het Hof et al., 1997), whereas the remaining RCT evaluated the effect of the intervention among subjects with CVD (Bahorun et al., 2010), diabetes (Hsu et al., 2011) or risk factors such as smoking habit (Princen et al., 1998) and obesity (Hsu et al., 2008).

None of the trials measured bioavailability and the more frequent limitations were the absence of double blinding (3/5), food antioxidant intake in subjects selection criteria (3/5) and wash out/run-in period (3/5). Only one study presented a high weight bias because randomization generated baseline levels for treated subjects and controls more different than the observed effect after follow up (Hsu et al., 2011). All studies adequately evaluated compliance but, as expected, blinding of subjects have been obtained only in studies with GTE (Hsu et al., 2008; Hsu et al., 2011).

### Meta-analysis

The number of subjects was 472 of which 309 received BT, GT or GTE and 163 placebo. No significant difference was detected with meta-analysis for the change of plasma concentration of UA between tea or extracts ingestion and control (Fig. 2A). The Pooled Estimate effect size was the same in fixed and random effect models (Fig. 2A) and similar in quality effect model [MD (95%CI): 0,0824 (-0,0866 to 0,2514)]. Low statistical heterogeneity was found (I<sup>2</sup> 0%,  $t^2$  0, Q 15%, Egger intercept -0.0135,  $p = 0.9798$ ), Funnel plot showed no asymmetric distribution of results and Trim-and-fill analysis did not suggest potential publication bias (Fig. 2B). A no significant decrease and increase, respectively, were detected with meta-analysis for the change of plasma concentration of UA between tea or extracts ingestion and control in subjects with baseline concentration over and under the cut off in fixed, random and quality effect models (Table 2). Also subgrouping for type of treatment revealed no significant effect size in fixed, random and quality effect models (Table 2). Furthermore, the exclusion of the study conducted

on healthy participants (van het Hof et al., 1997) did not significantly change (Table 2) the results obtained with meta analysis (Fig. 2A).

## CONCLUSION

In our meta-analysis, statistical significance has not been found, probably by reason of the low number of subjects, but different results have been found, after teas or tea extracts chronic consumption, in subjects with normal baseline levels and with hyperuricaemia, where increased and decreased UA levels have been observed, respectively (Table 2).

These opposite effects could be due to the dual effect on UA concentration of polyphenols of tea. The latter could prevent oxidation (Serafini et al., 2011a) but may also decrease the production and increase the excretion of UA (Roth et al., 2011; Fuchikami et al. 2006; Vaidyanathan and Walle 2001). Despite the higher effect of theaflavins as XO inhibitors in vitro (Lin et al., 2000; Dew et al., 2005), no significant differences have been observed in the effect size between BT or GT and GTE subgroups (Table 2).

However, out of one studies had UA as secondary outcome; therefore only this RCT (Bahorun et al., 2010) reported data for men and women separately, stratifying them for baseline levels.

This study with BT showed decreases of UA only in subjects (especially women) with high baseline levels and in men with baseline UA concentrations above 8mg/dl, the latter lowered after wash-out (BT  $7.3 \pm 1.7$  mg/dl and water  $8.0 \pm 2.0$  mg/dl) (Bahorun et al., 2010), suggesting an inhibitory effect on XO that persists after discontinuation of consumption. Accordingly to this hypothesis also in an uncontrolled trial GTE decreased UA after 7 days of wash-out, subsequent

to 7 days of supplementation, when catechins went down to pre-treatment concentrations (Kimura et al., 2002). Panza et al. (2008), reported a decrease of UA levels, after 7 days of ingestion of GT (600ml), and an inhibition of the exercise-induced activation of XO.

However, only one of the studies in bolus administration of tea or tea extracts, reported decreases in UA concentration (Rabovsky et al., 2006), whereas in the others increased (Henning et al., 2005; Natella et al., 2002) or unchanged (Jówko et al., 2012; Leenen et al., 2000; Müller et al., 2010; Van Amelsvoort et al. 2001; Kimura et al., 2002) UA levels were found. Moreover, Bahorun et al. (2010) in women observed decreases in UA after wash-out also in controls, suggesting differential gender effects of a high fluid intake.

Due to the limited number of studies, the major limitation of this meta-analysis is that we could not investigate sex, age, healthy status and body mass index differences between subjects or dose and duration of treatment effects. However, concerning the excluded studies both decreased (Kimura et al., 2002; Panza et al., 2008) or unchanged (Gomikawa et al., 2008; Li et al., 2010) UA levels have been reported after long term GT (Gomikawa et al., 2008; Panza et al., 2008) or GTE Kimura et al., 2002; Li et al., 2010) consumption in healthy subjects. The decreases were observed after 7 day of consumption (Kimura et al., 2002; Panza et al., 2008), while treatment of 2 weeks (Gomikawa et al., 2008) or 16 weeks (Li et al., 2010) were unable to change the UA concentration. Moreover, in an uncontrolled trial have been observed decreases of UA after 9 weeks of GT (100mg/day of total catechins) and increases of UA with catechin-enriched GT (400mg/day of total catechins), but these effects did not reach statistical significance (Sone et al., 2011). Therefore, longer or higher consumptions seem to be not related with higher effects.

Tea polyphenols could regulate gut microbioma (Axling et al., 2012), as well as phase I, II and III metabolism/transport systems (Chow et al., 2006; Chow et al., 2007; Vischini et al., 2011); all involved in their metabolism (Serafini et al., 2011; Henning et al., 2011).

From there, a critical point of this meta-analysis is that none of the selected studies reported data of tea flavanols or metabolites after the intervention. Besides, bioavailability in healthy and dys-metabolic subjects could profoundly differ because metabolic diseases and obesity are associated with dysbiosis (Hooper et al., 2012; Tremaroli and Bäckhed, 2012) and low grade chronic inflammation (Calder et al., 2011) and inflammatory cytokines modulate cytochrome P450 (Liptrott et al., 2009) and MRP (Kimura et al., 2006).

These evidence could be taken into account in the interpretation of data of these meta-analysis, considering that the subgroup with UA levels above the cut off comprise studies involving subjects with CVD (Bahorun et al., 2010) diabetes (Hsu et al. 2011) or obesity (Hsu et al., 2008). Of these studies only one, which reported better results on the effect on hyperuricaemia, recruited patients under treatment with drugs against hypertension or diabetes (Bahorun et al., 2010). Therefore, we cannot exclude a possible interaction of tea with other drugs at the level of the MRP (Vaidyanathan and Walle, 2001), involved in renal urate excretion (Sekine et al., 2006; El-Sheikh et al., 2008).

Hyperuricaemia is often associated with other diseases (Becker and Jolly, 2006; Hwu and Lin, 2010; Kanellis et al., 2004; Strazzullo and Puig, 2007; Zoppini et al., 2011), therefore considering that our data suggest that tea or tea extracts exerted differential effects in subjects with high or low UA concentration, RCT with UA levels and bioavailability as primary end

point, considering separately men and women with asymptomatic hyperuricaemia are needed in order to establish if tea flavonols could be useful in these subjects.

**Conflict-of-interest disclosure:** All authors declare that they have no conflicts of financial and commercial interest.

## References

- Axling U, Olsson C, Xu J, Fernandez C, Larsson S, Ström K, Ahrné S, Holm C, Molin G, Berger K. (2012) Green tea powder and *Lactobacillus plantarum* affect gut microbiota, lipid metabolism and inflammation in high-fat fed C57BL/6J mice. *Nutr Metab (Lond)*;9(1):105.
- Bahorun T, Luximon-Ramma A, Gunness TK, Sookar D, Bhoyroo S, Jugessur R, Reebye D, Googoolye K, Crozier A, Aruoma OI. (2010) Black tea reduces uric acid and C-reactive protein levels in humans susceptible to cardiovascular diseases. *Toxicology*;278(1):68-74.
- Becker MA, Jolly M. (2006) Hyperuricemia and associated diseases. *Rheum Dis Clin North Am*;32(2):275-93.
- Calder PC, Ahluwalia N, Brouns F, Buetler T, Clement K, Cunningham K, Esposito K, Jönsson LS, Kolb H, Lansink M, Marcos A, Margioris A, Matusheski N, Nordmann H, O'Brien J, Pugliese G, Rizkalla S, Schalkwijk C, Tuomilehto J, Wärnberg J, Watzl B, Winklhofer-Roob BM. (2011) Dietary factors and low-grade inflammation in relation to overweight and obesity. *Br J Nutr*;106(3):S5-78.
- Chow HH, Cai Y, Hakim IA, , Crowell JA, Shahi F, Brooks CA, Dorr RT, Hara Y, Alberts DS. (2003) Pharmacokinetics and safety of green tea polyphenols after multiple-dose administration of epigallocatechin gallate and polyphenon E in healthy individuals. *Clin Cancer Res*;9(9):3312-9.
- Chow HH, Hakim IA, Vining DR, Crowell JA, Cordova CA, Chew WM, Xu MJ, Hsu CH, Ranger-Moore J, Alberts DS. (2006) Effects of repeated green tea catechin administration on human cytochrome P450 activity. *Cancer Epidemiol Biomarkers Prev*;15(12):2473-6.



- Chow HH, Hakim IA, Vining DR, Crowell JA, Tome ME, Ranger-Moore J, Cordova CA, Mikhael DM, Briehl MM, Alberts DS. (2007) Modulation of human glutathione S-transferases by polyphenol intervention. *Cancer Epidemiol Biomarkers Prev.*;16(8):1662-6.
- Dew TP, Day AJ, Morgan MR. (2005) Xanthine oxidase activity in vitro: effects of food extracts and components. *J Agric Food Chem.*;53(16):6510-5.
- El-Sheikh AA, van den Heuvel JJ, Koenderink JB, Russel FG. (2008) Effect of hypouricaemic and hyperuricaemic drugs on the renal urate efflux transporter, multidrug resistance protein 4. *Br J Pharmacol.*;155(7):1066-75.
- Fink HA, Wilt TJ, Eidman KE, Garimella PS, MacDonald R, Rutks IR, Brasure M, Kane RL, Ouellette J, Monga M. (2013) Medical management to prevent recurrent nephrolithiasis in adults: a systematic review for an American College of Physicians Clinical Guideline. *Ann Intern Med.*;158(7):535-43.
- Fuchikami H, Satoh H, Tsujimoto M, Ohdo S, Ohtani H, Sawada Y. (2006) Effects of herbal extracts on the function of human organic anion-transporting polypeptide OATP-B. *Drug Metab Dispos.*;34(4):577-82.
- Gomikawa S, Ishikawa Y, Hayase W, Haratake Y, Hirano N, Matuura H, Mizowaki A, Murakami A, Yamamoto M. (2008) Effect of ground green tea drinking for 2 weeks on the susceptibility of plasma and LDL to the oxidation ex vivo in healthy volunteers. *Kobe J Med Sci.*;54(1):E62-72.

- Han SG, Han SS, Toborek M, Hennig B. (2012) EGCG protects endothelial cells against PCB 126-induced inflammation through inhibition of AhR and induction of Nrf2-regulated genes. *Toxicol Appl Pharmacol.*;261(2):181-8.
- Henning SM, Niu Y, Liu Y, Lee NH, Hara Y, Thames GD, Minutti RR, Carpenter CL, Wang H, Heber D. (2005) Bioavailability and antioxidant effect of epigallocatechin gallate administered in purified form versus as green tea extract in healthy individuals. *J Nutr Biochem.*;16(10):610-6.
- Henning SM, Wang P, Heber D. (2011) Chemopreventive effects of tea in prostate cancer: green tea versus black tea. *Mol Nutr Food Res.*;55(6):905-20.
- Hesse A, Siener R, Heynck H, Jahn A. (1993) The influence of dietary factors on the risk of urinary stone formation. *Scanning Microsc.*;7(3):1119-27.
- Hooper LV, Littman DR, Macpherson AJ. (2012) Interactions between the microbiota and the immune system. *Science.*;336(6086):1268-73.
- Hsu CH, Liao YL, Lin SC, Tsai TH, Huang CJ, Chou P. (2011) Does supplementation with green tea extract improve insulin resistance in obese type 2 diabetics? A randomized, double-blind, and placebo-controlled clinical trial. *Altern Med Rev.*;16(2):157-63.
- Hsu CH, Tsai TH, Kao YH, Hwang KC, Tseng TY, Chou P. (2008) Effect of green tea extract on obese women: a randomized, double-blind, placebo-controlled clinical trial. *Clin Nutr.*;27(3):363-70.
- Hu QH, Wang C, Li JM, Zhang DM, Kong LD. (2009) Allopurinol, rutin, and quercetin attenuate hyperuricemia and renal dysfunction in rats induced by fructose intake: renal organic ion transporter involvement. *Am J Physiol Renal Physiol.*;297(4):F1080-91.

- Hu QH, Zhang X, Wang X, Jiao RQ, Kong LD. (2012) Quercetin regulates organic ion transporter and uromodulin expression and improves renal function in hyperuricemic mice. *Eur J Nutr.*;51(5):593-606.
- Hwang IC, Suh SY, Suh AR, Ahn HY. (2011) The relationship between normal serum uric acid and nonalcoholic fatty liver disease. *J Korean Med Sci.*;26(3):386-91.
- Hwu CM, Lin KH. (2010) Uric acid and the development of hypertension. *Med Sci Monit.*;16(10):RA224-30.
- Itoh Y, Yasui T, Okada A, Tozawa K, Hayashi Y, Kohri K. (2005) Preventive effects of green tea on renal stone formation and the role of oxidative stress in nephrolithiasis. *J Urol.*;173(1):271-5.
- Jówko E, Sacharuk J, Balasinska B, Wilczak J, Charmas M, Ostaszewski P, Charmas R. (2012) Effect of a single dose of green tea polyphenols on the blood markers of exercise-induced oxidative stress in soccer players. *Int J Sport Nutr Exerc Metab.*;22(6):486-96.
- Kanellis J, Feig DI, Johnson RJ. (2004) Does asymptomatic hyperuricaemia contribute to the development of renal and cardiovascular disease? An old controversy renewed. *Nephrology (Carlton).*;9(6):394-9.
- Kawamoto R, Tabara Y, Kohara K, Kusunoki T, Abe M, Miki T. (2013) Serum uric acid is more strongly associated with impaired fasting glucose in women than in men from a community-dwelling population. *PLoS One.*;8(6):e65886.
- Kimura A, Ishida Y, Hayashi T, Wada T, Yokoyama H, Sugaya T, Mukaida N, Kondo T. (2006) Interferon-gamma plays protective roles in sodium arsenite-induced renal injury by up-

regulating intrarenal multidrug resistance-associated protein 1 expression. *Am J Pathol.*;169(4):1118-28.

Kimura M, Umegaki K, Kasuya Y, Sugisawa A, Higuchi M. (2002) The relation between single/double or repeated tea catechin ingestions and plasma antioxidant activity in humans. *Eur J Clin Nutr.*;56(12):1186-93.

Lahera V, Goicoechea M, de Vinuesa SG, Oubiña P, Cachofeiro V, Gómez-Campderá F, Amann R, Luño J. (2006) Oxidative stress in uremia: the role of anemia correction. *J Am Soc Nephrol.*;7(12 Suppl 3):S174-7.

Leenen R, Roodenburg AJ, Tijburg LB, Wiseman SA. (2000) A single dose of tea with or without milk increases plasma antioxidant activity in humans. *Eur J Clin Nutr.*;54(1):87-92.

Li L, Chen CY, Aldini G, Johnson EJ, Rasmussen H, Yoshida Y, Niki E, Blumberg JB, Russell RM, Yeum KJ. (2010) Supplementation with lutein or lutein plus green tea extracts does not change oxidative stress in adequately nourished older adults. *J Nutr Biochem.*;21(6):544-9.

Lin JK, Chen PC, Ho CT, Lin-Shiau SY. (2000) Inhibition of xanthine oxidase and suppression of intracellular reactive oxygen species in HL-60 cells by theaflavin-3,3'-digallate, (-)-epigallocatechin-3-gallate, and propyl gallate. *J Agric Food Chem.*;48(7):2736-43.

Liptrott NJ, Penny M, Bray PG, Sathish J, Khoo SH, Back DJ, Owen A. (2009) The impact of cytokines on the expression of drug transporters, cytochrome P450 enzymes and chemokine receptors in human PBMC. *Br J Pharmacol.*;156(3):497-508.

Łuczaj W, Skrzydlewska E. (2005) Antioxidative properties of black tea. *Prev Med.* 40(6):910-8.

Mandell BF. (2008) Clinical manifestations of hyperuricemia and gout. *Cleve Clin J Med.*;75 Suppl 5:S5-8.

Markel A. (2005) Allopurinol-induced DRESS syndrome. *Isr Med Assoc J.*;7(10):656-60.

Maron DJ, Lu GP, Cai NS, Wu ZG, Li YH, Chen H, Zhu JQ, Jin XJ, Wouters BC, Zhao J.

(2003) Cholesterol-lowering effect of a theaflavin-enriched green tea extract: a randomized controlled trial. *Arch Intern Med.*;163(12):1448-53.

Martinez-Hervas S, Real JT, Ivorra C, Priego A, Chaves FJ, Pallardo FV, Viña JR, Redon J,

Carmena R, Ascaso JF. (2010) Increased plasma xanthine oxidase activity is related to nuclear factor kappa beta activation and inflammatory markers in familial combined hyperlipidemia. *Nutr Metab Cardiovasc Dis.*; 20(10):734-9.

Mikuls TR, Farrar JT, Bilker WB, Fernandes S, Saag KG. (2005) Suboptimal physician

adherence to quality indicators for the management of gout and asymptomatic hyperuricaemia: results from the UK General Practice Research Database (GPRD). *Rheumatology (Oxford).*:44(8):1038-42.

Moher D, Liberati A, Tetzlaff J, Altman DG. PRISMA Group. (2009) Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med.*;6, e1000097.

Müller N, Ellinger S, Alteheld B, Ulrich-Merzenich G, Berthold HK, Vetter H, Stehle P. (2010)

Bolus ingestion of white and green tea increases the concentration of several flavan-3-ols in plasma, but does not affect markers of oxidative stress in healthy non-smokers. *Mol Nutr Food Res.*;54(11):1636-45.

Natella F, Nardini M, Giannetti I, Dattilo C, Scaccini C. (2002) Coffee drinking influences

plasma antioxidant capacity in humans. *J Agric Food Chem.*;50(21):6211-6.

- Nomura M, Ma W, Chen N, Bode AM, Dong Z. (2000) Inhibition of 12-O-tetradecanoylphorbol-13-acetate-induced NF-kappaB activation by tea polyphenols, (-)-epigallocatechin gallate and theaflavins. *Carcinogenesis*.;21(10):1885-90.
- Palermo CM, Westlake CA, Gasiewicz TA. (2005) Epigallocatechin gallate inhibits aryl hydrocarbon receptor gene transcription through an indirect mechanism involving binding to a 90 kDa heat shock protein. *Biochemistry*.;44(13):5041-52.
- Panza VS, Wazlawik E, Ricardo Schütz G, Comin L, Hecht KC, da Silva EL. (2008) Consumption of green tea favorably affects oxidative stress markers in weight-trained men. *Nutrition*.;24(5):433-42.
- Park AM, Dong Z. (2003) Signal transduction pathways: targets for green and black tea polyphenols. *J Biochem Mol Biol*.;36(1):66-77.
- Peluso I, Raguzzini A, Serafini M. (2013) Effect of flavonoids on circulating levels of TNF- $\alpha$  and IL-6 in humans: a systematic review and meta-analysis. *Mol Nutr Food Res*.;57(5):784-801.
- Princen HM, van Duyvenvoorde W, Buytenhek R, Blonk C, Tijburg LB, Langius JA, Meinders AE, Pijl H. (1998) No effect of consumption of green and black tea on plasma lipid and antioxidant levels and on LDL oxidation in smokers. *Arterioscler Thromb Vasc Biol*.;18(5):833-41.
- Rabovsky A, Cuomo J, Eich N. (2006) Measurement of plasma antioxidant reserve after supplementation with various antioxidants in healthy subjects. *Clin Chim Acta*.;371(1-2):55-60.

- Rodrigues SL, Baldo MP, Capingana P, Magalhães P, Dantas EM, Molina Mdel C, Salaroli LB, Morelato RL, Mill JG. (2012) Gender distribution of serum uric acid and cardiovascular risk factors: population based study. *Arq Bras Cardiol.*:98(1):13-21.
- Rondanelli M, Opizzi A, Solerte SB, Trotti R, Klersy C, Cazzola R. (2009) Administration of a dietary supplement (N-oleyl-phosphatidylethanolamine and epigallocatechin-3-gallate formula) enhances compliance with diet in healthy overweight subjects: a randomized controlled trial. *Br J Nutr.*;101(3):457-64.
- Roth M, Timmermann BN, Hagenbuch B. (2011) Interactions of green tea catechins with organic anion-transporting polypeptides. *Drug Metab Dispos.*;39(5):920-6.
- Sekine T, Miyazaki H, Endou H. (2006) Molecular physiology of renal organic anion transporters. *Am J Physiol Renal Physiol.*;290(2):F251-61.
- Serafini M, Del Rio D, Yao DN, Bettuzzi S, Peluso I. (2011a) Health Benefits of Tea. In: Benzie IFF, Wachtel-Galor S, editors. *Herbal Medicine: Biomolecular and Clinical Aspects. 2nd edition*. Boca Raton (FL): CRC Press; Chapter 12: 239-262.
- Serafini M, Miglio C, Peluso I, Petrosino T. (2011b) Modulation of plasma non enzymatic antioxidant capacity (NEAC) by plant foods: the role of polyphenols. *Curr Top Med Chem.*;11(14):1821-46.
- Shi Y, Evans JE, Rock KL. (2003) Molecular identification of a danger signal that alerts the immune system to dying cells. *Nature.*;425, 516-521.
- Sone T, Kuriyama S, Nakaya N, Hozawa A, Shimazu T, Nomura K, Rikimaru S, Tsuji I. (2011) Randomized controlled trial for an effect of catechin-enriched green tea consumption on

adiponectin and cardiovascular disease risk factors. *Food Nutr Res.*;55. doi: 10.3402/fnr.v55i0.8326.

Strazzullo P, Puig JG. (2007) Uric acid and oxidative stress: relative impact on cardiovascular risk? *Nutr Metab Cardiovasc Dis.*;17(6):409-14.

Sugihara K, Kitamura S, Yamada T, Ohta S, Yamashita K, Yasuda M, Fujii-Kuriyama Y. (2001) Aryl hydrocarbon receptor (AhR)-mediated induction of xanthine oxidase/xanthine dehydrogenase activity by 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Biochem Biophys Res Commun.*;281(5):1093-9.

Tausche AK, Jansen TL, Schröder HE, Bornstein SR, Aringer M, Müller-Ladner U. (2009) Gout--current diagnosis and treatment. *Dtsch Arztebl Int.*;106(34-35):549-55.

Tayar JH, Lopez-Olivo MA, Suarez-Almazor ME. (2012) Febuxostat for treating chronic gout. *Cochrane Database Syst Rev.*;11:CD008653.

Taylor EN, Curhan GC. (2006) Diet and fluid prescription in stone disease. *Kidney Int.* Sep;70(5):835-9.

Tremaroli V, Bäckhed F. (2012) Functional interactions between the gut microbiota and host metabolism. *Nature.*;489(7415):242-9.

Vaidyanathan JB, Walle T. (2001) Transport and metabolism of the tea flavonoid (-)-epicatechin by the human intestinal cell line Caco-2. *Pharm Res.*;18(10):1420-5.

Van Amelsvoort JM, Van Hof KH, Mathot JN, Mulder TP, Wiersma A, Tijburg LB. (2001) Plasma concentrations of individual tea catechins after a single oral dose in humans. *Xenobiotica.*;31(12):891-901.



van het Hof KH, de Boer HS, Wiseman SA, Lien N, Westrate JA, Tijburg LB. (1997)

Consumption of green or black tea does not increase resistance of low-density lipoprotein to oxidation in humans. *Am J Clin Nutr.*;66(5):1125-32.

Vischini G, Niscola P, Stefoni A, Farneti F. (2011) Increased plasma levels of tacrolimus after ingestion of green tea. *Am J Kidney Dis.*;58(2):329.

Wang CP, Wang X, Zhang X, Shi YW, Liu L, Kong LD. (2010) Morin improves urate excretion and kidney function through regulation of renal organic ion transporters in hyperuricemic mice. *J Pharm Pharm Sci.*;13(3):411-27.

Wang H, Wei Y, Kong X, Xu D. (2012) Effects of Urate-Lowering Therapy in Hyperuricemia on Slowing the Progression of Renal Function: A Meta-Analysis. *J Ren Nutr.*;Nov 3. pii: S1051-2276(12)00172-0.

Yin Z, Henry EC, Gasiewicz TA. (2009) (-)-Epigallocatechin-3-gallate is a novel Hsp90 inhibitor. *Biochemistry.*;48(2):336-45.

Yokozawa T, Rhyu DY, Cho EJ. (2004) (-)-Epicatechin 3-O-gallate ameliorates the damages related to peroxynitrite production by mechanisms distinct from those of other free radical inhibitors. *J Pharm Pharmacol.*;56(2):231-9.

Yu Z, Fong WP, Cheng CH. (2007) Morin (3,5,7,2',4'-pentahydroxyflavone) exhibits potent inhibitory actions on urate transport by the human urate anion transporter (hURAT1) expressed in human embryonic kidney cells. *Drug Metab Dispos.*;35(6):981-6.

Zhang ML, Gao YX, Wang X, Chang H, Huang GW. (2013) Serum uric acid and appropriate cutoff value for prediction of metabolic syndrome among Chinese adults. *J Clin Biochem Nutr.*;52(1):38-42.

Zoppini G, Targher G, Bonora E. (2011) The role of serum uric acid in cardiovascular disease in type 2 diabetic and non-diabetic subjects: a narrative review. *J Endocrinol Invest.*;34(11):881-6.

Table 1. RCT included in the meta-analyses

Reference	Subjects	Treatment (duration)	Baseline UA (reported effect)	Bias (Score range 0 - 1)
Bahorun 2010	CVD	600ml BT	(a) 4.4 mg/dl M ( $\leftrightarrow$ )	No food antioxidant intake in subjects
	n= 232, W 41%	(12 weeks)	(b) 6.1 mg/dl M ( $\leftrightarrow$ )	selection criteria,
	D 12%		(c) 8.4 mg/dl M ( $\downarrow$ )	No wash out/run-in period,
	BMI n.r.,		(d) 3.4 mg/dl W ( $\leftrightarrow$ )	No double blinding.
	Y 25-60		(e) 5.1 mg/dl W ( $\leftrightarrow$ )	(0.8)
			(f) 6.7 mg/dl W ( $\downarrow$ )	
Hsu 2008	Obeses	1.2g GTE	5.7 mg/dl W ( $\leftrightarrow$ )	No dietary record,
	n= 100, W 100%	(12 weeks)		No food antioxidant intake in subjects
	D 22%			selection criteria,
	BMI >26			No wash out/run-in period.
	Y 16-60			(0.79)
Hsu 2011	T2D	1.5g GTE	5.6 mg/dl M/W ( $\leftrightarrow$ )	No baseline comparability,
	n= 68, W 65%	(16 weeks)		No dietary record,

	D 15%			No food antioxidant intake in subjects
	BMI >26			selection criteria,
	Y 20-65			No wash out/run-in period.
				(0.49)
Princen 1998	Smokers	3.6 g GTE	4.5 mg/dl M/W (↔)	No double blinding,
	n= 64, W 50%	900 ml BT	4.4 mg/dl M/W (↔)	Supported by Unilever Research,
	D 8%	900 ml GT	4.4 mg/dl M/W (↔)	Products provided by T.J. Lipton.
	BMI < 28	(4 weeks)		(0.86)
	Y 22-46			
van het Hof 1997	Healthy	900ml BT	4.9 mg/dl M/W (↔)	No double blinding.
	n=45, W 46%	900ml GT	4.8 mg/dl M/W (↔)	(0.9)
	D 4%	(4 weeks)		
	BMI < 26			
	Y 37-52			

UA, uric acid; n, number of subjects (M: men, W: women); D: dropout (when not reported it has been calculated confronting the numbers in methods and in results); BMI: Body Mass Index ( $\text{kg/m}^2$ ); Y: years; CVD: Cardiovascular Disease; T2D: Type 2 Diabetes; BT: Black Tea; GT: Green Tea; GTE: Green Tea Extract.  $\leftrightarrow$ , no change;  $\downarrow$ , decrease. n.r, not reported.

**Table 2. Combined Effect Estimate on UA of studies sub-grouped for treatment and baseline levels.**

	<b>Studies/intervention</b>	<b>Fixed effect</b>	<b>Random effect</b>	<b>Quality effect</b>
	<b>(subjects)</b>	<b>MD (95%CI)</b>	<b>MD (95%CI)</b>	<b>MD (95%CI)</b>
		<b>p (I<sup>2</sup>%)</b>	<b>p (τ<sup>2</sup>)</b>	<b>p (Q%)</b>
<b>BT</b>	3/8 (266)	0.1186 (-0.0765 to 0.3136) 0.2337 (0%)	0.1186 (-0.0765 to 0.3136) 0.2337 (0)	0.1246 (-0.1056 to 0.3550) 0.2484 (15%)
<b>GT/GTE</b>	4/5 (206)	0.0351 (-0.1730 to 0.2433) 0.2433 (0%)	0.0351 (-0.1730 to 0.2433) 0.2433 (0)	0.0340 (-0.2131 to 0.2810) 0.7688 (15%)
<b>Baseline UA under cut off</b>	3/8 (224)	0.1078 (-0.0528 to 0.2684) 0.1884 (0%)	0.1078 (-0.0528 to 0.2684) 0.1884 (0)	0.1078 (0.0771 to 0.2926) 0.2188 (12%)
<b>Baseline UA over cut off</b>	3/5 (248)	-0.0239 (0.3311 to 0.2833) 0.8788 (0%)	-0.0239 (0.3311 to 0.2833) 0.8788 (0)	-0.0262 (-0.4416 to 0.3892) 0.885 (25%)

<b>Diseases or risk</b>	4/11 (427)	0.0602 (-0.1333 to	0.0602 (-0.1333 to	0.0634 (-0.1793 to
<b>factors</b>		0.2537)	0.2537)	0.3062)
		0.5419 (0%)	0.5419 (0%)	0.5648 (20%)

---

MD: mean difference, CI 95%: confidence intervals 95%. Cut off limits for UA baseline levels of 4.5 mg/dl for women, 6.1 mg/dl for men and 5.5 mg/dl for studies involving mixed populations.

Figure legends






Figure 1. Four-phase flow diagram of systematic review and meta-analysis.





**Figure 2.** Forest Plot of Fixed Effects estimate after black tea (BT), green tea (GT) and green tea extract (GTE) intervention (A). MD: mean difference. CI 95%: confidence intervals 95%. Trim-and-fill funnel plot (B). The black circles represent actual identified studies. MD, mean difference.