



Regulation of the intestinal tight junction by natural polyphenols: a mechanistic perspective

Guan Yang, Shima Bibi, Min Du, Takuya Suzuki & Mei-Jun Zhu

To cite this article: Guan Yang, Shima Bibi, Min Du, Takuya Suzuki & Mei-Jun Zhu (2016): Regulation of the intestinal tight junction by natural polyphenols: a mechanistic perspective, Critical Reviews in Food Science and Nutrition, DOI: [10.1080/10408398.2016.1152230](https://doi.org/10.1080/10408398.2016.1152230)

To link to this article: <http://dx.doi.org/10.1080/10408398.2016.1152230>



Accepted author version posted online: 23 Mar 2016.



Submit your article to this journal [↗](#)



View related articles [↗](#)



View Crossmark data [↗](#)

Regulation of the intestinal tight junction by natural polyphenols: a mechanistic perspective

Guan Yang¹, Shima Bibi¹, Min Du², Takuya Suzuki³, Mei-Jun Zhu^{1*}

¹*School of Food Science, Washington State University, Pullman, WA 99164, USA;* ²*Department of Animal Science, Washington State University, Pullman, WA 99164, USA;* ³*Department of Biofunctional Science and Technology, Hiroshima University, Higashi-Hiroshima, 739-8528, Japan*

(*): Corresponding author: Meijun Zhu, Ph.D., Associate Professor, School of Food Science, Washington State University, Pullman, WA 99163; Phone: (509) 335-4016; Fax: (509) 335-4815; E-mail: meijun.zhu@wsu.edu

Acknowledgements: We thank Mrs. Jeanene M. Deavila for her critical reading of this review paper. This activity was funded with Washington State University faculty seed grant and an Emerging Research Issues Internal Competitive Grant from the Washington State University, CAHNRS, ARC.

The authors declare no competing financial interest.

Abbreviations

AMPK, AMP-activated protein kinase; **AOM**, azoxymethane; **BTK**, Bruton's tyrosine kinase; **DAG**, diacylglycerol; **DSS**, dextran sodium sulfate; **ERK**, extracellular signal-regulated kinase; **GSE**, grape seed extract; **GSH-px**, glutathione peroxides; **IBD**, inflammatory bowel disease; **IFN- γ** , interferon gamma; **IKK**, I κ B kinase; **JAM**, junctional adhesion molecule; **IL**, interleukin; **JNK**, c-Jun N-terminal kinases; **MAPK**, mitogen-activated protein kinase; **MAPKAP**, MAPK activated protein; **MDCK**, Madin-Darby canine kidney; **MLCK**, Myosin light chain kinase; **MNK**, MAPK-interacting kinases; **NF- κ B**, nuclear factor kappa light-chain-enhancer of activated B cells; **PI3K**, phosphoinositide-3-kinase; **PKB**, protein kinase B; **PKC**, protein kinase C; **PLC**, phospholipase C; **PP2A**, protein phosphatase 2A; **PPAR- γ** , peroxisome proliferator activated receptor - γ ; **ROCK**, Rho-associated kinase; **ROS**, reactive oxygen species; **SOD**, superoxide dismutase; **STAT**, signal transducer and activator of transcription; **TEER**, the trans-epithelial electrical resistance; **TJ**, tight junction; **TK**, tyrosine kinase; **TLR**, Toll-like receptor; **TNF- α** , tumor necrosis factor- α ; **ZO**, zona occluden

Abstract

Impairment of the epithelial barrier function is closely linked to the pathogenesis of various gastrointestinal diseases, food allergies, type I diabetes and other systematic diseases. Plant-derived polyphenols are natural secondary metabolites and exert various physiological benefits including anti-inflammatory, anti-oxidative, anti-carcinogenic, and anti-aging effects. Recent studies also show the role of plant polyphenols in regulation of the intestinal barrier and prevention of intestinal inflammatory diseases. Here, we summarize the regulatory pathways and mediators linking polyphenols to their beneficial effects on tight junction and gut epithelial barrier functions, and provide useful information about using polyphenols as nutraceuticals for intestinal diseases.

Keywords: polyphenols; intestinal epithelium; barrier function; tight junction; signaling pathway

1. Introduction

Gut epithelium constitutes a single layer of tightly linked cells (Heath 2010). It is permeable to nutrients, minerals, water and selected antigens, while it fends off bacteria, viruses and antigenic materials. Impairment in the intestinal barrier integrity permits passage of normally excluded luminal substances, which elicits excessive immune response and hence inflammation (Suenaeert *et al.* 2002; Knight *et al.* 2007; Zeissig *et al.* 2007; Lee *et al.* 2010; Camilleri *et al.* 2012). Therefore, epithelial integrity and barrier function are critical for health (Hilsden *et al.* 1999; Groschwitz & Hogan 2009; Yu 2009; Camilleri *et al.* 2012; Vaarala 2012). To maintain integrity between epithelial cells, extensive tight junctions are formed, these regulate epithelial permeability (Groschwitz & Hogan 2009). A large number of tight junction proteins are involved in the formation of tight junction complexes, including intracellular proteins: zona occludens (ZO-1, ZO-2, and ZO-3), cingulin, 7H6 and ZA-1; and transmembrane proteins: occludin, claudin, junctional adhesion molecules and tricellulin (Laukoetter *et al.* 2006; Ulluwishewa *et al.* 2011; Suzuki 2013). Disturbances in production and formation of tight junction (TJ) complex negatively impact gut epithelial barrier function, which is involved in the pathogenesis of many diseases (Scaldaferri *et al.* 2012; Vaarala 2012).

The intestinal barrier integrity is dynamically regulated by a diverse array of environmental factors such as pathogens, luminal pro-inflammatory molecules, stresses and dietary factors (Suzuki 2013). For example, Enterohemorrhagic *Escherichia coli* (Philpott *et al.* 1998) and *Clostridium perfringens* (Singh *et al.* 2000) interact with the intestinal epithelial cells to disrupt

the TJ barrier. Environmental stresses, such as alcohol and acetaldehyde, might also disrupt the barrier integrity (Rao 1998; Keshavarzian *et al.* 2009). On the other hand, certain dietary factors exert protective effects on the intestinal barrier. Non-essential amino acids such as arginine and glutamine improve intestinal immunity (Yao *et al.* 2011; Ren *et al.* 2014a; Ren *et al.* 2014b; Ren *et al.* 2014c; Yin *et al.* 2014). Butyrate promotes the intestinal barrier function (Peng *et al.* 2009), and vitamin D enhances TJ formation and prevents dextran sodium sulfate (DSS)-induced epithelial disruption (Kong *et al.* 2008).

Polyphenols are bioactive compounds present abundantly in plant-derived foods. They are grossly classified into five groups based on their structures: phenolic acids, flavonoids, anthocyanins, stilbenes, and lignans (Tsao 2010). They are known for their anti-inflammatory and anti-oxidative activities (Gupta *et al.* 2014), as well as anti-diabetic, anti-carcinogenic, anti-muscle wasting and even anti-aging activities (Cao *et al.* 2008; Chis *et al.* 2009; Wang *et al.* 2009; Kiyici *et al.* 2010; Velmurugan *et al.* 2010b; Cuevas *et al.* 2011; Narita *et al.* 2011; Su & D'Souza 2011; Wang *et al.* 2014; Yang *et al.* 2015). Recent studies also show the beneficial effects of polyphenols on epithelial barrier function. Naringenin, a major polyphenol in citrus fruits, enhances TJ formation and barrier integrity in cultured intestinal Caco-2 cell monolayers (Noda *et al.* 2012). Naringenin also alleviates the DSS-induced colitis in BALB/c mice partially through improvement of the TJ barrier (Azuma *et al.* 2013). Quercetin (found in apples, onions, leafy green vegetables and tea) and kaempferol (found in tea, broccoli, cabbage, kale, beans, endive, leak, tomato, strawberries and grapes) regulate TJ protein expression and enhance intestinal Caco-2 cell monolayer barrier integrity (Suzuki & Hara 2009; Suzuki *et al.* 2011a; Noda *et al.* 2014). In Caco-2 cells, curcumin, which is the major anti-inflammatory polyphenol

in turmeric, ameliorates barrier disruption induced by hydrogen peroxide (H₂O₂) (Wang *et al.* 2012b) or inflammatory cytokines such as tumor necrosis factor α (TNF- α) (Ye *et al.* 2006) and interleukin (IL)-1 β production (Al-Sadi & Ma 2007). Similarly, the green tea flavonoid, epigallocatechin gallate, prevents interferon gamma (IFN- γ) (Watson *et al.* 2004), and *Staphylococcus* enterotoxin B (Watson *et al.* 2005) provoked intestinal T84 cell barrier dysfunction. Grape seed extract (GSE) that contains a mixture of polyphenols reduces the severity of selected disease markers in the proximal colon of DSS-induced colitis in rats (Cheah *et al.* 2013), and alleviates inflammatory response in TNBS-induced colitis in rats (Wang *et al.* 2011b) and IL-10-deficient mice (Wang *et al.* 2013; Bibi *et al.* 2016). GSE also increases occludin and ZO-1 expression in the gut epithelium of rats (Song *et al.* 2011a; Goodrich *et al.* 2012). Recently, our group reported that GSE supplementation ameliorates IBD indices in IL-10-deficient mice, which are associated with decreased inflammatory responses and reduced expression of claudin-2, a pore forming TJ protein, in colonic tissue (Wang *et al.* 2013). In addition, resveratrol, a polyphenol extracted from grape seed and skin, promotes the mRNA expression of tight junction proteins (Etxeberria *et al.* 2015). Currently, the exact mechanisms linking dietary polyphenols to gut epithelial health has not been established. Based on available literature, we briefly summarize possible regulatory pathways and mediators linking polyphenols to their beneficial effects on TJ protein expression and complex formation, and intestinal barrier integrity.

2. Polyphenols and signaling pathways regulating TJ formation

2.1 Polyphenols inhibit nuclear factor- κ B (NF- κ B) signaling

One important target of polyphenols' action *in vivo* is the inflammatory NF- κ B proteins, including p65 (Rel A), RelB, c-Rel, p50/p105 (NF- κ B1), and p52/p100 (NF- κ B2). In unstimulated cells, NF- κ B is sequestered in the cytoplasm as an inactive non-DNA-binding complex by partnering with inhibitor κ B proteins (I κ Bs), comprising I κ B α , I κ B β , I κ B γ , I κ B ϵ , Bcl-3, and precursors p100 and p105 (Caamano & Hunter 2002; Senftleben & Karin 2002; Santangelo *et al.* 2007). I κ B proteins undergo phosphorylation by the I κ B kinase (IKK). IKK contains two catalytic subunits, IKK α and IKK β , and the regulatory protein termed NF- κ B essential modifier (NEMO, also known as IKK γ) (Caamano & Hunter 2002; Senftleben & Karin 2002; Santangelo *et al.* 2007). Upon stimulation, IKK phosphorylates I κ B resulting in its degradation from the NF- κ B complex. The free NF- κ B can enter the nucleus to function as transcription factors that initiate the expression of specific genes, such as pro-inflammatory cytokines.

Polyphenols exert their anti-inflammatory activity by targeting different NF- κ B members (Velmurugan *et al.* 2010a; Biasi *et al.* 2011; Song *et al.* 2011b; Izzi *et al.* 2012; Wang *et al.* 2013). For example, dietary administration of 0.25 or 0.5% grape seed extract (GSE) at pre- and post-azoxymethane (AOM) treatment reduced ($P<0.001$) the AOM-induced NF- κ B (p65)-positive cells (Velmurugan *et al.* 2010a). Polyphenols also affect the NF- κ B pathway by inhibiting the IKK phosphorylation and/or preventing proteasomal degradation of I κ B. For instance, curcumin prevents phosphorylation of I κ B by inhibiting the activity of IKK (Joe *et al.* 2004). Quercetin also decreases the NF- κ B activity through inhibition of IKK phosphorylation and degradation of I κ B (Dias *et al.* 2005). It is also known that anthocyanins from blueberry,

blackberry, and blackcurrant attenuated lipopolysaccharide (LPS)-induced NF- κ B p65 translocation to the nucleus in macrophages (Lee *et al.* 2014)

Inflammatory cytokines, such as tumor necrosis factor- α (TNF- α), activate NF- κ B signaling and impair epithelial barrier function by TJ disassembly (Kisseleva *et al.* 2006; Amasheh *et al.* 2010; Petecchia *et al.* 2012; Rotches-Ribalta *et al.* 2012; Wang *et al.* 2012a; Zhang & Li 2012). Polyphenols improve barrier function partially by antagonizing the NF- κ B pathway induced by inflammatory cytokines. For example, berberine (found in European barberry, goldseal, goldthread, Oregon Grape, phellodendron and tree turmeric) supplementation ameliorated TNF- α induced claudin-1 disassembly and up-regulated claudin-2 expression in HT-29/B6 human colon monolayers. Furthermore, this effect is mimicked by BAY11-7082, a specific blocker of NF- κ B pathway, indicating the beneficial effects of berberine are mediated through the NF- κ B pathway (Amasheh *et al.* 2010). The role of NF- κ B in the impairment of the intestinal barrier was further supported by a recent study showing treatment with pyrrolidine dithiocarbamate (PDTC), which is an inhibitor of NF- κ B, upregulated the occludin and ZO-1 expression in DSS-induced colitis model (Yin *et al.* 2015b). Our previous study found that GSE decreases the protein expression of claudin-2 and increases the expression of claudin-1 in the colon of IL-10-deficient mice through down-regulation of NF- κ B signaling (Wang *et al.* 2013). More recently, Piegholdt *et al.* reported that the isoflavones, biochanin A (found in red clover, alfalfa sprouts, soy, peanuts, chick peas and other legumes) and prunetin (found in *Prunus yedoensis* (Yoshino cherry), improve transepithelial electrical resistance (TEER) and prevent TNF- α induced disruption of barrier integrity in Caco-2 cells (Piegholdt *et al.* 2014). These changes were associated with inhibition of the NF- κ B pathway (Piegholdt *et al.* 2014). Anthocyanin rich extract of raspberries

significantly attenuates inflammation via inhibition of the NF- κ B signaling, **demonstrated by** improved colonic histological architecture in **a DDS-induced colitis mouse** model (Li *et al.* 2014).

2.2 Polyphenols modulate the activity of multiple protein kinases

2.2.1 Mitogen-activated protein kinases (MAPK)

MAPK are a family of Ser/Thr specific protein kinases, which play fundamental roles in a wide range of cellular processes and are often **deregulated** in disease states. Mammals express at least four distinctly regulated MAPKs: extracellular signal-regulated kinases (ERKs 1/2), p38MAPKs (α , β , δ , and γ), c-Jun NH2-terminal kinases (JNKs-1/2/3), and extracellular signal-regulated kinase-5 (ERK5) or BMK (Santangelo *et al.* 2007). Generally, ERK1/2 **is activated** by mitogens and growth factors, and does not affect the inflammatory responses directly, while JNK cascades are usually triggered by stress and inflammation. The p38 plays an active signaling role in cellular responses to stress and inflammation (Kyriakis & Avruch 2001; Boncoeur *et al.* 2008). In addition, p38 directly binds and phosphorylates transcription factors, including NF- κ B and CDX-2, to induce inflammatory gene transcription (Hollenbach *et al.* 2004; Schindler *et al.* 2007; Broom *et al.* 2009). Members of the JNK family also play crucial roles in inflammation (Himes *et al.* 2006). JNK activates c-Jun, which forms hetero- or homodimers with activator protein-1 (AP-1) transcription factor, inducing the expression of inflammatory cytokines (Schmeck *et al.* 2006).

Polyphenols improve the barrier function through modulating MAPK signaling pathways (Morin 2005; Chuenkitiyanon *et al.* 2010; Shin *et al.* 2011). Anthocyanins are widely distributed polyphenolic compounds that can be found abundantly in fruits and vegetables. Anthocyanin rich

extract of raspberries prevents DSS-induced histological damage of the colon architecture by inhibiting MAPKs (Li *et al.* 2014). Quercetin, at 10 $\mu\text{mol/L}$, prevents the loss of ZO-1 and occludin induced by H_2O_2 through reducing p38 MAPK phosphorylation (Chuenkitiyanon *et al.* 2010). On the other hand, ERK 1/2 interacts directly with the C-terminal region of occludin and plays a crucial role in preventing EGF protection of TJs disrupted by H_2O_2 (Basuroy *et al.* 2006), acetaldehyde (Samak *et al.* 2011), and TNF- α (Piegholdt *et al.* 2014). Biochanin A and pruenetin were identified as potent inhibitors of ERK activation, which accounts for the barrier-improving ability of these isoflavones (Piegholdt *et al.* 2014). In HCT-116 human colon carcinoma cells, anthocyanin (45 $\mu\text{g/ml}$) treatment increases TEER, and decreases the expression of claudin-1, claudin-3 and claudin-4 (Shin *et al.* 2011), which is associated with p38 activation. Inhibition of the p38 MAPK attenuates the effects of anthocyanins on the expression of claudins, showing the importance of p38 signaling in mediating the beneficial effects of anthocyanins on epithelial barrier function (Shin *et al.* 2011). In short, these data suggest the mediatory roles of MAPK pathways in linking polyphenol compounds to the improvement of barrier function.

2.2.2 Phosphoinositide-3-Kinases (PI3K)/Akt

PI3K participates in numerous biological processes, including immune cell growth, differentiation, survival, proliferation, migration, and metabolism. In the immune system, impaired PI3K signaling leads to immunodeficiency, whereas unrestrained PI3K signaling contributes to autoimmunity and inflammation (Okkenhaug & Vanhaesebroeck 2003). By direct phosphorylation of phosphatidylinositol-4,5-bisphosphate ($\text{PtdIns}(4,5)\text{P}_2$), PI3K generates phosphatidylinositol-3,4,5-triphosphate ($\text{PtdIns}(3,4,5)\text{P}_3$), which in turn contributes to the

activation of various downstream targets, including Ser/Thr-specific protein kinase Akt (also known as protein kinase B).

PI3K signaling pathway regulates epithelial barrier function, primarily through regulating TJ expression (Li & Neu 2009; Mankertz *et al.* 2009; Suzuki *et al.* 2011b; Rosenthal *et al.* 2012). In addition, PI3K also regulates TJ assembly. A previous study showed that the PI3K is activated by oxidative stress, which results in increased intestinal permeability, while inhibition of PI3K reduces permeability (Sheth *et al.* 2003). Glutamine deprivation decreases the TEER level and increases the intestinal permeability (Wang *et al.* 2015), which was reversed by inhibition of PI3K (Li & Neu 2009). Moreover, the PI3K pathway is responsible for the up-regulation of claudin-2 expression induced by TNF- α (Mankertz *et al.* 2009; Amasheh *et al.* 2010), and IL-6 (Suzuki *et al.* 2011b). Limited evidence points to the PI3K/Akt signaling in linking polyphenols to the epithelial barrier function. The polyphenols, naringenin and quercetin have similarities in structure to the synthetic PI3K inhibitors and hence inhibit PI3K activity (Walker *et al.* 2000; Harmon & Patel 2003). In another *in vitro* study in HT-29/B6 human colon monolayers, berberine inhibits TNF- α induced activation of the PI3K/Akt pathway and reduces the claudin-2 protein expression (Amasheh *et al.* 2010). These studies indicate that polyphenols improve the barrier integrity partly through inhibition of PI3K/Akt pathway.

2.2.3. Protein kinase C (PKC)

The PKC family of Ser/Thr kinases regulates a number of fundamental biological processes in epithelial cells including barrier function (Suzuki *et al.* 2008; Suzuki *et al.* 2009; Suzuki & Hara 2009). Intestinal epithelial cells express a range of PKC isoforms involved in various

pathways, **i.e.** conventional (cPKC) isozymes (α , β I, β II, and γ), novel (nPKC) isozymes (δ , ϵ , η , μ , and θ), and atypical (aPKC) isozymes (ζ and ι/λ) (Song *et al.* 2001; Farhadi *et al.* 2006). These isoforms vary in their sensitivity to activators and cofactors, thus playing different roles in the regulation of intestinal barrier function (Farhadi *et al.* 2006). The role played by **the** PKC family on TJs is complex. Previously, it was reported that pretreatment with PKC selective antagonists, prevents the activation of down-stream signaling induced by Toll-like receptor 2 (TLR2), which greatly enhances TEER in Caco-2 and HT-29 monolayers through recruiting ZO-1 (Cario *et al.* 2004). The activation of PKC also induces the activation of MAP kinases, which results in the tight junction disruption in the corneal epithelial cells (Wang *et al.* 2004). Moreover, over-expression of nPKC δ on renal LLC-PK1 cells **has a** negative effect on the TJ barrier (Mullin *et al.* 1998), and **inhibition of nPKC δ by** quercetin promotes actin cytoskeletal association and expression of TJ proteins in Caco-2 **cells** (Suzuki & Hara 2009). Besides the intestinal barrier integrity, polyphenols also play a role in improving the blood-barrier function. Green tea polyphenols prevent **blood-brain barrier damage, exerting neuroprotective effects by** up-regulating the expression of claudin-5, occludin and ZO-1 as well as down-regulating PKC α expression (Liu *et al.* 2013). All these indicate that polyphenols inhibit PKC to improve epithelial barrier function.

2.2.4. Tyrosine kinases

Tyrosine kinases are key mediators of important cellular signaling pathways controlling cell proliferation and growth via activation of tyrosine kinase receptors (Lemmon & Schlessinger 2010; Segaliny *et al.* 2015). **This activation is considered** the starting point for many cellular

signaling pathways by recruiting enzymatic effectors (PLC γ , PI3K, etc.) and serving as the upstream of the signaling pathways including MAPK, PI3K, PLC γ , etc. (Lemmon & Schlessinger 2010; Segaliny *et al.* 2015). Accumulating evidences suggest the significant contribution of tyrosine kinase to TJ integrity (Rao *et al.* 1997; Atkinson & Rao 2001; Seth *et al.* 2004; Elias *et al.* 2009; Amasheh *et al.* 2010; Spalinger *et al.* 2013). Specifically, H₂O₂ administration induces tyrosine phosphorylation of several proteins (Rao *et al.* 1997) and acetaldehyde disrupts TJs through inducing phosphorylation of several proteins including ZO-1, E-cadherin and β -catenin at their tyrosine residues, which increases the paracellular permeability in Caco-2 and T84 cell monolayers (Atkinson & Rao 2001). In patients with active Crohn's disease, mRNA and protein levels of protein tyrosine phosphatase, nonreceptor type 22, were lower compared with the controls, suggesting the importance of the tyrosine phosphorylation in maintaining TJ integrity (Spalinger *et al.* 2013). Polyphenols have been reported to regulate the activity of tyrosine kinases in various studies. The administration of genistein, a soybean polyphenol, prevents acetaldehyde-induced permeability and TJ disruption by blocking tyrosine phosphorylation (Atkinson & Rao 2001). Moreover, berberine prevents TNF- α induced claudin-1 disassembly and claudin-2 up-regulation partially via down-regulation of tyrosine kinase pathway (Amasheh *et al.* 2010). In addition, biochanin A, and prunetin improved the barrier integrity partly through the down-regulation of ZO-1 tyrosine phosphorylation further indicating that polyphenols could exert beneficial effects on TJ integrity via inhibition of tyrosine kinase activity and tyrosine phosphorylation (Piegholdt *et al.* 2014).

2.2.5. Myosin light chain kinase (MLCK)

Myosin light chain kinases (MLCKs) are a family of soluble protein kinases that function principally to phosphorylate the regulatory myosin light chain (MLC-2) and thereby induce actomyosin contraction (Rigor *et al.* 2013). A number of studies have been conducted to understand the role of MLCK activity under disease conditions. It has been shown that increased MLCK activity decreases colonic epithelial barrier function (Ferrier *et al.* 2003). Inflammatory cytokines TNF- α and IFN- γ induced barrier loss, and increased MLCK phosphorylation. The TJ barrier was restored by the inhibition of MLCK *in vitro* (Zolotarevsky *et al.* 2002). MLCK inhibition also resulted in reduced ZO-1 exchange between tight junction and cytosolic pools, which suggests MLCK alters barrier regulation via a ZO-1 dependent process (Yu *et al.* 2010). Pharmacological inhibition of MLCK alleviates TJ barrier dysfunction (Scott *et al.* 2002; Ma *et al.* 2005; Gu *et al.* 2011). It is proposed that polyphenols could improve TJ integrity through down-regulation of MLCK activity (Gu *et al.* 2011; Cao *et al.* 2013). In a mouse model of endotoxemia, pretreatment with berberine for 7 days reduces small intestine permeability and prevents TJ disruption caused by LPS, concomitant with the decreased MLCK activity (Gu *et al.* 2011). Furthermore, berberine ameliorates IFN- γ and TNF- α induced intestinal epithelial barrier dysfunction in Caco-2 cell monolayers through suppression of MLCK-MLC2 signaling pathway (Cao *et al.* 2013). These studies show that, polyphenols inhibit MLCK to improve epithelial barrier function.

2.2.6 AMP-activated protein kinase (AMPK)

AMP-activated protein kinase (AMPK) is a Ser/Thr protein kinase that plays an important role in maintaining cellular energy balance (Gruzman *et al.* 2009). AMPK is activated by a

number of dietary factors including polyphenol compounds (Peng *et al.* 2009; Wang *et al.* 2011a; Castillo-Pichardo & Dharmawardhane 2012; Yang *et al.* 2012; Mukai *et al.* 2013; Zhao *et al.* 2015). For example, butyrate (Peng *et al.* 2009) and forskolin (Egawa *et al.* 2008) enhance the intestinal barrier via activation of AMPK *in vitro*. AMPK is activated during calcium-induced TJ assembly in Madin-Darby canine kidney (MDCK) cells, and up-regulation of AMPK activity facilitates TJ assembly (Zhang *et al.* 2006). Attenuation of AMPK is responsible for the impaired epithelial barrier function induced by IFN- γ (Scharl *et al.* 2009). All of the above evidences indicate the critical roles of AMPK in TJ integrity. Polyphenols also affect intestinal barrier function through AMPK modulation. It has recently been shown that theaflavins-3'-O-gallate significantly increased the TJ protein (occludin, claudin-1, and ZO-1) expression as well as the phosphorylation of AMPK (Park *et al.* 2015). Inhibition of AMPK completely abolished the beneficial effects of theaflavins-3'-O-gallate on the intestinal barrier, suggesting these effects were AMPK dependent (Park *et al.* 2015). 6-Gingerol, a major pungent phenolic component in ginger, has also been demonstrated to protect the DSS-induced intestinal inflammatory disorder through the activation of AMPK (Chang & Kuo 2015). However, our recent study showed that AMPK activity was heightened in IL-10-deficient mice, where severe inflammation occurs due to the absence of a key anti-inflammatory cytokine. GSE supplementation restored AMPK to levels that are associated with the improved barrier integrity (Yang *et al.* 2014). AMPK, as an intracellular energy sensor and regulator, contributes significantly to the maintenance of the epithelial barrier, which is energy-dependent (Novak & Mollen 2015). Hallmarks of mitochondrial dysfunction, including oxidative stress and impaired ATP production have been characterized in the IBD patients (Novak & Mollen 2015). Thus, this abnormal activation of

AMPK, in IL-10-deficient mice, could be due to the impairment in energetic metabolism elicited by acute inflammation, which warrants further research.

2.3 Polyphenols regulate the activity of key enzymes

2.3.1. Protein phosphatase 2A (PP2A)

PP2A comprises a family of Ser/Thr phosphatases. It has a broad range of cellular functions such as regulating cell survival (Strack *et al.* 2004), facilitating DNA repair (Chowdhury *et al.* 2005) and others. PP2A is composed of three subunits, scaffold subunit A, regulatory subunit B and catalytic subunit C. The functional diversity of PP2A is mediated by scaffold and regulatory subunits, which interact with a wide range of proteins (Janssens & Goris 2001). These interactions facilitate the cross-talk of PP2A with several cell signaling pathways including I κ B kinases, MAPK, Akt and PKC (Sontag *et al.* 1997; Van Kanegan *et al.* 2005; Grethe & Porn-Ares 2006). Interestingly, studies show that PP2A interacts with TJ proteins thus negatively regulating TJ assembly and function (Nunbhakdi-Craig *et al.* 2002; Seth *et al.* 2007; Sheth *et al.* 2009; Dunagan *et al.* 2012). Enhanced PP2A activity induces dephosphorylation of membrane TJ proteins, ZO-1, occludin, and claudin-1 at their serine residues, with an increasing paracellular permeability in MDCK cells (Nunbhakdi-Craig *et al.* 2002). This dephosphorylation is important in promoting TJ opening (Farshori & Kachar 1999; Clarke *et al.* 2000). Inhibition of PP2A and knockdown of PP2A-C α (the catalytic subunit of PP2A) attenuate the impairment of barrier function caused by H₂O₂ (Sheth *et al.* 2009). Polyphenols inhibit PP2A activity (Chen *et al.* 2013; Kiss *et al.* 2013), which improves TJ formation.

2.3.2. Rho family of small GTPases

The Rho family of small GTPases, consisting of Rho, Rac, and Cdc42, play an important role in the regulation of epithelial structure and function (Wittchen *et al.* 2003). Its downstream effector, Rho-associated kinase (ROCK) is a Ser/Thr kinase, which is important in regulating TJ permeability (Walsh *et al.* 2001). In cultured T84 cell monolayers, ROCK inhibition prevents proper localization of TJ proteins during TJ assembly, enhancing paracellular permeability (Walsh *et al.* 2001). ROCKs could also alter the cell-cell adhesion and actin cytoskeleton organization by phosphorylating MLC (Grassie *et al.* 2012). Moreover, a previous study has demonstrated that Rho GTP exchange factor ARHGEF11, mediates RhoA – MLC2 signaling pathways at cell-cell junctions, functioning in cooperation with ZO-1, therefore regulating the paracellular barrier and the organization of the apical junctional complex and peri-junctional actomyosin ring of epithelial cells (Itoh *et al.* 2012). Because polyphenols suppress Rho activity (Higashi *et al.* 2005; McLaughlin *et al.* 2006), polyphenols may decrease the TJ permeability through attenuating the Rho activity.

2.4 Polyphenols reduce reactive oxygen species (ROS)

The maintenance of redox homeostasis is of critical importance for normal cellular processes and organ function. This homeostasis is tightly regulated by the pro-oxidative and anti-oxidative systems. Oxidative stress is considered to be an imbalance between the production of ROS and their elimination through anti-oxidative mechanisms (Pi *et al.* 2010; Reuter *et al.* 2010; Kanninen *et al.* 2011). ROS production correlates with inflammation and alterations in several signal pathways including MAPK and NF- κ B (Morgan & Liu 2011; Liu *et al.* 2012; Park *et al.* 2012). H₂O₂ induced oxidative stress impairs TJs in the intestinal epithelium (Sheth *et al.* 2003;

Chuenkitiyanon *et al.* 2010; Duan *et al.* 2016). Dietary glutamate and aspartate are beneficial in alleviating the oxidative stress induced by diquat (Yin *et al.* 2015a) and H₂O₂ (Duan *et al.* 2016). Plant-derived polyphenols also exhibit anti-inflammatory and anti-oxidative effects on various disease models (Edgecombe *et al.* 2000; Cao *et al.* 2008; Chis *et al.* 2009; Cuevas *et al.* 2011; Nadour *et al.* 2012; Rajamurugan *et al.* 2012; Periasamy & Alshatwi 2013). Oregano essential oil, which is enriched with phenolic compounds, reduces the activation of superoxide dismutase (SOD) and glutathione peroxide (GSH-Px), and improves dysfunction of intestinal epithelial barrier in rats (Wei *et al.* 2015). These data suggest that polyphenols may improve the barrier function through the down-regulation of ROS.

3. Implications and Conclusions

Basic and clinical studies have shown that increased intestinal permeability is associated with various diseases, such as IBD, celiac disease, irritable bowel syndrome, and type I diabetes (Suzuki 2013). Dietary polyphenol supplementation might provide a potential therapy for these diseases. Interest in studying the mechanisms connecting polyphenol compounds and barrier integrity has grown in recent years (Table 1). However, in most cases, available studies were performed in cultured cells. Moreover, it is still unclear whether the polyphenols directly exert effects on epithelial development or mediates the epithelial formation through the regulation of TJs. Thus, further *in vivo* and mechanistic studies are needed, which will likely lead to the development of polyphenolic compounds as therapeutic and preventive approaches against the barrier-defect related diseases.

Additionally, existing data suggests that polyphenols exert their beneficial effects in TJ regulation via diversified pathways, which crosstalk with each other. For example, in intestinal cells, PKC activation is accompanied by progressively decreasing MLC2 phosphorylation. Meanwhile, an acute increase in MLCK phosphorylation concurrent with PKC activation leads to the decrease in MLC2 phosphorylation and improvement of barrier integrity (Turner *et al.* 1999). Furthermore, other signaling pathways and mediators may also interact with pathways, discussed in this review, which regulate TJ integrity and their regulatory roles in polyphenols' mediated TJ barrier function. Future investigations are required to elucidate the precise mechanisms involving these signaling pathways and their interactions which underlying the polyphenol-mediated protective effects on intestinal TJ formation and barrier function. Knowing these pathways and their cross-talks may also help to further understand the clinical use of the polyphenolic compounds in the prevention and treatment of diseases associated with intestinal barrier defects.

4. References

- Al-Sadi R.M. & Ma T.Y. (2007) IL-1beta causes an increase in intestinal epithelial tight junction permeability. *J Immunol* **178**, 4641-9.
- Amasheh M., Fromm A., Krug S.M., Amasheh S., Andres S., Zeitz M., Fromm M. & Schulzke J.D. (2010) TNFalpha-induced and berberine-antagonized tight junction barrier impairment via tyrosine kinase, Akt and NFkappaB signaling. *J Cell Sci* **123**, 4145-55.

- Atkinson K.J. & Rao R.K. (2001) Role of protein tyrosine phosphorylation in acetaldehyde-induced disruption of epithelial tight junctions. *Am J Physiol Gastrointest Liver Physiol* **280**, G1280-8.
- Azuma T., Shigeshiro M., Kodama M., Tanabe S. & Suzuki T. (2013) Supplemental naringenin prevents intestinal barrier defects and inflammation in colitic mice. *J Nutr* **143**, 827-34.
- Basuroy S., Seth A., Elias B., Naren A.P. & Rao R. (2006) MAPK interacts with occludin and mediates EGF-induced prevention of tight junction disruption by hydrogen peroxide. *Biochem J* **393**, 69-77.
- Biasi F., Astegiano M., Maina M., Leonarduzzi G. & Poli G. (2011) Polyphenol supplementation as a complementary medicinal approach to treating inflammatory bowel disease. *Curr Med Chem* **18**, 4851-65.
- Bibi S., Kang Y., Yang G. & Zhu M.-J. (2016) Grape seed extract improves small intestinal health through suppressing inflammation and regulating alkaline phosphatase in IL-10-deficient mice. *J Funct Foods* **20**, 245-52.
- Boncoeur E., Criq V.S., Bonvin E., Roque T., Henrion-Caude A., Gruenert D.C., Clement A., Jacquot J. & Tabary O. (2008) Oxidative stress induces extracellular signal-regulated kinase 1/2 mitogen-activated protein kinase in cystic fibrosis lung epithelial cells: Potential mechanism for excessive IL-8 expression. *Int J Biochem Cell Biol* **40**, 432-46.
- Broom O.J., Widjaya B., Troelsen J., Olsen J. & Nielsen O.H. (2009) Mitogen activated protein kinases: a role in inflammatory bowel disease? *Clin Exp Immunol* **158**, 272-80.
- Caamano J. & Hunter C.A. (2002) NF-kappaB family of transcription factors: central regulators of innate and adaptive immune functions. *Clin Microbiol Rev* **15**, 414-29.

- Camilleri M., Madsen K., Spiller R., Greenwood-Van Meerveld B. & Verne G.N. (2012) Intestinal barrier function in health and gastrointestinal disease. *Neurogastroenterol Motil* **24**, 503-12.
- Cao H., Urban J.F., Jr. & Anderson R.A. (2008) Cinnamon polyphenol extract affects immune responses by regulating anti- and proinflammatory and glucose transporter gene expression in mouse macrophages. *J Nutr* **138**, 833-40.
- Cao M., Wang P., Sun C., He W. & Wang F. (2013) Amelioration of IFN-gamma and TNF-alpha-induced intestinal epithelial barrier dysfunction by berberine via suppression of MLCK-MLC phosphorylation signaling pathway. *PLoS One* **8**, e61944.
- Cario E., Gerken G. & Podolsky D.K. (2004) Toll-like receptor 2 enhances ZO-1-associated intestinal epithelial barrier integrity via protein kinase C. *Gastroenterology* **127**, 224-38.
- Castillo-Pichardo L. & Dharmawardhane S.F. (2012) Grape polyphenols inhibit akt/mammalian target of rapamycin signaling and potentiate the effects of gefitinib in breast cancer. *Nutr Cancer* **64**, 1058-69.
- Chang K.W. & Kuo C.Y. (2015) 6-Gingerol modulates proinflammatory responses in dextran sodium sulfate (DSS)-treated Caco-2 cells and experimental colitis in mice through adenosine monophosphate-activated protein kinase (AMPK) activation. *Food Funct* **6**, 3334-41.
- Cheah K.Y., Bastian S.E., Acott T.M., Abimosleh S.M., Lymn K.A. & Howarth G.S. (2013) Grape seed extract reduces the severity of selected disease markers in the proximal colon of dextran sulphate sodium-induced colitis in rats. *Dig Dis Sci* **58**, 970-7.
- Chen W., Wang Z., Jiang C. & Ding Y. (2013) PP2A-Mediated Anticancer Therapy. *Gastroenterol Res Pract* **2013**, 675429.

- Chis I.C., Ungureanu M.I., Marton A., Simedrea R., Muresan A., Postescu I.D. & Decea N. (2009) Antioxidant effects of a grape seed extract in a rat model of diabetes mellitus. *Diab Vasc Dis Res* **6**, 200-4.
- Chowdhury D., Keogh M.C., Ishii H., Peterson C.L., Buratowski S. & Lieberman J. (2005) gamma-H2AX dephosphorylation by protein phosphatase 2A facilitates DNA double-strand break repair. *Mol Cell* **20**, 801-9.
- Chuenkitiyanon S., Pengsuparp T. & Jianmongkol S. (2010) Protective effect of quercetin on hydrogen peroxide-induced tight junction disruption. *Int J Toxicol* **29**, 418-24.
- Clarke H., Soler A.P. & Mullin J.M. (2000) Protein kinase C activation leads to dephosphorylation of occludin and tight junction permeability increase in LLC-PK1 epithelial cell sheets. *J Cell Sci* **113** (Pt 18), 3187-96.
- Cuevas V.M., Calzado Y.R., Guerra Y.P., Yera A.O., Despaigne S.J., Ferreiro R.M. & Quintana D.C. (2011) Effects of grape seed extract, vitamin C, and vitamin e on ethanol- and aspirin-induced ulcers. *Adv Pharmacol Sci* **2011**, 740687.
- Dias A.S., Porawski M., Alonso M., Marroni N., Collado P.S. & Gonzalez-Gallego J. (2005) Quercetin decreases oxidative stress, NF-kappaB activation, and iNOS overexpression in liver of streptozotocin-induced diabetic rats. *J Nutr* **135**, 2299-304.
- Duan J., Yin J., Ren W., Liu T., Cui Z., Huang X., Wu L., Kim S.W., Liu G., Wu X., Wu G., Li T. & Yin Y. (2016) Dietary supplementation with L-glutamate and L-aspartate alleviates oxidative stress in weaned piglets challenged with hydrogen peroxide. *Amino Acids* **48**, 53-64.

- Dunagan M., Chaudhry K., Samak G. & Rao R.K. (2012) Acetaldehyde disrupts tight junctions in Caco-2 cell monolayers by a protein phosphatase 2A-dependent mechanism. *Am J Physiol Gastrointest Liver Physiol* **303**, G1356-64.
- Edgecombe S.C., Stretch G.L. & Hayball P.J. (2000) Oleuropein, an antioxidant polyphenol from olive oil, is poorly absorbed from isolated perfused rat intestine. *J Nutr* **130**, 2996-3002.
- Egawa M., Kamata H., Kushiya A., Sakoda H., Fujishiro M., Horike N., Yoneda M., Nakatsu Y., Ying G., Jun Z., Tsuchiya Y., Takata K., Kurihara H. & Asano T. (2008) Long-term forskolin stimulation induces AMPK activation and thereby enhances tight junction formation in human placental trophoblast BeWo cells. *Placenta* **29**, 1003-8.
- Elias B.C., Suzuki T., Seth A., Giorgianni F., Kale G., Shen L., Turner J.R., Naren A., Desiderio D.M. & Rao R. (2009) Phosphorylation of Tyr-398 and Tyr-402 in occludin prevents its interaction with ZO-1 and destabilizes its assembly at the tight junctions. *J Biol Chem* **284**, 1559-69.
- Etxeberria U., Arias N., Boque N., Macarulla M.T., Portillo M.P., Martinez J.A. & Milagro F.I. (2015) Reshaping faecal gut microbiota composition by the intake of trans-resveratrol and quercetin in high-fat sucrose diet-fed rats. *J Nutr Biochem* **26**, 651-60.
- Farhadi A., Keshavarzian A., Ranjbaran Z., Fields J.Z. & Banan A. (2006) The role of protein kinase C isoforms in modulating injury and repair of the intestinal barrier. *J Pharmacol Exp Ther* **316**, 1-7.
- Farshori P. & Kachar B. (1999) Redistribution and phosphorylation of occludin during opening and resealing of tight junctions in cultured epithelial cells. *J Membr Biol* **170**, 147-56.

- Ferrier L., Mazelin L., Cenac N., Desreumaux P., Janin A., Emilie D., Colombel J.F., Garcia-Villar R., Fioramonti J. & Bueno L. (2003) Stress-induced disruption of colonic epithelial barrier: role of interferon-gamma and myosin light chain kinase in mice. *Gastroenterology* **125**, 795-804.
- Goodrich K.M., Fundaro G., Griffin L.E., Grant A., Hulver M.W., Ponder M.A. & Neilson A.P. (2012) Chronic administration of dietary grape seed extract increases colonic expression of gut tight junction protein occludin and reduces fecal calprotectin: a secondary analysis of healthy Wistar Furth rats. *Nutr Res* **32**, 787-94.
- Grassie M.E., Sutherland C., Ulke-Lemee A., Chappellaz M., Kiss E., Walsh M.P. & Macdonald J.A. (2012) Cross-talk between Rho-associated Kinase and Cyclic Nucleotide-dependent Kinase Signaling Pathways in the Regulation of Smooth Muscle Myosin Light Chain Phosphatase. *J Biol Chem* **287**, 36356-69.
- Grethe S. & Porn-Ares M.I. (2006) p38 MAPK regulates phosphorylation of Bad via PP2A-dependent suppression of the MEK1/2-ERK1/2 survival pathway in TNF-alpha induced endothelial apoptosis. *Cell Signal* **18**, 531-40.
- Groschwitz K.R. & Hogan S.P. (2009) Intestinal barrier function: molecular regulation and disease pathogenesis. *J Allergy Clin Immunol* **124**, 3-20.
- Gruzman A., Babai G. & Sasson S. (2009) Adenosine Monophosphate-Activated Protein Kinase (AMPK) as a New Target for Antidiabetic Drugs: A Review on Metabolic, Pharmacological and Chemical Considerations. *Rev Diabet Stud* **6**, 13-36.

- Gu L., Li N., Gong J., Li Q., Zhu W. & Li J. (2011) Berberine ameliorates intestinal epithelial tight-junction damage and down-regulates myosin light chain kinase pathways in a mouse model of endotoxemia. *J Infect Dis* **203**, 1602-12.
- Gupta S.C., Tyagi A.K., Deshmukh-Taskar P., Hinojosa M., Prasad S. & Aggarwal B.B. (2014) Downregulation of tumor necrosis factor and other proinflammatory biomarkers by polyphenols. *Arch Biochem Biophys* **559**, 91-9.
- Harmon A.W. & Patel Y.M. (2003) Naringenin inhibits phosphoinositide 3-kinase activity and glucose uptake in 3T3-L1 adipocytes. *Biochem Biophys Res Commun* **305**, 229-34.
- Heath J.K. (2010) Transcriptional networks and signaling pathways that govern vertebrate intestinal development. *Curr Top Dev Biol* **90**, 159-92.
- Higashi N., Kohjima M., Fukushima M., Ohta S., Kotoh K., Enjoji M., Kobayashi N. & Nakamuta M. (2005) Epigallocatechin-3-gallate, a green-tea polyphenol, suppresses Rho signaling in TWNT-4 human hepatic stellate cells. *J Lab Clin Med* **145**, 316-22.
- Hilsden R.J., Meddings J.B., Hardin J., Gall D.G. & Sutherland L.R. (1999) Intestinal permeability and postheparin plasma diamine oxidase activity in the prediction of Crohn's disease relapse. *Inflamm Bowel Dis* **5**, 85-91.
- Himes S.R., Sester D.P., Ravasi T., Cronau S.L., Sasmono T. & Hume D.A. (2006) The JNK are important for development and survival of macrophages. *J Immunol* **176**, 2219-28.
- Hollenbach E., Neumann M., Vieth M., Roessner A., Malfertheiner P. & Naumann M. (2004) Inhibition of p38 MAP kinase- and RICK/NF-kappaB-signaling suppresses inflammatory bowel disease. *Faseb Journal* **18**, 1550-2.

- Itoh M., Tsukita S., Yamazaki Y. & Sugimoto H. (2012) Rho GTP exchange factor ARHGEF11 regulates the integrity of epithelial junctions by connecting ZO-1 and RhoA-myosin II signaling. *Proc Natl Acad Sci U S A* **109**, 9905-10.
- Izzi V., Masuelli L., Tresoldi I., Sacchetti P., Modesti A., Galvano F. & Bei R. (2012) The effects of dietary flavonoids on the regulation of redox inflammatory networks. *Front Biosci* **17**, 2396-418.
- Janssens V. & Goris J. (2001) Protein phosphatase 2A: a highly regulated family of serine/threonine phosphatases implicated in cell growth and signalling. *Biochem J* **353**, 417-39.
- Joe B., Vijaykumar M. & Lokesh B.R. (2004) Biological properties of curcumin-cellular and molecular mechanisms of action. *Crit Rev Food Sci Nutr* **44**, 97-111.
- Kanninen K., White A.R., Koistinaho J. & Malm T. (2011) Targeting Glycogen Synthase Kinase-3beta for Therapeutic Benefit against Oxidative Stress in Alzheimer's Disease: Involvement of the Nrf2-ARE Pathway. *Int J Alzheimers Dis* **2011**, 985085.
- Keshavarzian A., Farhadi A., Forsyth C.B., Rangan J., Jakate S., Shaikh M., Banan A. & Fields J.Z. (2009) Evidence that chronic alcohol exposure promotes intestinal oxidative stress, intestinal hyperpermeability and endotoxemia prior to development of alcoholic steatohepatitis in rats. *J Hepatol* **50**, 538-47.
- Kiss A., Becsi B., Kolozsvari B., Komaromi I., Kover K.E. & Erdodi F. (2013) Epigallocatechin-3-gallate and penta-O-galloyl-beta-D-glucose inhibit protein phosphatase-1. *FEBS J* **280**, 612-26.

- Kisseleva T., Song L., Vorontchikhina M., Feirt N., Kitajewski J. & Schindler C. (2006) NF-kappaB regulation of endothelial cell function during LPS-induced toxemia and cancer. *J Clin Invest* **116**, 2955-63.
- Kiyici A., Okudan N., Gokbel H. & Belviranli M. (2010) The effect of grape seed extracts on serum paraoxonase activities in streptozotocin-induced diabetic rats. *J Med Food* **13**, 725-8.
- Knight A.K., Blazquez A.B., Zhang S., Mayer L., Sampson H.A. & Berin M.C. (2007) CD4 T cells activated in the mesenteric lymph node mediate gastrointestinal food allergy in mice. *Am J Physiol Gastrointest Liver Physiol* **293**, G1234-43.
- Kong J., Zhang Z., Musch M.W., Ning G., Sun J., Hart J., Bissonnette M. & Li Y.C. (2008) Novel role of the vitamin D receptor in maintaining the integrity of the intestinal mucosal barrier. *Am J Physiol Gastrointest Liver Physiol* **294**, G208-16.
- Kyriakis J.M. & Avruch J. (2001) Mammalian mitogen-activated protein kinase signal transduction pathways activated by stress and inflammation. *Physiol Rev* **81**, 807-69.
- Laukoetter M.G., Bruewer M. & Nusrat A. (2006) Regulation of the intestinal epithelial barrier by the apical junctional complex. *Curr Opin Gastroenterol* **22**, 85-9.
- Lee A.S., Gibson D.L., Zhang Y., Sham H.P., Vallance B.A. & Dutz J.P. (2010) Gut barrier disruption by an enteric bacterial pathogen accelerates insulinitis in NOD mice. *Diabetologia* **53**, 741-8.
- Lee S.G., Kim B., Yang Y., Pham T.X., Park Y.K., Manatou J., Koo S.I., Chun O.K. & Lee J.Y. (2014) Berry anthocyanins suppress the expression and secretion of proinflammatory mediators in macrophages by inhibiting nuclear translocation of NF-kappaB independent of NRF2-mediated mechanism. *J Nutr Biochem* **25**, 404-11.

- Lemmon M.A. & Schlessinger J. (2010) Cell signaling by receptor tyrosine kinases. *Cell* **141**, 1117-34.
- Li L., Wang L., Wu Z., Yao L., Wu Y., Huang L., Liu K., Zhou X. & Gou D. (2014) Anthocyanin-rich fractions from red raspberries attenuate inflammation in both RAW264.7 macrophages and a mouse model of colitis. *Sci Rep* **4**, 6234.
- Li N. & Neu J. (2009) Glutamine deprivation alters intestinal tight junctions via a PI3-K/Akt mediated pathway in Caco-2 cells. *J Nutr* **139**, 710-4.
- Liu C.M., Sun Y.Z., Sun J.M., Ma J.Q. & Cheng C. (2012) Protective role of quercetin against lead-induced inflammatory response in rat kidney through the ROS-mediated MAPKs and NF-kappaB pathway. *Biochim Biophys Acta* **1820**, 1693-703.
- Liu X., Wang Z., Wang P., Yu B., Liu Y. & Xue Y. (2013) Green tea polyphenols alleviate early BBB damage during experimental focal cerebral ischemia through regulating tight junctions and PKCalpha signaling. *BMC Complement Altern Med* **13**, 187.
- Ma T.Y., Boivin M.A., Ye D., Pedram A. & Said H.M. (2005) Mechanism of TNF- α modulation of Caco-2 intestinal epithelial tight junction barrier: role of myosin light-chain kinase protein expression. *Am J Physiol Gastrointest Liver Physiol* **288**, G422-30.
- Mankertz J., Amasheh M., Krug S.M., Fromm A., Amasheh S., Hillenbrand B., Tavalali S., Fromm M. & Schulzke J.D. (2009) TNFalpha up-regulates claudin-2 expression in epithelial HT-29/B6 cells via phosphatidylinositol-3-kinase signaling. *Cell Tissue Res* **336**, 67-77.
- McLaughlin N., Annabi B., Bouzeghrane M., Temme A., Bahary J.P., Moumdjian R. & Beliveau R. (2006) The Survivin-mediated radioresistant phenotype of glioblastomas is regulated by

- RhoA and inhibited by the green tea polyphenol (-)-epigallocatechin-3-gallate. *Brain Res* **1071**, 1-9.
- Morgan M.J. & Liu Z.G. (2011) Crosstalk of reactive oxygen species and NF-kappaB signaling. *Cell Res* **21**, 103-15.
- Morin P.J. (2005) Claudin proteins in human cancer: promising new targets for diagnosis and therapy. *Cancer Res* **65**, 9603-6.
- Mukai Y., Sun Y. & Sato S. (2013) Azuki bean polyphenols intake during lactation upregulate AMPK in male rat offspring exposed to fetal malnutrition. *Nutrition* **29**, 291-7.
- Mullin J.M., Kampherstein J.A., Laughlin K.V., Clarkin C.E., Miller R.D., Szallasi Z., Kachar B., Soler A.P. & Rosson D. (1998) Overexpression of protein kinase C-delta increases tight junction permeability in LLC-PK1 epithelia. *Am J Physiol* **275**, C544-54.
- Nadour M., Michaud P. & Moulti-Mati F. (2012) Antioxidant activities of polyphenols extracted from olive (*Olea europaea*) of chamlal variety. *Appl Biochem Biotechnol* **167**, 1802-10.
- Narita K., Hisamoto M., Okuda T. & Takeda S. (2011) Differential neuroprotective activity of two different grape seed extracts. *PLoS One* **6**, e14575.
- Noda S., Tanabe S. & Suzuki T. (2012) Differential effects of flavonoids on barrier integrity in human intestinal Caco-2 cells. *J Agric Food Chem* **60**, 4628-33.
- Noda S., Tanabe S. & Suzuki T. (2014) Quercetin increases claudin-4 expression through multiple transcription factors in intestinal Caco-2 cells. *J Funct Foods* **10**, 112-6.
- Novak E.A. & Mollen K.P. (2015) Mitochondrial dysfunction in inflammatory bowel disease. *Front Cell Dev Biol* **3**, 62.

Nunbhakdi-Craig V., Machleidt T., Ogris E., Bellotto D., White C.L., 3rd & Sontag E. (2002)

Protein phosphatase 2A associates with and regulates atypical PKC and the epithelial tight junction complex. *J Cell Biol* **158**, 967-78.

Okkenhaug K. & Vanhaesebroeck B. (2003) PI3K in lymphocyte development, differentiation and activation. *Nat Rev Immunol* **3**, 317-30.

Park H.Y., Kunitake Y., Hirasaki N., Tanaka M. & Matsui T. (2015) Theaflavins enhance intestinal barrier of Caco-2 Cell monolayers through the expression of AMP-activated protein kinase-mediated Occludin, Claudin-1, and ZO-1. *Biosci Biotech and Bioche* **79**, 130-7.

Park S.K., Dahmer M.K. & Quasney M.W. (2012) MAPK and JAK-STAT signaling pathways are involved in the oxidative stress-induced decrease in expression of surfactant protein genes. *Cell Physiol Biochem* **30**, 334-46.

Peng L., Li Z.R., Green R.S., Holzman I.R. & Lin J. (2009) Butyrate enhances the intestinal barrier by facilitating tight junction assembly via activation of AMP-activated protein kinase in Caco-2 cell monolayers. *J Nutr* **139**, 1619-25.

Periasamy V.S. & Alshatwi A.A. (2013) Tea polyphenols modulate antioxidant redox system on cisplatin-induced reactive oxygen species generation in a human breast cancer cell. *Basic Clin Pharmacol Toxicol* **112**, 374-84.

Petecchia L., Sabatini F., Usai C., Caci E., Varesio L. & Rossi G.A. (2012) Cytokines induce tight junction disassembly in airway cells via an EGFR-dependent MAPK/ERK1/2-pathway. *Lab Invest* **92**, 1140-8.

- Philpott D.J., McKay D.M., Mak W., Perdue M.H. & Sherman P.M. (1998) Signal transduction pathways involved in enterohemorrhagic *Escherichia coli*-induced alterations in T84 epithelial permeability. *Infect Immun* **66**, 1680-7.
- Pi J., Zhang Q., Fu J., Woods C.G., Hou Y., Corkey B.E., Collins S. & Andersen M.E. (2010) ROS signaling, oxidative stress and Nrf2 in pancreatic beta-cell function. *Toxicol Appl Pharmacol* **244**, 77-83.
- Piegholdt S., Pallauf K., Esatbeyoglu T., Speck N., Reiss K., Ruddigkeit L., Stocker A., Huebbe P. & Rimbach G. (2014) Biochanin A and prunetin improve epithelial barrier function in intestinal CaCo-2 cells via downregulation of ERK, NF-kappaB, and tyrosine phosphorylation. *Free Radic Biol Med* **70**, 255-64.
- Rajamurugan R., Selvaganabathy N., Kumaravel S., Ramamurthy C., Sujatha V. & Thirunavukkarasu C. (2012) Polyphenol contents and antioxidant activity of Brassica nigra (L.) Koch. leaf extract. *Nat Prod Res* **26**, 2208-10.
- Rao R.K. (1998) Acetaldehyde-induced increase in paracellular permeability in Caco-2 cell monolayer. *Alcohol Clin Exp Res* **22**, 1724-30.
- Rao R.K., Baker R.D., Baker S.S., Gupta A. & Holycross M. (1997) Oxidant-induced disruption of intestinal epithelial barrier function: role of protein tyrosine phosphorylation. *Am J Physiol* **273**, G812-23.
- Ren W., Chen S., Yin J., Duan J., Li T., Liu G., Feng Z., Tan B., Yin Y. & Wu G. (2014a) Dietary arginine supplementation of mice alters the microbial population and activates intestinal innate immunity. *J Nutr* **144**, 988-95.

- Ren W., Duan J., Yin J., Liu G., Cao Z., Xiong X., Chen S., Li T., Yin Y., Hou Y. & Wu G. (2014b) Dietary L-glutamine supplementation modulates microbial community and activates innate immunity in the mouse intestine. *Amino Acids* **46**, 2403-13.
- Ren W., Yin J., Wu M., Liu G., Yang G., Xion Y., Su D., Wu L., Li T., Chen S., Duan J., Yin Y. & Wu G. (2014c) Serum amino acids profile and the beneficial effects of L-arginine or L-glutamine supplementation in dextran sulfate sodium colitis. *PLoS One* **9**, e88335.
- Reuter S., Gupta S.C., Chaturvedi M.M. & Aggarwal B.B. (2010) Oxidative stress, inflammation, and cancer: how are they linked? *Free Radic Biol Med* **49**, 1603-16.
- Rigor R.R., Shen Q., Pivetti C.D., Wu M.H. & Yuan S.Y. (2013) Myosin light chain kinase signaling in endothelial barrier dysfunction. *Med Res Rev* **33**, 911-33.
- Rosenthal R., Gunzel D., Finger C., Krug S.M., Richter J.F., Schulzke J.D., Fromm M. & Amasheh S. (2012) The effect of chitosan on transcellular and paracellular mechanisms in the intestinal epithelial barrier. *Biomaterials* **33**, 2791-800.
- Rotches-Ribalta M., Andres-Lacueva C., Estruch R., Escribano E. & Urpi-Sarda M. (2012) Pharmacokinetics of resveratrol metabolic profile in healthy humans after moderate consumption of red wine and grape extract tablets. *Pharmacol Res* **66**, 375-82.
- Samak G., Aggarwal S. & Rao R.K. (2011) ERK is involved in EGF-mediated protection of tight junctions, but not adherens junctions, in acetaldehyde-treated Caco-2 cell monolayers. *Am J Physiol Gastrointest Liver Physiol* **301**, G50-9.
- Santangelo C., Vari R., Scazzocchio B., Di Benedetto R., Filesi C. & Masella R. (2007) Polyphenols, intracellular signalling and inflammation. *Ann Ist Super Sanita* **43**, 394-405.

- Scaldaferri F., Pizzoferrato M., Gerardi V., Lopetuso L. & Gasbarrini A. (2012) The gut barrier: new acquisitions and therapeutic approaches. *J Clin Gastroenterol* **46 Suppl**, S12-7.
- Scharl M., Paul G., Barrett K.E. & McCole D.F. (2009) AMP-activated protein kinase mediates the interferon-gamma-induced decrease in intestinal epithelial barrier function. *J Biol Chem* **284**, 27952-63.
- Schindler J.F., Monahan J.B. & Smith W.G. (2007) p38 pathway kinases as anti-inflammatory drug targets. *J Dent Res* **86**, 800-11.
- Schmeck B., Moog K., Zahlten J., van Laak V., N'Guessan P.D., Opitz B., Rosseau S., Suttorp N. & Hippenstiel S. (2006) Streptococcus pneumoniae induced c-Jun-N-terminal kinase- and AP-1 -dependent IL-8 release by lung epithelial BEAS-2B cells. *Respir Res* **7**, 98.
- Scott K.G., Meddings J.B., Kirk D.R., Lees-Miller S.P. & Buret A.G. (2002) Intestinal infection with Giardia spp. reduces epithelial barrier function in a myosin light chain kinase-dependent fashion. *Gastroenterology* **123**, 1179-90.
- Segaliny A.I., Tellez-Gabriel M., Heymann M.F. & Heymann D. (2015) Receptor tyrosine kinases: Characterisation, mechanism of action and therapeutic interests for bone cancers. *J Bone Oncol* **4**, 1-12.
- Senftleben U. & Karin M. (2002) The IKK/NF-kappaB pathway. *Crit Care Med* **30**, S18-S26.
- Seth A., Basuroy S., Sheth P. & Rao R.K. (2004) L-Glutamine ameliorates acetaldehyde-induced increase in paracellular permeability in Caco-2 cell monolayer. *Am J Physiol Gastrointest Liver Physiol* **287**, G510-7.

- Seth A., Sheth P., Elias B.C. & Rao R. (2007) Protein phosphatases 2A and 1 interact with occludin and negatively regulate the assembly of tight junctions in the CACO-2 cell monolayer. *J Biol Chem* **282**, 11487-98.
- Sheth P., Basuroy S., Li C., Naren A.P. & Rao R.K. (2003) Role of phosphatidylinositol 3-kinase in oxidative stress-induced disruption of tight junctions. *J Biol Chem* **278**, 49239-45.
- Sheth P., Samak G., Shull J.A., Seth A. & Rao R. (2009) Protein phosphatase 2A plays a role in hydrogen peroxide-induced disruption of tight junctions in Caco-2 cell monolayers. *Biochem J* **421**, 59-70.
- Shin D.Y., Lu J.N., Kim G.Y., Jung J.M., Kang H.S., Lee W.S. & Choi Y.H. (2011) Anti-invasive activities of anthocyanins through modulation of tight junctions and suppression of matrix metalloproteinase activities in HCT-116 human colon carcinoma cells. *Oncol Rep* **25**, 567-72.
- Singh U., Van Itallie C.M., Mitic L.L., Anderson J.M. & McClane B.A. (2000) CaCo-2 cells treated with *Clostridium perfringens* enterotoxin form multiple large complex species, one of which contains the tight junction protein occludin. *J Biol Chem* **275**, 18407-17.
- Song J.C., Hanson C.M., Tsai V., Farokhzad O.C., Lotz M. & Matthews J.B. (2001) Regulation of epithelial transport and barrier function by distinct protein kinase C isoforms. *Am J Physiol Cell Physiol* **281**, C649-61.
- Song P., Zhang R., Wang X., He P., Tan L. & Ma X. (2011a) Dietary grape-seed procyanidins decreased postweaning diarrhea by modulating intestinal permeability and suppressing oxidative stress in rats. *J Agric Food Chem* **59**, 6227-32.

- Song Y.A., Park Y.L., Yoon S.H., Kim K.Y., Cho S.B., Lee W.S., Chung I.J. & Joo Y.E. (2011b) Black tea polyphenol theaflavin suppresses LPS-induced ICAM-1 and VCAM-1 expression via blockage of NF-kappaB and JNK activation in intestinal epithelial cells. *Inflamm Res* **60**, 493-500.
- Sontag E., Sontag J.M. & Garcia A. (1997) Protein phosphatase 2A is a critical regulator of protein kinase C zeta signaling targeted by SV40 small t to promote cell growth and NF-kappaB activation. *EMBO J* **16**, 5662-71.
- Spalinger M.R., Lang S., Weber A., Frei P., Fried M., Rogler G. & Scharl M. (2013) Loss of Protein Tyrosine Phosphatase Nonreceptor Type 22 Regulates Interferon-gamma-Induced Signaling in Human Monocytes. *Gastroenterology* **144**, 978-88 e10.
- Strack S., Cribbs J.T. & Gomez L. (2004) Critical role for protein phosphatase 2A heterotrimers in mammalian cell survival. *J Biol Chem* **279**, 47732-9.
- Su X. & D'Souza D.H. (2011) Grape seed extract for control of human enteric viruses. *Appl Environ Microbiol* **77**, 3982-7.
- Suenaert P., Bulteel V., Lemmens L., Noman M., Geypens B., Van Assche G., Geboes K., Ceuppens J.L. & Rutgeerts P. (2002) Anti-tumor necrosis factor treatment restores the gut barrier in Crohn's disease. *Am J Gastroenterol* **97**, 2000-4.
- Suzuki T. (2013) Regulation of intestinal epithelial permeability by tight junctions. *Cell Mol Life Sci* **70**, 631-59.
- Suzuki T., Elias B.C., Seth A., Shen L., Turner J.R., Giorgianni F., Desiderio D., Guntaka R. & Rao R. (2009) PKC eta regulates occludin phosphorylation and epithelial tight junction integrity. *Proc Natl Acad Sci U S A* **106**, 61-6.

- Suzuki T. & Hara H. (2009) Quercetin enhances intestinal barrier function through the assembly of zonula [corrected] occludens-2, occludin, and claudin-1 and the expression of claudin-4 in Caco-2 cells. *J Nutr* **139**, 965-74.
- Suzuki T., Seth A. & Rao R. (2008) Role of phospholipase Cgamma-induced activation of protein kinase Cepsilon (PKCepsilon) and PKCbetaI in epidermal growth factor-mediated protection of tight junctions from acetaldehyde in Caco-2 cell monolayers. *J Biol Chem* **283**, 3574-83.
- Suzuki T., Tanabe S. & Hara H. (2011a) Kaempferol enhances intestinal barrier function through the cytoskeletal association and expression of tight junction proteins in Caco-2 cells. *J Nutr* **141**, 87-94.
- Suzuki T., Yoshinaga N. & Tanabe S. (2011b) Interleukin-6 (IL-6) regulates claudin-2 expression and tight junction permeability in intestinal epithelium. *J Biol Chem* **286**, 31263-71.
- Tsao R. (2010) Chemistry and biochemistry of dietary polyphenols. *Nutrients* **2**, 1231-46.
- Turner J.R., Angle J.M., Black E.D., Joyal J.L., Sacks D.B. & Madara J.L. (1999) PKC-dependent regulation of transepithelial resistance: roles of MLC and MLC kinase. *Am J Physiol* **277**, C554-62.
- Ulluwishewa D., Anderson R.C., McNabb W.C., Moughan P.J., Wells J.M. & Roy N.C. (2011) Regulation of tight junction permeability by intestinal bacteria and dietary components. *J Nutr* **141**, 769-76.
- Vaarala O. (2012) Is the origin of type 1 diabetes in the gut? *Immunol Cell Biol* **90**, 271-6.

- Van Kanegan M.J., Adams D.G., Wadzinski B.E. & Strack S. (2005) Distinct protein phosphatase 2A heterotrimers modulate growth factor signaling to extracellular signal-regulated kinases and Akt. *J Biol Chem* **280**, 36029-36.
- Velmurugan B., Singh R.P., Agarwal R. & Agarwal C. (2010a) Dietary-feeding of grape seed extract prevents azoxymethane-induced colonic aberrant crypt foci formation in fischer 344 rats. *Mol Carcinog* **49**, 641-52.
- Velmurugan B., Singh R.P., Kaul N., Agarwal R. & Agarwal C. (2010b) Dietary feeding of grape seed extract prevents intestinal tumorigenesis in APCmin/+ mice. *Neoplasia* **12**, 95-102.
- Walker E.H., Pacold M.E., Perisic O., Stephens L., Hawkins P.T., Wymann M.P. & Williams R.L. (2000) Structural determinants of phosphoinositide 3-kinase inhibition by wortmannin, LY294002, quercetin, myricetin, and staurosporine. *Mol Cell* **6**, 909-19.
- Walsh S.V., Hopkins A.M., Chen J., Narumiya S., Parkos C.A. & Nusrat A. (2001) Rho kinase regulates tight junction function and is necessary for tight junction assembly in polarized intestinal epithelia. *Gastroenterology* **121**, 566-79.
- Wang A., Liu M., Liu X., Dong L.Q., Glickman R.D., Slaga T.J., Zhou Z. & Liu F. (2011a) Up-regulation of adiponectin by resveratrol: the essential roles of the Akt/FOXO1 and AMP-activated protein kinase signaling pathways and DsbA-L. *J Biol Chem* **286**, 60-6.
- Wang B., Yang G., Liang X., Zhu M. & Du M. (2014) Grape seed extract prevents skeletal muscle wasting in interleukin 10 knockout mice. *BMC Complement Altern Med* **14**, 162.
- Wang H., Xue Y., Zhang H., Huang Y., Yang G., Du M. & Zhu M.J. (2013) Dietary grape seed extract ameliorates symptoms of inflammatory bowel disease in IL10-deficient mice. *Mol Nutr Food Res* **57**, 2253-7.

- Wang H., Zhang C., Wu G., Sun Y., Wang B., He B., Dai Z. & Wu Z. (2015) Glutamine enhances tight junction protein expression and modulates corticotropin-releasing factor signaling in the jejunum of weanling piglets. *J Nutr* **145**, 25-31.
- Wang H., Zhao J.X., Hu N., Ren J., Du M. & Zhu M.J. (2012a) Side-stream smoking reduces intestinal inflammation and increases expression of tight junction proteins. *World J Gastroenterol* **18**, 2180-7.
- Wang N., Wang G., Hao J., Ma J., Wang Y., Jiang X. & Jiang H. (2012b) Curcumin ameliorates hydrogen peroxide-induced epithelial barrier disruption by upregulating heme oxygenase-1 expression in human intestinal epithelial cells. *Dig Dis Sci* **57**, 1792-801.
- Wang Y., Zhang J., Yi X.J. & Yu F.S. (2004) Activation of ERK1/2 MAP kinase pathway induces tight junction disruption in human corneal epithelial cells. *Exp Eye Res* **78**, 125-36.
- Wang Y.H., Ge B., Yang X.L., Zhai J., Yang L.N., Wang X.X., Liu X., Shi J.C. & Wu Y.J. (2011b) Proanthocyanidins from grape seeds modulates the nuclear factor-kappa B signal transduction pathways in rats with TNBS-induced recurrent ulcerative colitis. *Int Immunopharmacol* **11**, 1620-7.
- Wang Y.J., Thomas P., Zhong J.H., Bi F.F., Kosaraju S., Pollard A., Fenech M. & Zhou X.F. (2009) Consumption of grape seed extract prevents amyloid-beta deposition and attenuates inflammation in brain of an Alzheimer's disease mouse. *Neurotox Res* **15**, 3-14.
- Watson J.L., Ansari S., Cameron H., Wang A., Akhtar M. & McKay D.M. (2004) Green tea polyphenol (-)-epigallocatechin gallate blocks epithelial barrier dysfunction provoked by IFN-gamma but not by IL-4. *Am J Physiol Gastrointest Liver Physiol* **287**, G954-61.

- Watson J.L., Vicario M., Wang A., Moreto M. & McKay D.M. (2005) Immune cell activation and subsequent epithelial dysfunction by Staphylococcus enterotoxin B is attenuated by the green tea polyphenol (-)-epigallocatechin gallate. *Cell Immunol* **237**, 7-16.
- Wei H.K., Chen G., Wang R.J. & Peng J. (2015) Oregano essential oil decreased susceptibility to oxidative stress-induced dysfunction of intestinal epithelial barrier in rats. *J Funct Foods* **18**, 1191-9.
- Wittchen E.S., Haskins J. & Stevenson B.R. (2003) NZO-3 expression causes global changes to actin cytoskeleton in Madin-Darby canine kidney cells: linking a tight junction protein to Rho GTPases. *Mol Biol Cell* **14**, 1757-68.
- Yang G., Wang H., Kang Y. & Zhu M.J. (2014) Grape seed extract improves epithelial structure and suppresses inflammation in ileum of IL-10-deficient mice. *Food Funct* **5**, 2558-63.
- Yang G., Xue Y., Zhang H., Du M. & Zhu M.J. (2015) Favourable effects of grape seed extract on intestinal epithelial differentiation and barrier function in IL10-deficient mice. *Br J Nutr*, 1-9.
- Yang T.P., Lee H.J., Ou T.T., Chang Y.J. & Wang C.J. (2012) Mulberry leaf polyphenol extract induced apoptosis involving regulation of adenosine monophosphate-activated protein kinase/fatty acid synthase in a p53-negative hepatocellular carcinoma cell. *J Agric Food Chem* **60**, 6891-8.
- Yao K., Guan S., Li T., Huang R., Wu G., Ruan Z. & Yin Y. (2011) Dietary L-arginine supplementation enhances intestinal development and expression of vascular endothelial growth factor in weanling piglets. *Br J Nutr* **105**, 703-9.

- Ye D., Ma I. & Ma T.Y. (2006) Molecular mechanism of tumor necrosis factor- α modulation of intestinal epithelial tight junction barrier. *Am J Physiol Gastrointest Liver Physiol* **290**, G496-504.
- Yin J., Liu M., Ren W., Duan J., Yang G., Zhao Y., Fang R., Chen L., Li T. & Yin Y. (2015a) Effects of dietary supplementation with glutamate and aspartate on diquat-induced oxidative stress in piglets. *PLoS One* **10**, e0122893.
- Yin J., Ren W., Duan J., Wu L., Chen S., Li T., Yin Y. & Wu G. (2014) Dietary arginine supplementation enhances intestinal expression of SLC7A7 and SLC7A1 and ameliorates growth depression in mycotoxin-challenged pigs. *Amino Acids* **46**, 883-92.
- Yin J., Wu M., Duan J., Liu G., Cui Z., Zheng J., Chen S., Ren W., Deng J., Tan X., Al-Dhabi N.A., Duraipandiyan V., Liao P., Li T. & Yulong Y. (2015b) Pyrrolidine Dithiocarbamate Inhibits NF-KappaB Activation and Upregulates the Expression of Gpx1, Gpx4, Occludin, and ZO-1 in DSS-Induced Colitis. *Appl Biochem Biotechnol* **177**, 1716-28.
- Yu D., Marchiando A.M., Weber C.R., Raleigh D.R., Wang Y., Shen L. & Turner J.R. (2010) MLCK-dependent exchange and actin binding region-dependent anchoring of ZO-1 regulate tight junction barrier function. *Proc Natl Acad Sci U S A* **107**, 8237-41.
- Yu L.C. (2009) The epithelial gatekeeper against food allergy. *Pediatr Neonatol* **50**, 247-54.
- Zeissig S., Burgel N., Gunzel D., Richter J., Mankertz J., Wahnschaffe U., Kroesen A.J., Zeitz M., Fromm M. & Schulzke J.D. (2007) Changes in expression and distribution of claudin 2, 5 and 8 lead to discontinuous tight junctions and barrier dysfunction in active Crohn's disease. *Gut* **56**, 61-72.

- Zhang L., Li J., Young L.H. & Caplan M.J. (2006) AMP-activated protein kinase regulates the assembly of epithelial tight junctions. *Proc Natl Acad Sci U S A* **103**, 17272-7.
- Zhang Y. & Li J. (2012) Carbachol ameliorates lipopolysaccharide-induced intestinal epithelial tight junction damage by down-regulating NF-kappabeta and myosin light-chain kinase pathways. *Biochem Biophys Res Commun* **428**, 321-6.
- Zhao Y., Sun Y., Ding Y., Wang X., Zhou Y., Li W., Huang S., Li Z., Kong L., Guo Q. & Lu N. (2015) GL-V9, a new synthetic flavonoid derivative, ameliorates DSS-induced colitis against oxidative stress by up-regulating Trx-1 expression via activation of AMPK/FOXO3a pathway. *Oncotarget* **6**, 26291-307.
- Zolotarevsky Y., Hecht G., Koutsouris A., Gonzalez D.E., Quan C., Tom J., Mrsny R.J. & Turner J.R. (2002) A membrane-permeant peptide that inhibits MLC kinase restores barrier function in in vitro models of intestinal disease. *Gastroenterology* **123**, 163-72.

Table 1. Summary of direct evidence of the mechanisms connecting the polyphenol compounds and barrier integrity.

Experimental setting	Polyphenol source	Signaling pathways	Response	Reference
TNF- α induced HT29/B6 cells	Berberine	NF- κ B↓ PI3K/Akt↓ tyrosine kinases↓	Ameliorated claudin-1 disassembly and claudin 2↓	(Amasheh <i>et al.</i> 2010)
IL-10-deficient mice	Grape seed extract	NF- κ B↓	Claudin-1↑ and claudin 2↓	(Wang <i>et al.</i> 2013; Yang <i>et al.</i> 2014)
TNF- α induced CaCO-2 cells	Biochanin A, prunetin	NF- κ B↓ ERK↓ tyrosine kinases↓	Improved TEER, Claudin 1→, occludin→, ZO-1→, E-cadherin →	(Piegholdt <i>et al.</i> 2014)
DDS-induced colitis in mice	Anthocyanin rich extract of raspberries	NF- κ B↓ MAPKs↓	Improved Colonic histological architecture	(Li <i>et al.</i> 2014)
Hydrogen peroxide induced ECV304	Quercetin	p38↓	ZO-1↑and occludin↑	(Chuenkitiyano <i>et al.</i> 2010)

monolayers				
HCT-116 human colon carcinoma cells	Anthocyanin	p38↑	Improved TEER, claudin-1↓, claudin-3↓ and claudin-4↓	(Shin <i>et al.</i> 2011)
Caco-2 cells	Quercetin	PKCδ↓	ZO-2↑, occludin↑, claudin-1↑, claudin-4↑	(Suzuki & Hara 2009)
Blood-brain barrier damage	Green tea polyphenols	PKCα↓	Claudin-5↑, occludin↑ and ZO-1↑	(Liu <i>et al.</i> 2013)
Acetaldehyde-induced Caco-2 cells	Genistein	Tyrosine kinases↓	Improved TEER, distribution of ZO-1 and occludin↑	(Atkinson & Rao 2001)
LPS challenged mouse	Berberine	MLCK↓	Distribution of claudin-1, claudin-4, ZO-1 and occludin↑ Intestinal permeability↓	(Gu <i>et al.</i> 2011)
IFN-γ and TNF-α Induced CaCO-2 cells	Berberine	MLCK↓	Distribution of claudin-1, ZO-1 and occludin↑ Intestinal permeability↓	(Cao <i>et al.</i> 2013)
	Theaflavins-	AMPK↑	Occludin↑, claudin-1↑,	(Park <i>et al.</i>

Caco-2 cells	3'-O-gallate		and ZO-1↑	2015)
DSS-treated Caco-2 monolayers	6-gingerol	AMPK↑	Improved TEER	(Chang & Kuo 2015)
IL-10-deficient mice	Grape seed extract	AMPK↓	Claudin-1↑ and claudin 2↓	(Yang <i>et al.</i> 2014)
Diquat-induced rats	Oregano essential oils	SOD and GSH-Px↓	ZO-1↑and occludin↑	(Wei <i>et al.</i> 2015)

↑, increase or improve; ↓, decrease; →, not change.