



Critical Reviews in Food Science and Nutrition

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/bfsn20>

Meta-Review of Protein Network Regulating Obesity Between Validated Obesity Candidate Genes in the White Adipose Tissue of High-Fat Diet-Induced Obese C57BL/6J Mice

Eunjung Kim^{a e}, Eun Jung Kim^b, Seung-Won Seo^c, Cheol-Goo Hur^d, Robin A McGregor^e & Myung-Sook Choi^e

^a Department of Food Science and Nutrition, Catholic University of Daegu, Gyeongsan, Republic of Korea

^b Department of Pharmacology and Dental Therapeutics, Chonnam National University School of Dentistry, Gwangju, Republic of Korea

^c University of Science and Technology, Daejeon, Korea

^d Bioinformatics Research Center, Korea Research Institute of Bioscience and Biotechnology, Daejeon, Republic of Korea

^e Center for Food & Nutritional Genomics, Department of Food Sciences and Nutrition, Kyungpook National University, Daegu, Republic of Korea

Accepted author version posted online: 08 Mar 2013. Published online: 05 Feb 2014.

To cite this article: Eunjung Kim, Eun Jung Kim, Seung-Won Seo, Cheol-Goo Hur, Robin A McGregor & Myung-Sook Choi (2014) Meta-Review of Protein Network Regulating Obesity Between Validated Obesity Candidate Genes in the White Adipose Tissue of High-Fat Diet-Induced Obese C57BL/6J Mice, Critical Reviews in Food Science and Nutrition, 54:7, 910-923, DOI: [10.1080/10408398.2011.619283](https://doi.org/10.1080/10408398.2011.619283)

To link to this article: <http://dx.doi.org/10.1080/10408398.2011.619283>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

Mini Review Paper

Meta-Review of Protein Network Regulating Obesity Between Validated Obesity Candidate Genes in the White Adipose Tissue of High-Fat Diet-Induced Obese C57BL/6J Mice

EUNJUNG KIM,^{1,5} EUN JUNG KIM,² SEUNG-WON SEO,³ CHEOL-GOO HUR,⁴
ROBIN A MCGREGOR,⁵ and MYUNG-SOOK CHOI⁵

¹Department of Food Science and Nutrition, Catholic University of Daegu, Gyeongsan, Republic of Korea

²Department of Pharmacology and Dental Therapeutics, Chonnam National University School of Dentistry, Gwangju, Republic of Korea

³University of Science and Technology, Daejeon, Korea

⁴Bioinformatics Research Center, Korea Research Institute of Bioscience and Biotechnology, Daejeon, Republic of Korea

⁵Center for Food & Nutritional Genomics, Department of Food Sciences and Nutrition, Kyungpook National University, Daegu, Republic of Korea

Worldwide obesity and related comorbidities are increasing, but identifying new therapeutic targets remains a challenge. A plethora of microarray studies in diet-induced obesity models has provided large datasets of obesity associated genes. In this review, we describe an approach to examine the underlying molecular network regulating obesity, and we discuss interactions between obesity candidate genes. We conducted network analysis on functional protein–protein interactions associated with 25 obesity candidate genes identified in a literature-driven approach based on published microarray studies of diet-induced obesity. The obesity candidate genes were closely associated with lipid metabolism and inflammation. Peroxisome proliferator activated receptor gamma (Pparg) appeared to be a core obesity gene, and obesity candidate genes were highly interconnected, suggesting a coordinately regulated molecular network in adipose tissue. In conclusion, the current network analysis approach may help elucidate the underlying molecular network regulating obesity and identify anti-obesity targets for therapeutic intervention.

Keywords Obesity, nutrigenomics, high-fat, C57BL/6J mouse, adipose tissue

INTRODUCTION

Obesity is a global health problem characterized by excess body fat, and is associated with many chronic diseases including type 2 diabetes mellitus (T2DM), hypertension, dyslipidemia, and certain types of cancer (Friedman, 2003; Adan et al., 2008; Bessesen, 2008; Hill et al., 2008). Despite recent advances in obesity research, the mechanisms underlying the pathogenesis

of obesity are not fully understood. Obesity can develop due to a chronic imbalance between energy intake and energy expenditure, leading to greater energy storage and increased adipose tissue mass (Spiegelman and Flier, 2001).

Adipose tissue was for many decades regarded as a passive lipid and energy storage depot releasing fatty acids (FAs) and glycerol to be used as an energy source in other oxidative tissues in times of fasting or starvation (Flier, 2004). In the fed state adipose tissue primarily absorbs circulating FAs and triglycerides (TGs) (Frayn, 2002; Sharma and Staels, 2007), hence protecting other tissues from excessive FA flux (Frayn et al., 2005). More recently, it has been established that adipose tissue is a much

Address correspondence to Myung-Sook Choi, Ph.D., Department of Food Science and Nutrition, Kyungpook National University, 1370 San-Kyuk Dong, Buk-Ku, Daegu 702-701, Republic of Korea. E-mail: mschoi@knu.ac.kr

more dynamic endocrine organ, capable of producing a wide range of hormones and cytokines, called adipokines (Kershaw and Flier, 2004; Hauner, 2005). So far, more than 100 adipokines have been identified and these molecules can affect inflammatory processes, endocrine and metabolic processes, and cardiovascular disease development. Hence, adipose tissue appears to play an integral part in both energy homeostasis and inflammation (Rajala and Scherer, 2003; Guerre-Millo, 2004; Trayhurn and Wood, 2004).

Recent development of microarray technology has now made it possible to analyze integrated system changes in the transcriptome underlying the development of obesity (Kim and Park, 2008; Lopez et al., 2003). However, despite the enormous amount of gene expression data available from obese animal models and obese humans, the disparity between microarray studies has impeded the coordinated interpretation of the changes in the transcriptome underlying obesity.

We, therefore, extensively mined published microarray datasets of gene expression changes in white adipose tissue (WAT) of high-fat diet (HFD) induced obese C57BL/6J mice. We selected 25 obesity candidate genes from these microarray studies, which have been independently validated. In the following, we provide a systemic overview of functional protein–protein interactions of these obesity candidate genes to better understand the pathophysiological changes contributing to the development of the obese phenotype.

Identification of Obesity Candidate Genes in White Adipose Tissue

Firstly, we identified adipose tissue microarray studies of high-fat diet fed C57BL/6J mice in the PubMed database. From these microarray studies, we extracted genes which were reported as differentially expressed following HFD feeding. From this gene list, we then searched for independent evidence that these genes are differentially expressed in obesity based on other analytical methods such as reverse transcription-polymerase chain reaction or northern blot analysis.

Twenty-five genes were selected and their specific cellular functions are summarized in Table 1. Of 25 genes, 6 genes were associated with lipid metabolism, 3 genes were associated with inflammation, and 7 genes were associated with signal transduction and transport. Overall, the expression of genes involved in FA synthesis [*acetyl CoA carboxylase (Acaca)*, *fatty acid synthase (Fasn)*, *stearoyl coenzyme A desaturase 1 (Scd1)*, and *stearoyl coenzyme A desaturase 2 (Scd2)*] and lipid uptake [*lipoprotein lipase (Lpl)*] were significantly increased in the adipose tissue of HFD fed mice. The expression of 3-hydroxy-3-methylglutaryl CoA synthase 1 (*Hmgcs1*) involved in steroid synthesis was downregulated probably because high sterol intake inhibits *Hmgcs1* expression by negative feedback inhibition.

On the other hand, genes involved in immune and inflammation [*interleukin 6 (Il6)*, *chemokine (C-C motif) ligand 2*

(*Ccl2*, *MCP-1*), and *tumor necrosis factor alpha (Tnfa)*] were all upregulated. In addition, the expression of signal transduction and transporter-associated genes [*fatty acid binding protein 4 (Fabp4)*, *complement factor D (Cfd)*, *leptin (Lep)*, and *peroxisome proliferator activated receptor γ (Pparg)*] were upregulated; however, *adiponectin (Adipoq)* expression was downregulated.

HFD feeding also affects the expression of glucose metabolism-associated genes. For example, increased expression of *glucose-6-phosphate dehydrogenase 2 (G6pd2)* in adipose tissue appears to be related with insulin resistance, while decreased expression of *glucose transporter 4 (Slc2a4)* and *sterol regulatory element binding transcription factor 1 (Srebfl)* suggests some impairment of the insulin signaling pathway in HFD fed mice. Downregulated *glycerol-3-phosphate dehydrogenase 1 (Gpd1)* suggests energy production by glycolysis and gluconeogenesis is decreased by HFD feeding.

Furthermore, the expression of genes associated with xenobiotic metabolism, involved in the detoxification of oxidative stress [*catalase (Cat)*] was inhibited by HFD. Other obesity candidate genes identified in adipose tissue of HFD fed mice included *cathepsin D (Ctsd)*, *CCAAT/enhancer binding protein α (Cebpa)*, *plasminogen activator inhibitor 1 (Serpine 1, PAI-1)*, *uncoupling protein 1 (Ucp1)*, and *uncoupling protein 2 (Ucp2)* which were all upregulated, whereas *superoxide dismutase 1 (Sod1)* was downregulated.

Analysis of Functional Protein Interactions Between Obesity Candidate Genes

Next, we used a network approach based on protein–protein interactions to determine functional connections between obesity candidate genes. Protein–protein interactions were extracted from the STRING database and a network was built using the active prediction method (confidence score above 0.8, network depth 2). We found 8 of 25 obesity genes were closely connected to each other in one-depth condition in which a protein is directly connected with another protein (pink block; Fig. 1). Furthermore, all nonconnected genes in the one-depth condition (17 genes) were found to be related when the analysis was extended to the two-depth condition. The network analysis suggested a high number of interactions are predicted to occur between the identified obesity candidate genes.

Pparg is a Core Gene in the Functional Protein Interactions of Obesity Candidate Genes

Pparg emerged as a central node among the selected obesity candidate genes in adipose tissue. *Pparg* belongs to the ligand-activated nuclear hormone receptor family and acts as a transcription factor in heterodimer formation with retinoid X receptor to modulate gene expression involved in adipocyte differentiation, FA uptake and storage (Ferre, 2004; Anghel and

Table 1 Lipid metabolism-associated genes differentially expressed in adipose tissue of HF-induced obese C57BL/6J mice

| Gene name | Expression | Gene ontology (process) | References |
|--|------------|---|--|
| Acetyl CoA carboxylase (Acaca) | ↑ | Acetyl-CoA metabolic process Fatty acid biosynthetic process Lipid homeostasis Lipid metabolic process Response to organic cyclic substance tissue homeostasis | (Al-Hasani and Joost, 2005; Srivastava et al., 2006; Kim and Kim, 2009) |
| Fatty acid synthase (Fasn) | ↑ | Fatty acid biosynthetic process Metabolic process Oxidation reduction | (Soukas et al., 2000; Harada et al., 2003; Al-Hasani and Joost, 2005; Srivastava et al., 2006; Mong et al., 2011) |
| 3-hydroxy-3-methylglutaryl CoA synthase 1 (Hmgcs1) | ↓ | Cholesterol biosynthetic process Isoprenoid biosynthetic process Lipid biosynthetic process Metabolic process Response to organic nitrogen Steroid biosynthetic process Sterol biosynthetic process | (Soukas et al., 2000; Harada et al., 2003; Mong et al., 2011) |
| Lipoprotein lipase (Lpl) | ↑ | Lipid metabolic process Positive regulation of macrophage derived foam cell differentiation Triglyceride biosynthetic process Triglyceride catabolic process | (Harada et al., 2003; Al-Hasani and Joost, 2005; Srivastava et al., 2006) |
| Stearoyl coenzyme A desaturase 1 (Scd1) | ↑ | Brown fat cell differentiation Cholesterol esterification Fatty acid biosynthetic process Lipid metabolic process Oxidation reduction White fat cell differentiation | (Soukas et al., 2000; Soukas et al., 2001; Al-Hasani and Joost, 2005; Flachs et al., 2005; Nikonova et al., 2008) |
| Stearoyl coenzyme A desaturase 2 (Scd2) | ↑ | Fatty acid biosynthetic process Lipid metabolic process Oxidation reduction | (Soukas et al., 2000; Soukas et al., 2001) |
| Interleukin 6 (Il6) | ↑ | Glucagon secretion Immune response Interleukin-6-mediated signaling pathway Regulation of apoptosis Regulation of cell proliferation Regulation of cell shape | (Weisberg et al., 2003; Chen et al., 2005; Cole et al., 2010; Li et al., 2010; Kang et al., 2010; Orellana-Gavalda et al., 2011) |
| chemokine (C-C motif) ligand 2 (Ccl2, MCP-1) | ↑ | Chemokine-mediated signaling pathway Chemotaxis Cytokine-mediated signaling pathway Immune response Inflammatory response | (Chen et al., 2005; Kanda et al., 2006; Fukushima et al., 2009; Cole et al., 2010; Kang et al., 2010; Orellana-Gavalda et al., 2011) |
| Tumor necrosis factor alpha (Tnfa) | ↑ | Apoptosis Defense response Humoral immune response Immune response Inflammatory response | (Weisberg et al., 2003; Harada et al., 2003; Chen et al., 2005; Kim et al., 2008; Hagiwara et al., 2009; Cole et al., 2010; Li et al., 2010; Kang et al., 2010; Orellana-Gavalda et al., 2011) |
| Fatty acid binding protein 4 (Fabp4) | ↑ | Brown fat cell differentiation Cholesterol homeostasis Cytokine production Fatty acid metabolic process Transport White fat cell differentiation | (Harada et al., 2003; Al-Hasani and Joost, 2005; Soukas et al., 2000; Kim et al., 2008; Kim and Park, 2010) |
| Adiponectin (Adipoq) | ↓ | Adiponectin-mediated signaling pathway Brown fat cell differentiation Fatty acid beta-oxidation Fatty acid oxidation Glucose homeostasis | (Barnea et al., 2006; Hosogai et al., 2007; Hagiwara et al., 2009; Soukas et al., 2000; Soukas et al., 2001; Cole et al., 2010; Kang et al., 2010) |
| Complement factor D (Cfd) | ↑ | Notch signaling pathway Complement activation, alternative pathway Innate immune response Proteolysis | (Soukas et al., 2000; Soukas et al., 2001; Harada et al., 2003; Al-Hasani and Joost, 2005; Flachs et al., 2005; Chen et al., 2005) |

(Continued on next page)

Table 1 Lipid metabolism-associated genes differentially expressed in adipose tissue of HF-induced obese C57BL/6J mice (*Continued*)

| Gene name | Expression | Gene ontology (process) | References |
|---|------------|--|---|
| Glucose transporter 4 (Slc2a4) | ↓ | Brown fat cell differentiation Carbohydrate transport Glucose homeostasis Glucose transport Hexose transmembrane transport Ion transmembrane transport Proton transport Transmembrane transport | (Moraes et al., 2003; Al-Hasani and Joost, 2005; Kang et al., 2010) |
| Leptin (Lep) | ↑ | Bile acid metabolic process Central nervous system neuron development Cholesterol metabolic process Eating behavior Glucose metabolic process Hormone metabolic process Insulin secretion Lipid metabolic process | (Harada et al., 2003; Van Schothorst et al., 2005; Marrades et al., 2006; Hosogai et al., 2007; Fukushima et al., 2009; Hagiwara et al., 2009; An et al., 2010) |
| Peroxisome proliferator-activated receptor γ (Pparg) | ↑ | Cell fate commitment Fat cell differentiation Fatty acid oxidation Inflammatory response Long-chain fatty acid transport Response to lipid Signal transduction Transcription White fat cell differentiation | (Soukas et al., 2000; Morton et al., 2004; Hosogai et al., 2007; Kim et al., 2008; Kim and Kim, 2009; Kim and Park, 2010) |
| Sterol regulatory element-binding transcription factor 1 (Srebf1) | ↓ | Cellular response to starvation Cholesterol metabolic process Insulin receptor signaling pathway Lipid metabolic process Regulation of fatty acid metabolic process Regulation of heart rate by chemical signal Regulation of insulin secretion Regulation of transcription Response to glucose stimulus Steroid metabolic process Transcription | (Nadler et al., 2000; Soukas et al., 2000; Al-Hasani and Joost, 2005; Srivastava et al., 2006; Kim et al., 2008) |
| Glucose-6-phosphate dehydrogenase 2 (G6pd2) | ↑ | Carbohydrate metabolic process Glucose metabolic process Metabolic process Oxidation reduction | (Soukas et al., 2000; Soukas et al., 2001; Orellana-Gavalda et al., 2011) |
| Glycerol-3-phosphate dehydrogenase 1 (Gpd1) | ↓ | NADH metabolic process NADH oxidation Carbohydrate metabolic process Gluconeogenesis Glycerol-3-phosphate catabolic process Glycerolipid metabolic process | (Harada et al., 2003; Kim and Kim, 2009; Balwierz et al., 2009; Kalupahana et al., 2010) |
| Catalase (Cat) | ↓ | Aerobic respiration Cholesterol metabolic process Hydrogen peroxide catabolic process Response to oxidative stress Response to reactive oxygen species Triglyceride metabolic process | (Rong et al., 2007; Hagiwara et al., 2009; Coate and Huggins, 2010) |
| Cathepsin D (Ctsd) | ↑ | Autophagic vacuole assembly Proteolysis | (Nadler et al., 2000; Tanaka et al., 2003; Meugnier et al., 2007; Li et al., 2010) |
| CCAAT/enhancer binding protein α (Cebpa) | ↑ | Brown fat cell differentiation Cell maturation Cytokine-mediated signaling pathway Fat cell differentiation Regulation of cell proliferation Regulation of transcription Regulation of transcription, DNA-dependent | (Soukas et al., 2000; Harada et al., 2003; Kim et al., 2008) |

(Continued on next page)

Table 1 Lipid metabolism-associated genes differentially expressed in adipose tissue of HF-induced obese C57BL/6J mice (*Continued*)

| Gene name | Expression | Gene ontology (process) | References |
|---|------------|--|--|
| Superoxide dismutase 1, soluble (Sod1) | ↓ | Transcription White fat cell differentiation Activation of MAPK activity Anti-apoptosis Glutathione metabolic process Hydrogen peroxide biosynthetic process Response to oxidative stress Response to superoxide Superoxide anion generation Superoxide metabolic process | (Rong et al., 2007; Hagiwara et al., 2009; Hwang et al., 2009) |
| Plasminogen activator inhibitor 1 (Serpine 1, PAI-1) | ↑ | Negative regulation of Endopeptidase activity Positive regulation of coagulation Positive regulation of receptor-mediated endocytosis Regulation of angiogenesis Regulation of cell proliferation wound healing | (Moraes et al., 2003; Al-Hasani and Joost, 2005; Chen et al., 2005; Hosogai et al., 2007; Hagiwara et al., 2009) |
| Uncoupling protein 1 (mitochondrial, proton carrier) (Ucp1) | ↑ | Brown fat cell differentiation Mitochondrial transport Proton transport Transmembrane transport Transport | (Rong et al., 2007; Nikonova et al., 2008; Fukushima et al., 2009; Kim and Park, 2010) |
| Uncoupling protein 2 (mitochondrial, proton carrier) (Ucp2) | ↑ | mitochondrial transport response to superoxide transmembrane transport transport | (Harada et al., 2003; Morton et al., 2004; Fukushima et al., 2009; Orellana-Gavaldà et al., 2011) |

Wahli, 2007). There are two isoforms of *Pparg* which exist dependent on alternative promoter usage and splicing; *Pparg1* and *Pparg2*. *Pparg1* is ubiquitously found in WAT, brown adipose tissue (BAT), macrophages, liver, skeletal muscle, and other tissues (Chawla et al., 1994; Escher et al., 2001), whereas, *Pparg2* is mainly expressed in WAT (Rahimian et al., 2001; Tontonoz et al., 1994). There have been many studies on the role of *Pparg* in adipose tissue. For example, treatment of Zucker rats with thiazolidinedione (TZD), which is a specific and synthetic ligand of *Pparg* is reported to lead to reduced visceral adipocyte size (Ferre, 2004). In addition, the use of TZD in T2DM patients is reported to lower plasma glucose concentrations via activation of *Pparg* (Boden et al., 2003; Gastaldelli et al., 2007). Specific deletion of the *Pparg* gene in adipocytes causes markedly reduced adipocytes number, elevated plasma TG, and FFA, while plasma leptin and adiponectin are decreased (He et al., 2003). However, complete depletion of adipose tissue, as in the case of fatless mice, causes a phenotype characterized by higher insulin resistance, glucose intolerance, hyperphagia, hyperlipidemia, and fatty liver (Gavrilova et al., 2003). Furthermore, *Pparg* knock-out mice display similar phenotypes to fatless mice (Jones et al., 2005). These findings suggest that *Pparg* plays an important role in adipocyte differentiation and adipogenesis, as well as maintaining metabolic homeostasis.

Cebpa, is a transcription factor which appears to act coordinately with *Pparg* during adipogenesis (Clarke et al., 1997; Freytag et al., 1994; Farmer, 2005). The differentiation of preadipocytes into mature adipocytes is highly dependent on the expression of *Pparg* and *Cebpa* (Rosen et al., 2000). When

3T3-L1 preadipocytes are treated with differentiation inducing agents a rapid induction of C/EBP β and C/EBP δ occurs followed by increased expression of *Pparg* and *Cebpa* (Morrison and Farmer, 2000). Both *Pparg* and *Cebpa* are necessary for the differentiation of fibroblasts into adipocytes (Park et al., 2004). Past studies have established *Pparg* and *Cebpa* are responsible for coordinately directing the adipogenesis program governing adipose specific gene expression as well as the activation of other transcription factors (Wu et al., 1999).

Tcfap2a, also referred to as C/EBP undifferentiated protein has been identified as a *Cebpa* repressor (Jiang et al., 1998; Holt and Lane, 2001). In preadipocytes, *Cebpa* gene expression is repressed by *Tcfap2a* and repression is relieved during adipocyte differentiation by downregulation of *Tcfap2a* (Jiang et al., 1998). Furthermore, during differentiation of 3T3-L1 preadipocytes into mature adipocytes, the expression of *Cebpa* is increased and the kinetics of this process parallels the decline of *Tcfap2a* protein (Holt and Lane, 2001), indicating *Tcfap2a* is a suppressor of adipogenesis and fat cell development.

In addition to the adipocyte differentiation, *Pparg* plays a role in directing lipid storage in mature adipocytes (Rosen and Spiegelman, 2001; Way et al., 2001). *Pparg* promotes free fatty acid (FFAs) release from circulating lipoproteins and enhances FFA binding and uptake by regulating *Lpl* and *Fabp4* expression (Schoonjans et al., 1996; Ferre, 2004; Schachtrup et al., 2004). *Pparg* also promotes esterification of FFA to TG by regulating the expression of enzymes such as phosphoenol pyruvate carboxykinase (Olswang et al., 2002; Tontonoz et al., 1995), *Gpd1*, and diacylglycerol O acyltransferase (Anghel and Wahli, 2007).




Figure 1 Functional protein–protein interaction between obesity candidate genes in the adipose tissue of high-fat diet induced obese C57BL/6J mice. Each circle represents one gene in the data set. There are 25 genes in this subcomponent of the protein interaction. Circles with solid lines denote proteins in the one-depth condition (pink box), where circles with dotted lines denote proteins in the two-depth condition. The circles colored pink, yellow, and green denote obesity candidate genes, core obesity candidate gene, and the genes linking between nondirected obesity candidate genes in the one-depth condition, respectively. Arrows indicate up- or downregulation of gene expression upon HFD consumption. *Acaca*, acetyl-coenzyme A carboxylase; *Adipoq*, adiponectin; *Cat*, catalase; *Ctsd*, cathepsin D; *Cebpa*, CCAAT/enhancer-binding protein alpha; *Ccl2*, chemokine (C-C motif) ligand 2; *Cfd*, complement factor D; *Fabp4*, fatty acid binding protein 4; *Fasn*, fatty acid synthase; *G6pd2*, glucose-6-phosphate dehydrogenase 2; *Slc2a4*, glucose transporter 4; *Gpd1*, glycerol-3-phosphate dehydrogenase 1; *Hmgcs1*, 3-hydroxy-3-methylglutaryl-coenzyme A synthase 1; *Il6*, interleukin 6; *Lep*, leptin; *Lpl*, lipoprotein lipase; *Pparg*, peroxisome proliferator-activated receptor gamma; *PAI-1*, plasminogen activator inhibitor type 1; *Scd1*, stearoyl coenzyme A desaturase 1; *Srebf1*, sterol regulatory element-binding transcription factor 1; *Sod1*, superoxide dismutase 1; *Tnf*, tumor necrosis factor alpha; *Ucp1*, uncoupling protein 1; *Ucp2*, uncoupling protein 2. (Color figure available online.)

Natural ligands of *Pparg* include FAs such as oleic acid, linoleic acid, eicosapentaenoic acid, and arachidonic acid. Therefore, *Pparg* can act as a lipid sensor and modulate gene expression in response to changing FFA availability (Dreyer et al., 1992; Issemann and Green, 1990). Furthermore, *Pparg* also stimulates de novo FFA synthesis by regulating directly or indirectly the expression of *Fasn*, *Acaca*, and *Scd1* (Fig. 1). *Scd1* is expressed mainly in liver and adipose tissue, while adipose *Scd2* expression is strongly induced in HFD fed mice (Kaestner et al., 1989). Interestingly, loss of *Scd2* in mature adipocytes is reported to be associated with decreased *Pparg* protein level. In addition, *Scd2* depletion has been shown to decrease the metabolic labeling of *Pparg* protein but not the protein degradation rate, indicating that *Pparg* protein translation may be inhibited by *Scd2* (Christianson et al., 2008). It is clear from the obesity candidate gene network (Fig. 1) that *Pparg* regulates a sophisticated network of other transcription factors and their corepressors and coactivators such as *Tcfap2a*, *Ppara*, *Ppard*, and nuclear receptor subfamily 1 (*Nr1h3*). Therefore, *Pparg* is likely to have widespread effects on many more genes in adipose tissue, but these remain to be characterized.

Functional Protein Interactions of Adipokines in Obesity

Adipose tissue integrates hormonal signals from different body parts and responds to extracellular stimuli by secreting its own signaling molecules called adipokines, which effect the systemic regulation of energy homeostasis including lipid and glucose metabolism (Wajchenberg, 2000). Since adipose tissue is a mixture of adipocytes, macrophages, and stromal cells, a large proportion of adipokines are also released from cells other than adipocytes with the exception of adiponectin and leptin (Fain et al., 2004). Deregulation of adipokine secretion can lead to the development of metabolic syndrome, characterized by obesity and altered metabolism (Bergman et al., 2001; Grundy, 2004).

So far the most studied adipokine is a small 16 kDa polypeptide *Lep* (from the Greek *leptos*, meaning thin). It is produced mainly by adipocytes, and shows a strong positive correlation with the mass of adipose tissue. *Lep* directly binds to its receptors and inhibits appetite but the absence of *Lep* or its receptor leads into uncontrolled food intake and obesity (Halaas et al., 1995). However, this does not apply to all the cases. In several

obesity models, despite high levels of endogenous *Lep* or treatment with exogenous *Lep*, there was no improvement of obesity (Frederich et al., 1995). In fact, it has often been found that the level of circulating *Lep* is even higher in obese people compared to that of nonobese people (Considine et al., 1996). One explanation for these observations is that high sustained *Lep* concentrations result in *Lep* resistance similar to insulin resistance in T2DM. In contrast to *Lep*, *Ghrl* is a hormone mainly produced in the stomach that increases food intake by directly stimulating neuropeptide Y/agouti-related protein (NPY/AgRP) neurons in the arcuate nucleus of the hypothalamus (ARH) (Grove and Cowley, 2005). Since the levels of *Ghrl* increase greatly before meals and the levels decrease after meals, it has been suggested to be the gut signal that initiates a meal. It, therefore, is considered the counterpart of *Lep* which directly inhibits ARH NPY/AgRP through the *Lep* receptor and induces satiation (Cowley and Grove, 2004).

Adiponectin encoded by the *Adipoq* gene is one of the major adipokines secreted by adipocytes (Chandran et al., 2003). Plasma *Adipoq* is inversely correlated with the mass of adipose tissue, and *Adipoq* gene expression is reportedly down-regulated in obesity (Ukkola and Santaniemi, 2002). In addition, weight reduction significantly increases circulating *Adipoq* levels in humans (Coppola et al., 2009). In obese mice *Adipoq* reportedly stimulates the AMP activated protein kinase signaling pathway in skeletal muscle and the liver, thereby regulating insulin sensitivity, FA oxidation, and glucose metabolism (Yamauchi et al., 2002). Several studies suggest *Adipoq* plays a role in regulating insulin sensitivity. For example, plasma *Adipoq* is reported to be associated with visceral fat accumulation and insulin resistance in Japanese men with T2DM (Yatagai et al., 2003). Also it has been reported that plasma *Adipoq* decreases in parallel with reduced insulin sensitivity before T2DM development in Rhesus monkeys (Hotta et al., 2001). As shown in Fig. 1, decreased expression of *Adipoq* was also associated with decreased *Slc2a4* expression. In fact, *Adipoq* has been reported to increase insulin stimulated glucose uptake, via increased *Slc2a4* expression and recruitment of *Slc2a4* to the plasma membrane (Al-Hasani and Joost, 2005). Hypoadiponectinemia is reportedly more closely related to insulin resistance than the degree of obesity, and furthermore *Adipoq* appears to be an independent risk factor for metabolic syndrome (Renaldi et al., 2009; Weyer et al., 2001). *Adipoq* is also an important mediator of *Pparg* agonist-mediated insulin sensitivity improvement as well because administration of TZD increases *Adipoq* (Kamei et al., 2006).

Functional Protein Interactions of Inflammation-Associated Genes in Obesity

Some of the obesity candidate genes were associated with inflammation, which is in agreement with past studies suggesting a close interrelationship between obesity and inflammation (Wellen and Hotamisligil, 2003; Wellen and Hotamisligil,

2005), although the precise mechanisms have not been fully elucidated. *Cfd* also known as adipsin is a serine protease secreted by adipocytes. Since it is involved in the alternative complement pathway of the complement system, a high level of expression in adipocytes suggests a role for adipose tissue in immune system as well. It has been shown that *Cfd* also stimulates glucose transport for TG storage and inhibits lipolysis in adipocytes (Ronti et al., 2006). Macrophages are part of the innate immune defense system and significantly increased macrophage infiltration to adipose tissue is reported (Weisberg et al., 2003; Xu et al., 2003). Many macrophage and inflammation-related genes were also reported to be dramatically increased in WAT in an animal model of obesity (Xu et al., 2003). Colocalization of metabolically active adipocytes and macrophages in adipose tissue appears to amplify cytokine production, while infiltration of inflammatory cells leads to adipocyte dysfunction (Wellen and Hotamisligil, 2003). One study suggests that *Adipoq* may have anti-inflammatory effects by inhibiting the production of inflammation-associated cytokines such as *Tnfa* and *Il6* (Zhao et al., 2005).

Tnfa is a 26 kDa transmembrane protein that is cleaved to a smaller active 17 kDa protein (Anghel and Wahli, 2007). It is known to be involved in systemic inflammation; however, adipocyte *Tnfa* production exerts only a local effect since it can not be secreted. Hence, systemic *Tnfa* primarily originates from macrophages (Weisberg et al., 2006). Although there is no clear relationship between the levels of plasma *Tnfa* and obesity, the expression of *Tnfa* in WAT correlates well with obesity and insulin resistance (Bullo et al., 2002; Winkler et al., 2003). Chronic exposure of mice to *Tnfa* induces insulin resistance and decreases adipose tissue mass, whereas, *Tnfa* gene knockout improves insulin sensitivity in obese mice (Prins et al., 1997). *Tnfa* regulates expression of other adipokines in WAT as well. For example, the expression of *adipoq* is downregulated by *Tnfa*, whereas *Il6* is upregulated by *Tnfa* (Zhang et al., 1990). *Adipoq* inhibits *Tnfa* production, therefore increased *Tnfa* may actually result in an amplification of the inflammatory environment leading to adipocyte dysfunction. *Il6* is a multifunctional cytokine involved in cell-to-cell signal transduction in the immune system. It is abundantly expressed in many different cell types including immune cells, fibroblasts, and many endocrine cells (Tracey and Cerami, 1993; Van Snick, 1990). Majority of plasma *Il6*, however, is produced by adipocytes, especially visceral adipocytes (Anghel and Wahli, 2007). Levels of plasma *Il6* are reported positively correlated with obesity and insulin resistance, while decreased *Il6* improves insulin sensitivity similar to *Tnfa* (Bastard et al., 2002).

Tnfa also influences the level of *PAI-1* and *Ccl2*. *PAI-1* is a serine protease inhibitor protein (*serpin*) that is a major inhibitor of tissue plasminogen activator and urokinase which activates plasminogen activator. Thus, increased expression of *PAI-1* causes inhibition of fibrinolysis and the physiological breakdown of blood clots. *PAI-1* is mainly produced by the endothelium, but is also expressed and secreted by WAT in rodents and humans (Alessi et al., 2000; Pandey et al., 2005).

Circulating levels of PAI-1 are correlated with obesity and insulin resistance (Juhan-Vague et al., 2003; Alessi and Juhan-Vague, 2006); however, the underlying mechanisms are not well understood. It has been suggested that the inhibition of fibrinolysis by PAI-1 may be responsible for the high incidence of cardiovascular diseases associated with obesity (Skurk and Hauner, 2004). *Ccl2*, also known as monocyte chemoattractant protein-1 (MCP-1), is a small cytokine belonging to the C-C chemokine family. It recruits various inflammatory cells to the sites of inflammation (Carr et al., 1994; Xu et al., 1996) and the level of *Ccl2* is known to be increased in obesity (Weisberg et al., 2006). Increased expression of *Ccl2* in adipocytes is thought to be associated with the inflammatory response of adipose tissue in obesity. Chemokine (C-C motif) receptor 2 (CCR2) knock-out mice are characterized by reduced food intake, adipose tissue mass and macrophage numbers, combined with lower adipose tissue inflammation (Weisberg et al., 2006). In addition, CCR2 knock-out mice show elevated adiponectin, reduced steatosis, and improved insulin sensitivity (Weisberg et al., 2006). On the contrary, overexpression of *Ccl2* in adipose tissue results in increased macrophage infiltration (Kamei et al., 2006). TZD treatment in adipose tissue culture decreases *Ccl2* levels and similar effects are observed in T2DM patients as well (Bruun et al., 2005; Pfutzner et al., 2005). These findings suggest that activation of *Pparg* by *Pparg* agonists exerts beneficial effects on the production of a whole range of adipokines including *adipoq*, *Tnfa*, *Il6*, *PAI-1*, and *Ccl2*.

Functional Protein Interactions of Lipid Metabolism-Associated Genes in Obesity

Among the obesity candidate genes, some were associated with lipid and cholesterol metabolism. *Hmgcs1* is the first and rate-limiting enzyme of sterol biosynthesis which catalyzes acetyl-CoA to 3-hydroxy-3-methylglutaryl-CoA (HMG-CoA). In the adipocytes of HFD fed animals, the expression of *Hmgcs1* is downregulated, which has been suggested to be due to a feedback mechanism to maintain cholesterol homeostasis (Goldstein and Brown, 1990). ATP-binding cassette transporter 1 (*Abca1*), also known as the cholesterol efflux regulatory protein is a member of the ATP-binding cassette (ABC) transporters superfamily and is tightly regulated by transcriptional and posttranscriptional mechanisms (Luciani et al., 1994). *Abca1* is a major transporter responsible for phospholipid and cholesterol efflux (Schmitz and Langmann, 2001), which also regulates HDL formation (Oram and Vaughan, 2000). Mutations of *Abca1* therefore have been associated with familial HDL deficiency (Ordovas, 2000) and the pathogenesis of atherosclerosis (Guo et al., 2006). In obesity, the expression of *Abca1* is increased probably to maintain cholesterol homeostasis in adipocytes and to offer protective effects against atherosclerosis (Ferre, 2004).

Ctsd is a lysosomal aspartic protease and activated at pH 5 in the endosome of hepatocytes (Authier et al., 2002). Recently, it has been reported that *Ctsd* expression is upregu-

lated during adipocyte differentiation, and in adipose tissue of obese C57BL/6J mice as well as obese humans (Masson et al., 2011). Moreover, suppression of *Ctsd* expression in preadipocytes leads to lipid-depletion and inhibits the expression of adipocyte differentiation markers including *Pparg*, hormone sensitive lipase, and *Fabp4* (Masson et al., 2011). Interestingly, inhibition of *Ctsd* activity or expression reduced *Abca1* expression, whereas, the expression of *Ctsd* increased both *Abca1* mRNA expression and protein abundance (Haidar et al., 2006), which suggests that *Ctsd* plays an important role in intracellular cholesterol trafficking and *Abca1*-mediated cholesterol efflux. *Srebf1*, also known as sterol regulatory element binding protein 1 (Srebp1) is a member of the basic helix-loop-helix-leucine zipper transcription factor family (Yokoyama et al., 1993). *Srebf1* binds to the sterol regulatory element-1 (SRE1) in the promoter region of various genes involved in sterol biosynthesis and activates gene transcription. *Srebf1* is synthesized as a precursor protein in the endoplasmic reticulum. Thereafter *Srebf1* is cleaved, activated, and translocates to the nucleus. Sterol reportedly inhibits the cleavage of the precursor *Srebf1* and thereby reduces gene transcription. *Adipoq* administration suppresses *Srebp1c* expression in hepatocytes (Awazawa et al., 2009), whereas 3T3-L1 fibroblasts overexpressing *adipoq* exhibit more prolonged and robust *Srebp1c* expression (Al-Hasani and Joost, 2005). Seo et al. also showed that adenoviral overexpression of *Srebp1c* elevated *adipoq* mRNA and protein abundance in 3T3-L1 adipocytes (Seo et al., 2004). Furthermore, *Srebp1c* may play an important role in the production of a potent *Pparg* ligand (Kim et al., 1998). *G6pd2* is the first and rate-limiting enzyme of the pentose phosphate pathway (Spolarics, 1999). It has been reported that *G6pd2* is highly expressed in the adipocytes of obese animals and ectopic overexpression of *G6pd2* resulted in dysregulated lipid metabolism and insulin resistance (Park et al., 2005).

Functional Protein Interactions of Energy Metabolism and Oxidative Stress-Associated Genes in Obesity

Another interesting obesity candidate gene involved in energy metabolism was *Ucps*, which encodes a mitochondrial transmembrane protein that uncouples oxidative phosphorylation from ATP generation and dissipates energy as heat. *Ucp1* expression is known to be restricted to BAT (Nicholls et al., 1978). However, recent evidence showed that chronic treatment of *Pparg* agonist strongly induces *Ucp1* gene expression in WAT as well (Kim et al., 1998; Petrovic et al., 2010). Nevertheless, *Ucp1* induction in WAT alone is not sufficient to increase energy expenditure (Sell et al., 2004) as these cells retain WAT characteristics (Petrovic et al., 2010). Activation of *Ppara* leads to an increase in *Ucp2* expression and enzymes involved in β -oxidation. Unlike *Ucp1*, *Ucp2* is ubiquitously expressed and *Ucp2* expression is also increased by TZDs in adipocytes (Camirand et al., 1998). *Ucp* is an interesting obesity candidate gene, because potentially therapeutic activation of *Ucp* could




Figure 2 The schematic diagram of gene expression and regulation in the adipose tissue of obese mice. The expression of genes and the effect of gene expression is color coded. Red color: upregulation. Blue color: downregulation. (Color figure available online.)

help oxidize stored FFA in obese individuals, reducing lipid droplet and adipose tissue size.

Some obesity candidate genes were also associated with oxidative stress. In adipose tissue, increased influx of FFAs, glucose, and inflammatory cytokines leads to increased oxidation of energy substrates and ceramide formation, which can lead to generation of mitochondrial reactive oxygen species (ROS) causing oxidative stress (Evans et al., 2002; Evans et al., 2005; Eriksson, 2007). Emerging evidence suggests that intracellular accumulation of ROS is associated with insulin resistance and metabolic syndrome (Evans et al., 2002; Robertson, 2004; Evans et al., 2005). Furthermore, oxidative stress is also reported to be increased in patients with abdominal obesity (Fujita et al., 2006). *Sod1* is one of the three superoxide dismutases and converts harmful superoxide radicals to molecular oxygen and hydrogen peroxide. Hydrogen peroxide is also a harmful byproduct of normal metabolism and it must be quickly converted into less harmful substances (Zelko et al., 2002). *Cat*, a commonly found enzyme in nearly all aerobic organisms, degrades hydrogen peroxide to water and oxygen. Since the expression of both *Sod1* and *Cat* is downregulated, it is thought that this further exacerbates existing oxidative stress and insulin resistance. Whether obesity candidate genes associated with oxidative stress are useful therapeutic targets has not yet been established. It would be interesting to observe whether transgenic or knockout mice have altered responses to diet-induced obesity.

Our network analysis revealed interplay between obesity candidate genes with a variety of functions apparently coor-

inated by *Pparg* and *Cebpa*. Excessive fat intake increases the size and mass of adipose tissue and increases FA uptake (*Fabp4* and *Lpl*) (Fig. 2). Enhanced secretion of inflammatory cytokines (*Tnfa*, *Il6*, *Ccl2*, *Ctsd*, and *Serpine*) and decreased secretion of adipokine (*Adipoq*) affect food intake (*Lep* and *Ghrl*), insulin sensitivity (*Slc2a4* and *G6pd2*), FA synthesis (*Acaca*, *Fasn*, *Scd1*, and *Scd2*), FA oxidation (*Ucp-1* and *Ucp-2*), glucose metabolism (*Gpd1*), and inflammation. Oxidative stress is increased as a byproduct and this is further exacerbated by downregulation of detoxifying genes (*Sod* and *Cat*). In future studies, it may be worthwhile to use a similar approach to identify tissue specific obesity candidate genes.

The present meta-review of obesity candidate genes in adipose tissue is based on genes altered in diet-induced obesity, rather than genes underlying genetic predisposition to obesity. Fat mass- and obesity-associated gene (FTO) was one of the earliest obesity candidate genes identified by large genome-wide association studies (Frayling et al., 2007), but is primarily associated with genetic predisposition to obesity rather than diet induced obesity. Studies have shown in humans whom have at least one FTO risk allele, increased intake of food, impaired satiety, and increased susceptibility to T2DM (Speakman et al., 2008; Haupt et al., 2009; Tanofsky-Kraff et al., 2009). Furthermore in mice the expression of FTO gene is also modulated by food intake, body weight, and fat mass (Fischer et al., 2009; Church et al., 2010). However, the interactions between FTO and other obesity candidate genes still remain to be elucidated.

CONCLUDING REMARKS

Despite the enormous amount of microarray data obtained from diet induced obesity models, systematic approaches for interpreting findings from these studies are still lacking. In this review, we discussed findings using a literature-driven network analysis approach to search for obesity candidate genes in published microarray studies, which revealed the following:

1. Upregulated genes were associated with FA synthesis (*Acaca*, *Fasn*, *Lpl*, *Scd1*, and *Scd2*), inflammation (*Tnfa*, *Il6*, *Ccl2*, *Ctsd*, and *Serpine*), signal transduction and transporters (*Fabp4*, *Cfd*, *Lep*, *Pparg*, and *Cebpa*), and energy homeostasis (*Gpd1*, *Ucp-1*, and *Ucp-2*).
2. Downregulated genes include Adipoq and genes associated with sterol biosynthesis (*Hmgcs1* and *Srebf1*), insulin sensitivity (*Slc2a4* and *G6pd2*), and oxidative stress (*Sod* and *Cat*).
3. *Pparg* appeared to be a core obesity gene, which interacts with lipid metabolism and inflammation genes.

An important drawback using the literature-driven network analysis approach described is it will tend to identify only genes which have strong evidence of differential expression in diet-induced obesity. Nevertheless, it can help identify interactions between genes with strong evidence of dysregulation in diet induced obesity. If we assume that the next generation of obesity drug, functional food, or food components will be designed to inhibit an obesity gene/protein, it is really essential to look at how gene products (e.g., proteins) interact. So we know the potential side-effects of blocking a gene or protein to prevent an individual developing obesity. On top of that, it is also essential to develop better ways to coordinately interpret findings from past and future microarray studies, which will help identify therapeutic targets for the treatment of obesity.

ACKNOWLEDGMENT

This work was supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (MEST) (No. 2011-0000920). The authors have declared no conflict of interest.

ABBREVIATIONS

| | |
|--------|--------------------------------------|
| Acaca | Acetyl-Coenzyme A carboxylase |
| Adipoq | Adiponectin |
| Cat | Catalase |
| Ctsd | Cathepsin D |
| Cebpα | CCAAT/enhancer binding protein alpha |
| Ccl2 | Chemokine (C-C motif) ligand 2 |
| Cfd | Complement factor D |

| | |
|--------|--|
| Fabp4 | Fatty acid binding protein 4 |
| Fasn | Fatty acid synthase |
| G6pd2 | Glucose-6-phosphate dehydrogenase 2 |
| Slc2a4 | Glucose transporter 4 |
| Gpd1 | Glycerol-3-phosphate dehydrogenase 1 |
| Hmgcs1 | 3-hydroxy-3-methylglutaryl-coenzyme A synthase 1 |
| Il6 | Interleukin 6 |
| Lep | Leptin |
| Lpl | Lipoprotein lipase |
| Pparg | Peroxisome proliferator activated receptor gamma |
| PAI-1 | Plasminogen activator inhibitor type 1 |
| Scd1 | Stearoyl coenzyme A desaturase 1 |
| Srebf1 | Sterol regulatory element binding transcription factor 1 |
| Sod1 | Superoxide dismutase 1 |
| Tnfa | Tumor necrosis factor alpha |
| Ucp1 | Uncoupling protein 1 |
| Ucp2 | Uncoupling protein 2 |

REFERENCES

- Adan, R. A., Vanderschuren, L. J. and la Fleur, S. E. (2008). Anti-obesity drugs and neural circuits of feeding. *Trends. Pharmacol. Sci.* **29**:208–217.
- Al-Hasani, H. and Joost, H. G. (2005). Nutrition/diet-induced changes in gene expression in white adipose tissue. *Best. Pract. Res. Clin. Endocrinol. Metab.* **19**:589–603.
- Alessi, M. C., Bastelica, D., Morange, P., Berthet, B., Leduc, I., Verdier, M., Geel, O. and Juhan-Vague, I. (2000). Plasminogen activator inhibitor 1, transforming growth factor-beta1, and BMI are closely associated in human adipose tissue during morbid obesity. *Diabetes.* **49**:1374–1380.
- Alessi, M. C. and Juhan-Vague, I. (2006). PAI-1 and the metabolic syndrome: links, causes, and consequences. *Arterioscler. Thromb. Vasc. Biol.* **26**:2200–2207.
- An, H. J., Rim, H. K., Suh, S. E., Jeong, H. J., Um, J. Y., Hong, S. H. and Kim, H. M. (2010). Gamiwalbitang, composed of four herbs, controls body weight increase and lipid level elevation induced by a high-fat diet in mice. *Immunopharmacol. Immunotoxicol.* **32**:307–312.
- Anghel, S. I. and Wahli, W. (2007). Fat poetry: a kingdom for PPAR gamma. *Cell. Res.* **17**:486–511.
- Authier, F., Metioui, M., Fabrega, S., Kouach, M. and Briand, G. (2002). Endosomal proteolysis of internalized insulin at the C-terminal region of the B chain by cathepsin D. *J. Biol. Chem.* **277**:9437–9446.
- Awazawa, M., Ueki, K., Inabe, K., Yamauchi, T., Kaneko, K., Okazaki, Y., Bardeesy, N., Ohnishi, S., Nagai, R. and Kadowaki, T. (2009). Adiponectin suppresses hepatic SREBP1c expression in an AdipoR1/LKB1/AMPK dependent pathway. *Biochem. Biophys. Res. Commun.* **382**:51–56.
- Balwierz, A., Polus, A., Razny, U., Wator, L., Dyduch, G., Tomaszewska, R., Scherneck, S., Joost, H. and Dembinska-Kiec, A. (2009). Angiogenesis in the New Zealand obese mouse model fed with high fat diet. *Lipids. Health. Dis.* **8**:13.
- Barnea, M., Shamay, A., Stark, A. H. and Madar, Z. (2006). A high-fat diet has a tissue-specific effect on adiponectin and related enzyme expression. *Obesity (Silver Spring).* **14**:2145–2153.
- Bastard, J. P., Maachi, M., Van Nhieu, J. T., Jardel, C., Bruckert, E., Grimaldi, A., Robert, J. J., Capeau, J. and Hainque, B. (2002). Adipose tissue IL-6 content correlates with resistance to insulin activation of glucose uptake both in vivo and in vitro. *J. Clin. Endocrinol. Metab.* **87**:2084–2089.
- Bergman, R. N., Van Citters, G. W., Mittelman, S. D., Dea, M. K., Hamilton-Wessler, M., Kim, S. P. and Ellmerer, M. (2001). Central role of the adipocyte in the metabolic syndrome. *J. Invest. Med.* **49**:119–126.

- Bessesen, D. H. (2008). Update on obesity. *J. Clin. Endocrinol. Metab.* **93**:2027–2034.
- Boden, G., Cheung, P., Mozzoli, M. and Fried, S. K. (2003). Effect of thiazolidinediones on glucose and fatty acid metabolism in patients with type 2 diabetes. *Metabolism*. **52**:753–759.
- Bruun, J. M., Lihn, A. S., Pedersen, S. B. and Richelsen, B. (2005). Monocyte chemoattractant protein-1 release is higher in visceral than subcutaneous human adipose tissue (AT): implication of macrophages resident in the AT. *J. Clin. Endocrinol. Metab.* **90**:2282–2289.
- Bullo, M., Garcia-Lorda, P., Peinado-Onsurbe, J., Hernandez, M., Del Castillo, D., Argiles, J. M. and Salas-Salvado, J. (2002). TNF α expression of subcutaneous adipose tissue in obese and morbid obese females: relationship to adipocyte LPL activity and leptin synthesis. *Int. J. Obes. Relat. Metab. Disord.* **26**:652–658.
- Camirand, A., Marie, V., Rabelo, R. and Silva, J. E. (1998). Thiazolidinediones stimulate uncoupling protein-2 expression in cell lines representing white and brown adipose tissues and skeletal muscle. *Endocrinology*. **139**:428–431.
- Carr, M. W., Roth, S. J., Luther, E., Rose, S. S. and Springer, T. A. (1994). Monocyte chemoattractant protein 1 acts as a T-lymphocyte chemoattractant. *Proc. Natl. Acad. Sci. USA*. **91**:3652–3656.
- Chandran, M., Phillips, S. A., Ciaraldi, T. and Henry, R. R. (2003). Adiponectin: more than just another fat cell hormone? *Diabetes. Care*. **26**:2442–2450.
- Chawla, A., Schwarz, E. J., Dimaculangan, D. D. and Lazar, M. A. (1994). Peroxisome proliferator-activated receptor (PPAR) γ : adipose-predominant expression and induction early in adipocyte differentiation. *Endocrinology*. **135**:798–800.
- Chen, A., Mumick, S., Zhang, C., Lamb, J., Dai, H., Weingarth, D., Mudgett, J., Chen, H., MacNeil, D. J., Reitman, M. L. and Qian, S. (2005). Diet induction of monocyte chemoattractant protein-1 and its impact on obesity. *Obes. Res.* **13**:1311–1320.
- Christianson, J. L., Nicoloso, S., Straubhaar, J. and Czech, M. P. (2008). Stearoyl-CoA desaturase 2 is required for peroxisome proliferator-activated receptor γ expression and adipogenesis in cultured 3T3-L1 cells. *J. Biol. Chem.* **283**:2906–2916.
- Church, C., Moir, L., McMurray, F., Girard, C., Banks, G. T., Teboul, L., Wells, S., Bruning, J. C., Nolan, P. M., Ashcroft, F. M. and Cox, R. D. (2010). Overexpression of Fto leads to increased food intake and results in obesity. *Nat. Genet.* **42**:1086–1092.
- Clarke, S. L., Robinson, C. E. and Gimble, J. M. (1997). CAAT/enhancer binding proteins directly modulate transcription from the peroxisome proliferator-activated receptor γ 2 promoter. *Biochem. Biophys. Res. Commun.* **240**:99–103.
- Coate, K. C. and Huggins, K. W. (2010). Consumption of a high glycemic index diet increases abdominal adiposity but does not influence adipose tissue pro-oxidant and antioxidant gene expression in C57BL/6 mice. *Nutr. Res.* **30**:141–150.
- Cole, B. K., Keller, S. R., Wu, R., Carter, J. D., Nadler, J. L. and Nunemaker, C. S. (2010). Valsartan protects pancreatic islets and adipose tissue from the inflammatory and metabolic consequences of a high-fat diet in mice. *Hypertension*. **55**:715–721.
- Considine, R. V., Sinha, M. K., Heiman, M. L., Kriauciunas, A., Stephens, T. W., Nyce, M. R., Ohannesian, J. P., Marco, C. C., McKee, L. J., Bauer, T. L. and Caro, J. F. (1996). Serum immunoreactive-leptin concentrations in normal-weight and obese humans. *N. Engl. J. Med.* **334**:292–295.
- Coppola, A., Marfella, R., Coppola, L., Tagliamonte, E., Fontana, D., Liguori, E., Cirillo, T., Cafiero, M., Natale, S. and Astarita, C. (2009). Effect of weight loss on coronary circulation and adiponectin levels in obese women. *Int. J. Cardiol.* **134**:414–416.
- Cowley, M. A. and Grove, K. L. (2004). Ghrelin—satisfying a hunger for the mechanism. *Endocrinology*. **145**:2604–2606.
- Dreyer, C., Krey, G., Keller, H., Givel, F., Helftenbein, G. and Wahli, W. (1992). Control of the peroxisomal β -oxidation pathway by a novel family of nuclear hormone receptors. *Cell*. **68**:879–887.
- Eriksson, J. W. (2007). Metabolic stress in insulin's target cells leads to ROS accumulation — a hypothetical common pathway causing insulin resistance. *FEBS Lett.* **581**:3734–3742.
- Escher, P., Braissant, O., Basu-Modak, S., Michalik, L., Wahli, W. and Desvergne, B. (2001). Rat PPARs: quantitative analysis in adult rat tissues and regulation in fasting and refeeding. *Endocrinology*. **142**:4195–4202.
- Evans, J. L., Goldfine, I. D., Maddux, B. A. and Grodsky, G. M. (2002). Oxidative stress and stress-activated signaling pathways: a unifying hypothesis of type 2 diabetes. *Endocr. Rev.* **23**:599–622.
- Evans, J. L., Maddux, B. A. and Goldfine, I. D. (2005). The molecular basis for oxidative stress-induced insulin resistance. *Antioxid. Redox. Signal.* **7**:1040–1052.
- Fain, J. N., Madan, A. K., Hiler, M. L., Cheema, P. and Bahouth, S. W. (2004). Comparison of the release of adipokines by adipose tissue, adipose tissue matrix, and adipocytes from visceral and subcutaneous abdominal adipose tissues of obese humans. *Endocrinology*. **145**:2273–2282.
- Farmer, S. R. (2005). Regulation of PPAR γ activity during adipogenesis. *Int. J. Obes. (Lond)*. **29**(Suppl 1):S13–S16.
- Ferre, P. (2004). The biology of peroxisome proliferator-activated receptors: relationship with lipid metabolism and insulin sensitivity. *Diabetes*. **53**(Suppl 1):S43–S50.
- Fischer, J., Koch, L., Emmerling, C., Vierkotten, J., Peters, T., Bruning, J. C. and Ruther, U. (2009). Inactivation of the Fto gene protects from obesity. *Nature*. **458**:894–898.
- Flachs, P., Horakova, O., Brauner, P., Rossmeisl, M., Pecina, P., Franssen-van Hal, N., Ruzickova, J., Sponarova, J., Drahotova, Z., Vlcek, C. et al. (2005). Polyunsaturated fatty acids of marine origin upregulate mitochondrial biogenesis and induce β -oxidation in white fat. *Diabetologia*. **48**:2365–2375.
- Flier, J. S. (2004). Obesity wars: molecular progress confronts an expanding epidemic. *Cell*. **116**:337–350.
- Frayling, T. M., Timpson, N. J., Weedon, M. N., Zeggini, E., Freathy, R. M., Lindgren, C. M., Perry, J. R., Elliott, K. S., Lango, H., Rayner, N. W., Shields, B., Harries, L. W., Barrett, J. C., Ellard, S., Groves, C. J., Knight, B., Patch, A. M., Ness, A. R., Ebrahim, S., Lawlor, D. A., Ring, S. M., Ben-Shlomo, Y., Jarvelin, M. R., Sovio, U., Bennett, A. J., Melzer, D., Ferrucci, L., Loos, R. J., Barroso, I., Wareham, N. J., Karpe, F., Owen, K. R., Cardon, L. R., Walker, M., Hitman, G. A., Palmer, C. N., Doney, A. S., Morris, A. D., Smith, G. D., Hattersley, A. T., McCarthy, M. I. (2007) A common variant in the FTO gene is associated with body mass index and predisposes to childhood and adult obesity. *Science*. **316**:889–894.
- Frayn, K. N. (2002). Adipose tissue as a buffer for daily lipid flux. *Diabetologia*. **45**:1201–1210.
- Frayn, K. N., Fielding, B. A. and Karpe, F. (2005). Adipose tissue fatty acid metabolism and cardiovascular disease. *Curr. Opin. Lipidol.* **16**:409–415.
- Frederich, R. C., Hamann, A., Anderson, S., Lollmann, B., Lowell, B. B. and Flier, J. S. (1995). Leptin levels reflect body lipid content in mice: evidence for diet-induced resistance to leptin action. *Nat. Med.* **1**:1311–1314.
- Freytag, S. O., Paielli, D. L. and Gilbert, J. D. (1994). Ectopic expression of the CCAAT/enhancer-binding protein α promotes the adipogenic program in a variety of mouse fibroblastic cells. *Genes. Dev.* **8**:1654–1663.
- Friedman, J. M. (2003). A war on obesity, not the obese. *Science*. **299**:856–858.
- Fujita, K., Nishizawa, H., Funahashi, T., Shimomura, I. and Shimabukuro, M. (2006). Systemic oxidative stress is associated with visceral fat accumulation and the metabolic syndrome. *Circ. J.* **70**:1437–1442.
- Fukushima, Y., Kasuga, M., Nakao, K., Shimomura, I. and Matsuzawa, Y. (2009). Effects of coffee on inflammatory cytokine gene expression in mice fed high-fat diets. *J. Agric. Food Chem.* **57**:11100–11105.
- Gastaldelli, A., Ferrannini, E., Miyazaki, Y., Matsuda, M., Mari, A. and DeFronzo, R. A. (2007). Thiazolidinediones improve β -cell function in type 2 diabetic patients. *Am. J. Physiol. Endocrinol. Metab.* **292**:E871–E883.
- Gavrilova, O., Haluzik, M., Matsusue, K., Cutson, J. J., Johnson, L., Dietz, K. R., Nicol, C. J., Vinson, C., Gonzalez, F. J. and Reitman, M. L. (2003). Liver peroxisome proliferator-activated receptor γ contributes to hepatic steatosis, triglyceride clearance, and regulation of body fat mass. *J. Biol. Chem.* **278**:34268–34276.
- Goldstein, J. L. and Brown, M. S. (1990). Regulation of the mevalonate pathway. *Nature*. **343**:425–430.

- Grove, K. L. and Cowley, M. A. (2005). Is ghrelin a signal for the development of metabolic systems? *J. Clin. Invest.* **115**:3393–3397.
- Grundy, S. M. (2004). Obesity, metabolic syndrome, and cardiovascular disease. *J. Clin. Endocrinol. Metab.* **89**:2595–2600.
- Guerre-Millo, M. (2004). Adipose tissue and adipokines: for better or worse. *Diabetes. Metab.* **30**:13–19.
- Guo, Z. G., Wu, P. S., Li, J. H. and Lai, W. Y. (2006). [Modulation of inflammatory cytokines by ATP-binding cassette A1 in THP-1 macrophages]. *Nan. Fang. Yi. Ke. Da. Xue. Xue. Bao.* **26**:1269–1272.
- Hagiwara, S., Gohda, T., Tanimoto, M., Ito, T., Murakoshi, M., Ohara, I., Yamazaki, T., Matsumoto, M., Horikoshi, S., Funabiki, K. and Tomino, Y. (2009). Effects of pyridoxamine (K-163) on glucose intolerance and obesity in high-fat diet C57BL/6J mice. *Metabolism.* **58**:934–945.
- Haidar, B., Kiss, R. S., Sarov-Blat, L., Brunet, R., Harder, C., McPherson, R. and Marcel, Y. L. (2006). Cathepsin D, a lysosomal protease, regulates ABCA1-mediated lipid efflux. *J. Biol. Chem.* **281**:39971–39981.
- Halaas, J. L., Gajiwala, K. S., Maffei, M., Cohen, S. L., Chait, B. T., Rabinowitz, D., Lallone, R. L., Burley, S. K. and Friedman, J. M. (1995). Weight-reducing effects of the plasma protein encoded by the obese gene. *Science.* **269**:543–546.
- Harada, K., Shen, W. J., Patel, S., Natu, V., Wang, J., Osuga, J., Ishibashi, S. and Kraemer, F. B. (2003). Resistance to high-fat diet-induced obesity and altered expression of adipose-specific genes in HSL-deficient mice. *Am. J. Physiol. Endocrinol. Metab.* **285**:E1182–E1195.
- Hauner, H. (2005). Secretory factors from human adipose tissue and their functional role. *Proc. Nutr. Soc.* **64**:163–169.
- Haupt, A., Thamer, C., Staiger, H., Tschritter, O., Kirchhoff, K., Machicao, F., Haring, H. U., Stefan, N. and Fritsche, A. (2009). Variation in the FTO gene influences food intake but not energy expenditure. *Exp. Clin. Endocrinol. Diabetes.* **117**:194–197.
- He, W., Barak, Y., Hevener, A., Olson, P., Liao, D., Le, J., Nelson, M., Ong, E., Olefsky, J. M. and Evans, R. M. (2003). Adipose-specific peroxisome proliferator-activated receptor gamma knockout causes insulin resistance in fat and liver but not in muscle. *Proc. Natl. Acad. Sci. USA.* **100**:15712–15717.
- Hill, J. O., Peters, J. C., Catenacci, V. A. and Wyatt, H. R. (2008). International strategies to address obesity. *Obes Rev.* **9**(Suppl 1):41–47.
- Holt, E. H. and Lane, M. D. (2001). Downregulation of repressive CUP/AP-2 isoforms during adipocyte differentiation. *Biochem. Biophys. Res. Commun.* **288**:752–756.
- Hosogai, N., Fukuhara, A., Oshima, K., Miyata, Y., Tanaka, S., Segawa, K., Furukawa, S., Tochino, Y., Komuro, R., Matsuda, M. and Shimomura, I. (2007). Adipose tissue hypoxia in obesity and its impact on adipocytokine dysregulation. *Diabetes.* **56**:901–911.
- Hotta, K., Funahashi, T., Bodkin, N. L., Ortmeier, H. K., Arita, Y., Hansen, B. C. and Matsuzawa, Y. (2001). Circulating concentrations of the adipocyte protein adiponectin are decreased in parallel with reduced insulin sensitivity during the progression to type 2 diabetes in rhesus monkeys. *Diabetes.* **50**:1126–1133.
- Hwang, I. K., Kim, I. Y., Kim, Y. N., Yi, S. S., Park, I. S., Min, B. H., Doo, H. K., Ahn, S. Y., Kim, Y. S., Lee, I. S., Yoon, Y. S. and Seong, J. K. (2009). Comparative study on high fat diet-induced 4-hydroxy-2E-nonenal adducts in the hippocampal CA1 region of C57BL/6N and C3H/HeN mice. *Neurochem. Res.* **34**:964–972.
- Issemann, I. and Green, S. (1990). Activation of a member of the steroid hormone receptor superfamily by peroxisome proliferators. *Nature.* **347**:645–650.
- Jiang, M. S., Tang, Q. Q., McLenithan, J., Geiman, D., Shillinglaw, W., Henzel, W. J. and Lane, M. D. (1998). Derepression of the C/EBPalpha gene during adipogenesis: identification of AP-2alpha as a repressor. *Proc. Natl. Acad. Sci. USA.* **95**:3467–3471.
- Jones, J. R., Barrick, C., Kim, K. A., Lindner, J., Blondeau, B., Fujimoto, Y., Shiota, M., Kesterson, R. A., Kahn, B. B. and Magnuson, M. A. (2005). Deletion of PPARgamma in adipose tissues of mice protects against high fat diet-induced obesity and insulin resistance. *Proc. Natl. Acad. Sci. USA.* **102**:6207–6212.
- Juhan-Vague, I., Alessi, M. C., Mavri, A. and Morange, P. E. (2003). Plasminogen activator inhibitor-1, inflammation, obesity, insulin resistance and vascular risk. *J. Thromb. Haemost.* **1**:1575–1579.
- Kaestner, K. H., Ntambi, J. M., Kelly, T. J., Jr. and Lane, M. D. (1989). Differentiation-induced gene expression in 3T3-L1 preadipocytes. A second differentially expressed gene encoding stearoyl-CoA desaturase. *J. Biol. Chem.* **264**:14755–14761.
- Kalupahana, N. S., Claycombe, K., Newman, S. J., Stewart, T., Siriwardhana, N., Matthan, N., Lichtenstein, A. H. and Moustaid-Moussa, N. (2010). Eicosapentaenoic acid prevents and reverses insulin resistance in high-fat diet-induced obese mice via modulation of adipose tissue inflammation. *J. Nutr.* **140**:1915–1922.
- Kamei, N., Tobe, K., Suzuki, R., Ohsugi, M., Watanabe, T., Kubota, N., Ohtsuka-Kawatari, N., Kumagai, K., Sakamoto, K., Kobayashi, M. M., Yamauchi, T., Ueki, K., Oishi, Y., Nishimura, S., Manabe, I., Hashimoto, H., Ohnishi, Y., Ogata, H., Tokuyama, K., Tsunoda, M., Ide, T., Murakami, K., Nagai, R. and Kadowaki, T. (2006). Overexpression of monocyte chemoattractant protein-1 in adipose tissues causes macrophage recruitment and insulin resistance. *J. Biol. Chem.* **281**:26602–26614.
- Kanda, H., Tateya, S., Tamori, Y., Kotani, K., Hiasa, K., Kitazawa, R., Kitazawa, S., Miyachi, H., Maeda, S., Egashira, K. and Kasuga, M. (2006). MCP-1 contributes to macrophage infiltration into adipose tissue, insulin resistance, and hepatic steatosis in obesity. *J. Clin. Invest.* **116**:1494–1505.
- Kang, J. H., Goto, T., Han, I. S., Kawada, T., Kim, Y. M. and Yu, R. (2010). Dietary capsaicin reduces obesity-induced insulin resistance and hepatic steatosis in obese mice fed a high-fat diet. *Obesity (Silver Spring).* **18**:780–787.
- Kershaw, E. E. and Flier, J. S. (2004). Adipose tissue as an endocrine organ. *J. Clin. Endocrinol. Metab.* **89**:2548–2556.
- Kim, A. and Park, T. (2010). Diet-induced obesity regulates the galanin-mediated signaling cascade in the adipose tissue of mice. *Mol. Nutr. Food Res.* **54**:1361–1370.
- Kim, J. B., Wright, H. M., Wright, M. and Spiegelman, B. M. (1998). ADD1/SREBP1 activates PPARgamma through the production of endogenous ligand. *Proc. Natl. Acad. Sci. USA.* **95**:4333–4337.
- Kim, K. Y., Lee, H. N., Kim, Y. J. and Park, T. (2008). Garcinia cambogia extract ameliorates visceral adiposity in C57BL/6J mice fed on a high-fat diet. *Biosci. Biotechnol. Biochem.* **72**:1772–1780.
- Kim, M. J. and Kim, H. K. (2009). Perilla leaf extract ameliorates obesity and dyslipidemia induced by high-fat diet. *Phytother. Res.* **23**:1685–1690.
- Kim, Y. J. and Park, T. (2008). Genes are differentially expressed in the epididymal fat of rats rendered obese by a high-fat diet. *Nutr. Res.* **28**:414–422.
- Li, P., Lu, M., Nguyen, M. T., Bae, E. J., Chapman, J., Feng, D., Hawkins, M., Pessin, J. E., Sears, D. D., Nguyen, A. K., Amidi, A., Watkins, S. M., Nguyen, U. and Olefsky, J. M. (2010). Functional heterogeneity of CD11c-positive adipose tissue macrophages in diet-induced obese mice. *J. Biol. Chem.* **285**:15333–15345.
- Lopez, I. P., Marti, A., Milagro, F. I., Zulet Md Mde, L., Moreno-Aliaga, M. J., Martinez, J. A. and De Miguel, C. (2003). DNA microarray analysis of genes differentially expressed in diet-induced (cafeteria) obese rats. *Obes. Res.* **11**:188–194.
- Luciani, M. F., Denizot, F., Savary, S., Mattei, M. G. and Chimini, G. (1994). Cloning of two novel ABC transporters mapping on human chromosome 9. *Genomics.* **21**:150–159.
- Marrades, M. P., Milagro, F. I., Martinez, J. A. and Moreno-Aliaga, M. J. (2006). Differential expression of aquaporin 7 in adipose tissue of lean and obese high fat consumers. *Biochem. Biophys. Res. Commun.* **339**:785–789.
- Masson, O., Prebois, C., Derocq, D., Meulle, A., Dray, C., Daviaud, D., Quilliot, D., Valet, P., Muller, C. and Liadet-Coopman, E. (2011). Cathepsin-D, a key protease in breast cancer, is up-regulated in obese mouse and human adipose tissue, and controls adipogenesis. *PLoS One.* **6**:e16452.
- Meugnier, E., Bossu, C., Olié, M., Jeanne, S., Michaut, A., Sothier, M., Brozek, J., Rome, S., Laville, M. and Vidal, H. (2007). Changes in gene expression in skeletal muscle in response to fat overfeeding in lean men. *Obesity (Silver Spring).* **15**:2583–2594.
- Mong, M. C., Chao, C. Y. and Yin, M. C. (2011). Histidine and carnosine alleviated hepatic steatosis in mice consumed high saturated fat diet. *Eur. J. Pharmacol.* **653**:82–88.
- Moraes, R. C., Blondet, A., Birkenkamp-Demtroeder, K., Tirard, J., Orntoft, T. F., Gertler, A., Durand, P., Naville, D. and Begeot, M. (2003). Study of the alteration of gene expression in adipose tissue of diet-induced obese mice

- by microarray and reverse transcription-polymerase chain reaction analyses. *Endocrinology*. **144**:4773–4782.
- Morrison, R. F. and Farmer, S. R. (2000). Hormonal signaling and transcriptional control of adipocyte differentiation. *J. Nutr.* **130**:3116S–3121S.
- Morton, N. M., Paterson, J. M., Masuzaki, H., Holmes, M. C., Staels, B., Fievet, C., Walker, B. R., Flier, J. S., Mullins, J. J. and Seckl, J. R. (2004). Novel adipose tissue-mediated resistance to diet-induced visceral obesity in 11 beta-hydroxysteroid dehydrogenase type 1-deficient mice. *Diabetes*. **53**:931–938.
- Nadler, S. T., Stoehr, J. P., Schueler, K. L., Tanimoto, G., Yandell, B. S. and Attie, A. D. (2000). The expression of adipogenic genes is decreased in obesity and diabetes mellitus. *Proc. Natl. Acad. Sci. USA*. **97**:11371–11376.
- Nicholls, D. G., Bernson, V. S. and Heaton, G. M. (1978). The identification of the component in the inner membrane of brown adipose tissue mitochondria responsible for regulating energy dissipation. *Experientia. Suppl.* **32**:89–93.
- Nikonova, L., Koza, R. A., Mendoza, T., Chao, P. M., Curley, J. P. and Kozak, L. P. (2008). Mesoderm-specific transcript is associated with fat mass expansion in response to a positive energy balance. *Faseb. J.* **22**:3925–3937.
- Olswang, Y., Cohen, H., Papo, O., Cassuto, H., Croniger, C. M., Hakimi, P., Tilghman, S. M., Hanson, R. W. and Reshef, L. (2002). A mutation in the peroxisome proliferator-activated receptor gamma-binding site in the gene for the cytosolic form of phosphoenolpyruvate carboxykinase reduces adipose tissue size and fat content in mice. *Proc. Natl. Acad. Sci. USA*. **99**:625–630.
- Oram, J. F. and Vaughan, A. M. (2000). ABCA1-mediated transport of cellular cholesterol and phospholipids to HDL apolipoproteins. *Curr. Opin. Lipidol.* **11**:253–260.
- Ordovas, J. M. (2000). ABC1: the gene for Tangier disease and beyond. *Nutr. Rev.* **58**:76–79.
- Orellana-Gavalda, J. M., Herrero, L., Malandrino, M. I., Paneda, A., Sol Rodriguez-Pena, M., Petry, H., Asins, G., Van Deventer, S., Hegardt, F. G. and Serra, D. (2011). Molecular therapy for obesity and diabetes based on a long-term increase in hepatic fatty-acid oxidation. *Hepatology*. **53**:821–832.
- Pandey, M., Loskutoff, D. J. and Samad, F. (2005). Molecular mechanisms of tumor necrosis factor-alpha-mediated plasminogen activator inhibitor-1 expression in adipocytes. *Faseb. J.* **19**:1317–1319.
- Park, B. H., Qiang, L. and Farmer, S. R. (2004). Phosphorylation of C/EBPbeta at a consensus extracellular signal-regulated kinase/glycogen synthase kinase 3 site is required for the induction of adiponectin gene expression during the differentiation of mouse fibroblasts into adipocytes. *Mol. Cell. Biol.* **24**:8671–8680.
- Park, J., Rho, H. K., Kim, K. H., Choe, S. S., Lee, Y. S. and Kim, J. B. (2005). Overexpression of glucose-6-phosphate dehydrogenase is associated with lipid dysregulation and insulin resistance in obesity. *Mol. Cell. Biol.* **25**:5146–5157.
- Petrovic, N., Walden, T. B., Shabalina, I. G., Timmons, J. A., Cannon, B. and Nedergaard, J. (2010). Chronic peroxisome proliferator-activated receptor gamma (PPARgamma) activation of epididymally derived white adipocyte cultures reveals a population of thermogenically competent, UCP1-containing adipocytes molecularly distinct from classic brown adipocytes. *J. Biol. Chem.* **285**:7153–7164.
- Pfutzner, A., Marx, N., Lubben, G., Langenfeld, M., Walcher, D., Konrad, T. and Forst, T. (2005). Improvement of cardiovascular risk markers by pioglitazone is independent from glycemic control: results from the pioneer study. *J. Am. Coll. Cardiol.* **45**:1925–1931.
- Prins, J. B., Niesler, C. U., Winterford, C. M., Bright, N. A., Siddle, K., O'Rahilly, S., Walker, N. I. and Cameron, D. P. (1997). Tumor necrosis factor-alpha induces apoptosis of human adipose cells. *Diabetes*. **46**:1939–1944.
- Rahimian, R., Masih-Khan, E., Lo, M., van Breemen, C., McManus, B. M. and Dube, G. P. (2001). Hepatic over-expression of peroxisome proliferator activated receptor gamma2 in the ob/ob mouse model of non-insulin dependent diabetes mellitus. *Mol. Cell. Biochem.* **224**:29–37.
- Rajala, M. W. and Scherer, P. E. (2003). Minireview: The adipocyte—at the crossroads of energy homeostasis, inflammation, and atherosclerosis. *Endocrinology*. **144**:3765–3773.
- Renaldi, O., Pramono, B., Sinorita, H., Purnomo, L. B., Asdie, R. H. and Asdie, A. H. (2009). Hypoadiponectinemia: a risk factor for metabolic syndrome. *Acta. Med. Indones.* **41**:20–24.
- Robertson, R. P. (2004). Chronic oxidative stress as a central mechanism for glucose toxicity in pancreatic islet beta cells in diabetes. *J. Biol. Chem.* **279**:42351–42354.
- Rong, J. X., Qiu, Y., Hansen, M. K., Zhu, L., Zhang, V., Xie, M., Okamoto, Y., Mattie, M. D., Higashiyama, H., Asano, S., Strum, J. C. and Ryan, T. E. (2007). Adipose mitochondrial biogenesis is suppressed in db/db and high-fat diet-fed mice and improved by rosiglitazone. *Diabetes*. **56**:1751–1760.
- Ronti, T., Lupattelli, G. and Mannarino, E. (2006). The endocrine function of adipose tissue: an update. *Clin. Endocrinol. (Oxf)*. **64**:355–365.
- Rosen, E. D. and Spiegelman, B. M. (2001). PPARgamma: a nuclear regulator of metabolism, differentiation, and cell growth. *J. Biol. Chem.* **276**:37731–37734.
- Rosen, E. D., Walkey, C. J., Puigserver, P. and Spiegelman, B. M. (2000). Transcriptional regulation of adipogenesis. *Genes. Dev.* **14**:1293–1307.
- Schachtrup, C., Emmeler, T., Bleck, B., Sandqvist, A. and Spener, F. (2004). Functional analysis of peroxisome-proliferator-responsive element motifs in genes of fatty acid-binding proteins. *Biochem. J.* **382**:239–245.
- Schmitz, G. and Langmann, T. (2001). Structure, function and regulation of the ABC1 gene product. *Curr. Opin. Lipidol.* **12**:129–140.
- Schoonjans, K., Peinado-Onsurbe, J., Lefebvre, A. M., Heyman, R. A., Briggs, M., Deeb, S., Staels, B. and Auwerx, J. (1996). PPARalpha and PPARgamma activators direct a distinct tissue-specific transcriptional response via a PPRE in the lipoprotein lipase gene. *Embo. J.* **15**:5336–5348.
- Sell, H., Berger, J. P., Samson, P., Castriota, G., Lalonde, J., Deshaies, Y. and Richard, D. (2004). Peroxisome proliferator-activated receptor gamma agonism increases the capacity for sympathetically mediated thermogenesis in lean and ob/ob mice. *Endocrinology*. **145**:3925–3934.
- Seo, J. B., Moon, H. M., Noh, M. J., Lee, Y. S., Jeong, H. W., Yoo, E. J., Kim, W. S., Park, J., Youn, B. S., Kim, J. W., Park, S. D. and Kim, J. B. (2004). Adipocyte determination- and differentiation-dependent factor 1/sterol regulatory element-binding protein 1c regulates mouse adiponectin expression. *J. Biol. Chem.* **279**:22108–22117.
- Sharma, A. M. and Staels, B. (2007). Review: Peroxisome proliferator-activated receptor gamma and adipose tissue—understanding obesity-related changes in regulation of lipid and glucose metabolism. *J. Clin. Endocrinol. Metab.* **92**:386–395.
- Skurk, T. and Hauner, H. (2004). Obesity and impaired fibrinolysis: role of adipose production of plasminogen activator inhibitor-1. *Int. J. Obes. Relat. Metab. Disord.* **28**:1357–1364.
- Soukas, A., Cohen, P., Socci, N. D. and Friedman, J. M. (2000). Leptin-specific patterns of gene expression in white adipose tissue. *Genes. Dev.* **14**:963–980.
- Soukas, A., Socci, N. D., Saatkamp, B. D., Novelli, S. and Friedman, J. M. (2001). Distinct transcriptional profiles of adipogenesis in vivo and in vitro. *J. Biol. Chem.* **276**:34167–34174.
- Speakman, J. R., Rance, K. A. and Johnstone, A. M. (2008). Polymorphisms of the FTO gene are associated with variation in energy intake, but not energy expenditure. *Obesity. (Silver Spring)*. **16**:1961–1965.
- Spiegelman, B. M. and Flier, J. S. (2001). Obesity and the regulation of energy balance. *Cell*. **104**:531–543.
- Spolarics, Z. (1999). A carbohydrate-rich diet stimulates glucose-6-phosphate dehydrogenase expression in rat hepatic sinusoidal endothelial cells. *J. Nutr.* **129**:105–108.
- Srivastava, R. A., Jahagirdar, R., Azhar, S., Sharma, S. and Bisgaier, C. L. (2006). Peroxisome proliferator-activated receptor-alpha selective ligand reduces adiposity, improves insulin sensitivity and inhibits atherosclerosis in LDL receptor-deficient mice. *Mol. Cell. Biochem.* **285**:35–50.
- Tanaka, T., Yamamoto, J., Iwasaki, S., Asaba, H., Hamura, H., Ikeda, Y., Watanabe, M., Magoori, K., Ioka, R. X., Tachibana, K., Watanabe, Y., Uchiyama, Y., Sumi, K., Iguchi, H., Ito, S., Doi, T., Hamakubo, T., Naito, M., Auwerx, J., Yanagisawa, M., Kodama, T. and Sakai, J. (2003). Activation of peroxisome proliferator-activated receptor delta induces fatty acid beta-oxidation in skeletal muscle and attenuates metabolic syndrome. *Proc. Natl. Acad. Sci. USA*. **100**:15924–15929.
- Tanofsky-Kraff, M., Han, J. C., Anandalingam, K., Shomaker, L. B., Wolkoff, L. E., Kozlosky, M., Elliott, C., Ranzenhofner, L. M., Roza, C. A., Yanovski,

- S. Z. and Yanovski, J. A. (2009). The FTO gene rs9939609 obesity-risk allele and loss of control over eating. *Am. J. Clin. Nutr.* **90**:1483–1488.
- Tontonoz, P., Hu, E., Devine, J., Beale, E. G. and Spiegelman, B. M. (1995). PPAR gamma 2 regulates adipose expression of the phosphoenolpyruvate carboxykinase gene. *Mol. Cell. Biol.* **15**:351–357.
- Tontonoz, P., Hu, E., Graves, R. A., Budavari, A. I. and Spiegelman, B. M. (1994). mPPAR gamma 2: tissue-specific regulator of an adipocyte enhancer. *Genes. Dev.* **8**:1224–1234.
- Tracey, K. J. and Cerami, A. (1993). Tumor necrosis factor, other cytokines and disease. *Annu. Rev. Cell Biol.* **9**:317–343.
- Trayhurn, P. and Wood, I. S. (2004). Adipokines: inflammation and the pleiotropic role of white adipose tissue. *Br. J. Nutr.* **92**:347–355.
- Ukkola, O. and Santaniemi, M. (2002). Adiponectin: a link between excess adiposity and associated comorbidities? *J. Mol. Med.* **80**:696–702.
- Van Schothorst, E. M., Franssen-van Hal, N., Schaap, M. M., Pennings, J., Hoebee, B. and Keijer, J. (2005). Adipose gene expression patterns of weight gain suggest counteracting steroid hormone synthesis. *Obes. Res.* **13**:1031–1041.
- Van Snick, J. (1990). Interleukin-6: an overview. *Annu. Rev. Immunol.* **8**:253–278.
- Wajchenberg, B. L. (2000). Subcutaneous and visceral adipose tissue: their relation to the metabolic syndrome. *Endocr. Rev.* **21**:697–738.
- Way, J. M., Harrington, W. W., Brown, K. K., Gottschalk, W. K., Sundseth, S. S., Mansfield, T. A., Ramachandran, R. K., Willson, T. M. and Klierer, S. A. (2001). Comprehensive messenger ribonucleic acid profiling reveals that peroxisome proliferator-activated receptor gamma activation has coordinate effects on gene expression in multiple insulin-sensitive tissues. *Endocrinology*. **142**:1269–1277.
- Weisberg, S. P., Hunter, D., Huber, R., Lemieux, J., Slaymaker, S., Vaddi, K., Charo, I., Leibel, R. L. and Ferrante, A. W., Jr. (2006). CCR2 modulates inflammatory and metabolic effects of high-fat feeding. *J. Clin. Invest.* **116**:115–124.
- Weisberg, S. P., McCann, D., Desai, M., Rosenbaum, M., Leibel, R. L. and Ferrante, A. W., Jr. (2003). Obesity is associated with macrophage accumulation in adipose tissue. *J. Clin. Invest.* **112**:1796–1808.
- Wellen, K. E. and Hotamisligil, G. S. (2003). Obesity-induced inflammatory changes in adipose tissue. *J. Clin. Invest.* **112**:1785–1788.
- Wellen, K. E. and Hotamisligil, G. S. (2005). Inflammation, stress, and diabetes. *J. Clin. Invest.* **115**:1111–1119.
- Weyer, C., Funahashi, T., Tanaka, S., Hotta, K., Matsuzawa, Y., Pratley, R. E. and Tataranni, P. A. (2001). Hypoadiponectinemia in obesity and type 2 diabetes: close association with insulin resistance and hyperinsulinemia. *J. Clin. Endocrinol. Metab.* **86**:1930–1935.
- Winkler, G., Kiss, S., Keszthelyi, L., Sapi, Z., Ory, I., Salamon, F., Kovacs, M., Vargha, P., Szekeres, O., Speer, G., Karadi, I., Sikter, M., Kaszas, E., Dworak, O., Gero, G. and Cseh, K. (2003). Expression of tumor necrosis factor (TNF)-alpha protein in the subcutaneous and visceral adipose tissue in correlation with adipocyte cell volume, serum TNF-alpha, soluble serum TNF-receptor-2 concentrations and C-peptide level. *Eur. J. Endocrinol.* **149**:129–135.
- Wu, Z., Rosen, E. D., Brun, R., Hauser, S., Adelmant, G., Troy, A. E., McKeon, C., Darlington, G. J. and Spiegelman, B. M. (1999). Cross-regulation of C/EBP alpha and PPAR gamma controls the transcriptional pathway of adipogenesis and insulin sensitivity. *Mol. Cell.* **3**:151–158.
- Xu, H., Barnes, G. T., Yang, Q., Tan, G., Yang, D., Chou, C. J., Sole, J., Nichols, A., Ross, J. S., Tartaglia, L. A. and Chen, H. (2003). Chronic inflammation in fat plays a crucial role in the development of obesity-related insulin resistance. *J. Clin. Invest.* **112**:1821–1830.
- Xu, L. L., Warren, M. K., Rose, W. L., Gong, W. and Wang, J. M. (1996). Human recombinant monocyte chemotactic protein and other C-C chemokines bind and induce directional migration of dendritic cells in vitro. *J. Leukoc. Biol.* **60**:365–371.
- Yamauchi, T., Kamon, J., Minokoshi, Y., Ito, Y., Waki, H., Uchida, S., Yamashita, S., Noda, M., Kita, S., Ueki, K. et al. (2002). Adiponectin stimulates glucose utilization and fatty-acid oxidation by activating AMP-activated protein kinase. *Nat. Med.* **8**:1288–1295.
- Yatagai, T., Nagasaka, S., Taniguchi, A., Fukushima, M., Nakamura, T., Kuroe, A., Nakai, Y. and Ishibashi, S. (2003). Hypoadiponectinemia is associated with visceral fat accumulation and insulin resistance in Japanese men with type 2 diabetes mellitus. *Metabolism*. **52**:1274–1278.
- Yokoyama, C., Wang, X., Briggs, M. R., Admon, A., Wu, J., Hua, X., Goldstein, J. L. and Brown, M. S. (1993). SREBP-1, a basic-helix-loop-helix-leucine zipper protein that controls transcription of the low density lipoprotein receptor gene. *Cell*. **75**:187–197.
- Zelko, I. N., Mariani, T. J. and Folz, R. J. (2002). Superoxide dismutase multi-gene family: a comparison of the CuZn-SOD (SOD1), Mn-SOD (SOD2), and EC-SOD (SOD3) gene structures, evolution, and expression. *Free. Radic. Biol. Med.* **33**:337–349.
- Zhang, Y. H., Lin, J. X. and Vilcek, J. (1990). Interleukin-6 induction by tumor necrosis factor and interleukin-1 in human fibroblasts involves activation of a nuclear factor binding to a kappa B-like sequence. *Mol. Cell. Biol.* **10**:3818–3823.
- Zhao, T., Hou, M., Xia, M., Wang, Q., Zhu, H., Xiao, Y., Tang, Z., Ma, J. and Ling, W. (2005). Globular adiponectin decreases leptin-induced tumor necrosis factor-alpha expression by murine macrophages: involvement of cAMP-PKA and MAPK pathways. *Cell. Immunol.* **238**:19–30.