



Fillet Proximate Composition, Lipid Quality, Yields and Organoleptic Quality of Mediterranean Farmed Marine Fish: A Review with Emphasis on New Species

Kriton Grigorakis

To cite this article: Kriton Grigorakis (2015): Fillet Proximate Composition, Lipid Quality, Yields and Organoleptic Quality of Mediterranean Farmed Marine Fish: A Review with Emphasis on New Species, Critical Reviews in Food Science and Nutrition, DOI: [10.1080/10408398.2015.1081145](https://doi.org/10.1080/10408398.2015.1081145)

To link to this article: <http://dx.doi.org/10.1080/10408398.2015.1081145>



Accepted author version posted online: 15 Oct 2015.



Submit your article to this journal [↗](#)



Article views: 30



View related articles [↗](#)



View Crossmark data [↗](#)

**FILLET PROXIMATE COMPOSITION, LIPID QUALITY, YIELDS AND
ORGANOLEPTIC QUALITY OF MEDITERRANEAN FARMED MARINE FISH: A
REVIEW WITH EMPHASIS ON NEW SPECIES**

Kriton Grigorakis

Institute of Marine Biology, Biotechnology and Aquaculture, Hellenic Centre for Marine
Research, Agios Kosmas Hellinikon, 16777, Athens, Greece

Corresponding Author Email: Tel:+30 210 9856735; Fax: + 30 210 9829239; Email:
kgrigo@hcmr.gr

Abstract

Species diversification in Mediterranean mariculture involves various important fish that contribute to the diet of many human populations. These include meagres (Sciaenidae), flatfishes, mullets, and various sparids. Their quality aspects are discussed in this review (yields, fillet proximate composition and lipid quality). Their filleting yield is mostly 40-45%. The viscerosomatic index ranges from 1.5 to 14%, depending on species. Flatfishes' and meagres' low muscle fat contents, differentiate them from the rest of the farmed species. Farmed fish contain high n-3 PUFA (12.3–36.3% vs. 5.48-37.2% in the wild) and have higher muscle fat and n-6 PUFA contents (mainly 18:2n-6) than their wild counterparts. The aquaculture management, diet, and season can affect fillet composition and fatty acids, while season (i.e. food availability

and maturation) largely affects lipid quality in wild fish. Data on the sensory quality of Mediterranean farmed species mainly limit to whether specific management differentiates the sensory quality; thus further development of sensory analysis tools is required. Observations on the quality features in farmed Mediterranean fish indicate that species diversification can also provide product diversification, based on different commercial weights and fillet quality specifications.

Keywords

quality, fat, filleting yield, fatty acids, sensory quality

Introduction

The farmed finfish production worldwide was 66.6 million metric tons in 2012 (FAO, 2014). The Mediterranean area is an important contributor to world aquaculture, with two countries, Egypt and Spain being among the top 12 and 20 world aquaculture producers, respectively (FAO, 2013; FAO, 2014). The Mediterranean marine finfish production has been growing, to reach a share of 36% of its total aquaculture output vs. 48% for freshwater fish and 14% for molluscs. Specifically, marine finfish production in the Mediterranean area has grown from 61,024 tons in 1995 to 436,401 tons in 2007 (Barazi-Yeroulanos, 2010). The spark for the continuous growth of the contemporary Mediterranean mariculture was in the late 70s or early 80s. The Mediterranean marine finfish farming all these years has been dominated by two species, the European sea bass (*Dicentrarchus labrax*) and the gilthead sea bream (*Sparus aurata*). These two species are industrially produced and account for approximately 52% of the total marine finfish production in the area (Table 1). The other dominating species, the flathead grey mullet (*Mugil cephalus*), is traditionally semi-intensively farmed in brackish water ponds, and its production is mainly based on wild fry collection (Saleh, 2008). However, there have been some serious fry overfishing issues and recent pressure towards wild fry banning (McGrath, 2012).

In the 90s, the ranching of Atlantic bluefin tuna, *Thunnus thynnus*, was introduced in the area, based on wild stocks capture, since no integrated production could be achieved. The latter has led to almost crashing of the species' stocks in the Mediterranean (MacKenzie et al., 2009). Other sparid species (Table 1) have been farmed with variable success. The market saturation for gilthead sea bream and sea bass and the crisis of this sector during the 90s has led to a persistent

recession situation that was enhanced by a general economic crisis (Cardia and Lovatelli, 2007; Perdicaris and Paschos, 2011). Species diversification has been considered a possible way out from this situation (Cardia and Lovatelli, 2007). Among the candidates, various species, mainly from Sciaenidae and Caragidae families, have been proposed and produced nowadays, with meagre (*Argyrosomus regius*) being the most successful one (in terms of production and know-how) (Table 1). Among, the flatfish, the most commercialized in Mediterranean countries, is turbot (*Psetta maxima*), mainly produced in Spain since the 90's. Its farming is integrated and intensive rearing is mostly land-based (FAO 2005-2015).

Regarding their commercial characteristics, these fish species have various commercialization sizes and forms. The grey mullet, besides being sold as whole fish at sizes of about 300-800g, is used for its roe to produce a highly valued traditional product named bottarga (Barra et al., 2008). The Scieanidae species that include the meagre, the brown meagre (*Sciaena umbra*) and the shi drum (*Umbrina cirrosa*) are fast growers that are usually commercialized in sizes bigger than 1 Kg. Although they are usually commercialized as whole, for larger fish various forms like cuts and fillets can be available (Monfort, 2010). Also, small quantities of frozen fish, smoked fillets and sushi sales have been reported for meagre (Monfort 2010). Usual commercialization weights for the various sparid species are similar to those for gilthead sea bream, i.e. 350-500g (Hernández et al., 2001; Barazi-Yeroulanos, 2010) Therefore, they are sold in fresh, whole or gutted forms. The Carangidae family members have large commercialization sizes, and forms similar to the tuna fish, e.g. 3.5-5.5 Kg for greater amberjack (*Seriola dumerili*), sold either in sushi markets or in various cuts (Nakada, 2008). Flatfish, on the other hand, are sold in sizes

starting from 125g up to 8Kg as whole or fillets, fresh or frozen (Howell, 1997; Imsland et al., 2003).

One significant aspect of the market fate of farmed fish is their quality as perceived by the consumer. Two important aspects in the quality-defining set of parameters, are the nutritional and the technical quality of the fish. The former refers to the nutritional value of the food, while the latter to the processing technical losses and the edible yields. The nutritional significance of fish mainly refers to the contents in polyunsaturated fatty acids and their numerous health benefits (Kris-Etherton et al., 2003; Nicholson et al., 2013). This also reflects to the nutritional status image that the fish has among the consumer (Vanhonacker et al., 2013). The technical quality, on the other side, is mainly an economic aspect and is of interest for both consumers and processors. Furthermore, the organoleptic quality of the fish, i.e. the human sensory impression, is among the capital factors for purchasing those (Grigorakis, 2007).

The aim of this study is to critically review the aforementioned quality aspects for new Mediterranean farmed fish, along with a comparison with their wild counterparts where possible. The importance of examining the quality aspects of these species lies within their universal significance as food, since they are part of the diet, not only in the Mediterranean area but also in many other countries. Besides, they have a great potential for expansion in their production and consumption (Barazi-Yeroulanos, 2010).

Although there is plenty of literature directly comparing farmed and wild-caught gilthead sea bream and sea bass for their quality features (indicatively: Krajnovic-Ozretic et al., 1994; Alasalvar et al., 2002; Grigorakis et al., 2002; Mnari et al., 2007; Periago et al., 2005), there is a

scarcity in similar studies for the rest of the Mediterranean fish. Existing data refer only to some of the farmed species and to specific geographic areas (Rueda et al., 1997; Rueda et al., 2001; Cejas et al., 2004; Álvarez et al., 2009; Álvarez et al., 2009b; Dincer et al., 2010). A comparison by reviewing existing literature would elucidate actual differences in quality of farmed and wild Mediterranean fish. Additionally inter-species quality comparisons would allow a better knowledge of how quality differentiates among them and how these individualities could be of commercial advantage. Since there was a similar attempt in the past for gilthead sea bream and sea bass (Grigorakis, 2007; Arechavala-Lopez et al., 2013), this study will mainly focus on the rest of the Mediterranean farmed fish species.

Methods

Data and calculations

The somatic yields that were reviewed or calculated from the respective literature were the following:

$$\text{Filleting yield (FY)} = \left[100 \times (\text{fillet weight} / \text{body weight}) \right],$$

$$\text{Viscerosomatic index (VSI)} = \left[100 \times (\text{total viscera weight} / \text{body weight}) \right],$$

$$\text{Hepatosomatic index (HSI)} = \left[100 \times (\text{liver weight} / \text{body weight}) \right] \text{ and}$$

$$\text{Condition index (CI)} = \left[100 \times \text{body weight (g)} / \text{body length}^3 (\text{cm}^3) \right]$$

The fillet quality was assessed in terms of total composition (moisture, protein, ash and fat percentages) and in terms of fatty acid composition. For the latter, the percentages of the main fatty acid groups were calculated, i.e. saturates (SUFA), monounsaturates (MUFA), n-3 polyunsaturates (n-3 PUFA) and n-6 polyunsaturates (n-6 PUFA), as well as the two important n-3 fatty acids, the eicosapentaenoic (20:5n-3, EPA) and docohexaenoic (22:6n-3, DHA).

Besides, for estimation of the actual nutritional value of n-3 PUFA their total contents and the contents of EPA and DHA in 100g fish fillet were calculated as:

$$(\text{Fatty acid \%} \times \text{fillet fat \%}) / 100.$$

For all of the above factors, the weighed mean was calculated in the wild and farmed fish.

Study restrictions and limitations

It is important to notice that all data considered within this study refer to fish near or at commercial weights. Furthermore, for species with wider geographical distribution, the data for wild fish limits only in Mediterranean populations, since the environment has an important impact in all biological parameters and therefore in the end product quality (Love, 1992). Thus, taking into account fish of the same species but different origins (e.g. the Atlantic, the Black Sea or the Red Sea) was considered to be misleading.

Among the farmed species in the Mediterranean, the Atlantic bluefin tuna was not taken into account for this study since its aquaculture is far from achieving a complete cycle production, includes only fattening and is based on feeding with fresh fish instead of commercial diets (Grigorakis and Rigos, 2011; Hattour and Couchet, 2014).

The red drum (*Sciaenops ocellatus*) was not included in this study either. Although its farming has been established in Israel since the '90s (Diamant, 1998), this decision was due to the facts that this species does not naturally inhabit the Mediterranean (FishBase, 2011) and that all sources referring to its quality features refer to geographic origins other than the Mediterranean (Craig et al., 2000; Recks and Seaborn, 2008; Li et al., 2013).

A small part of the literature referring to the quality of the studied species was not taken into account for not providing information that allow calculations on whole wet fillet basis (Hernández et al., 2001; Hernández et al., 2003; Bonaldo et al., 2004; Rondán et al., 2004; Lloret et al., 2005; Testi et al., 2006; Álvarez et al., 2009; Álvarez et al., 2009b; Haouas et al., 2010; Nogales Merida et al., 2011; Valente et al., 2011), and for calculating fatty acid compositions of lipid fractions without providing the relative proportions of these lipid fractions (Regost et al., 2003; Varljen et al., 2003).

For all studied parameters weighed means were calculated, taking into account the literature average values and the number of individuals analyzed in each study. In cases that sample size was not evident (Rueda et al., 1997; Soriguer et al., 1997; El-Dakar et al., 2007; Koubaa et al., 2011), a $n=3$ was assumed on the basis that this is the minimum size for statistical analysis in the respective studies.

Metanalysis

For comparisons of average values \bar{X} between farmed (F) and wild (W) individuals of the same species, effect size was evaluated by the use of Hedges d (Hedges and Olkin 1985):

$$d = \frac{\bar{X}_F - \bar{X}_W}{s_p} J \quad (1)$$

where the correction for bias

$$J = 1 - \frac{3}{4(n_F + n_W) - 9} \quad (2)$$

and

$$s_p = \sqrt{\frac{(n_F - 1)se_F^2 + (n_W - 1)se_W^2}{n_F + n_W - 2}} \quad (3)$$

The asymptotic standard error (se) of the effect size was calculated as following:

$$s_e = \sqrt{\frac{n_F + n_W}{n_F n_W} + \frac{d^2}{2(n_F + n_W - 2)}} \quad (4)$$

Precision of d with 95% confidence intervals was evaluated as $d - 1.96se$ to $d + 1.96se$.

Biometric parameters and yields

The filleting yield is an important parameter, especially for species that filleting is among their custom processing, because it describes their actual edible gain. The respective data for the Mediterranean farmed fish is actually limited to some species, mainly meagre, dentex (*Dentex dentex*) and shi drum (Table 2). For the rest of the species, lack of data for some of them (like porgies *Pagellus* sp., annular seabream *Diplodus annularis* and mullets *Mugil* sp. and *Liza* sp.) may be due to the fact that they are not customly commercialized in filleted forms. However, for some other such as the flatfishes (*Solea* sp., *Psetta maxima*), red seabream (*Pagrus* sp.), dentex, greater amberjack and brown meagre, filleting is of interest and the collection of respective data should be within the aims of future research. Existing data show that the filleting yields for most species are around 40-45 % with few exceptions of extremely high and low yields reported for sharpsnout seabream (*Diplodus puntazzo*) and white seabream (*Diplodus sargus*), respectively (Table 2).

The viscerosomatic index (VSI) is a measure for the estimation of the visceral fat deposition and together with the hepatosomatic index (HSI) that represents the hepatic weight (Table 2), consist a way to evaluate the technical loss from fish gutting. The species with the lowest visceral losses are the flatfishes (*Solea senegalensis* and *Psetta maxima*), as becomes evident from the VSI values in Table 2. Also the Sciaenidae family species (*Argyrosomus regius*, *Sciaena umbra*, *Umbrina cirrosa*) appear to have low visceral losses (Table 2). The obvious reason for that is their low visceral fat deposition (Poli et al., 2003; García Mesa et al., 2014).

The condition index (CI) is indicative of the feeding condition of the fish and has been shown to increase in well-fed fish, like in the intensive farming-originated fish in comparison with extensively farmed and wild fish (Floss et al., 2002; Grigorakis, 2007; Piccolo et al., 2007; Martelli et al., 2013).

When concerning the CI, only intra-species comparisons are meaningful, since possible differences between species is most likely to be due to their different allometry. The latter for instance, becomes evident by the significantly lower condition indexes found for turbot and Senegalese sole than for the rest of the species (Table 2), obviously attributed to the different body geometry.

Metanalysis for the technical yields was possible only for VSI in the case of sharpsnout sea bream, where data are available in sufficient numbers for both wild and cultured counterparts (Table 2). A Hedges d index of 12.87 indicated statistically significant difference ($p < 0.01$) with

farmed fish exhibiting higher VSI. This can be explained as a higher visceral fat deposition for farmed fish, a fact that was also confirmed in one study that directly compared wild and farmed sharpsnout sea bream (Rueda et al., 2001).

Technical yields of Mediterranean species seem to be affected in various ways by environmental parameters, feeding and ploidy, and although the respective research is sporadic and sometimes contradictory, this is summarized in Table 3.

Technical yields, being of importance because they define the edible part of the products, show differentiations based on species, season and dietary history. Data on filleting yields are limited only to some of the species. For many species that filleting is expected to be of commercial interest, a monitoring of filleting yield is required in the future. Although sporadic data are available on the effects of dietary treatments and season in the fillet percentage, an important issue that needs to be addressed is the effect of fish size. This will be particularly useful in defining the best sizes for optimizing the filleting in each species. The herein results (Table 2) indicate that a technical loss of more than 50% during filleting, should be counterbalanced by the added value of the produced fillet. In regard with the filleting yield differentiation, none of the studied species seems to be more advantageous to others and therefore factors, other than technical ones, would determine economic efficiency.

The viscerosomatic index, defining the yield of eviscerated fish, seems to be lower for two fish categories, the flatfishes and the meagres. This can be advantageous for the commercialization of these fish in gutted forms. The condition index is meaningful for intra-species comparisons and can be potentially used as a tool for traceability issues between wild and farmed fish.

Fillet composition and fatty acids

The fillet composition is an important quality aspect since it largely defines the fish nutritional value and is also closely associated with its sensory attributes in the way they are perceived by the consumer (Grigorakis, 2007; Grigorakis, 2010). In general, with few exceptions, there is sufficient data on the fillet composition of the Mediterranean farmed fish species and in particular with their fillet fat.

The only important exception is that of wild meagre, where data is limited to those deriving from Sinanoglou et al. (2014). However, there is a serious probability that these authors have mistaken their so-assumed wild specimens. This can be postulated by the fact that they mentioned a fatty acid profile with high n6 fatty acids and in specifically of the 18:2n6 (linoleic acid) which is scarce in marine food chain and characterizes farmed fish due to its dietary terrestrial plant oil origin (Tocher, 2003; Linder et al., 2010). A potential explanation is that the authors have been probably provided with farmed escapees.

In general, the fillet protein of all fish species is quite similar, nearing a 20% of total fillet constituents. Fillet protein is generally believed to be stable in fully grown fish and not to be influenced by external parameters (Love, 1992; Grigorakis, 2010). However, there have been some cases where seasonal changes have been reported for wild fish populations (Gökçe et al., 2004; Saoud et al., 2008). Reduction of muscle protein in adult fish has been mentioned in cases of mobilization under prolonged fasting (Love, 1992).

The fillet fat ranges between the various fish species. Among them, there are two categories, the flatfishes (*Psetta maxima*, *Solea senegalensis*, *Solea Solea*) and the Sciaenidae family (*Argyrosomus regius*, *Sciaena umbra* and *Umbrina cirrosa*) that have low fillet fat when compared to the rest of the fish (Table 4).

The meta-analysis results for fillet fat content comparisons between wild and farmed counterparts include dentex, sharpsnout sea bream, red porgy (*Pagrus pagrus*), common sole and brown meagre (Table 5). In all cases, the farmed counterparts were found to have increased muscle fat comparing to the wild ones. The higher fat contents in farmed animals have been confirmed by direct comparisons in individual studies for dentex (Dincer et al., 2010), sharpsnout sea bream (Rueda et al., 2001; Dincer et al., 2010), white seabream (Cejas et al., 2004), blackspot seabream (Álvarez et al., 2009), red porgy (Rueda et al., 1997; Loukas et al., 2010), brown meagre (Cakli et al., 2006; Dincer et al., 2010) and greater amberjack (Haouas et al., 2010; Rodriguez-Barreto et al., 2012).

The fillet fat even within the same species is highly dependable to various internal and external parameters, and therefore effects of season, for both wild and farmed counterparts, and feeding characteristics and intensity for farmed fish have been demonstrated in Mediterranean species (Table 6)

The fatty acid profiles of all studied Mediterranean fish are rich in n-3 polyunsaturated fatty acids (Table 7). Results of metanalysis, available for dentex, sharpsnout sea bream, red porgy, brown meagre and common sole (Table 5) indicated differences in fatty acid profiles between

farmed and wild counterparts. The general pattern observed is the higher n-6 content of the former ones, with the only exception of the red porgy where higher n-6 levels have been found in wild fish. The fact that the one out two studies contributed to the metanalysis, found almost double n-6 contents in wild fish (Rueda et al., 1997) is responsible for this. The former authors although found higher 18:2n6 contents in farmed fish, in agreement with the general rule, they mentioned 9 times higher 20:4n6 in wild individuals.

In studies directly comparing wild and farmed individuals, significantly higher DHA contents have been observed in farmed counterparts for brown meagre and dentex (Dincer et al., 2010) and for sharpsnout sea bream in one case (Piccolo et al., 2007), while the opposite trend has been mentioned for sharpsnout sea bream in a second case (Dincer et al., 2010), for red porgy (Loukas et al., 2010) and greater amberjack (Haouas et al., 2010). EPA has been found to be higher in farmed fish for red porgy (Loukas et al., 2010), dentex, brown meagre (Dincer et al., 2010) and greater amberjack (Haouas et al., 2010) but similar (Piccolo et al., 2007) or higher (Dincer et al., 2010) in wild fish for sharpsnout sea bream.

When fatty acids are expressed as contents per 100g fish fillet, it becomes evident that Mediterranean fish species are highly nutritious and that their consumption can easily fulfill the n-3 PUFA daily needs (Table 8). It is worth mentioning that even species with low fat contents (flatfish and meagres) can cover a significant part of these needs. A comparison in total n-3, EPA and DHA contents between farmed and wild counterparts is always in favor of the former ones, obviously due to their higher fillet fat. This contradicts the general consumer impressions who perceive the farmed fish as of inferior nutritional quality (Claret et al., 2014).

The general rule of muscle fatty acids reflecting the dietary ones, applies for all Mediterranean farmed fish. Within these frames, inclusion of plant oils such as safflower oil (Altundag et al., 2014), soybean oil (Regost et al., 2003; Piedecausa et al., 2007), linseed oil (Regost et al., 2003; Piedecausa et al., 2007) or mixtures of vegetable oils (Valente et al., 2011) have shown to increase the n6 contents. Inclusion of animal fat (such as lard) to replace fish oil, leads into increase of muscle SFA and reduction of n3 PUFA and n3/n6 ratio (Nogales Merida et al., 2011b). For the herein studied species, fasting has been evaluated in common sole; muscle saturated fatty acids increased during starvation while PUFAs decreased, opposing the general rule that implies preservation of PUFAs and consumption of SFA with fasting (Fonseca et al., 2013).

Although, it is very difficult to make direct correlations of flesh fatty acids in wild fish with feeding habits, a very interesting recent observation indicated the depletion of PUFAs, EPA, DHA and ARA in particular, in wild white seabream due to consumption of the invasive algae *Caulerpa racemosa* (Felline et al., 2014)

Rearing system is, in some cases, influential to the muscle fatty acids, like one study that showed fatty acid differentiation between cage-reared and tank-reared meagre receiving the same diet (Martelli et al., 2013). It is not exactly clear what the causes of these differences. Perhaps the stocking density is one of the regulating factors in these cases (Piccolo et al., 2008), although this has not been always confirmed (Roncarati et al., 2006). Usually, the stocking density seems to have a more profound effect in the liver fatty acids, as a result to respective metabolic adaptations (Montero et al., 1999; Karakatsouli et al., 2007). The stocking density-derived

differences, however, can be potentially attributed to feed intake differences (Lund et al., 2013). The other potential regulating factor in different rearing systems is the water temperature, or the general seasonal differences, which is discussed in the following.

Season may have an impact on the muscle fatty acids of temperate water fish. An increase of the unsaturation level, primarily in the polar lipid fraction, has been associated to the water temperature drop, in order to maintain the membrane fluidity (Hazel and Prosser, 1974; Love, 1992; Delgado et al., 1994). Furthermore, a seasonal depletion in MUFA, and in particular in oleic acid (18:1n9) is correlated with mobilization during gonadal development and this is of importance in species that maturation occurs at commercial size (Sargent, 1995; Özyurt and Polat, 2006). In wild fish, seasonality in fatty acids, has been noticed for several Mediterranean populations (Gökçe et al., 2004; Cakli et al., 2006; Özyurt et al., 2005; Özoğul et al., 2011b).

Studies referring to farmed Mediterranean populations, in specific gilthead sea bream, common sea bass and meagre (Senso et al., 2007; Yildiz et al., 2006; Yildiz et al., 2008; García Mesa et al., 2014), indicated minor or absolute absence of season-related fatty acid changes. García Mesa et al. (2014) justified this lack of markedly seasonal effects, to the high winter temperatures occurring in the Mediterranean. Nevertheless, a common weakness of the majority of the existing studies, is that seasonal effect is not individually studied and that other parameters, such as growth (Hernández et al., 2003; Poli et al., 2003; García Mesa et al., 2014), fish size (Martelli et al., 2013b) or variable genetic origin (Cardinal et al., 2011) interfere. Therefore, the examination of seasonal impact would require a comparison in fish of similar weights, dietary

history and genetic uniformity, at different year intervals. Chatzifotis et al. (2004) having studied the fatty acid changes of dentex from the age of 8 to 36 months, i.e. for a period longer than two years, provided some data that allow conclusions in regard with the seasonal effect. Thus, irrespective to the fish weight (unfortunately not provided in the respective reference), a similar annual pattern can be observed with the fish always exhibiting a minimum of DHA and n3 fatty acids in February and a maximum in October, while the exactly opposite pattern applied for total MUFA (Fig. 1). On the other side, no specific pattern was observed for the rather stable levels of SFA, total n6 and EPA. These findings contradict the absence of season-related changes mentioned by the aforementioned studies. These also imply that EPA which is conserved, unlike DHA that fluctuates, plays an important biological role.

Based on the fillet composition, it becomes evident that species can be distinguished in low fat species and fat accumulating ones, with the former ones including the meagres and the flatfishes. In all species, the general rule of higher fillet fat contents in farmed specimens has been confirmed for all species (Table 4).

The fatty acid profiles showed differentiation between wild and farmed specimens for all species (where data are sufficient). These fatty acid differences can be justified based, firstly on the general rule of dietary fatty acids reflection in fish flesh, and secondly on the seasonal changes related to temperature adaptations and gonadal maturation. The daily needs of an adult in n-3 PUFA are covered in most cases even by the consumption of 100g fish, due to the high unsaturation level of the fatty acids in all studied species. Farmed fish retain a nutritional advantage over wild ones due to their higher n-3 contents.

Organoleptic quality of Mediterranean farmed fish

Sporadic data occur in aspects of the organoleptic quality of the Mediterranean farmed species.

Acceptability comparisons have been made between two different farmed sparidae species, the sharpsnout sea bream and gilthead sea bream (Hernández et al., 2001b), which were always in favour of the gilthead sea bream probably due to its familiarity for the consumers. However, the hedonic test participants were well disposed towards both species.

Difference tests and blind acceptability studies have been also conducted between meagre of different sizes (Gonçalves et al., 2011; Giogios et al., 2013; Ribeiro et al., 2013; Saavedra et al., 2015) mainly to resolve the market rumor of quality inferiority in fish weighing less than 1 Kg (Monfort, 2010; FAO, 2013; Ribeiro et al., 2013). All respective studies showed similarities in their findings, i.e. significant sensory difference between smaller (<1Kg) and larger fish (>1Kg), high acceptability for fish with weights ranging between 500g and 1Kg although preference remained higher for larger fish. The assumption that the major quality feature responsible for better acceptability of large fish is the size-dependent texture differentiation, was supported in some cases (Gonçalves et al., 2011; Ribeiro et al., 2013) but not in other (Giogios et al., 2013; Saavedra et al., 2015).

Organoleptic comparisons between wild and cultured fish have been taken place for gilthead sea bream and sea bass in some cases and these have been reviewed in the past (Grigorakis, 2007; Arechavala-Lopez et al., 2013), while sporadic data occur for other species, for the brown meagre in specifically (Cakli et al., 2006). In all cases wild fish are characterized by reduced fatiness of their fillet, perceived flavour differences, and a darker appearance of the muscle

(Cakli et al., 2006; Grigorakis, 2007; Arechavala-Lopez et al., 2013). Besides wild and farmed fish sensory differences, the rearing environment can also have an impact on fish sensory properties. In particular freshwater and seawater-reared red drum was found different in aspects of colour and texture; seawater-rearing resulted into lower hardness and color intensity (Klanian and Alonso, 2013).

A number of studies attempt to evaluate the impact of the diet on the sensory properties of produced fish. Fish oil and fishmeal substitution by plant raw materials (Izquierdo et al., 2005; Izquierdo et al., 2005; Cabral et al., 2013; Matos et al., 2012; Moreira et al., 2014) and the differentiation of dietary fat levels (Lopparelli et al., 2004) mostly resulted in minor or no sensory changes, mainly limiting in fillet color alterations (Izquierdo et al., 2005; Segato et al., 2005b; Matos et al., 2012). Otherwise, no other impact was observed in aspects of aroma or mouth sensation (texture or flavor of the fish). In some studies, however, dietary interventions have been found to affect sensory-perceived texture in case of fishmeal substitution (Hernández et al., 2007) or odour intensity in cases of dietary fat elevation (Segato et al., 2008). What can be assumed is that the species-characteristic fat accumulation and muscle microstructure, as well as the differential dietary requirements may differentiate the way that each species' sensory quality responds to the dietary treatments.

The existing data, so far provide evidence that sensory quality of fish can be altered subject to fish feeding history, somatic size and rearing environment. However, the sporadic results and the absence of analytic tools, i.e. descriptive analysis methods for describing Mediterranean fish species, do not allow safe conclusions on the degree and direction of these organoleptic changes.

Overall critical conclusion-recommendations

The Mediterranean farmed fish species offer a perspective upon which aquaculture diversification can be based. Their somatic yields and their fillet composition are important determinants of their overall quality and of the alternative ways of their commercialization.

The filleting yield for most of the farmed species is within a range of 40-45%. Herein data indicated that the viscerosomatic index differs among species, with flatfishes and Sciaenidae species exhibiting low visceral losses when compared to the rest of the Mediterranean farmed species. It is therefore suggested that the representatives of these two families have an advantage when commercialized in gutted forms, due to the low technical loss.

The condition index is related to the body geometry of each species, can be indicative of the feeding condition of the fish, and can be also a useful tool to discriminate wild from farmed counterparts of the same species.

In aspects of fillet composition, present review indicated that some flatfishes and meagres have very low muscle fat contents. This feature differentiates them from the rest of the farmed species. All farmed fish that were herein reviewed, have been found to contain high n-3 PUFA. Farmed fish differ in their fatty acid composition from their wild counterparts. The most characteristic differences are the significantly higher muscle fat contents and the higher n-6 (mainly 18:2n-6) contents in farmed fish. The aquaculture management, dietary treatment and season can impact

on farmed fish fillet composition and fatty acid quality, while season largely affects lipid quality in wild fish through food availability and maturation process.

We herein indicated that for some species that their aquaculture is long established such as the meagres, there is lack of comparisons with the respective wild individuals. Their farming and commercialization is largely based on the well established knowledge for gilthead sea bream and sea bass. This results into defective perception or speculations on their actual dietary needs and their produced quality capacities. The establishment of future comparisons with wild fish would elucidate answers in respect to quality individualities (e.g. due to different fatty acid metabolism in meagres (Monroig et al., 2013)) and the “ideal” in their quality.

Contrary to the fillet proximate composition and the fat qualities that received large attention by scientists, the sensory characteristics of the vast majority of the Mediterranean farmed species have not been described systematically. Most of the existing literature focuses on examining whether specific management or dietary treatments differentiate the sensory quality of the fish when compared to control groups. The development of sensory description tools for the herein studied species, such as descriptive analysis, should be within future research priorities in order to gain knowledge of their sensory characteristics and factors affecting them.

Based on the herein observations of the quality features in the farmed Mediterranean fish, it becomes evident that species diversification can also provide products diversification. First of all, different species can be commercialized in various forms. The fish with larger commercial

weights, such as the meagres or the greater amberjack can provide a wider variety of products, based on fillets or cuts.

The Mediterranean farmed fish can also provide flesh with different nutritional specifications. Within these frames, the muscle fat is a determinant of the commercialization potential of each species. Fish of low fillet fat, such as flatfish and meagres, independently of their commercial sizes, can be recommended in low fat diets. Those species that accumulate muscular fat, like the medium-fat species that include all the sparids and the common sea bass, can be proposed as good candidate foods for n-3 rich diets since their EPA and DHA contents largely exceed the minimum daily recommended intake. The confirmed difference in the fillet fat of wild and cultured specimens, for all species, can be used in advantage of the fattier farmed fish, considering the high unsaturation level in the fatty acids of all species.

Aknowledgements

This review was conducted within the frames of DIVERSIFY project. This project is funded from the European Union's Seventh Framework Programme for research, technological development and demonstration (KBBE-2013-07 single stage, GA 603121, DIVERSIFY).

References

- Alasalvar, C., Taylor, K.D.A., Zubcov, E., Shahidi, F., and Alexis, M. (2002). Differentiation of cultured and wild sea bass (*Dicentrarchus labrax*): total lipid content, fatty acid and trace mineral composition. *Food Chem.* **79**: 145–150.
- Altundag, M.S., Tiril, S.U., and Ozdemir, A. (2014). Effects of safflower oil supplementation in diet on growth performance and body fatty acid composition of turbot (*Psetta maxima*). *Aquac. Int.* **22**: 597-605.
- Álvarez, V., Medina, I., Prego, R., and Auburg, S.P. (2009). Lipid and mineral distribution in different zones of farmed and wild blackspot seabream (*Pagellus bogaraveo*). *Eur. J. Lipid Sci. Technol.* **111**: 957-966.
- Álvarez, V., Trigo, M., Lois, S., Fernández, D., Medina, I., and Aubourg, S.P. (2009b). Comparative lipid composition study in farmed and wild blackspot seabream (*Pagellus bogaraveo*). *Czech J. Food Sci. Technol.* **27 (SPEC. ISS.)**: S274-S276.
- Arechavala-Lopez, P., Fernandez-Jover, D., Black, K., Ladoukakis, E., Bayle-Sempere, J.T., Sanchez-Jerez, P., and Dempster, T. (2013). Differentiating the wild or farmed origin of Mediterranean fish: a review of tools for sea bream and sea bass. *Rev. Aquac.* **5**: 137-157.
- Barra, A., Garau, V.L., Dessi, S., Sarais, G., Cereti, E., Arlorio, M., Coisson, J.D., and Cabras, P. (2008). Chemical characterization and DNA tracking of Sardinian botargo by *Mugil cephalus* from different geographical origins. *J. Agric. Food Chem.* **56**: 10847-10852.

Barazi-Yeroulanos, L. (2010). Synthesis of Mediterranean marine finfish aquaculture – a marketing and promotion strategy. Studies and Reviews., pp. 198. **In:** 88, F.N. Ed. General Fisheries Commission for the Mediterranean. FAO No 88. FAO, Rome.

Bonaldo, A., Rizzi, L., Badiani, A., Testi, S., and Gatta, P.P. (2004). Growth response and body composition of sharpsnout sea bream (*Diplodus puntazzo*) fed a high energy diet with different protein levels. *Ital. J. Anim. Sci.* **3**: 235-242.

Cabral, E.M., Fernandes, T.J.R., Campos, S.T., Castro-Cunha, M., Oliveira, M.B.P.P., Cunha, L.M., and Valente, L.M.P. (2013). Replacement of fish meal by plant protein sources up to 75% induces good growth performance without affecting flesh quality in on-growing Senegalese sole. *Aquaculture* **380-383**: 130-138.

Cakli, Ş., Dincer, T., Cadun, A., Saka, Ş., and Firan, A. (2006). Nutrient content comparison of the new culture species brown meagre (*Sciaena umbra*). *Archiv Lebensmittelhyg.* **57**: 80-84.

Cakli, S., Kilinc, B., Dincer, T., and Tolasa, S. (2008). Shelf life of new culture species (*Diplodus puntazzo*) in refrigerator. *J. Muscle Foods* **19**: 315-332

Cardia, F. and Lovatelli, A. (2007). A review of cage aquaculture: Mediterranean sea. **In:**, Cage Aquaculture: regional reviews and global overview, pp 159-187. Halwart, M., Soto, D., and Arthur, J.R., eds, FAO Fisheries Technical Paper 498, Rome, Italy.

Cardinal, M., Cornet, J., Donnay-Moreno, C., Gouygou, J.P., Bergé, J.P., Rocha, E. Soares, S., Escórcio, C., Borges, P., and Valente, L.M.P. (2011). Seasonal variation of physical, chemical and sensory characteristics of sea bream (*Sparus aurata*) reared under intensive conditions in Southern Europe. *Food Control* **22**: 574-585.

Cejas, J.R., Almansa, E., Jérez, S., Bolaños, A., Samper, M., and Lorenzo, A. (2004). Lipid and fatty acid composition of muscle and liver from wild and captive mature female broodstocks of white seabream, *Diplodus sargus*. *Comp. Biochem. Physiol. Part B: Biochem. Molec. Biol.* **138**: 91-102.

Chatzifotis, S., Muje, P., Pavlidis, M., Ågren, J., Paalavuo, M., and Mölsä, H. (2004). Evolution of tissue composition and serum metabolites during gonadal development in the common dentex (*Dentex dentex*). *Aquaculture* **236**: 557-573.

Chen, L, and Opara, U.L. (2013). Texture measurement approaches in fresh and processed foods - A review. *Food Res. Int.* **51**: 823-835.

Claret, A., Guerrero, L., Ginés, R., Grau, A., Hernández, M.D., Aguirre, E., Peleteiro, J.B., Fernández-Pato, C., and Rodríguez-Rodríguez, C. (2014). Consumer beliefs regarding farmed versus wild fish. *Appetite* **79**: 25-31.

Craig, S.R., MacKenzie, D.S., Jones, G., and Gatlin III, D.M. (2000). Seasonal changes in the reproductive condition and body composition of free-ranging red drum, *Sciaenops ocellatus*. *Aquaculture*. **190**: 89-102.

de Almeida Ozório, R.O., Andrade, C., Timóteo, V.M.F.A., da Conceição, L.E.C., and Valente, L.M.P. (2009). Effects of feeding levels on growth response, body composition, and energy expenditure in blackspot seabream, *Pagellus bogaraveo*, juveniles. *J. World Aquac. Soc.* **40**: 95-103.

Delgado, A., Estevez, A., Hortelano, P., and Alexandre, M.J. (1994). Analyses of fatty acids from different lipids in liver and muscle of sea bass (*Dicentrarchus labrax* L.). influence of temperature and fasting. *Comp. Biochem. Physiol. - A: Physiol.* **108**: 673-680.

Diamant, A. (1998). Red drum *Sciaenops ocellatus* (Sciaenidae), a recent introduction to Mediterranean mariculture, is susceptible to *Myxidium leei* (Myxosporidia). *Aquaculture* **162**: 33-39.

Dincer, T., Cakli, S., and Cadun, A. (2010). Comparison of proximate and fatty acid composition of the flesh of wild and cultured fish species *Archiv Lebensmittelhyg.* **61**: 12-17.

El-Dakar, A.Y., Shalaby, S.M., and Saoud, I.P., (2007). Assessing the use of a dietary probiotic/prebiotic as an enhancer of spinefoot rabbitfish *Siganus rivulatus* survival and growth. *Aquac. Nutr.* **13**: 407-412.

El-Sebaiy, L.A., Metwalli, S.M., and Khalil, M.E. (1987). Phospholipid changes in muscles of plathead grey mullet (*Mugil cephalus*) during frozen storage. *Food Chem.* **26**: 85-96.

Ersoy, B., Çelik, M., Yanar, Y., Küçülgülmez, A., and Sangün, L. (2008). Comparison of the proximate compositions and fatty acid profiles of gilthead sea bream (*Sparus aurata*) and Sole (*Solea solea*). *Asian J. Chem.* **20**: 1251-1259.

FAO (2005-2015) Cultured Aquatic Species Information Programme. *Psetta maxima*. Cultured Aquatic Species Information Programme. (Rodríguez Villanueva, J. L. and Fernández Souto, B). In: *FAO Fisheries and Aquaculture Department* [online]. Rome. Updated 4 May 2005. [Cited 7 July 2015]. http://www.fao.org/fishery/culturedspecies/Psetta_maxima/en

FAO (2010). FAO yearbook. Fisheries and Aquaculture statistics 2009: Aquaculture production, pp 231.

FAO (2013). FAO Fisheries and Aquaculture Department: Global Aquaculture Production Statistics for the year 2011 [online]. <ftp://ftp.fao.org/FI/news/GlobalAquacultureProductionStatistics2011.pdf>

FAO (2014). The state of world fisheries and aquaculture: Opportunities and challenges. [online].
<http://www.fao.org/3/a-i3720e.pdf>

Felline, S., Mollo, E., Ferramosca, A., Zara, V., Regoli, F., Corbi, S., and Terlizzi, A. (2014).
 Can a marine pest reduce the nutritional value of Mediterranean fish flesh? *Marine Biol.* **161**:
 1275-1283.

Figueiredo-Silva, A.C., Corraze, G., Borges, P., and Valente, L.M.P. (2010). Dietary
 protein/lipid level and protein source effects on growth, tissue composition and lipid metabolism
 of blackspot seabream (*Pagellus bogaraveo*). *Aquac. Nutr.* **16**: 173-187.

FishBase (2011) World Wide Web electronic publication. Froese, R. and D. Pauly. (eds).
www.fishbase.org, (last assessed 18/10/2013)

Flos, R., Reig, L., Oca, J., and Ginovart, M. (2002). Influence of marketing and different land-
 based systems on gilthead sea bream (*Sparus aurata*) quality. *Aquac. Int.* **10**: 189-206.

Fonseca, V.F., Narciso, L., and Cabral, H.N. (2013). Lipid-based indicators of nutritional
 condition in juvenile sole *Solea solea*. *J. Appl. Ichthyol.* **29**: 154-162.

García, J.R., Kalinowski, C.T.H., Izquierdo, M.S.L., and Robaina, L.E.R. (2010). Marine and
 freshwater crab meals in diets for red porgy (*Pagrus pagrus*): effect on growth, fish composition
 and skin colour. *Aquac. Res.* **41**: 1759-1769.

García Mesa, S., Suárez, M.D., Rincón Cervera, M.A., Guil Guerrero, J.L., González, G.,
 Cárdenas, S., and García Gallego, M. (2014). Time course of muscle fatty acid composition of
 cultured meagre (*Argyrosomus regius*) during the first sixteen months of a cage culture. *Grasas y*
Aceites **65** in press, doi: <http://dx.doi.org/10.3989/gya.049813>.

Giogios, I., Grigorakis, K., and Kalogeropoulos, N. (2013). Organoleptic and chemical quality of farmed meagre (*Argyrosomus regius*) as affected by size. *Food Chem.* **141**: 3153-3159.

Gökçe, M.A., Taşbozan, O., Çelik, M., and Tabakoğlu, Ş.S. (2004). Seasonal variations in proximate and fatty acid compositions of female common sole (*Solea solea*). *Food Chem.* **88**: 419-423.

Gonçalves, A., Sá, M.d., Pousão-Ferreira, P., Ribeiro, L., Nunes, L., and Bandarra, N. (2011). Sensory quality and nutritional value of farmed meager (*Argyrosomus regius*). EAS2011 “Mediterranean Aquaculture 2020”. European Aquaculture Society, Rhodes, Greece pp. 345.

Grigorakis, K., Alexis, M.N., Taylor, K.D.A., and Hole, M. (2002). Comparison of wild and cultured gilthead sea bream; composition, appearance and seasonal alterations. *Int. J. Food Sci. Technol.* **37**: 477–484.

Grigorakis, K., and Alexis, M. (2005). Effects of fasting on the meat quality and fat deposition of commercial-size farmed gilt-head bream (*Sparus aurata* L.) fed different dietary regimes. *Aquac. Nutr.* **11: 341-344**

Grigorakis, K. (2007). Compositional and organoleptic quality of farmed and wild gilthead sea bream (*Sparus aurata*) and sea bass (*Dicentrarchus labrax*) and factors affecting it: A review. *Aquaculture* **272**: 55-75.

Grigorakis, K. (2010) Tailorizing quality: effects of aquaculture handling techniques and nutrition on fish quality. **In:** Seafood Quality, Safety and Health Effects. pp. 82-95. Alasalvar, C., Shahidi, F., Miyashita, K., and Wanasundara, U., Eds, WILEY-BLACKWELL Publishing U.K.

- Grigorakis, K., Fountoulaki, E., Vasilaki, A., Mittakos, I., and Nathanailides, C. (2011). Lipid quality and filleting yield of reared meagre (*Argyrosomus regius*). *Int. J. Food Sci. Technol.* **46**: 711-716.
- Grigorakis K., and Rigos, G. (2011). Aquaculture effects of environmental and public welfare - the case of Mediterranean mariculture. *Chemosphere* **855**: 899-919.
- Haouas, W.G., Zayene, N., Guerbej, H., Hammami, M., and Lotfi, A. (2010). Fatty acids distribution in different tissues of wild and reared *Seriola dumerili*. *Int. J. Food Sci. Technol.* **45**: 1478-1485.
- Hattour, A., and Kouched, W. (2014). Temporal distribution of size and weight of fattened bluefin tuna (*Thunnus thynnus* L.) from Tunisian farms (2005-2010). *Mediterr. Mar. Sci.* **15**: 115-125
- Hazel, J.R., and Prosser, C.L. (1974). Molecular mechanisms of temperature compensation in poikilotherms. *Physiol. Rev.* **54**: 620-677.
- Hedges, L.V., and Olkin, I. (1985). Statistical methods for meta-analysis. Academic Press, San Diego, CA, USA.
- Hernández, M.D., Egea, M.A., Rueda, F.M., Aguado, F., Martínez, F.J., and García, B. (2001). Effects of commercial diets with different P/E ratios on sharpsnout seabream (*Diplodus puntazzo*) growth and nutrient utilization. *Aquaculture* **195**: 321-329.
- Hernández, M.D., Martínez, F.J., and García, B. (2001b). Sensory evaluation of farmed sharpsnout seabream (*Diplodus puntazzo*). *Aquac. Int.* **9**: 519-529.

- Hernández, M.D., Egea, M.A., Rueda, F.M., Martínez, F.J., and García García, B. (2003). Seasonal condition and body composition changes in sharpsnout seabream (*Diplodus puntazzo*) raised in captivity. *Aquaculture* **220**: 569-580.
- Hernández, M.D., Martínez, F.J., Jover, M., and García García, B. (2007). Effects of partial replacement of fish meal by soybean meal in sharpsnout seabream (*Diplodus puntazzo*) diet. *Aquaculture* **263**: 159-167.
- Hernández, M.D., López, M.B., Álvarez, A., Ferrandini, E., García García, B., and Garrido, M.D. (2009). Sensory, physical, chemical and microbiological changes in aquacultured meagre (*Argyrosomus regius*) fillets during ice storage. *Food Chem.* **114**: 237-245.
- Hornung, H., Sukenik, A., and Gablielides, G.P. (1994). Distribution and composition of fatty acids in muscle lipids of inshore fish and deep water sharks from the Eastern *Mediterranean*. *Mar. Poll. Bull.* **28**: 448-450.
- Howell, B.R. (1997) A re-appraisal of the potential of the sole, *Solea solea* (L.), for commercial cultivation. *Aquaculture* **155**: 359-369.
- Imslund, A.K., Foss, A., Conceição, L.E.C., Dinis, M.T., Delbare, D., Schram, E., Kamstra, A., Rema, P., and White, P. (2003). A review of the culture potential of *Solea solea* and *S. senegalensis*. *Rev. Fish Biol. Fisher.* **13**: 379-408.
- Izquierdo, M.S., Montero, D., Robaina, L., Caballero, M.J., Rosenlund, G., and Gines, R. (2005). Alterations in fillet fatty acid profile and flesh quality in gilthead seabream (*Sparus aurata*) fed vegetable oils for long term period. Recovery of fatty acid profiles by fish oil feeding. *Aquaculture* **250**: 431-444.

Kalinowski, C.T., Socorro, J., and Robaina, L.E. (2015). Effect of dietary canthaxanthin on the growth and lipid composition of red porgy (*Pagrus pagrus*). *Aquac. Res.* **46**: 893-900

Kamdem, S.S., Vernocchi, P., Maffei, M., Belletti, N., Gardini, F., Guerzoni, M.E., and Lanciotti, R. (2008). Assessment of safety, nutritional, and spoilage characteristics of different lagoon grey mullets (*Liza ramada*, *Liza aurata*, and *Liza saliens*). *J. Food Protect.* **71**: 2572-2577.

Karakatsouli, N., Papoutsoglou, S.E., and Manoleosos, G. (2007). Combined effects of rearing density and tank colour on the growth and welfare of juvenile white sea bream *Diplodus sargus* L. in a recirculating water system. *Aquac. Res.* **38**: 1152-1160.

Ketata Khitouni, I., Abdelmouleh, A. Bouain, A., and Boudhrioua Mihoubi, N. (2010). Variations of the chemical composition of five coastal catch fish species of the Gulf of Gabes (Tunisia). *Cybium* **34**: 175-183.

Klanian, M.G., and Alonso, M.G. (2013). Sensory characteristics and nutritional value of red drum *Sciaenops ocellatus* reared in freshwater and seawater conditions. *Aquac. Res.* In press, doi:10.1111/are.12307

Koubaa A., Abdelmouleh, A., Bouain, and A., Mihoubi, N.B. (2011). Experimental and statistical investigations of the global chemical composition of six trawling fish of the Gulf of Gabès (Mediterranean sea). *J. Mar. Biol. Assoc. U.K.* **91**: 1689-1695.

Koubaa, A., Abdelmouleh, A., Bouain, A., and Boudhrioua Mihoubi, N. (2013). Effect of Cooking Methods on Nutritional Profile of Common Pandora (*Pagellus erythrinus*) from the Mediterranean Sea. *J. Food Proc. Preserv.* **38**: 1682-1689

- Krajnovic-Ozretic, M., Najdek, M., and Ozretic, B. (1994). Fatty acids in liver and muscle of farmed and wild sea bass (*Dicentrarchus labrax* L.). *Comp. Biochem. Physiol.* **109A**: 611–617.
- Kris-Etherton, P.M., Harris, W.S., and Appel, L.J. (2003). Fish consumption, fish oil, ω -3 fatty acids and cardiovascular disease. *Atherosclerosis, Thromb. Vasc. Biol.* **23**: 20-31.
- Li, T., Li, J., Hu, W., and Li, X. (2013). Quality enhancement in refrigerated red drum (*Sciaenops ocellatus*) fillets using chitosan coatings containing natural preservatives. *Food Chem.* **138**: 821-826.
- Linder, M., Belhaj, N., Sautot, P., and Tehrani, E.A. (2010). From krill to whale: An overview of marine fatty acids and lipid compositions. *OCL – Ol Corps Gras Li.* **17**: 194-204.
- Lloret, J., Galzin, R., Gil De Sola, L., Souplet, A., and Demestre, M. (2005). Habitat related differences in lipid reserves of some exploited fish species in the north-western Mediterranean continental shelf. *J. Fish Biol.* **67**: 51-65.
- Lopparelli, R.M., Segato, S., Corato, A., Fasolato, L., and Andrighetto, I. (2004). Sensory evaluation of sea bass (*Dicentrarchus labrax* L.) fed two diets differing in fat content. *Vet. Res. Commun.* **28** : 225–227.
- Loukas, V., Dimizas, C., Sinanoglou, V.J., and Miniadis-Meimaroglou, S. (2010). EPA, DHA, cholesterol and phospholipid content in *Pagrus pagrus* (cultured and wild), *Trachinus draco* and *Trigla lyra* from Mediterranean Sea. *Chem. Phys. Lipid.* **163**: 292–299
- Love, R.M. (1992). Biochemical dynamics and the quality of fresh and frozen fish. **In**: Fish Processing Technology. pp. 1–30. Hall, G.M., Ed., Balckie Academic, NY.
- Lund, I., Steenfeldt, S.J., Herrmann, B., and Pedersen, P.B. (2013). Feed intake as explanation for density related growth differences of common sole *Solea solea*. *Aquac. Res.* **44**: 367-377.

MacKenzie, B.R., Mosegaard, H., and Rosenberg, A.A. (2009). Impending collapse of bluefin tuna in the northeast Atlantic and Mediterranean. *Conserv. Lett.* **2**: 26-35.

Martelli, R., Parisi, G., Lupi, P., Bonelli, A., Zotte, A.D., and Franci, O. (2013). Effect of rearing system on body traits and fillet quality of meagre (*Argyrosomus regius*, Asso 1801) chilled for a short time. *Ital. J. Anim. Sci.* **12e30**: 186-195.

Martelli, R., Zotte, A.D., Bonelli, A., Lupi, P., Franci, O., and Parisi, G. (2013b). Macronutrient and fatty acid profiles of meagre (*Argyrosomus regius*) fillets as influenced by harvesting time and boiling. *Ital. J. Anim. Sci.* **12e88**: 538-545.

Matos, E., Gonçalves, A., Bandarra, N., Colen, R., Nunes, L.M., Valente, L.M.P., Dinis, M.T., and Dias, J. (2012). Plant proteins and vegetable oil do not have detrimental effects on post-mortem muscle instrumental texture, sensory properties and nutritional value of gilthead seabream. *Aquaculture* **358-359**: 205-212.

McGrath, C. (2012). Fishermen blame aquaculture for diminishing wild fry population. *Egypt Independent* **24.02.2012** <http://www.egyptindependent.com/news/fishermen-blame-aquaculture-diminishing-wild-fry-population>

Mnari, A., Bouhlel, I., Chraief, I., Hammami, M., Romdhane, M.S., El Cafsi, M., and Chaouch, A. (2007). Fatty acids in muscles and liver of Tunisian wild and farmed gilthead sea bream, *Sparus aurata*. *Food Chem.* **100**: 1393-1397.

Monfort, M.C. (2010). Present market situation and prospects of meagre (*Argyrosomus regius*), as an emerging species in Mediterranean aquaculture. Studies and Reviews. General Fisheries Commission for the Mediterranean. No. 89. Rome, FAO. 2010. 28p. (available in <http://www.fao.org/docrep/013/i1675e/i1675e.pdf>).

Montero, D., Izquierdo, M.S., Tort, L., Robaina, L., and Vergara, J.M. (1999). High stocking density produces crowding stress altering some physiological and biochemical parameters in gilthead seabream, *Sparus aurata*, juveniles. *Fish Physiol. Biochem.* **20**: 53-60.

Moreira, N. Soares, S., Valente, L.M.P., Castro-Cunha, M., Cunha, L.M., and Gueades de Pinho, P. (2014). Effect of two experimental diets (protein and lipid vegetable oil blends) on the volatile profile of Senegalese sole (*Solea senegalensis* Kaup, 1858) muscle. *Food Chem.* **153**: 327-333.

Monroig, Ó., Tocher, D.R., Hontoria, F., and Navarro, J.C. (2013). Functional characterisation of a Fads2 fatty acyl desaturase with $\Delta 6/\Delta 8$ activity and an Elovl5 with C16, C18 and C20 elongase activity in the anadromous teleost meagre (*Argyrosomus regius*). *Aquaculture* **412-413**: 14-22.

Nakada, M. (2008). Capture-based aquaculture of yellowtail. In: Capture-based aquaculture. Global overview pp. 199–215. Lovatelli, A., and Holthus, P.F., Eds. FAO Fisheries Technical Paper. No. 508. Rome, FAO. (In <http://www.fao.org/3/a-i0254e/i0254e10.pdf>).

Nathanailides, C., Kokokiris, L., Caripoglou, C., Kanlis, G., Logothetis, P., Mittakos, I, and Lenas, D. (2013). Muscle aerobic capacity and filleting yield of farmed fish species in the Mediterranean Sea. *J. Agric. Sci. Technol.* **B3**: 685-688.

Nevigado, T., Masci, M., Orban, E., Di Lena, G., Casini, I., and Caproni, R. (2012). Analysis of fatty acids in 12 Mediterranean fish species: Advantages and limitations of a new GC-FID/GC–MS based technique. *Lipids* **47**: 741-753.

Nicholson, T., Khademi, H., and Moghadasian, M.H. (2013). The role of marine n-3 fatty acids in improving cardiovascular health: a review. *Food Funct.* **4**: 357-365.

Nogales Mérida, S., Jover Cerdá, M., Martínez Llorens, S., and Tomás Vidal, A. (2011). A study of partial replacement of fish meal with sunflower meal on growth, amino acid retention, and body composition of sharpsnout seabream, *Diplodus puntazzo* (Actinopterygii: Perciformes: Sparidae). *Acta Ichthyol. Pisc.* **41**: 47-54.

Nogales-Mérida, S., Tomás-Vidal, A., Cerdá, M.J., and Martínez-Llorens, S. (2011b). Growth performance, histological alterations and fatty acid profile in muscle and liver of sharp snout sea bream (*Diplodus puntazzo*) with partial replacement of fish oil by pork fat. *Aquac. Int.* **19**: 917-927.

Nortvedt, R., and Tuene, S. (1998). Body composition and sensory assessment of three weight groups of Atlantic halibut (*Hippoglossus hippoglossus*) fed three pellet sizes and three dietary fat levels. *Aquaculture* **161**: 295-313.

Orban, E., Di Lena, G., Ricelli, A., Paoletti, F., Casini, I., Gambelli, L., and Carponi, R. (2000). Quality characteristics of sharpsnout sea bream (*Diplodus puntazzo*) from different intensive rearing systems. *Food Chem.* **70**: 27-32.

Özden, Ö., and Erkan, N. (2008). Comparison of biochemical composition of three aqua cultured fishes (*Dicentrarchus labrax*, *Sparus aurata*, *Dentex dentex*) *Int. J. Food Sci. Nutr.* **59**: 545-557.

Özoğul, Y., and Özoğul, F. (2007). Fatty acid profiles of commercially important fish species from the Mediterranean, Aegean and Black Seas. *Food Chem.* **100**: 1634-1638.

- Özoğul, Y., Özoğul, F., Ççek, E., Polat, A., and Kuley, E. (2008). Fat content and fatty acid compositions of 34 marine water fish species from the Mediterranean Sea., *Int. J. of Food Sci. Nutr.* **60**: 464 – 475.
- Özoğul, Y., Boğa, E.K., Tokur, B., and Özoğul, F. (2011). Changes in biochemical, sensory and microbiological quality indices of common sole (*Solea solea*) from the Mediterranean sea, during ice storage. *Turk. J. Fish. Aquat. Sci.* **11**: 243-251.
- Özoğul, Y., Polat, A., Uçak, I., and Özoğul, F. (2011b). Seasonal fat and fatty acids variations of seven marine fish species from the Mediterranean Sea. *Eur. J. Lipid Sci. Technol.* **113** : 1491-1498.
- Özyurt, G., Polat, A., and Özkütük, S. (2005). Seasonal changes in the fatty acids of gilthead sea bream (*Sparus aurata*) and white sea bream (*Diplodus sargus*) captured in Iskenderun Bay, Eastern Mediterranean coast of Turkey. *Eur. Food Res. Technol.* **220**: 120-124.
- Özyurt, G., and Polat, A. (2006). Amino acid and fatty acid composition of wild sea bass (*Dicentrarchus labrax*): A seasonal differentiation. *Eur. Food Res. Technol.* **222**: 316-320.
- Palmegiano, G.B., Costanzo, M.T., Daprà, F., Gai, F., Galletta, M.G., Maricchiolo, G., Micale, V., Peiretti P.G, and Genovese, L. (2007). Rice protein concentrate meal as potential dietary ingredient in practical diets for blackspot seabream (*Pagellus bogaraveo*). *J. Anim. Physiol. Anim. Nutr.* **91**: 235-239.

Perdicaris, C., and Paschos, I. (2011). Aquaculture and fisheries crisis within the global crisis. Interiencia 36: 76-80.

Pérez, M.J., Rodríguez, C., Cejas, J.R., Martín, M.V., Jerez, S., and Lorenzo, A. (2007). Lipid and fatty acid content in wild white seabream (*Diplodus sargus*) broodstock at different stages of the reproductive cycle. *Comp. Biochem. Physiol. - B Biochem Molec. Biol.* **146**: 187-196.

Pérez-Jiménez, A., Hidalgo, M.C., Morales, A.E., Arizcun, M., Abellán, E., and Gardenete, G. (2009). Use of different combinations of macronutrients in diets for dentex (*Dentex dentex*). Effects on intermediary metabolism. *Comp. Biochem Physiol. A* **152**: 314-321.

Periago, M.J., Ayala, M.D., Lopez-Albors, O., Abdel, I., Martinez, C., García-Alcazar, A., Ros, G., and Gil, F. (2005). Muscle cellularity and flesh quality of wild and farmed sea bass, *Dicentrarchus labrax* L. *Aquaculture* **249**: 175–188.

Piccolo, G., De Riu, N., Tulli, F., Cappuccinelli, R., Marono, S., and Moniello, G. (2007). Somatic indexes, chemical-nutritive characteristics and metal content in caught and reared sharpsnout seabream (*Diplodus puntazzo*). *Ital. J. Anim. Sci.* **6**: 351-360.

Piccolo, G., Marono, S., Bovera, F., Tudisco, R., Caricato, G., and Nizza, A. (2008). Effect of stocking density and protein/fat ratio of the diet on the growth of Dover sole (*Solea solea*). *Aquac. Res.* **39**: 1697-1704.

Piccolo, G., Centoducati, G., Bovera, F., Marrone, R., and Nizza, A. (2013). Effects of mannan oligosaccharide and inulin on sharpsnout seabream (*Diplodus puntazzo*) in the context of partial fish meal substitution by soybean meal. *Ital. J. Anim. Sci.* **12**: 133-138.

Piedecausa, M.A., Mazón, M.J., García García, B., and Hernández, M.D. (2007). Effects of total replacement of fish oil by vegetable oils in the diets of sharpsnout seabream (*Diplodus puntazzo*). *Aquaculture* **263** : 211-219.

Poli, B.M., Parisi G., Zampacavallo, G., Iurzan, F., Meccati, M., Lupi, P., and Bonelli, A. (2003). Preliminary results on quality and quality changes in reared meagre (*Argyrosomus regius*): body and fillet traits and freshness changes in refrigerated commercial-size fish. *Aquac. Int.* **11**: 301-311.

Prato, E., and Biandolino, F. (2012). Total lipid content and fatty acid composition of commercially important fish species from the Mediterranean, Mar Grande Sea. *Food Chem.* **131**: 1233-1239.

Recks, M.A., and Seaborn, G.T. (2008). Variation in fatty acid composition among nine forage species from a southeastern US estuarine and nearshore coastal ecosystem. *Fish Physiol. Biochem.* **34**: 275-287.

Regost, C., Arzel, J., Robin, J., Roselund, G., and Kaushik, S.J. (2003). Total replacement of fish oil by soybean or linseed oil with a return to fish oil in turbot (*Psetta maxima*) 1. Growth performance, flesh fatty acid profile, and lipid metabolism. *Aquaculture* **217**: 465-482.

Ribeiro, L., Soares, L., Quental- Ferreira, H., Gonçalves, A., and Pousão-Ferreira, P. (2013). Portuguese research studies meagre production in earthen ponds. *Glob. Aquac. Advoc.* **16**: 38-40.

Rodríguez-Barreto, D., Jerez, S., Cejas, J.R., Martin, M.V., Acosta, N.G., Bolaños, A., and Lorenzo, A. (2012). *Comparative study of lipid and fatty acid composition in different tissues*

of wild and cultured female broodstock of greater amberjack (Seriola dumerili). Aquaculture **360**: 1-9.

Roncarati, A., Melotti, P., Dees, A., Mordenti, O., and Angellotti, L. (2006). Welfare status of cultured seabass (*Dicentrarchus labrax* L.) and seabream (*Sparus aurata* L.) assessed by blood parameters and tissue characteristics. *J. Appl. Ichthyol.* **22**: 225-234.

Rondán, M., Hernández, Ma.D., Egea, Ma.A., García, B., Jover, M., Rueda, F.M., and Martínez, F.J. (2004). Effects of fishmeal replacement with soybean meal as protein source, and protein replacement with carbohydrates as an alternative energy source on sharpsnout sea bream, *Diplodus puntazzo*, fatty acid profile. *Aquac. Res.* **35**: 1220-1227.

Rondán, M., Hernández, Ma.D., Egea, Ma.A., García, B., Rueda, F.M., and Martínez, F.J. (2004b). Effect of feeding rate on fatty acid composition of sharpsnout seabream (*Diplodus puntazzo*). *Aquac. Nutr.* **10**: 301-307

Rueda, F.M., López, J.A., Martínez, F.J., Zamora, S., Divanach, P., and Kentouri, M. (1997). Fatty acids in muscle of wild and farmed red porgy, *Pagrus pagrus*. *Aquac. Nutr.* **3**: 161-165

Rueda, F.M., Hernández, M.D., Egea, M.A., Aguado, F., García, B., and Martínez, F.J. (2001). Differences in tissue fatty acid composition between reared and wild sharpsnout sea bream, *Diplodus puntazzo* (Cetti, 1777). *Brit. J. Nutr.* **86**: 617-622.

Rueda, F.M., Martínez, F.J., Zamora, S., and Divanach, P. (2008). Effect of fasting and refeeding on growth and body composition of red porgy, *Pagrus pagrus* L. *Aquac. Res.* **29**: 447-452.

Saavedra, M., Pereira, T.G., Grade, A., Barbeiro, M., Pousão-Ferreira, P., Quental-Ferreira, H., Nunes, M.L., Bandarra, N., and Gonçalves, A. (2015). Farmed meagre, *Argyrosomus regius* of

three different sizes: what are the differences in flesh quality and muscle cellularity? *Int. J. Food Sci. Technol.* In press, DOI: 10.1111/ijfs.12769

Saleh, M. (2008). Capture-based aquaculture of mullets in Egypt. **In:** Capture-based aquaculture. Global overview. pp. 109–126. Lovatelli, A., and Holthus, P.F., Eds., FAO Fisheries Technical Paper No. 508, FAO, Rome.

Saoud, I.P., Batal, M., Ghanawi, J., and Lebbos N. (2008). Seasonal evaluation of nutritional benefits of two fish species in the eastern Mediterranean Sea. *Int. J. Food Sci. Technol.* **43**: 538-542.

Sargent, J.R. (1995). Origins and functions of lipids in fish eggs: nutritional implications. **In:** Broodstock management and egg and larval quality. pp. 353–372. Bromage, N.R., Roberts, R.J., Eds., Blackwell Sci., Oxford.

Segato, S., Loparelli R.M., Borgoni, N., Zanella L., Corato, A., and Andrighetto, I. (2005). Effect of dietary crude fat to NFE ratio on growth, feed efficiency and quality traits in juvenile shi drum (*Umbrina cirrosa*). *Cah. Opt. Méditer.* **63**: 27-34.

Segato, S., Corato, A., Fasolato, L., and Andrighetto, I. (2005b). Effect of the partial replacement of fish meal and oil by vegetable products on performance and quality traits of juvenile shi drum (*Umbrina cirrosa* L.) *Ital. J. Anim. Sci.* **4**: 159-166.

Segato, S., Fasolato, L., Bertotto, D., Libertini, A., Balzan, S., Corato, A., and Novelli, E. (2007). Effect of triploidy on quality traits of shi drum (*Umbrina cirrosa* L.) until the second rearing year. *Aquac. Res.* **38**: 59-65.

Segato, S., Fasolato, L., Balzan, S., Elia, C.A., Novelli, E., and Andrighetto, I. (2008). Effect of dietary EE/NFE ratio on sensorial traits of shi drum. *Acta Agric. Sloven.* **91 (SUPPL. 2)**: 123-127.

Şengör, G.F., Özden, Ö., Erkan, N., Tüter, M. and Aksoy, H.A. (2003). Fatty acid compositions of flathead grey mullet (*Mugil cephalus* L., 1758) fillet, raw and beewaxed caviar oils. *Turk. J. Fish. Aquat. Sci.* **3**: 93-96.

Senso, L., Suárez, M.D., Ruiz-Cara, T., and García-Gallego, M. (2007). On the possible effects of harvesting season and chilled storage on the fatty acid profile of the fillet of farmed gilthead sea bream. *Food Chem.* **101**: 298–307.

Sevgili, H., Kurtoğlu, A., Oikawa, M., Fedekar, D., Emre, N., and Takeno, N. (2014). Evaluation of nutritional values of selected commercial fish meal sources in turbot (*Psetta maxima*) diets. *Aquac. Res.* doi:10.1111/are.12389

Sinanoglou, V.J., Proestos, C., Lantzouraki D.Z., Calokerinos, A.C., and Miniadis-Meimaroglou, S. (2014). *Lipid evaluation of farmed and wild meager (Argyrosomus regius). Eur. J. Lipid Sci. Technol. 116: 134-143.*

Soriguer, F., Serna, S., Valverde, E., Hernando, J., Martín-Reyes, A., Soriguer, M., Pareja, A., Tinahones, F., and Esteva, I. (1997). *Lipid, protein, and calorie content of different Atlantic and Mediterranean fish, shellfish, and molluscs commonly eaten in the south of Spain. Eur. J. Epidemiol. 13: 451-463.*

Suárez, M.D., Martínez, T.F., Abellán, E., Arizcun, M., Pérez-Jiménez, A., Hidalgo, M.C., and Cardenete, G. (2009). The effects of the diet on flesh quality of farmed dentex (*Dentex dentex*). *Aquaculture 288: 106-113.*

Suárez, M.D., Cervera, M.Á.R., Abellán, E., Morales, A.E., Gallego, M.G., and Cardenete, G. (2010). Influence of starvation on flesh quality of farmed dentex, *Dentex dentex*. *J. World Aquac. Soc. 41: 490-505.*

Tejada, M., De Las Heras, C., and Kent, M. (2007). Changes in the quality indices during ice storage of farmed Senegalese sole (*Solea senegalensis*). *Eur. Food Res. Technol. 225: 225-232.*

Testi, S., Bonaldo, A., Gatta, P.P., and Badiani, A. (2006). Nutritional traits of dorsal and ventral fillets from three farmed fish species. *Food Chem. 98: 104–111.*

Thakur, D.P., Morioka, K., Itoh, N., Wada, M., and Itoh, Y. (2009) Muscle biochemical constituents of cultured amberjack *Seriola dumerili* and their influence on raw meat texture. *Fish. Sci. 75: 1489-1498.*

Tocher, D.R. (2003). Metabolism and functions of lipids and fatty acids in teleost fish. *Rev. Fish. Sci.* **11**: 107-184.

Valente, L.M.P., Olmedo, M., Borges, P., Soares, S., Gomes, E.F.S., Álvarez-Blázquez, B., Pazos, G., and Linares, F. (2010). Effects of carbohydrate sources on growth, body composition and tissue lipid deposition of blackspot seabream, *Pagellus bogaraveo* (Brunnich). *J. Anim. Physiol. Anim. Nutr.* **94**: 212-219.

Valente, L.M.P., Linares, F., Villanueva, J.L.R., Silva, J.M.G., Espe, M., Escórcio, C., Pires, M.A., Saavedra, M.J., Borges, P., Medales, F., Álvarez-Blázquez, B., and Peleteiro, J.B. (2011). Dietary protein source or energy levels have no major impact on growth performance, nutrient utilisation or flesh fatty acids composition of market-sized Senegalese sole. *Aquaculture* **318**: 128-137.

Vanhonacker, F., Pieniak, Z., and Verbeke, W. (2013). European consumer image of farmed fish, wild fish, seabass and seabream. *Aquac. Int.* **21**: 1017-1033.

Varljen, J., Šulić, S., Brmalj, J., Batičić L., Obersnel, V., and Kapović, M. (2003). Fatty acid composition of *D. vulgaris* and *C. conger*. *Food Technol. Biotech.* **41**: 149–156.

Yildiz, M., Şener, E., and Tumur, M. (2006). The effects of seasons and different feeds on fatty acid composition in fillets of cultured gilthead sea bream (*Sparus aurata* L.) and European Sea Bass (*Dicentrarchus labrax* L.) in Turkey. *Turk. J. Vet. Anim. Sci.* **30**: 133–141.

Yildiz, M., Şener, E., and Timur, M. (2008). Effects of differences in diet and seasonal changes on the fatty acid composition in fillets from farmed and wild sea bream (*Sparus aurata* L.) and sea bass (*Dicentrarchus labrax* L.). *Int. J. Food Sci. Technol.* **43**: 853–858.

Table 1: Mediterranean marine finfish aquaculture production in tonnes. The groupers, *Epinephelus spp.* and wreckfish *Polyprion americanus* are not included since no commercial production has been recorded in the Mediterranean. Data refer to 2009 official production numbers (except † that refer to 2003 and ‡ to 2008 last maximum recorded quantities). Source: FAO, 2010.

Common name	Scientific name	Main producers in the Mediterranean	Total Medit. production	Total world production
Grey mullet, striped mullet	<i>Mugil cephalus</i>	Egypt	213,194	221,978
Other mullets	Mugilidae	Italy, Spain	390	9,078
Sea bass	<i>Dicentrarchus labrax</i>	Turkey, Greece, Spain	111,764	112,183
Gilthead sea bream	<i>Sparus aurata</i>	Greece, Turkey, Spain	133,124	136,070
Common sole	<i>Solea solea</i>	Italy, Spain	16	30
Senegalese sole	<i>Solea senegalensis</i>	Spain	63	63
Turbot	<i>Psetta maxima</i>	Spain	7,188	69,006
Brown meagre	<i>Sciaena umbra</i>	Spain	30†	30†
Shi drum	<i>Umbrina cirrosa</i>	Italy	45‡	45‡

Meagre	<i>Argyrosomus regius</i>	Egypt, Spain, France	4,068	4,112
Red drum	<i>Sciaenops ocellatus</i>	Israel	459	51,476
Blackspot seabream	<i>Pagellus bogaraveo</i>	Spain	183	183
Common pandora	<i>Pagellus erythrinus</i>	Greece	42	42
White sea bream	<i>Diplodus sargus</i>	Greece, Italy	104	105
Two-banded sea bream	<i>Diplodus vulgaris</i>	Italy	18‡	18‡
Sharpsnout seabream	<i>Diplodus puntazzo</i>	Italy	51	51
Common dentex	<i>Dentex dentex</i>	Boznia, Spain	10	10
Red Porgy	<i>Pagrus pagrus</i>	Cyprus, Greece	23†	23†
Other porgies	<i>Pagrus</i> spp.	Cyprus	10	10
Other sparids	Sparidae	Italy	65	42,827
Marbled spinefoot (or rabbitfish)	<i>Siganus rivulatus</i>	Cyprus	2	2
Atlantic bluefin tuna	<i>Thunnus, thynnus</i>	Croatia, Malta	1,994	1,999
Greater amberjack	<i>Seriola dumerili</i>	Spain	1	4

Table 2: Technological and somatometric characteristics of farmed Mediterranean fish species.

Values are weighed means. In parenthesis the total number of studies taken into account and the total number of fish that contributed to the weighed mean.

species		FY	VSI	HSI	CI	References
<i>Argyrosomus regius</i>	Farmed	43.3 (6/266)	4.96 (4/204)	1.50 (3/194)	1.05 (2/154)	Poli et al., 2003; Grigorakis et al., 2011; Giogios et al., 2013; Martelli et al., 2013; Nathanailides et al., 2013; Sinanoglou et al, 2013
<i>Dentex dentex</i>	Farmed	44.7 (2/306)	5.16 (2/306)	1.85 (3/342)	1.51 (3/342)	Pérez-Jimenez et al., 2009; Suárez et al., 2009; Suárez et al., 2010
<i>Diplodus annularis</i>	wild	-	7.32 (1/16)	1.28 (1/16)	-	Ketata Khituni et al., 2010
<i>Diplodus puntazzo</i>	Farmed	51.3 (1/54)	3.93 (6/215)	1.45 (6/232)	2.03 (6/237)	Hernández et al., 2001; Nogales-Merida et al., 2001b; Rueda et al., 2001; Hernández et al., 2003; Piccolo et al., 2007; Piedecausa et al., 2007; Piccolo et al., 2013
	wild	-	3.15 (2/65)	1.95 (1/60)	1.93 (1/60)	Rueda et al., 2001; Piccolo et al., 2007
<i>Diplodus</i>	wild	29.3	-	-	-	Saoud et al., 2008

<i>sargus</i>		(1/120)				
<i>Liza aurata</i>	wild	-	9.19 (1/13)	2.26 (1/13)	-	Ketata Khituni et al., 2010
<i>Pagellus bogaraveo</i>	farmed	-	8.41 (3/108)	1.24 (4/136)	2.70 (3/132)	Palmegiano et al., 2007; de Almeida Ozório et al., 2009; Figueiredo Silva et al., 2010; Valente et al., 2010.
<i>Pagrus pagrus</i>	farmed	-	6.81 (3/95)	1.64 (3/95)	3.13 (1/45)	Rueda et al., 1998; García et al., 2010; Kalinowski et al., 2015
<i>Psetta maxima</i>	farmed	-	3.66 (2/105)	1.57 (2/105)	1.42 (2/81)	Regost et al., 2003; Altundag et al., 2014; Sevgili et al., 2014
<i>Siganus rivulatus</i>	farmed	-	14 (1/25)	3.46 (1/25)	2.46 (1/25)	El-Dakar et al., 2007
	wild	36.7 (1/120)	-	-	-	Saoud et al., 2008
<i>Solea senegalensis</i>	farmed	-	1.49 (2/228)	1.17 (2/228)	1.51 (1/180)	Valente et al., 2011 ; Cabral et al., 2013
<i>Umbrina cirrosa</i>	farmed	40.5 (2/132)	4.19 (3/156)	2.45 (2/48)	2.12 (3/156)	Segato et al., 2005; Segato et al., 2005b; Segato et al., 2007

Table 3: Effects of influencing parameters in the technical yields and condition index of Mediterranean farmed species, or wild counterparts.

Species	Influencing Parameter	Effect	References
Filleting yield			
<i>Diplodus puntazzo</i>	inclusion of vegetable oil in diet	increase	Piedecausa et al., 2007
<i>Dentex dentex</i>	various dietary protein and fat levels	none	Suárez et al., 2009
	5 weeks starvation	increase	Suárez et al., 2010
<i>Siganus rivulatus</i> (wild)	Season	lowest yields in March & October	
<i>A. regius</i>	Fish growth / season	none	Saoud et al., 2008
VSI			
<i>Sciaena umbra</i>	dietary fat increase	increase	Segato et al., 2005
	inclusion of vegetable oil in diet	increase	Segato et al., 2005b
<i>Solea senegalensis</i>	Fish meal substitution by plant meals	increase	Valente et al., 2011; Cabral et al., 2013
<i>Dentex dentex</i> , <i>Diplodus puntazzo</i> , <i>Pagellus</i>	Fish meal and fish oil substitution by plant raw materials	none	Palmegiano et al., 2007; Suárez et al., 2009; Nogales-Merida et al.,

<i>bogaraveo</i>			2011, Piccolo et al., 2013
<i>Sciaena umbra</i>	triploidy	increase	Segato et al., 2007
<i>A. regius</i>	season	variable seasonality	Poli et al., 2003; Martelli et al., 2013
CI			
<i>Pagellus bogaraveo</i>	feeding intensity increase	increase	deAlmeida Ozório et al., 2009
<i>Dentex dentex</i>	5 weeks starvation	decrease	Suárez et al., 2010
<i>Argyrosomus regius</i>	Fish growth / season	none	Poli et al., 2003; Martelli et al., 2013
<i>Dentex dentex</i> , <i>Diplodus puntazzo</i> , <i>Pagrus pagrus</i> , <i>Solea senegalensis</i> , <i>Umbrina cirrosa</i> ,	diet composition	none	Hernández et al., 2001; Segato et al., 2005; Palmegiano et al., 2007; Piedecausa et al., 2007; Pérez-Jimenez et al., 2009; Suárez et al., 2009; García et al., 2010; Valente et al., 2011; Nogales-Merida et al., 2013; Piccolo et al., 2013

<i>Sciaena umbra</i>	triploidy	decrease	Segato et al., 2007
<i>Diplodus puntazzo</i>	season	Increase when closing reproductive season (September)	Hernández et al., 2003

Table 4: Fillet proximate composition of farmed Mediterranean fish species. Values are weighed means. In parenthesis the total number of studies taken into account and the total number of fish that contributed to the weighed mean.

species		Moisture %	Protein %	Ash %	Fat %	References
<i>Argyrosomus regius</i>	Farmed	72.8 (6/212)	20.5 (3/120)	1.34 (4/150)	2.12 (7/218)	Poli et al., 2003; Hernández et al., 2009; Grigorakis et al., 2011; Nevigado et al., 2012; Giogios et al., 2013; Martelli et al., 2013; Sinanoglou et al., 2013
<i>Dentex dentex</i>	Farmed	73.1 (4/329)	20.2 (4/329)	2.94 (2/36)	3.78 (4/329)	Ozden and Erkan, 2008; Suárez et al., 2009; Suárez et al., 2010; Dincer et al., 2010
	wild	75.2 (1/5)	22.1 (2/8)	-	1.91 (2/8)	Soriguer et al., 1997; Dincer et al., 2010
<i>Diplodus annularis</i>	wild	-	-	-	0.92 (1/3)	Özoğul et al., 2008
<i>Diplodus puntazzo</i>	Farmed	70.5 (3/16)	19.2 (3/16)	1.23 (2/11)	8.39 (4/20)	Orban et al., 2000; Rueda et al., 2001; Cakli et al., 2008; Dincer et al., 2010

	wild	76.7 (1/5)	19.2 (1/5)	-	2.80 (3/13)	Rueda et al., 2001; Özoğul et al., 2008; Dincer et al., 2010
<i>Diplodus sargus</i>	farmed	76.4 (1/9)	-	-	5.82 (1/9)	Cejas et al., 2004
	wild	77.4 (2/129)	19.3 (2/129)	1.76 (2/129)	1.51 (5/141)	Hornung et al., 1994; Soriguer et al., 1997; Özyurt et al., 2005; Saoud et al., 2008
<i>Diplodus vulgaris</i>	wild	76.7 (1/10)	-	-	2.27 (3/23)	Varljen et al., 2003; Özoğul et al., 2008; Prato and Biandolino 2012
<i>Epinephelus auneus</i>	wild	77.6 (1/18)	19.7 (1/18)	1.57 (1/18)	0.77 (1/18)	Özoğul et al., 2011b
<i>Liza aurata</i>	wild	74.5 (1/9)	20.1 (1/9)	1.42 (1/9)	4.51 (3/22)	Kamden et al., 2008; Özoğul et al., 2008; Prato and Biandolino 2012
<i>Liza ramada</i>	wild	74.4 (1/9)	19.8 (1/9)	1.58 (1/9)	3.25 (3/15)	Kamden et al., 2008; Özoğul et al., 2008 ; Nevigado et al., 2012
<i>Liza saliens</i>	wild	76.0 (1/9)	18.5 (1/9)	1.28 (1/9)	2.87 (2/12)	Kamden et al., 2008; Özoğul et al., 2008

<i>Mugil cephalus</i>	farmed	76 (1/6)	-	-	10.0 (1/6)	El-Sebaiy et al., 1987
	wild	-	-	-	2.10 (2/7)	Özoğul and Özoğul 2007; Özoğul et al., 2008
<i>Pagellus bogaraveo</i>	farmed	72.2 (1/36)	19.8 (1/36)	1.66 (1/36)	7.26 (4/120)	Palmegiano et al., 2007; de Almeida Ozório et al., 2009; Figueiredo Silva et al., 2010; Valente et al., 2010.
	wild	-	21.1 (1/3)	-	4.32 (1/3)	Soriguer et al., 1997
<i>Pagellus erythrinus</i>	wild	75.9 (2/36)	18.1 (2/36)	1.54 (2/36)	3.86 (3/40)	Özoğul and Özoğul 2007; Koubaa et al., 2011; Koubaa et al., 2014
<i>Pagrus pagrus</i>	farmed	74.7 (1/24)	21.6 (1/24)	1.70 (1/24)	4.19 (3/47)	Rueda et al., 1997; Loukas et al., 2010; Kalinowski et al., 2015
	wild	-	-	-	0.85 (2/24)	Rueda et al., 1997; Loukas et al., 2010
<i>Psetta maxima</i>	farmed	77.4 (2/24)	20.1 (2/24)	-	1.57 (3/27)	Regost et al., 2003; Nevigado et al., 2012; Altundag et al., 2013

	wild	-	20.4 (1/3)	-	0.97 (1/3)	Soriguer et al., 1997
<i>Sciaena umbra</i>	farmed	74.1 (2/45)	20.2 (2/45)	-	3.18	Cakli et al., 2006; Dincer et al., 2010
	wild	74.7 (1/40)		-	1.40 (1/40)	Cakli et al., 2006
<i>Seriola dumerili</i>	farmed	71.7 (2/39)	23.2 (1/30)	-	6.65 (2/39)	Thakur et al., 2009; Rodriguez-Bareto et al., 2012
	wild	76.7 (1/9)	-	-	3.64 (1/9)	Rodriguez-Bareto et al., 2012
<i>Siganus rivulatus</i>	wild	76.7 (1/120)	19.7 (1/120)	1.39 (1/120)	1.95 (1/120)	Saoud et al., 2008
<i>Solea senegalensis</i>	farmed	-	19.5 (1/36)	-	1.47 (2/84)	Valente et al., 2011; Cabral et al., 2013
	wild	76.5 (1/5)	20.6 (1/5)	1.24 (1/5)	1.43(1/5)	Tejada et al., 2007
<i>Solea Solea</i>	farmed	76.9 (1/48)	19.6 (1/48)	1.15 (1/48)	2.65 (2/51)	Piccolo et al., 2008; Nevigado et al., 2012;
	wild	78.5 (4/70)	19.2 (4/70)	1.27 (4/70)	0.54 (5/74)	Gökçe et al., 2004; Özoğul and Özoğul, 2007; Ersoy et

						al., 2008; Özoğul et al., 2011; Özoğul et al., 2011b
<i>Umbrina cirrosa</i>	farmed	75.9 (4/136)	20.8 (4/136)	1.26 (3/84)	1.65 (4/136)	Segato et al., 2005; Segato et al., 2005b; Segato et al., 2007; Segato et al., 2008
	wild	-	-	-	0.92 (1/3)	Özoğul et al., 2008

Table 5: Metanalysis results: Hedges d index for detection of differences between wild and farmed counterparts. The * and ** denote statistically significant differences ($p < 0.05$ and $p < 0.01$, respectively).

Species	Muscle Fat	SFA	MUFA	n-3 PUFA	n-6 PUFA	EPA	DHA
<i>Dentex dentex</i>	17.8**	2.90**	5.15**	7.15**	10.5**	1.59*	7.40**
<i>Diplodus puntazzo</i>	5.77**	18.7**	0.27	7.49**	7.55**	2.90**	3.05**
<i>Pagrus pagrus</i>	26.4**	9.79	8.88**	4.00**	4.28**	2.42**	10.2**
<i>Sciaena umbra</i>	12.1**	26.6**	8.81**	13.3**	111.1**	-	-
<i>Solea Solea</i>	17.4**	8.31**	21.8**	0.89**	7.55**	6.26**	5.11**

Table 6: Factors influencing the fillet fat of Mediterranean farmed species, or wild counterparts.

Species	Influencing Parameter	Effect	References
Mugilidae, Sparidae	Fish size	Increase with size	Grigorakis, 2007; Ketata Kitouni et al., 2010
Wild populations of: <i>Dicentrarchus</i> <i>labrax</i> , <i>Diplodus</i> <i>sargus</i> , <i>Sciaena</i> <i>umbra</i> , <i>Siganus</i> <i>rivulatus</i> , <i>Sparus</i> <i>aurata</i> , <i>Solea solea</i>	Season	Increasing in the summer months, due to intensive feeding and elevated food availability. Maximum at the end of the warm period (late summer to mid autumn). Depletion in cold months.	Gökçe et al., 2004; Özyurt et al., 2005; Cakli et al., 2006; Özyurt and Polat, 2006; Senso et al., 2007; Ersoy et al., 2008; Saoud et al., 2008; Özoğul et al., 2011b
Farmed <i>Argyrosomus</i> <i>regius</i> , <i>Scieana</i> <i>umbra</i> , <i>Seriola</i> <i>dumerili</i>	Season	Depletion in cold months.	Poli et al., 2003; Cakli et al., 2006; Thakur et al., 2009
<i>Solea senegalensis</i> ,	Dietary fat	none	Segato et al., 2005;

<i>Solea solea</i> , <i>Umbrina cirrosa</i>			Piccolo et al., 2008; Valente et al., 2011
<i>Hippoglossus hippoglossus</i>	Dietary fat	Increase with fat level	Nortvendt and Tuene, 1998
<i>Pagellus bogaraveo</i> , <i>Psetta maxima</i> , <i>Solea senegalensis</i> , <i>Umbrina cirrosa</i>	fish meal and fish oil substitution by plant materials	none	Regost et al., 2003; Segato et al., 2005b; Palmegiano et al., 2007; Figueiredo-Silva et al., 2010; Cabral et al., 2013
<i>Psetta maxima</i> , <i>Solea senegalensis</i>	fish meal and fish oil substitution by plant materials	Increase in plant diets	Valente et al., 2010; Altundag et al., 2014
<i>Pagellus bogaraveo</i> , <i>Sparus aurata</i>	Feeding intensity	Increase or increase tendency in highly-fed fish	Flos et al., 2002; deAlmeida Ozorio et al., 2009
<i>Dentex dentex</i> , <i>Sparus aurata</i>	fasting	Decrease in prolonged fasting	Grigorakis & Alexis, 2005; Suárez et al., 2010.

Table 7: Fillet fatty acid composition (as % of total fatty acids) of farmed Mediterranean fish species. Values are weighed means. In parenthesis the total number of studies taken into account and the total number of fish that contributed to the weighed mean.

species		SFA	MUFA	n-3 PUFA	n-6 PUFA	EPA	DHA	References
<i>Argyrosomus regius</i>	Farmed	31.2 (7/192)	29.8 (7/192)	26.3 (6/174)	11.4 (6/174)	7.36 (6/174)	14.1 (6/174)	Hernández et al., 2009; Grigorakis et al., 2011; Nevigado et al., 2012; Giogios et al., 2013; Martelli et al., 2013; Sinanoglou et al., 2013
<i>Dentex dentex</i>	Farmed	31.3 (4/83)	27.4 (4/83)	25.9 (3/65)	8.02 (3/65)	4.76 (4/83)	16.5 (4/83)	Chatzifotis et al., 2004; Ozden and Erkan, 2008; Dincer et al.,

								2010; Suárez et al., 2010
	wild	37.1 (2/8)	34.8 (2/8)	14.1 (2/8)	2.98 (2/8)	4.38 (2/8)	11.7 (2/8)	Sorigueret al., 1997; Dincer et al., 2010
<i>Diplodus annularis</i>	wild	38.4 (1/3)	31.0 (1/3)	12.7 (1/3)	3.9 (1/3)	3.44 (1/3)	8.79 (1/3)	Özoğul et al., 2008
<i>Diplodus puntazzo</i>	Farmed	27.0 (7/123)	35.6 (7/123)	25.0 (7/123)	12.4 (7/123)	5.81 (7/123)	9.65 (7/123)	Orban et al., 2000; Rueda et al., 2001; Rondán et al., 2004b; Piccolo et al., 2007; Piedecausa et al., 2007; Dincer et al., 2010; Piccolo et al., 2013
	wild	34.0 (4/73)	35.0 (4/73)	20.8 (4/73)	6.5 (4/73)	5.39 (4/73)	7.70 (4/73)	Rueda et al., 2001; Piccolo et al., 2007;

								Özoğul et al., 2008; Dincer et al., 2010
<i>Diplodus sargus</i>	farmed	31.8 (1/9)	23.1 (1/9)	36.3 (1/9)	5.99 (1/9)	6.04 (1/9)	25.1 (1/9)	Cejas et al., 2004
	wild	37.7 (3/129)	25.4 (3/129)	16.4 (3/129)	7.59 (3/129)	5.45 (3/129)	9.91 (3/129)	Hornung et al., 1994; Özyurt et al., 2005; Özyurt et al., 2006; Özoğul et al., 2008;
<i>Diplodus vulgaris</i>	wild	37.2 (2/13)	25.2 (2/13)	27.1 (2/13)	7.06 (2/13)	6.95 (2/13)	18.3 (2/13)	Özoğul et al., 2008; Prato and Biandolino 2012
<i>Epinephelus auneus</i>	wild	30.9 (1/18)	15.1 (1/18)	34.2 (1/18)	7.08 (1/18)	4.60 (1/18)	29.3 (1/18)	Özoğul et al., 2011b
<i>Liza aurata</i>	wild	33.8 (3/22)	42.4 (3/22)	13.1 (3/22)	6.98 (3/22)	5.16 (3/22)	6.04 (3/22)	Kamden et al., 2008; Özoğul et al.,

								2008; Prato and Biandolino, 2012
<i>Liza ramada</i>	wild	27.8 (3/15)	40.4 (3/15)	19.7 (3/15)	6.15 (3/15)	10.5 (3/15)	5.21 (3/15)	Kamden et al., 2008; Özoğul et al., 2008; Nevigado et al., 2012
<i>Liza saliens</i>	wild	27.3 (2/12)	40.8 (2/12)	18.3 (2/12)	7.81 (2/12)	13.2 (2/12)	2.60 (2/12)	Kamden et al., 2008; Özoğul et al., 2008
<i>Mugil cephalus</i>	wild	36.6 (2/7)	23.0 (2/7)	18.4 (2/7)	6.98 (2/7)	9.98 (2/7)	6.40 (2/7)	Özoğul and Özoğul 2007; Özoğul et al., 2008
<i>Pagellus bogaraveo</i>	farmed	30.3 (2/48)	34.2 (2/48)	18.5 (2/48)	9.40 (2/48)	4.82 (2/48)	9.71 (2/48)	Palmegiano et al., 2007; Figueiredo Silva et al.,

								2010
	wild	30.6 (1/3)	30.6 (1/3)	26.3 (1/3)	3.17 (1/3)	8.6 (1/3)	13.5 (1/3)	Soriguer et al., 1997
<i>Pagellus erythrinus</i>	wild	52.7 (2/37)	38.8 (2/37)	5.48 (2/37)	1.57 (2/37)	1.18 (2/37)	2.95 (2/37)	Özoğul and Özoğul, 2007; Koubaa et al., 2014
<i>Pagrus pagrus</i>	farmed	31.2 (3/47)	42.1 (3/47)	20.7 (3/47)	5.84 (3/47)	5.58 (3/47)	12.8 (3/47)	Rueda et al., 1997; Loukas et al., 2010; Kalinowski et al., 2015
	wild	36.9 (2/24)	29.7 (2/24)	25.6 (2/24)	7.01 (2/24)	4.12 (2/24)	20.5 (2/24)	Rueda et al., 1997; Loukas et al., 2010
<i>Psetta maxima</i>	farmed	22.1 (2/9)	21.9 (2/9)	26.2 (2/9)	16.9 (2/9)	6.02 (2/9)	16.8 (2/9)	Nevigado et al., 2012; Altundag et al., 2013
	wild	28.8 (1/3)	23.6 (1/3)	37.2 (1/3)	4.15 (1/3)	8.00 (1/3)	24.5 (1/3)	Soriguer et al., 1997
<i>Sciaena</i>	farmed	41.5	37.8	12.3	8.14	-	-	Cakli et al.,

<i>umbra</i>		(2/45)	(2/45)	(2/45)	(2/45)			2006; Dincer et al., 2010
	wild	52.1 (2/45)	33.3 (2/45)	8.99 (2/45)	2.85 (2/45)	3.21 (1/5)	6.45 (1/5)	Cakli et al., 2006; Dincer et al., 2010
<i>Seriola dumerili</i>	farmed	31.6 (2/19)	27.4 (2/19)	25.5 (2/19)	14.0 (2/19)	7.57 (2/19)	12.8 (2/19)	Haouas et al., 2010; Rodriguez-Bareto et al., 2012
	wild	36.0 (1/9)	34.0 (1/9)	24.4 (1/9)	4.80 (1/9)	2.66 (1/9)	18.8 (1/9)	Rodriguez-Bareto et al., 2012
<i>Solea senegalensis</i>	farmed	28.1 (2/84)	24.5 (2/84)	32.1 (2/84)	10.2 (2/84)	3.65 (2/84)	20.8 (2/84)	Valente et al., 2011; Cabral et al., 2013
<i>Solea Solea</i>	farmed	25.5 (2/51)	34.4 (2/51)	20.5 (2/51)	9.37 (2/51)	4.70 (2/51)	13.1 (2/51)	Piccolo et al., 2008; Nevigado et al., 2012
	wild	29.2 (4/74)	16.9 (4/74)	24.0 (4/74)	7.32 (4/74)	3.90 (4/74)	19.5 (4/74)	Gökçe et al., 2004; Özoğul

								and Özoğul, 2007; Ersoy et al., 2008; Özoğul et al., 2011b
<i>Umbrina cirrosa</i>	wild	40.9 (1/3)	23.0 (1/3)	24.0 (1/3)	1.69 (1/3)	6.52 (1/3)	17.0 (1/3)	Özoğul et al., 2008

Table 8: n-3 PUFA nutritional value of farmed Mediterranean fish species. Values are expressed as g/ 100g fish fillet. In the third column appears what % of daily recommended intake they represent, based on EFSA recommendation for healthy adults*.

Species		n-3 PUFA	EPA+DHA	EPA+DHA (% Daily recommended intake)
<i>Argyrosomus regius</i>	Farmed	0.56	0.45	181.9
<i>Dentex dentex</i>	Farmed	0.98	0.80	320.8
	wild	0.27	0.31	122.7
<i>Diplodus puntazzo</i>	Farmed	2.10	1.30	518.8
	wild	0.58	0.37	147.0
<i>Diplodus sargus</i>	farmed	2.11	1.82	726.8
	wild	0.25	0.23	92.6
<i>Diplodus vulgaris</i>	wild	0.62	0.57	229.1
<i>Epinephelus auneus</i>	wild	0.26	0.26	104.4
<i>Liza aurata</i>	wild	0.59	0.50	201.9
<i>Liza ramada</i>	wild	0.64	0.51	203.9
<i>Liza saliens</i>	wild	0.53	0.45	181.6
<i>Mugil cephalus</i>	wild	0.39	0.35	138.0
<i>Pagellus bogaraveo</i>	farmed	1.35	1.06	422.0
	wild	1.14	0.95	381.9

<i>Pagellus erythrinus</i>	wild	0.21	0.16	63.8
<i>Pagrus pagrus</i>	farmed	0.87	0.77	307.0
	wild	0.22	0.21	83.1
<i>Psetta maxima</i>	farmed	0.41	0.36	143.3
	wild	0.36	0.32	126.1
<i>Sciaena umbra</i>	farmed	0.39	-	-
	wild	0.13	0.14	54.2
<i>Seriola dumerili</i>	farmed	1.70	1.35	540.9
	wild	0.89	0.78	312.4
<i>Solea senegalensis</i>	farmed	0.47	0.36	143.8
<i>Solea solea</i>	farmed	0.54	0.47	189.0
	wild	0.13	0.13	51.0
<i>Umbrina cirrosa</i>	wild	0.22	0.22	86.7

*Source : EFSA, 2010, <http://www.efsa.europa.eu/en/press/news/nda100326.htm>




Fig. 1: Annual fluctuations of fatty acids (expressed in percentages of total fatty acids) in dentex *Dentex dentex* during growth. Data obtained from Chatzifotis et al. (2004)