



Impact of consumption of repeatedly heated cooking oils on the incidence of various cancers- A critical review

Kumar Ganesan, Kumeshini Sukalingam & Baojun Xu

To cite this article: Kumar Ganesan, Kumeshini Sukalingam & Baojun Xu (2017): Impact of consumption of repeatedly heated cooking oils on the incidence of various cancers- A critical review, Critical Reviews in Food Science and Nutrition, DOI: [10.1080/10408398.2017.1379470](https://doi.org/10.1080/10408398.2017.1379470)

To link to this article: <http://dx.doi.org/10.1080/10408398.2017.1379470>



Accepted author version posted online: 19 Sep 2017.



Submit your article to this journal [↗](#)



View related articles [↗](#)



View Crossmark data [↗](#)

Impact of consumption of repeatedly heated cooking oils on the incidence of various cancers- A
critical review

Kumar Ganesan, Kumeshini Sukalingam, Baojun Xu*

Food Science and Technology Program, Beijing Normal University – Hong Kong Baptist
University United International College, Zhuhai 519085, China

*Corresponding author: Prof. Baojun Xu Corresponding author address: 28, Jinfeng Road,
Tangjiawan, Zhuhai, Guangdong 519085, China, Email: baojunxu@uic.edu.hk. Tel: 86-756-
3620636; Fax: 86-756-3620882

Abstract

Repeated heating of vegetable oils at high temperatures during cooking is a very common cooking practice. Repeatedly heated cooking oils (RCO) can generate varieties of compounds, including polycyclic aromatic hydrocarbons (PAH), some of which have been reported as carcinogenic. RCO is one of the commonly consumed cooking and frying medium. These RCO consumption and inhalation of cooking fumes can pose a serious health hazard. Taking into account exploratory study, the present review aims to provide the consumption of RCO and its fumes cause the high incidence of genotoxic, mutagenic, tumorigenic and various cancers. The information on RCO and its fumes were collected through a library database and electronic search (ScienceDirect, PubMed, and Google Scholar). Remarkable studies demonstrated that the health adverse effects of RCO and its cooking fumes have been often attributed to their detrimental properties and ease to genotoxic, mutagenic and carcinogenic activities. RCO and its cooking fumes were found to enhance the incidence of aberrant cells, including breaks, fragments, exchanges and multiple chromosomal damages and micronuclei in a dose-dependent manner. Furthermore, the large consumption of RCO has been associated with a number of malignancies, including lung, colorectal, breast, and prostate cancers. The present review provides additional insights into the polluting features of PAHs produced various cancers via cooking activities in indoor environments.

Keywords

Repeatedly heated vegetable oils; polycyclic aromatic hydrocarbons; genotoxicity; mutagenicity; carcinogenicity; cancers

Abbreviation

AGEs	advanced glycation end products
ALA	alpha-linolenic acid
ALP	alkaline phosphatase
AST	aspartate transaminase
ATP	adenosine triphosphate
BMI	basal metabolic index
CI	confidence interval
CVD	cardiovascular diseases
DHA	docosahexaenoic acid
DNA	deoxy ribonucleic acids
FFA	free fatty acids
GC-MS	gas chromatography- mass spectroscopy
GGT	gamma glutamyl transferase
GLA	gamma-linolenic acid
GPx	glutathione peroxidase
GSH	glutathione
GSR	glutathione reductase
GST	glutathione-s-transferase
HCA	heterocyclic amines
HPLC	high performance liquid chromatography

IFN interferon

IL interleukin

MDA malondialdehyde

MTT (3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide

MUFA monounsaturated fatty acids

NF- κ B nuclear factor kappa B transcription factor

OR odds ratio

PAH polycyclic aromatic hydrocarbons

PCa prostate cancer

PCNA -proliferating cell nuclear antigen

PM particulate matter

PUFA polyunsaturated fatty acids

RCO repeatedly heated cooking oils

ROS reactive oxygen species

TBARS thiobarbituric acid reactive substances

TGF transforming growth factor

TGL triglycerides

TUNEL terminal deoxynucleotidyl transferase dUTP nick end labeling

UV ultra violet

VEGF vascular endothelial growth factor

VEGFR vascular endothelial growth factor receptor

Introduction

Vegetable oil is one of the main dietary components in the day to life food consumption and is used in nearly all types of food preparations including frying, baking, sauteing, dressing, marinating, and extrusion cooking. They are generally obtained from oil seeds (i.e. mustard, sunflower, cottonseed, corn, and coconut), food legumes (i.e. soybean, peanut), nuts (i.e. almond) or the soft substances of fruits (i.e. olives). They are primarily composed of triglycerides and serves as main sources of fat consists of three fatty acids and one molecule of glycerol (Sayon-Orea et al., 2015). The minor components are free fatty acids, fat-soluble vitamins, pigments, phospholipids, waxes, sterols and fatty alcohols. Vegetable oils contain different kinds of fatty acids and their compositions are widely varied; however, one type of fatty acid is generally predominant over the other fatty acids. The chief fatty acid in olive oil is oleic acid, monounsaturated fatty acids; and the predominant fatty acid in sunflower oil is linoleic acid, polyunsaturated fatty acids (Foster et al., 2009). Physical and chemical characteristic features of vegetable oils are influenced by the quantity of fatty acid and the place, where they positioned on the glycerol (American Oil Chemists' Society, 2006).

Deep frying of vegetable oils is one of the most common methods of food preparation worldwide. The practice of reusing oils for repeated frying is also common in an attempt to cost-effective. When oils are used in frying, the temperature ranges between 170° to 220°C and it undergoes physicochemical changes such as oxidation, hydrolysis, cyclization, and polymerization and eventually, it degrades to volatile compounds (Ku et al., 2014). These repeatedly heating alter manifestation of the oil with high viscosity, color darkening, froth and smoke point reduction, which makes more harmful when these oils consume along with the food. Many studies showed that the reheated oils cause genotoxic (Dung et al., 2006), mutagenic and carcinogenic potential (Srivastava et al., 2010a, 2010b). Further, the studies demonstrated that the deleterious health effects of consumption of reheated oil cause increased blood pressure (Psaltopoulou et al., 2004; Leong et al., 2009; 2012), cardiovascular diseases (Ng et al., 2012; 2014a, 2014b), atherosclerosis (Adam et al., 2008), endothelial dysfunction (Lopez-Garcia et al., 2005), impaired vasorelaxation responses (Owu et al., 1997), hypertension (Soriguer et al., 2003) and elevated LDL, and lipid peroxidation (Siti et al., 2008).

2. Nutritional and health benefits of cooking oil

Cooking oil is a crucial part of our day to day life in the diet possess the primary sources of lipid, which gives energy (9 calories/gram) to the human and a chief ingredient of biomembrane (Vaskova and Buckova, 2015) and building blocks of many lipid hormones (Falade et al., 2017). In addition, the nutritional and health benefits of these cooking oils are massive and this can be endorsed to their individual ingredients such as fatty acids in the amount of saturated to unsaturated fats; and monounsaturated to polyunsaturated fats. In addition, these fatty acids possess an assortment quantity of natural antioxidants such as vitamin A, E, and carotenoids that protect cells from free radical damage and to prevent the incidence of oxidative rancidity in the frying oil (Fahy et al., 2009).

Polyunsaturated fatty acids (PUFAs) are present in excellent quantities in vegetable oils such as sunflower oil (Das, 2008) and offer as major integral part in mitochondrial membranes. The effectiveness of electron transport system is terminated due to the inadequacy of PUFA. PUFA, in general, are anti-atherogenic properties that reduce serum cholesterol and triglycerides (Endo and Arita, 2016). The biological importance of prostaglandins, thromboxanes, and leukotrienes are produced by arachidonic acid, one of the major PUFA constitutes 10–15% of the total fatty acids in the membranes. Generally, PUFAs enhances cementing constituents and fluidity in the membrane and protects the cell. Docosahexaenoic acid (DHA) is usually synthesized by alpha-linolenic acid (ALA), another chief PUFA that found in soybeans, flax seeds, walnuts, and spinach (Lv et al., 2016). Gamma-linolenic acid (GLA) prevents CVD through expanding the blood vessels, lowering BP, and avoids the occurrence of atherosclerosis (Kitamura et al., 2011).

In addition, GLA inhibits the augmentation of tumors and various cancers. The dietary sources of GLA are primarily obtained from plant seed including primrose, black currant, and borage. Biologically significant PUFAs are eicosapentaenoic acid (EPA) and DHA synthesized from ALA, jointly called as omega-3 fatty acids. The main sources of ALA in the diet are walnuts and hazelnuts (Calder, 2012). Intake of these omega-3 fatty acids is clinically important, which reduces the activities of white blood cells and generate mediators of inflammation. Furthermore, they reduce the ability of blood platelets to release thromboxanes and stimulate the

blood clotting mechanism. Hence, omega-3 fatty acids serve as the precursors for potent anti-inflammatory lipids (Anand et al., 2008), powerful cell signaling role and thus acting as a reservoir of the biomolecules (Healy et al., 2000).

Monounsaturated fatty acids (MUFAs) are chiefly present in the vegetable oil such as peanut oil, mustard oil, olive oil and canola oil (Guthrie and Pcciano, 1995), which prevents coronary heart disease (Hu and Willett, 2002). It is well known to have a favorable effect on the blood lipid profile and thus reduce the risk of CVD (Demonty et al., 2006; Metcalf et al., 2007). In addition, these fatty acids are less likely to react with ROS and more stable in oxidative stress settings when compared with PUFA (Diniz et al., 2004). However, the experimental studies associations between dietary consumption of MUFAs and risk of coronary heart disease have been still uncertain (Jakobsen et al., 2004).

3. Deep fried cooking oil

Frying is a commonly employed method for preparing food as it can make the food very appetizing and aromatic, which provides delicious taste. The common issue raised in cooking oil nowadays is repeatedly heating and used it over several times. This practice may not be regular in residential cooking, however, it is more common in hawkers which can aid them to reduce their expenses and earn more income. Normally, cooking oil can be obtained from plant or animal sources. The most commonly used cooking oils are peanut oil, corn oil, palm oil, coconut oil, sunflower oil, and lard. During cooking, hydrolysis of oil occurs in the beginning when moist food is fried in hot oil. This reaction enhances the acid value of oil because of fatty acid production from triglycerides. Oxidation of oil during frying is the key of concern. Oxidation occurs due to reaction with the atmospheric oxygen. Auto-oxidation can also occur though the oil is not heated, this process is supported by external temperature or exposed to UV light (Gotoh et al., 2007). Three phases that occur during oxidation of oil are initiation, propagation and termination phases (Gotoh et al., 2007). These processes generate more free radicals such as lipid peroxides, PAH, aldehydes, ketones, alcohols, and acids (Romero et al., 2006; Choe and Min, 2007). Oxidation mechanism is more prone to occur in unsaturated fatty acid compared to saturated form (Mensah and Obeng, 2013). When the oil is heated, thermal dissociation occurs in unsaturated fatty acid containing double bonds, which allows removal of the hydrogen atom to

form alkyl radical or lipid radical and this process occurs in initiation phase (Gotoh et al., 2007, Good, 2012). In the propagation phase, unstable lipid radical become more unstable free radicals; they are generally called as peroxy lipid free radicals. In the termination phase, unstable two free radicals react and form non-radical species (Romero et al., 2006). The physicochemical changes during thermal oxidation of cooking oils are depicted in **Table 1**.

4. Generation of carcinogenic agents and fumes

Frying is a process that persuades a multitude of biochemical reactions in the hot medium and produces an abundance of chemical compounds (Belitz et al., 2004). These degraded compounds may include FFA, aldehydes, alkanes, 4-hydroxy nonenal, hydro-peroxide volatile compounds, and polymerized triglycerides (Choe and Min, 2007). The quantities of degraded products elevate with the length of heating time at high temperature. Thus, the more the oil is heated, the more deprivation takes place and the more noxious compounds and lipid peroxidation products are formed in the cooking oil (Lapointe et al., 2006; Romero et al., 2006). Several factors can influence the quality value of cooking oil during heating, including aeration, temperature, duration of the heating, the type of oil, the saturation ratio of the oil, and the presence of a catalyst (Gupta, 2005; Falade and Oboh, 2015). The degree of changes in the physicochemical properties of the oil dictates the quality of the oil for human consumption.

Research studies have indicated that RCO is generally used 3--6 times before being discarded as waste (Mensah and Obeng, 2013). RCO lowers the smoke point, which is the temperature at which the oil breaks down creating acrolein, an obnoxious smelling compound; and visible cooking fumes are produced (Good, 2012). In addition, almost sixteen PAHs were recognized in fumes released during deep-frying of rapeseed, soybean, peanut, and olive oil (Yao et al., 2015). Furthermore, Li et al. (1992) also recognized five PAHs (benzo(a)pyrene, dibenzo(a,h)anthracene, benzo(a)anthracene, dibenzo(a,b)anthracene, and benzo(e)pyrene) in six cooking oil fumes. Zhao et al. (2006) also found that cooking oil increases lung cancer susceptibility in China due to frequent deep-frying fumes and deprived ventilation in kitchens. Chen et al. (2012) also identified 21 PAHs and Jørgensen et al. (2013) found 32 PAHs from deep-frying oil medium. In addition, Dost and Ideli (2012) identified 9 PAHs in fumes produced when meat and fish were roasted using three different edible oils. See and Balasubramanian

(2008) also examined the chemical characteristics of fine particulate matter (PM_{2.5}) emitted when plain tofu was cooked using various cooking methods with cooking oil. The list of PAHs in cooking oil fumes (Li et al., 1992; 2003; Chen et al., 2012; Yao et al., 2015) are depicted in **Figure 1**.

PAHs are organic compounds, containing two or more fused benzene rings composed of hundreds of chemical groups (Abdel-Shafy and Mansour, 2016), and they are consistent organic pollutants exhibit teratogenic, genotoxic (Srivastava et al., 2010a, 2010b), carcinogenic (Sinha et al., 1999; Ramesh et al., 2004; Pandey et al., 2006), and mutagenic effects in living organisms (Wornat et al. 2001; Dung et al., 2006). Increase in mutagenicity is proportionally related to its frying time (Chen et al., 2003). Humans are exposed to PAHs largely through breathing and ingestion (Chen et al. 2012), and these PAHs threaten human health to cause lung, stomach, esophagus, and skin cancers (Wu and Yen, 2004; Li et al. 2011). Pandit et al. (2001) also reported that RCO generates a diversity of volatile, semi-volatile, and particulate matters. The epidemiological study was done by Lopez- Abente et al. (2001) suggested that people exposed to PAH are at the risk of developing cancers. Another epidemiological study has also indicated that a high intake of RCO containing saturated fat increases the risk of cancers (Houthuijzen, 2016).

Degraded compounds from RCO may form DNA adducts, which have been proposed as predictive biomarkers of human cancers (Korsh et al., 2015). Results of this study revealed the presence of six times higher amount of total PAH in RCO compared with the fresh cooking oil. The greater amounts of PAH are reported to have significant carcinogenic and mutagenic potential (Isidori and Parrella, 2009), which are dependent on the number of cycles used for heating (Pandey et al., 2006). The genotoxicity and carcinogenicity assays performed to detect both the chromosomal aberration and micronuclei induction in the bone marrow (Shukla et al., 2003).

By using these two assays in this study, RCO was found to enhance the incidence of aberrant cells, including breaks, fragments, exchanges and multiple chromosomal damages and micronuclei in a dose-dependent manner. Chromosomal aberration and micronuclei induction by PAH (Shukla et al., 2003) are well documented. The results of this study also suggest that RCO

may contain a high amount of PAH (Marsili et al., 2001). Repeatedly heated cooking oil emits PAHs and subsequent causes of cancer are depicted in **Figure 2**.

5. Mutagenicity and genotoxicity effect of PAHs in RCO

Benzo[a]pyrene was classified as group 1 carcinogens (IARC 2012); dibenz[a,h]anthracene was originally classified as “probable” human carcinogens (2A); whereas, naphthalene, benzo(a)anthracene, benzo[b]fluoranthene, benzo[k]fluoranthene, and indeno[1,2,3-cd]pyrene were deemed of “possible” human carcinogens (2B) (IARC 2010). Dibenz[a,h]anthracene was detected in nearly all samples collected in peanut and olive oil cooking fumes. Regarding “possible” human carcinogen (2B) compounds, naphthalene, benzo(a)anthracene were found in all the samples collected in cooking fumes. Indeno[1,2,3-cd]pyrene and benzo[b]fluoranthene were observed in all deep-fried food samples.

PAHs are highly lipid soluble and thus readily absorbed in the gastrointestinal mammals. They are rapidly distributed in a variety of tissues with the marked tendency for localization in body fat (Abdel-Shafy and Mansour, 2016). These carcinogenic PAHs are genotoxic and induce mutations that initiate cancer. These progressions of cancer chemically induce by enzymes in metabolites that react with DNA, leading to mutations. When the DNA sequence is altered in genes that regulate cell replication, cancer can result. PAHs can bind to DNA at specific sites, forming bulky complexes called DNA adducts that can be stable or unstable (*Henkler et al., 2012*). Stable adducts may lead to DNA replication errors, while unstable adducts react with DNA strand, removing a purine base (*Henkler et al., 2012*). Such mutations if they are not repaired, can transform genes encoding for normal cell signaling proteins into cancer-causing oncogenes (Baird et al., 2005). Low molecular weight PAHs are more potent as co-carcinogens during the promotional stage of cancer. In this stage, first cells are removed from growth suppressing signals from its neighboring cells and begin to clonally replicate (*Ramesh et al., 2004*). Low molecular weight PAHs that have bay or bay like regions can dysregulate gap junction channels, interfering with intracellular communication, and affect mitogen-activated protein kinases that activate transcription factors involved in cell proliferation (*Ramesh et al., 2004*).

The bulk intake of these fried foods has been related with various malignancies, including colon, rectum, breast, kidney and pancreas cancers (Gonzales et al., 2014). These malignancies may be due to the reduction of omega-3: omega- 6 proportions, and intake of frying oil by-products (Ansorena et al., 2010). The bulk consumption of fried food was related with a 1.3- to 2.3-fold increased risk of prostate cancer (Li et al., 2014; Stott-Miller et al., 2013). *In-vitro* and *in vivo* genotoxicity and mutagenicity effect of RCO are depicted in **Table 2**.

6. Tumorigenic effect of PAHs in RCO

Extensive and systematic animal studies on the tumorigenicity effect of PAH, possess diol epoxides, which are main mutagenic and carcinogenic species (Conney et al., 2001). These metabolites are effectively changed by epoxide ring by electrophilic carbonium ions that strongly bind to nucleophilic sites of DNA and in proteins. Furthermore, these oxidative metabolites produce BaP-7, 8-dihydrodiol-9, 10-epoxide (BPDE) from BaP-7,8-dihydrodiol and found in almost all tissues, while examined. This metabolite is the only isomer with high tumorigenic potential and is the predominantly bound to DNA (Conney et al., 2001). *In vitro* and *in vivo* studies have also shown the cytotoxicity effect of BaP. Various other PAHs cause very severe, long-lasting hyperplasia and other adverse effects similar to the neoplastic changes (Charles and Adaku, 2010). In addition, BaP and various other PAHs administered through oral cause tumors in the forestomach, lungs, liver, pancreas, and breast of rodents, while mono- and dinitro pyrenes also cause tumors in breast and pituitary (Imaida et al., 1991). The impacts of the tumorigenic effect of RCO are depicted in **Table-3**.

7. Impact of RCO on various cancers

The impacts of covalent binding of PAHs to DNA in the various tissues of rodents have been extensively studied in *in vivo* and *in-vitro* (Boström et al., 2002). Experimental animal studies have demonstrated that (+)-anti-BPDE-DNA adducts are the primary adducts formed in lung, liver, heart, pancreas, stomach, skin and kidney. However, the studies using systemic administration in rodents, the quantities of DNA adducts were higher in liver, lung, and spleen. On the other hand, DNA adducts were lower in kidney and stomach (Ross et al. 1990, 1991; Rojas et al., 2001; Pauk et al., 2013). According to the epidemiological studies, DNA adducts and its several factors associated with an increased risk of cancer and are related with high risk

of atherosclerosis. This is because of somatic mutation involved in the formation of the atherosclerotic plaque (Pulliero et al., 2015). PAHs, including dibenz[a,h]anthracene, dibenz[a,c]anthracene, 7,12-dimethylbenz[a]anthracene and BaP, were shown to act as initiators and/or accelerators of various cancers in chickens, pigeons and mice (Wakabayashi, 1990). Newborn mice are very susceptible to the carcinogenic action of PAHs. When the administration of BaP and other PAHs in rats through intraperitoneal or subcutaneous cause liver and lung tumors within 6 months (Platt et al., 1990). In addition, nitro-PAHs cause leukemia, breast and colorectal cancer (Imaida et al., 1992). Intrapulmonary injection of BaP and other PAHs also produced lung tumors in rats and hamsters (Hartung et al., 1990). However, the significance of this pathway for PAH carcinogenicity in experimental animals and humans are as yet unclear (Cavalieri and Rogan, 1996). The impacts of RCO on various cancers are depicted in **Table-4**.

7.1. Lung cancer

The basis of the epidemiological studies, cooking fumes exposure to women from cooking oils showed to be a primary risk factor for various cancers including lung cancer (Purcaro et al., 2006; Yao et al., 2015). A study in Taiwan associated with three different cooking oils and their fumes were analyzed. Various PAHs including (benzo(a)pyrene (B(a)P), benz(a)anthracene (B(a)A), and dibenz(a,h)anthracene (DB(ah)A) were detected in all samples. A concentration of DB[a,h]A and B[a]A were 1.9 and 2.2 $\mu\text{g}/\text{m}^3$ in fumes from lard oil, 2.1 and 2.3 $\mu\text{g}/\text{m}^3$ in soybean oil, 1.8 and 1.3 $\mu\text{g}/\text{m}^3$ in peanut oil, respectively. Benzo[a]pyrene (B[a]P) was identified in fume samples of soybean and peanut oil, in concentrations of 19.6 and 18.3 $\mu\text{g}/\text{m}^3$ respectively. This experimental observation concluded that women exposed to the emitted fumes of RCO are at increased risk of lung cancer (Chiang et al., 1997). Earlier studies have also suggested a positive correlation between the emitted fumes of RCO and high risk of lung cancer (Wu et al., 1998; Purcaro et al., 2006; Yao et al., 2015). Another investigation related to population-based case-control study among Chinese women in Shanghai, there was an increased risk of cancer in mammary gland related with high intake of deep-fried red meat (OR = 1.92; 95% CI, 1.30–2.83) and deep-fried fish (OR = 1.52; 95% CI, 1.05–2.22) (Dai et al., 2002). Consumption of these deep-fried food has been highly connected with the risk of pancreatic (Ji et

al., 1995), lung (Huang et al., 1992), oral/pharyngeal, esophageal (Galeone et al., 2005), and laryngeal (Bosetti et al., 2002) cancers.

7.2. Breast cancer

Evidence from some animal and epidemiological studies have suggested an increased risk of breast cancer associated with high intake of meat (De Stefani et al., 1997; Zheng et al., 1998; Sinha et al., 2000; Dai et al., 2002). In addition, the studies revealed the breast cancer is caused due to high intake of plant seed cooking oils (D'Avanzo et al., 1991; Favero et al., 1998; Zock et al., 1998). It has been reported that RCO not only produces fumes containing mutagenic compounds (i.e. benzene, acrolein, 1,3-butadiene, and formaldehyde), but also produces non-volatile hazardous compounds (hydroperoxides, *trans*-fatty acids, and aldehydes) (Grandgirard et al., 1984, Goburdhun et al., 2001). Both these hydroperoxides and aldehydes are endogenous reactive metabolites and possess mutagenic and carcinogenic properties (Gupta et al., 1999). Other epidemiological studies have also reported that a positive correlation between the intake of oil containing linoleic and linolenic acid and breast cancer risk (Klein et al., 2000; Pala et al., 2001). In Western countries, hydrogenated oils have been generally used for home cooking and food formulation and food processing in food industry have *trans*- fatty acids. These *trans*- fatty acids could be a high-risk factor for breast cancer (Kohlmeier et al., 1997).

7.3. Colorectal cancer

Normally, heterocyclic amines (HCAs) have been generated during the frying of protein rich food including meat, eggs and fish and HCAs have been proved as bacterial mutagens and animal carcinogens (Pfau and Marquardt, 2001). In an experimental animal study, these amines cause cancer especially in the organs such as breast, colorectal, stomach and pancreas (Galeone et al., 2007). An American case-control study also further reported that a positive relationship between pan-fried red meat intake and colon cancer risk (Butler et al., 2003), and other epidemiological studies also investigated a direct relationship with colorectal adenomas and cancer (Sinha et al., 2001; Navarro et al., 2004). In a cohort study on Swedish women, including 741 cases of colorectal cancer, there was a connection between the quantities of acrylamide intake and colorectal cancer (Mucci et al., 2006).

7.4. Prostate cancer

In a population-based study of 1096 controls, 717 localized and 1140 advanced cases, it was observed an increased risk of prostate cancer (PCa) relation with the intake of pan-fried meat (OR = 1.4; 95% CI, 1.0–1.8) adjusted for age, BMI, total intake of calorie, and family history (Joshi et al., 2012). In addition, the study had further reported the more relative risks of PCa connected with high intake of fish cooked with high-temperature cooking methods (Joshi et al., 2012). The carcinogen, acrylamide, is produced in an enormous quantity during deep-frying, especially carbohydrate-rich foods such as potatoes (Tareke et al., 2002). The positive correlation between acrylamide and PCa risk have been reported in two case-control studies and four cohort studies (Lipworth et al., 2012).

Normally, foods cooked with high temperature produces high levels of advanced glycation end products (AGEs) (Uribarri et al., 2010). For instance, a chicken breast deep-fried for 20 min generated nine times more quantities of AGEs as a chicken breast boiled for 60 min (Uribarri et al., 2010). These AGEs have been linked to increased oxidative stress and proinflammatory effects (Uribarri et al., 2007). These inflammatory effects may also play a vital role in PCa (De Marzo et al., 2007), and further *in vitro* studies have also suggested that the interactions between AGEs and the receptor of AGE could be a driver in PCa progression (Elangovan et al., 2012). Furthermore, the study associated with fast food restaurants and cooking methods, where the French fries prepared by commercial vegetable oils such as sunflower seed, canola, soybean, cottonseed, and palm oil (Jahren and Schubert, 2010). These oils are normally containing a high amount of omega-6 fatty acids, which have also been reported to induce proliferation of human prostate tumor (Hughes-Fulford et al., 2006). In animal study, the intake of omega-6-rich diets resulted in prostate tumor growth and development (Kelavkar et al., 2009). In human-based studies reported a higher risk of PCa associated with high intake of omega-6 fatty acids (Neuhouser et al., 2007).

8. Public awareness and legislation

An adequate quantity of oil is regularly required for deep-frying of the food materials. Therefore the person may regularly keep the used frying oil for future use. This practice usually maintains at household kitchens as well as various small scaled industrial sectors. Reusing this

frying oil is thought to limit the cost of food preparation without contemplating the potential toxic effects on human health. Despite the fact that about 70% of populations in Kuala Lumpur, Malaysia realize that the quality of cooking oil does not stay the same after deep-frying (Azman et al., 2012), reusing frying oil is still generally practiced in the population in Kuala Lumpur.

Public awareness with respect to the toxic of using RCO is still not satisfactory in developing and under-developed nations. About 60% of people have conceded utilizing still the same part of RCO (Phiri et al., 2006; Azman et al., 2012; Abdullah et al., 2010). A chemical investigation of the oils demonstrated the presence of FFA is almost thrice in the ranges of 0.84–1.4112 compared with the fresh oil (0.42). The peroxide value in RCO raise to 14.7–16.6 compared with 9.0 in the fresh oil (Phiri et al., 2006). In a Japanese survey, about 90% of the respondents had not been aware of food quality and safety issues of RCO (Phiri et al., 2006). Another study conducted in Japan found that restaurant frying oils were used at 180 °C for 3 h /day for five continuous days before disposing of those oils (Totani et al., 2006).

The impacts of heating process on the safety of cooking oils have drawn enormous attention from the researchers worldwide. During the recent 7th International Symposium on Deep-Fat Frying held in San Francisco, USA, scientists have reaffirmed that the best indices for evaluation of the used oils are total polar materials and polymeric TGL by using recognized methods (European Federation for the Science, Technology of Lipids, 2013). Peroxide value, FFA value, and anisidine value should not be used as regulatory indices when it comes to monitoring and comparing the degree of degradation of different frying oils. The cut-off point for rejection of reused frying oil has been proposed to be set at a content range of 20–27% for polar compounds (Paul and Mittal, 1997), which with the implementation of Hazard Analysis and Critical Control Point regulation has been employed in the national food laws in several European countries (Dobargarnes and Márquez-Ruiz, 1998).

Nonetheless, the index does not speak the careful bunch of products from lipid peroxidation as 25% of polar compounds in cooking oil relate to a considerably higher content of TGL oligomers (Bastida et al., 2002). Furthermore, some secondary oxidation products may have potential harmful to the human health. A study has found significant concentrations of toxic aldehydes present in the oil albeit the 25% limit for polar compounds is reached in the oil

(Guillén et al., 2012). These findings recommend that the limit may overlook and permits alternative arrangement of possible hazardous compounds. Since the products of lipid oxidation are nearly connected with CVD risk factors and cancers, the quality of the vegetable oil consumed daily unquestionably affects the cardiovascular system and progression of tumor/malignancies in various peripheral organs.

9. Conclusion and future directions

On the basis of the data discussed in the present review, consumption of diets containing RCO could be harmful and cause various cancers in peripheral organs. The nutritive and protective benefits of vegetable oils deteriorate when the oils are repeatedly exposed to extreme heat, air, and moisture, during food preparation, particularly in deep-fat frying. Through a complex series of reaction that occurs during deep-fat frying, various oxidation products including PAHs are formed, affecting the quality of oil physically and chemically. The data in this review suggest the deleterious effects of heated oils and their fumes produce various PAHs, they are mutagenic, genotoxic and carcinogenic agents. Studies report that RCO and its fumes may increase the risk of cancer via DNA mutations that initiate cancer. These progressions of cancer chemically induce by enzymes that react with DNA, leading to mutations. When the DNA sequence is altered in genes that regulate cell replication, cancer can result. However, the issue on re-using oil in food preparation has not been paid enough attention as the economic perspective of the food industry or the society is given more priority than health perspective of the consumers. Public education, awareness, and stringent laws are needed to contain the usage of thermally oxidized oil in food industry and households. Finally, though a substantial amount of *in vitro* and *in vivo* animal studies, the effects of heated oil in humans are still vaguely understood or contradictory. Since oil is an essential dietary component in daily meals, the mechanisms by which heated oil exerts its effects in humans merit further investigation in long-term, larger cohort studies to get ascertainable evidence as well as to increase awareness among the people.

Conflicts of Interest

The authors declare no conflict of interest.

Acknowledgments

The work was jointly supported by two grants UIC 201624 and UIC 201714 from Beijing Normal University-Hong Kong Baptist University United International College, Zhuhai, Guangdong, China.

Author Contributions

K.S. and K.G. conceived, designed and wrote the review; B.X. critically read and improved the manuscript.

References

- Abdel-Shafy, H.I. and Mansour, M.S.M. (2016). A review on polycyclic aromatic hydrocarbons: Source, environmental impact, effect on human health and remediation. *Egyptian J. Petroleum* **25**: 107--123
- Abdullah, A., Suondoh, M.S., Xuan, C.S., Abdul Patah, N., Mokhtar, K., Mohd Fahami, N.A., Saad, Q.H.M., Yusof, K. and Jaarin, K. (2015). Awareness regarding the usage of repeatedly heated cooking oil in Kuala Lumpur, Malaysia. *Int. Med. J.* **17**(4): 310–311.
- Adam, S.K., Sulaiman, N.A., Mat Top A.G. and Jaarin, K. (2007). Heating reduces vitamin E content in palm and soy oils. *Malays. J. Biochem. Mol. Biol.* **15** (2): 76--79.
- Adam, S.K., Soleiman, I.N., Umar, N.A., Mokhtar, N., Mohamed, N. and Jaarin, K. (2008). Effects of repeatedly heated palm oil on serum lipid profile: Lipid peroxidation and homocysteine levels in a postmenopausal rat model. *Mcgill J. Med.* **11**: 145--151.
- Agbaria, R., Gabarin, A., Dahan, A. and Ben-Shabat, S. (2015). Anticancer activity of *Nigella sativa* (black seed) and its relationship with the thermal processing and quinone composition of the seed. *Drug Des. Devel. Ther.* **9**: 3119--3124.
- Aladedunye, F.A. and Przybylski, R. (2009). Degradation and nutritional quality changes of oil during frying. *J. Am. Oil Chem. Soc.* **86**: 149–156.
- Alía, M., Ramos, S., Mateos, R., Granado-Serrano, A.B., Bravo, L. and Goya, L. (2006). Quercetin protects human hepatoma HepG2 against oxidative stress induced by tert-butyl hydroperoxide. *Toxicol. Appl. Pharmacol.* **212**(2): 110--118.
- American Oil Chemists' Society, (2006). Institute of Shortening and Edible Oils: Food Fats and Oils, 9th edition. 1750 New York Avenue, NW, Suite 120 Washington, DC on the Internet at <http://www.iseo.org/foodfats.htm>
- Anand, R.G., Alkadri, M., Lavie, C.J. and Milani, R.V. (2008). The role of fish oil in arrhythmia prevention. *J. Cardiopulm. Rehabil. Preven.* **28**: 92–98.
- Ansorena, D., Guembe, A., Mendizábal, T. and Astiasarán, I. (2010). Effect of fish and oil nature on frying process and nutritional product quality. *J. Food Sci.* **75**(2): H62-67.
- Arora, A. and Shukla, Y. (2004). Induction of preneoplastic altered hepatic foci following dietary sulphur supplementation. *Hum. Exp. Toxicol.* **23**(5): 229--234.

- Azman, A., Mohd Shahrul, S., Chan, S.X., Noorhazliza, A.P., Khairunnisak, M., Nur Azlina, M.F., Qodriyah, H.M., Kamisah, Y. and Jaarin, K. (2012). Level of knowledge, attitude and practice of night market food outlet operators in Kuala Lumpur regarding the usage of repeatedly heated cooking oil. *Med. J. Malaysia* **67**(1): 97–107.
- Baird, W.M., Hooven, L.A. and Mahadevan, B. (2005). Carcinogenic polycyclic aromatic hydrocarbon-DNA adducts and mechanism of action. *Environ. Mol. Mut.* **45** (2-3): 110--114.
- Bastida, S. and Sánchez-Muniz, F.J. (2002). Polar content vs. TAG oligomer content in the frying life assessment of monounsaturated and polyunsaturated oils used in deep-frying. *J. Am. Oil Chem. Soc.* **79**(5): 447--451.
- Belitz, H.-D., Grosch, W. and Schieberle, P. (2004). Food Chemistry, 3rd ed. Springer-Verlag, Berlin. pp. 218--219.
- Bosetti, C., Talamini, R., Levi, F., Negri, E., Franceschi, S., Airoldi, L. and La Vecchia, C. (2002). Fried foods: A risk factor for laryngeal cancer? *Br. J. Cancer* **87**(11):1230–1233.
- Boström, C.E., Gerde, P., Hanberg, A., Jernström, B., Johansson, C., Kyrklund, T., Rannug, A., Törnqvist, M., Victorin, K. and Westerholm, R. (2002). Cancer risk assessment, indicators, and guidelines for polycyclic aromatic hydrocarbons in the ambient air. *Environ. Health Perspect.* **110** (Suppl 3): 451--488.
- Brenes, M., García, A., Dobarganes, M.C., Velasco, J. and Romero, C.N. (2002). Influence of thermal treatments simulating cooking processes on the polyphenol content in virgin olive oil. *J. Agric. Food Chem.* **50**: 5962–5967.
- Butler, L.M., Sinha, R., Millikan, R.C., Martin, C.F., Newman, B., Gammon, M.D., Ammerman, A.S. and Sandler, R.S. (2003). Heterocyclic amines, meat intake, and association with colon cancer in a population-based study. *Am. J. Epidemiol.* **157**: 434--445.
- Cai, Y., Lv, J., Zhang, W. and Zhang, L. (2012). Dietary exposure estimates of 16 polycyclic aromatic hydrocarbons (PAHs) in Xuanwei and Fuyuan, counties in a high lung cancer incidence area in China. *J. Environ. Monit.* **14**(3): 886--892.
- Calder, P.C. (2012). Mechanisms of action of (n-3) fatty acids. *J. Nutr.* **142**: 592S–599S.

- Cao, W., Wan, Y., Liang, Z.H., Duan, Y.Y., Liu, X., Wang, Z.M., Liu, Y.Y., Zhu, J., Liu, X.T. and Zhang, H.X. (2010). Heated lipiodol as an embolization agent for transhepatic arterial embolization in VX2 rabbit liver cancer model. *Eur. J. Radiol.* **73**(2): 412--419.
- Cao, W., Xu, X., Zhang, J. and Duan, Y. (2013). Tumor angiogenesis after heated lipiodol infusion via the hepatic artery in a rabbit model of VX2 liver cancer. *PLoS One* **8**(4): e61583.
- Carrasco-Pancorbo, A., Cerretani, L., Bendini, A., Segura-Carretero, A., Lercker, G., and Fernaández-Gutiérrez, A. (2007). Evaluation of the influence of thermal oxidation on the phenolic composition and on the antioxidant activity of extra-virgin olive oils. *J. Agric. Food Chem.* **55**: 4771--4780.
- Catsburg, C., Kim, R.S., Kirsh, V.A., Soskolne, C.L., Kreiger, N. and Rohan, T.E. (2015). Dietary patterns and breast cancer risk: A study in 2 cohorts. *Am. J. Clin. Nutr.* **101**(4): 817--823.
- Cavalieri, E.L. and Rogan, E.G. (1996). Central role of radical cations in metabolic activation of polycyclic aromatic hydrocarbons. *Xenobiotica*, **25**: 677--688.
- Chang, L.W., Lo, W.S. and Lin, P. (2005). Trans, trans-2,4-decadienal, a product found in cooking oil fumes, induces cell proliferation and cytokine production due to reactive oxygen species in human bronchial epithelial cells. *Toxicol. Sci.* **87**(2): 337--343.
- Chang, Y.C. and Lin, P. (2008). Trans, trans-2,4-decadienal induced cell proliferation via p27 pathway in human bronchial epithelial cells. *Toxicol. Appl. Pharmacol.* **228**(1): 76--83.
- Charles, I.O. and Adaku, N.A. (2010). Polycyclic aromatic hydrocarbons (PAHs) and benzene, toluene, ethylbenzene, and xylene (BTEX) contamination of soils in automobile mechanic workshops in Port-Harcourt Metropolis, Rivers State, Nigeria. *J. Am. Sci.* **6**(9): 242--246
- Chen, H., Yang, M. and Ye, S. (1992). A study on genotoxicity of cooking fumes from rapeseed oil. *Biomed. Environ. Sci.* **5**(3): 229--235.
- Chen, J.W., Wang, S.L., Hsieh, D.P.H., Yang, H.H. and Lee, H.L. (2012). Carcinogenic potencies of polycyclic aromatic hydrocarbons for back-door neighbours of restaurants with cooking emissions. *Sci. Total Environ.* **417--418**:68--75.

- Chen, Y.C., Chiu, C.P. and Chen, B.H. (2003). Mutagenicity of fumes from fried chicken legs. *J. Food Prot.* **66**: 1269–1276.
- Cherng, S.H., Huang, K.H., Yang, S.C., Wu, T.C., Yang, J.L. and Lee, H. (2002). Human 8-oxoguanine DNA glycosylase 1 mRNA expression as an oxidative stress exposure biomarker of cooking oil fumes. *J. Toxicol. Environ. Health A.* **65**(3-4): 265--278.
- Chiang, T.A., Wu, P.F., Ying, L.S., Wang, L.F. and Ko, Y.C. (1999a). Mutagenicity and aromatic amine content of fumes from heated cooking oils produced in Taiwan. *Food Chem. Toxicol.* **37**(2-3): 125--134.
- Chiang, T.A., Wu, P.F. and Ko, Y.C. (1999b). Identification of carcinogens in cooking oil fumes. *Environ. Res.* **81**(1): 18--22.
- Chiang, T.A., Wu, P.F., Wang, L.F., Lee, H., Lee, C.H. and Ko, Y.C. (1997). Mutagenicity and polycyclic aromatic hydrocarbon content of fumes from heated cooking oils produced in Taiwan. *Mutat. Res.* **381**(2): 157--161.
- Choe, E. and Min, D.B. (2007). Chemistry of deep-fat frying oils. *J. Food Sci.* **72**: R77–R86.
- Conney, A.H., Chang, R.L., Cui, X.X., Schiltz, M., Yagi, H., Jerina, D.M. and Wei, S.J. (2001). Dose-dependent differences in the profile of mutations induced by carcinogenic (R,S,S,R) bay- and fjord-region diol epoxides of polycyclic aromatic hydrocarbons. *Adv. Exp. Med. Biol.* **500**: 697--707.
- Cottet, V., Touvier, M., Fournier, A., Touillaud, M.S., Lafay, L., Clavel-Chapelon, F. and Boutron-Ruault, M.C. (2009). Postmenopausal breast cancer risk and dietary patterns in the E3N-EPIC prospective cohort study. *Am. J. Epidemiol.* **170**(10): 1257--1267.
- D’Avanzo, B., Negri, E., Gramenzi, A., Franceschi, S., Parazzini, F., Boyle, P., and La Vecchia, C. (1991). Fats in seasoning and breast cancer risk: An Italian case-control study. *Eur. J. Cancer* **27**: 420–423.
- Dai, Q., Shu, X.O., Jin, F., Gao, Y.T., Ruan, Z.X. and Zheng, W. (2002). Consumption of animal foods, cooking methods, and risk of breast cancer. *Cancer Epidemiol. Biomarkers Prev.* **11**(9): 801–808.
- Das, U.N. (2008). Essential fatty acids and their metabolites could function as endogenous HMG-CoA reductase and ACE enzyme inhibitors, antiarrhythmic, antihypertensive,

- antiatherosclerotic, anti-inflammatory, cytoprotective, and cardioprotective molecules. *Lipids Health Dis.* **7**: 37.
- Datta, K. and Biswas, J. (2009). Influence of dietary habits, physical activity and affluence factors on breast cancer in East India: A case-control study. *Asian Pac. J. Cancer Prev.* **10**(2): 219--222.
- De Marzo, A.M., Platz, E.A., Sutcliffe, S., Xu, J., Gronberg, H., Drake, C.G., Nakai, Y., Isaacs, W.B. and Nelson, W.G. (2007). Inflammation in prostate carcinogenesis. *Nat Rev Cancer* **7**(4): 256--269.
- De Stefani, E., Ronco, A., Mendilaharsu, M., Guidobono, M. and Deneo-Pellegrini, H. (1997). Meat intake, heterocyclic amines, and risk of breast cancer: A case-control study in Uruguay. *Cancer Epidemiol. Biomark. Prev.* **6**: 573--581.
- Dobargarnes, M.C. and Márquez-Ruiz, G. (1998). Regulation of used frying fats and validity of quick tests for discarding the fats. *Grasas Aceites* **49**(3--4): 331--335.
- Dost, K. and Ideli, C. (2012). Determination of polycyclic aromatic hydrocarbons in edible oils and barbecued food by HPLC/UV-Vis detection. *Food Chem.* **133**:193--199.
- Duan, X., Shen, G., Yang, H., Tian, J., Wei, F., Gong, J. and Zhang, J.J. (2016). Dietary intake polycyclic aromatic hydrocarbons (PAHs) and associated cancer risk in a cohort of Chinese urban adults: Inter- and intra-individual variability. *Chemosphere* **144**: 2469--2475.
- Dung, C.H., Wu, S.C. and Yen, G.C. (2006). Genotoxicity and oxidative stress of the mutagenic compounds formed in fumes of heated soybean oil, sunflower oil, and lard. *Toxicol. In Vitro* **20** (4): 439--447.
- Eder, K. and Kirchgessner, M. (1998). The effect of dietary vitamin E supply and a moderately oxidized oil on activities of hepatic lipogenic enzymes in rats. *Lipids.* **33**(3): 277--283.
- Elangovan, I., Thirugnanam, S., Chen, A., Zheng, G., Bosland, M.C., Kajdacsy-Balla, A. and Gnanasekar, M. (2012). Targeting receptor for advanced glycation end products (RAGE) expression induces apoptosis and inhibits prostate tumor growth. *Biochem. Biophys. Res. Commun.* **417**(4): 1133--1138.

- Endo, J. and Arita, M. (2016). Cardioprotective mechanism of omega-3 polyunsaturated fatty acids. *J. Cardiol.* **67**(1): 22--27.
- European Federation for the Science, Technology of Lipids. (2013). 7th international symposium on deep-fat frying. Optimum drying for safe and improved quality fried foods practical information for the foodservice and industrial frying [San Francisco].
- Fahy, E., Subramaniam, S., Murphy, R.C., Nishijima, M., Raetz, C.R., Shimizu, T., Spener, F., van Meer, G., Wakelam, M.J. and Dennis, E.A. (2009). Update of the LIPID MAPS comprehensive classification system for lipids. *J. Lipid Res.* **50**(Suppl): S9–S14.
- Falade, A.O., Oboh G., Ademiluyi A.O. and Odubango O.V. (2015). Consumption of thermally oxidized palm oil diets alters biochemical indices rats. *Beni Suef Univ J. Basic Appl. Sci.* **4**(2): 150–156.
- Falade, A.O., Oboh. G. and Okoh, A.L. (2017). Potential health implications of the consumption of thermally-oxidized cooking oils – a review. *Pol. J. Food Nutr. Sci.* **67**(2): 95–105.
- Falade, A.O. and Oboh, G. (2015). Thermal oxidation induces lipid peroxidation and changes in the physicochemical properties and β -carotene content of Arachis oil. *Int. J. Food Sci.* **2015**: 806524.
- Favero, A., Parpinel, M. and Franceschi, S. (1998). Diet and risk of breast cancer: Major findings from an Italian case-control study. *Biomed. Pharmacother.* **52**: 109–115.
- Foster, R., Williamson, C.S. and Lunn, J. (2009). BRIEFING PAPER: Culinary oils and their health effects. *Nutr. Bull.* **34**: 4–47
- Galeone, C., Pelucchi, C., Talamini, R., Levi, F., Bosetti, C., Negri, E., Franceschi, S. and La Vecchia, C. (2005). Role of fried foods and oral/pharyngeal and oesophageal cancers. *Br. J. Cancer.* **92**(11): 2065–2069.
- Galeone, C., Talamini, R., Levi, F., Pelucchi, C., Negri, E., Giacosa, A., Montella, M., Franceschi, S. and La Vecchia, C. (2007). Fried foods, olive oil and colorectal cancer. *Ann. Oncol.* **18**(1): 36--39
- Ghorbani, Z., Hekmatdoost, A., Zinab, H.E., Farrokhzad, S., Rahimi, R., Malekzadeh, R. and Pourshams, A. (2015). Dietary food groups intake and cooking methods associations with pancreatic cancer: A case-control study. *Indian J. Gastroenterol.* **34**(3): 225--232.

- Goburdhun, D., Jhaumeer-Laulloo, S. B. and Musruck, R. (2001). Evaluation of soybean oil quality during conventional frying by FTIR and some chemical indexes. *Int. J. Food Sci. Nutr.* **52**: 31–42.
- Gonzales, J.F., Barnard, N.D., Jenkins, D.J., Lanou, A.J., Davis, B., Saxe, G. and Levin, S. (2014). Applying the precautionary principle to nutrition and cancer. *J. Am. Coll. Nutr.* **33**: 239–246.
- Good, J. (2012). Healthiest cooking oil chart with smoke points, baseline of health foundation. The Baseline of Health Foundation. <https://jonbarron.org/diet-and-nutrition/healthiest-cooking-oil-chart-smoke-points>.
- Gotoh, N., AI-Iwasawa, Watanabe, H., Osato, R. and Wada, S. (2007). Oxidation of fats and oils in instant noodles stored under various conditions. *J. Food Lipids.* **14**: 350--365.
- Goya, L., Mateos, R. and Bravo, L. (2007). Effect of the olive oil phenol hydroxytyrosol on human hepatoma HepG2 cells. Protection against oxidative stress induced by tert-butylhydroperoxide. *Eur. J. Nutr.* **46**(2): 70--78.
- Grandgirard, A., Sebedio, J. L. and Fleury, J. (1984). Geometrical isomerization of linolenic acid during heat-treatment of vegetable-oils. *J. Am. Oil Chem. Soc.* **61**: 1563–1568.
- Guillén, M.D. and Uriarte. P.S. (2012). Aldehydes contained in edible oils of a very different nature after prolonged heating at frying temperature: Presence of toxic oxygenated α,β unsaturated aldehydes. *Food Chem.* **131**(3): 915–926.
- Gupta, M. K. (2005). Frying oil. In: Shahidi F., editor. *Edible Oil and Food Products: Products and Applications*. Hoboken, NJ, USA: John Wiley & Sons. pp. 1–31
- Gupta, R. C. and Lutz, W. K. (1999). Background DNA damage for endogenous and unavoidable exogenous carcinogens: A basis for spontaneous cancer incidence? *Mutat. Res.* **424**: 1–8.
- Hageman, G., Hermans, R., ten Hoor, F. and Kleinjans, J. (1990). Mutagenicity of deep-frying fat, and evaluation of urine mutagenicity after consumption of fried potatoes. *Food Chem. Toxicol.* **28**(2): 75--80.
- Hageman, G., Kikken, R., Ten Hoor, F. and Kleinjans, J. (1988). Assessment of mutagenic activity of repeatedly used deep-frying fats. *Mutat. Res.* **204**: 593–604.

- Hageman, G., Verhagen, H., Schutte, B. and Kleinjans, J. (1991). Biological effects of short-term feeding to rats of repeatedly used deep-frying fats in relation to fat mutagen content. *Food Chem. Toxicol.* **29**(10): 689--698.
- Hanf, V. and Gonder, U. (2005). Nutrition and primary prevention of breast cancer: Foods, nutrients and breast cancer risk. *Eur. J. Obstet. Gynecol. Reprod. Biol.* **123**(2):139-149.
- Hartung, R.W., Brune, H., Grimmer, G., Germann, P., Timm, J. and Wosniok, W. (1990). Evaluation of the carcinogenic potency of 4 environmental polycyclic aromatic compounds following intrapulmonary application in rats. *Exp. Pathol.* **40**: 221--227.
- Healy, D.A., Wallace, F.A., Miles, E.A., Calder, P.C. and Newsholm, P. (2000). Effect of low-to-moderate amounts of dietary fish oil on neutrophil lipid composition and function. *Lipids* **35**: 763--768.
- Henkler, F., Stolpmann, K. and Luch, A. (2012). Exposure to polycyclic aromatic hydrocarbons: Bulky DNA adducts and cellular responses. *EXS.* **101**: 107--131.
- Hori, H., Takayanagi, T., Kamada, Y., Shimoyoshi, S., Ono, Y., Kitagawa, Y., Shibata, H., Nagao, M., Fujii, W. and Sakakibara, Y. (2011). Genotoxicity evaluation of sesamin and episesamin. *Mutat Res.* **719**(1-2): 21--28.
- Houthuijzen, J.M. (2016). For better or worse: FFAR1 and FFAR4 signaling in cancer and diabetes. *Mol. Pharmacol.* **90**(6): 738--743.
- Huang, C., Zhang, X., Qiao, Z., Guan, L., Peng, S., Liu, J., Xie, R. and Zheng, L. (1992). A case-control study of dietary factors in patients with lung cancer. *Biomed. Environ. Sci.* **5**(3): 257--265.
- Hughes-Fulford, M., Li, C.F., Boonyaratanakornkit, J. and Sayyah, S. (2006). Arachidonic acid activates phosphatidylinositol 3-kinase signaling and induces gene expression in prostate cancer. *Cancer Res.* **66**(3): 1427--1433.
- Hung, H.S., Wu, W.J., Cheng, Y.W., Wu, T.C., Chang, K.L. and Lee, H. (2007). Association of cooking oil fumes exposure with lung cancer: Involvement of inhibitor of apoptosis proteins in cell survival and proliferation in vitro. *Mutat Res.* **628**(2):107-116.
- Ibáñez, R., Agudo, A., Berenguer, A., Jakszyn, P., Tormo, M.J., Sánchez, M.J., Quirós, J.R., Pera, G., Navarro, C., Martinez, C., Larrañaga, N., Dorronsoro, M., Chirlaque, M.D.,

- Barricarte, A., Ardanaz, E., Amiano, P. and González, C.A. (2005). Dietary intake of polycyclic aromatic hydrocarbons in a Spanish population. *J. Food Prot.* **68**: 2190–2195.
- Imaida, K., Lee, M.S., Wang, C.Y. and King, C.M. (1991). Carcinogenicity of dinitro pyrenes in the weanling female CD rat. *Carcinogenesis*, **12**: 1187–1191.
- Imaida, K., Uneyama, C., Ogasawara, H., Hayashi, S., Fukuhara, K., Miyata, N. and Takahashi, M. (1992). Induction of colon adenocarcinomas in CD rats and lung adenomas in ICR mice by 6-nitrochrysene: Comparison of carcinogenicity and aryl hydrocarbon hydroxylase induction in the target organs of each species. *Cancer Res.* **52**: 1542–1545.
- Isidori, M. and Parrella, A. (2009). Genotoxicity of aqueous extract from heated cooking oils and its suppression by Lactobacilli. *Food Sci. Technol. Int.* **15**(3): 267–273
- Jahren, A.H. and Schubert, B.A. (2010). Corn content of French fry oil from national chain vs. small business restaurants. *Proc. Natl. Acad. Sci. USA.* **107**(5): 2099–2101.
- Ji, B.T., Chow, W.H., Gridley, G., McLaughlin, J.K., Dai, Q., Wacholder, S., Hatch, M.C., Gao, Y.T. and Fraumeni, J.F. Jr. (1995). Dietary factors and the risk of pancreatic cancer: A case-control study in Shanghai China. *Cancer Epidemiol. Biomarkers Prev.* **4**(8): 885–893.
- Jørgensen, R.B., Strandberg, B., Sjaastad, A.K., Johansen, A. and Svendsen, K. (2013) Simulated restaurant cook exposure to emissions of PAHs, mutagenic aldehydes, and particles from frying bacon. *J. Occup. Environ. Hyg.* **10**:122–131.
- Joshi, A.D., Corral, R., Catsburg, C., Lewinger, J.P., Koo, J., John, E.M., Ingles, S.A. and Stern, M.C. (2012). Red meat and poultry, cooking practices, genetic susceptibility and risk of prostate cancer: Results from a multiethnic case-control study. *Carcinogenesis* **33**(11): 2108–2118
- Joshi, A.D., John, E.M., Koo, J., Ingles, S.A. and Stern, M.C. (2012). Fish intake, cooking practices, and risk of prostate cancer: Results from a multi-ethnic case-control study. *Cancer Causes Control* **23**(3): 405–420.
- Kasamatsu, T., Ogura, R., Ikeda, N., Morita, O., Saigo, K., Watabe, H., Saito, Y. and Suzuki, H. (2005). Genotoxicity studies on dietary diacylglycerol (DAG) oil. *Food Chem Toxicol.* **43**(2): 253–260.

- Katragadda, H.R., Fullana A.S., Sidhu S. and Carbonell-Barrachina, A.A. (2010). Emissions of volatile aldehydes from heated cooking oils. *Food Chem.* **120**: 59–65.
- Kelavkar, U.P., Hutzley, J., McHugh, K., Allen, K.G. and Parwani, A. (2009). Prostate tumor growth can be modulated by dietarily targeting the 15-lipoxygenase-1 and cyclooxygenase-2 enzymes. *Neoplasia* **11**(7): 692–699.
- Kim, D.J., Lee, K.K., Han, B.S., Ahn, B., Bae, J.H. and Jang, J.J. (1994). Biphasic modifying effect of indole-3-carbinol on diethylnitrosamine-induced preneoplastic glutathione S-transferase placental form-positive liver cell foci in Sprague-Dawley rats. *Jpn J Cancer Res.* **85**(6): 578–583.
- Kitamura, K., Shibata, R., Tsuji, Y., Shimano, M., Inden, Y. and Murohara T. (2011). Eicosapentaenoic acid prevents atrial fibrillation associated with heart failure in a rabbit model. *Am. J. Physiol. Heart Circ. Physiol.* **300**: H1814–H1821.
- Klein, V., Chajes, V., Germain, E., Schulgen, G., Pinault, M., Malvy, D., Lefrancq, T., Fignon, A., Le Floch, O., Lhuillery, C. and Bougnoux, P. Low α -linolenic acid content of adipose breast tissue is associated with an increased risk of breast cancer. *Eur. J Cancer* **36**: 335–340.
- Ko, Y.C., Cheng, L.S., Lee, C.H., Huang, J.J., Huang, M.S., Kao, E.L., Wang, H.Z. and Lin, H.J. (2000). Chinese food cooking and lung cancer in women nonsmokers. *Am. J. Epidemiol.* **151**(2): 140–147.
- Ko, Y.C., Lee, C.H., Chen, M.J., Huang, C.C., Chang, W.Y., Lin, H.J., Wang, H.Z. and Chang, P.Y. (1997). Risk factors for primary lung cancer among non-smoking women in Taiwan. *Int. J. Epidemiol.* **26**(1): 24–31.
- Kohlmeier, L., Simonsen, N., van't Veer, P., Strain, J. J., Martin-Moreno, J. M., Margolin, B., Huttunen, J. K., Fernandez-Crehuet, N. J., Martin, B. C., Thamm, M., Kardinaal, A. F. and Kok, F. J. (1997). Adipose tissue *trans* fatty acids and breast cancer in the European community multicenter study on antioxidants, myocardial infarction, and breast cancer. *Cancer Epidemiol. Biomark. Prev.* **6**: 705–710.

- Kojima, R., Okada, E., Ukawa, S., Mori, M., Wakai, K., Date, C., Iso, H. and Tamakoshi, A. (2017). Dietary patterns and breast cancer risk in a prospective Japanese study. *Breast Cancer* **24**(1): 152--160.
- Konings, E.J., Baars, A.J., van Klaveren, J.D., Spanjer, M.C., Rensen, P.M., Hiemstra, M., van Kooij, J.A. and Peters, P.W. (2003). Acrylamide exposure from foods of the Dutch population and an assessment of the consequent risks. *Food Chem. Toxicol.* **41**(11): 1569--1579.
- Korsh, J., Shen, A., Aliano, K. and Davenport, T. (2015). Polycyclic aromatic hydrocarbons and breast cancer: A review of the literature. *Breast Care (Basel)*. **10**(5): 316--318.
- Ku, S.K., Ruhaifi, M.S.M., Fatin, S.S., Saffana, M., Anna, K.T., Das, S. and Kamsiah, J. (2014). The harmful effects of consumption of repeatedly heated edible oils: A short review. *Clin. Ter.* **165**(4):217-222.
- Lapointe, A., Couillard, C. and Lemieux, S. (2006). Effects of dietary factors on oxidation of low-density lipoprotein particles. *J. Nutr. Biochem.* **17**(10): 645--658.
- Lee, C.H., Yang, S.F., Peng, C.Y., Li, R.N., Chen, Y.C., Chan, T.F., Tsai, E.M., Kuo, F.C., Huang, J.J., Tsai, H.T., Hung, Y.H., Huang, H.L., Tsai, S. and Wu, M.T. (2010). The precancerous effect of emitted cooking oil fumes on precursor lesions of cervical cancer. *Int. J. Cancer* **127**(4): 932--941.
- Lee, T. and Gany, F. (2013). Cooking oil fumes and lung cancer: A review of the literature in the context of the U.S. population. *J. Immigr. Minor Health.* **15**(3): 646--652.
- Leong, X.F., Salimon, J., Mustafa, M.R. and Jaarin, K. (2012). Effect of repeatedly heated palmolein on blood pressure-regulating enzymes activity and lipid peroxidation in rats. *Mal. J. Med. Sci.* **19**: 20--29.
- Leong, X.F., Najib, M.N.M., Das, S., Mustafa, M.R. and Jaarin, K. (2009). Intake of repeatedly heated palm oil causes elevation in blood pressure with impaired vasorelaxation in rats. *Tohoku J. Exp. Med.* **219**: 71--78.
- Li, C.T., Lin, Y.C., Lee, W.J. and Tsai, P.J. (2003) Emission of polycyclic aromatic hydrocarbons and their carcinogenic potencies from cooking sources to the urban atmosphere. *Environ. Health Perspect.* **111**: 483--487

- Li, J.W., Wang, X.G. and Jin, Q.Z. (2011). The sources, detection, and control of benzo[a]pyrene in edible oil. *China Grease* **36**: 7–11.
- Li, M.L., Lin, J., Hou, J.G., Xu, L., Cui, X.G., Xu, X.X., Yu, Y.W., Han, X., Wang, G.M., Guo, J.M., Xu, D.F., Thompson, T.C., Cao, G.W. and Zhang, H.W. (2014). Environmental and psycho-social factors related to prostate cancer risk in the Chinese population: A case–control study. *Biomed. Environ. Sci.* **27**:707–717
- Li, S., Wang, Y., Zhang, J. and Zhang, X. (1999). Study on the chemical components of edible oil fume in kitchen and its genotoxicity on *Drosophila*. *Wei Sheng Yan Jiu* **28**: 15–17.
- Li, S.G., Pan, D.H. and Wang, G.X. (1992). Analysis of polycyclic aromatic hydrocarbons in cooking oil smoke. *Chin. J. Pub. Health* **11**: 235–236.
- Lin, P.P., Yang, M.H., Liao, P.C., Wu, H.Y., Chang, L.W., Tsai, H.T. and Tyan, Y.C. (2008). Proteomic analysis of proteins associated with tt-DDE induced toxicity in BEAS-2B cells. *Biochem. Biophys. Res. Commun.* **376**(3): 519--524.
- Lipworth, L., Sonderman, J.S., Tarone, R.E. and McLaughlin, J.K. (2012). Review of epidemiologic studies of dietary acrylamide intake and the risk of cancer. *Eur. J. Cancer Prev.* **21**(4): 375–386.
- Liu, P., Holman, C.D., Jin, J. and Zhang, M. (2015). Diet and risk of adult leukemia: A multicenter case-control study in China. *Cancer Causes Control.* **26**(8): 1141--1151.
- Liu, Q., Sasco, A.J., Riboli, E. and Hu, M.X. (1993). Indoor air pollution and lung cancer in Guangzhou, People’s Republic of China. *Am. J. Epidemiol.* **137**(2): 145--154.
- Lodovici, M., Dolaro, P., Casalini, C., Ciappellano, S. and Testolin, G. (1995). Polycyclic aromatic hydrocarbon contamination in the Italian diet. *Food Addit Contam.* **12**: 703–713.
- Lopez-Abente, G., Sanz-Anquela, J.M. and Gonzalez, C.A. (2001). Consumption of wine stored in leather wine bottles and incidence of gastric cancer. *Arch. Environ. Health* **56**: 559–561.
- Lopez-Garcia, E., Schulze, M.B., Meigs, J.B., Manson, J.E., Rifai, N., Stampfer, M.J., Willett, W.C. and Hu, F.B. (2005). Consumption of trans- fatty acids is related to plasma biomarkers of inflammation and endothelial dysfunction. *J. Nutr.* **135**: 562–566.

- Lüchtenborg, M., Weijenberg, M.P., de Goeij, A.F., Wark, P.A., Brink, M., Roemen, G.M., Lentjes, M.H., de Bruïne, A.P., Goldbohm, R.A., van't Veer, P. and van den Brandt, P.A. (2005). Meat and fish consumption, APC gene mutations and hMLH1 expression in colon and rectal cancer: A prospective cohort study (The Netherlands). *Cancer Causes Control*. **16**(9): 1041--1054.
- lv, M.Y., Zhang, X., Ren, H.R., Liu, L., Zhao, Y.M., Wang, Z., Wu, Z.L., Liu, L.M. and Xu, H.J. (2016). A rapid method to authenticate vegetable oils through surface-enhanced Raman scattering. *Sci Rep*. **6**: 23405.
- Marinova, E.M., Seizova, K.A., Totseva, I.R., Panayotova, S.S., Marekov, I.N. and Momchilova, S.M. (2012). Oxidative changes in some vegetable oils during heating at frying temperature. *Bulg. Chem. Commun.* **44** (1): 57–63.
- Marsili, L., Caruso, A., Fossi, M.C., Zanardelli, M., Politi, E. and Focardi, S. (2001) Polycyclic aromatic hydrocarbons (PAHs) in subcutaneous biopsies of Mediterranean cetaceans. *Chemosphere* **44**: 147–154.
- Mensah, E. and Obeng, G.Y. (2013). Assessment and projection of waste vegetable oils from hotels for biodiesel feedstock and CO₂ savings in Ghana. *Int J Engg Res Sci Technol*. **2**(4): 1--12.
- Mojska, H., Gielecińska, I., Szponar, L. and Oltarzewski, M. (2010). Estimation of the dietary acrylamide exposure of the Polish population. *Food Chem. Toxicol.* **48**(8-9): 2090--2096.
- Moreira, R.G., Castell-Perez, M.E. and Barrufet, M.A. (1999). Deep-fat frying: Fundamentals and applications. Springer US, Springer-Verlag US, pp.350. ISBN 978-0-8342-1321-0.
- Mucci, L.A., Adami, H.O. and Wolk, A. (2006). Prospective study of dietary acrylamide and risk of colorectal cancer among women. *Int. J. Cancer* **118**: 169–173.
- Nakagawa, K., Hidaka, T., Kitano, M., Asakura, M., Kamigaito, T., Noguchi, T. and Hosoe, K. (2008). Genotoxicity studies on licorice flavonoid oil (LFO). *Food Chem. Toxicol.* **46**(7): 2525--2532.
- Navarro, A., Munoz, S.E., Lantieri, M.J., del Pilar Diaz, M., Cristaldo, P.E., de Fabro, S.P. and Eynard, A.R. (2004). Meat cooking habits and risk of colorectal cancer in Cordoba, Argentina. *Nutrition* **20**: 873–877.

- Naz, S., Siddiqi, R., Sheikh, H. and Sayeed, S.A. (2005). Deterioration of olive, corn and soybean oils due to air, light, heat, and deep-frying. *Food Res. Int.* **38**: 127–134.
- Neuhouser, M.L., Barnett, M.J., Kristal, A.R., Ambrosone, C.B., King, I., Thornquist, M. and Goodman, G. (2007). (n-6) PUFA increase and dairy foods decrease prostate cancer risk in heavy smokers. *J. Nutr.* **137**(7): 1821–1827.
- Ng, C.Y., Leong, X.F., Masbah, N., Adam, S.K., Kamisah, Y. and Jaarin, K. (2014a). Heated vegetable oils and cardiovascular disease risk factors, *Vasc. Pharmacol.* **61**: 1–9.
- Ng, C.Y., Leong, X.F., Masbah, N., Adam, S.K., Kamisah, Y. and Jaarin, K. (2014b). Reprint of heated vegetable oils and cardiovascular disease risk factors, *Vasc. Pharmacol.* **62**: 38–46.
- Ng, C.Y., Kamisah, Y., Othman, F. and Jaarin, K. (2012). The role of repeatedly heated soybean oil in the development of hypertension in rats: Association with vascular inflammation. *Int. J. Exp. Pathol.* **93**: 377–387.
- Nie, J., Shi, J., Duan, X., Wang, B., Huang, N. and Zhao, X. (2014). Health risk assessment of dietary exposure to polycyclic aromatic hydrocarbons in Taiyuan, China. *J. Environ. Sci. (China)*. **26**(2): 432–439.
- Obboh, G., Falade, A.O. and Ademiluyi, A.O. (2014). Effect of thermal oxidation on the physicochemical properties, malondialdehyde and carotenoid contents of palm oil. *Riv. Ital. Sostanze Gr.* **91**(1): 59–65
- Ohba, S., Nishi, M. and Miyake, H. (1996). Eating habits and pancreas cancer. *Int J Pancreatol.* **20**(1): 37–42.
- Orozco-Solano, M.I., Priego-Capote, F. and Luque de Castro, M.D. (2013). Analysis of esterified and nonesterified fatty acids in serum from obese individuals after intake of breakfasts prepared with oils heated at frying temperature. *Anal. Bioanal. Chem.* **405**(18): 6117–6129.
- Owu, D.U., Orie, N.N. and Osim, E.E. (1997). Altered responses of isolated aortic smooth muscle following chronic ingestion of palm oil diets in rats. *Afr. J. Med. Med. Sci.* **26**: 83–86.

- Pala, V., Krogh, V., Muti, P., Chajes, V., Riboli, E., Micheli, A., Saadatian, M., Sieri, S. and Berrino, F. (2001). Erythrocyte membrane fatty acids and subsequent breast cancer: A prospective Italian study. *J. Natl. Cancer Inst.* **93**: 1088–1095.
- Pandey, M.K., Dhawan, A. and Das, M. (2006). Induction of p53, p21waf1, ornithine decarboxylase activity, and DNA damage leading to cell-cycle arrest and apoptosis following topical application of repeated fish fried oil extract to mice. *Mol. Carcinog.* **45**: 805–813.
- Pandit, G.G., Srivastava, P.K. and Mohan Rao, A.M. (2001). Monitoring of indoor volatile organic compounds and polycyclic aromatic hydrocarbons arising from kerosene cooking fuel. *Sci. Total Environ.* **279**:159–165
- Pauk, N., Klimesova, S., Kara, J., Topinka, J. and Labaj, J. (2013). The relevance of monitoring of antibodies against the polycyclic aromatic hydrocarbon (PAH) and PAH-DNA adducts in serum in relation to lung cancer and chronic obstructive pulmonary disease (COPD). *Neoplasma* **60**(2): 182--187.
- Paul, S. and Mittal, G.S. (1997). Regulating the use of degraded oil/fat in deep-fat/oil food frying. *Crit. Rev. Food Sci. Nutr.* **37**(7): 635–662.
- Pawlega, J., Rachtan, J. and Dyba, T. (1996). Dietary factors and risk of prostate cancer in Poland. Results of case–control study. *Neoplasma* **43**:61–63.
- Pellizzari, E.D., Michael, L.C., Thomas, K.W., Shields, PG. and Harris, C. (1995). Identification of 1,3-butadiene, benzene, and other volatile organics from wok oil emissions. *J. Expo. Anal. Environ. Epidemiol.* **5**(1): 77--87.
- Pfau, W. and Marquardt, H. (2001). Cell transformation in vitro by food-derived heterocyclic amines Trp-P-1, Trp-P-2 and N(2)-OH-PhIP. *Toxicol.* **166**: 25–30.
- Pham, T.M., Fujino, Y., Kikuchi, S., Tamakoshi, A., Matsuda, S. and Yoshimura, T. (2010). Dietary patterns and risk of stomach cancer mortality: The Japan collaborative cohort study. *Ann. Epidemiol.* **20**(5): 356--363.
- Phiri, G., Mumba, P. and Mangwera, A. (2006). The quality of cooking oil used in informal food processing in Malawi: A preliminary study. *Int. J. Consum. Stud.* **30**(6): 527–532.

- Platt, K.L., Pfeiffer, E., Petrovic, P., Friesel, H., Beermann, D., Hecker, E. and Oesch, F. (1990). Comparative tumorigenicity of picene and dibenz[a,h]anthracene in the mouse. *Carcinogenesis*, **11**: 1721–1726.
- Psaltopoulou, T, Naska, A., Orfanos, P., Trichopoulos, D., Mountokalakis, T. and Trichopoulou, A. (2004). Olive oil, the Mediterranean diet, and arterial blood pressure: The Greek European Prospective Investigation into Cancer and Nutrition(EPIC) study. *Am. J. Clin. Nutr.* **80**:1012–1018.
- Pu, Y., Yin, L., Yuan, H. and Liang, G. (2002). Oxidative stress of cooking oil fume on rat type II lung cells. *Wei Sheng Yan Jiu.* **31**(2):85-87.
- Pulliero, A., Godschalk, R., Andreassi, M.G., Curfs, D., Van Schooten, F.J. and Izzotti, A. (2015). Environmental carcinogens and mutational pathways in atherosclerosis. *Int. J. Hyg. Environ. Health.* **218**(3): 293--312.
- Purcaro, G., Navas, J.A., Guardiola, F., Conte, L.S. and Moret, S. (2006). Polycyclic aromatic hydrocarbons in frying oils and snacks. *J. Food Prot.* **69**(1): 199--204.
- Qu, Y.H., Xu, G.X., Zhou, J.Z., Chen, T.D., Zhu, L.F., Shields, P.G., Wang, H.W. and Gao, Y.T. (1992). Genotoxicity of heated cooking oil vapors. *Mutat. Res.* **298**(2):105-111.
- Ramesh, S.A., Walker, D.B., Hood, M.D, Guillén, M.D., Schneider, K. and Weyand, E.H. (2004). Bioavailability and risk assessment of orally ingested polycyclic aromatic hydrocarbons. *Int. J. Toxicol.* **23**: 301–333.
- Ramesh, A., Walker, S.A., Hood, D.B., Guillén, M. D., Schneider, K. and Weyand, E. H. (2004). Bioavailability and risk assessment of orally ingested polycyclic aromatic hydrocarbons. *Int. J. Toxicol.* **23**(5): 301–333.
- Rojas, M., Godschalk, R., Alexandrov, K., Cascorbi, I., Kriek, E., Ostertag, J., Van Schooten, F.J. and Bartsch, H. (2001). Myeloperoxidase--463A variant reduces benzo[a]pyrene diol epoxide DNA adducts in skin of coal tar treated patients. *Carcinogenesis* **22**(7):1015-1018.
- Romero, A., Bastida, S. and Sanchez-Muniz, F.J. (2006). Cyclic fatty acid monomer formation in domestic frying of frozen foods in sunflower oil and high oleic acid sunflower oil without oil replenishment. *Food Chem. Toxicol.* **44**: 1674--1681.

- Ross, J., Nelson, G., Erexson, G., Kligerman, A., Earley, K., Gupta, R.C. and Nesnow, S. (1991). DNA adducts in rat lung, liver and peripheral blood lymphocytes produced by i.p. administration of benzo[a]pyrene metabolites and derivatives. *Carcinogenesis*, **12**: 1953–1955.
- Ross, J., Nelson, G., Kligerman, A., Erexson, G., Bryant, M., Earley, K., Gupta, R. and Nesnow, S. (1990). Formation and persistence of novel benzo(a)pyrene adducts in rat lung, liver, and peripheral blood lymphocyte DNA. *Cancer Research*, **50**: 5088–5094.
- Rossell, J.B. (1998). Industrial Frying Process. *Grasas y Aceites* **49**(3-4): 282--295.
- Sadoudi, R., Ammouche, A. and Ali, A.D. (2014). Thermal oxidative alteration of sunflower oil. *Afr. J. Food Sci.* **8**(3): 116–121
- Sayon-Orea, C., Carlos, S. and Martínez-Gonzalez, M.A. (2015). Does cooking with vegetable oils increase the risk of chronic diseases?: A systematic review. *Br. J. Nutr.* **113**(Suppl 2): S36-48.
- See, S.W. and Balasubramanian, R. (2008). Chemical characteristics of fine particles emitted from different gas cooking methods. *Atmos. Environ.* **42**: 8852–8862
- Seow, A., Poh, W.T., Teh, M, Eng, P., Wang, Y.T., Tan, W.C., Yu, M.C. and Lee, H.P. (2000). Fumes from meat cooking and lung cancer risk in Chinese women. *Cancer Epidemiol. Biomarkers Prev.* **9**(11): 1215--1221.
- Shukla, Y., Arora, A. and Taneja, P. (2003). Antigenotoxic potential of certain dietary constituents. *Teratog. Carcinog. Mutagen* **1**: 323–335.
- Shukla, Y. and Arora, A. (2003). Enhancing effects of mustard oil on preneoplastic hepatic foci development in Wistar rats. *Hum. Exp. Toxicol.* **22**(2): 51--55.
- Shukla, Y., Kalra, N., Katiyar, S., Siddiqui, I.A. and Arora, A. (2004). Chemopreventive effect of indole-3-carbinol on induction of preneoplastic altered hepatic foci. *Nutr. Cancer.* **50**(2): 214--220.
- Sinha, R., Chow, W.H., Kulldorff, M., Denobile, J., Butler, J., Garcia-Closas, M., Weil, R., Hoover, R.N. and Rothman, N. (1999). Well-done, grilled red meat increases the risk of colorectal adenomas. *Cancer Res.* **59**: 4320–4324.

- Sinha, R., Kulldorff, M., Chow, W.H., Denobile, J. and Rothman, N. (2001). Dietary intake of heterocyclic amines, meat derived mutagenic activity, and risk of colorectal adenomas. *Cancer Epidemiol. Biomarkers Prev.* **10**: 559–562.
- Sinha, R., Gustafson, D. R., Kulldorff, M., Wen, W. Q., Cerhan, J. R. and Zheng, W. (2000). 2-Amino-1-methyl-6-phenylimidazo[4,5-*b*]pyridine, a carcinogen in high-temperature-cooked meat, and breast cancer risk. *J. Natl. Cancer Inst.* **92**: 1352–1354.
- Siti, K.A., Srijit, D., Ima, N.S., Nor, A.U. and Kamsiah, J. (2008). Consumption of repeatedly heated soy oil increases the serum parameter related to atherosclerosis in ovariectomized rats. *Tohoku J. Exp. Med.* **215**: 219–226
- Somi, M.H., Mousavi, S.M., Naghashi, S., Faramarzi, E., Jafarabadi, M.A., Ghojzade, M., Majidi, A. and Naseri Alavi, S.A. (2015). Is there any relationship between food habits in the last two decades and gastric cancer in North-Western Iran? *Asian Pac. J. Cancer Prev.* **16**(1): 283–290.
- Soriguer, F., Rojo-Martinez, G., Dobarganes, M.C., García- Almeida, J.M., Esteva, I., Beltrán, M., Ruiz De Adana, M.S., Tinahones, F., Gómez-Zumaquero, J.M., García-Fuentes, E. and González-Romero, S. (2003). Hypertension is related to the degradation of dietary frying oils. *Am. J. Clin. Nutr.* **78**: 1092–1097.
- Srivastava, S., Singh, M., George, J., Bhui, K., Murari Saxena, A. and Shukla, Y. (2010a). Genotoxic and carcinogenic risks associated with the dietary consumption of repeatedly heated coconut oil. *Br. J. Nutr.* **104** (9):1343–1352.
- Srivastava, S., Singh, M., George, J., Bhui, K. and Shukla, Y. (2010b). Genotoxic and carcinogenic risks associated with the consumption of repeatedly boiled sunflower oil. *J. Agric. Food Chem.* **58** (20):11179–11186.
- Stott-Miller, M., Neuhouser, M.L. and Stanford, J.L. (2013). Consumption of deep-fried foods and risk of prostate cancer. *Prostate* **73**(9): 960--969.
- Svensson, K., Abramsson, L., Becker, W., Glynn, A., Hellenäs, K.E, Lind, Y. and Rosén, J. (2003). Dietary intake of acrylamide in Sweden. *Food Chem. Toxicol.* **41**(11):1581-1586.
- Takamatsu, S., Matsui, O., Gabata, T., Kobayashi, S., Okuda, M., Ougi, T., Ikehata, Y., Nagano, I. and Nagae, H. (2008). Selective induction hyperthermia following transcatheter arterial

- embolization with a mixture of nano-sized magnetic particles (ferucarbotran) and embolic materials: Feasibility study in rabbits. *Radiat. Med.* **26**(4): 179--187.
- Takayama, S., Monma, Y., Tsubota-Utsugi, M., Nagase, S., Tsubono, Y., Numata, T., Toyoshima, M., Utsunomiya, H., Sugawara, J. and Yaegashi, N. (2013). Food intake and the risk of endometrial endometrioid adenocarcinoma in Japanese women. *Nutr Cancer* **65**(7): 954--960.
- Tareke, E., Rydberg, P., Karlsson, P., Eriksson, S. and Tornqvist, M. (2002). Analysis of acrylamide, a carcinogen formed in heated foodstuffs. *J. Agric. Food Chem.* **50**(17): 4998--5006
- Topinka, J., Milcova, A., Schmuczerova, J., Mazac, M., Pechout, M. and Vojtisek-Lom, M. (2012). Genotoxic potential of organic extracts from particle emissions of diesel and rapeseed oil powered engines. *Toxicol. Lett.* **212**(1): 11--17.
- Totani, N., Ohno, C. and Yamaguchi A. (2006). Is the frying oil in deep-fried foods safe? *J. Oleo. Sci.* **55**(9): 449--456.
- Tung, Y.H., Ko, J.L., Liang, Y.F., Yin, L., Pu, Y. and Lin, P. (2001). Cooking oil fume-induced cytokine expression and oxidative stress in human lung epithelial cells. *Environ. Res.* **87**(1): 47--54.
- Uribarri, J., Cai, W., Peppia, M., Goodman, S., Ferrucci, L., Striker, G. and Vlassara, H. (2007). Circulating glycotoxins and dietary advanced glycation endproducts: Two links to inflammatory response, oxidative stress, and aging. *J. Gerontol. Biol. Sci. Med. Sci.* **62**(4): 427--433.
- Uribarri, J., Woodruff, S., Goodman, S., Cai, W., Chen, X., Pyzik, R., Yong, A., Striker, G.E. and Vlassara, H. (2010). Advanced glycation end products in foods and a practical guide to their reduction in the diet. *J. Am. Diet Assoc.* **110**(6): 911--16.e12.
- Vaskova, H. and Buckova, M. (2015). Thermal degradation of vegetable oils: Spectroscopic measurement and analysis. *Procedia Eng.* **100**: 630--635.
- Velema, J.P., Ferrera, A., Figueroa, M., Bulnes, R., Toro, L.A., de Barahona, O., Claros, J.M. and Melchers, W.J. (2002). Burning wood in the kitchen increases the risk of cervical neoplasia in HPV-infected women in Honduras. *Int J Cancer* **97**(4): 536--541.

- Wakabayashi, K. (1990). International commission for protection against environmental mutagens and carcinogens. ICPEMC Working Paper 7/1/3. Animal studies suggesting involvement of mutagen/carcinogen exposure in atherosclerosis. *Mutat. Res.* **239**: 181–187.
- Williams, G.M., Aardema, M.J., Long, P.H., Thompson, E.D. and Allgood, G.S. (1996). Genotoxicity and subchronic toxicity studies with heated olestra. *Food Chem Toxicol.* **34**(10): 941–950.
- Wornat, M.J., Ledesma, E.B., Sandrowitz, A.K., Roth, M.J., Dawsey, S.M., Qiao, Y.L. and Chen, W. (2001). Polycyclic aromatic hydrocarbons identified in soot extracts from domestic coal-burning stoves of Henan Province, China. *Environ. Sci. Technol.* **35**:1943–1952
- Wu, P.F., Chiang, T.A., Wang, L.F., Chang, C.S. and Ko, Y.C. (1998). Nitro-polycyclic aromatic hydrocarbon contents of fumes from heated cooking oils and prevention of mutagenicity by catechin. *Mutat. Res.* **403**(1-2): 29--34.
- Wu, S.C. and Yen, G.C. (2004). Effects of cooking oil fumes on the genotoxicity and oxidative stress in human lung carcinoma (A-549) cells. *Toxicol In Vitro* **18**: 571–580.
- Wu, S.C., Yen, G.C. and Sheu, F. (2001). Mutagenicity and identification of mutagenic compounds of fumes obtained from heating peanut oil. *J. Food Prot.* **64**(2): 240--245.
- Yao, Z., Li, J., Wu, B., Hao, X., Yin, Y. and Jiang, X. (2015). Characteristics of PAHs from deep-frying and frying cooking fumes. *Environ. Sci. Pollut. Res. Int.* **22**(20): 16110--16120.
- Young, S.C., Chang, L.W., Lee, H.L., Tsai, L.H., Liu, Y.C. and Lin, P. (2010). DNA damages induced by trans, trans-2,4-decadienal (tt-DDE), a component of cooking oil fume, in human bronchial epithelial cells. *Environ. Mol. Mutagen.* **51**(4): 315--321.
- Zhang, H., Wang, G. and Tan, W. (2002). Study on the effects of cooking oil fume condensate on the DNA integrity. *Wei Sheng Yan Jiu.* **31**(4): 238--240.
- Zhang, L., Lv, J. and Liao, C. (2012). Dietary exposure estimates of 14 trace elements in Xuanwei and Fuyuan, two high lung cancer incidence areas in China. *Biol. Trace Elem. Res.* **146**(3): 287--292.

- Zhao, Y., Wang, S.X., Aunan, K., Seip, H.M. and Hao, J.M. (2006). Air pollution and lung cancer risks in China—a meta-analysis. *Sci. Total Environ.* **366**: 500–513.
- Zheng, W., Blot, W.J., Shu, X.O., Diamond, E.L., Gao, Y.T., Ji, B.T. and Fraumeni, J.F. Jr. (1992). Risk factors for oral and pharyngeal cancer in Shanghai, with emphasis on diet. *Cancer Epidemiol. Biomarkers Prev.* **1**(6): 441–448.
- Zheng, W., Gustafson, D. R., Sinha, R., Cerhan, J. R., Moore, D., Hong, C. P., Anderson, K. E., Kushi, L. H., Sellers, T. A., and Folsom, A. R. (1998). Well-done meat intake and the risk of breast cancer. *J. Natl. Cancer Inst.* **90**: 1724–1729.
- Zhong, L., Goldberg, M.S., Gao, Y.T. and Jin, F. (1999a). Lung cancer and indoor air pollution arising from Chinese-style cooking among nonsmoking women living in Shanghai, China. *Epidemiol.* **10**(5): 488–494.
- Zhong, L., Goldberg, M.S., Parent, M.E. and Hanley, J.A. (1999b). Risk of developing lung cancer in relation to exposure to fumes from Chinese-style cooking. *Scand. J. Work Environ. Health.* **25**(4): 309–316.
- Zock, P. L. and Katan, M. B. (1998). Linoleic acid intake and cancer risk: A review and meta-analysis. *Am. J. Clin. Nutr.* **68**: 142–153.

Table 1: Physico-chemical changes during thermal oxidation of various cooking oils.

Vegetable oil	Temperature (°C)	Duration of heating	Physico-chemical changes	References
Canola oil	185 & 215	7 h/day	1. Decrease total polar compounds and anisidine value.	Aladedunye and Przybylski, 2009
			2. Increase vitamin E degradation	
Coconut, safflower, canola & olive oil	180, 210, 240, 270	6 h	Formation of acrolein increased with temperature.	Katragadda et al., 2010
Olive oil	180	1.5 – 25 h	1. Decrease the quantity of hydroxytyrosol & tyrosol like substances.	Brenes et al., 2002
			2. Degradation of vitamin E and glyceridic fractions.	
			3. Losses in polyphenols.	
	180	30–180 min	Decrease concentration of hydroxytyrosol, elenolic acid, decarboxymethyl oleuropein aglycon, and oleuropein aglycon	Carrasco-Pancorbo et al., 2007
Olive, corn, soybean oils	180	30, 60, 90 min	Increase peroxide value, <i>p</i> -anisidine value, FFA	Naz et al., 2005
Palm oil and soybean oil	180	Heated once & 5 times (10 mins)	Decrease vitamin E content and various isomers of vitamin E	Adam et al., 2007
Palm oil	30–320	0–20 min	Increase MDA content, decrease carotenoid content	Oboh et al., 2014
Peanut oil	220	20 min	1. Increase acid, peroxide value, MDA content.	Falade and Oboh, 2015
			2. Decrease of total carotenoid content	
Sunflower oil	100	52 h	Decrease amount of linoleic acid	Sadoudi et al., 2014
Sunflower, grapeseed, soybean, corn & olive oil	180	50 h	Increase quantity of conjugated trienes and total polar components.	Marinova et al., 2012

Table 2: *In-vitro* and *in vivo* genotoxicity and mutagenicity effect of RCO

Model	Deep fried oil	Negative control	Investigation	Results	References
Wistar albino rats and Salmonella tester strain TA97, TA100	coconut oil and palm oil	mutagenic to strain TA100	TBARS, hydroperoxides, mutagenicity of urine and faeces, alkaline phosphatase, cell proliferative indices in the gastro-intestinal tract	Genotoxicity and mutagenicity effect	Hageman et al., 1991
Human lung carcinoma pulmonary type II-like epithelium cell (A-549 cell)	soybean oil, sunflower oil, and lard	trans-trans-2,4-decadienal, t-t-2,4-DDE; trans-trans-2,4-nonadienal, t-t-2,4-NDE; trans-2-decenal, t-2-DCA and trans-2-undecenal, t-2-UDA	GSH, GST, GSR and ROS assay	Genotoxicity and mutagenicity effect	Dung et al., 2006
Wistar albino rats and Salmonella tester strain TA97, TA100	coconut oil and palm oil	mutagenic to strain TA100	TBARS, mutagenicity of urine and faeces, cell proliferative indices in the gastro-intestinal tract	Genotoxicity and mutagenicity effect	Hageman et al., 1990
Wistar albino rats	soybean oil and coconut oil	-	fatty acid synthase, acetyl-CoA-carboxylase, glucose-6-phosphate dehydrogenase, 6-phosphogluconate dehydrogenase, and ATP citrate lyase, TGL, TBARS,	Genotoxicity and mutagenicity effect	Eder and Kirchgessner, 1998
Salmonella tester strain TA97, TA100	Peanut oil	trans-trans-2,4-decadienal, trans-trans-2,4-nonadienal, trans-2-decenal, trans-2-undecenal	smoke point, unsaturated fatty acids, fume formation, and assay of mutagenicity effect	mutagenicity effect	Wu et al., 2001
In vitro	Lard, soybean oil, peanut oil	polycyclic aromatic hydrocarbons, (benzo(a)pyrene, benz(a)anthracene, and dibenz(a,h)anthracene	Assay of mutagenicity effect	mutagenicity effect	Wu et al., 1998
Human and Salmonella tester strain TA97	Lard, soybean oil, peanut oil	dibenz[a,h]anthracene and benz[a]anthracene	Assay of mutagenicity effect	mutagenicity effect	Chiang et al., 1997
Human	Safflower, olive, coconut, mustard, and corn	enzo[a]pyrene, dibenz[a,h]anthracene, benzo[b]fluoranthene and benzo[a]anthracene	HPLC, GC-MS assay, Ames test, sister chromatid exchange, and SOS chromotest	Genotoxicity	Chiang et al., 1999b
Wistar albino rats	Coconut oil	diethylnitrosamine	SOD, CAT, GGT, GST	Genotoxicity and mutagenicity effect	Srivastava et al., 2010a
Wistar albino rats	Sunflower oil	polycyclic aromatic hydrocarbons	SOD, CAT, GGT, GST. ATP, ALP glucose-6-phosphatase	Genotoxicity and mutagenicity effect	Srivastava et al., 2010b
Wistar albino rats	Soybean oil	cooking oil fume	MTT reduction assay, type II pneumocytes were studied by modified alkaline single-cell gel using a electrophoresis assay (comet	DNA damage and genotoxicity effect	Zhang et al., 2002

			assay),		
Mice	Rapeseed oil, Peanut oil, and lard	cooking oil fume	Salmonella mutation assay, SV50 forward mutation assay, and sister chromatid exchange assay, as well as the micronucleus assay in mouse bone marrow	Mutagenicity effect	Qu et al., 1992
Salmonella typhimurium tester strains TA98 and TA104	Chinese rapeseed, refined Canola, soybean, and peanut oil	cooking oil fume	Salmonella mutation assay	Mutagenicity effect	Shields et al., 1995
Human	Peanut, soybean, and Canola oils	cooking oil fume	Determination of 1,3-butadiene, benzene, and a series of aldehydes, olefins, and saturated hydrocarbons	Genotoxicity effect	Pellizzari et al., 1995
Wistar albino rats, hamster, mice	Canola, soybean, and peanut oil	cooking oil fume	Salmonella mutagenesis (Ames test); forward mutagenesis of mouse lymphoma cells at the thymidine kinase locus; unscheduled DNA synthesis in rat hepatocytes; and clastogenicity in cultured Chinese hamster ovary cells	Genotoxicity effect	Williams et al., 1996
Salmonella typhimurium tester strains TA98 and TA104; Chinese hamster lung cells (CHL/IU)	Canola, soybean, and peanut oil	cooking oil fume	Bacterial reverse mutation assay (Ames test), the chromosomal aberration assay in cultured Chinese hamster lung cells and a bone marrow micronucleus assay	Genotoxicity effect	Kasamatsu et al., 2005
Sprague-Dawley rats and ICR mice	Sesame seeds and oil	cooking oil fume	Bacterial reverse mutation assay (Ames test), the chromosomal aberration assay in cultured Chinese hamster lung cells and a bone marrow micronucleus assay	Genotoxicity effect	Hori et al., 2011
Sprague-Dawley rats and ICR mice	Licorice flavonoid oil	cooking oil fume	Bacterial reverse mutation assay (Ames test), the chromosomal aberration assay in cultured Chinese hamster lung cells and a bone marrow micronucleus assay	Genotoxicity effect	Nakagawa et al., 2008
Sprague-Dawley rats	Rapeseed oil	cooking oil fume	Rat liver microsomal S9 fraction	Genotoxicity effect	Topinka et al., 2012

Table 3: Impact of RCO on tumourigenic effect

Model	Deep fried oil	Negative control	Investigation	Results	References
Wistar albino rats	Mustard oil	Diethylnitrosamine	GST, GGT, liver weight	Tumourigenic effect	Shukla and Arora, 2003
Wistar albino rats	Mustard oil	Diethylnitrosamine	GST, GGT, liver weight alkaline phosphatase, adenosine triphosphatase, glucose-6-phosphatase	Tumourigenic effect	Shukla and Arora, 2003
Wistar albino rats	Mustard oil	Diethylnitrosamine	GST, GGT, liver weight ALP, adenosine triphosphatase, glucose-6-phosphatase	Tumourigenic effect	Kim et al., 1994; Shukla et al., 2004
Rabbit	soybean, and peanut oil	Heated lipiodol	Vascular endothelial growth factor receptor (VEGFR) and vascular endothelial growth factor (VEGF) expression levels, Proliferating cell nuclear antigen (PCNA) expression	Enhance tumor angiogenesis	Cao et al., 2013
Rabbit	Canola oil	Ferucarbotran and lipiodol	Apoptosis assay by TUNEL staining	Enhance tumor angiogenesis	Takamatsu et al., 2008
Rabbit	Canola, soybean, and peanut oil	Heated doxorubicin-lipiodol	serum AST levels	Enhance tumor angiogenesis	Cao et al., 2010
Human bronchial epithelial cells (BEAS-2B cells)	Peanut oil	Trans, trans-2,4-decadienal	ROS production, GSH/GSSG ratio, cell proliferation, and expression of TNF α and IL-1 β	Tumourigenic and genotoxicity effect	Chang et al., 2005
Human bronchial epithelial cells (BEAS-2B cells)	Soybean and peanut oil	Trans, trans-2,4-decadienal	Vitamin C and N-acetylcysteine, downstream targets of p27, including CDK4, cyclin D1 and phosphorylated-Rb proteins	Tumourigenic and genotoxicity effect	Chang and Lin, 2008
Human bronchial epithelial cells (BEAS-2B cells)	Canola oil	Trans, trans-2,4-decadienal	endonuclease III/formamidopyrimidine-DNA glycosylase, nucleotide excision repair enzymes	Tumourigenic and genotoxicity effect	Young et al., 2010
Human bronchial epithelial cells (BEAS-2B cells)	Canola, soybean, and peanut oil	Trans, trans-2,4-decadienal	Oxidative stress markers	Tumourigenic and genotoxicity effect	Lin et al., 2008

Table 4: Impact of RCO on various cancers

Model	Deep fried oil	Negative control	Investigation	Results	References
Human lung carcinoma pulmonary type II-like epithelium cell (A-549 cell)	Soybean oil, sunflower seed oil, and lard	Trans-trans-2,4-decadienal, t-t-2,4-DDE; trans-trans-2,4-nonadienal, t-t-2,4-NDE; trans-2-decenal, t-2-DCA and trans-2-undecenal, t-2-UDA	GSH, GST, GSR and ROS assay	Women exposed to emitted fumes from cooking oil cause at higher risk of lung cancer	Dung et al., 2006
Human lung carcinoma pulmonary type II-like epithelium cell (A-549 cell)	Soybean oil and sunflower oil	Trans-trans-2,4-decadienal, t-t-2,4-DDE	GSH, GST, GSR, ROS	Fumes from cooking oil cause at higher risk of lung cancer	Wu and Yen, 2004
human hepatoma HepG2 cells	Olive oil	Tertbutyl hydroperoxide	GSH, GST, GPx, MDA	Liver cancer	Alfa et al., 2006; Goya et al., 2007
Human	Sunflower oil, lord oil	Salmonella tester strain TA97	HPLC, GC-MS, and assay of 2-naphthylamine and 4-aminobiphenyl	Bladder cancer	Chiang et al., 1999a
Human	Vegetable oils	Cooking oil fumes	Analysis of DNA damage	Lung cancer	Lee and Gany, 2013
Human	Rapeseed oil	Cooking oil fumes	Exposure to indoor air pollutants from Chinese-style cooking was ascertained through in-person interviews	Lung cancer	Zhong et al., 1999a
Human	Vegetable oils	Cooking oil fumes	Analysis of DNA damage	lung cancer	Liu et al., 1993
Mouse colon carcinoma (MC38) cells and Hodgkin's lymphoma (L428) cells.	Vegetable oils	Cooking oil	Analysis of NF-κB pathway	Colon cancer	Agbaria et al., 2015
Human obese	Sunflower seed oil and virgin olive oil	Dimethylsiloxane	Assay of omega-3 and omega-6 EFAs	Vascular cancer	Orozco-Solano et al., 2013
Human lung adenocarcinoma CL-3 cells	Soybean oil	Cooking oil fumes and H ₂ O ₂	Assay human 8-oxoguanine DNA glycosylase 1	Lung cancer	Cherng et al., 2002
Human lung adenocarcinoma CL-3 cells	Rapeseed oil	Cooking oil fumes	Assay of TGF beta1, TGF beta2, IL-6, IL-8, and IFN-gamma gene expressions with the RT-PCR method;	Lung cancer	Tung et al., 2001

			intracellular peroxide formation by dichlorofluorescein method		
Human lung adenocarcinoma A549 cells	Rapeseed oil	Benzo[a]pyrene (BaP) and 2,4-decadienal	Assay of IAP2 and phosphorylated Akt proteins, Flow cytometry and terminal deoxynucleotidyl transferase-mediated dUTP nick end labeling (TUNEL) analysis	Lung cancer	Hung et al., 2007
Wistar albino rats	Soybean oil	Cooking oil fume	Assay of MDA, N-acetylcysteine, GSH	Lung cancer	Pu et al., 2002
Human	Vegetable oils	Cooking oil fume	Epidemiological study	Lung cancer	Zhong et al., 1999b
Human	Vegetable oils	Cooking oil fume	Epidemiological study	Lung cancer	Purcaro et al., 2006
Salmonella TA98 and mouse	Rapeseed oil	Cooking oil fume	Ames test, SCE/V79 in vitro and mice micronucleus in vivo test	Lung cancer	Chen et al., 1992
Human	Vegetable oils	Cooking oil fume	Pap smear screening and biopsy examination, community-based case-control study	Cervical cancer	Velema et al., 2002; Lee et al., 2010
Human	Vegetable oils	Cooking oil fume	Histological analysis	Lung cancer	Ko et al., 1997; 2000
Human	Vegetable oils	Cooking oil fume	Histological analysis	Lung cancer	Seow et al., 2000
Wistar albino rats	Mustard oil	Diethylnitrosamine	GST, GGT, liver weight ALP, adenosine triphosphatase, glucose-6-phosphatase	Lung cancer	Kim et al., 1994; Shukla et al., 2004
Human	Vegetable oils	French fries, fried chicken, fried fish, doughnuts and snack chips	Structured questionnaire, BMI,	Prostate cancer	Stott-Miller et al., 2013
Human	Vegetable oils	French fries, fried chicken, fried fish,	Structured questionnaire, BMI,	Prostate cancer	Pawlega et al., 1996
Human	Vegetable oils	French fries, fried chicken, fried fish,	Structured questionnaire, BMI,	Prostate cancer	Li et al., 2014
Human	Olive oil	Fried foods	Structured questionnaire, BMI,	Colorectal cancer	Galeone et al., 2007
Human	Olive oil	Fried fishes	Structured questionnaire	Prostate cancer	Joshi et al., 2012
Human	Vegetable oils	Fried bread, rice, and red meat and deep fried vegetables	Structured questionnaire-gender, age, body mass index, years of education, diabetes and alcohol history, smoking status, and opium use	Pancreatic cancer	Ghorbani et al., 2015
Human	Soybean oil	Fried fishes	Structured questionnaire	Breast cancer	Dai et al., 2002
Human	Vegetable oils	Fried foods	Structured questionnaire	Gastric cancer	Somi et al., 2015
Human	Vegetable oils	potato crisps, chips (deep-	Structured questionnaire	Gastric cancer	Konings et al., 2003; Svensson

		fried), cocktail snacks, and gingerbread			et al., 2003; Mojska et al., 2010
Human	Vegetable oils	Fried foods	Structured questionnaire	Pharyngeal cancer	Zheng et al., 1992
Human	Vegetable oils	Fried foods	Structured questionnaire	Pharyngeal cancer	Ji et al., 1995
Human	Vegetable oils	Tofu, deep-fried tofu, raw fish, and tempura,	Structured questionnaire	Pancreas cancer	Ohba et al., 1996
Human	Vegetable oils	benzo(a)pyrene-containing barbecued, smoked or deep-fried meats	Dietary intake exposure assessment, and intra-individual variance	Lung cancer	Cai et al., 2012; Zhang et al., 2012; Nie et al., 2014; Duan et al., 2016
Human	Vegetable oils	Fried foods	Structured questionnaire and hematological profile	leukemia	Liu et al., 2015
Human	Vegetable oils	Fried foods	Structured questionnaire and hematological profile	Breast cancer	Kojima et al., 2017
Human	Vegetable oils	Fried foods	Structured questionnaire	Stomach cancer	Pham et al., 2010
Human	Vegetable oils	Fried foods	Structured questionnaire	Breast cancer	Cottet et al., 2009; Catsburg et al., 2015
Human	Vegetable oils	Deep fried vegetables	Structured questionnaire	Pancreas cancer	Ghorbani et al., 2015
Human	Vegetable oils	Deep fried foods	Structured questionnaire	endometrial adenocarcinoma	Takayama et al., 2013
Human	Vegetable oils	Deep fried foods	Structured questionnaire	Breast cancer	Datta and Biswas, 2009
Human	Vegetable oils	Deep fried foods	Structured questionnaire	Breast cancer	Hanf and Gonder, 2005
Human	Vegetable oils	Deep fried meats	Structured questionnaire	colon and rectal cancer	Lüchtenborg et al., 2005




Figure 1: List of PAHs in cooking oil fumes (Li et al., 1992, 2003; Chen et al., 2012; Yao et al., 2015)




Figure 2: Repeatedly heated cooking oil emits PAHs and subsequent causes of cancer