



Critical Reviews in Food Science and Nutrition

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/bfsn20>

ONION: NATURE PROTECTION AGAINST PHYSIOLOGICAL THREATS

Hafiz Ansar Rasul Suleria^a, Masood Sadiq Butt^a, Faqir Muhammad Anjum^a, Farhan Saeed^a & Nauman Khalid^b

^a National Institute of Food Science and Technology, University of Agriculture, Faisalabad, Pakistan

^b Department of Global Agriculture, Graduate School of Agriculture and Life Sciences, University of Tokyo, 1-1-1 Yayoi, Bunkyo-ku, Tokyo, 113-8657, Japan

Accepted author version posted online: 14 Jun 2013.

To cite this article: Hafiz Ansar Rasul Suleria, Masood Sadiq Butt, Faqir Muhammad Anjum, Farhan Saeed & Nauman Khalid (2013): ONION: NATURE PROTECTION AGAINST PHYSIOLOGICAL THREATS, Critical Reviews in Food Science and Nutrition, DOI:10.1080/10408398.2011.646364

To link to this article: <http://dx.doi.org/10.1080/10408398.2011.646364>

Disclaimer: This is a version of an unedited manuscript that has been accepted for publication. As a service to authors and researchers we are providing this version of the accepted manuscript (AM). Copyediting, typesetting, and review of the resulting proof will be undertaken on this manuscript before final publication of the Version of Record (VoR). During production and pre-press, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal relate to this version also.

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.tandfonline.com/page/terms-and-conditions>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

ONION: NATURE PROTECTION AGAINST PHYSIOLOGICAL THREATS

**Hafiz Ansar Rasul Suleria¹, Masood Sadiq Butt¹, Faqir Muhammad Anjum¹, Farhan
Saeed¹ and Nauman Khalid^{2*}**

¹ National Institute of Food Science and Technology, University of Agriculture, Faisalabad
Pakistan

²Department of Global Agriculture, Graduate School of Agriculture and Life Sciences,
University of Tokyo, 1-1-1 Yayoi, Bunkyo-ku, Tokyo, 113-8657, Japan

Corresponding Author

Nauman Khalid^{2*}

Department of Global Agriculture

Graduate School of Agriculture and Life Sciences

The University of Tokyo, Japan

nauman_khalid120@yahoo.com

+81-80-3385-0786

ABSTRACT

Onion (Allium cepa L.) is found in various regions of Europe, North America, Asia, and Africa. It is one of the classic examples of allium species used not only for culinary preparations but also for medicinal purposes. Onion with a variety of purpose, often used as a raw material in many dishes and accepts almost all of the traditions and culture. Owing to its storage characteristics and durability of the shipping, Onions have been traded more widely than most vegetables. The pungent fractions of the garlic are mostly sulfur containing moieties while its two chemical groups have marked effect on human health. These are flavonoids and ALK (EN)-based cysteine sulfoxides (ACSOs). Compounds in onions have been reported with a range of health benefits, including anti-cancer properties, anti-platelet activity, antithrombotic activity, antiasthmatic and antibiotic effects.

KEYWORDS: *Allium cepa, Physico-chemical characteristics, nutritional benefits, functional benefits, health promoting potentials*

BACKGROUND

Medicinal spices are used to enhance flavor and aroma in many parts of the world. The economy from spices, accounting for about \$ 2 billion tons (Srinivasan, 2005). Possible medical benefits of spices include the risk of atherosclerosis, cardiovascular disease, cancer and diabetes

(Ali et al., 2000; Ashraf et al., 2005; Banerjee et al., 2003; Cazzola et al., 2011; Lai and Roy, 2004). Herbs are nature's gift to mankind and an herbal revitalization is blooming across the world. Medicinal herbs contain substances that are well known to modern and ancient civilizations for their healing properties. They have been important to mankind, both socially and economically, for thousands of years (Butt et al. 2009).

Over the centuries, species in the genus *Allium* have attained reputations and special attention in the tradition of many cultures as formidable prophylactic and therapeutic medicinal agents (Kendler, 1987). It is cited in the Egyptian Codex Ebers, a 35 century old document, useful in the treatment of heart disorders, tumors, worms, bites, and other ailments (Fenwick and Hanley, 1985). Hippocrates and Pliny the Elder were promoters of *Allium's* medicinal virtues, Charak, the father of Ayurvedic medicine, claimed that garlic and onion maintains the fluidity of blood and strengthens the heart (Chutani and Bordia, 1981).

World onion production is increased by at least 25%, than in the past 10 years, the current output of about 44 million tons, making it the second most important horticultural crop after tomatoes (FAO Statistics, 2011). Onion has also been utilized as a medicinal agent according to sources dating from ancient times. Mixed with honey, it was used to treat such disparate conditions as visual disorders and dog bites (Fenwick and Hanley, 1985). According to Corzo-Martinez et al. (2007) garlic and onion intake offer protection against cancer development, and its utilization as therapeutic agents seems to be very safe. Onions contain a flavonoid called quercetin, which delays oxidative damage in cells and other bodily tissues (National Onion Association, 2011). Quercetin is the most abundant flavonoid in the human diet, and is mainly found in onions (Duthie and Dobson, 1998). Studies done at Wageningen Agricultural

University, the Netherlands, showed twice absorption of quercetin from onion relative to tea, and thrice as compare to apple (National Onion Association, 2011).

In this global village, health claims regarding the potential benefits of onion or its various preparations has been approved and are now available in the market as dietary supplement, herbal products and getting a wide range of popularity now a days in many cultures for their hypolipidemic and procirculatory effects. Onion and its various preparations are imperative functional foods with diverse health benefits. Evidence also ensures its possible applications in cancer therapy, reactive oxygen species (ROS) associated diseases and certainly in immunonutrition (Butt et al., 2009; Butt et al., 2007). This review is also an effort to clarify the importance of onion and its constituents as nature's gift to fight against various physiological threats.

CLASSIFICATION AND CHEMISTRY

There are about 660 species in the genus *Allium* that are mostly found in Asia but with some species in Africa, Central and South America. Onions are believed to have originated from Afghanistan/Iran/USSR region and are now grown in over 175 countries worldwide. Onions (*Allium cepa*) belong to the family Liliaceae (Butt et al., 2009), although they are listed by some authors in different families like Alliaceae. Onions are perennial crop, which may be red, white or yellow in color, and consumed in its tender state, raw, ripe, and pickled or in form of powder (FAO, 2011). The plants bear small flowers that are usually white or purple. Onions are highly valued for their flavor and nutritional value and preserved in the form of pickles. Its leaves are also used in salads and soups.

Insert Figure 1 here

Chemical composition of Allium cepa

Onion has an approximately 90% water with high content of dietary fibre and sugars. In terms of vitamins and minerals content onion has low sodium content and a high content of vitamin B₆, folic acid, calcium, magnesium, phosphorus and potassium. By difference, onion has low content of lipids and among the amino acid content only arginine and glutamic acid are extraordinary.

Insert Table 1 and 2 here

Flavonoids, Quercetin and quercetin glucosides

Onion is the richest source of dietary flavonoids and contributes to a large extent to the overall intake of flavonoids. Two flavonoid subgroups are found in onion, the anthocyanins, which impart a red to purple color to some varieties and the flavonols such as quercetin and its derivatives responsible for the yellow and brown skins of different varieties (Griffiths et al., 2002). Flavonols are the predominant pigments of onions. Only compounds belonging to the flavonols, the anthocyanins, and the dihydroflavonols have been reported to occur in onion bulbs. At least 25 different flavonols have been characterized, and quercetin derivatives are the most important ones in all onion cultivars. Their glycosyl moieties are almost exclusively glucose, which is mainly attached to the 4', 3, and/or 7-positions of the aglycones. Analogous derivatives of kaempferol and isorhamnetin have been identified as minor onion pigments (Slimestad et al., 2007).

Quercetin 3, 4'-diglucoside (QDG) and quercetin 4'-glucoside (QMG) are in most cases reported as the main onion flavonols in literature (Caridi et al., 2007). These glucosides of quercetin represent about the 90% of the overall contents in different *Allium* species (Bonaccorsi

et al., 2008). Significant differences in the levels and ratios of the two compounds were identified among red ('Redwing' QDG 191 mg/100 g dry weight-dw and QMG 85 mg/100 g dw), brown ('Cream Gold' QDG 153 mg/100 g dw, QMG 58 mg/100 g dw), and white onion varieties ('Spanish white' QDG < 1 mg/100 g dw, QMG < 1 mg/100 g dw) (Caridi et al., 2007). Likewise, early studies showed the presence of predominantly cyanidin 3-glucoside, with lesser amounts of cyanidin 3-laminaribioside and other minor unidentified cyanidin, peonidin and pelargonidin glycosides. (Terahara et al., 1994) determined the anthocyanins in the Japanese cultivar Kurenai having cyanidin 3-glucoside, cyaniding 3-laminaribioside and their 6''-malonyl derivatives. Fossen et al. (1996) reported four major and six minor anthocyanins content in different cultivars like Red Baron, Tropea and Comred (grown in Norway) including the 3-malonylglucoside, 3 dimalonylglucoside and 3,5-diglucoside derivatives of cyanidin, peonidin 3,5-diglucosides and two 3-glycosylated derivatives of pelargonidin. In red onion, grown in Canada and the USA (Mambo, Red Jumbo, Red Bone and Red Granex), the main anthocyanins were cyanidin 3-glucoside, cyaniding 3-laminaribioside, cyanidin 3-(6''-malonylglucoside) and cyanidin 3-(6''- malonyllaminaribioside) (Donner et al., 1997). Minor anthocyanins were shown to be cyanidin 3 (3''-malonylglucoside), peonidin 3-glucoside, peonidin 3-malonylglucoside and cyaniding 3-dimalonyllaminaribioside. These differences in anthocyanin composition are probably due to genetic variations.

Insert Figure 2 here

Organosulfur compounds (OSCs)

Allium plants contain high concentrations of alk(en)yl cysteine sulfoxides (ACSOs) (Lee et al., 2011). Among them, onions, shallots, and leeks contain the methyl, propyl, and prop-1-

enyl cysteine sulfoxides. When the onion tissue is disrupted by cutting, crushing or chewing, the ACSOs are enzymatically degraded by the enzyme allinase to iminopropionic acid and alk(en)yl cysteine sulfenic acids. The iminopropionic acid spontaneously hydrolyses to form ammonia and pyruvic acid. The sulfenic acids decompose spontaneously afterwards, methyl and propyl sulfenic acids yield mainly thiosulfinates, while prop-1-enyl sulfenic acid forms both, the corresponding thiosulfinate and thiopropanal S-oxide, the onion lachrymatory factor (Block et al., 1993).

Onion pungency is caused by a wide range of sulfur compounds that causes burning sensation in the back of the mouth and throat. A convenient method to measure onion pungency is to analyze pyruvic acid, which is formed in stoichiometric amount to the thiosulfinates. Pyruvic acid has been shown to correlate well with flavour perception. The balance between level of pungency and level of sugars determines the perception of sweetness in an onion. High pungency can mask a high level of sugars so that the onion is not perceived as sweet. Also, onions with low pungency and low sugar content can be perceived as bland. Ideally, a sweet onion would have a high level of sugars and low pungency (Vagen and Slimestad, 2008).

Thiosulfinates are unstable, particularly on heating, and break down to a complex mixture of compounds, in which mono-, di-, tri- and tetra-sulfides predominate (Munday and Munday, 2004; Rose et al., 2005). Dipropyl disulfide, dipropyl trisulfide, and propenyl disulfides are the major constituents of onion volatiles, although many others compounds have been identified, including dipropyl sulfide and dipropenyl sulfide (Munday and Munday, 2001).

Insert Figure 3 here

Recently Yamazaki et al. (2011) determined 11 sulfur containing flavor precursors in onion including S-alk(en)yl-L-cysteine derivatives, methiin, alliin, isoalliin, cycloalliin, deoxyalliin, N-(gamma-glutamyl)-S-methyl-L-cysteine, N-(gamma-glutamyl)-S-(2-propenyl)-L-cysteine, N-(gamma-glutamyl)-S-(E-1-propenyl)-L-cysteine (Glu-PEC), N-(gamma-glutamyl)-S-(2-propenyl)-L-cysteine sulfoxide, N-(gamma-glutamyl)-S-(E-1-propenyl)-L-cysteine sulfoxide (Glu-PECSO) and S-(2-carboxypropyl) glutathione.

Fructans and fructooligosaccharides (FOS)

Bulb dry matter content is an important quality parameter of onion, also significant in the onion dehydration industry as it directly relates to the energy needed for drying. About 65 to 80% of bulb dry matter consists of non-structural carbohydrates (Benkeblia, 2005; Kahane et al., 2001). Onion's predominant non-structural carbohydrates are glucose, fructose, sucrose and low-molecular-weight fructans, while starch and raffinose are absent. Fructans, also known as fructooligosaccharides (FOS), are polyfructosyl sucroses of varying molecular size that constitute the main carbohydrate reserve of onion. Fructans accumulate during bulbing followed by catabolism during growth and sprout development of the bulbs (Benkeblia, 2005).

It is generally accepted that FOS is a common name only for fructose oligomers that are mainly composed of kestose (GF2), nystose (GF3) and fructofuranosylnystose (GF4), in which fructosyl units (F) are bound by β -linkage at the position of sucrose (glucose+fructose-GF) respectively. There is a clear predominance of kestose (GF2) in every onion tissue and no occurrence of highly polymerized fructans. The tissues richest in fructans are the fleshy layers,

so that the outer two fleshy layers turn out to be the best onion by-product as a possible fructan source (Jaime et al., 2001a; Jaime et al., 2002).

The fructan degree of polymerization (DP) level in onion is mostly in between 3 and 15. Short chain fructans, with a degree of polymerization less than 5, are potentially used as natural low-calorie sweeteners. Onion bulbs with fructans of a high DP may be used for lipid replacement with consequential health benefits (Kahane et al., 2001). Onion showed a better soluble/insoluble dietary fibre (SDF:IDF) ratio than other vegetables that are connected with different metabolic and physiological effect. SDF increases the viscosity of the stomach contents, thereby allowing down-mixing and absorption of nutrients, whereas IDF reduces intestinal transit time and increases the bulk of the food mass (Jaime et al., 2002). Fructans could act as osmoregulators due to their solubility in water inside the vacuole. Fructans act stimulating the growth of specific microorganisms in the colon (*Bifidobacteria* and *Lactobacilli*) with a general positive health effect, including on colonic inflammation (Ernst and Feldheim, 2000; Roberfroid, 2007). Administration of FOS significantly lowered fasting glycemia and total cholesterol, increasing the intestinal absorption and bone density of calcium and magnesium.

Other onion bioactive compounds

Recent literature reported that several interesting novel compounds have been isolated from onion. Among them, saponins and peptides have been isolated and studied for their potentially beneficial health perspectives. A number of different saponins have been identified in *Allium species*, with processing give rise to different saponins again (Corea et al., 2005; Lanzotti, 2006). 5-hydroxy-3-methyl-4-propylsulfanyl-5H-furan-2-one, and four others compounds, were isolated and confirmed to be quinone reductase and glutathione *S*-transferase inducers *in vitro*

(Xiao and Parkin, 2007), therefore they could act as chemopreventive agents. Several research reported antifungal, antitumor, cytotoxicity, blood coagulability, antispasmodic and cholesterol-lowering effects of saponins isolated from onion and garlic (Lanzotti, 2006).

Four furostanol saponins, two of which were new compounds, named ceparoside A and ceparoside B were isolated from the seeds of *Allium cepa* L. (Yuan et al., 2008). Other new saponins were found by Corea et al. (2005) they were reported to possess antispasmodic activity in the guinea pig isolated ileum such an effect might contribute to explaining the traditional use of onion in the treatment of disturbances of the gastrointestinal tract. Recently, it was also reported that gamma-glutamyl peptide from onion (Wetli et al., 2004) inhibits the development and activity of osteoclasts *in vitro* (Langos et al., 2007).

HEALTH BENEFITS OF ONIONS

Allium vegetables health properties have been supported by numerous *in vitro*, *in vivo*, and *ex-vivo* studies. Particularly, onion has been described to have several health benefits related to its antioxidant, anticarcinogenic, hypolipidemic, hypoglycaemic, and antiaggregatory effects. From medical and nutritionally point of view, it has to be considered that the onion used as a food or a food ingredient in many dishes also exerts a wide an array of medicinal effects that are very interesting for its human health potentials. It has been described that a diet rich in *Allium* vegetables, including onion, would lead to several health benefits that could be helpful in the prevention of two of the more relevant and prevalent diseases nowadays such as CVD and cancer insurgences.

Onion and cancer

Research has indicated that onions may have a role in the prevention of a wide range of different cancers, including colorectal, stomach, liver, renal, lung, bladder, breast, ovarian, brain and oesophagus cancer. A large and recent European study, published in 2006 in the American Journal of Clinical Nutrition, found that moderate frequency of onion consumption protected against colorectal, laryngeal and oesophageal cancers. More frequent consumption was even strongly protective and was also significant for oral cavity and oesophageal but not for prostate, breast or renal cell cancers (Galeone et al., 2006). Hsing et al. (2002) also showed the anti-tumour effects of onions, with men consuming 10 g of onions per day being 70% less likely to develop prostate cancer than those consuming less than 2 g of onions per day. The organosulfur compounds in onions proved to be strong anticarcinogens in cell experiments, animals and human trials (Chu et al., 2002; Fukushima et al., 1997; Hatono et al., 1996; Munday and Munday, 2001). This is thought to be partially because of their role in the activation of detoxifying enzymes, which remove potentially cancer-causing substances.

In last years, scientists have focused their research activities on the prevention of cancer by nutrition with a diet containing high proportion of vegetables and fruits. Reviews concerning the effects of the *Allium species* were published by (Block, 1996; Corzo-Martinez et al., 2007; Kendler, 1987; Krishnaswamy and Polasa, 1995; Steinmetz and Potter, 1996; Wargovich and Uda, 1996). Onions are rich in organoselenium compounds, which may help to prevent cancer. Selenium is usually fixed to sulphur-containing amino acid derivatives. Quercetin and its derivatives, which are also typical constituents of onions are of great interest, because of their anticarcinogenic properties (Formica and Regelson, 1995).

The association between consumption of *Allium* vegetables and risk for cancer has been assessed in several epidemiologic studies, mainly case-control (Bianchini and Vainio, 2001; Galeone et al., 2006). In general, these studies are more consistent in reporting a protective effect of onion in gastric cancer (Zhou et al., 2011). However, onion consumption has been also consistently related with a decreased colorectal cancer risk. In addition, onion consumption was reported to decrease the risk for the cancer of the lung (Sankaranarayanan et al., 1994) and of the brain (Hu et al., 1999) in case-control studies. Onion consumption was significantly inversely correlated with the risk of the stomach cancer. Recently, (Gonzalez et al., 2006; Gonzalez et al., 2011) observed in the European Prospective Investigation into Cancer and Nutrition (EPIC-EURGAST) a probable protective effect of total vegetables and *Allium* vegetables intake on the intestinal type of gastric cancer and cervical cancer. Most of the case-control studies concerning onion were conducted in China (Gao et al., 1999; You et al., 1989) and several of them in Asia and Europe (Boeing et al., 1991; Gonzalez et al., 1991; Hansson et al., 1993; Tuyns et al., 1992). A cohort study was conducted in The Netherlands (Dorant et al., 1996). The antimutagenic effect of ethanolic extracts of onion and garlic against the mutagenicity of *N*-nitrosamines was evaluated by the Ames test (Ikken et al., 1999). Onion showed a greater effect than garlic. Onion consumption reduced the risk of carcinoma in the non-cardial part of the stomach (Challier et al., 1998). Onions also provide protection against aorta and liver cancer recently identified by (Gorinstein et al., 2011). Similarly, it also provide protection against breast cancer (Viry et al., 2011).

Diets rich in fruit and deep-yellow vegetables, dark-green vegetables, onions and garlic are modestly associated with reduced risk of colorectal adenoma, a precursor of colorectal cancer

(Millen et al., 2007). Decreased risk for colorectal cancer with the consumption of onion was generally found in case-control studies. The effect was particularly significant for consumption of cooked onions and leeks in Belgium (Tuyns et al., 1988), for a combination of garlic, onions, and pepper in Argentina (Iscovich et al., 1992). In a case-control study in Australia (Steinmetz and Potter, 1993) it was reported a lower risk for both sexes, with a more pronounced decrease for women and also for cancer of the proximal compared with the distal colon. However, a cohort study in The Netherlands showed no significant effect of consumption of onions, leeks, and garlic supplements (Dorant et al., 1996).

Insert Table 4 here

In addition to organo-sulphur compounds, eruboside-B, a steroid saponin isolated from garlic bulb, and organo- Se compounds are largely responsible for the anticarcinogenic activity. Se-enriched garlic and onion have higher anticarcinogenic activity than the common plants (El-Bayoumy et al., 1996; Matsuura, 1997). This increased effect of cancer prevention is achieved at least partly by S substitution with Se. The pure Se-compounds have proved to be superior anticancer agents than their corresponding S-analogues. For example, diallyl selenide is at least 300 times more active than DAS in the reduction of tumors of mammal cancer (El-Bayoumy et al., 2006). The two major Se-compounds possessing anticancer activity in onion are g-glutamyl-Se-methyl selenocysteine and Se-methyl selenocysteine, being Se-methyl selenocysteine and Se-allyl selenocysteine the most chemopreventive Se-compounds (Block et al., 2001). Quercetin and kaempferol, from onion, possess anticarcinogenic properties. Particularly, they have antineoplastic effects by inhibiting bioactivating enzymes (Lautraite et al., 2002), by inducing

detoxifying enzymes and apoptosis (Brisdelli et al., 2007), due to their antioxidant and anti-inflammatory activities (Raso et al., 2001).

Several epidemiological studies have found inverse associations between lung cancer risk and onion intake, probably due to its high content of flavonoids (Le Marchand et al., 2000). Quercetin enhances bioavailability of some anticancer drugs, as Tamoxifen, a non-steroidal antiestrogen for treating and preventing breast cancer, by promoting their intestinal absorption and reducing their metabolism (Shin et al., 2006; Wu et al., 2005). Large-scale gene expression analysis in combination with functional assays yields a considerable amount of information on anticarcinogenic potential of all these active components. Thus, for example, data from cDNA array studies revealed that the antiproliferative effects of DADS may be related to changes in gene expression of aggrecan 1, tenascin R, vitronectin and cadherin 5 (Knowles and Milner, 2003). A recent study (Van Erk et al., 2005) showed the broad range effects at gene expression level, elucidated that quercetin exerted effect on colon cancer cells *in vitro*. Likewise, it has been recently elucidated the response of garlic and its components depends on the consumer's genetic backgrounds (nutrigenetic effects), DNA methylation and histone regulation (nutritional epigenomic effects), ability to induce or repress gene expression patterns (nutritional transcriptomics effects), occurrence and activity of specific proteins (nutriproteomic effects), and/or dose and temporal changes in cellular small-molecularweight compounds (metabolomics effects).

Onion and cardiovascular disease (CVD) prevention

CVD include coronary heart disease (heart attacks), cerebrovascular disease, raised blood pressure (hypertension), peripheral artery disease, rheumatic heart disease, congenital heart disease and heart failure. If current trends are allowed to continue, by 2015 an estimated 20 million people will die from CVD (mainly from heart attacks and strokes). Therefore, CVD have a major impact on the mortality and quality of life on human populations across the world, despite improvements in lifestyle and innovations in the prevention and treatment of CVD in previous decades (Wensing et al., 2009). CVD risk factors are mainly determined by uncontrollable causes (heredity, gender and age) and lifestyle-related causes (smoking, physical inactivity, stress and unhealthy diet), which are possible to be modified.

The study by Galeone et al. (2009), the first from Mediterranean countries, suggested that a diet rich in onions may have a favorable effect on the risk of acute myocardial infarction; therefore these vegetables could be useful in a CVD preventive diet. Several biomarkers are measured to predict CVD events including blood lipids levels (LDL-cholesterol and triglycerides), fibrinogen (a marker of thrombosis and inflammation), D-dimer (a marker of thrombosis), plasminogen-activator inhibitor type 1 (a marker of fibrinolytic potential and endothelial function), high-sensitivity creactive protein (CRP) (inflammation marker), homocysteine (a marker of endothelial function and oxidant stress), B-type and N-terminal pro-atrial natriuretic peptides, serum aldosterone, plasma renin (markers of neurohormonal activity), and urinary albumin-to-creatinine ratio (a marker of glomerular endothelial function) (Kannel, 2005; Wang et al., 2006). Alterations in lipid profiles, diabetes, hypertension, and obesity are risk factors conventionally associated to the early appearance of CVD. Onion has been described to have hypolipidemic, hypoglycaemic, and antithrombotic effects and therefore could be useful

in the CVD prevention. Focusing on onion lipid lowering effects, this vegetable has been reported to exert moderately hypolipidemic effects in experimental animal such as healthy pigs fed a high fat diet and consequently reduction of CVD and obesity (Gabler et al., 2006; Ostrowska et al., 2004).

Among bioactive compounds involved in onion hypolipidemic effects, quercetin has shown to have the ability to reduce serum cholesterol levels and arteriosclerosis severity (Glasser et al., 2002). A recent study by Kumari and Augusti (2007) also proclaimed for the lipid lowering action of the S-methyl cysteine sulfoxide (SMCS) isolated from *Allium cepa* L. Onion has also been reported to have hypoglycaemic effects (Srinivasan, 2005). It was inferred that beneficial ameliorating influence of dietary onion on diabetic nephropathy may be mediated through onion's ability to lower blood cholesterol levels and to reduce lipid peroxidation, dietary onion caused significant beneficial modulation of the progression of renal lesions in diabetic rats (Babu and Srinivasan, 1999). Other rat studies have assessed onion hypoglycaemic effects (El-Demerdash et al., 2005).

Recently, Lee et al. (2008) showed that onion skin was effective in controlling hyperglycemia in animal models of type 2 diabetes mellitus, at least in part by inhibiting alpha-glucosidase activity. Thrombosis complications play a major role in CVD. Blood clot formation depends on an intricate series of events involving platelets, other cells, and the activation of specific blood proteins, known as coagulation factors. A thrombus is a blood clot formed when there is an imbalance in the blood coagulation system that can block the flow of blood through a vein or artery, and can detach from the vessel wall to become a life-threatening embolus when it lodges in the lungs or other vital organs. Blood clots in coronary arteries cause acute coronary

syndrome and blood clots that form in the heart are the major cause of stroke in people with atrial fibrillation. Onion inhibits platelet aggregation *in vitro* and *in vivo* (Ali et al., 1999; Ali et al., 2000; Briggs et al., 2001; Hubbard et al., 2006; Jung et al., 2002). The mechanism by which onion exerts its antithrombotic effect has been shown to involve the inhibition of thromboxane A₂ formation, potent inducer of platelet aggregation (Moon et al., 2000).

The antiplatelet activity observed in onion is influenced by genotype, environmental factors and genotypically determined sulfur content of the bulb (Goldman et al., 1996; Sance et al., 2008) having onion α -sulfinil-disulfides (cepaenes) a demonstrated antithrombotic activity (Block et al., 1997). Another recent study by Sakakibara et al. (2008) suggested that onion exerted antidepressant-like activity in a behavioural model that acted independently of the hypothalamic-pituitary-adrenal axis. Similarly, Brankovic et al. (2011) studied the acute effect of ethanol extracts ginkgo (*Ginkgo biloba* L.), garlic (*Allium sativum* L.), and onion on arterial blood pressure (BP), and heart rate (HR) in anesthetized normotensive rats. They concluded that the most effective in reducing arterial BP and HR is extract of garlic and onion.

Antiplatelet or antithrombotic effect

The main focus of recent research is on antiplatelet activity of onion extracts. Reducing platelet aggregation has a preventive effect on some cardiovascular disorders, such as atherosclerosis. In addition, onion extracts have some lipid-lowering effects. The antithrombotic potential of aqueous extracts from onion and garlic was evaluated firstly by Bordia et al. (1996). Extracts were administered orally and intraperitoneally to rats. A relatively low dose of the aqueous garlic extract (50 mg kg⁻¹ body weight) decreased thromboxane-B₂ levels significantly,

regardless of the mode of administration. Onion extracts were effective at higher concentrations (500 mg kg⁻¹ body weight). Boiling the extracts before application resulted in an almost complete loss of activity. Cooking may therefore cause the decomposition of the potent antithrombotic components in alliums.

The effects of aqueous extracts of raw and boiled onion and garlic on collagen induced platelet aggregation were studied *in vitro* (Ali et al., 1999; Ali et al., 2000). The concentrations of onion and garlic required for a 50% inhibition of the platelet aggregation were calculated to be 90 and 7 mg/ml plasma (rabbit), respectively. Boiled extracts showed a reduced inhibitory effect. Goldman et al. (1996) investigated the antiplatelet activity of extracts of four bulbonion cultivars (mild cvs 'Exhibition' and 'MSU8155B' and pungent cvs 'W434B' and 'W420B'), grown at two different locations in the USA. The highest antiplatelet activity was observed in the two pungent cultivars, i.e. onion bulbs with high sulphur content exhibited a significantly greater antiplatelet activity than those containing low levels of sulphur. Significantly greater activity was determined for three out of four cultivars grown in Oregon compared with those from Wisconsin. During postharvest cold storage at 4°C, antiplatelet activity was increased by about 60% across all cultivars, reaching a maximum at 90 days of storage. However, changes in pungency were not correlated with changes in antiplatelet activity (Debaene et al., 1999). Juice prepared from a flowering umbel of the onion genotype 'W420B' had antiplatelet activity 336% higher than that of juice obtained from onion bulbs (Goldman et al., 1996).

The active principle of onion and related species in terms of inhibition of platelet aggregation is not yet identified but previous studies showed that antithrombotic action of onion aqueous extracts was because of inhibition of thromboxane formation, potent inducers of platelet

aggregation (Moon et al., 2000). However, some studies have been carried out with isolated compounds (Makheja and Bailey, 1990). Allicin, which occurs more abundantly in garlic than in onion, and adenosine, both inhibited platelet aggregation without affecting cyclo-oxygenase and lipoxygenase metabolites of arachidonic acid. The trisulphides investigated inhibited platelet aggregation, as well as thromboxane synthesis, along with the induction of lipoxygenase metabolites. The observed *in vivo* antiplatelet effects appear to be more attributable to adenosine than to allicin and alk(en)yl polysulphides of onion. Several epidemiologic studies have reported that antiplatelet activity of onion is considered to be a property of organo-sulphur compounds. In particular, a class of α -sulphinyl-disulphides (cepaenes) found in onion extracts has demonstrated antithrombotic activity (Block et al., 1997). These compounds have structural similarity to ajoene, considered the major antiplatelet compound in garlic extracts. In addition, other non-sulphur compounds, such as b-chlorogenic acid and quercetin, have also been shown to inhibit platelet aggregation (Rahman and Lowe, 2006).

Antiplatelet activity is substantially affected by genotype, environment and storage duration of vegetable. It has been reported that onions and garlic's antiplatelet activity was determined by the native concentration of organo-sulphur compounds and genotypically (Goldman et al., 1996), being garlic 13 folds more potent than onion (Effendy et al., 1997).

Bone strengthening Properties

Onion consumption has also been reported to be involved in the bone metabolism and in the behaviour as a possible antidepressant agent. A recent study by Matheson et al. (2009) reported that onion consumption seems to have a beneficial effect on bone density in perimenopausal and postmenopausal women. Furthermore, older women who consume onions

most frequently may decrease their risk of hip fracture by more than 20% *versus* those who never consume onions. Prevention of low bone mass is important to reduce the incidence of osteoporotic fractures. Onion retains its bone resorption inhibitory activity in the rat when added to a vegetarian diet (Muhlbauer et al., 2002).

Antimicrobial and Antifungal Properties

Onions have been shown to possess antibacterial and antifungal properties (Kyung, 2011; Mohamed, 2010; Santas et al., 2010). Volatile oil of onion has been shown to be highly effective against Gram positive bacteria, dermatophytic fungi, growth and aflatoxin production of *Aspergillus* fungi genera including *Aspergillus niger*, *Brettanomyces anomalus*, *Candida albicans*, *C. lipolytica*, *Cladosporium werneckii*, *Fusarium oxysporium*, *Geotribum candidum* and *Saccharomyces cerevisiae*. Onion oils and aqueous extracts were almost ineffective against Gram-negative bacteria (Griffiths et al., 2002). Onion extracts inhibit oral bacteria causing dental caries (Kim, 1997). In contrast, comparable garlic preparations were active against Gram-negative bacteria (Dankert et al., 1979; Elnima et al., 1983; Yoshida et al., 1999a; Yoshida et al., 1999b). Aqueous extract or the juice of onion has been reported to inhibit *in vitro* growth of *Escherichia coli*, *Serratia marcescens*, *Streptococcus species*, *Acetobacillus odontolyticus*, *Pseudomonas aeruginosa* and *Salmonella Typhosa*, *Streptococcus mutans*, *Streptococcus sobrinus*, *Porphyromonas gingivalis* and *Prevotella intermedia* (Bakri and Douglas, 2005). A petroleum ether extract of onion inhibited the *in vitro* growth of *Clotridium paraputrificum* and *Staphylococcus aureus*.

Similarly, Onion extracts are effective against many yeast species and their essential oil inhibits the dermatophytic fungi (Kunicka-Styczynska, 2011; Zohri et al., 1995). The active compounds of onion destroy fungal cells decreasing the oxygen uptake, reducing cellular growth, inhibiting the synthesis of lipids, proteins and nucleic acids, changing the lipid profile of the cell membrane and inhibiting the synthesis of the fungal cell wall (Gupta and Porter, 2001).

The main active antifungal agents from onion extracts are the breakdown products of allicin, including diallyl trisulphide (DATS, DADS, DAS) and ajoene, which have a greater antifungal effect than allicin (Tansey and Appleton, 1975). An antifungal compound, fistulosin (octadecyl 3-hydroxyindole), has been isolated from welsh onion (*A. fistulosum*), which showed a high activity towards *Fusarium oxysporum* inhibiting primarily the protein synthesis (Phay et al., 1999). In addition to sulphur compounds, a great variety of antifungal proteins and peptides have been isolated from several *Allium* species (Lam et al., 2000; Wang and Ng, 2001) such as allicepin, a novel isolated antifungal peptide from onion bulbs (Wang and Ng, 2004). Finally, it is necessary to consider certain steroid saponins, such as eruboside-B, isolated from the garlic bulb that also exhibit antifungal activity for *Candida albicans* (Matsuura et al., 1988).

Welsh onion extracts have been reported to exert more inhibitory activity towards aflatoxin production than the preservatives sorbate and propionate at pH values near 6.5, even at concentrations 3-10 folds higher than maximum level used in foods. In addition to inhibitory effects against pathogenic bacteria, onions have been found to promote beneficial microorganisms. Onions contain fructo-oligosaccharide (FOS), probiotics which are non-digestible ingredients fermented by bifido bacteria in the body that help to maintain the health of

the gut and colon. Onions contain 2.8% FOS as compared to 1.0% FOS in garlic, 0.7% in rye, and 0.3% in bananas.

Insert Table 5 here

Chemical characterization of sulphur compounds has proven the main active ingredient for antimicrobial activity (Rose et al., 2005). In addition some proteins, saponins and phenolic compounds can also contribute to this activity (Griffiths et al., 2002). Recently it was found that onion peel could be a good source of functional biomaterial for pharmaceutical industries if prepared by sub critical water extraction (SWE) because onions have strong antimicrobial activity against *Bacillus cereus* (Kim et al., 2011).

Antiparasitic activity

Onion extracts are effective against *Opalina ranarum*, *Opalina dimidicita*, *Balantidium entozoon*, *Entamoeba histolytica*, *Tripanosoma brucei*, *Leishmania*, *Leptomonas* and *Crithidia* (Reuter et al., 1996). In China, DATS, an allicin breakdown product, easily synthesized and more stable than the extremely volatile allicin, is commercially available as a preparation, called Dasuansu, prescribed for the treatment of giardiasis and infections by *E. histolytica* and *Trichomonas vaginalis* (Lun et al., 1994). Allicin, ajoene and other organo-sulphur compounds from garlic and onion are also effective antiprotozoals. Antiparasitic properties of onion extracts towards different strains of *Leishmania* and *T. vaginalis* have been reported as well (Saleheen et al., 2004). Similarly administration of onion was beneficial in protecting susceptible hosts against opportunistic zoonotic parasites such as *Cryptosporidium parvum* (Nadia et al., 2011). Onion and garlic intake can significantly cause reduction of worm burden, hepatic and intestinal

eggs and oogram count in mice infected with *Schistosoma mansoni* recently reported by (Mantawy et al., 2011).

Antiviral activity

Quercetin, the major onion flavonoid, possesses antiviral activity and enhances the bioavailability of some antiviral drugs (Wu et al., 2005). Lectins are heterogeneous group of glycoproteins with the ability to recognize and bind specifically to carbohydrate ligands. Chen et al. (2011) studied the antiviral activity of extract of five *Allium* plants (shallots, garlic, onions, leeks, and green onions) in comparison with compounds of quercetin, zalcitabine (ddC), allicin, and ribavirin against adenovirus. With regard to the antiviral activity of ddC, both the MU and PRD (Plaque reduction) methods showed a high correlation with no significant difference. Most of the *Allium* plants tested were non-toxic to human lung carcinoma (A549) cells, and shallots exhibited the highest level of antiviral activity for both ADV41 and ADV3, followed by garlic and onions.

Antioxidant activity

Oxidation of DNA, proteins and lipids by reactive oxygen species (ROS) plays an important role in a wide range of common diseases, including cancer and cardiovascular, inflammatory and neurodegenerative diseases, such as Alzheimer's disease and other age-related degenerative conditions (Borek, 1997; Richardson, 1993).

Flavonoids, ubiquitous in the plant kingdom, have been widely studied for their antioxidative effects (Hertog and Katan, 1998; Rice-Evans et al., 1995). Onions are known to contain anthocyanins and the flavonols quercetin and kaempferol (Bilyk et al., 1984; Rhodes and Price, 1996) and both have antioxidant activity. Onion is one of the major sources of dietary

flavonoids in many countries, which are present either as sugar conjugates or as aglycones. The antioxidative effects of consumption of onions have been associated with a reduced risk of neurodegenerative disorders (Shutenko et al., 1999), many forms of cancer (Hertog and Katan, 1998; Kawaii et al., 1999), cataract formation (Sanderson et al., 1999) and ulcer development (Suzuki et al., 1998). Data from a range of *in vitro* testing methods suggested onions have moderate levels of antioxidant activity in comparison with other vegetables (Halvorsen et al., 2002; Pellegrini et al., 2003; Wu et al., 2004).

The major flavonoid found in onion is quercetin, present in conjugated form, as quercetin 4-O- β -glycopyranoside, quercetin 3,4-O- β -diglycopyranoside, and quercetin 3,7,4-O- β -triglycopyranoside (Koncic and Jug, 2011; Sellappan and Akoh, 2002). The dry outer layers of onion, which are wasted before food processing such as cooking, contain large amounts of quercetin, quercetin glycoside and their oxidative products (Gu' lsen et al., 2007), which are effective antioxidants against non enzymatic lipid peroxidation and oxidation of low density lipoproteins (LDL). Quercetin and its dimerized compound showed the highest antioxidative activity, which is comparable to that of α -tocopherol (Ly et al., 2005). It has been confirmed that its absorption is low in contrast to other dietary antioxidants such as vitamins C and E, limiting its capability to act as antioxidant in plasma *in vivo* (Lotito and Frei, 2006).

Onions and Eye health

The carotenoids in spring onions may protect against macular degeneration. Some epidemiologic evidence does suggested that lutein and zeaxanthin protect against age-related eye disease (Mares-Perlman et al., 2002; Sies and Stahl, 2003). Lower risk of eye disease has been found in conjunction with consumption of foods rich in lutein and zeaxanthin (Mares-Perlman et

al., 2002), higher levels of lutein and zeaxanthin in the blood and higher levels of lutein and zeaxanthin in the retina (Beatty et al., 2001; Bone et al., 2000). However, these relationships were not observed in other studies, or were only observed in subgroups of the study population (Mares-Perlman et al., 2002). Mares-Perlman et al. (2002) described findings with respect to the relationship between lutein and zeaxanthin and reducing cataract risk as “somewhat consistent”. Two studies showed a higher incidence of cataracts in those in the lowest quintile of lutein and zeaxanthin intake compared with the highest, and three prospective studies found that those in the highest quintiles had a 20–50% lower risk of experiencing cataract problems. Although concentrations are generally highest in ocular tissue, a number of studies have established the presence of lutein and zeaxanthin in serum and body tissues. Their antioxidant activity has led to speculation that higher consumption of these chemicals lead to higher levels in body tissues, and that this may lower the risk of chronic disease. Lutein is more widely dispersed in the body than zeaxanthin and it is possible that, along with other carotenoids with antioxidant activity, it may confer protection against diseases such as cancer and cardiovascular disease as well as positively affecting immune function.

Daily consumption of more than 500 ml of tea, a large source of quercetin, was associated with decreased risk of cataracts (Robertson et al., 1991). It has been reported that the percentage of quercetin absorbed from onions is approximately twice that from tea (de Vries et al., 1998). Therefore, high daily intake of onions may provide some protection against the risk of cataract formation.

Anti-inflammatory effects

Quercetin's anti-inflammatory effect on prostaglandins, leukotrienes, histamine release and subsequent anti-asthmatic activity has been investigated (Wagner et al., 1990). Inflammation is part of the body's natural immune response to trauma. Thiosulfinates and capsaenins responsible for the anti-inflammatory activities also cause inhibition of the immune response (Chisty et al., 1996; Dorsch and Wagner, 1991). The organosulfur compounds of onions also have been credited with anti-asthmatic effects (Augusti, 1996). Thiosulfinates formed from onion tissue degradation (i.e. chopping) have been credited with inhibition of arachidonic acid metabolic pathways and subsequent anti-inflammatory and anti-asthmatic effects (Wagner et al., 1990). Saponins have also been shown to have anti-inflammatory activity (Sparg et al., 2004).

Anti-diabetic effect

Significant research has been done on the effect of onion consumption on diabetic conditions. Two organosulfur compounds were linked to significant amelioration of weight loss, hyperglycemia, low liver protein and glycogen, and other characteristics of diabetes mellitus in rats. Similarly, Suresh Babu and Srinivasan (1997) found that 3% onion powder reduced hyperglycemia, circulating lipid peroxides and blood cholesterol (LDL-VLDL exclusively). Analysis of the effects of quercetin on human diabetic lymphocytes showed a significant increase in protection against DNA damage from hydrogen peroxide at the tissue level (Lean et al., 1999). Further human studies are needed to assess the ability of a high flavonoid diet to attenuate diabetic conditions. Similarly the antihyperglycaemic effect of 12 edible plants was studied on 27 healthy rabbits (Roman-Ramos et al., 1995). Apart from cauliflower, only onion

and garlic decreased the hyperglycaemic peak, which was induced by the glucose tolerance test. In further studies, alloxan-diabetic rats were treated with S-methyl-L-cysteine sulfoxide (typical for many *Allium* species) and S-allyl-L-cysteine sulfoxide (typical for garlic) at a concentration of 200 mg kg⁻¹ body weight per day (Kumari and Augusti, 2007; Kumari et al., 1995). Both compounds significantly lowered blood glucose levels. Medium levels of FOS (1-kestose (1-kestotriose; GF-2), nystose (1,1-kestotetraose; GF-3), and 1F-beta-fructofuranosylnystose (1,1,1-kestopentaose; GF-4) and sugars (glucose, fructose and sucrose) have been identified in red onions that have significant role in protection against diabetes (Judprasong et al., 2011).

Effect on memory

There has been recent interest in the effects of allium-derived compounds on memory impairment. An animal study showed onion extract and a compound found in onions, di-n-propyl trisulfide, improved memory function in a mouse model and demonstrated that its efficacy was due to antioxidant activity (Nishimura et al., 2006).

Effects on the respiratory system

Certain onion-derived compounds, in particular thiosulfinates and cepaenes, showed remarkable *in vitro* inhibitory effect of cyclooxygenase and lipoxygenase mediated reactions which initiated eicosanoid metabolism and lead to bronchial restriction. Therefore, these compounds have antiasthmatic activity (Wagner et al., 1990). It has been described that saturated thiosulphinates are less active than unsaturated ones and that cepaenes are more active than thiosulphinates. In further studies, guinea-pigs were chosen as the model for investigating the effect of organosulphur compounds on platelet activating factor (PAF)-induced bronchial

constriction (Dorsch, 1996; Dorsch et al., 1987b). Furthermore, a chloroform extract (at 20 mg kg⁻¹ body weight) was more active than the lyophilized onion juice (at 100 mg kg⁻¹ body weight). Thiosulphinates and cepaenes were identified as the active compounds that has been investigated in a previous study by (Dorsch et al., 1987a).

In a human study, adult volunteers suffering from allergic bronchial asthma were treated with 100 ml of a 5% ethanol onion extract prepared from 100 g sliced onions 1 h prior to an allergen inhalation test (Dorsch et al., 1987a). Immediate and late bronchial reactions were markedly reduced after onion pretreatment.

ADVERSE EFFECTS AND TOXICITY

The adverse effects of *Allium* species in humans can be most common, less frequent and rare (Tattelman, 2005). The most common side effects, produced by intake of small amounts of onion, are bad breath and body odour. However, several studies have demonstrated that consumption of excessive amounts onion, especially when the stomach is empty, can cause other less frequent undesirable effects, such as gastrointestinal upsets (burning sensation and diarrhoea), flatulence and changes in the intestinal flora (Ackermann et al., 2001). Within the group of rare effects, dermatological problems in susceptible individuals (allergic dermatitis, burns and blisters), produced by the direct application on skin of fresh or crushed garlic and onion (Davis, 2005; Friedman et al., 2006).

Onion preparations used as complementary medicine with drugs are heavily scrutinized due to their ability to stimulate P450 enzymes in the liver, which are responsible for metabolising exogenous chemical compounds. Several recent studies have reported that concurrent intake of onion, with a high quercetin content, markedly decreases the oral bioavailability of cyclosporin,

with subsequent organ transplant rejection (Yang et al., 2006). Studies to determine the influence of quercetin coadministration on plasma concentrations of the protease inhibitor saquinavir (antiviral HIV drug) have also been carried out. However, more studies are needed to determine if saquinavir intracellular concentrations are altered by coadministration of quercetin, because of substantial interand intrasubject variability (DiCenzo et al., 2006).

The intake of onion and their derivatives should be controlled, including ingested dose, long-term medication and safety and effectiveness of chosen preparation, with the purpose of minimizing the risk of adverse side effects (Davis, 2005). In spite of this, onion use as therapeutic agent seems to be very safe, since all adverse effects previously described appear with an excessive and prolonged consumption.

FACTORS DECREASING HEALTH BENEFITS OF ONIONS

Tannins and anthocyanins from the skin of red onion have been reported to have antioxidant activity (Augusti, 1996), but in one study no appreciable amounts remained in the edible portion once the outer skin had been removed (Rhodes and Price, 1996). However, this is not true for all varieties, with some still containing appreciable amounts. In peeled Tropea Red onions the edible portion contained only 27% of the anthocyanins, although 79% of the flavonols. Quercetin content is highest in the dry skin and decreases from the outer to inner rings (Patil and Pike, 1995). Thus, peeling may significantly reduce the flavonoid content (especially anthocyanins and to a lesser extent flavonols) and hence some of the health benefits of onions.

In contrast, fructans are richest in the fleshy layers (Jaime et al., 2001b), as are sulfur compounds. Chopping may also affect the phytochemical content. As mentioned above, many of the sulfur compounds that have health benefits are not formed until the onion tissue is chopped.

However, if left too long these compounds can be changed further and lose activity. Rhodes and Price, (1997)(Rhodes and Price, 1996) showed that quercetin 3,4'-diglucoside was rapidly degraded in macerated tissues (50% decline after 5 hours), being converted to the quercetin monoglycoside and free quercetin. All these compounds have antioxidant activity so this feature of onions might not be affected by chopping. In a different study (Makris and Rossiter, 2001), chopping was shown to have no significant effect on flavonol content or antioxidant activity. Ewald et al. (1999) showed the greatest loss of flavonoids in onion occurred during the pre-processing step when the onion was peeled, trimmed and chopped before blanching.

Ioku et al. (2001) measured the effects of various cooking methods on the flavonoid content in onion. Microwave cooking without water retained both flavonoids and ascorbic acid. Frying did not affect flavonoid intake while boiling of onions led to about a 30% loss of quercetin glycosides, which transfers to the boiling water (flavonoids are water-soluble). (Crozier et al., 1997) also examined the effects of cooking on onions and found boiling reduced flavonoid content significantly, while microwaving had slightly less of an effect and frying resulted in the lowest loss. Makris and Rossiter (2001) showed a flavonol loss of 20% on boiling and antioxidant activity also decreased.

Adam et al. (2000) and Benítez et al. (2011) examined quality changes in onion during drying. The results showed that drying temperatures above 65°C exerted a pronounced influence on colour. The pyruvate content decreased with increasing temperature and slice thickness. The sugar content was also significantly influenced by the drying temperature. The rate of ascorbic acid degradation decreased with increasing temperature and slice thickness. Because carotenoids present in spring onions are fat-soluble, they are best absorbed in the body if accompanied by

some form of oil or fat in the meal. Chopping and cooking assists in releasing carotenoids from the food matrix and this also increases their bioavailability.

CONSULSION

Many health benefits have been reported for onion. Extract of this spice exhibited a significant antibiotic activity. That could be a replacer of modern antibiotics. Antiasthmatic, antidiabetic and a weak antiplatelet activity were proved for *A. cepa*. Cardiovascular benefit of this excellent medicinal plant is well known since old days. 40 to 100g of *Allium cepa* is the recommended intake and it must be incorporated in daily diet in order to compete with onset of different diseases.

PICTURE CAPTION

Figure 1: Classification of *Allium cepa*

Figure 2: Flavonoids, Quercetin and quercetin glucosides structures present in *Allium cepa*

Figure 3: Formation of different organo-sulphur compounds in processed onions

TABLE CAPION

Table 1: Proximate analysis, vitamin contents and minerals present in 100g of raw onion

Table 2: Lipids and amino acids present in 100 g of raw onions

Table 3: Major functions of Onion (*Allium cepa*) in human body

Table 4: Overview of cancer prevention by *Allium cepa*

Table 5: Overview of antibacterial, antifungal and antiparasitic activity of onions

REFERENCES

- Ackermann, R. T., Mulrow, C. D., Ramirez, G., Gardner, C. D., Morbidoni, L., and Lawrence, V. A. (2001). Garlic shows promise for improving some cardiovascular risk factors. *Arch Int Med.* **161**: 813-824.
- Adam, E., Muhlbauer, W., Esper, A., Wolf, W., and Spiess, W. (2000). Quality changes of onion (*Allium cepa* L.) as affected by the drying process. *Die Nahrung.* **44**: 32-37.
- Ali, M., Bordia, T., and Mustafa, T. (1999). Effect of raw versus boiled aqueous extract of garlic and onion on platelet aggregation. *Prostaglandins Leukot Essent Fatty Acids.* **60**: 43-47.
- Ali, M., Thomson, M., and Afzal, M. (2000). Garlic and onions: their effect on eicosanoid metabolism and its clinical relevance. *Prostaglandins Leukot Essent Fatty Acids.* **62**: 55-73.
- Ashraf, M. Z., Hussain, M. E., and Fahim, M. (2005). Antiatherosclerotic effects of dietary activities of onion (*Allium cepa* L.) oil. *Microbiol Res.* **150**: 167-172.
- Augusti, K. (1996). Therapeutic values of onion and garlic. *Indian J Exp Biol.* **34**: 634-640.

- Babu, P. S., and Srinivasan, K. (1999). Renal lesions in streptozotocin-induced diabetic rats maintained on onion and capsaicin containing diets. *J Nutr Biochem.* **10**: 477-483.
- Bakri, I. M., and Douglas, C. W. I. (2005). Inhibitory effect of garlic extract on oral bacteria. *Arc Oral Biol.* **50**: 645-651.
- Banerjee, S. K., Mukherjee, P. K., and Maulik, S. K. (2003). Garlic as an antioxidant: the good, the bad and the ugly. *Phytother Res.* **17**: 97-106.
- Beatty, S., Murray, I. J., Henson, D. B., Carden, D., Koh, H. H., and Boulton, B. B. (2001). Macular pigment and risk for age-related macular degeneration in subjects from a Northern European population. *Invest Ophthalmol Visual Sci.* **42**: 439-446.
- Benítez, V., Mollá, E., María, A., Martín-Cabrejas, Aguilera, Y., Francisco, J., López-Andréu, and Esteban, R. M. (2011). Effect of sterilisation on dietary fibre and physicochemical properties of onion by-products. *Food Chem.* **127**: 501-507.
- Benkeblia, N. (2005). Free-radical scavenging capacity and antioxidant properties of some selected onions (*Allium cepa* L.) and garlic (*Allium sativum* L.) extracts. *Brazilian Arch Biol Technol.* **48**: 753-759.

Bianchini, F., and Vainio, H. (2001). Allium vegetables and organosulfur compounds: Do they help prevent cancer? *Environ Health Pers* **109**: 893-902.

Bilyk, A., Cooper, P. L., and Saper, G. M. (1984). Varietal difference in distribution of quercetin and kaempferol in onion (*Allium cepa* L.). *J Agric Food Chem.* **32**: 274-276.

Block, E. (1996). Recent results in the organosulphur and organoselenium chemistry of genus *Allium* and Brassica plants. Relevance for cancer prevention. *Adv Exp Med Bio.* **401**: 155-169.

Block, E., Birringer, M., Jiang, W., Nakahodo, T., Thompson, H. J., and Toscano, H. e. a. (2001). Chemistry of analogous organoselenium and organosulfur compounds. *J Agric Food Chem.* **49**: 458.

Block, E., Gulati, H., Putman, D., Sha, D. Y., You, N. N., and Zhao, S. H. (1997). Allium chemistry: Synthesis of 1-[alk(en)ylsulfinyl]propyl alk(en)yl disulfides (cepaenes), antithrombotic flavorants from homogenates of onion (*Allium cepa*). *J Agric Food Chem.* **45**: 4412-4422.

Block, E., Naganathan, S., Putman, D., and Zhao, S. H. (1993). Organosulfur chemistry of garlic and onion: Recent results. *Pure Appl Chem.* **65**: 625-632.

- Boeing, H., Jedrychowski, W., Wahrendorf, J., Popiela, T., Tobiaszadamczyk, B., and Kulig, A. (1991). Dietary risk-factors in intestinal and diffuse types of stomach-cancer - a multicenter case-control study in Poland. *Cancer Causes Con.* **2**: 227-233.
- Bonaccorsi, P., Caristi, C., Gargiulli, C., and Leuzzi, U. (2008). Flavonol glucosides in *Allium* species: A comparative study by means of HPLC-DAD-ESI-MS-MS. *Food Chem.* **107**: 1668-1669.
- Bone, R. A., Landrum, J. T., Dixon, Z., Chen, Y. M., and Llerena, C. M. (2000). Lutein and zeaxanthin in the eyes, serum and diet of human subjects. *Exp Eye Res.* **71**: 239-245.
- Bordia, T., Mohammed, N., Thomson, M., and Ali, M. (1996). An evaluation of garlic and onion as antithrombotic agents. *Prostaglandins Leukot Essent Fatty Acids.* **54**: 183-186.
- Borek, C. (1997). Antioxidants and cancer. *Sci Med.* **4**: 51-62.
- Brankovic, S., Radenkovic, M., Kitic, D., Veljkovic, S., Ivetic, V., Pavlovic, D., and Miladinovic, B. (2011). Comparison of the Hypotensive and Bradycardic Activity of Ginkgo, Garlic, and Onion Extracts. *Clin Exp Hyp.* **33**: 95-99.
- Briggs, W. H., Folts, J. D., Osman, H. E., and Goldman, I. L. (2001). Administration of raw onion inhibits platelet-mediated thrombosis in dogs. *J Nutr.* **131**: 2619-2622.

- Brisdelli, F., Coccia, C., Cinque, B., Cifone, M. G., and Bozzi, A. (2007). Induction of apoptosis by quercetin: different response of human chronic myeloid (K562) and acute lymphoblastic (HSB-2) leukemia cells. *Mol Cell Biochem.* **296**: 137-149.
- Butt, M. S., Sultan, M., Butt, M. S., and Iqbal, J. (2009). Garlic: Nature's Protection Against Physiological Threats. *Crit Rev Food Sci Nutr.* **49**: 538-551.
- Butt, M. S., Sultan, M. T., Anjum, F. M., Sharif, M. K., and Khan, S. H. (2007). Functional foods: emerging trend in nutritional support programs. **In:** International Symposium on Emerging Trends in Food Science & Technology. National Institute of Food Science and Technology, University of Agriculture, Faisalabad, Pakistan.
- Caridi, D., Trenerry, V. C., Rochfort, S., Duong, S., Laughler, D., and Jones, R. (2007). Profiling and quantifying quercetin glucosides in onion (*Allium cepa* L.) varieties using capillary zone electrophoresis and high performance liquid chromatography. *Food Chem.* **105**: 691-699.
- Cazzola, R., Camerotto, C., and Cestaro, B. (2011). Anti-oxidant, anti-glycant, and inhibitory activity against alpha-amylase and alpha-glucosidase of selected spices and culinary herbs. *Int J Food Sci Nutr.* **62**: 175-184.

- Challier, B., Perarnau, J. M., and Viel, J. F. (1998). Garlic, onion and cereal fibre as protective factors for breast cancer: a French case-control study. *Eur J Epidemiol.* **14**: 737-747.
- Chen, C.-H., Chou, T.-W., Cheng, L.-H., and Ho, C.-W. (2011). In vitro anti-adenoviral activity of five *Allium* plants. *J Taiwan Inst Chem Eng.* **42**: 228-232.
- Chisty, M., Quddus, R., Islam, B., and Khan, B. (1996). Effect of onion extract on immune response in rabbits. *Bangladesh Med Rese Council Bull.* **22**: 81-85.
- Chu, Y. F., Sun, J., Wu, X., and Liu, R. H. (2002). Antioxidant and antiproliferative activities of common vegetables. *J Agric Food Chem.* **50**: 6910-6916.
- Chutani, S. K., and Bordia, A. (1981). The effect of fried vs. raw garlic on Iibrinolytic activity in man. *Atherosclerosis.* **38**: 417-421.
- Corea, G., Fattorusso, E., Lanzotti, V., Capasso, R., and Izzo, A. A. (2005). Antispasmodic saponins from bulbs of red onion, *Allium cepa* L. var. Tropea. *J Agri Food Chem.* **53**: 935-940.
- Corzo-Martinez, M., Corzo, N., and Villamiel, M. (2007). Biological properties of onions and garlic. *Trends Food Sci Technol.* **18**: 609-625.

- Crozier, A., Lean, M. E. J., McDonald, M. S., and Black, C. (1997). Quantitative analysis of the flavonoid content of commercial tomatoes, onions, lettuce, and celery. *J Agric Food Chem.* **45**: 590-595.
- Dankert, J., Tromp, T. F., de Vries, H., and Klasen, H. J. (1979). Antimicrobial activity of crude juices of *Allium ascalonicum*, *Allium cepa* and *Allium sativum*. Zentralblatt für Bakteriologie, Parasitenkunde, Infektionskrankheiten und Hygiene, 1,. *Originale A* **245**: 229-239.
- Davis, S. R. (2005). An overview on the antifungal properties of allicin and its breakdown products e the possibility of a safe and effective antifungal properties. *Mycoses*: 2.
- de Vries, J., Hollman, P., Meyboom, S., Buysman, M., Zock, P., Van Staveren, W., and Katan, M. (1998). Plasma concentrations and urinary excretion of the antioxidant flavonoids quercetin and kaempferol as biomarkers for dietary intake. *Am J Clin Nutr.* **68**: 60-65.
- Debaene, J. E. P., Goldman, I. L., and Yandell, B. S. (1999). Postharvest flux and genotype environment effects for onion-induced antiplatelet activity, pungency, and soluble solids in long-day onion during postharvest cold storage. *J Am Soc Hort Sci.* **124**: 366-372.
- DiCenzo, R., Frerichs, V., Larppanichpoonphol, P., Predko, L., Chen, A., and Reichman, R. e. a. (2006). Effect of quercetin on the plasma and intracellular concentrations of saquinavir in healthy adults. *Pharmacother.* **26**: 1255-1261.

- Donner, H., Gao, L., and Mazza, G. (1997). Separation and characterization of simple and malonylated anthocyanins in red onions, *Allium cepa* L. *Food Res Int.* **30**: 637-643.
- Dorant, E., Van Den Brandt, P., and Goldbohm, R. A. (1996). A prospective cohort study on the relationship between onion and leek consumption, garlic supplement use and the risk of colorectal carcinoma in The Netherlands. *Carcinogenesis.* **17**: 477-484.
- Dorsch, W. (1996). *Allium cepa* L. (onion): Part 2. Chemistry, analysis and pharmacology. *Phytomedicine.* **3**: 391-397.
- Dorsch, W., Adelman-Grill, B., Bayer, T., Ettl, M., Hein, G., Jaggy, H., Ring, J., Scheftner, P., Schneider, T., and Wagner, H. (1987a). Zwiebelextrakte als Asthma-Therapeutika? *Allergologie* **10**: 316-326.
- Dorsch, W., Ettl, M., Hein, G., Scheftner, P., Weber, J., Bayer, T., and Wagner, H. (1987b). Antiasthmatic effects of onions. Inhibition of platelet-activating factor-induced bronchial obstruction by onion oils. *Int Arch Aller App Immunol.* **82**: 535-536.
- Dorsch, W., and Wagner, H. (1991). New antiasthmatic drugs from traditional medicine? . *Int Arch Aller App Immunol.* **94**: 262-265.

Duthie, S. J., and Dobson, V. L. (1998). Dietary flavanoids protect human colonocyte DNA from oxidative attack in vitro. *Eur J Nutr.* **38**: 28-34.

Effendy, J. L., Simmons, D. L., Campbell, G. R., and Campbell, J. H. (1997). The effect of aged garlic extract “Kyolic”, on the development of experimental atherosclerosis. *Atherosclerosis.* **132**: 37-42.

El-Bayoumy, K., Chae, Y. H., Upadhyaya, P., and Ip, C. (1996). Chemoprevention of mammary cancer by diallyl selenide, a novel organoselenium compound. *Anticancer Res.* **16**: 2911-2918.

El-Bayoumy, K., Sinka, R., Pinto, J. T., and Rivlin, R. S. (2006). Cancer chemoprevention by garlic and garlic-containing sulfur and selenium compounds. *J Nutr.* **136**: 864-869.

El-Demerdash, F. M., Yousef, M. I., and Abou El-Naga, N. I. (2005). Biochemical study on the hypoglycemic effects of onion and garlic in alloxan-induced diabetic rats. *Food Chem Toxicol.* **43**: 57-63.

Elnima, E. I., Ahmed, S. A., Mekkawi, A. G., and Mossa, J. S. (1983). The antimicrobial activity of garlic and onion extracts. *Die Pharmazie.* **38**: 747-748.

Ernst, M., and Feldheim, W. (2000). Fructans in higher plants and in human nutrition. *J App Bot.* **74**: 5-9.

Ewald, C., Fjellkner-Modig, S., Johansson, K., Sjöholm, I., and Akesson, B. (1999). Effect of processing on major flavonoids in processed onions, green beans, and peas. *Food Chem.* **64**: 231-235.

FAO (2011). Onions: Post-Harvest Operation. Food and Agriculture Organization of the United Nations.

FAO Statistics (2011). Productions, crops.

Fenwick, G. R., and Hanley, A. B. (1985). The genus *Allium*. Part 1. *Crit Rev Food Sci Nutr.* **22**: 199-271.

Formica, J. V., and Regelson, W. (1995). Review of the biology of quercetin and related bioflavonoids. *Food Chem Toxicol.* **33**: 1061-1080.

Fossen, T., Andersen, O. M., Ovstedal, D. O., Pedersen, A. T., and Raknes, A. (1996). Characteristic anthocyanin pattern from onions and other *Allium* spp. *J Food Sci.* **61**: 4.

Friedman, T., Shalom, A., and Westreich, M. (2006). Self-inflicted garlic burns: our experience and literature review. *Int J Dermatol.* **45**: 1161-1163.

Fukushima, S., Takada, N., Hori, T., and Wanibuchi, H. (1997). Cancer prevention by organosulfur compounds from garlic and onion. *J Cell Biochem. (Suppl)***27**: 100-105.

Gabler, N. K., Osrowska, E., Imsic, M., Eagling, D. R., Jois, M., Tatham, B. G., and Dunshea, F. R. (2006). Dietary onion intake as part of a typical high fat diet improves indices of cardiovascular health using the mixed sex pig model. *Plant Foods Human Nutr.* **61**: 179-185.

Galeone, C., Pelucchi, C., Levi, F., Negri, E., Franceschi, S., Talamini, R., Giacosa, A., and La Vecchia, C. (2006). Onion and garlic use and human cancer. *Am J Clin Nutr.* **84**: 1027-1032.

Galeone, C., Tavani, A., Pelucchi, C., Negri, E., and La Vecchia, C. (2009). Allium vegetable intake and risk of acute myocardial infarction in Italy. *Eur J Nutr.* **48**: 120-123.

Gao, C. M., Takezaki, T., Ding, J. H., Li, M. S., and Tajima, K. (1999). Protective effect of allium vegetables against both esophageal and stomach cancer: A simultaneous casereferent study of a high-epidemic area in Jiangsu province, China. *Jpn J Cancer Res.* **90**: 614-621.

Glasser, G., Graefe, E. U., Struck, F., Veit, M., and Gebhardt, R. (2002). Comparison of antioxidative capacities and inhibitory effects on cholesterol biosynthesis of quercetin and potential metabolites. *Phytomedicine*. **9**: 33-40.

Goldman, I. L., Kopelberg, M., Debaene, J. E. P., and Schwartz, B. S. (1996). Antiplatelet activity in onion (*Allium cepa*) is sulfur dependent. *Thrombosis Haemostasis*. **76**: 450-452.

Gonzalez, C. A., Pera, G., Agudo, A., Bueno-De-Mesquita, H. B., Ceroti, M., Boeing, H., Schulz, M., Del Giudice, G., Plebani, M., Carneiro, F., Berrino, F., Sacerdotde, C., Tumino, R., Panico, S., Berglund, G., Siman, H., Hallmans, G., Stenling, R., Martinez, C., Dorronsoro, M., Barricarte, A., Navarro, C., Quiros, J. R., Allen, N., Key, T. J., Bingham, S., Day, N. E., Linseisen, J., Nagel, G., Overvad, K., Jensen, M. K., Olsen, A., Tjonneland, A., Buchner, F. L., Peeters, P. H., Numans, M. E., Clavel-Chapelon, F., Boutron-Ruault, M. C., Roukos, D., Trichopolou, A., Psaltopoulou, T., Lund, E., Casagrande, C., Slimani, N., Jenab, M., and Riboli, E. (2006). Fruit and vegetable intake and the risk of stomach and oesophagus adenocarcinoma in the European Prospective Investigation into Cancer and Nutrition (EPIC-EURGAST). *Int J Cancer*. **118**: 2559-2566.

Gonzalez, C. A., Sanz, J. M., Marcos, G., Pita, S., Brullet, E., Saigi, E., Badia, A., and Riboli, E.

(1991). Dietary factors and stomach-cancer in Spain - a multicenter case-control study.

Int J Cancer. **49**: 513-519.

Gonzalez, C. A., Travier, N., Lujan-Barroso, L., Castellsague, X., Xavier Bosch, F., Roura, E.,

Bueno-de-Mesquita, H. B., Palli, D., Boeing, H., Pala, V., Sacerdote, C., Tumino, R.,

Panico, S., Manjer, J., Dillner, J., Hallmans, G., Kjellberg, L., Sanchez, M.-J., Altzibar, J.

M., Barricarte, A., Navarro, C., Rodriguez, L., Allen, N., Key, T. J., Kaaks, R.,

Rohrmann, S., Overvad, K., Olsen, A., Tjonneland, A., Munk, C., Kjaer, S. K., Peeters,

P. H. M., van Duijnhoven, F. J. B., Clavel-Chapelon, F., Boutron-Ruault, M.-C.,

Trichopoulou, A., Benetou, V., Naska, A., Lund, E., Engeset, D., Skeie, G., Franceschi,

S., Slimani, N., Rinaldi, S., and Riboli, E. (2011). Dietary factors and in situ and invasive

cervical cancer risk in the European prospective investigation into cancer and nutrition

study. *Int J Cancer*. **129**: 449-459.

Gorinstein, S., Leontowicz, H., Leontowicz, M., Najman, K., Biecki, W., Ham, K.-S., Kang, S.-

G., Paredes-Lopez, O., Leticia Martinez-Ayala, A., and Trakhtenberg, S. (2011). Aorta

and Liver Changes in Rats Fed Cholesterol-Containing and Raw Vegetable-

Supplemented Diets: Experiments in Vitro and in Vivo. *J Agric Food Chem*. **59**: 7441-

7451.

Griffiths, G., Trueman, L., Crowther, T., Thomas, B., and Smith, B. (2002). Onions-a global benefit to health. *Phytother Res.* **16**: 603-615.

Guñ Isen, A., Makris, D. P., and Kefalas, P. (2007). Biomimetic oxidation of quercetin: isolation of a naturally occurring quercetin heterodimer and evaluation of its in vitro antioxidant properties. *Food Res Int.* **40**: 7-14.

Gupta, N., and Porter, T. D. (2001). Garlic and garlic-derived compounds inhibit human squalene monooxygenase. *J Nutr.* **131**: 1662-1667.

Halvorsen, B. L., Holte, K., Myhrstad, M. C. W., Barikmo, I., Hvattum, E., Remberg, S. F., Wold, A.-B., Haffner, K., Baugerod, H., Andersen, L. F., Moskaug, J. O., Jacobs, D. R., and Blomhoff, R. (2002). A systematic screening of total antioxidants in dietary plants. *J Nutr.* **132**: 461-471.

Hansson, L. E., Nyren, O., Bergstrom, R., Wolk, A., Lindgren, A., Baron, J., and Adami, H. O. (1993). Diet and risk of gastric-cancer - a population-based case-control study in Sweden. *Int J Cancer.* **55**: 181-189.

Hatono, S., Jimenez, A., and Wargovich, M. J. (1996). Chemopreventive effect of S-allylcysteine and its relationship to the detoxification enzyme glutathione S-transferase. *Carcinogenesis* **17**: 1041-1044.

- Hertog, M., and Katan, M. (1998). Quercetin in foods, cardiovascular disease, and cancer. **In:** Flavonoids in health and disease, pp. 447-467. Marcel Dekker Inc., New York, USA.
- Hsing, A. W., Chokkalingam, A. P., Gao, Y. T., Madigan, M. P., Deng, J., Gridley, G., and Fraumeni, J. F. J. (2002). Allium vegetables and risk of prostate cancer, a population-based study. *J Nat Cancer Inst.* **94**: 1648-1651.
- Hu, J. F., La Vecchia, C., Negri, E., Chatenoud, L., Bosetti, C., Jia, X. Y., Liu, R. Z., Huang, G. R., Bi, D. Z., and Wang, C. X. (1999). Diet and brain cancer in adults: A case-control study in northeast China. *Int J Cancer.* **81**: 20-23.
- Hubbard, G. P., Wolfram, S., de Vos, R., Bovy, A., Gibbins, J. M., and Lovegrove, J. A. (2006). Ingestion of onion soup high in quercetin inhibits platelet aggregation and essential components of the collagen-stimulated platelet activation pathway in man: a pilot study. *Br J Nutr.* **96**: 482-488.
- Ikken, Y., Morales, P., Martinez, A., Marin, M. L., Haza, A. I., and Cambero, M. I. (1999). Antimutagenic effect of fruit and vegetable ethanolic extracts against N-nitrosamines evaluated by the Ames test. *J Agric Food Chem.* **47**: 3257-3264.
- Ioku, K., Aoyama, Y., Tokuno, A., Terao, J., Nakatani, N., and Takei, Y. (2001). Various cooking methods and the flavonoid content in onion. *J Nutr Sci Vitaminol.* **47**: 78-83.

- Iscovich, J. M., Labbe, K. A., Castelleto, R., Calzona, A., Bernedo, A., Chopita, N. A., Jmelnitzsky, A., and Kaldor, J. (1992). Colon cancer in Argentina I. Risk from intake of dietary items. *Int J Cancer*. **51**: 851-857.
- Jaime, L., Martin-Cabrejas, M. A., Molla, E., Lopez-Andreu, F. J., and Esteban, R. M. (2001a). Effect of storage on fructan and fructooligosaccharide of onion (*Allium cepa* L.) *J Agric Food Chem*. **49**: 982-988.
- Jaime, L., Martínez, F., Martín-Cabrejas, M. A., Mollá, E., López-Andréu, F. J., Waldron, K. W., and Esteban, R. M. (2001b). Study of total fructan and fructooligosaccharide content in different onion tissues. *J Sci Food Agric*. **81**: 177-182.
- Jaime, L., Mollá, E., Fernandez, A., Martín-Cabrejas, M. A., López-Andréu, F. J., and Esteban, R. M. (2002). Structural carbohydrate differences and potential source of dietary fiber of onion (*Allium cepa* L.) tissues. *J Agric Food Chem*. **50**: 122-128.
- Judprasong, K., Tanjor, S., Puwastien, P., and Sungpuag, P. (2011). Investigation of Thai plants for potential sources of inulin-type fructans. *J Food Comp Anal*. **24**: 642-649.

- Jung, Y. S., Kim, M. H., Lee, S. H., Baik, E. J., Park, S. W., and Moon, C. H. (2002). Antithrombotic effect of onion in streptozotocin-induced diabetic rat. *Prostaglandins Leukot Essent Fatty Acids*. **66**: 453-458.
- Kahane, R., Vialle-Guérin, E., Boukema, I., Tzanoudakis, D., Bellamy, C., Chamaux, C., and Kik, C. (2001). Changes in non-structural carbohydrate composition during bulbing in sweet and high-solid onions in field experiments. *Environ Exp Bot*. **45**: 73-83.
- Kannel, W. B. (2005). Overview of hemostatic factors involved in atherosclerotic cardiovascular disease. *Lipids*. **40**: 1215-1220.
- Kawaii, S., Tomono, Y., Katase, E., Ogawa, K., and Yano, M. (1999). Antiproliferative activity of flavonoids on several cancer cell lines. *Biosci Biotechnol Biochem*. **63**: 896-899.
- Kendler, B. S. (1987). Garlic (*allium-sativum*) and Onion (*allium-cepa*) - a review of their relationship to cardiovascular-disease. *Prevent Med*. **16**: 670-685.
- Kim, J. H. (1997). Anti-bacterial action of onion (*Allium cepa* L.) extracts against oral pathogenic bacteria. *J Nihon Uni Sch Dent*. **39**: 136-141.
- Kim, W. J., Lee, K. A., Kim, K.-T., Chung, M.-S., Cho, S., and Paik, H.-D. (2011). Antimicrobial Effects of Onion (*Allium cepa* L.) Peel Extracts Produced via Subcritical

Water Extraction against *Bacillus cereus* Strains as Compared with Ethanolic and Hot Water Extraction. *Food Sci Biotechnol.* **20**: 1101-1106.

Knowles, L. M., and Milner, J. A. (2003). Diallyl disulphide induces ERK phosphorylation and alters gene expression profiles in human colon tumor cells. *The J Nutr.* **133**: 2901-2906.

Koncic, M. Z., and Jug, M. (2011). Antioxidant and Bioadhesive Properties of Onions (*Allium* L., Alliaceae) Processed Under Acidic Conditions. *Inter J Food Prop.* **14**: 92-101.

Krishnaswamy, K., and Polasa, K. (1995). Diet, nutrition and cancer – the Indian scenario. *Indian J Med Res.* **102**: 200-209.

Kumari, K., and Augusti, K. T. (2007). Lipid lowering effect of S-methyl cysteine sulfoxide from *Allium cepa* Linn in high cholesterol diet fed rats. *J Ethnopharmacol.* **109**: 367-371.

Kumari, K., Mathew, B. C., and Augusti, K. T. (1995). Antidiabetic and hypolipidemic effects of S-methyl cysteine sulfoxide isolated from *Allium cepa* Linn. *Indian J Biochem Biophy.* **32**: 49-54.

Kunicka-Styczynska, A. (2011). Activity of essential oils against food-spoiling yeast. A review. *Flavour Fragrance J.* **26**: 326-328.

- Kyung, K. H. (2011). Antimicrobial properties of allium species. *Curr Opin Biotechnol.* **23**: 1-6.
- Lai, P. K., and Roy, J. (2004). Antimicrobial and chemopreventive properties of herbs and spices. *Curr Med Chem.* **11**: 1451-1460.
- Lam, Y. W., Wang, H. X., and Ng, T. B. (2000). A robust cysteine deficient chitinase-like antifungal protein from inner shoots of the edible chive *Allium tuberosum*. *Biochem Biophys Res Comm.* **279**: 74-80.
- Langos, M., Hofstetter, W., Dolder, S., Felix, R., Muhlbauer, R. C., and Brenneisen, R. (2007). A gamma-glutamyl peptide from onion inhibits the development and activity of osteoclasts in vitro. *Calcified Tissue Int.* **80**: 91-92.
- Lanzotti, V. (2006). The analysis of onion and garlic. *J Chromat A.* **1112**: 3-22.
- Lautraite, S., Musonda, A. C., Doehmer, J., Edwards, G. O., and Chipman, J. K. (2002). Flavonoids inhibit genetic toxicity produced by carcinogens in cells expressing CYP1A2 and CYP1A1. *Mutagenesis.* **17**: 45-53.
- Le Marchand, L., Murphy, S. P., Hankin, J. H., Wilkens, L. R., and Kolonel, L. N. (2000). Intake of flavonoids and lung cancer. *J Nat Cancer Inst.* **92**: 154-160.

- Lean, M., Noroozi, M., I, K., Burns, J., Talwar, D., Satter, N., and Crozier, A. (1999). Dietary flavonoids protect diabetic human lymphocytes against oxidant damage to DNA. *Diabetes*. **48**: 176-181.
- Lee, S., Kim, J. N., Choung, D.-H., and Lee, H. K. (2011). Facile Synthesis of trans-S-1-Propenyl-L-Cysteine Sulfoxide (Isoalliin) in Onions (*Allium cepa*). *Bull Korean Chem Soc*. **32**: 319-320.
- Lee, S. U., Lee, J. H., Choi, S. H., Lee, J. S., Ohnisi-Kameyama, M., Kozukue, N., Levin, C. E., and Friedman, M. (2008). Flavonoid content in fresh, home-processed, and light-exposed onions and in dehydrated commercial onion products. *J Agric Food Chem*. **56**: 8541-8548.
- Lotito, S. B., and Frei, B. (2006). Consumption of flavonoid-rich foods and increased plasma antioxidant capacity in humans: cause, consequence, or epiphenomenon? . *Free Rad Biol Med*. **41**: 1727-1746.
- Lun, Z. R., Burri, C., Menzinger, M., and Kaminsky, R. (1994). Antiparasitic activity of diallyl trisulfide (Dasuansu) on human and animal pathogenic protozoa (*Trypanosoma* sp., *Entamoeba histolytica* and *Giardia lamblia*) in vitro. *Annales de la Societe Belge de Medecine Tropicale*. **74**: 51-59.

- Ly, T. N., Hazama, C., Shimoyamada, M., Ando, H., Kato, K., and Yamauchi, R. (2005). Antioxidative compounds from the outer scales of onion. *J Agric Food Chem.* **53**: 8183-8189.
- Makheja, A. N., and Bailey, J. M. (1990). Antiplatelet constituents of garlic and onion. *Agents and Actions.* **29**: 360-363.
- Makris, D. P., and Rossiter, J. T. (2001). Domestic processing of onion bulbs (*Allium cepa*) and asparagus spears (*Asparagus officinalis*), Effect on flavonol content and antioxidant status. *J Agric Food Chem.* **49**: 3216-3222.
- Mantawy, M. M., Ali, H. F., and Rizk, M. Z. (2011). THERAPEUTIC EFFECTS OF *Allium sativum* AND *Allium cepa* IN *Schistosoma mansoni* EXPERIMENTAL INFECTION. *Revista Do Instituto De Medicina Tropical De Sao Paulo.* **53**: 155-163.
- Mares-Perlman, J. A., Millen, A. E., Ficek, T. L., and Hankinson, S. E. (2002). The body of evidence to support a protective role for lutein and zeaxanthin in delaying chronic diseases. Overview. *J Nutr.* **132**: 518S-524S.
- Matheson, E. M., Mainous, A. G., and Carnemolla, M. A. (2009). The association between onion consumption and bone density in perimenopausal and postmenopausal non-Hispanic white women 50 years and older. . *Menopause.* **16**: 756-759.

Matsuura, H. (1997). Phytochemistry of garlic horticultural and processing procedures. **In:** Neutraceuticals: Designer foods III. Garlic, soy and licorice, pp. 55-59. Lachance, P. A., and Trumbull, C. T. (Eds.).

Matsuura, H., Ushiroguchi, T., Itakura, Y., Hayashi, H., and Fuwa, T. I. (1988). A furostanol glycoside from garlic bulbs of *Allium sativum*. *Chem Pharm Bull.* **36**: 3659-3663.

Millen, A. E., Subar, A. F., Graubard, B. I., Peters, U., Hayes, R. B., Weissfeld, J. L., Yokochi, L. A., Ziegler, R. G., and for the PLCO Cancer Screening Project Team (2007). Fruit and vegetable intake and prevalence of colorectal adenoma in a cancer screening trial. *Am J Clin Nutr.* **86**: 1754-1764.

Mohamed, E. F. (2010). Antiviral properties of garlic cloves juice compared with onion bulbs juice against potato virus Y (PVY). *J Am Sci.* **6**: 302-310.

Moon, C. H., Jung, Y. S., Kim, M. H., Lee, S. H., Baik, E. J., and Park, S. W. (2000). Mechanism for antiplatelet effect of onion: AA release inhibition, thromboxane A2 synthase inhibition and TXA2/PGH2 receptor blockade. *Prostaglandins Leukot Essent Fatty Acids.* **62**: 277-283.

Muhlbauer, R. C., Lozano, A., and Reinli, A. (2002). Onion and a mixture of vegetables, salads, and herbs affect bone resorption in the rat by a mechanism independent of their base excess. *J Bone Mineral Res.* **17**: 1230-1236.

Munday, R., and Munday, C. M. (2001). Relative activities of organosulfur compounds derived from onions and garlic in increasing tissue activities of quinone reductase and glutathione transferase in rat tissues. *Nutr Cancer Int J.* **40**: 205-210.

Munday, R., and Munday, C. M. (2004). Induction of phase II enzymes by aliphatic sulfides derived from garlic and onions: An overview. **In:** Quinones and Quinone Enzymes, pp. 449-456. Academic Press Inc, San Diego.

Nadia, M. T., El Ezz, A., Khalil, F. A. M., and Shapaan, M. R. (2011). Therapeutic Effect of Onion (*Allium cepa*) and Cinnamon (*Cinnamomum zeylanicum*) Oils on Cryptosporidiosis in Experimentally Infected Mice. *Global Veterinaria.* **7**: 179-183.

National Onion Association (2011). All about Onions. National Onion Association.

Nishimura, H., Higuchi, O., Tateshita, K., Tomobe, K., Okuma, Y., and Nomura, Y. (2006). Antioxidative activity and ameliorative effects of memory impairment of sulfurcontaining compounds in *Allium* species. *Biofactors.* **26**: 135-146.

Ostrowska, E., Gabler, N. K., Sterling, S. J., Tatham, B. G., Jones, R. B., Eagling, D. R., Jois, M., and Dunshea, F. R. (2004). Consumption of brown onions (*Allium cepa* var. cavalier and var. density) moderately modulates blood lipids, haematological and haemostatic variables in healthy pigs. . *Br J Nutr.* **91**: 211-218.

Patil, B. S., and Pike, L. M. (1995). Distribution of quercetin content in different rings of various coloured onion (*Allium cepa* L.) cultivars. *J Hort Sci.* **70**: 643-650.

Pellegrini, N., Serafini, M., Colombi, B., Del Rio, D., Salvatore, S., Bianchi, M., and Brigheni, F. (2003). Total antioxidant capacity of plant foods, beverages and oils consumed in Italy assessed by three different in vitro assays. *J Nutr.* **133**: 2812-2819.

Phay, N., Higashiyama, T., Tsuji, M., Matsuura, H., Fukushi, Y., and Yokota, A. e. a. (1999). An antifungal compound from roots of Welsh onion. *Phytochem.* **52**: 271-274.

Rahman, K., and Lowe, G. M. (2006). Garlic and cardiovascular disease: a critical review. *Journal of Nutrition.* **136**: 736S-740S.

Raso, G. M., Meli, R., Di Carlo, G., Pacilio, M., and Di Carlo, R. (2001). Inhibition of inducible nitric oxide synthase and cyclooxygenase-2 expression by flavonoids in macrophage J774A.1. *Life Sciences.* **68**: 921-931.

Reuter, H. D., Koch, H. P., and Lawson, L. D. (1996). Therapeutic effects and applications of garlic and its preparations. **In:** Garlic, the science and therapeutic application of *Allium sativum* L. and related species pp. 135-213. Koch, H. P., and Lawson, L. D. (Eds.), Williams and Wilkins, Baltimore.

Rhodes, M., and Price, K. (1996). Analytical problems in the study of flavonoid compounds in onions. *Food Chemistry*. **57**: 113-117.

Rice-Evans, C. A., Miller, N. J., Bolwell, P. G., Bramley, P. M., and Pridham, J. B. (1995). The relative antioxidant activities of plant-derived polyphenolic flavonoids. *Free Radical Res.* **22**: 375-383.

Richardson, S. J. (1993). Free radicals in the genesis of Alzheimer's disease. *Ann New York Acad Sci.* **695**: 73-76.

Roberfroid, M. B. (2007). Inulin-Type Fructans: Functional Food Ingredients. *J Nutr.* **137**: 2493-2503.

Robertson, J. M., Donner, A. P., and Trevithick, J. R. (1991). A possible role for vitamin C and E in cataract prevention. *Am J Clin Nutr.* **53**: 346S-351S.

Roman-Ramos, R., Flores-Saenz, J. L., and Alarcon-Aguilar, F. J. (1995). Anti-hyperglycemic effect of some edible plants. *J Ethnopharmacol.* **48**: 25-32.

Rose, P., Whiteman, M., Moore, P. K., and Zhu, Y. Z. (2005). Bioactive Salk(en)yl cysteine sulfoxide metabolites in the genus *Allium*: the chemistry of potential therapeutic agents. *Nat Prod Rep.* **22**: 351-368.

Sakakibara, H., Yoshino, S., Kawai, Y., and Terao, J. (2008). Antidepressant-Like Effect of Onion (*Allium cepa* L.) powder in a rat behavioral model of depression. *Biosci Biotechnol Biochem.* **72**: 94-100.

Saleheen, D., Ali, S. A., and Yasinza, M. M. (2004). Antileishmanial activity of aqueous onion extract in vitro. *Fitoterapia.* **75**: 9-13.

Sance, M. M., Gonzalez, R. E., Soto, V. C., and Galmarini, C. R. (2008). Relationships between antiplatelet activity, dry matter content and flavor in onion cultivars. *J Food Agri Environ.* **6**: 41-46.

Sanderson, J., Mclauchlin, W., and Williamson, G. (1999). Quercetin inhibits hydrogen peroxide-induced oxidization of the rat lens. *Free Rad Biol Med.* **26**: 639-645.

Sankaranarayanan, R., Varghese, C., Duffy, S. W., Padmakumary, G., Day, N. E., and Nair, M. K. (1994). A case-control study of diet and lung-cancer in Kerala, South-India. *Int J Cancer.* **58**: 644-649.

- Santas, J., Almajano, M. P., and Carbo, R. (2010). Antimicrobial and antioxidant activity of crude onion (*Allium cepa* L.) extracts. *Int J Food Sci Technol.* **45**: 403-409.
- Sellappan, S., and Akoh, C. (2002). Flavonoids and antioxidant capacity of Georgia-grown *Vidalia* onions. *J. Agric. Food Chem.* **50**: 5338-5342.
- Shin, S. C., Choi, J. S., and Li, X. (2006). Enhanced bioavailability of tamoxifen after oral administration of tamoxifen with quercetin in rats. *Int J Pharm.* **313**: 144-149.
- Shutenko, Z., Henry, Y., Pinard, E., Seylaz, J., Potier, P., Berthet, F., Girard, P., and Sercombe, R. (1999). Influence of the antioxidant quercetin in vivo on the level of nitric oxide determined by electron paramagnetic resonance in rat brain during global ischemia and reperfusion. *Biochem Pharmacol.* **57**: 199-208.
- Sies, H., and Stahl, W. (2003). Non-nutritive bioactive constituents of plants, lycopene, lutein and zeaxanthin. *Int J Nutr Res.* **73**: 95-100.
- Slimestad, R., Fossen, T., and Vagen, I. M. (2007). Onions: A source of unique dietary flavonoids. *J Agric Food Chem.* **55**: 10067-10080.

Sparg, S. G., Light, M. E., and van Staden, J. (2004). Biological activities and distribution of plant saponins. *J Ethnopharmacol.* **94**: 219-243.

Srinivasan, K. (2005). Plant foods in the management of diabetes mellitus: Spices as beneficial antidiabetic food adjuncts. *Int J Food Sci Nutr.* **56**: 399-414.

Steinmetz, K. A., and Potter, J. D. (1993). Food-group consumption and colon cancer in the adelaide case-control study I Vegetables and fruit. *Int J Cancer.* **53**: 711-719.

Steinmetz, K. A., and Potter, J. D. (1996). Vegetables, fruit, and cancer prevention: a review. *Journal of the American Dietetic Association.* **96**: 1027-1039.

Suresh Babu, P., and Srinivasan, K. (1997). Influence of dietary capaicin and onion on the metabolic abnormalities associated with streptozotocin induced diabetes mellitus. *Mol CellBiochem.* **175**: 49-57.

Suzuki, Y., Masashi, I., Segami, T., and Ito, M. (1998). Anti-ulcer effects of antioxidant, quecetin, alpha-tocopherol, nifedipine and tetracycline in rats. *Jpn J Pharmacol.* **78**: 435-441.

Tansey, M. R., and Appleton, J. A. (1975). nhibition of fungal growth by garlic extract. *I Mycologia.* **67**: 409-413.

- Tattelman, E. M. D. (2005). Health effects of garlic. *American Family Physician*. **72**: 103-106.
- Terahara, N., Yamaguchi, M., and Honda, T. (1994). Malonylated anthocyanins from bulbs of red onion, *Allium cepa* L. *Biosci Biotechnol Biochem*. **58**: 1324-1325.
- Tuyns, A. J., Kaaks, R., and Haelterman, M. (1988). Colorectal-cancer and the consumption of foods - a case-control study in Belgium. *Nutr Cancer Int J*. **11**: 189-204.
- Tuyns, A. J., Kaaks, R., Haelterman, M., and Riboli, E. (1992). Diet and gastric-cancer - A casecontrol study in Belgium. *Int J Cancer*. **51**: 1-6.
- Vagen, I. M., and Slimestad, R., 88:404-411. (2008). Amount of characteristic compounds in 15 cultivars of onion (*Allium cepa* L.) in controlled field trials. *J Sci Food Agric*. **88**: 404-411.
- Van Erk, M. J., Roepman, P., van der Lende, T. R., Stierum, R. H., Aarts, J. M. M. J. G., and van Bladeren, P. J. e. a. (2005). Integrated assessment by multiple gene expression analysis of quercetin bioactivity on anticancer-related mechanisms in colon cancer cells in vitro. *Eur J Nutr*. **44**: 143-156.
- Viry, E., Anwar, A., Kirsch, G., Jacob, C., Diederich, M., and Bagrel, D. (2011). Antiproliferative effect of natural tetrasulfides in human breast cancer cells is mediated

through the inhibition of the cell division cycle 25 phosphatases. *Inter J Oncol.* **38**: 1103-1111.

Wagner, H., Dorsch, H., Bayer, T., Breu, W., and Willer, F. (1990). Antiasthmatic effects of onions, inhibition of 5-lipoxygenase and cyclooxygenase in vitro by thiosulfinates and “Cepaenes.”. *Prostaglandins Leukot Essent Fatty Acids* **39**: 59-62.

Wang, H. X., and Ng, T. B. (2001). Purification of allivin, a novel antifungal protein from bulbs of the round-cloved garlic. *Life Sci.* **70**: 357-365.

Wang, H. X., and Ng, T. B. (2004). Isolation of allicepin, a novel antifungal peptide from onion (*Allium cepa*) bulbs. *J Peptide Sci.* **10**: 173-177.

Wang, T. J., Gona, P., Larson, M. G., Tofler, G. H., Levy, D., Newton-Cheh, C., Jacques, P. F., Rifai, N., Selhub, J., Robins, S. J., Benjamin, E. J., D'Agostino, R. B., and Vasan, R. S. (2006). Multiple biomarkers for the prediction of first major cardiovascular events and death. *New Eng J Med.* **355**: 2631-2639.

Wargovich, M. J., and Uda, N. (1996). Allium vegetables and the potential for chemoprevention of cancer. *Adv Exp Med Biol.* **401**: 171-177.

- Wensing, M., Ludt, S., Campbell, S., van Lieshout, J., Volbracht, E., and Grol, R. (2009). European practice Assessment of cardiovascular risk management (EPA Cardio): protocol of an international observational study in primary care. *Implementation Sci.* **4**.
- Wetli, H. A., Brenneisen, R., Tschudi, I., Bigler, P., Sprang, T., Schurch, S., and Muhlbauer, R. C. (2004). Gamma-glutamyl-peptide isolated from onion by bioassay guided fractionation inhibits resorption activity of osteoclast **In:** 26th Annual Meeting of the American Society for Bone and Mineral Research, pp. S314-S314. Seattle, WA.
- Wu, C. P., Calcagno, A. M., Hladky, S. B., Ambudkar, S. V., and Barrand, M. A. (2005). Modulatory effects of plant phenols on human multidrug-resistance proteins 1, 4 and 5 (ABCC1, 4 and 5). *FEBS Journal.* **272**: 4725-4740.
- Wu, X., Beecher, G. R., Holden, J. M., Haytowitz, D. B., Gebhardt, S. E., and Prior, R. L. (2004). Lipophilic and hydrophilic antioxidant capacities of common foods in the United States. *J Agric Food Chem.* **52**: 12.
- Xiao, H., and Parkin, K. L. (2007). Isolation and identification of potential cancer chemopreventive agents from methanolic extracts of green onion (*Allium cepa*). *Phytochem.* **68**: 1059-1067.

- Yamazaki, Y., Iwasaki, K., Mikami, M., and Yagihashi, A. (2011). Distribution of Eleven Flavor Precursors, S-Alk(en)yl-L-Cysteine Derivatives, in Seven Allium Vegetables. *Food Sci Technol Res.* **17**: 55-62.
- Yang, C. Y., Chao, P. D. L., Hou, Y. C., Tsai, S. Y., Wen, K. C., and Hsiu, S. L. (2006). Marked decrease of cyclosporin bioavailability caused by coadministration ginkgo and onion in rats. *Food Chem Toxicol.* **44**: 1572-1578.
- Yoshida, H., Katsuzaki, M., Ohta, R., Ishikawa, K., Fukuda, H., Fujino, T., and Suzuki, A. (1999a). Antimicrobial activity of the thiosulfinates isolated from oil-macerated garlic extract. *Biosci Biotechnol Biochem.* **63**: 591-594.
- Yoshida, H., Katsuzaki, M., Ohta, R., Ishikawa, K., Fukuda, H., Fujino, T., and Suzuki, A. (1999b). An organosulfur compound isolated from oil-macerated garlic extract, and its antimicrobial effect. *Biosci Biotechnol Biochem.* **63**: 588-590.
- You, W. C., Blot, W. J., Chang, Y. S., Ershow, A., Yang, Z. T., An, Q., Henderson, B. E., Fraumeni, J. F., and Wang, T. G. (1989). Allium vegetables and reduced risk of stomach-cancer. *J NatCancer Inst.* **81**: 162-164.
- Yuan, L., Ji, T. F., Wang, A. G., Yang, J. B., and Su, Y. L. (2008). Two new furostanol saponins from the seeds of *Allium cepa* L. . *Chinese Chem Lett.* **19**: 461-464.

Zhou, Y., Zhuang, W., Hu, W., Liu, G.-J., Wu, T.-X., and Wu, X.-T. (2011). Consumption of Large Amounts of Allium Vegetables Reduces Risk for Gastric Cancer in a Meta-analysis. *Gastroenterol.* **141**: 80-89.

Zohri, A. N., Abdel-Gawad, K., and Saber, S. (1995). Antibacterial, antidermatophytic and antitoxigenic activities of onion (*Allium cepa* L.) oil. . *Microbiol Res.* **150**: 167-172.

Kingdom	Plantae
Clade	Angiosperms
Clase	Monocots
Order	Asparagales
Family	Amaryllidaceae
Subfamily	Allioieae
Genus	<i>Allium</i>
Species	<i>A. cepa</i>



Scientific Classification


Allium cepa L. var.Figure 1: Classification of *Allium cepa*

Figure 2: Flavonoids, Quercetin and quercetin glucosides structures present in *Allium cepa*




Figure 3: Formation of different organo-sulphur compounds in processed onions (Corzo-Martinez et al., 2007)

PROXIMATES	MEAN (g)	VITAMINS	MEAN (mg)	MINERALS	MEAN (mg)
Water	89.11	Vitamin C	7.4	Calcium	23
Protein	1.1	Thiamin	0.046	Iron	0.21
Total lipids	0.1	Riboflavin	0.027	Magnesium	10
Ash	0.35	Niacin	0.116	Phosphorus	29
Carbohydrates	0.34	Pantothenic acid	0.123	Potassium	146
Fibre	1.7	Vitamin B-6	0.12	Sodium	4
Sugar total	4.24	Folate	19 mcg	Zinc	0.17
Sucrose	0.99	Choline	6.1	Copper	0.039
Dextrose	1.97	Betaine	0.1	Manganese	0.129
Fructose	1.29	β-Carotene	1 µg	Selenium	0.5 µg
Energy	40 Kcal	Vitamin A	2 IU	-	-
-	-	Lutein+zeaxanthin	4 µg	-	-
-	-	Vitamin E	0.02	-	-
-	-	Vitamin K	0.4 µg	-	-

Table 1: Proximate analysis, vitamin contents and minerals present in 100g of raw onion
(USDA, 2008)

LIPIDS	MEAN (g)	AMINO ACIDS	MEAN (g)
Myristic acid	0.004	Tryptophan	0.014
Palmitic acid	0.034	Threonine	0.021
Stearic acid	0.004	Isoleucine	0.014
Total Saturated acids	0.042	Leucine	0.025
Oleic acid	0.013	Lysine	0.039
Linoleic acid	0.013	Methionine	0.002
Linolenic acid	0.004	Cystine	0.004
Total monounsaturated acids	0.013	Phenylalanine	0.025
Total polyunsaturated acids	0.017	Tyrosine	0.014
Phytosterols	15 mg	Valine	0.021

-	-	Arginine	0.104
-	-	Histidine	0.014
-	-	Alanine	0.021
-	-	Aspartic acid	0.091
-	-	Glutamic acid	0.258
-	-	Glycine	0.025
-	-	Proline	0.012
-	-	Serine	0.021

Table 2: Lipids and amino acids present in 100 g of raw onions (USDA, 2008)

NAME	FUNCTION
Vitamin A	Important for normal vision and eye health
Retinol (animal origin)	Involved in gene expression, embryonic development
Some carotenoids (plant origin, converted to retinol in the body)	and growth and health of new cells Assists in immune function
	May protect against cancers and atherosclerosis
Vitamin C	Necessary for healthy connective tissues – tendons, ligaments, cartilage, wound healing and healthy teeth
Ascorbic acid	Assists in iron absorption
	A protective antioxidant - may protect against some cancers
	Involved in hormone and neurotransmitter synthesis
Vitamin E	Non-specific chain-breaking antioxidant
alpha-tocopherols and tocotrienols	Reduces peroxidation of fatty acids
	May protect against atherosclerosis
Thiamin	Coenzyme in the metabolism of carbohydrates and branched-chain amino acids
Vitamin B1	Needed for nerve transmission
	Involved in formation of blood cells
Riboflavin	Important for skin and eye health
Vitamin B2	Coenzyme in numerous cellular redox reactions involved in energy metabolism, especially from fat and protein
Niacin	Coenzyme or cosubstrate in many biological reduction and oxidation reactions required for energy metabolism and fat synthesis and breakdown
Vitamin B3	Reduces LDL cholesterol and increases HDL cholesterol
Nicotinic acid, nicotinamide	

Vitamin B6	Coenzyme in nucleic acid metabolism,
Pyridoxine, pyridoxal, pyridoxamine	neurotransmitter synthesis, haemoglobin synthesis.
	Involved in neuronal excitation
	Reduces blood homocysteine levels
	Prevents megaloblastic anaemia
Folate	Coenzyme in DNA synthesis and amino acid
Generic term for large group of compounds including folic acid and pterylpolyglutamates	synthesis. Important for preventing neural tube defects
	Key role in preventing stroke and heart disease, including reducing blood homocysteine levels with vitamin B12
	May protect against colonic and rectal cancer
Calcium	Structural component of bones and teeth
	Role in cellular processes, muscle contraction, blood clotting, enzyme activation, nerve function
Copper	Aids in utilization of iron stores, lipid, collagen, pigment
	Role in neurotransmitters synthesis
Iron	Component of haemoglobin and myoglobin in blood, needed for oxygen transport
	Role in cellular function and respiration
Magnesium	Component of bones
	Role in enzyme, nerve, heart functions, and protein synthesis
Manganese	Aids in brain function, collagen formation, bone structure, growth, urea synthesis, glucose and lipid metabolism and central nervous system functioning
Potassium	Major ion of intracellular fluid
	Maintains water, electrolyte and pH balances
	Role in cell membrane transfer and nerve impulse transmission
Phosphorus	Structural component of bone, teeth, cell membranes, phospholipids, nucleic acids, nucleotide enzymes, cellular energy metabolism
	pH regulation
	Major ion of intracellular fluid and constituent of many essential compounds in body and processes

Zinc	Major role in immune system
	Required for numerous enzymes involved in growth and repair
	Involved in sexual maturation
	Role in taste, smell functions

Table 3: Major functions of Onion (*Allium cepa*) in human body (BUPA, 2006; FAO, 2009)

EFFECT	AUTHOR	MODEL
Antimutagenic effects	Ikken et al., 1999	Ames test
Antimutagenic effects	Kato et al., 1998	Salmonella strain
Protective effects	Gao et al., 1999	Human case-referent study
Stomach-carcinoma protection	Darant et al., 1996	Human case- control study
Breast- cancer protection	Challier et al., 1998	Human case-control study
Induction of Phase II enzymes	Guyinnet et al., 1999	Wistar rats
Chemopreventive activity	Siess et al., 1997	Rats
Chemopreventive activity	Takada et al., 1997	Ito rat-liver test
Lung cancer protection	Sankaranarayanan et al., 1994	Human case-control study
Brain tumor protection	Hu et al., 1999	Human case-control study
Gastric cancer protection	González et al., 2006	Human case-control study
colorectal cancer protection	Millen et al., 2007	Human case-control study
Prostate cancer protection	Hsing et al., 2002	Human case-control study
Tumors protection	Shrivastava and Ganesh, 2010	Rats

Table 4: Overview of cancer prevention by *Allium cepa*

EFFECT	AUTHOR	MODEL
Antifungal activity	Yin and Tsao, 1999	Aspergillus species
Antifungal activity	Mahmoudabadi et al., 2009	Scopulariopsis sp.
Antifungal activity	Abdel-Gawad and Saber, 1995	dermatophytic fungi

Antibacterial activity	Kim, 1997	Oral Pathogenic bacteria
Antibacterial activity	Zohri et al., 1995	Bacterial sps
Antibacterial activity	Elnima et al., 1983	several bacterial sps
Antibacterial activity	Abubakar, 2009	Mycobacterium tuberculosis
Antiparasitic activity	Reuter et al., 1996	various parasites
Antiparasitic activity	Nadia et al, 2011	<i>Cryptosporidium parvum</i>

Table 5: Overview of antibacterial, antifungal and antiparasitic activity of onions