



Recent Developments in the Food Quality Detected by Non-Invasive Nuclear Magnetic Resonance Technology

Kai Fan & Min Zhang

To cite this article: Kai Fan & Min Zhang (2018): Recent Developments in the Food Quality Detected by Non-Invasive Nuclear Magnetic Resonance Technology, Critical Reviews in Food Science and Nutrition, DOI: [10.1080/10408398.2018.1441124](https://doi.org/10.1080/10408398.2018.1441124)

To link to this article: <https://doi.org/10.1080/10408398.2018.1441124>



Accepted author version posted online: 16 Feb 2018.



Submit your article to this journal [↗](#)



View related articles [↗](#)



View Crossmark data [↗](#)

Publisher: Taylor & Francis

Journal: *Critical Reviews in Food Science and Nutrition*

DOI: <https://doi.org/10.1080/10408398.2018.1441124>

Recent developments in the food quality detected by non-invasive nuclear magnetic resonance technology

Kai Fan^{a, b}, Min Zhang^{a, c*}

^a State Key Laboratory of Food Science and Technology, Jiangnan University, 214122 Wuxi, Jiangsu, China

^b International Joint Laboratory on Food Safety, Jiangnan University, 214122 Wuxi, Jiangsu, China

^c Jiangsu Province Key Laboratory of Advanced Food Manufacturing Equipment and Technology, Jiangnan University, China

*Corresponding author: Dr. Min Zhang, Professor of School of Food Science and Technology, Jiangnan University, 214122 Wuxi, P. R. China

Tel.: +86-510-85877225; Fax: +86-510-85877225

E-mail: min@jiangnan.edu.cn

Abstract

Nuclear magnetic resonance (NMR) is a rapid, accurate and non-invasive technology and widely used to detect the quality of food, particularly to fruits and vegetables, meat and aquatic products. This review is a survey of recent developments in experimental results for the quality of food on various NMR technologies in processing and storage over the past decade. Following a discussion of the quality discrimination and classification of food, analysis of food compositions and detection of physical, chemical, structural and microbiological properties of food are outlined. Owing to high cost, low detection limit and sensitivity, the professional knowledge involved and the safety issues related to the maintenance of the magnetic field, so far the practical applications are limited to detect small range of food. In order to promote applications for a broader range of foods further research and development efforts are needed to overcome the limitations of NMR in the detection process. The needs and opportunities for future research and developments are outlined.

Keywords Nuclear magnetic resonance, Relaxation, Composition, Quality, Food

Introduction

Nuclear magnetic resonance (NMR) is a phenomenon of energy exchange with an alternating magnetic field by nuclei of fixed magnetic moments such as ^1H , ^{13}C , ^{23}Na , ^{31}P , ^{15}N (Laghi et al., 2014; Gudjonsdottir et al., 2015). NMR in food science was first used to determine moisture of food in the 1950s (Zhu, 2017). With the development and improvement of NMR technology in food, NMR is applied to different food fields including quantitative analysis (Chen et al., 2010; del Campo et al., 2010), conformational analysis (Vilén et al., 2010; Youssef et al., 2017), nutritional or functional aspects (Caligiani et al., 2010a; Lee et al., 2010), quality detection (Alonsosalces et al., 2010; Di et al., 2011) and process control (Gudjonsdottir et al., 2010). The advantages of NMR are that there is no need to separate different food ingredients, a small amount of sample pretreatment and preparation, as well as repeatability, non-destructive and quantitative (Kirtil and Oztop, 2015). However, the detection limits and sensitivity of NMR need to be further improved compared to gas chromatography, liquid chromatograph mass spectrometer, inductively coupled plasma mass spectrometry, liquid chromatography electrospray ionization and mass spectrometry technology (Marccone et al., 2013; Herrero et al., 2010; Le Bourse et al., 2010).

NMR technology has two types of high resolution NMR (HR-NMR) and low field NMR (LF-NMR) basing on NMR spectrum and magnetic resonance imaging (MRI) techniques (Kirtil et al., 2017). NMR technology was widely applied for food science in recent years because of the need for effective quality control analysis and the growing demand for technological and product innovation in the food industry (Spyros and Dais, 2012; Marcone et al., 2013). NMR spectroscopy was proposed for the determination of relaxation time. NMR relaxation time is used not only to measure chemical compositions, but also to evaluate the quality of food (Mariette, 2009). MRI is also a powerful tool to observe the interior structure information of food and monitor physical and chemical properties in food processing and storage (Marcone et al., 2013; Lagnika et al., 2013; Jiang et al., 2013; Trimigno et al., 2015; Xu et al., 2015). Therefore, NMR relaxation and NMI technology are very popular in food-related applications.

The advances in the application of NMR have been reviewed in food including wine (Ogrinc et al., 2003), dairy products (Karoui and Debaerdemaeker, 2007) or other applications such as food traceability (Mannina et al., 2012), food authenticity (Moore et al., 2012), study on the relationship between water distribution and the quality parameters in fruits and vegetables and meat (Pearce et al., 2011; Butz et al., 2010; Islam et al., 2015; Su et al., 2017). However, these discussions had a certain limited coverage in NMR application. Much progress in

recent years has been made to study NMR application in the food science. Therefore, the aims of this paper were to present an overview of the recent developments in the quality of food detected by NMR and prospects further research.

Quality discrimination and classification by NMR

The properties of food quality generally concerned by consumers including flavor, type, origin, year and so on (Consonni et al., 2011; Consonni et al., 2008; de Oliveira et al., 2014). These are important properties for evaluating food quality. NMR technology has been a valid method for molecular structure identification and research of food components, and plays an important role in quantitative analysis (Corsaro et al., 2016; Ribeiro et al., 2014a; Panarese et al., 2012). NMR combined with common multivariate methods including principal component analysis (PCA) and partial least square discriminant analysis (PLS-DA) was applied to identify and characterize food components by the collection and analysis of one-dimension (1D), two-dimensional (2D) and multidimensional NMR spectrum (Deborde et al., 2017; Kirtil et al., 2017; Panarese et al., 2012). Recently, NMR technology is the most common method in the field of food classification and discrimination applications (Ribeiro et al., 2014b; Santos et al., 2016; Yang et al., 2009; Boffo et al., 2009; Hachem et al., 2016; Zhang et al., 2012b; Shao and

Li, 2010). According to the specific standards of different foods, NMR can identify type, origin, date of manufacture, preservation time of fruit juice, meat and aquatic products, as shown in Table 1.

The nutrients and sensory attributes from different kinds of fruits are different. The NMR technique can be applied to identify the fruit juice and classify fruits cultivars, thus guaranteeing authenticity of fruit juice consumed by the large number of the consumers (Ogrinc et al., 2003). Cuny et al. (2008) presented that ^1H NMR spectroscopy combined with different chemometric analyses were used to discriminate orange juice from grapefruit juice. They obtained that independent-component analysis (ICA) as a useful chemometric tool gave better classification results than PCA basing on ^1H NMR spectra information. As reported by Clausen et al. (2011), juices from seven sour cherry clones/cultivars were classified by sensory evaluation and high-resolution ^1H NMR spectroscopy combined PCA and PLS. They obtained that malic acid was the most important metabolite and highly correlated with sweetness and sourness by PCA and PLS combined with 2D NMR spectra such as ^1H - ^1H COSY and ^1H - ^{13}C HSQC. On the other hand, Koda et al. (2012) presented that band-selective excited low-field ^1H NMR spectra combined with PCA was used to identify five mango cultivars (a total of 55). They found that arginine, histidine, phenylalanine, glutamine, shikimic acid and trigonelline were important components for five mango cultivars classified by F2-selective 2D NMR spectra including TOCSY, DQF-COSY, NOESY, ^1H - ^{13}C HSQC and ^1H - ^{13}C CT-HMBC. Similarly, ^1H NMR technology combined with PCA and PLS was used not

only to discriminate orange varieties but also to evaluate degradation of fresh orange juice with the changes of storage temperature and time for 24 h (de Oliveira et al., 2014). Results showed that formic, fumaric and acetic acids were produced in orange juice after storage 24 h, and succinic and lactic acids and ethanol in orange juice increased at the early stage of storage. Moreover, five orange juice varieties can be effectively identified basing on PCA analysis. Therefore, NMR coupled with chemometric techniques can be used to distinguished different varieties of fruit juice.

NMR technology has been explored to discriminate traceability and geographical origin of meat products such as beef. As reported by Jung et al. (2010), ^1H NMR assisted with PCA and orthogonal projection to latent structure-discriminant analysis (OPLS-DA) was applied to discriminate geographical origin. They found that beef extracts originated from Australia, Korea, New Zealand and United States were significantly difference. In addition, the succinate and amino acids were major metabolites, among which amino acids included isoleucine, leucine, methionine, tyrosine, and valine for discriminating the origin of beef according to ^1H NMR spectra and OPLS-DA analysis. Zanardi et al. (2013) presented that ^1H NMR combined with stepwise linear discriminant analysis (sLDA) and artificial neural network (ANN) was applied for classifying 90 irradiated or 30 non-irradiated beef samples. They obtained that both irradiated and non-irradiated beef can be classified by sLDA and ANN analyses. Thus, NMR could be suitable for identifying the irradiation dose in meat. On the other hand, time-domain ^1H Nuclear

Magnetic Resonance (TD-NMR) using CPMG and CWFP sequences was applied to discriminate 99 beef samples basing on race and sex of various bull (Santos et al., 2014). They obtained that TD-NMR combined with PLS-DA can provide a reliable identification for tracing beef samples. Therefore, NMR combined reliable analysis methods can be suitable to discriminate meat sample.

NMR technology has also been extended to discriminate authenticity and origin of aquatic products such as salmon, hake and prawn. Masoum et al. (2007) presented that the origin of salmon was classified by using ^1H NMR and support vector machines (SVMs). Results showed that ^1H NMR spectroscopy combined with SVMs can provide a validated method to discriminate wild and farmed salmon and reduce the possibility of fraud from salmon origins. As reported by Greiff et al. (2014), low-field NMR transverse relaxation time (T_2) was suitable to distinguish 0-3.0% salt hake basing on PCA. They obtained that ^1H LF-NMR can be used to study low-salt products. On the other hand, Li et al. (2018) investigated that LF-NMR spectroscopy and MRI was used to detect the hydrocolloids injected prawns. They found that T_2 fitting curves can be successfully applied for distinguishing adulterated prawns and formal ones with PCA. Therefore, NMR is a powerful technique to distinguish aquatic products.

NMR technology has been used to discriminate authenticity and origin of honey. Schievano et al. (2010) presented that the botanical origin of unifloral and polyfloral honey was identified by using ^1H NMR. They obtained that the time of NMR analysis required only 30 min and the yield of correct honey identification was 100%. On the other hand, honey adulteration increased with adding different commercial sugar syrups by producers because of limited production of honey. Ribeiro et al. (2014a) reported that the possibility of differentiating the botanical origin of honeys was evaluated by using LF-NMR. They established the relationship between T_2 parameters and physicochemical data of honey from 8 botanical sources including water content, water activity, pH and color. Results showed that good linear correlations were obtained between the T_2 and T_{21} parameters and physicochemical data. Similarly, Ribeiro et al. (2014b) investigated honey adulteration by high fructose corn syrup concentration from 0% to 100% using LF-NMR. Results showed that NMR T_2 parameters increased with increasing high fructose syrup concentration in adulterated honey. Linear correlations were observed between the T_2 , T_{21} and T_{22} parameters and physicochemical data. Therefore, LF-NMR can be used to discriminate and classify honeys by their botanical origin.

Composition analyzed by NMR

The energy and nutrition for people are mainly originated from food. All kinds of food have complex compositions including water, fat, protein, carbohydrates, cellulose and inorganic salts, which contain rich protons (Shao and Li, 2011; Kovrija and Rondeau-Mouro, 2017; Li et al., 2012b; Petrov et al., 2008; Nowacka et al., 2014; Aursand et al., 2010; Gudjonsdottir et al., 2015; McDonnell et al., 2013; Gudjónsdóttir et al., 2011a; Wang et al., 2016). NMR technology can rapidly and accurately analyze the change of components without changing the original state of foods. NMR technology in recent years has been widely applied to analyze water, fat and protein from food (Petrov et al., 2008; Sekiyama et al., 2012; Zhang and McCarthy, 2013; Tao and Ngadi, 2017; Niu et al., 2014; Khan et al., 2016; Castell-Palou et al., 2011; Mati et al., 2015; Yang et al., 2018).

Water analysis

Water distribution and water content are closely related to the quality of food in production, processing, storage and transportation (Halder et al., 2011; Mikac et al., 2015; Hong et al., 2009; Xu et al., 2016; Otero and Préstamo, 2009; Fundo et al., 2016; Aguiló-Aguayo et al., 2014; Santagapita et al., 2012; Geng et al., 2015; Sanchez-Alonso et al., 2014; Lv et al., 2016). Water distribution and water content in food can be measured by using NMR technology. According to NMR parameters including longitudinal relaxation time (T_1) and transverse relaxation

time (T_2) and MRI, water signal and can reflect the quantification of water content and water distribution in food such as fruits and vegetables, meat and aquatic products, as shown in Table 2.

NMR was widely used to study water distribution of fruits and vegetables. In terms of fruits, Cheng et al. (2014) investigated that LF-NMR and MRI were applied to analyze water status in osmotically dehydrated strawberry treated by ultrasound and pulsed vacuum. They found that the vacuolar space decreased and cytoplasm intercellular space increased basing on T_2 and MRI analysis. As reported by Tylewicz et al. (2016), TD-NMR was applied to assess water distribution of freeze-dried apple treated by pulsed electric fields at $0.3\text{--}1.2\text{ kV cm}^{-1}$ for 5, 10 and 15 pulses. Results showed that water molecules had high T_1 values in vacuoles and extracellular spaces. Similarly, Traffano-Schiffo et al. (2017) studied that water distribution in osmotic dehydrated kiwifruit pretreated by pulsed electric fields at $100\text{--}400\text{ V/cm}$ for 60 pulse was observed by using TD-NMR. They obtained that TD-NMR can quantify water molecules basing on water state of kiwifruit tissue, and obtain sorption isotherms of adsorbed water by applying BET model. In terms of vegetables, Xin et al. (2013) reported that LF-NMR and MRI was used to analyze water state in broccoli treated by ultrasound-assisted osmotic dehydration using trehalose. Results showed that less mobile water in cytoplasm (T_{21}) and high mobile water in vacuole (T_{22}) were decreased by ultrasound-assisted osmotic dehydration within 30 min. Meanwhile, MRI signals would present a bright “water strip”. Similarly, as reported by Xu et al. (2017), whose studied water distribution by LF-NMR and

MRI techniques in the broccoli during hot-air drying. On the other hand, Lv et al. (2017) investigated that LF-NMR was used to measure online water status of six kinds of vegetables dried by microwave vacuum drying. Results showed that there is no single empirical model fitting for signal amplitude (A_2) and moisture content (M_1) of mushroom, carrot, potato, lotus, edamame and corn. However, there was a significant linear relationship between A_2 and M_1 in different microwave vacuum drying stages for each vegetable. Therefore, NMR can be used to quantify and observe water status and distribution.

NMR as a fast noninvasive tool has been applied to understand water mobility and distribution in meat and aquatic products. For example, Carneiro et al. (2016) investigated that water mobility of salted sardines muscle was detected by using LF-NMR during storage at 30 °C. Results suggested that T_2 data displayed three components: pool I (11.2-17.5 ms), pool II (35.3-44.7 ms) and pool III (161.7-256.5 ms). On the other hand, Shao et al. (2016) studied that protein hydration states were measured for mobility water in meat batters by added water and fat. They found that there were three relaxation components (T_{2b} , T_{21} and T_{22}) in meat and fat groups by using LF-NMR. T_{2b} and T_{21} (0-10 ms) were protons in macromolecules and combination with macromolecules, respectively. T_{22} (40-70 ms) was water or fat combined with sol matrix of meat batters. After meat batters were heated, T_{23} (200-300 ms) was observed. In addition, T_{22} of meat added fat was higher than that of meat added water. However, T_{23} of meat added water was higher than that of meat added fat. Similarly, as reported by Ojha et al. (2017), LF-NMR

was used for evaluating the changes of water distribution in the drying of beef jerky by ultrasound pretreatment at 25, 33 and 45 kHz for 30 min. Results showed that there were three distinct peaks, which were respectively bound water (T_{2b} with the relaxation time of 0-10 ms), water from dense myofibrillar protein matrix (T_{21} with the relaxation time of 10-100 ms) and free water (T_{22} with the relaxation time of >100 ms). Therefore, LF-NMR can be used to detect water mobility in meat and aquatic products.

Fat analysis

Fat for the human body plays an important role in food composition (Elmsjo et al., 2015). On the one hand, fat is an essential component of the cell membrane and maintenance of body moisture, which is also a lot of hormone-generating material in the human bodies (Blair et al., 2016). On the other hand, the high intake of fat can cause obesity, coronary heart disease, and other health problems (Sun et al., 2017). Therefore, it is very important to detect the content and composition of fat. NMR spectroscopy can be used to analyze the content and composition of fat in foods (Youssef and Barbut, 2009; Fang et al., 2013; Colnago et al., 2011; Nascimento et al., 2017; Castell-Palou et al., 2012; Fry et al., 2017; Tao and Ngadi, 2017; Miklos et al., 2014; Ritota et al., 2010; Pereira et al., 2013a). Recently, some researchers have applied NMR to measure the content and fatty acid composition of the fat in animal products. For instance, Nestor et al. (2010) investigated

that high resolution magic angle spinning (HR-MAS) NMR was applied for quantifying small metabolites in Arctic char muscle. They found that HR-MAS NMR can obtain total n-3 fatty acid, EPA and DHA contents. In addition, they also observed the most abundant creatine/phosphocreatine, anserine and taurine. Similarly, Manzano Maria et al. (2010) studied that conjugated linoleic acids (CLA) in beef was measured by ^1H NMR spectra. Results showed that ^1H NMR was at least 10 times faster than GC for measurement of CLA content. Moreover, NMR can also detect CLA content in dairy and other meats products. Stefanova et al. (2011) reported that ^1H -NMR spectroscopy was used for examining fatty acid compositions of chicken meat irradiated by gamma-ray at 0.5-15.0 kGy. They found that there was a dose-effect relationship of irradiation by ^1H -NMR. As reported by Miklos et al. (2015), fat mobility in gels adding lard, diacylglycerols or sunflower oil was measured by LF-NMR. Results found that there was no significance in fat mobility between lard and diacylglycerols. However, sunflower oil has a large influence in fat mobility. On the other hand, Zang et al. (2017) investigated that fat content of small yellow croaker was measured during drying according to LF-NMR and multivariate analyses including PCA and PLS. They obtained that fat content was 0.43-1.72 g and accurately predicted by LF-NMR combined with PLS. Therefore, NMR can be effectively used to quantify fat content and analyze fatty acid compositions.

Protein analysis

Protein is an important component of food. NMR has been used for the qualitative and quantitative analysis of protein and amino acids in food (Consonni et al., 2011; Carneiro et al., 2016; Lee et al., 2010; Ritota et al., 2010; Gudjónsdóttir et al., 2011b). For example, Gudjónsdóttir et al. (2011a) presented that the relationship between NMR parameters and protein content was analyzed in rehydrated cod. Results showed that T_1 and T_{21} were the most significant for changes and denaturation of protein. Additionally, T_1 and T_{21} had significantly correlated with the non-protein nitrogen content (free amino acids, peptides, trimethylamine oxide, and trimethylamine). As reported by Sanchez-Alonso et al. (2014), LF-NMR was performed to analyze quality of hake muscle in ice for 3 and 14 days prior to different freezing methods (nitrogen, air blast and walk-in) and frozen storage temperature (-10 and -20 °C for 5 days, 8 and 18 weeks). Results found that T_{21} time constant decreased in hake muscle at -10 °C for 18 weeks due to protein denaturation and aggregation. Similarly, Carneiro et al. (2016) investigated that LF-NMR was used to measure the quality changes of salted sardines induced by water mobility during storage at 30 °C. They found that decarboxylase enzymes increased in salted sardines due to protein fraction denatures and degrades causing an increase in free amino acids. On the other hand, NMR combined with PCA was used to evaluate the changes of amino acids from orange juice treated by atmospheric cold plasma (ACP) and ozone (Alves Filho et al., 2016). Results showed that ^1H NMR spectrum for orange juice after ACP processing displayed high concentration of several amino acids including proline at 2.06 and 2.34 ppm, dimethylproline at 2.06 ppm, glutamic acid at 2.18 ppm, and of arginine at 1.94

ppm. ^1H NMR spectrum for orange juice after ozone processing presented high amount of tyrosine at 6.96 ppm and 7.24 ppm. Therefore, NMR is an effective tool to obtain qualitative and quantitative information on proteins.

Quality detected by NMR

The quality changes of food would take place in processing and storage (Gudjonsdottir et al., 2015; Carneiro et al., 2016). In order to evaluate physical, chemical, structural and microbiological properties changes, some researchers have used NMR to detect the quality of different food (del Campo et al., 2010; Gudjónsdóttir et al., 2011c; Zhang et al., 2012a; Caligiani et al., 2010b; Sekiyama et al., 2012; Fundo et al., 2016; Geng et al., 2015; Bizzani et al., 2017; Pereira et al., 2013b; Li et al., 2014).

Physical properties

Physical properties including color, water activity, texture, pH and water holding capacity are very important indicators for the quality of food in processing and storage (Pearce et al., 2011; Corsaro et al., 2016). Some researchers have used NMR to detect physical parameters (Li et al., 2012a; Pereira et al., 2013a; Geng et al., 2015; Li et al., 2014).

The color is an essential physical indicator from the consumer point of view. Li et al. (2012a) studied that the relationship between T_{21} and brightness (L^* value) was analyzed for thawed pork in Group A (38.68 ms for T_{21}), B (42.32 ms for T_{21}) and C (44.53 ms for T_{21}) by LF-NMR. Results presented that T_{21} was positive and significant correlation with L^* value ($P < 0.05$). As presented by Carneiro et al. (2013), the color parameters (L^* , a^* and b^*) of frozen shrimp with added sodium polyphosphate was investigated by using LF-NMR. They found that T_{21} was significant correlation with the L^* ($r = -0.82$), a^* ($r = -0.82$) and b^* ($r = -0.79$). Additionally, T_{21} were correlated with a lower intensity of a^* and b^* . Similarly, Li et al. (2017) presented that color changes in frozen shrimps were analyzed by LF-NMR. They observed that T_2 (T_{2b} and T_{21}) with exception for T_{22} was significantly correlated with L^* and b^* . However, T_2 was insignificant correlation with a^* . On the other hand, Fundo et al. (2016) investigated that a relationship between T_2 and total color difference (TCD) was established. Results showed that TCD decreased with an increase in T_2 because free water in cells can promote the biochemical reaction causing color changes. Therefore, NMR can reflect color change of food.

Water activity is important parameter for the microbiological stability of food. Aursand et al. (2008) presented the relationship between T_2 parameters and water activity during salted cod and salmon. They found that T_{21} has a good linear correlation ($F \leq 0.05$) with water activity basing on LF-NMR T_{21} parameter. Similarly, Gudjónsdóttir et al. (2011a) also analyzed LF-NMR parameters (T_1 , T_{21} and T_{22}) correlation with

water activity in the injected and brined pre-salted cod muscle. Results showed that T_1 , T_{21} and T_{22} presented significantly correlation with water activity. On the other hand, As presented by Fundo et al. (2016), the relationship between NMR parameters and water activity was established for fresh-cut 'Rocha' pear during 7 days of refrigerated storage. Results showed that an increase in maximum value for T_2 was with an increase in water activity. Therefore, NMR can be used to predict water activity in food processing and storage.

Another relevant physical parameter of food quality is texture. Sanchez-Alonso et al. (2012) investigated that shear stress was assessed by LF-NMR in hake during frozen storage at $-10\text{ }^{\circ}\text{C}$ for 6 months. They found that the shear stress values showed significant correlation with T_{21} by bi-exponential or CONTIN analysis. However, there was poor correlation with shear stress. Therefore, authors considered that T_{22} could be more heterogeneous than T_{21} . In addition, relaxation time of other water populations could also be closer to that of free water. Similarly, Li et al. (2014) presented that shear force of chicken jerky was measured during drying process by LF-NMR. They obtained that the signal area of the immobilized water per mass (A_2) was notable correlations with the shear force. The signal area of total water content per mass (A) and the shear force had significantly correlation. On the other hand, As presented by Geng et al. (2015), the correlations of NMR T_2 parameters and texture parameters (hardness, chewiness, cohesiveness and springiness) was showed in dried sea cucumber during rehydration process. Results demonstrated that both hardness and chewiness had significant correlations with T_2 parameters. However, there was a good

correlation between cohesiveness and T_{21} . The springiness had insignificantly correlated with T_2 parameters. Li et al. (2017) also presented that the correlations of NMR T_2 data and texture parameters were established in frozen shrimp muscle. They also observed that the hardness, gumminess, and chewiness were significant correlations with T_2 data with exception of T_{22} ($p < 0.01$ or $p < 0.05$). Therefore, NMR can be used to evaluate the texture change of food.

The change of pH can reflect the quality changes of food. Gudjónsdóttir et al. (2011a) presented that the correlation was studied between the NMR relaxation parameters and the muscle pH in cod during the salting and rehydration processes. They observed that the muscle pH had significantly correlated with T_1 . The transversal water distribution (A_{21}/A_{22}) was also significantly correlated with muscle pH. The similar results were reported by Gudjónsdóttir et al. (2011c), they found that the NMR T_{22} data was significant correlation with the muscle pH in brine injected cod loins during superchilled storage. The NMR T_1 , T_{21} and T_{22} data had also significant correlation with the muscle pH in low salt cod loins. As presented by Li et al. (2012a), They investigated the correlation of NMR T_{21} component and the muscle pH in thawed pork. Results showed that NMR T_{21} was significantly correlated with muscle pH ($p < 0.05$). On the other hand, Carneiro et al. (2013) studied that the correlation of NMR T_2 data and muscle pH was described in frozen shrimp treated by sodium tripolyphosphate. They observed that T_2

relaxation time ($r=0.87$ for T_{21} ; $r=0.70$ for T_{22}) had a good correlation with muscle pH. Therefore, NMR was a good way to indicate pH change of food.

Water holding capacity (WHC) can reflect the water and affect economic value of food. Gudjónsdóttir et al. (2011a) investigated that the relationship between NMR relaxation parameters and WHC was analyzed in cod during salting and rehydration process. They found that T_1 , T_{21} and T_{22} were significant correlations with WHC in cod muscle. The similar results were reported by Gudjónsdóttir et al. (2011c), They also observed that the relaxation times T_{21} and T_{22} had significant correlations with the WHC in brine injected and high salt cod loins during superchilled storage. T_1 , T_{21} and transversal water distribution (A_{21}/A_{22}) had also significant correlations with the WHC in modified atmosphere packed cod. There was significantly correlation between T_{21} and the WHC in brined cod. According to Sanchez-Alonso et al. (2012), they studied the relationship between LF-NMR parameters and the WHC in hake during frozen storage at $-10\text{ }^{\circ}\text{C}$ for 6 months. Results showed that T_{21} and T_{22} had significant correlations with the WHC by bi-exponential analysis ($p<0.10$). On the other hand, Carneiro et al. (2013) studied that the relationship of LF-NMR relaxation parameters and the WHC was established in frozen shrimp muscle treated by sodium tripolyphosphate. They obtained that T_{21} and T_{22} showed good correlations with the WHC ($r=0.90$ for T_{21} ; $r=0.81$ for T_{22}). As presented by Gudjonsdottir et al.

(2015), they investigated the correlation of LF-NMR relaxation times and the WHC in cod during the salting process. The relaxation time T_{21} was significant correlation with the WHC ($p < 0.05$). Therefore, NMR can be suitable to estimate the water holding capacity of food.

Chemical properties

Chemical compounds from food could be changed in processing and storage. Some researchers have used NMR to detect the chemical compounds of food (Alves Filho et al., 2016; Bizzani et al., 2017; Ritota et al., 2010). For instance, Otero and Préstamo (2009) presented that HR-MAS NMR was employed to analyze sugars and content and organic acids in control and pressurized strawberry at low pressures (100-200 MPa). They observed that there were no significant changes of organic acids in pressurized strawberries in comparison to the control ones basing on ^1H HR-MAS NMR spectrum region of 0-3 ppm. However, there was a significant hydrolysis of sucrose detected by ^1H HR-MAS NMR spectrum. The major sugars content of strawberries decreased in the region of 3-5 ppm causing an increase in glucose and fructose. Similarly, Alves Filho et al. (2016) also presented that HR-NMR combined with PCA was applied for evaluating composition changes of orange juice treated by atmospheric cold plasma and ozone. Results showed that there were no significant changes on sucrose, β -glucose, fructose and citric acid in the orange juice by quantitative analysis. As reported by Soares et al. (2017), ^1H HR-NMR spectroscopy coupled with PCA and

PLS-DA was used to estimate chemical composition changes in passion fruit juice at 85 and 140 °C for 4, 15, 30, and 60 s. Results showed that the degradation of sucrose in passion fruit juice occurred causing an increase in glucose and fructose and the formation of 5-(hydroxymethyl)-2-furfural (HMF) basing on ^1H HR-NMR spectroscopy combined with chemometric analysis at 140 °C for 60 s. Furthermore, according to the concentration change of HMF monitored by ^1H NMR profile thermally treated in fruit juice, the possibility of overheating damages was easily detected. Francini et al. (2017) investigated that quantitative ^1H HR-NMR (^1H HR-qNMR) spectroscopy with PCA was used to identify and quantify several polyphenols including catechin, epicatechin and chlorogenic acid in different apple varieties after drying. Results showed that the signals of several polyphenol compounds appeared in the NMR spectrum region of 6.0-8.5 ppm. ^1H HR-qNMR spectra combined with PCA provided the phenolic compounds information of different dried apple. Thus, the phenolic contents of different apple products can be detected to improve the nutraceutical value. Therefore, NMR was a valid way to evaluate chemical properties of food.

Structural properties

The tissue structure changes of food were induced in processing and storage. MRI was widely used to visualize the difference in tissue structure. Some researchers have applied MRI to observe water status and water distribution related to internal tissue structure changes in

food processing and storage (Otero and Préstamo, 2009; Gonzalez et al., 2010; Zhang and McCarthy, 2012; Xu et al., 2017). For instance, Zhang and McCarthy (2012) presented that the tissue structure differences of healthy pomegranate and pomegranate with black heart were reflected by using MRI during postharvest storage. They found that the T_2 weighted MRI image can visualize all arils within the rind in healthy pomegranate. However, the arils region of pomegranate with black heart had some void areas and a decrease in signal intensity for portions. Thus, MRI can recognize black heart region in pomegranate tissues to ensure postharvest quality of fruit. As reported by Xin et al. (2013), water-selective transverse images showed different water distribution and migration in untreated and osmotically treated broccoli by using MRI. They found that a bright image signal was obtained in untreated broccoli. There was a 'water strip' in broccoli osmotically treated by ultrasound before 30 min. However, 'water strip' disappeared and MRI images darkened in broccoli osmotically treated by ultrasound after 30 min. MRI results suggested that the ultrasound treatment for long time destroyed cell structure in broccoli causing water and solid loss. The similar results were presented by Cheng et al. (2014), MRI provide visual images on water distribution and status in the cellular and intercellular spaces of untreated and osmotically dehydrated strawberry. Therefore, MRI can reflect water distribution and status related to cell structure change of strawberry during processing. Geng et al. (2015) studied that MRI was used for analyzing interior structure of dried sea cucumber during the rehydration. Results showed that T_1 and T_2 weighted MRI images visualize interior structure changes of lightly dried sea

cucumber during presoaking process for 48 h and rehydration for 144 h. T_1 and T_2 weighted MRI images respectively showed bound water distribution and free water distribution in rehydrated sea cucumber. They found that water distribution was relatively uniform at presoaking time for 24h, and free water population was sufficient at rehydration time for 96 h. Therefore, MRI can provide internal tissue structure information to monitor the quality of food.

Microbiological properties

The product safety is the major factor which determines the product's suitability and market value. The microbiological growth is responsible for many deteriorative processes. Thus, the control of microbiological quality of foods is very critical. Some researchers have used NMR to detect microbiological quality of foods in processing and storage (Duarte et al., 2006; Gudjónsdóttir et al., 2011c). For example, Gudjónsdóttir et al. (2011c) investigated that the correlation of LF-NMR T_2 relaxation times and total viable counts (TVC) of H_2S -producing bacteria was established in brine injected and air-stored cod loins during superchilled storage. They found that the relaxation times (T_1 , T_{21} and T_{22}) were significant correlations with TVC in the brine injected cod loins. T_1 was also significant correlations with TVC of H_2S -producing bacteria in the air-stored cod loins. Therefore, fish quality deterioration can be monitored by using NMR technology. On the other hand, water

dynamics can affect microbiological quality of food. Fundo et al. (2016) studied T_2 parameter of fresh-cut pears during 7 days of refrigerated storage by solid state HR-NMR. Results showed that the T_2 maximum values changed to low values at 3 days. This demonstrated that the subcellular structure was disrupted through the wounding and a decrease in water activity led to water being use for microbiological growth. Thus, microbiological quality of food was evaluated basing on food water dynamics studied by NMR technology.

Potential application of online monitoring by NMR

The traditional analytical methods were difficult to be used for online monitoring in food processing because of long time and sample damage. Many non-invasive detection methods such as light transmission, X-ray transmission and ultrasonic have been applied to assess the quality of food. However, these methods were difficult to provide specific information about complex properties of food (Marcone et al., 2013). Recently, NMR was applied to achieve food online processing. For Example, Gudjonsdottir et al. (2011) presented that online control during shrimp processed by prebrining with polyphosphates and freezing was investigated by LF-NMR. They found that LF-NMR can rapidly detect physicochemical properties changes of shrimp including muscle pH, protein content, phosphate levels, moisture content and water-holding capacity. They also found that measurement settings, number of sample replicates, size of analyzing surface and suitable probe

selection need to be further optimized in shrimp processing by using NMR online control. On the other hand, Lv et al. (2017) presented that moisture contents of vegetables were measured online during microwave vacuum drying by LF-NMR method. They found that the relationship between signal amplitude A_{21} , A_{22} , A_{23} and A_2 and moisture content of fresh vegetables (mushroom, carrot, potato, lotus, edamame and corn) could not be fitted by using a single mathematical model. This method can measure moisture content of six kinds of vegetables online. However, the results from NMR could not be predicted all fresh vegetables. They also pointed out that online measuring efficiency should be improved needing further research to industrial application. Therefore, NMR is very potential to achieve online monitoring in food processing.

Conclusion and future research

Basing on the literature, it is clear that NMR technology is an effective and non-invasive tool to detect the authenticity, composition and quality of food such as fruits and vegetables, meat and aquatic products in processing and storage. NMR combined with multivariate analysis can rapidly identify and classify foods. NMR can be also used to analyze water, fat and protein composition in different food. Moreover, NMR can detect physical properties including color, water activity, texture, pH and WHC changes of foods. Meanwhile, NMR can provide quantitative information on chemical composition of foods. MRI can visualize internal tissue structure change of foods. From microbiological

quality analysis, the relationship between NMR parameters and microbiological growth was established to evaluated quality deterioration of food. However, there are still some drawbacks needing further research to successfully apply NMR in the quality detection of the industry. The future research and developments for NMR technology in food analysis area can be as follows: (1) NMR should hyphenated other spectroscopy methods such as NIR spectroscopy, size exclusion chromatographic, ultrasound, mass spectrometry, scattering techniques and rheometry to improve more accuracy, sensitivity and stability in quality detection process. (2) The developments of NMR hardware and user-friendly software are helpful to achieve simple operation of equipment and ensure the reliability of the obtaining data for food researchers. (3) A comprehensive database for NMR should be established for specific food products considering richness and diversity of food types. (4) Novel NMR sequences and automated data processing procedures should be created to achieve high throughput.

Acknowledgments

We acknowledge the financial support from National Key R&D Program of China (Contract No. 2017YFD0400901), Jiangsu Province (China) Agricultural Innovation Project (Contract No. CX(17)2017), Jiangsu Province(China) “Collaborative Innovation Center for Food Safety and

Quality Control” Industry Development Program, Jiangsu Province (China) Infrastructure Project (Contract No. BM2014051) all of which enabled us to carry out this study.

References

- Aguiló-Aguayo, I., Downey, G., Keenan, D. F., Lyng, J. G., Brunton, N., and Rai, D. K. (2014). Observations on the water distribution and extractable sugar content in carrot slices after pulsed electric field treatment. *Food Research International*. **64**:18-24.
- Alonsosalces, R. M., Héberger, K., Holland, M. V., Morenorojas, J. M., Mariani, C., Bellan, G., Reniero, F., and Guillou, C. (2010). Multivariate analysis of NMR fingerprint of the unsaponifiable fraction of virgin olive oils for authentication purposes. *Food Chemistry*. **118** (4):956-965.
- Alves Filho, E. G., Almeida, F. D., Cavalcante, R. S., de Brito, E. S., Cullen, P. J., Frias, J. M., Bourke, P., Fernandes, F. A., and Rodrigues, S. (2016). (1)H NMR spectroscopy and chemometrics evaluation of non-thermal processing of orange juice. *Food Chemistry*. **204**:102-107.

- Aursand, I. G., Erikson, U., and Veliyulin, E. (2010). Water properties and salt uptake in Atlantic salmon fillets as affected by ante-mortem stress, rigor mortis, and brine salting: A low-field ^1H NMR and $^1\text{H}/^{23}\text{Na}$ MRI study. *Food Chemistry*. **120** (2):482-489.
- Aursand, I. G., Gallartjornet, L., Erikson, U., Axelson, D. E., and Rustad, T. (2008). Water Distribution in Brine Salted Cod (*Gadus morhua*) and Salmon (*Salmo salar*): A Low-Field ^1H NMR Study. *Journal of Agricultural & Food Chemistry*. **56** (15):6252-6260.
- Bizzani, M., Flores, D. W. M., Colnago, L. A., and Ferreira, M. D. (2017). Non-invasive spectroscopic methods to estimate orange firmness, peel thickness, and total pectin content. *Microchemical Journal*. **133**:168-174.
- Blair, H. C., Sepulveda, J., and Papachristou, D. J. (2016). Nature and Nurture in Atherosclerosis: The Roles of Acylcarnitine and Cell Membrane-Fatty Acid Intermediates. *Vascular Pharmacology*. **78**:17.
- Boffo, E. F., Tavares, L. A., Ferreira, M. M. C., and Ferreira, A. G. (2009). Classification of Brazilian vinegars according to their ^1H NMR spectra by pattern recognition analysis. *LWT - Food Science and Technology*. **42** (9):1455-1460.

- Butz, P., Hofmann, C., and Tauscher, B. (2010). Recent developments in noninvasive techniques for fresh fruit and vegetable internal quality analysis. *Journal of Food Science*. **70** (9):R131-R141.
- Caligiani, A., Palla, G., Maietti, A., Cirlini, M., and Brandolini, V. (2010a). ¹H NMR fingerprinting of soybean extracts, with emphasis on identification and quantification of isoflavones. *Nutrients*. **2** (3):280-289.
- Caligiani, A., Palla, G., Maietti, A., Cirlini, M., and Brandolini, V. (2010b). ¹H NMR fingerprinting of soybean extracts, with emphasis on identification and quantification of isoflavones. *Nutrients*. **2** (3):280-289.
- Carneiro, C. d. S., Mársico, E. T., Ribeiro, R. d. O. R., Conte-Júnior, C. A., Mano, S. B., Augusto, C. J. C., and Oliveira de Jesus, E. F. (2016). Low-Field Nuclear Magnetic Resonance (LF NMR ¹H) to assess the mobility of water during storage of salted fish (*Sardinella brasiliensis*). *Journal of Food Engineering*. **169**:321-325.
- Carneiro, C. d. S., Mársico, E. T., Ribeiro, R. d. O. R., Conte Júnior, C. A., Álvares, T. S., and Jesus, E. F. O. d. (2013). Studies of the effect of sodium tripolyphosphate on frozen shrimp by physicochemical analytical methods and Low Field Nuclear Magnetic Resonance (LF ¹H NMR). *LWT - Food Science and Technology*. **50** (2):401-407.

Castell-Palou, A., Rosselló, C., Femenia, A., Bon, J., and Simal, S. (2011). Moisture profiles in cheese drying determined by TD-NMR:

Mathematical modeling of mass transfer. *Journal of Food Engineering*. **104** (4):525-531.

Castell-Palou, A., Rosselló, C., Femenia, A., and Simal, S. (2012). Simultaneous Quantification of Fat and Water Content in Cheese by TD-NMR.

Food and Bioprocess Technology. **6** (10):2685-2694.

Chen, F. L., Wei, Y. M., and Zhang, B. (2010). Characterization of water state and distribution in textured soybean protein using DSC and NMR.

Journal of Food Engineering. **100** (3):522-526.

Cheng, X.-f., Zhang, M., Adhikari, B., and Islam, M. N. (2014). Effect of power ultrasound and pulsed vacuum treatments on the dehydration

kinetics, distribution, and status of water in osmotically dehydrated strawberry: a combined NMR and DSC study. *Food Bioprocess*

Technol. **7** (10):2782-2792.

Clausen, M. R., Pedersen, B. H., Bertram, H. C., and Kidmose, U. (2011). Quality of sour cherry juice of different clones and cultivars (*Prunus*

cerasus L.) determined by a combined sensory and NMR spectroscopic approach. *Journal of Agricultural and Food Chemistry*. **59**

(22):12124-12130.

- Colnago, L. A., Azeredo, R. B., Marchi Netto, A., Andrade, F. D., and Venancio, T. (2011). Rapid analyses of oil and fat content in agri-food products using continuous wave free precession time domain NMR. *Magnetic Resonance in Chemistry*. **49 Suppl 1**:S113-120.
- Consonni, R., Cagliani, L. R., Benevelli, F., Spraul, M., Humpfer, E., and Stocchero, M. (2008). NMR and Chemometric methods: A powerful combination for characterization of Balsamic and Traditional Balsamic Vinegar of Modena. *Analytica Chimica Acta*. **611** (1):31-40.
- Consonni, R., Cagliani, L. R., Guantieri, V., and Simonato, B. (2011). Identification of metabolic content of selected Amarone wine. *Food Chemistry*. **129** (2):693-699.
- Corsaro, C., Cicero, N., Mallamace, D., Vasi, S., Naccari, C., Salvo, A., Giofrè, S. V., and Dugo, G. (2016). HR-MAS and NMR towards Foodomics. *Food Research International*. **89**:1085-1094.
- Cuny, M., Vigneau, E., Le Gall, G., Colquhoun, I., Lees, M., and Rutledge, D. N. (2008). Fruit juice authentication by ^1H NMR spectroscopy in combination with different chemometrics tools. *Analytical and Bioanalytical Chemistry*. **390** (1):419-427.

- de Oliveira, C. R., Carneiro, R. L., and Ferreira, A. G. (2014). Tracking the degradation of fresh orange juice and discrimination of orange varieties: an example of NMR in coordination with chemometrics analyses. *Food Chemistry*. **164**:446-453.
- Deborde, C., Moing, A., Roch, L., Jacob, D., Rolin, D., and Giraudeau, P. (2017). Plant metabolism as studied by NMR spectroscopy. *Progress in Nuclear Magnetic Resonance Spectroscopy*. **102-103**:61-97.
- del Campo, G., Berregi, I., Caracena, R., and Zuriarrain, J. (2010). Quantitative determination of caffeine, formic acid, trigonelline and 5-(hydroxymethyl)furfural in soluble coffees by ^1H NMR spectrometry. *Talanta*. **81** (1-2):367-371.
- Di, A. C., Callao, M. P., and Ruisánchez, I. (2011). ^1H NMR and UV-visible data fusion for determining Sudan dyes in culinary spices. *Talanta*. **84** (3):829-833.
- Duarte, I. F., Delgadillo, I., and Gil, A. M. (2006). Study of natural mango juice spoilage and microbial contamination with *Penicillium expansum* by high resolution ^1H NMR spectroscopy. *Food Chemistry*. **96** (2):313-324.

- Elmsjö, A., Rosqvist, F., Engskog, M. K., Haglöf, J., Kullberg, J., Iggman, D., Johansson, L., Ahlström, H., Arvidsson, T., Riserus, U., and Pettersson, C. (2015). NMR-based metabolic profiling in healthy individuals overfed different types of fat: links to changes in liver fat accumulation and lean tissue mass. *Nutr Diabetes*. **5**:e182.
- Fang, G., Goh, J. Y., Tay, M., Lau, H. F., and Li, S. F. (2013). Characterization of oils and fats by ¹H NMR and GC/MS fingerprinting: classification, prediction and detection of adulteration. *Food Chemistry*. **138** (2-3):1461-1469.
- Francini, A., Romeo, S., Cifelli, M., Gori, D., Domenici, V., and Sebastiani, L. (2017). (1)H NMR and PCA-based analysis revealed variety dependent changes in phenolic contents of apple fruit after drying. *Food Chemistry*. **221**:1206-1213.
- Fry, C. G., Hofstetter, H., and Bowman, M. D. (2017). A First Laboratory Utilizing NMR for Undergraduate Education: Characterization of Edible Fats and Oils by Quantitative ¹³C NMR. *Journal of Chemical Education*. **94** (9):1319-1323.
- Fundo, J. F., Galvis-Sanchez, A., Madureira, A. R., Carvalho, A., Feio, G., Silva, C. L. M., and Quintas, M. A. C. (2016). NMR water transverse relaxation time approach to understand storage stability of fresh-cut 'Rocha' pear. *LWT - Food Science and Technology*. **74**:280-285.

- Geng, S., Wang, H., Wang, X., Ma, X., Xiao, S., Wang, J., and Tan, M. (2015). A non-invasive NMR and MRI method to analyze the rehydration of dried sea cucumber. *Analytical Methods*. **7** (6):2413-2419.
- Gonzalez, M. E., Barrett, D. M., McCarthy, M. J., Vergeldt, F. J., Gerkema, E., Matser, A. M., and Van As, H. (2010). (1)H-NMR study of the impact of high pressure and thermal processing on cell membrane integrity of onions. *Journal of Food Science*. **75** (7):E417-425.
- Greiff, K., Fuentes, A., Aursand, I. G., Erikson, U., Masot, R., Alcaniz, M., and Barat, J. M. (2014). Innovative nondestructive measurements of water activity and the content of salts in low-salt hake minces. *Journal of Agricultural and Food Chemistry*. **62** (12):2496-2505.
- Gudjónsdóttir, M., Arason, S., and Rustad, T. (2011a). The effects of pre-salting methods on water distribution and protein denaturation of dry salted and rehydrated cod – A low-field NMR study. *Journal of Food Engineering*. **104** (1):23-29.
- Gudjónsdóttir, M., Karlsdóttir, M. G., Arason, S., and Rustad, T. (2011b). Injection of fish protein solutions of fresh saithe (*Pollachius virens*) fillets studied by low field Nuclear Magnetic Resonance and physicochemical measurements. *Journal of Food Science and Technology*. **50** (2):228-238.

- Gudjónsdóttir, M., Lauzon, H. L., Magnússon, H., Sveinsdóttir, K., Arason, S., Martinsdóttir, E., and Rustad, T. (2011c). Low field Nuclear Magnetic Resonance on the effect of salt and modified atmosphere packaging on cod (*Gadus morhua*) during superchilled storage. *Food Research International*. **44** (1):241-249.
- Gudjonsdottir, M., Gunnlaugsson, V. N., Finnbogadottir, G. A., Sveinsdottir, K., Magnusson, H., Arason, S., and Rustad, T. (2010). Process control of lightly salted wild and farmed Atlantic cod (*Gadus morhua*) by brine injection, brining, and freezing--a low field NMR study. *Journal of Food Science*. **75** (8):E527-E536.
- Gudjonsdottir, M., Jonsson, A., Bergsson, A. B., Arason, S., and Rustad, T. (2011). Shrimp processing assessed by low field nuclear magnetic resonance, near infrared spectroscopy, and physicochemical measurements--the effect of polyphosphate content and length of prebrining on shrimp muscle. *Journal of Food Science*. **76** (4):E357-367.
- Gudjonsdottir, M., Traore, A., Jonsson, A., Karlsdóttir, M. G., and Arason, S. (2015). The effects of pre-salting methods on salt and water distribution of heavily salted cod, as analyzed by $(1)\text{H}$ and $(23)\text{Na}$ MRI, $(23)\text{Na}$ NMR, low-field NMR and physicochemical analysis. *Food Chemistry*. **188**:664-672.

- Hachem, R., Assemat, G., Martins, N., Balayssac, S., Gilard, V., Martino, R., and Malet-Martino, M. (2016). Proton NMR for detection, identification and quantification of adulterants in 160 herbal food supplements marketed for weight loss. *Journal of Pharmaceutical and Biomedical Analysis*. **124**:34-47.
- Halder, A., Datta, A. K., and Spanswick, R. M. (2011). Water transport in cellular tissues during thermal processing. *AIChE Journal*. **57** (9):2574-2588.
- Herrero, M., Simó, C., Garcíacañas, V., Fanali, S., Cifuentes, A., and Cifuentes, A. (2010). Chiral capillary electrophoresis in food analysis. *Electrophoresis*. **31** (13):2106-2114.
- Hong, Y.-S., Cho, J.-H., Kim, N.-R., Lee, C., Cheong, C., Hong, K. S., and Lee, C.-H. (2009). Artifacts in the measurement of water distribution in soybeans using MR imaging. *Food Chemistry*. **112** (1):267-272.
- Islam, M. N., Zhang, M., Fang, Z., and Sun, J. (2015). Direct contact ultrasound assisted freezing of mushroom (*Agaricus bisporus*): Growth and size distribution of ice crystals. *International Journal of Refrigeration*. **57**:46-53.

Jiang, H., Zhang, M., Mujumdar, A. S., and Lim, R. X. (2013). Analysis of Temperature Distribution and SEM Images of Microwave Freeze Drying Banana Chips. *Food & Bioprocess Technology*. **6** (5):1144-1152.

Jung, Y., Lee, J., Kwon, J., Lee, K. S., Ryu, D. H., and Hwang, G. S. (2010). Discrimination of the geographical origin of beef by (1)H NMR-based metabolomics. *Journal of Agricultural and Food Chemistry*. **58** (19):10458-10466.

Karoui, R., and Debaerdemaeker, J. (2007). A review of the analytical methods coupled with chemometric tools for the determination of the quality and identity of dairy products. *Food Chemistry*. **102** (3):621-640.

Khan, M. I. H., Wellard, R. M., Nagy, S. A., Joardder, M. U. H., and Karim, M. A. (2016). Investigation of bound and free water in plant-based food material using NMR T 2 relaxometry. *Innovative Food Science & Emerging Technologies*. **38**:252-261.

Kirtil, E., Cikrikci, S., McCarthy, M. J., and Oztop, M. H. (2017). Recent advances in time domain NMR & MRI sensors and their food applications. *Current Opinion in Food Science*. **17**:9-15.

- Kirtil, E., and Oztop, M. H. (2015). ¹H Nuclear magnetic resonance relaxometry and magnetic resonance imaging and applications in food science and processing. *Food Engineering Reviews*. **8** (1):1-22.
- Koda, M., Furihata, K., Wei, F., Miyakawa, T., and Tanokura, M. (2012). Metabolic discrimination of mango juice from various cultivars by band-selective NMR spectroscopy. *Journal of Agricultural and Food Chemistry*. **60** (5):1158-1166.
- Kovrlja, R., and Rondeau-Mouro, C. (2017). Multi-scale NMR and MRI approaches to characterize starchy products. *Food Chemistry*. **236**:2-14.
- Laghi, L., Picone, G., and Capozzi, F. (2014). Nuclear magnetic resonance for foodomics beyond food analysis. *TRAC Trends in Analytical Chemistry*. **59**:93-102.
- Lagnika, C., Zhang, M., and Mothibe, K. J. (2013). Effects of ultrasound and high pressure argon on physico-chemical properties of white mushrooms (*Agaricus bisporus*) during postharvest storage. *Postharvest Biology & Technology*. **82** (4):87-94.
- Le Bourse, D., Jegou, S., Conreux, A., Villaine, S., and Jeandet, P. (2010). Review of preparative and analytical procedures for the study of proteins in grape juice and wine. *Analytica Chimica Acta*. **667** (1-2):33-42.

- Lee, J. E., Lee, B. J., Chung, J. O., Hwang, J. A., Lee, S. J., Lee, C. H., and Hong, Y. S. (2010). Geographical and climatic dependencies of green tea (*Camellia sinensis*) metabolites: a (1)H NMR-based metabolomics study. *Journal of Agricultural and Food Chemistry*. **58** (19):10582-10589.
- Li, C., Liu, D., Zhou, G., Xu, X., Qi, J., Shi, P., and Xia, T. (2012a). Meat quality and cooking attributes of thawed pork with different low field NMR T(21). *Meat Sci*. **92** (2):79-83.
- Li, J., Kang, J., Wang, L., Li, Z., Wang, R., Chen, Z. X., and Hou, G. G. (2012b). Effect of water migration between arabinoxylans and gluten on baking quality of whole wheat bread detected by magnetic resonance imaging (MRI). *Journal of Agricultural and Food Chemistry*. **60** (26):6507-6514.
- Li, M., Li, B., and Zhang, W. (2018). Rapid and non-invasive detection and imaging of the hydrocolloid-injected prawns with low-field NMR and MRI. *Food Chemistry*. **242**:16-21.
- Li, M., Wang, H., Zhao, G., Qiao, M., Li, M., Sun, L., Gao, X., and Zhang, J. (2014). Determining the drying degree and quality of chicken jerky by LF-NMR. *Journal of Food Engineering*. **139**:43-49.

- Li, X.-x., Liu, S., Su, W., Cai, L., and Li, J. (2017). Physical quality changes of precooked Chinese shrimp *Fenneropenaeus chinensis* and correlation to water distribution and mobility by low-field NMR during frozen storage. *Journal of Food Processing and Preservation*. **41** (6):e13220.
- Lv, W., Zhang, M., Bhandari, B., Li, L., and Wang, Y. (2017). Smart NMR method of measurement of moisture content of vegetables During microwave vacuum drying. *Food and Bioprocess Technology*. **10** (12):2251-2260.
- Lv, W., Zhang, M., Bhandari, B., Yang, Z., and Wang, Y. (2016). Analysis of drying properties and vacuum-impregnated qualities of edamame (Glycine max (L.) Merrill). *Drying Technology*. **35** (9):1075-1084.
- Mannina, L., Sobolev, A. P., and Viel, S. (2012). Liquid state ^1H high field NMR in food analysis. *Progress in Nuclear Magnetic Resonance Spectroscopy*. **66**:1-39.
- Manzano Maria, R., Colnago, L. A., Aparecida Forato, L., and Bouchard, D. (2010). Fast and simple nuclear magnetic resonance method to measure conjugated linoleic acid in beef. *Journal of Agricultural and Food Chemistry*. **58** (11):6562-6564.

Marcone, M. F., Wang, S., Albabish, W., Nie, S., Somnarain, D., and Hill, A. (2013). Diverse food-based applications of nuclear magnetic resonance (NMR) technology. *Food Research International*. **51** (2):729-747.

Mariette, F. (2009). Investigations of food colloids by NMR and MRI. *Current Opinion in Colloid & Interface Science*. **14** (3):203-211.

Masoum, S., Malabat, C., Jalali-Heravi, M., Guillou, C., Rezzi, S., and Rutledge, D. N. (2007). Application of support vector machines to ^1H NMR data of fish oils: methodology for the confirmation of wild and farmed salmon and their origins. *Analytical and Bioanalytical Chemistry*. **387** (4):1499-1510.

Mati, M., Staruch, L., and Šoral, M. (2015). Use of NMR spectroscopy in the analysis of carnosine and free amino acids in fermented sausages during ripening. *Chemical Papers*. **69** (10).

McDonnell, C. K., Allen, P., Duggan, E., Arimi, J. M., Casey, E., Duane, G., and Lyng, J. G. (2013). The effect of salt and fibre direction on water dynamics, distribution and mobility in pork muscle: a low field NMR study. *Meat Sci*. **95** (1):51-58.

Mikac, U., Sepe, A., and Sersa, I. (2015). MR microscopy for noninvasive detection of water distribution during soaking and cooking in the common bean. *Magnetic Resonance Imaging*. **33** (3):336-345.

Miklos, R., Cheong, L.-Z., Xu, X., Lametsch, R., and Larsen, F. H. (2015). Water and Fat Mobility in Myofibrillar Protein gels Explored by Low-Field NMR. *Food Biophysics*. **10** (3):316-323.

Miklos, R., Mora-Gallego, H., Larsen, F. H., Serra, X., Cheong, L. Z., Xu, X., Arnau, J., and Lametsch, R. (2014). Influence of lipid type on water and fat mobility in fermented sausages studied by low-field NMR. *Meat Sci*. **96** (1):617-622.

Moore, J. C., Spink, J., and Lipp, M. (2012). Development and application of a database of food ingredient fraud and economically motivated adulteration from 1980 to 2010. *Journal of Food Science*. **77** (4):R118-R126.

Nascimento, P. A., Barsanelli, P. L., Rebellato, A. P., Pallone, J. A., Colnago, L. A., and Pereira, F. M. (2017). Time-Domain Nuclear Magnetic Resonance (TD-NMR) and Chemometrics for Determination of Fat Content in Commercial Products of Milk Powder. *Journal of AOAC International*. **100** (2):330-334.

- Nestor, G., Bankefors, J., Schlechtriem, C., Brannas, E., Pickova, J., and Sandstrom, C. (2010). High-resolution (1)H magic angle spinning NMR spectroscopy of intact Arctic char (*Salvelinus Alpinus*) muscle. Quantitative analysis of n-3 fatty acids, EPA and DHA. *Journal of Agricultural and Food Chemistry*. **58** (20):10799-10803.
- Niu, L., Li, J., Chen, M.-S., and Xu, Z.-F. (2014). Determination of oil contents in Sacha inchi (*Plukenetia volubilis*) seeds at different developmental stages by two methods: Soxhlet extraction and time-domain nuclear magnetic resonance. *Industrial Crops and Products*. **56**:187-190.
- Nowacka, M., Tylewicz, U., Laghi, L., Dalla Rosa, M., and Witrowa-Rajchert, D. (2014). Effect of ultrasound treatment on the water state in kiwifruit during osmotic dehydration. *Food Chemistry*. **144**:18-25.
- Ogrinc, N., Kosir, I. J., Spangenberg, J. E., and Kidric, J. (2003). The application of NMR and MS methods for detection of adulteration of wine, fruit juices, and olive oil. A review. *Analytical and Bioanalytical Chemistry*. **376** (4):424-430.
- Ojha, K. S., Kerry, J. P., and Tiwari, B. K. (2017). Investigating the influence of ultrasound pre-treatment on drying kinetics and moisture migration measurement in *Lactobacillus sakei* cultured and uncultured beef jerky. *LWT - Food Science and Technology*. **81**:42-49.

- Otero, L., and Préstamo, G. (2009). Effects of pressure processing on strawberry studied by nuclear magnetic resonance. *Innovative Food Science & Emerging Technologies*. **10** (4):434-440.
- Panarese, V., Laghi, L., Pisi, A., Tylewicz, U., Rosa, M. D., and Rocculi, P. (2012). Effect of osmotic dehydration on Actinidia deliciosa kiwifruit: A combined NMR and ultrastructural study. *Food Chemistry*. **132** (4):1706-1712.
- Pearce, K. L., Rosenvold, K., Andersen, H. J., and Hopkins, D. L. (2011). Water distribution and mobility in meat during the conversion of muscle to meat and ageing and the impacts on fresh meat quality attributes--a review. *Meat Sci*. **89** (2):111-124.
- Pereira, F. M., Bertelli Pflanzner, S., Gomig, T., Lugnani Gomes, C., de Felicio, P. E., and Colnago, L. A. (2013a). Fast determination of beef quality parameters with time-domain nuclear magnetic resonance spectroscopy and chemometrics. *Talanta*. **108**:88-91.
- Pereira, F. M. V., Carvalho, A. d. S., Cabeça, I. F., and Colnago, L. A. (2013b). Classification of intact fresh plums according to sweetness using time-domain nuclear magnetic resonance and chemometrics. *Microchemical Journal*. **108**:14-17.

Petrov, O. V., Hay, J., Mastikhin, I. V., and Balcom, B. J. (2008). Fat and moisture content determination with unilateral NMR. *Food Research International*. **41** (7):758-764.

Ribeiro, R. d. O. R., Mársico, E. T., Carneiro, C. d. S., Monteiro, M. L. G., Conte Júnior, C. A., Mano, S., and de Jesus, E. F. O. (2014a).

Classification of Brazilian honeys by physical and chemical analytical methods and low field nuclear magnetic resonance (LF 1H NMR).

LWT - Food Science and Technology. **55** (1):90-95.

Ribeiro, R. d. O. R., Mársico, E. T., Carneiro, C. d. S., Monteiro, M. L. G., Júnior, C. C., and Jesus, E. F. O. d. (2014b). Detection of honey

adulteration of high fructose corn syrup by Low Field Nuclear Magnetic Resonance (LF 1H NMR). *Journal of Food Engineering*.

135:39-43.

Ritota, M., Marini, F., Sequi, P., and Valentini, M. (2010). Metabolomic characterization of Italian sweet pepper (*Capsicum annum* L.) by means of

HRMAS-NMR spectroscopy and multivariate analysis. *Journal of Agricultural and Food Chemistry*. **58** (17):9675-9684.

Sanchez-Alonso, I., Martinez, I., Sanchez-Valencia, J., and Careche, M. (2012). Estimation of freezing storage time and quality changes in hake

(*Merluccius merluccius*, L.) by low field NMR. *Food Chemistry*. **135** (3):1626-1634.

- Sanchez-Alonso, I., Moreno, P., and Careche, M. (2014). Low field nuclear magnetic resonance (LF-NMR) relaxometry in hake (*Merluccius merluccius*, L.) muscle after different freezing and storage conditions. *Food Chemistry*. **153**:250-257.
- Santagapita, P., Laghi, L., Panarese, V., Tylewicz, U., Rocculi, P., and Rosa, M. D. (2012). Modification of transverse NMR relaxation times and water diffusion coefficients of kiwifruit pericarp tissue subjected to osmotic dehydration. *Food and Bioprocess Technology*. **6** (6):1434-1443.
- Santos, P. M., Corrêa, C. C., Forato, L. A., Tullio, R. R., Cruz, G. M., and Colnago, L. A. (2014). A fast and non-destructive method to discriminate beef samples using TD-NMR. *Food Control*. **38**:204-208.
- Santos, P. M., Pereira-Filho, E. R., and Colnago, L. A. (2016). Detection and quantification of milk adulteration using time domain nuclear magnetic resonance (TD-NMR). *Microchemical Journal*. **124**:15-19.
- Schievano, E., Peggion, E., and Mammi, S. (2010). ¹H Nuclear Magnetic Resonance Spectra of Chloroform Extracts of Honey for Chemometric Determination of Its Botanical Origin. *Journal of Agricultural & Food Chemistry*. **58** (1):57-65.

- Sekiyama, Y., Horigane, A. K., Ono, H., Irie, K., Maeda, T., and Yoshida, M. (2012). T2 distribution of boiled dry spaghetti measured by MRI and its internal structure observed by fluorescence microscopy. *Food Research International*. **48** (2):374-379.
- Shao, J.-H., Deng, Y.-M., Song, L., Batur, A., Jia, N., and Liu, D.-Y. (2016). Investigation the effects of protein hydration states on the mobility water and fat in meat batters by LF-NMR technique. *LWT - Food Science and Technology*. **66**:1-6.
- Shao, X., and Li, Y. (2010). Classification and prediction by LF NMR. *Food and Bioprocess Technology*. **5** (5):1817-1823.
- Shao, X., and Li, Y. (2011). Application of low-field NMR to analyze water characteristics and predict unfrozen water in blanched sweet corn. *Food and Bioprocess Technology*. **6** (6):1593-1599.
- Soares, M. V., Alves Filho, E. G., Silva, L. M., Novotny, E. H., Canuto, K. M., Wurlitzer, N. J., Narain, N., and de Brito, E. S. (2017). Tracking thermal degradation on passion fruit juice through Nuclear Magnetic Resonance and chemometrics. *Food Chemistry*. **219**:1-6.
- Spyros, A., and Dais, P. (2012). NMR Spectroscopy in Food Analysis. *Progress in Nuclear Magnetic Resonance Spectroscopy*. **54** (3–4):195-207.

- Stefanova, R., Vasilev, N. V., and Vassilev, N. G. (2011). ^1H -NMR spectroscopy as an alternative tool for the detection of γ -ray irradiated meat. *Food Analytical Methods*. **4** (3):399-403.
- Su, Y., Zhang, M., Fang, Z., and Zhang, W. (2017). Analysis of dehydration kinetics, status of water and oil distribution of microwave-assisted vacuum frying potato chips combined with NMR and confocal laser scanning microscopy. *Food Research International*. **101**:188-197.
- Sun, Q., Li, M., Yang, X., Xu, X., Wang, J., and Zhang, J. (2017). Dietary salean reverts partially the metabolic gene expressions and NMR-based metabolomic profiles from high-fat-diet-induced obese rats. *Journal of Nutritional Biochemistry*. **47**:53-62.
- Tao, F., and Ngadi, M. (2017). Applications of spectroscopic techniques for fat and fatty acids analysis of dairy foods. *Current Opinion in Food Science*. **17**:100-112.
- Traffano-Schiffo, M. V., Laghi, L., Castro-Giráldez, M., Tylewicz, U., Rocculi, P., Ragni, L., Dalla Rosa, M., and Fito, P. J. (2017). Osmotic dehydration of organic kiwifruit pre-treated by pulsed electric fields and monitored by NMR. *Food Chemistry*. **236**:87-93.

- Trimigno, A., Marincola, F. C., Dellarosa, N., Picone, G., and Laghi, L. (2015). Definition of food quality by NMR-based foodomics. *Current Opinion in Food Science*. **4**:99-104.
- Tylewicz, U., Aganovic, K., Vannini, M., Toepfl, S., Bortolotti, V., Dalla Rosa, M., Oey, I., and Heinz, V. (2016). Effect of pulsed electric field treatment on water distribution of freeze-dried apple tissue evaluated with DSC and TD-NMR techniques. *Innovative Food Science & Emerging Technologies*. **37**:352-358.
- Vilén, E. M., Lundqvist, L. C. E., Jouanneau, D., Helbert, W., and Sandström, C. (2010). NMR Study on Hydroxy Protons of kappa- and kappa/mu-Hybrid Carrageenan Oligosaccharides: Experimental Evidence of Hydrogen Bonding and Chemical Exchange Interactions in kappa/mu Oligosaccharides. *Biomacromolecules*. **11**.
- Wang, L., Zhang, M., Bhandari, B., and Gao, Z. (2016). Effects of malondialdehyde-induced protein modification on water functionality and physicochemical state of fish myofibrillar protein gel. *Food Research International*. **86**:131-139.
- Xin, Y., Zhang, M., and Adhikari, B. (2013). Effect of trehalose and ultrasound-assisted osmotic dehydration on the state of water and glass transition temperature of broccoli (*Brassica oleracea* L. var. botrytis L.). *Journal of Food Engineering*. **119** (3):640-647.

- Xu, B.-g., Zhang, M., Bhandari, B., Sun, J., and Gao, Z. (2016). Infusion of CO₂ in a solid food: A novel method to enhance the low-frequency ultrasound effect on immersion freezing process. *Innovative Food Science & Emerging Technologies*. **35**:194-203.
- Xu, B. G., Zhang, M., Bhandari, B., Cheng, X. F., and Sun, J. (2015). Effect of Ultrasound Immersion Freezing on the Quality Attributes and Water Distributions of Wrapped Red Radish. *Food & Bioprocess Technology*. **8** (6):1366-1376.
- Xu, F., Jin, X., Zhang, L., and Chen, X. D. (2017). Investigation on water status and distribution in broccoli and the effects of drying on water status using NMR and MRI methods. *Food Research International*. **96**:191-197.
- Yang, F., Zhang, M., Bhandari, B., and Liu, Y. (2018). Investigation on lemon juice gel as food material for 3D printing and optimization of printing parameters. *LWT - Food Science and Technology*. **87**:67-76.
- Yang, S. O., Kim, M. S., Liu, K. H., Auh, J. H., Kim, Y. S., Kwon, D. Y., and Choi, H. K. (2009). Classification of fermented soybean paste during fermentation by ¹H nuclear magnetic resonance spectroscopy and principal component analysis. *Bioscience Biotechnology and Biochemistry*. **73** (3):502-507.

Youssef, M. K., and Barbut, S. (2009). Effects of protein level and fat/oil on emulsion stability, texture, microstructure and color of meat batters.

Meat Sci. **82** (2):228-233.

Youssof, L., Lallemand, L., Giraud, P., Soule, F., Bhaw-Luximon, A., Meilhac, O., D'Hellencourt, C. L., Jhurry, D., and Couprie, J. (2017).

Ultrasound-assisted extraction and structural characterization by NMR of alginates and carrageenans from seaweeds. *Carbohydrate Polymers.* **166**:55-63.

Zanardi, E., Caligiani, A., Padovani, E., Mariani, M., Ghidini, S., Palla, G., and Ianieri, A. (2013). Detection of irradiated beef by nuclear magnetic resonance lipid profiling combined with chemometric techniques. *Meat Sci.* **93** (2):171-177.

Zang, X., Lin, Z., Zhang, T., Wang, H., Cong, S., Song, Y., Li, Y., Cheng, S., and Tan, M. (2017). Non-destructive measurement of water and fat contents, water dynamics during drying and adulteration detection of intact small yellow croaker by low field NMR. *Journal of Food Measurement and Characterization.* **11** (4):1550-1558.

Zhang, J., Liu, Y., Wang, N., and Ruan, R. (2012a). NMR technique application in evaluating the quality of navel orange during storage. *Procedia Engineering.* **37**:234-239.

Zhang, L., and McCarthy, M. J. (2012). Black heart characterization and detection in pomegranate using NMR relaxometry and MR imaging.

Postharvest Biology and Technology. **67**:96-101.

Zhang, L., and McCarthy, M. J. (2013). NMR study of hydration of navy bean during cooking. *LWT - Food Science and Technology*. **53**

(2):402-408.

Zhang, Q., Saleh, A. S. M., and Shen, Q. (2012b). Discrimination of edible vegetable oil adulteration with used frying oil by low field nuclear magnetic resonance. *Food and Bioprocess Technology*. **6** (9):2562-2570.

Zhu, F. (2017). NMR spectroscopy of starch systems. *Food Hydrocolloids*. **63**:611-624.

List of Tables

Table 1 Recent studies on discrimination and classification of food by NMR

Table 2 Recent studies on application of NMR in water analysis

Table 1 Recent studies on discrimination and classification of food by NMR

Sample	NMR type	Analytical method	Research aim	Reference
Sour cherry juice	High-resolution ^1H NMR, 600.13 MHz	1D ^1H NMR spectra, 2D NMR (^1H - ^1H COSY and ^1H - ^{13}C HSQC) spectra, sensory evaluation, PCA and PLS	To classify juices from seven sour cherry clones/cultivars	Clausen et al. (2011)
Mango	1D and 2D ^1H NMR, 500 MHz	Low-field 1D ^1H NMR spectra, 2D NMR (TOCSY, DQF-COSY, NOESY, ^1H - ^{13}C HSQC and ^1H - ^{13}C CT-HMBC) spectra, PCA	To discriminate five mango cultivars	Koda et al. (2012)
Orange juice	^1H NMR, 400 MHz	^1H NMR spectrum, PCA and PLS	To identify five different orange varieties	de Oliveira et al. (2014)
Beef	^1H NMR, 600 MHz	^1H NMR spectroscopy, 2D NMR (^1H - ^1H TOCSY, ^1H - ^{13}C HMBC and ^1H - ^{13}C HSQC) spectra, PCA and OPLS-DA	To differentiate geographical origin of beef	Jung et al. (2010)

	^1H NMR, 600 MHz	^1H NMR spectra, sLDA and ANN	To classify irradiated or non-irradiated beef	Zanardi et al. (2013)
	Time-domain ^1H NMR, 85 MHz	TD-NMR, CPMG and CWFP pulse sequences, SIMCA, KNN and PLS-DA	To discriminate beef basing on bull race and sex	Santos et al. (2014)
Hake	Low-field ^1H NMR, 20 MHz	Transverse relaxation time (T_2), PCA	To distinguish 0-3% salt hake	Greiff et al. (2014)
Prawn	Low-field ^1H NMR, MRI, 21.16 MHz	Spin-spin relaxation time (T_2), PCA	To detect adulterated prawns	Li et al. (2018)

Table 2 Recent studies on application of NMR in water analysis

Sample	NMR type	Main results	Reference
Strawberry	^1H HR-MAS NMR, 500 MHz; MRI, 200 MHz	T_1 , T_2 data and MRI maps clearly indicated the changes of water molecules caused by 100-200 MPa pressure in strawberry.	Otero and Préstamo (2009)
Pear	Solid state NMR, 300 MHz	There are 3 kinds of water in the cells after processing of fresh-cut pear. T_2 values are respectively 10 ms, 187 ms and 3 s for cellular wall, cytoplasm and vacuole.	Fundo et al. (2016)
Kiwifruit	TD-NMR, 20 MHz	T_2 signals by TD-NMR can quantify water distribution in osmotic dehydrated kiwifruit treated by pulsed electric fields.	Traffano-Schiffo et al. (2017)
Broccoli	LF-NMR, 23.2 MHz	T_2 with three water fractions and MRI provide the changes of water status and distribution in broccoli during hot-air drying, which showed water mobility and water loss.	Xu et al. (2017)
Pork	LF-NMR, 23.4 MHz	T_2 can identify three water components (7–13 ms for T_{2b} ,	McDonnell et al. (2013)

		39-54 ms for T_{21} and 133-177 ms for T_{22}), all of which were significantly affected by NaCl concentration.	
Beef	LF-NMR, 23.2 MHz	Water distribution presented three peaks (T_{2b} , T_{21} and T_{22}) during drying of beef by T_2 analysis.	Ojha et al. (2017)
Salmon	LF ^1H NMR, 20 MHz	Water distribution of salmon has significant difference by T_2 analysis in ante-mortem handling, fillet location and brine salting.	Aursand et al. (2010)
Cod	LF-NMR, 20 MHz	Two water populations (23.1-94.2 ms for T_{21} and 126-210 ms for T_{22}) were observed according to T_2 data	Gudjónsdóttir et al. (2011a)
	LF-NMR, 20 MHz	A major band (T_{21} , 47-60 ms) of fresh cod muscle was presented basing on T_2 data. T_{21} of freezing cod muscle had no change at -20 °C and decreased at -18 °C during frozen storage.	Sanchez-Alonso et al. (2014)
	LF ^1H NMR, 20 MHz	Water distribution was the most homogeneous in cod with injection of salt and phosphates by T_2 analysis.	Gudjonsdottir et al. (2015)
Sea cucumber	LF-NMR, 23.2 MHz	Water distribution basing on T_2 data revealed the water uptake process during rehydration of dried sea cucumber.	Geng et al. (2015)