



Comparative effects of different dietary approaches on blood pressure in hypertensive and pre-hypertensive patients: a systematic review and network meta-analysis

Lukas Schwingshackl, Anna Chaimani, Carolina Schwedhelm, Estefania Toledo, Marina Pünsch, Georg Hoffmann & Heiner Boeing

To cite this article: Lukas Schwingshackl, Anna Chaimani, Carolina Schwedhelm, Estefania Toledo, Marina Pünsch, Georg Hoffmann & Heiner Boeing (2018): Comparative effects of different dietary approaches on blood pressure in hypertensive and pre-hypertensive patients: a systematic review and network meta-analysis, Critical Reviews in Food Science and Nutrition, DOI: [10.1080/10408398.2018.1463967](https://doi.org/10.1080/10408398.2018.1463967)

To link to this article: <https://doi.org/10.1080/10408398.2018.1463967>



View supplementary material [↗](#)



Accepted author version posted online: 02 May 2018.



Submit your article to this journal [↗](#)



View related articles [↗](#)



View Crossmark data [↗](#)

Comparative effects of different dietary approaches on blood pressure in hypertensive and pre-hypertensive patients: a systematic review and network meta-analysis

Lukas Schwingshackl^{1,2}, Anna Chaimani^{3,4,5}, Carolina Schwedhelm^{1,2}, Estefania Toledo^{6,7,8}, Marina Püsch¹ Georg Hoffmann⁹, Heiner Boeing^{1,2}

¹ Department of Epidemiology German Institute of Human Nutrition Potsdam-Rehbruecke (DIfE), Arthur-Scheunert-Allee 114-116, 14558 Nuthetal, Germany

² NutriAct-Competence Cluster Nutrition Research Berlin-Potsdam, Nuthetal, Germany

³ Paris Descartes University, Paris, France

⁴ INSERM, UMR1153 Epidemiology and Statistics, Sorbonne Paris Cité Research Center (CRESS), METHODS Team, Paris, France

⁵ Cochrane France, Paris, France

⁶ Department of Preventive Medicine and Public Health, Universidad de Navarra, Calle Irunlarrea 1, 31008 Pamplona, Spain

⁷ Centro del Investigación Biomédica en Red de la Fisiopatología de la Obesidad y Nutrición (CIBEROBN), Av. Monforte de Lemos 3-5, 28029 Madrid, Spain

⁸ IdiSNA, Navarra Institute for Health Research, Calle Irunlarrea 3, 31007 Pamplona, Spain

⁹ Department of Nutritional Sciences, University of Vienna, Althanstraße 14, 1090 Vienna, Austria

Corresponding author: Lukas Schwingshackl, PhD Arthur-Scheunert-Allee 114-116; 14558 Nuthetal, Germany T: +49 (0)33200 88-2723 lukas.schwingshackl@dife.de

Email: anna.chaimani@parisdescartes.fr carolina.schwedhelm@dife.de
etoledo@unav.es

marina.puensch@gmail.com

georg.hoffmann@univie.ac.at

boeing@dife.de

Authors' last names: Schwingshackl, Chaimani, Schwedhelm, Toledo, Püsch, Hoffmann,

Boeing

Running title: Network meta-analysis on diet and blood pressure

Financial support: No funding to declare

List of abbreviations: BMI: Body Mass Index; DASH: Dietary approaches to Stop Hypertension; DBP: Diastolic blood pressure; HP: high protein; LC: low-carbohydrate; LF: low-fat; LGI/LGL: low-glycaemic index/load; MD: mean difference; SBP: Systolic blood pressure; SUCRA: surface under the cumulative ranking curves;

Competing interests: Authors declare to have no competing of interest

Funding: No funding received

Abstract

Background: Pairwise meta-analyses have shown beneficial effects of individual dietary approaches on blood pressure but their comparative effects have not been established.

Objective: Therefore we performed a systematic review of different dietary intervention trials and estimated the aggregate blood pressure effects through network meta-analysis including hypertensive and pre-hypertensive patients.

Design: PubMed, Cochrane CENTRAL, and Google Scholar were searched until June 2017. The inclusion criteria were defined as follows: i) Randomized trial with a dietary approach; ii) hypertensive and pre-hypertensive adult patients; and iii) minimum intervention period of 12 weeks. In order to determine the pooled effect of each intervention relative to each of the other intervention for both diastolic and systolic blood pressure (SBP and DBP), random effects network meta-analysis was performed.

Results: A total of 67 trials comparing 13 dietary approaches (DASH, low-fat, moderate-carbohydrate, high-protein, low-carbohydrate, Mediterranean, Palaeolithic, vegetarian, low-GI/GL, low-sodium, Nordic, Tibetan, and control) enrolling 17,230 participants were included. In the network meta-analysis, the DASH, Mediterranean, low-carbohydrate, Palaeolithic, high-protein, low-glycaemic index, low-sodium, and low-fat dietary approaches were significantly more effective in reducing SBP (-8.73 to -2.32 mmHg) and DBP (-4.85 to -1.27 mmHg) compared to a control diet. According to the SUCRAs, the DASH diet was ranked the most effective dietary approach in reducing SBP (90%) and DBP (91%), followed by the Palaeolithic, and the low-carbohydrate diet (ranked 3rd for SBP) or the Mediterranean diet (ranked 3rd for DBP). For most comparisons, the credibility of evidence was rated very low to moderate, with the exception for the DASH vs. the low-fat dietary approach for which the quality of evidence was rated high.

Conclusion: The present network meta-analysis suggests that the DASH dietary approach might be the most effective dietary measure to reduce blood pressure among hypertensive and pre-hypertensive patients based on high quality evidence.

Keywords: dietary approaches, hypertension, network meta-analysis, blood pressure, evidence synthesis

ACCEPTED MANUSCRIPT

Background

According to the World Health Organization, approximately 1 billion adults worldwide suffer from hypertension, and the prevalence is increasing (NCD-Risk-Factor-Collaboration, 2017; World-Health-Organization, 2012). Hypertension is the most important modifiable risk factor for premature cardiovascular disease worldwide. Nearly 50% of all ischemic heart diseases and stroke events are attributable to hypertension (Lawes et al., 2008).

Dietary modifications play an important role in the prevention and management of hypertension. Evidence from a recent dose-response meta-analysis of prospective cohort studies indicates that optimal consumption of certain food groups (whole grains, fruit, nuts, legumes, dairy products, red and processed meat and sugar sweetened beverages) is associated with as strong reduction in hypertension risk (Schwingshackl et al., 2017b). According to the most recent guidelines by the American Heart Association, hypertensive and pre-hypertensive patients should follow dietary modifications (alcohol moderation, sodium reduction, and emphasis on increased consumption of fresh fruits, vegetables, and low-fat dairy products) (Eckel et al., 2014; Smith et al., 2011). However, considering various guidelines, dietary recommendations for the prevention and management of hypertension are still not comprehensive (Eckel et al., 2014; Mancia et al., 2013b; Smith et al., 2011).

Pairwise meta-analyses comparing Dietary approaches to stop hypertension (DASH) (Saneei et al., 2014), combined dietary approaches (Ndanuko et al., 2016), or lower sodium intake with usual care/control diets, indicate modest reductions in blood pressure (Aburto et al., 2013). Due to the high prevalence of hypertension, and the preventive effects of dietary factors, one of the most important questions that remain to be answered is which dietary approach offers the

greatest benefits. For answering this question, a promising approach is network meta-analysis (Leucht et al., 2016; Mavridis et al., 2015; Salanti, 2012). The methodology of network meta-analysis is an extension of the standard pairwise meta-analysis that enables a simultaneous comparison of multiple interventions forming a connected network while preserving the internal randomization of individual trials. This systematic review and network meta-analysis aims to establish a clinically meaningful hierarchy of different dietary approaches on blood pressure in hypertensive patients in a systematic review through the synthesis of available evidence from randomized trials.

Methods and design

The study protocol was registered in PROSPERO International Prospective Register of Systematic Reviews (<https://www.crd.york.ac.uk/prospero/>, identifier CRD42016049243) and published a priori (Schwingshackl et al., 2017a). The present systematic review adheres to the PRISMA guidelines and the corresponding extension for network meta-analyses (Chaimani et al., 2017; Hutton et al., 2015).

Search strategy

Literature searches were performed in PubMed, Cochrane Central Register of Controlled Trials (CENTRAL), and Google Scholar until June 2017, by applying no language restriction using a pre-defined search strategy (**Supplemental Appendix 1**).

Moreover, the reference lists from the identified articles were screened to search for additional relevant studies.

Eligibility criteria

As previously described the following inclusion criteria were established (Schwingshackl et al., 2017a):

- i) Randomized comparison/controlled design (parallel or cross-over) between different dietary approaches (Schwingshackl et al., 2018):
 - a. DASH diet: high intake of fruits & vegetables, low-fat dairy products, and whole grains, and low in sodium (Appel et al., 1997)
 - b. Mediterranean dietary pattern: high consumption of fruit, vegetables, olive oil, legumes, cereals, fish, and moderate intake of red wine during meals (Schwingshackl and Hoffmann, 2014a, c, 2015a, 2016; Schwingshackl et al., 2015b);
 - c. Low carbohydrate (LC) diet: <25% carbohydrates of total energy intake; high intake of animal and/ or plant protein; often high intake of fat (Schwingshackl and Hoffmann, 2013a);
 - d. Palaeolithic Diet: lean meat, fish, fruit, leafy and cruciferous vegetables, root vegetables, eggs and nuts, while excluding dairy products, cereal grains, beans, refined fats, sugar, candy, soft drinks, beer and extra addition of salt (Jonsson et al., 2009);
 - e. Moderate-carbohydrate diet: 25-45% carbohydrates of total energy intake; 10-20% protein intake (Schwingshackl and Hoffmann, 2013a);
 - f. High protein (HP) diet: >20% protein intake of total energy intake; high intake of animal and/ or plant protein; <35% fat (Schwingshackl and Hoffmann, 2013b);

- g. Nordic Diet: whole-grain products, abundant use of berries, fruit and vegetables, rapeseed oil, three fish meals per week, low-fat dairy products and avoidance of sugar-sweetened products (Uusitupa et al., 2013);
 - h. Tibetan Diet: high-protein and vitamin-rich food, preferably cooked and warm food (von Haehling et al., 2013);
 - i. Low fat (LF) diet: <30% fat of total energy intake; high intake of cereals & grains; 10-15% protein intake (Schwingshackl and Hoffmann, 2013a, 2014b);
 - j. Low glycaemic index/load (LGI/LGL) diet (Schwingshackl et al., 2015a; Schwingshackl and Hoffmann, 2013c);
 - k. Vegetarian/Vegan diet: no meat and fish/ no animal products (Haider et al., 2017)
 - l. Low-sodium diet (Whelton et al., 1998);
 - m. Control diet: usual diet.
- ii) Minimum intervention period of 12 weeks;
 - iii) Participants with a mean age ≥ 18 years;
 - iv) Hypertension or pre-hypertension (high normal blood pressure: 130-139 mmHg for systolic and/or 85-89 for diastolic blood pressure) defined according to the European Society of Hypertension and the European Society of Cardiology (Mancia et al., 2013a);
 - v) The outcomes include systolic (SBP) and diastolic blood pressure (DBP) (mmHg).

The following studies were excluded:

- i) Randomized trials including pregnant women, children, and adolescents
- ii) Intervention studies solely based on dietary supplements or single foods;
- iii) Intervention studies using dietary supplements as placebo;
- iv) Studies with an exercise/medication (Schwingshackl et al., 2013; Schwingshackl et al., 2014) co-intervention that was not applied in all the intervention/control groups;
- v) Interventions based on very low energy diets (i.e. <600 kcal/day);

Data extraction

After determination of the study selection, two reviewers extracted the following characteristics: name of first author, year of publication, country, sample size, study design, mean baseline Body Mass Index (BMI), mean baseline SBP and DBP, mean baseline age, % female, study duration, description of the different dietary arms, type of diet (energy restricted, ad libitum, iso-caloric), drop outs, % hypertension, % presence of comorbidities (obesity, hyperlipidaemia, type 2 diabetes, metabolic syndrome, coronary heart disease), % antihypertensive medication, and % lipid lowering medication. Outcome data include: post-intervention values with corresponding standard deviations for SBP and DBP.

Risk of bias assessment

One author assessed the risk of bias of the included trials, by applying the Cochrane risk of bias tool (Higgins et al., 2011). The following items were assessed: random sequence generation and allocation concealment, blinding of participants and personnel, incomplete outcome data, and selective reporting.

Studies were classified as being at low risk of bias (if at least three out of a maximum of five items were rated as low risk; and maximum one item rated with a high risk of bias), high risk of bias (if at least two out of a maximum of five items were rated as high risk), and moderate/unclear risk (all other studies).

Dealing with missing data

We contacted authors to receive missing outcome data (1 author sent additional data). If the post-intervention values with the corresponding standard deviations were not available, the change scores with the corresponding standard deviations were used, according to the guidelines of the Cochrane Handbook (Higgins and Green, 2011).

Evaluation of synthesis assumptions

Data synthesis

Description of the available data

Direct comparisons between different dietary approaches were illustrated by using a network diagram (Chaimani et al., 2013). Moreover, the contribution matrix was used to identify the direct comparisons with greater influence in the network relative effects (Chaimani et al., 2013; Krahn et al., 2013).

Assessment of transitivity

To evaluate the assumption of transitivity we compared the distribution of the potential effect modifiers (age, study length, Body Mass Index (BMI)) across the available direct comparisons.

Statistical analysis

In order to quantify the pooled relative effect of each dietary approach against every other dietary approach in terms of the post-intervention values (or changes from baseline values) of the different dietary measures we performed a random effects network meta-analysis for SBP and DBP. We performed network meta-analysis to synthesize all the available evidence. Compared to the standard pairwise meta-analysis model, network meta-analysis methods are extensions that enable a simultaneous comparison of multiple interventions forming a connected network while preserving the internal randomization of individual trials.

Random effects network meta-analysis for SBP and DBP were performed to estimate all possible pairwise relative effects and obtain a relative ranking of the different dietary approaches. The summary mean differences with their 95% CI were presented in a League table. We estimated the relative ranking of the different dietary approaches for SBP and DBP (Salanti et al., 2011) with using the distribution of the ranking probabilities and the surface under the cumulative ranking curves (SUCRA). For SBP and DBP, we assumed a common network-specific heterogeneity parameter and estimated the predictive intervals to assess how much this heterogeneity affects the relative effects with respect to the additional uncertainty anticipated in future studies (Riley et al., 2011). Statistical analyses were performed by using Stata 14.0 (StataCorp., 2015) (*network* package (White, 2015)) and produced presentation tools with the *network graphs* package (Chaimani and Salanti, 2015).

Assessment of inconsistency

To evaluate the presence of statistical inconsistency, the loop-specific approach (Bucher et al.,

1997) (detection of loops of evidence that might present important inconsistency), and the side-splitting approach (Dias et al., 2010) (detect comparisons for which direct estimates disagree with indirect evidence from the entire network) were applied. Global methods jointly investigate the presence of inconsistency from all possible sources in the entire network simultaneously. For this purpose, we used the design-by-treatment interaction model (Higgins et al., 2012; Jackson et al., 2014).

Subgroup and sensitivity analyses

Subgroup analyses according to hypertension status, study length (≥ 12 months vs. < 12 months), and sample size (≥ 100 vs. < 100) were performed for SBP and DBP. We also conducted a sensitivity analysis by analysing only studies considered being at low risk of bias. Additionally, we ran meta-regressions by participant and study characteristics for SBP and DBP: age, study length, and mean differences in weight change (by comparing study arms within each trial).

Small study effects and publication bias

The presence of small-study effects for each SBP and DBP was evaluated by drawing comparison-adjusted funnel plot that accounts for the fact that different studies compare different sets of interventions (Chaimani et al., 2013). Funnel plots included all comparisons of dietary approaches compared to a low-fat or control diet.

Credibility of the evidence

To assess the credibility of evidence from the network meta-analysis, we used the GRADE system extended for network meta-analysis following the approach suggested by Salanti et al.

(Supplemental Appendix 2) (Salanti et al., 2014).

ACCEPTED MANUSCRIPT

Results

Of a total of 2,949 records identified in the literature search, 115 full text articles were assessed in detail (**Supplemental Figure 1**). Of these, 48 were excluded, for the exclusion criteria summarized in **Supplemental Table 1, Supplemental References**.

Overall 67 trials met the eligibility criteria and were included in the meta-analysis (**Table 1**). 17,230 pre-hypertensive or hypertensive patients were enrolled, and the trials were published between 1981 and 2016. Eighteen trials were conducted in North America, 29 trials in Europe, 4 trials in Asia, and 15 trials in Australia and New Zealand, and one study in South America. Study length ranged between 3 and 48 months; the patients' mean age was between 23.6 and 71 years, and their BMI between 23.6 and 45.4 kg/m². The general and specific study characteristics are summarized in **Supplemental Table 2-3**.

Twenty-three trials were judged to be low risk of bias, five trials to be high risk of bias, and 39 trials were classified as moderate/unclear risk of bias studies. With regard to the specific items of the risk of bias assessment tool by the Cochrane Collaboration, 46% of the included studies indicate a low risk of bias for random-sequence generation, 21% for allocation concealment, 0% for blinding, 60% for incomplete data outcome, and 96% for selective reporting (**Supplemental Figure 2**).

The definition of the different intervention diets was heterogeneous for the intensity and type of dietary approach (i.e. group meeting, dietary counselling) and for the prescribed diets (ad-libitum, iso-caloric, hypocaloric). Across the included low-fat dietary approaches the overall amount of fat intake varied between ~10-15% of total energy intake. **We thus had to harmonize the single trials and formed classes of dietary approaches** (Schwingshackl et al., 2017a).

Figure 1 shows the network diagram of direct comparison for SBP with the number of studies reflected by the size of the edges, and the number of patients reflected by the size of the nodes. The comparisons with the largest amount of trials include: HP vs. LF (Brinkworth et al., 2004a, Brinkworth et al., 2004b; Campbell and Meckling, 2012; Dalle Grave et al., 2013; Dansinger et al., 2005; Delbridge et al., 2009; Kim et al., 2014; Krebs et al., 2012; Luger et al., 2013; Tang et al., 2013; Watson et al., 2016; Wycherley et al., 2010) (n=12), low-sodium vs. control (Alli et al., 1992; Chalmers et al., 1986; Costa et al., 1981; Dodson et al., 1989; Erwieman et al., 1984; Jula and Karanko, 1994; Makela et al., 2008; Silman et al., 1983; Whelton, 1997; Whelton et al., 1998) (n=10), LF vs. control (Anderssen et al., 1995; Andrews et al., 2011; Ard et al., 2016; Coppel et al., 2010; Gordon et al., 1997; Heilbronn et al., 1999; Rock et al., 2014; Uusitupa et al., 1993; Watkins et al., 2003) (n=9), and LC vs. LF dietary approaches (Daly et al., 2006; Guldbrand et al., 2012; Iqbal et al., 2010; Liu et al., 2013; Stern et al., 2004; Tay et al., 2008; Tay et al., 2015; Veum et al., 2017) (n=8). **Table 2** shows the percentage of statistical contribution coming from direct and indirect comparisons for each dietary approach compared to each other. It was shown that most of the contribution to the study effects came from indirect comparisons. Direct comparisons dominated the comparisons of DASH, Vegetarian, Mediterranean, HP, moderate-carbohydrate, LC, LGI/LGG, Palaeolithic, Tibetan, and a control diet with a LF diet for both outcomes. In general, there are some important differences in the examined effect modifiers across comparisons for BMI and study duration. For some comparisons, though, we did not have enough studies and we could not test transitivity appropriately (**Supplemental Figure 3-5**).

Table 3 summarizes the pooled estimates of SBP and DBP for the comparison of every dietary

approach using both direct and indirect evidence. The DASH, Mediterranean, LC, Palaeolithic, HP, LGI/LGL, low-sodium, and LF dietary approaches were more effective in reducing SBP and DBP compared to a control diet. The DASH diet was more effective in reducing SBP compared to a LF (mean difference (MD): -5.05 mmHg, 95% -7.08, -3.03), Mediterranean (MD: -3.31 mmHg, 95% -6.20, -0.42), HP (MD: -4.78 mmHg, 95% -7.72, -1.85), moderate-carbohydrate (MD: -5.57 mmHg, 95% -8.75, -2.39), LGI/LGL (MD: -4.92 mmHg, 95% -8.55, -1.30), Nordic (MD: -4.74 mmHg, 95% -9.11, -0.37), and low-sodium dietary approaches (MD: -4.42 mmHg, 95% -7.12, -1.72), and more effective in reducing DBP compared to a LF diet (MD: -3.10 mmHg, 95% -4.52, -1.68), moderate-carbohydrate (MD: -3.05 mmHg, 95% -5.37, -0.74), Nordic (MD: -3.22 mmHg, 95% -6.14, -0.31), Tibetan (MD: -3.20 mmHg, 95% -6.32, -0.08) and low-sodium diet (MD: -1.91 mmHg, 95% -3.70, -0.12). The Mediterranean dietary approach was more effective in reducing DBP compared to a LF diet (MD: -1.54 mmHg, 95% -2.95, -0.13). According to the SUCRAs, the DASH diet was ranked the most effective dietary approach in reducing SBP (90%) and DBP (91%), followed by the Palaeolithic, and the low-carbohydrate diet (ranked 3rd for SBP) or the Mediterranean diet (ranked 3rd for DBP) (**Supplemental Table 4-5**).

The rankograms showed **uncertainty in ranking (similar distribution of rank probabilities across many possible ranks indicates uncertain ranking for that dietary approach)** for the HP, LC, LGI/LGL, moderate-carbohydrate, Mediterranean, low-sodium, Nordic, and Tibetan dietary approach compared to the other dietary approaches (**Supplemental Figure 6-7**).

The side-splitting approach suggested no significant inconsistency for SBP and DBP (**Supplemental Table 6-7**). The loop-specific approach identified two-loops (control – LF –

moderate-carbohydrate; and LF – LC – Mediterranean) for SBP presenting statistical inconsistency, and one-loop for DBP (LF – LC – LGI/GL) (**Supplemental Figure 8-9**). The design-by-treatment model did not suggest the presence of statistical inconsistency for SBP ($p=0.65$), and DBP ($p=0.83$). The important inconsistency in the loop specific approach might be explained by several differences across LF dietary approaches (hypocaloric if compared to a control diet; often iso-caloric if compared to other dietary approaches), differences in total fat to carbohydrate intake ratio, and fatty acids composition among moderate carbohydrate approaches, and LC dietary approaches (larger weight loss compared to other interventions).

Subgroup and sensitivity analysis

In sensitivity analysis including only studies with a low risk of bias, the results of the primary analysis for the best ranked dietary approach, i.e. the DASH diet could be confirmed (**Table 4**). No significant effects for all other dietary approaches were observed in the low risk of bias sensitivity analysis.

The a priori planned subgroup analysis for age was not possible to conduct since only 8 trials included participants >60 years of age. In the subgroup analyses comparing longer-term (≥ 12 months) vs. shorter-term trials (<12 months), most of the results of the primary analysis were not confirmed among longer-term studies. Moreover, it seems that trials with a smaller sample size (<100) yielded more significant results compared to studies with a larger sample size (≥ 100) (**Supplemental Table 8-15**).

Due to the low number of studies presenting results for hypertensive patients ($n=12$), it was not possible to conduct sensitivity analysis.

In univariate meta-regression analysis we could show that mean reduction in SBP and DBP was larger in short-term trial, and among trials including younger participants. Moreover, we could show that larger differences in body weight change between study arms within trials were associated with a stronger impact on SBP and DBP, compared to smaller differences (**Supplemental Figure 10-15**), thereby showing that weight loss is a major contributing factor for blood pressure control.

Small study effects

The comparison-adjusted funnel plots for SBP and DBP appeared asymmetric suggesting that small studies tend to favour the dietary approaches when compared to control dietary approaches, and slightly asymmetric for low fat diets (**Supplemental Figure 16-19**).

Credibility of the evidence

For most comparisons in the network meta-analysis, the credibility of evidence was rated very low to moderate with the exception for the DASH vs. LF dietary approach the quality of evidence was rated high (**Supplemental Figure 20, Supplemental Appendix 2**).

Discussion

In this meta-analysis of 67 trials including 17230 hypertensive and pre-hypertensive patients we compared the effects of 13 different dietary approaches (DASH, Control, Low-Fat, Moderate-carbohydrate, High-protein, Low-carbohydrate, Mediterranean, Palaeolithic, Vegetarian, Low-GI/GL, Low-sodium, Nordic, and Tibetan) using network meta-analysis. According to the SUCRAs, the DASH diet was ranked the most effective dietary approach in reducing SBP (90%) and DBP (91%), followed by the Palaeolithic, and the low-carbohydrate diet (ranked 3rd for SBP) or the Mediterranean diet (ranked 3rd for DBP). The DASH, Mediterranean, low-carbohydrate, Palaeolithic, high-protein, low-glycaemic index, low-sodium, and low-fat dietary approach were significantly more effective in reducing SBP (-8.73 to -2.32 mmHg) and DBP (-4.85 to -1.27 mmHg) compared to a control diet. For most comparisons, the credibility of evidence was rated very low to moderate, with the exception for the DASH vs. the low-fat dietary approach the quality of evidence was rated high.

Compared to a LF or a control diet, the DASH dietary approach reduced SBP by approximately 5-7 mmHg, and DBP by 3-4 mm Hg. With respect to other therapeutic options, a recent meta-analysis investigating the effect of different exercise modalities resulted in some less pronounced effect of aerobic exercise on blood pressure (Cornelissen and Smart, 2013). Moreover, the blood-pressure lowering effect of the DASH dietary pattern in our meta-analysis was comparable to drug monotherapy (Elmer et al., 2006).

The importance of blood pressure reduction, as shown for several dietary approaches is strengthened by a large meta-analysis of epidemiological studies which have shown that a decrease of approximately 10 mmHg reduction in SBP was inversely associated with risk of

cardiovascular disease events by 20%, coronary heart disease by 17%, stroke by 27%, heart failure by 28% and all-cause mortality by 13% (Ettehad et al., 2016), whereas a 5 mmHg decrease in DBP reduces the risk of ischemic heart disease by 20%, and the risk of stroke by 32% (Law et al., 2003). Another meta-analysis stressed the fact that even a small decline in SBP of about 2 mm Hg will be accompanied by a 10% lower risk of death due to stroke or 7% due to ischemic heart disease (Lewington et al., 2002). Albeit the overall net effect was modest for several dietary approaches, it is important to note that the control/low-fat groups also experienced some benefit and the results were incremental reductions experienced by those groups adopting dietary interventions.

Although this is the first network meta-analysis to assess the comparative effects of different dietary approaches, several previous pairwise meta-analyses have been published, either including only one dietary approach or mixing normotensive and hypertensive patients. A recent standard pairwise meta-analysis of 24 randomized trials including 23858 normotensive and hypertensive participants showed that the DASH, the Mediterranean, a low-sodium and a low-calorie diet were all effective in reducing blood pressure, whereas the Mediterranean diet did not significantly decrease SBP compared to control diets (Gay et al., 2016). Similar to our findings, previous meta-analyses reported a strong reduction in SBP and DBP by the DASH diet (Ndanuko et al., 2016; Saneei et al., 2014; Siervo et al., 2015).

The DASH dietary pattern is based on several food groups like whole grains, fruits, vegetables, dairy products (mainly low-fat), and nuts and lower intakes of red and processed meat as well as and sweets (Schwingshackl and Hoffmann, 2015b). Optimal consumption of risk-decreasing foods (whole grains, fruits, nuts, legumes and dairy products) resulted in a 41% reduction of

hypertension compared to non-consumption of these foods (Schwingshackl et al., 2017b). One potential explanation for the superiority of the DASH diet compared to the Mediterranean diet is the recommendation to increase dairy products and at the same time to lower sodium intake (Sacks et al., 2001) in the former regime, whereas the latter discourages higher intakes of dairy products and does not consider sodium in the diet (Trichopoulou et al., 2003). The original DASH diet was tested independent of sodium intake (Appel et al., 1997). However, 6 out of the included 7 trials investigating the DASH approach in the present NMA tested the effects of the refined DASH diet implementing sodium restriction, whereas one trial did not provide the corresponding information (Edwards et al., 2011).

Anti-hypertensive effects of DASH dietary components are biologically plausible, given the high intake of constituents like phytochemicals and nutrients such as magnesium, potassium, calcium, lactotripeptides, selenium, zinc, antioxidants, vitamins, unsaturated fatty acid, and fibre which have been shown to lower blood pressure (Alonso et al., 2006; Appel et al., 2006; Fekete et al., 2015; Han et al., 2017; Houston, 2011; Schwingshackl et al., 2011; Whelton et al., 2005).

Whereas the evidence for an antihypertensive effect of the DASH diet is highly consistent, inconsistent results were reported for the Mediterranean Diet. One recent meta-analysis of six trials including more than 7,000 participants found insufficient evidence to suggest that the Mediterranean Diet decreased blood pressure (Nissensohn et al., 2016). In contrast, another recent meta-analysis showed that the Mediterranean dietary approach was effective in reducing SBP and DBP (Ndanuko et al., 2016).

Across all dietary approaches, only the low-sodium diet was rated with a high quality of meta-evidence. The blood pressure lowering effect observed in the present meta-analysis is consistent

with several pairwise meta-analyses of randomized trials (Aburto et al., 2013). In line with our and previous findings the European Society of Hypertension and Cardiology rated the recommendation for the prevention and management of hypertension of a maximum daily salt intake of 5-6 gram with the highest level of evidence (Mancia et al., 2013b).

Previous meta-analyses reported conflicting results regarding the blood pressure lowering effects of LC and HP diets (Bueno et al., 2013; Mansoor et al., 2016; Santesso et al., 2012; Schwingshackl and Hoffmann, 2013b; Wycherley et al., 2012). Although in the present network meta-analysis the LC and the HP diets were more effective in reducing blood pressure compared to control diets, the effects compared to LF diets were not statistically significant.

The results based on the Palaeolithic, Vegetarian, Nordic, and Tibetan dietary approach should be interpreted with caution, since only a very low number of trials (≤ 2) were available. Nevertheless, our findings for the Palaeolithic diet are in line with a previous pairwise meta-analysis showing a reduction in SBP (Manheimer et al., 2015). Blood pressure reductions of the Vegetarian diet and the Nordic diet as reported by previous meta-analyses could not be confirmed (Ndanuko et al., 2016; Yokoyama et al., 2014) in pre-hypertensive and hypertensive individuals. Similar to our findings, a recent meta-analysis comparing LGI/LGL vs. high-GI diets showed evidence of a blood pressure reducing effect of low GI/GL diets (Evans et al., 2017).

Beside diet and physical activity, but very likely mediated via these measures, weight loss is the third big lifestyle pillar in the prevention and management of hypertension (Mancia et al., 2013a). In a comprehensive dose-response meta-analysis, each kilogram of weight loss showed a reduction of approximately 1 mmHg of diastolic and systolic blood pressure, respectively (Neter et al., 2003). In line with these observations, the meta-regression analysis showed that larger

differences in body weight changes between arms within trials were associated with a stronger impact on SBP and DBP, compared to smaller differences.

Strengths & Limitations

The present systematic review has several strengths including the application of network meta-analyses methodology, which simultaneously combined direct and indirect evidence, the stringent inclusion and exclusion criteria, the high number of included dietary approaches, participants and trials, the comprehensive literature search, the assessment of the risk of bias, subgroup and sensitivity analyses, the a priori published systematic review protocol, the quality of meta-evidence assessment, and the overall low observed heterogeneity.

Several limitations should be considered when interpreting the results. First, all analyses were based on the original intended randomized design, not by adherence to the actual dietary approach and/or macronutrient composition and caloric intake consumed. This means that although participants were randomized to various dietary approaches, details on their actual adherence to the dietary program were not accounted for in the analyses. Second, 44 of 67 trials were rated as unclear or high risk of bias mostly due to lack of allocation concealment and blinding. The risk of bias sensitivity analysis was based on 23 trials, and only the primary results for the DASH dietary pattern could be confirmed. Third, was the substantial clinical heterogeneity for the included populations in terms of age, sex, study methods, study duration, sample size, and differences in food intake included across the various dietary pattern. Blood pressure reduction was lower in long-term studies, suggesting that participant's adherence decline over time. Moreover, larger size studies (sample size: ≥ 100) had a lower effect on blood pressure reduction compared to studies with a smaller sample size (< 100). Fourth, due to the low

number of studies including hypertensive patients, we were not able to run a sensitivity network meta-analysis. Finally, several nodes were not well connected, due to the low number of trials and this should be taken into account when interpreting the results of the present network meta-analysis.

Conclusion

Our network meta-analysis suggest that the DASH dietary approach might be the most effective dietary approach to improve blood pressure in pre-hypertensive and hypertensive patients based on high quality evidence. The findings of the present network meta-analysis have important clinical and public health implications, suggesting that dietary modifications characterized by high consumption of fruit and vegetables, whole grains, legumes, nuts and seeds, and dairy products and low consumption of red and processed meat, sugar sweetened beverages and sodium are an effective method for controlling blood pressure within pre-hypertensive and hypertensive populations.

Declarations

Ethics approval and consent to participate: NA

Consent for publication: NA

Availability of data and material: NA

Competing interests: Authors declare to have no competing of interest

Funding: No funding received

Authors' contributions: LS, AC, HB, GH contributed to the conception and design of the systematic review and meta-analysis. LS, MP, HB, were involved in the acquisition and analysis of the data. LS, CS, HB, interpreted the results. LS, AC, GH, CS, ET, HB, drafted this manuscript. All authors provided critical revisions of the protocol and approved submission of the final manuscript.

Acknowledgements: We thank Thomas Wolever, PhD for sending us additional data for the corresponding meta-analysis.

References

- Aburto, N. J., Ziolkovska, A., Hooper, L., Elliott, P., Cappuccio, F. P., and Meerpohl, J. J. (2013). Effect of lower sodium intake on health: systematic review and meta-analyses. *BMJ*. **346**: f1326.
- Alli, C., Avanzini, F., Bettelli, G., Bonati, M., Colombo, F., Corso, R., Di Tullio, M., Gentile, M. G., Sangalli, L., Taioli, E., and et al. (1992). Feasibility of a long-term low-sodium diet in mild hypertension. *J Hum Hypertens*. **6**: 281-286.
- Alonso, A., Beunza, J. J., Bes-Rastrollo, M., Pajares, R. M., and Martinez-Gonzalez, M. A. (2006). Vegetable protein and fiber from cereal are inversely associated with the risk of hypertension in a Spanish cohort. *Arch Med Res*. **37**: 778-786.
- Anderssen, S., Holme, I., Urdal, P., and Hjermann, I. (1995). Diet and exercise intervention have favourable effects on blood pressure in mild hypertensives: the Oslo Diet and Exercise Study (ODES). *Blood Press*. **4**: 343-349.
- Andrews, R. C., Cooper, A. R., Montgomery, A. A., Norcross, A. J., Peters, T. J., Sharp, D. J., Jackson, N., Fitzsimons, K., Bright, J., Coulman, K., England, C. Y., Gorton, J., McLenaghan, A., Paxton, E., Polet, A., Thompson, C., and Dayan, C. M. (2011). Diet or diet plus physical activity versus usual care in patients with newly diagnosed type 2 diabetes: the Early ACTID randomised controlled trial. *Lancet*. **378**: 129-139.

- Appel, L. J., Brands, M. W., Daniels, S. R., Karanja, N., Elmer, P. J., and Sacks, F. M. (2006). Dietary approaches to prevent and treat hypertension: a scientific statement from the American Heart Association. *Hypertension*. **47**: 296-308.
- Appel, L. J., Champagne, C. M., Harsha, D. W., Cooper, L. S., Obarzanek, E., Elmer, P. J., Stevens, V. J., Vollmer, W. M., Lin, P. H., Svetkey, L. P., Stedman, S. W., and Young, D. R. (2003). Effects of comprehensive lifestyle modification on blood pressure control: main results of the PREMIER clinical trial. *JAMA*. **289**: 2083-2093.
- Appel, L. J., Moore, T. J., Obarzanek, E., Vollmer, W. M., Svetkey, L. P., Sacks, F. M., Bray, G. A., Vogt, T. M., Cutler, J. A., Windhauser, M. M., Lin, P. H., and Karanja, N. (1997). A clinical trial of the effects of dietary patterns on blood pressure. DASH Collaborative Research Group. *N Engl J Med*. **336**: 1117-1124.
- Ard, J. D., Gower, B., Hunter, G., Ritchie, C. S., Roth, D. L., Goss, A., Wingo, B. C., Bodner, E. V., Brown, C. J., Bryan, D., Buys, D. R., Haas, M. C., Keita, A. D., Flagg, L. A., Williams, C. P., and Locher, J. L. (2016). Effects of Calorie Restriction in Obese Older Adults: The CROSSROADS Randomized Controlled Trial. *J Gerontol A Biol Sci Med Sci*. **73**: 73-80.
- Azadbakht, L., Mirmiran, P., Esmailzadeh, A., Azizi, T., and Azizi, F. (2005). Beneficial effects of a Dietary Approaches to Stop Hypertension eating plan on features of the metabolic syndrome. *Diabetes Care*. **28**: 2823-2831.

- Blumenthal, J. A., Babyak, M. A., Hinderliter, A., Watkins, L. L., Craighead, L., Lin, P. H., Caccia, C., Johnson, J., Waugh, R., and Sherwood, A. (2010). Effects of the DASH diet alone and in combination with exercise and weight loss on blood pressure and cardiovascular biomarkers in men and women with high blood pressure: the ENCORE study. *Arch Intern Med.* **170**: 126-135.
- Brehm, B. J., Lattin, B. L., Summer, S. S., Boback, J. A., Gilchrist, G. M., Jandacek, R. J., and D'Alessio, D. A. (2009). One-year comparison of a high-monounsaturated fat diet with a high-carbohydrate diet in type 2 diabetes. *Diabetes Care.* **32**: 215-220.
- Brinkworth, G. D., Noakes, M., Keogh, J. B., Luscombe, N. D., Wittert, G. A., and Clifton, P. M. (2004a). Long-term effects of a high-protein, low-carbohydrate diet on weight control and cardiovascular risk markers in obese hyperinsulinemic subjects. *Int J Obes Relat Metab Disord.* **28**: 661-670.
- Brinkworth, G. D., Noakes, M., Parker, B., Foster, P., and Clifton, P. M. (2004b). Long-term effects of advice to consume a high-protein, low-fat diet, rather than a conventional weight-loss diet, in obese adults with type 2 diabetes: one-year follow-up of a randomised trial. *Diabetologia.* **47**: 1677-1686.
- Bucher, H. C., Guyatt, G. H., Griffith, L. E., and Walter, S. D. (1997). The results of direct and indirect treatment comparisons in meta-analysis of randomized controlled trials. *J Clin Epidemiol.* **50**: 683-691.

- Bueno, N. B., de Melo, I. S., de Oliveira, S. L., and da Rocha Ataíde, T. (2013). Very-low-carbohydrate ketogenic diet v. low-fat diet for long-term weight loss: a meta-analysis of randomised controlled trials. *Br J Nutr.* **110**: 1178-1187.
- Campbell, D. D., and Meckling, K. A. (2012). Effect of the protein:carbohydrate ratio in hypoenergetic diets on metabolic syndrome risk factors in exercising overweight and obese women. *Br J Nutr.* **108**: 1658-1671.
- Chaimani, A., Caldwell, D. M., Li, T., Higgins, J. P., and Salanti, G. (2017). Additional considerations are required when preparing a protocol for a systematic review with multiple interventions. *J Clin Epidemiol.* **83**: 65-74.
- Chaimani, A., Higgins, J. P., Mavridis, D., Spyridonos, P., and Salanti, G. (2013). Graphical tools for network meta-analysis in STATA. *PLoS One.* **8**: e76654.
- Chaimani, A., and Salanti, G. (2015). Visualizing assumptions and results in network meta-analysis: The network graphs package. *Stata Journal.* **15**: 905-950.
- Chalmers, J., Morgan, T., Doyle, A., Dickson, B., Hopper, J., Mathews, J., Matthews, G., Moulds, R., Myers, J., Nowson, C., and et al. (1986). Australian National Health and Medical Research Council dietary salt study in mild hypertension. *J Hypertens Suppl.* **4**: S629-637.

- Clifton, P. M., Noakes, M., and Keogh, J. B. (2004). Very low-fat (12%) and high monounsaturated fat (35%) diets do not differentially affect abdominal fat loss in overweight, nondiabetic women. *J Nutr.* **134**: 1741-1745.
- Coppell, K. J., Kataoka, M., Williams, S. M., Chisholm, A. W., Vorgers, S. M., and Mann, J. I. (2010). Nutritional intervention in patients with type 2 diabetes who are hyperglycaemic despite optimised drug treatment--Lifestyle Over and Above Drugs in Diabetes (LOADD) study: randomised controlled trial. *BMJ.* **341**: c3337.
- Cornelissen, V. A., and Smart, N. A. (2013). Exercise training for blood pressure: a systematic review and meta-analysis. *J Am Heart Assoc.* **2**: e004473.
- Costa, F. V., Ambrosioni, E., Montebugnoli, L., Paccaloni, L., Vasconi, L., and Magnani, B. (1981). Effects of a low-salt diet and of acute salt loading on blood pressure and intralymphocytic sodium concentration in young subjects with borderline hypertension. *Clin Sci (Lond).* **61 Suppl 7**: 21s-23s.
- Dalle Grave, R., Calugi, S., Gavasso, I., El Ghoch, M., and Marchesini, G. (2013). A randomized trial of energy-restricted high-protein versus high-carbohydrate, low-fat diet in morbid obesity. *Obesity (Silver Spring).* **21**: 1774-1781.
- Daly, M. E., Paisey, R., Paisey, R., Millward, B. A., Eccles, C., Williams, K., Hammersley, S., MacLeod, K. M., and Gale, T. J. (2006). Short-term effects of severe dietary

- carbohydrate-restriction advice in Type 2 diabetes--a randomized controlled trial. *Diabet Med.* **23**: 15-20.
- Dansinger, M. L., Gleason, J. A., Griffith, J. L., Selker, H. P., and Schaefer, E. J. (2005). Comparison of the Atkins, Ornish, Weight Watchers, and Zone diets for weight loss and heart disease risk reduction: a randomized trial. *JAMA.* **293**: 43-53.
- Delbridge, E. A., Prendergast, L. A., Pritchard, J. E., and Proietto, J. (2009). One-year weight maintenance after significant weight loss in healthy overweight and obese subjects: does diet composition matter? *Am J Clin Nutr.* **90**: 1203-1214.
- Deluis, D. A., Sagrado, M. G., Aller, R., Izaola, O., and Conde, R. (2010). Effects of C358A missense polymorphism of the degrading enzyme fatty acid amide hydrolase on weight loss, adipocytokines, and insulin resistance after 2 hypocaloric diets. *Metabolism.* **59**: 1387-1392.
- Dias, S., Welton, N. J., Caldwell, D. M., and Ades, A. E. (2010). Checking consistency in mixed treatment comparison meta-analysis. *Stat Med.* **29**: 932-944.
- Dodson, P. M., Beevers, M., Hallworth, R., Webberley, M. J., Fletcher, R. F., and Taylor, K. G. (1989). Sodium restriction and blood pressure in hypertensive type II diabetics: randomised blind controlled and crossover studies of moderate sodium restriction and sodium supplementation. *BMJ.* **298**: 227-230.

Eckel, R. H., Jakicic, J. M., Ard, J. D., de Jesus, J. M., Houston Miller, N., Hubbard, V. S., Lee, I. M., Lichtenstein, A. H., Loria, C. M., Millen, B. E., Nonas, C. A., Sacks, F. M., Smith, S. C., Jr., Svetkey, L. P., Wadden, T. A., Yanovski, S. Z., Kendall, K. A., Morgan, L. C., Trisolini, M. G., Velasco, G., Wnek, J., Anderson, J. L., Halperin, J. L., Albert, N. M., Bozkurt, B., Brindis, R. G., Curtis, L. H., DeMets, D., Hochman, J. S., Kovacs, R. J., Ohman, E. M., Pressler, S. J., Sellke, F. W., Shen, W. K., Smith, S. C., Jr., and Tomaselli, G. F. (2014). 2013 AHA/ACC guideline on lifestyle management to reduce cardiovascular risk: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *Circulation*. **129**: S76-99.

Edwards, K. M., Wilson, K. L., Sadjia, J., Ziegler, M. G., and Mills, P. J. (2011). Effects on blood pressure and autonomic nervous system function of a 12-week exercise or exercise plus DASH-diet intervention in individuals with elevated blood pressure. *Acta Physiol (Oxf)*. **203**: 343-350.

Elmer, P. J., Obarzanek, E., Vollmer, W. M., Simons-Morton, D., Stevens, V. J., Young, D. R., Lin, P. H., Champagne, C., Harsha, D. W., Svetkey, L. P., Ard, J., Brantley, P. J., Proschian, M. A., Erlinger, T. P., and Appel, L. J. (2006). Effects of comprehensive lifestyle modification on diet, weight, physical fitness, and blood pressure control: 18-month results of a randomized trial. *Ann Intern Med*. **144**: 485-495.

- Erwtaman, T. M., Nagelkerke, N., Lubsen, J., Koster, M., and Dunning, A. J. (1984). Beta blockade, diuretics, and salt restriction for the management of mild hypertension: a randomised double blind trial. *Br Med J (Clin Res Ed)*. **289**: 406-409.
- Esposito, K., Maiorino, M. I., Ciotola, M., Di Palo, C., Scognamiglio, P., Gicchino, M., Petrizzo, M., Saccomanno, F., Beneduce, F., Ceriello, A., and Giugliano, D. (2009). Effects of a Mediterranean-style diet on the need for antihyperglycemic drug therapy in patients with newly diagnosed type 2 diabetes: a randomized trial. *Ann Intern Med*. **151**: 306-314.
- Esposito, K., Marfella, R., Ciotola, M., Di Palo, C., Giugliano, F., Giugliano, G., D'Armiento, M., D'Andrea, F., and Giugliano, D. (2004). Effect of a mediterranean-style diet on endothelial dysfunction and markers of vascular inflammation in the metabolic syndrome: a randomized trial. *JAMA*. **292**: 1440-1446.
- Ettehad, D., Emdin, C. A., Kiran, A., Anderson, S. G., Callender, T., Emberson, J., Chalmers, J., Rodgers, A., and Rahimi, K. (2016). Blood pressure lowering for prevention of cardiovascular disease and death: a systematic review and meta-analysis. *Lancet*. **387**: 957-967.
- Evans, C. E., Greenwood, D. C., Threapleton, D. E., Gale, C. P., Cleghorn, C. L., and Burley, V. J. (2017). Glycemic index, glycemic load, and blood pressure: a systematic review and meta-analysis of randomized controlled trials. *Am J Clin Nutr*. **105**: 1176-1190.

- Fekete, A. A., Givens, D. I., and Lovegrove, J. A. (2015). Casein-derived lactotripeptides reduce systolic and diastolic blood pressure in a meta-analysis of randomised clinical trials. *Nutrients*. **7**: 659-681.
- Frisch, S., Zittermann, A., Berthold, H. K., Gotting, C., Kuhn, J., Kleesiek, K., Stehle, P., and Kortke, H. (2009). A randomized controlled trial on the efficacy of carbohydrate-reduced or fat-reduced diets in patients attending a telemedically guided weight loss program. *Cardiovasc Diabetol*. **8**: 36.
- Gay, H. C., Rao, S. G., Vaccarino, V., and Ali, M. K. (2016). Effects of Different Dietary Interventions on Blood Pressure: Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Hypertension*. **67**: 733-739.
- Gordon, N. F., Scott, C. B., and Levine, B. D. (1997). Comparison of single versus multiple lifestyle interventions: are the antihypertensive effects of exercise training and diet-induced weight loss additive? *Am J Cardiol*. **79**: 763-767.
- Guldbrand, H., Dizdar, B., Bunjaku, B., Lindstrom, T., Bachrach-Lindstrom, M., Fredrikson, M., Ostgren, C. J., and Nystrom, F. H. (2012). In type 2 diabetes, randomisation to advice to follow a low-carbohydrate diet transiently improves glycaemic control compared with advice to follow a low-fat diet producing a similar weight loss. *Diabetologia*. **55**: 2118-2127.

- Haider, L. M., Schwingshackl, L., Hoffmann, G., and Ekmekcioglu, C. (2017). The effect of vegetarian diets on iron status in adults: A systematic review and meta-analysis. *Critical Reviews in Food Science and Nutrition*. **0**: 1-16.
- Han, H., Fang, X., Wei, X., Liu, Y., Jin, Z., Chen, Q., Fan, Z., Aaseth, J., Hiyoshi, A., He, J., and Cao, Y. (2017). Dose-response relationship between dietary magnesium intake, serum magnesium concentration and risk of hypertension: a systematic review and meta-analysis of prospective cohort studies. *Nutr J*. **16**: 26.
- Heilbronn, L. K., Noakes, M., and Clifton, P. M. (1999). Effect of energy restriction, weight loss, and diet composition on plasma lipids and glucose in patients with type 2 diabetes. *Diabetes Care*. **22**: 889-895.
- Higgins, J. P., Altman, D. G., Gotzsche, P. C., Juni, P., Moher, D., Oxman, A. D., Savovic, J., Schulz, K. F., Weeks, L., and Sterne, J. A. (2011). The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ*. **343**: d5928.
- Higgins, J. P., Jackson, D., Barrett, J. K., Lu, G., Ades, A. E., and White, I. R. (2012). Consistency and inconsistency in network meta-analysis: concepts and models for multi-arm studies. *Res Synth Methods*. **3**: 98-110.
- Higgins, J. P. T., and Green, S. (2011). Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011. Available from www.cochrane-handbook.org.

Houston, M. C. (2011). The importance of potassium in managing hypertension. *Curr Hypertens Rep.* **13**: 309-317.

Hutton, B., Salanti, G., Caldwell, D. M., Chaimani, A., Schmid, C. H., Cameron, C., Ioannidis, J. P., Straus, S., Thorlund, K., Jansen, J. P., Mulrow, C., Catala-Lopez, F., Gotzsche, P. C., Dickersin, K., Boutron, I., Altman, D. G., and Moher, D. (2015). The PRISMA extension statement for reporting of systematic reviews incorporating network meta-analyses of health care interventions: checklist and explanations. *Ann Intern Med.* **162**: 777-784.

Iqbal, N., Vetter, M. L., Moore, R. H., Chittams, J. L., Dalton-Bakes, C. V., Dowd, M., Williams-Smith, C., Cardillo, S., and Wadden, T. A. (2010). Effects of a low-intensity intervention that prescribed a low-carbohydrate vs. a low-fat diet in obese, diabetic participants. *Obesity (Silver Spring).* **18**: 1733-1738.

Itsiopoulos, C., Brazionis, L., Kaimakamis, M., Cameron, M., Best, J. D., O'Dea, K., and Rowley, K. (2011). Can the Mediterranean diet lower HbA1c in type 2 diabetes? Results from a randomized cross-over study. *Nutr Metab Cardiovasc Dis.* **21**: 740-747.

Jackson, D., Barrett, J. K., Rice, S., White, I. R., and Higgins, J. P. (2014). A design-by-treatment interaction model for network meta-analysis with random inconsistency effects. *Stat Med.* **33**: 3639-3654.

Jonsson, T., Granfeldt, Y., Ahren, B., Branell, U. C., Palsson, G., Hansson, A., Soderstrom, M., and Lindeberg, S. (2009). Beneficial effects of a Paleolithic diet on cardiovascular risk

factors in type 2 diabetes: a randomized cross-over pilot study. *Cardiovasc Diabetol.* **8**: 35.

Jula, A. M., and Karanko, H. M. (1994). Effects on left ventricular hypertrophy of long-term nonpharmacological treatment with sodium restriction in mild-to-moderate essential hypertension. *Circulation.* **89**: 1023-1031.

Kim, H. H., Kim, Y. J., Lee, S. Y., Jeong, D. W., Lee, J. G., Yi, Y. H., Cho, Y. H., Choi, E. J., and Kim, H. J. (2014). Interactive effects of an isocaloric high-protein diet and resistance exercise on body composition, ghrelin, and metabolic and hormonal parameters in untrained young men: A randomized clinical trial. *J Diabetes Investig.* **5**: 242-247.

Kirpizidis, H., Stavratsi, A., and Geleris, P. (2005). Assessment of quality of life in a randomized clinical trial of candesartan only or in combination with DASH diet for hypertensive patients. *J Cardiol.* **46**: 177-182.

Klemsdal, T. O., Holme, I., Nerland, H., Pedersen, T. R., and Tonstad, S. (2010). Effects of a low glycemic load diet versus a low-fat diet in subjects with and without the metabolic syndrome. *Nutr Metab Cardiovasc Dis.* **20**: 195-201.

Krahn, U., Binder, H., and König, J. (2013). A graphical tool for locating inconsistency in network meta-analyses. *BMC Med Res Methodol.* **13**: 35.

- Krebs, J. D., Elley, C. R., Parry-Strong, A., Lunt, H., Drury, P. L., Bell, D. A., Robinson, E., Moyes, S. A., and Mann, J. I. (2012). The Diabetes Excess Weight Loss (DEWL) Trial: a randomised controlled trial of high-protein versus high-carbohydrate diets over 2 years in type 2 diabetes. *Diabetologia*. **55**: 905-914.
- Law, M., Wald, N., and Morris, J. (2003). Lowering blood pressure to prevent myocardial infarction and stroke: a new preventive strategy. *Health Technol Assess*. **7**: 1-94.
- Lawes, C. M., Vander Hoorn, S., and Rodgers, A. (2008). Global burden of blood-pressure-related disease, 2001. *Lancet*. **371**: 1513-1518.
- Lean, M. E., Han, T. S., Prvan, T., Richmond, P. R., and Avenell, A. (1997). Weight loss with high and low carbohydrate 1200 kcal diets in free living women. *Eur J Clin Nutr*. **51**: 243-248.
- Leucht, S., Chaimani, A., Cipriani, A. S., Davis, J. M., Furukawa, T. A., and Salanti, G. (2016). Network meta-analyses should be the highest level of evidence in treatment guidelines. *Eur Arch Psychiatry Clin Neurosci*. **266**: 477-480.
- Lewington, S., Clarke, R., Qizilbash, N., Peto, R., and Collins, R. (2002). Age-specific relevance of usual blood pressure to vascular mortality: a meta-analysis of individual data for one million adults in 61 prospective studies. *Lancet*. **360**: 1903-1913.

- Lima, S. T., da Silva Nalin de Souza, B., Franca, A. K., Salgado Filho, N., and Sichieri, R. (2013). Dietary approach to hypertension based on low glycaemic index and principles of DASH (Dietary Approaches to Stop Hypertension): a randomised trial in a primary care service. *Br J Nutr.* **110**: 1472-1479.
- Liu, X., Zhang, G., Ye, X., Li, H., Chen, X., Tang, L., Feng, Y., Shai, I., Stampfer, M. J., Hu, F. B., and Lin, X. (2013). Effects of a low-carbohydrate diet on weight loss and cardiometabolic profile in Chinese women: a randomised controlled feeding trial. *Br J Nutr.* **110**: 1444-1453.
- Luger, M., Holstein, B., Schindler, K., Kruschitz, R., and Ludvik, B. (2013). Feasibility and efficacy of an isocaloric high-protein vs. standard diet on insulin requirement, body weight and metabolic parameters in patients with type 2 diabetes on insulin therapy. *Exp Clin Endocrinol Diabetes.* **121**: 286-294.
- Ma, Y., Olendzki, B. C., Merriam, P. A., Chiriboga, D. E., Culver, A. L., Li, W., Hebert, J. R., Ockene, I. S., Griffith, J. A., and Pagoto, S. L. (2008). A randomized clinical trial comparing low-glycemic index versus ADA dietary education among individuals with type 2 diabetes. *Nutrition.* **24**: 45-56.
- Makecia, P., Vahlberg, T., Kantola, I., Vesalainen, R., and Jula, A. (2008). The effects of a 6-month sodium restriction on cardiac autonomic function in patients with mild to moderate essential hypertension. *Am J Hypertens.* **21**: 1183-1187.

Mancia, G., Fagard, R., Narkiewicz, K., Redon, J., Zanchetti, A., Bohm, M., Christiaens, T., Cifkova, R., De Backer, G., Dominiczak, A., Galderisi, M., Grobbee, D. E., Jaarsma, T., Kirchhof, P., Kjeldsen, S. E., Laurent, S., Manolis, A. J., Nilsson, P. M., Ruilope, L. M., Schmieder, R. E., Sirnes, P. A., Sleight, P., Viigimaa, M., Waeber, B., and Zannad, F. (2013a). 2013 ESH/ESC Guidelines for the management of arterial hypertension: the Task Force for the management of arterial hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). *J Hypertens.* **31**: 1281-1357.

Mancia, G., Fagard, R., Narkiewicz, K., Redon, J., Zanchetti, A., Bohm, M., Christiaens, T., Cifkova, R., De Backer, G., Dominiczak, A., Galderisi, M., Grobbee, D. E., Jaarsma, T., Kirchhof, P., Kjeldsen, S. E., Laurent, S., Manolis, A. J., Nilsson, P. M., Ruilope, L. M., Schmieder, R. E., Sirnes, P. A., Sleight, P., Viigimaa, M., Waeber, B., Zannad, F., Redon, J., Dominiczak, A., Narkiewicz, K., Nilsson, P. M., Burnier, M., Viigimaa, M., Ambrosioni, E., Caulfield, M., Coca, A., Olsen, M. H., Schmieder, R. E., Tsioufis, C., van de Borne, P., Zamorano, J. L., Achenbach, S., Baumgartner, H., Bax, J. J., Bueno, H., Dean, V., Deaton, C., Erol, C., Fagard, R., Ferrari, R., Hasdai, D., Hoes, A. W., Kirchhof, P., Knuuti, J., Kolh, P., Lancellotti, P., Linhart, A., Nihoyannopoulos, P., Piepoli, M. F., Ponikowski, P., Sirnes, P. A., Tamargo, J. L., Tendera, M., Torbicki, A., Wijns, W., Windecker, S., Clement, D. L., Coca, A., Gillebert, T. C., Tendera, M., Rosei, E. A., Ambrosioni, E., Anker, S. D., Bauersachs, J., Hitij, J. B., Caulfield, M., De Buyzere, M., De Geest, S., Derumeaux, G. A., Erdine, S., Farsang, C., Funck-Brentano, C., Gerc, V., Germano, G., Gielen, S., Haller, H., Hoes, A. W., Jordan, J., Kahan, T., Komajda, M.,

- Lovic, D., Mahrholdt, H., Olsen, M. H., Ostergren, J., Parati, G., Perk, J., Polonia, J., Popescu, B. A., Reiner, Z., Ryden, L., Sirenko, Y., Stanton, A., Struijker-Boudier, H., Tsioufis, C., van de Borne, P., Vlachopoulos, C., Volpe, M., and Wood, D. A. (2013b). 2013 ESH/ESC guidelines for the management of arterial hypertension: the Task Force for the Management of Arterial Hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). *Eur Heart J* **34**: 2159-2219.
- Manheimer, E. W., van Zuuren, E. J., Fedorowicz, Z., and Pijl, H. (2015). Paleolithic nutrition for metabolic syndrome: systematic review and meta-analysis. *Am J Clin Nutr*. **102**: 922-932.
- Mansoor, N., Vinknes, K. J., Veierod, M. B., and Retterstol, K. (2016). Effects of low-carbohydrate diets v. low-fat diets on body weight and cardiovascular risk factors: a meta-analysis of randomised controlled trials. *Br J Nutr*. **115**: 466-479.
- Mavridis, D., Giannatsi, M., Cipriani, A., and Salanti, G. (2015). A primer on network meta-analysis with emphasis on mental health. *Evid Based Ment Health*. **18**: 40-46.
- Mellberg, C., Sandberg, S., Ryberg, M., Eriksson, M., Brage, S., Larsson, C., Olsson, T., and Lindahl, B. (2014). Long-term effects of a Palaeolithic-type diet in obese postmenopausal women: a 2-year randomized trial. *Eur J Clin Nutr*. **68**: 350-357.

NCD-Risk-Factor-Collaboration (2017). Worldwide trends in blood pressure from 1975 to 2015:

a pooled analysis of 1479 population-based measurement studies with 19.1 million participants. *Lancet*. **389**: 37-55.

Ndanuko, R. N., Tapsell, L. C., Charlton, K. E., Neale, E. P., and Batterham, M. J. (2016).

Dietary Patterns and Blood Pressure in Adults: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Adv Nutr*. **7**: 76-89.

Neter, J. E., Stam, B. E., Kok, F. J., Grobbee, D. E., and Geleijnse, J. M. (2003). Influence of

weight reduction on blood pressure: a meta-analysis of randomized controlled trials. *Hypertension*. **42**: 878-884.

Nicholson, A. S., Sklar, M., Barnard, N. D., Gore, S., Sullivan, R., and Browning, S. (1999).

Toward improved management of NIDDM: A randomized, controlled, pilot intervention using a lowfat, vegetarian diet. *Prev Med*. **29**: 87-91.

Nissensohn, M., Roman-Vinas, B., Sanchez-Villegas, A., Piscopo, S., and Serra-Majem, L.

(2016). The Effect of the Mediterranean Diet on Hypertension: A Systematic Review and Meta-Analysis. *J Nutr Educ Behav*. **48**: 42-53.e41.

Nowson, C. A., Worsley, A., Margetison, C., Jorna, M. K., Godfrey, S. J., and Booth, A. (2005).

Blood pressure change with weight loss is affected by diet type in men. *Am J Clin Nutr*. **81**: 983-989.

- Philippou, E., Bovill-Taylor, C., Rajkumar, C., Vampa, M. L., Ntatsaki, E., Brynes, A. E., Hickson, M., and Frost, G. S. (2009). Preliminary report: the effect of a 6-month dietary glycemic index manipulation in addition to healthy eating advice and weight loss on arterial compliance and 24-hour ambulatory blood pressure in men: a pilot study. *Metabolism*. **58**: 1703-1708.
- Poulsen, S. K., Crone, C., Astrup, A., and Larsen, T. M. (2015). Long-term adherence to the New Nordic Diet and the effects on body weight, anthropometry and blood pressure: a 12-month follow-up study. *Eur J Nutr*. **54**: 67-76.
- Riley, R. D., Higgins, J. P., and Deeks, J. J. (2011). Interpretation of random effects meta-analyses. *BMJ*. **342**: d549.
- Rock, C. L., Flatt, S. W., Pakiz, B., Taylor, K. S., Leone, A. F., Brelje, K., Heath, D. D., Quintana, E. L., and Sherwood, N. E. (2014). Weight loss, glycemic control, and cardiovascular disease risk factors in response to differential diet composition in a weight loss program in type 2 diabetes: a randomized controlled trial. *Diabetes Care*. **37**: 1573-1580.
- Sacks, F. M., Svetkey, L. P., Vollmer, W. M., Appel, L. J., Bray, G. A., Harsha, D., Obarzanek, E., Conlin, P. R., Miller, E. R., 3rd, Simons-Morton, D. G., Karanja, N., and Lin, P. H. (2001). Effects on blood pressure of reduced dietary sodium and the Dietary Approaches to Stop Hypertension (DASH) diet. DASH-Sodium Collaborative Research Group. *N Engl J Med*. **344**: 3-10.

- Salanti, G. (2012). Indirect and mixed-treatment comparison, network, or multiple-treatments meta-analysis: many names, many benefits, many concerns for the next generation evidence synthesis tool. *Res Synth Methods*. **3**: 80-97.
- Salanti, G., Ades, A. E., and Ioannidis, J. P. (2011). Graphical methods and numerical summaries for presenting results from multiple-treatment meta-analysis: an overview and tutorial. *J Clin Epidemiol*. **64**: 163-171.
- Salanti, G., Del Giovane, C., Chaimani, A., Caldwell, D. M., and Higgins, J. P. (2014). Evaluating the quality of evidence from a network meta-analysis. *PLoS One*. **9**: e99682.
- Saneei, P., Salehi-Abargouei, A., Esmailzadeh, A., and Azadbakht, L. (2014). Influence of Dietary Approaches to Stop Hypertension (DASH) diet on blood pressure: a systematic review and meta-analysis on randomized controlled trials. *Nutr Metab Cardiovasc Dis*. **24**: 1253-1261.
- Santesso, N., Akl, E. A., Bianchi, M., Mente, A., Mustafa, R., Heels-Ansdell, D., and Schunemann, H. J. (2012). Effects of higher- versus lower-protein diets on health outcomes: a systematic review and meta-analysis. *Eur J Clin Nutr*. **66**: 780-788.
- Schwingshackl, L., Chaimani, A., Hoffmann, G., Schwedhelm, C., and Boeing, H. (2017a). Impact of different dietary approaches on blood pressure in hypertensive and prehypertensive patients: protocol for a systematic review and network meta-analysis. *BMJ Open*. **7**: e014736.

- Schwingshackl, L., Chaimani, A., Hoffmann, G., Schwedhelm, C., and Boeing, H. (2018). A network meta-analysis on the comparative efficacy of different dietary approaches on glycaemic control in patients with type 2 diabetes mellitus. *Eur J Epidemiol.* **33**: 157-170.
- Schwingshackl, L., Dias, S., Strasser, B., and Hoffmann, G. (2013). Impact of different training modalities on anthropometric and metabolic characteristics in overweight/obese subjects: a systematic review and network meta-analysis. *PLoS One.* **8**: e82853.
- Schwingshackl, L., Hobl, L. P., and Hoffmann, G. (2015a). Effects of low glycaemic index/low glycaemic load vs. high glycaemic index/ high glycaemic load diets on overweight/obesity and associated risk factors in children and adolescents: a systematic review and meta-analysis. *Nutr J.* **14**: 87.
- Schwingshackl, L., and Hoffmann, G. (2013a). Comparison of effects of long-term low-fat vs high-fat diets on blood lipid levels in overweight or obese patients: a systematic review and meta-analysis. *J Acad Nutr Diet.* **113**: 1640-1661.
- Schwingshackl, L., and Hoffmann, G. (2013b). Long-term effects of low-fat diets either low or high in protein on cardiovascular and metabolic risk factors: a systematic review and meta-analysis. *Nutr J.* **12**: 48.
- Schwingshackl, L., and Hoffmann, G. (2013c). Long-term effects of low glycemic index/load vs. high glycemic index/load diets on parameters of obesity and obesity-associated risks: a systematic review and meta-analysis. *Nutr Metab Cardiovasc Dis.* **23**: 699-706.

Schwingshackl, L., and Hoffmann, G. (2014a). Adherence to Mediterranean diet and risk of cancer: a systematic review and meta-analysis of observational studies. *Int J Cancer*. **135**: 1884-1897.

Schwingshackl, L., and Hoffmann, G. (2014b). Comparison of the long-term effects of high-fat v. low-fat diet consumption on cardiometabolic risk factors in subjects with abnormal glucose metabolism: a systematic review and meta-analysis. *Br J Nutr*. **111**: 2047-2058.

Schwingshackl, L., and Hoffmann, G. (2014c). Mediterranean dietary pattern, inflammation and endothelial function: a systematic review and meta-analysis of intervention trials. *Nutr Metab Cardiovasc Dis*. **24**: 929-939.

Schwingshackl, L., and Hoffmann, G. (2015a). Adherence to Mediterranean diet and risk of cancer: an updated systematic review and meta-analysis of observational studies. *Cancer Med*. **4**: 1933-1947.

Schwingshackl, L., and Hoffmann, G. (2015b). Diet quality as assessed by the Healthy Eating Index, the Alternate Healthy Eating Index, the Dietary Approaches to Stop Hypertension score, and health outcomes: a systematic review and meta-analysis of cohort studies. *J Acad Nutr Diet*. **115**: 780-800.e785.

Schwingshackl, L., and Hoffmann, G. (2016). Does a Mediterranean-Type Diet Reduce Cancer Risk? *Current Nutrition Reports*. **5**: 9-17.

- Schwingshackl, L., Missbach, B., Dias, S., Konig, J., and Hoffmann, G. (2014). Impact of different training modalities on glycaemic control and blood lipids in patients with type 2 diabetes: a systematic review and network meta-analysis. *Diabetologia*. **57**: 1789-1797.
- Schwingshackl, L., Missbach, B., Konig, J., and Hoffmann, G. (2015b). Adherence to a Mediterranean diet and risk of diabetes: a systematic review and meta-analysis. *Public Health Nutr*. **18**: 1292-1299.
- Schwingshackl, L., Schwedhelm, C., Hoffmann, G., Knuppel, S., Iqbal, K., Andriolo, V., Bechthold, A., Schlesinger, S., and Boeing, H. (2017b). Food Groups and Risk of Hypertension: A Systematic Review and Dose-Response Meta-Analysis of Prospective Studies. *Adv Nutr*. **8**: 793-803.
- Schwingshackl, L., Strasser, B., and Hoffmann, G. (2011). Effects of monounsaturated fatty acids on cardiovascular risk factors: a systematic review and meta-analysis. *Ann Nutr Metab*. **59**: 176-186.
- Shai, I., Schwarzfuchs, D., Henkin, Y., Shahar, D. R., Witkow, S., Greenberg, I., Golan, R., Fraser, D., Bolotin, A., Vardi, H., Tangi-Rozental, O., Zuk-Ramot, R., Sarusi, B., Brickner, D., Schwartz, Z., Sheiner, E., Marko, R., Katorza, E., Thiery, J., Fiedler, G. M., Bluher, M., Stumvoll, M., and Stampfer, M. J. (2008). Weight loss with a low-carbohydrate, Mediterranean, or low-fat diet. *N Engl J Med*. **359**: 229-241.

Siervo, M., Lara, J., Chowdhury, S., Ashor, A., Oggioni, C., and Mathers, J. C. (2015). Effects of the Dietary Approach to Stop Hypertension (DASH) diet on cardiovascular risk factors: a systematic review and meta-analysis. *Br J Nutr.* **113**: 1-15.

Silman, A. J., Locke, C., Mitchell, P., and Humpherson, P. (1983). Evaluation of the effectiveness of a low sodium diet in the treatment of mild to moderate hypertension. *Lancet.* **1**: 1179-1182.

Smith, S. C., Jr., Benjamin, E. J., Bonow, R. O., Braun, L. T., Creager, M. A., Franklin, B. A., Gibbons, R. J., Grundy, S. M., Hiratzka, L. F., Jones, D. W., Lloyd-Jones, D. M., Minissian, M., Mosca, L., Peterson, E. D., Sacco, R. L., Spertus, J., Stein, J. H., and Taubert, K. A. (2011). AHA/ACCF Secondary Prevention and Risk Reduction Therapy for Patients with Coronary and other Atherosclerotic Vascular Disease: 2011 update: a guideline from the American Heart Association and American College of Cardiology Foundation. *Circulation.* **124**: 2458-2473.

StataCorp. (2015). Stata Statistical Software: Release 14. College Station, TX: StataCorp LP.

Stern, L., Iqbal, N., Seshadri, P., Chicano, K. L., Daily, D. A., McGrory, J., Williams, M., Gracely, E. J., and Samaha, F. F. (2004). The effects of low-carbohydrate versus conventional weight loss diets in severely obese adults: one-year follow-up of a randomized trial. *Ann Intern Med.* **140**: 778-785.

- Tang, M., Armstrong, C. L., Leidy, H. J., and Campbell, W. W. (2013). Normal vs. high-protein weight loss diets in men: effects on body composition and indices of metabolic syndrome. *Obesity (Silver Spring)*. **21**: E204-210.
- Tay, J., Brinkworth, G. D., Noakes, M., Keogh, J., and Clifton, P. M. (2008). Metabolic effects of weight loss on a very-low-carbohydrate diet compared with an isocaloric high-carbohydrate diet in abdominally obese subjects. *J Am Coll Cardiol*. **51**: 59-67.
- Tay, J., Luscombe-Marsh, N. D., Thompson, C. H., Noakes, M., Buckley, J. D., Wittert, G. A., Yancy, W. S., Jr., and Brinkworth, G. D. (2015). Comparison of low- and high-carbohydrate diets for type 2 diabetes management: a randomized trial. *Am J Clin Nutr*. **102**: 780-790.
- Toledo, E., Hu, F. B., Estruch, R., and Salas-Salvad6, J., et al. (2013). Effect of the Mediterranean diet on blood pressure in the PREDIMED trial: results from a randomized controlled trial. *BMC Med*. **11**: 207.
- Trichopoulou, A., Costacou, T., Bamia, C., and Trichopoulos, D. (2003). Adherence to a Mediterranean diet and survival in a Greek population. *N Engl J Med*. **348**: 2599-2608.
- Uusitupa, M., Hermansen, K., Savolainen, M. J., Schwab, U., Kolehmainen, M., Brader, L., Mortensen, L. S., Cloetens, L., Johansson-Persson, A., Onning, G., Landin-Olsson, M., Herzig, K. H., Hukkanen, J., Rosqvist, F., Iggman, D., Paananen, J., Pulkki, K. J., Siloaho, M., Dragsted, L., Barri, T., Overvad, K., Bach Knudsen, K. E., Hedemann, M.

- S., Arner, P., Dahlman, I., Borge, G. I., Baardseth, P., Ulven, S. M., Gunnarsdottir, I., Jonsdottir, S., Thorsdottir, I., Oresic, M., Poutanen, K. S., Riserus, U., and Akesson, B. (2013). Effects of an isocaloric healthy Nordic diet on insulin sensitivity, lipid profile and inflammation markers in metabolic syndrome -- a randomized study (SYSDIET) *J Intern Med.* **274**: 52-66.
- Uusitupa, M., Laitinen, J., Siitonen, O., Vanninen, E., and Pyorala, K. (1993). The maintenance of improved metabolic control after intensified diet therapy in recent type 2 diabetes. *Diabetes Res Clin Pract.* **19**: 227-238.
- Veum, V. L., Laupsa-Borge, J., Eng, O., Rostrup, E., Larsen, T. H., Nordrehaug, J. E., Nygard, O. K., Sagen, J. V., Gudbrandsen, O. A., Dankel, S. N., and Mellgren, G. (2017). Visceral adiposity and metabolic syndrome after very high-fat and low-fat isocaloric diets: a randomized controlled trial. *Am J Clin Nutr.* **105**: 85-99.
- von Haehling, S., Stellos, K., Qusar, N., Gawaz, M., and Bigalke, B. (2013). Weight reduction in patients with coronary artery disease: comparison of Traditional Tibetan Medicine and Western diet. *Int J Cardiol.* **168**: 1509-1515.
- Walker, K. Z., O'Dea, K., Nicholson, G. C., and Muir, J. G. (1995). Dietary composition, body weight, and NIDDM. Comparison of high-fiber, high-carbohydrate, and modified-fat diets. *Diabetes Care.* **18**: 401-403.

- Watkins, L. L., Sherwood, A., Feinglos, M., Hinderliter, A., Babyak, M., Gullette, E., Waugh, R., and Blumenthal, J. A. (2003). Effects of exercise and weight loss on cardiac risk factors associated with syndrome X. *Arch Intern Med.* **163**: 1889-1895.
- Watson, N., Dyer, K., Buckley, J., Brinkworth, G., Coates, A., Parfitt, G., Howe, P., Noakes, M., and Murphy, K. (2016). Effects of Low-Fat Diets Differing in Protein and Carbohydrate Content on Cardiometabolic Risk Factors during Weight Loss and Weight Maintenance in Obese Adults with Type 2 Diabetes. *Nutrients.* **8**.
- Westman, E. C., Yancy, W. S., Jr., Mavropoulos, J. C., Marquart, M., and McDuffie, J. R. (2008). The effect of a low-carbohydrate, ketogenic diet versus a low-glycemic index diet on glycemic control in type 2 diabetes mellitus. *Nutr Metab (Lond).* **5**: 36.
- Whelton, P. e. a. (1997). Effects of weight loss and sodium reduction intervention on blood pressure and hypertension incidence in overweight people with high-normal blood pressure. The Trials of Hypertension Prevention, phase II. The Trials of Hypertension Prevention Collaborative Research Group. *Arch Intern Med.* **157**: 657-667.
- Whelton, P. K., Appel, L. J., Espeland, M. A., Applegate, W. B., Ettinger, W. H., Jr., Kostis, J. B., Kumanyika, S., Lacy, C. R., Johnson, K. C., Folmar, S., and Cutler, J. A. (1998). Sodium reduction and weight loss in the treatment of hypertension in older persons: a randomized controlled trial of nonpharmacologic interventions in the elderly (TONE). TONE Collaborative Research Group. *JAMA.* **279**: 839-846.

Whelton, S. P., Hyre, A. D., Pedersen, B., Yi, Y., Whelton, P. K., and He, J. (2005). Effect of dietary fiber intake on blood pressure: a meta-analysis of randomized, controlled clinical trials. *J Hypertens.* **23**: 475-481.

White, I. R. (2015). Network meta-analysis. *Stata Journal.* **15**: 951-985.

World-Health-Organization (2012). Raised Blood Pressure: Situation and Trends. Global Health Observatory http://www.who.int/gho/ncd/risk_factors/blood_pressure_prevalence/en/ (accessed 04.02.2016).

Wycherley, T. P., Moran, L. J., Clifton, P. M., Noakes, M., and Brinkworth, G. D. (2012). Effects of energy-restricted high-protein, low-fat compared with standard-protein, low-fat diets: a meta-analysis of randomized controlled trials. *Am J Clin Nutr.* **96**: 1281-1298.

Wycherley, T. P., Noakes, M., Clifton, P. M., Cleanthous, X., Keogh, J. B., and Brinkworth, G. D. (2010). A high-protein diet with resistance exercise training improves weight loss and body composition in overweight and obese patients with type 2 diabetes. *Diabetes Care.* **33**: 969-976.

Yokoyama, Y., Nishimura, K., Barnard, N. D., Takegami, M., Watanabe, M., Sekikawa, A., Okamura, T., and Miyamoto, Y. (2014). Vegetarian diets and blood pressure: a meta-analysis. *JAMA Intern Med.* **174**: 577-587.

(Alli et al., 1992; Anderssen et al., 1995; Andrews et al., 2011; Appel et al., 2003; Ard et al., 2016; Azadbakht et al., 2005; Blumenthal et al., 2010; Brehm et al., 2009; Brinkworth et al., 2004a; Brinkworth et al., 2004b; Campbell and Meckling, 2012; Chalmers et al., 1986; Clifton et al., 2004; Coppell et al., 2010; Costa et al., 1981; Dalle Grave et al., 2013; Daly et al., 2006; Dansinger et al., 2005; Delbridge et al., 2009; Deluis et al., 2010; Dodson et al., 1989; Edwards et al., 2011; Erwtaman et al., 1984; Esposito et al., 2009; Esposito et al., 2004; Frisch et al., 2009; Gordon et al., 1997; Guldbrand et al., 2012; Heilbronn et al., 1999; Iqbal et al., 2010; Itsiopoulos et al., 2011; Jonsson et al., 2009; Jula and Karanko, 1994; Kim et al., 2014; Kirpizidis et al., 2005; Klemsdal et al., 2010; Krebs et al., 2012; Lean et al., 1997; Lima et al., 2013; Liu et al., 2013; Luger et al., 2013; Ma et al., 2008; Makela et al., 2008; Mellberg et al., 2014; Nicholson et al., 1999; Nowson et al., 2005; Philippou et al., 2009; Poulsen et al., 2015; Rock et al., 2014; Shai et al., 2008; Silman et al., 1983; Stern et al., 2004; Tang et al., 2013; Tay et al., 2008; Tay et al., 2015; Toledo et al., 2013; Uusitupa et al., 2013; Uusitupa et al., 1993; Veum et al., 2017; von Haehling et al., 2013; Walker et al., 1995; Watkins et al., 2003; Watson et al., 2016; Westman et al., 2008; Whelton, 1997; Whelton et al., 1998; Wycherley et al., 2010)

Table 1: References of the 67 trials included in the present network meta-analysis

DBP																									
DASH	0	10	0	10	0	10	0	10	0	10	0	10	0	10	0	10	0	10	59	41	60	40			
	%	0	%	0	%	0	%	0	%	0	%	0	%	0	%	0	%	0	%	%	%	%			
	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%			
0	10	Mediterranean	3	65	0	10	0	10	0	10	0	10	0	10	0	10	0	10	87	13	0	10			
%	0		5	%	%	0	%	0	%	0	%	0	%	0	%	0	%	0	%	%	%	0			
%	%		%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%			
0	10	3	66	Low-Carb	0	10	0	10	0	10	0	10	1	84	0	10	0	10	0	10	72	28	0	10	
%	0	4	%		%	0	%	0	%	0	%	0	6	%	%	0	%	0	%	0	%	%	0		
%	%	%	%		%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%		
0	10	0	10	0	10	Palaeolithic	0	10	0	10	0	10	0	10	0	10	0	10	0	10	10	0	0	10	
%	0	%	0	%	0		%	0	%	0	%	0	%	0	%	0	%	0	%	0	0	%	%	0	
%	%	%	%	%	%		%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	
0	10	0	10	0	10	0	10	High-Protein	0	10	0	10	0	10	0	10	0	10	0	10	10	0	0	10	
%	0	%	0	%	0	%	0		%	0	%	0	%	0	%	0	%	0	%	0	0	%	%	0	
%	%	%	%	%	%	%	%		%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	
0	10	0	10	0	10	0	10	0	10	Vegetarian	0	10	0	10	0	10	0	10	0	10	10	0	0	10	
%	0	%	0	%	0	%	0	%	0		%	0	%	0	%	0	%	0	%	0	0	%	%	0	
%	%	%	%	%	%	%	%	%	%		%	%	%	%	%	%	%	%	%	%	%	%	%	%	
0	10	0	10	0	10	0	10	0	10	0	10	Moderate Carb	0	10	0	10	0	10	0	10	79	21	27	73	
%	0	%	0	%	0	%	0	%	0	%	0		%	0	%	0	%	0	%	0	%	%	%	%	
%	%	%	%	%	%	%	%	%	%	%	%		%	%	%	%	%	%	%	%	%	%	%	%	
0	10	0	10	1	84	0	10	0	10	0	10	Low GI/GL	0	10	0	10	0	10	0	10	90	10	0	10	
%	0	%	0	6	%	%	0	%	0	%	0		%	0	%	0	%	0	%	0	%	%	%	0	
%	%	%	%	%	%	%	%	%	%	%	%		%	%	%	%	%	%	%	%	%	%	%	%	
0	10	0	10	0	10	0	10	0	10	0	10	0	10	Nordic	0	10	0	10	0	10	0	10	10	0	
%	0	%	0	%	0	%	0	%	0	%	0	%	0		%	0	%	0	%	0	%	0	0	%	
%	%	%	%	%	%	%	%	%	%	%	%	%	%		%	%	%	%	%	%	%	%	%	%	
0	10	0	10	0	10	0	10	0	10	0	10	0	10	0	10	Tibetan	0	10	10	0	0	10	0	10	
%	0	%	0	%	0	%	0	%	0	%	0	%	0	%	0		%	0	0	%	%	%	%	0	
%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%		%	%	%	%	%	%	%	%	
0	10	0	10	0	10	0	10	0	10	0	10	0	10	0	10	0	10	Low-Sodium	0	10	10	0	0	10	
%	0	%	0	%	0	%	0	%	0	%	0	%	0	%	0	%	0		%	0	0	%	0	%	
%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%		%	%	%	%	%	%	
6	23	8	13	7	27	10	0	10	0	10	8	19	9	10	0	10	10	0	0	10	Low-Fat	74	26	0	10
2	%	7	%	3	%	0	%	0	%	0	1	%	0	%	%	0	0	%	0	%		%	%	%	%
%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%		%	%	%	%

5	43	0	10	0	10	0	10	0	10	0	10	2	73	0	10	10	0	0	10	10	0	73	27	Control
7	%	%	0	%	0	%	0	%	0	%	0	7	%	%	0	0	%	%	0	%	%	%	%	
%			%		%		%		%		%	%			%	%		%	%					
SBP																								

Table 2: Percentage contribution of each direct estimate derived from direct (blue) and indirect (red) comparisons (the colour corresponds to the percentage of contribution). The values above the diet classes correspond to the percentage contribution of direct and indirect comparisons between the row and columns for DBP (eg, the percentage contribution of direct comparisons for DBP between Mediterranean and Low-Carb diet is 35%, and 65% for the indirect comparisons). The values below the diet classes correspond to the percentage contribution of direct and indirect comparisons between the column and the rows for SBP (eg, the percentage contribution of direct comparisons for SBP between Mediterranean and Low-Carb diet is 34%, and 66% for the indirect comparisons).

DASH, Dietary Approaches to Stop Hypertension; DBP, diastolic blood pressure; SBP, systolic blood pressure.

DBP (mmHg)												
DASH	-1.56 (-3.55,0.44)	-1.74 (-3.92,0.43)	0.48 (-3.72,4.68)	-1.91 (-3.92,0.10)	-5.90 (-19.60,7.80)	-3.05 (-5.37,-0.74)	-1.60 (-4.05,0.85)	-3.22 (-6.14,-0.31)	-3.20 (-6.32,-0.08)	-1.91 (-3.70,-0.12)	-3.10 (-4.52,-1.68)	-4.37 (-5.78,-2.97)
-3.31 (-6.20,-0.42)	Mediterranean	-0.18 (-2.11,1.75)	2.04 (-2.16,6.24)	-0.35 (-2.35,1.65)	-4.34 (-18.04,9.36)	-1.50 (-3.85,0.86)	-0.04 (-2.47,2.39)	-1.66 (-4.80,1.47)	-1.64 (-4.75,1.47)	-0.35 (-2.47,1.76)	-1.54 (-2.95,0.13)	-2.81 (-4.63,1.00)
-2.86 (-5.96,0.25)	0.45 (-2.35,3.26)	Low-Carb	2.22 (-2.07,6.51)	-0.17 (-2.35,2.01)	-4.16 (-17.89,9.57)	-1.31 (-3.82,1.19)	0.14 (-2.34,2.63)	-1.48 (-4.73,1.77)	-1.46 (-4.69,1.77)	-0.17 (-2.45,2.11)	-1.56 (-3.01,0.30)	-2.63 (-4.64,-0.62)
1.35 (-6.31,9.01)	4.66 (-3.02,12.34)	4.21 (-3.55,11.97)	Palaeolithic	-2.39 (-6.59,1.82)	-6.38 (-20.57,7.81)	-3.53 (-7.92,0.85)	-2.08 (-6.51,2.35)	-3.70 (-8.55,1.11)	-3.68 (-8.51,1.11)	-2.39 (-6.66,1.88)	-3.58 (-7.53,0.38)	-4.85 (-8.97,-0.73)
-4.78 (-7.72,-1.85)	-1.47 (-4.44,1.50)	-1.92 (-5.11,1.26)	-6.13 (-13.83,1.56)	High-Protein	-3.99 (-17.69,9.71)	-1.15 (-3.51,1.22)	0.31 (-2.14,2.76)	-1.31 (-4.45,1.82)	-1.29 (-4.41,1.83)	-0.00 (-2.14,2.14)	-1.19 (-2.61,0.23)	-2.46 (-4.28,-0.64)
-0.65 (-17.15,15.84)	2.66 (-13.85,19.16)	2.20 (-14.34,18.75)	-2.00 (-19.97,15.96)	4.13 (-12.38,20.64)	Vegetarian	2.24 (-10.92,16.60)	4.30 (-9.47,18.07)	2.68 (-11.24,16.59)	2.70 (-11.21,16.61)	3.99 (-9.74,17.71)	2.80 (-10.83,16.43)	1.53 (-12.15,15.20)
-5.57 (-8.75,-2.39)	-2.26 (-5.54,1.03)	-2.71 (-6.19,0.76)	-6.92 (-14.74,0.90)	-0.79 (-4.11,2.53)	-4.91 (-21.48,11.66)	Moderate-Carb	1.46 (-1.29,4.20)	-0.17 (-3.48,3.14)	-0.14 (-3.50,3.22)	1.14 (-1.24,3.52)	-0.04 (-1.93,1.85)	-1.32 (-3.42,0.78)
-4.92 (-8.55,-1.30)	-1.61 (-5.25,2.03)	-2.07 (-5.74,1.61)	-6.27 (-14.26,1.71)	-0.14 (-3.83,3.55)	-4.27 (-20.92,12.38)	0.65 (-3.29,4.59)	Low GI/GL	-1.62 (-5.06,1.82)	-1.60 (-5.02,1.82)	-0.31 (-2.88,2.26)	-1.50 (-3.49,0.49)	-2.77 (-5.06,-0.48)
-4.74 (-9.11,-0.37)	-1.43 (-6.11,3.26)	-1.88 (-6.70,2.94)	-6.09 (-14.59,2.41)	0.04 (-4.66,4.75)	-4.08 (-20.99,12.82)	0.83 (-3.98,5.65)	0.19 (-4.98,5.35)	Nordic	0.02 (-3.91,3.95)	1.31 (-1.49,4.11)	0.12 (-2.67,2.91)	-1.15 (-3.70,1.41)
-3.95	-0.64	-1.10	-5.30	0.83	-3.30	1.61	0.97	0.78	Tibetan	1.29	0.10	-1.17

(-8.48,0.57)	(-5.19,3.91)	(-5.79,3.59)	(-13.73,3.12)	(-3.75,5.40)	(-20.17,13.57)	(-3.17,6.40)	(-4.07,6.01)	(-5.05,6.61)		(-1.92,4.49)	(-2.68,2.88)	(-4.17,1.83)
-4.42 (-7.12,-1.72)	-1.11 (-4.26,2.03)	-1.57 (-4.92,1.79)	-5.77 (-13.55,2.00)	0.36 (-2.86,3.58)	-3.77 (-20.32,12.78)	1.15 (-2.24,4.53)	0.50 (-3.38,4.38)	0.31 (-3.93,4.56)	-0.47 (-5.19,4.25)	Low-Sodium	-1.19 (-2.79,0.41)	-2.46 (-3.59, -1.32)
-5.05 (-7.08,-3.03)	-1.74 (-3.82,0.34)	-2.20 (-4.56,0.17)	-6.40 (-13.80,0.99)	-0.27 (-2.40,1.86)	-4.40 (-20.77,11.97)	0.51 (-2.03,3.06)	-0.13 (-3.14,2.88)	-0.32 (-4.51,3.88)	-1.10 (-5.15,2.95)	-0.63 (-3.06,1.80)	Low-Fat	-1.27 (-2.41,-0.14)
-7.38 (-9.43,-5.33)	-4.07 (-6.72,-1.41)	-4.52 (-7.41,-1.64)	-8.73 (-16.31,-1.15)	-2.60 (-5.29,0.10)	-6.72 (-23.18,9.73)	-1.81 (-4.69,1.07)	-2.46 (-5.89,0.97)	-2.64 (-6.50,1.22)	-3.42 (-7.80,0.95)	-2.96 (-4.74,-1.17)	-2.32 (-3.98,-0.67)	Control
SBP (mmHg)												

Table 3: League table: The values above the diet classes correspond to the difference in mean (95% CI) in DBP (mmHg) between the row and columns (e.g, the mean difference in average DBP between DASH and Low-Fat diets diet is -3.10 mmHg). The value below the diet classes correspond to the difference in mean in SBP (mmHg) between the column and the row (e.g, the mean difference in average SBP between DASH and Low-Fat diet is -5.05 mmHg).

DASH, Dietary Approaches to Stop Hypertension; DBP, diastolic blood pressure; SBP, systolic blood pressure.

DBP (mmHg)										
DASH	-1.91 (-4.00, 0.18)	-2.59 (-5.05, -0.13)	0.82 (-5.55, 7.19)	-1.83 (-4.55, 0.88)	-4.01 (-7.76, -0.27)	-2.80 (-5.98, 0.38)	-1.09 (-4.90, 2.72)	-3.42 (-5.83, -1.02)	-3.18 (-4.89, -1.47)	-3.79 (-5.72, -1.86)

-4.15 (-7.46, -0.85)	Mediterranean	-0.68 (-2.62, 1.26)	2.73 (-3.55, 9.00)	0.07 (-2.41, 2.56)	-2.10 (-5.79, 1.58)	-0.89 (-3.84, 2.06)	0.82 (-3.06, 4.69)	-1.52 (-4.02, 0.99)	-1.27 (-2.59, 0.05)	-1.88 (-3.93, 0.16)
-4.24 (-8.09, -0.39)	-0.08 (-3.18, 3.01)	Low-Carb	3.41 (-2.99, 9.81)	0.76 (-2.04, 3.55)	-1.42 (-5.32, 2.47)	-0.21 (-3.24, 2.83)	1.50 (-2.58, 5.57)	-0.83 (-3.64, 1.97)	-0.59 (-2.42, 1.24)	-1.20 (-3.61, 1.21)
3.17 (-11.04, 17.38)	7.32 (-6.81, 21.46)	7.41 (-6.86, 21.67)	Palaeolithic	-2.65 (-9.15, 3.84)	-4.83 (-11.86, 2.20)	-3.62 (-10.30, 3.07)	-1.91 (-9.05, 5.22)	-4.24 (-10.74, 2.25)	-4.06 (-10.13, 2.13)	-4.61 (-10.94, 1.72)
-7.12 (-11.76, -2.47)	-2.96 (-7.36, 1.43)	-2.88 (-7.69, 1.93)	-10.29 (-24.78, 4.20)	High-Protein	-2.18 (-6.23, 1.87)	-0.96 (-4.39, 2.46)	0.74 (-3.48, 4.97)	-1.59 (-4.61, 1.43)	-1.35 (-5.49, 0.80)	-1.96 (-4.61, 0.70)
-6.96 (-12.50, -1.42)	-2.81 (-8.32, 2.71)	-2.73 (-8.58, 3.13)	-10.13 (-25.00, 4.74)	0.16 (-6.23, 6.54)	Moderate-Carb	1.21 (-3.13, 5.56)	2.92 (-1.87, 7.71)	0.59 (-3.18, 4.36)	0.83 (-2.61, 4.27)	0.22 (-3.27, 3.71)
-2.96 (-8.16, 2.24)	1.19 (-3.72, 6.10)	1.27 (-3.69, 6.24)	-6.13 (-20.81, 8.54)	4.15 (-1.76, 10.06)	4.00 (-2.79, 10.79)	Low GI/GL	1.71 (-2.80, 6.22)	-0.63 (-4.03, 2.78)	-0.38 (-3.04, 2.27)	-0.99 (-4.08, 2.09)
-3.18 (-9.51, 3.15)	0.97 (-5.47, 7.41)	1.05 (-5.67, 7.78)	-6.35 (-21.59, 8.89)	3.94 (-3.26, 11.13)	3.78 (-3.72, 11.28)	-0.22 (-7.78, 7.34)	Nordic	-2.33 (-5.92, 1.25)	-2.09 (-5.73, 1.55)	-2.70 (-5.99, 0.59)
-6.15 (-10.01, -2.29)	-2.00 (-6.04, 2.04)	-1.92 (-6.41, 2.57)	-9.32 (-23.71, 5.06)	0.96 (-4.20, 6.13)	0.81 (-4.77, 6.39)	-3.19 (-8.84, 2.46)	-2.97 (-9.04, 3.09)	Low sodium	0.24 (-1.88, 2.37)	-0.37 (-1.80, 1.07)
-5.83 (-8.44, -3.22)	-1.68 (-3.81, 0.46)	-1.59 (-4.49, 1.30)	-9.00 (-22.97, 4.97)	1.29 (-2.56, 5.14)	1.13 (-3.97, 6.23)	-2.87 (-7.35, 1.61)	-2.65 (-8.73, 3.44)	0.32 (-3.12, 3.77)	Low-Fat	-0.61 (-2.18, 0.96)
-6.88 (-9.89, -3.87)	-2.73 (-5.97, 0.51)	-2.65 (-6.43, 1.14)	-10.05 (-24.24, 4.13)	0.24 (-4.33, 4.80)	0.08 (-4.95, 5.11)	-3.92 (-9.03, 1.20)	-3.70 (-9.26, 1.86)	-0.73 (-3.14, 1.68)	-1.05 (-3.51, 1.41)	Control
SBP (mmHg)										

Table 4: League table (Low risk of bias sensitivity analysis): The values above the diet classes correspond to the difference in mean (95% CI) in DBP (mmHg) between the row and columns (eg, the mean difference in average DBP between DASH and Low-Fat diets is -3.18 mmHg).

The value below the diet classes correspond to the difference in mean in SBP (mmHg) between

the column and the row (eg, the mean difference in average SBP between DASH and Low-Fat diets is -5.83 mmHg);

DASH, Dietary Approaches to Stop Hypertension; DBP, diastolic blood pressure; SBP, systolic blood pressure.




Figure 1: Network diagram for diastolic blood pressure: The size of the nodes is proportional to the total number of participants allocated to each dietary approach and the thickness of the lines proportional to the number of studies evaluating each direct comparison.