



## Critical Reviews in Food Science and Nutrition

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/bfsn20>

### Immunity: Plants as Effective Mediators

M. Tauseef Sultan<sup>a</sup>, Masood Sadiq Butt<sup>xs</sup>, Mir M. Nasir Qayyum<sup>c</sup> & Hafiz Ansar Rasul Suleria<sup>b</sup>

<sup>a</sup> Department of Food Sciences, Faculty of Agricultural Sciences and Technology, Bahauddin Zakariya University, Multan, Pakistan

<sup>b</sup> National Institute of Food Science & Technology, University of Agriculture, Faisalabad, 38000, Pakistan

<sup>c</sup> Department of Food Technology, Karakoram International University, Gilgit-Baltistan, Pakistan

Accepted author version posted online: 14 May 2013. Published online: 24 Feb 2014.

To cite this article: M. Tauseef Sultan, Masood Sadiq Butt<sup>xs</sup>, Mir M. Nasir Qayyum & Hafiz Ansar Rasul Suleria (2014) Immunity: Plants as Effective Mediators, Critical Reviews in Food Science and Nutrition, 54:10, 1298-1308, DOI: [10.1080/10408398.2011.633249](https://doi.org/10.1080/10408398.2011.633249)

To link to this article: <http://dx.doi.org/10.1080/10408398.2011.633249>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

# Immunity: Plants as Effective Mediators

M. TAUSEEF SULTAN,<sup>1</sup> MASOOD SADIQ BUTT,<sup>2</sup> MIR M. NASIR QAYYUM,<sup>3</sup>  
and HAFIZ ANSAR RASUL SULERIA<sup>2</sup>

<sup>1</sup>Department of Food Sciences, Faculty of Agricultural Sciences and Technology, Bahauddin Zakariya University, Multan, Pakistan

<sup>2</sup>National Institute of Food Science & Technology, University of Agriculture, Faisalabad 38000, Pakistan

<sup>3</sup>Department of Food Technology, Karakoram International University, Gilgit-Baltistan, Pakistan

*In the domain of nutrition, exploring the diet–health linkages is major area of research. The outcomes of such interventions led to widespread acceptance of functional and nutraceutical foods; however, augmenting immunity is a major concern of dietary regimens. Indeed, the immune system is incredible arrangement of specific organs and cells that enabled humans to carry out defense against undesired responses. Its proper functionality is essential to maintain the body homeostasis. Array of plants and their components hold immunomodulating properties. Their possible inclusion in diets could explore new therapeutic avenues to enhanced immunity against diseases. The review intended to highlight the importance of garlic (*Allium sativum*), green tea (*Camellia sinensis*), ginger (*Zingiber officinale*), purple coneflower (*Echinacea*), black cumin (*Nigella sativa*), licorice (*Glycyrrhiza glabra*), *Astragalus* and *St. John's wort* (*Hypericum perforatum*) as natural immune boosters. These plants are bestowed with functional ingredients that may provide protection against various menaces. Modes of their actions include boosting and functioning of immune system, activation and suppression of immune specialized cells, interfering in several pathways that eventually led to improvement in immune responses and defense system. In addition, some of these plants carry free radical scavenging and anti-inflammatory activities that are helpful against cancer insurgence. Nevertheless, interaction between drugs and herbs/botanicals should be well investigated before recommended for their safe use, and such information must be disseminated to the allied stakeholders.*

**Keywords** Immunomodulation, medicinal plants, nutraceuticals, phytochemicals

## INTRODUCTION

The tenet “Let food be thy medicine and medicine be thy food,” exposed by Hippocrates nearly 2500 years ago, gained renewed interests that resulted in coinage of terms like pharma, nutraceuticals, and functional foods. These terms are not only gaining popularity among researchers, but consumers are also interested toward natural diet based regimens. These natural products act as immune boosters thus improving the quality of life (Ares et al., 2009; Sultan et al., 2009). Augmenting immunity is a major concern of diet-based therapies to cure various disorders, and researchers are thriving for immune boosters (Bourgeon et al., 2007; Butt et al., 2009).

The immune system is an incredibly intricate network of specialized cells that prevents infections and diseases by engulfing, modulating, and moderating malignant and foreign cells. Hu-

man's immune system is composed of organs like spleen and thymus, whereas additionally lymph nodes and bone marrow also contribute in immune system by producing and storing specific immune cells (Chaouat et al., 2007). Immune cells are of two major types, i.e., B cells and T cells. B cells are responsible for producing antibodies (immunoglobulins) that are proteins designed to recognize and mark a specific antigens, whereas T cells are charged moieties with destroying antigens tagged with an antibody (Chanana et al., 2007; Zhang et al., 2007).

T cells play critical role in controlling the adaptive immune functions, and their responses could be used to develop protective vaccines and may induce tolerance to antigens causing inappropriate immune responses, e.g., autoimmune diseases (Cooper and Alder, 2006; Li et al., 2007). In addition, phagocytes such as granulocytes, macrophages and natural killer cells (NK cells) release pyrogens and interferons that act as immunoregulatory moieties (Currier and Miller, 2002; Fauci et al., 2005). Chemical mediators like cytokines (monokines and lymphokines) are also effective in regulating immune responses.

Address correspondence to Dr. M. Tauseef Sultan, Department of Food Sciences, Faculty of Agricultural Sciences and Technology, Bahauddin Zakariya University, Multan, Pakistan. E-mail: tauseefsultan@bzu.edu.pk

Some other mediators like tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ), interleukins (IL), chemokines, and interferon bodies (IFN) also contribute in proper functioning of immune system (Timár et al., 2007).

Interest in medicinal plants has burgeoned owing to increased efficiency of new phytochemicals and the growing awareness among consumers regarding natural product consumption. A larger number of plants and their isolated constituents have shown health benefits, including antioxidant, anti-inflammatory, anticancer, antimicrobial, and immunomodulatory effects (Butt et al., 2009; Sultan et al., 2009; Butt and Sultan, 2011). Plants produce thousand of bioactive compounds, and their complexity must be well investigated (Schwager et al., 2008). Several plants and their metabolites have health-promoting properties, i.e., vitamins, antioxidants, dietary fibers, phytosterols, flavonoids, omega-3-fatty acids, etc. (Butt et al., 2008; Wang et al., 2009). Human diet containing these phytochemicals possess antimutagenic and immunomodulatory potential and have shown anti-inflammatory, antistress, and anticancer properties. It is important for clinicians to document use of herbal medicines as part of the patient's drug profile (Ramaa et al., 2006; Roller et al., 2007).

Summarizing nutraceutical potential of all herbs is out of scope for this paper, thus emphasis paid on some of promising plants like garlic (*Allium sativum*), green tea (*Camellia sinensis*), ginger (*Zingiber officinale*), purple coneflower (*Echinacea*), black cumin (*Nigella sativa*), licorice (*Glycyrrhiza glabra*), *As-tragalus* and St. John's wort (*Hypericum perforatum*) as natural immune boosters.

### Garlic

Garlic (*Allium sativum* L.) was originated from Central Asia over 6000 years ago and holds potential to be employed as medicinal agent (Butt et al., 2009). The health benefits of garlic appear to be true today as diet-based therapy and its use as a dietary supplement is recommended in many countries (Raman et al., 2008). Most of its biological effects are attributed to organosulfur compounds including, diallyl sulfide (DAS), diallyl disulfide (DADS),  $\delta$ -glutamyl-S-allyl-L-cysteines, S-allylmercaptocysteine (SAMC), S-allyl-L-cysteine sulfoxides, and nonstarch polysaccharides (Clement and Venkatesh, 2010; Liang et al., 2011). Various garlic preparations such as aged garlic extract (AGE) and aqueous extract, garlic oil are being sold in the market with distinct health claims (Chandrashekar et al., 2011).

Immunomodulatory properties of garlic and its bioactive profile include immune stimulation, reduce platelet aggregation, and chemoprevention of cancer (Gamboa-Léon et al., 2007; Wojcikowski et al., 2007). Garlic and AGE can scavenge free radicals and act as successful antioxidant. In addition, they also have the potency to enhance and regulate the activities of the antioxidant enzymes, such as glutathione peroxidase, glutathione S-transferase (GST), catalase, acid soluble sulfhydryl (-SH) cytochrome b5 and cytochrome P450, and superoxide dismutase

(Borek, 2001; Kris-Etherton et al., 2002; Zhou et al., 2007; Nagaraj et al., 2012)

The role of garlic in immunonutrition is multifarious both as immunostimulant and immunosuppressant. The consumption of garlic results in improvement in hematological attributes, e.g., total white blood cell (WBC) count as well as homeostasis characteristics (Wilson and Demmig-Adams, 2007). Consumption of garlic may increase the production and release of nitric oxide (NO) that is further responsible for enhanced release of IFN- $\alpha$  in humans, beneficial against viral, or proliferative diseases (Bhattacharyya et al., 2007). The overall improvement in immune system could be attributed to the activation of specific responses and detoxification mechanisms (Chandrashekar and Venkatesh, 2009; Zamani et al., 2009). Proliferation of lymphocytes; macrophage phagocytosis; infiltration of macrophages and lymphocytes; and stimulating release of IL-2, TNF- $\alpha$ , and IFN- $\gamma$  are some possible routes of garlic for its immunomodulatory properties. In addition, it can enhance NK cell and lymphokine-activated killer cell activities that represent effective stimulation of the immune responses (Iciek et al., 2009; Kim and Kwon, 2009).

Inhibiting growth of cancerous cells is perhaps the most remarkable beneficial action of garlic. It has been reported from the last decade that garlic and garlic preparations may possess anticancer activities against skin, colon, prostate, mammary carcinoma, lungs, gastric cancer, etc. (Wu et al., 2009; Altonsy and Andrews, 2011; Wang et al., 2012). The development of successful strategy employing dietary chemopreventive agent reside in their mode of actions. Certainly, it does not appear that a single mechanism could account for the observed protection based on the variety of carcinogens (Galeone et al., 2006; Chen et al., 2012). Organosulfur compounds of garlic inhibit carcinogen activation/formation, boost phase-II detoxifying processes, cause cell cycle arrest mostly in G2/M phase, stimulate the mitochondrial apoptotic pathway, increase acetylation of histones, and suppression of tumor proliferation (Iciek et al., 2009; Bat-Chen et al., 2010; Altonsy and Andrews, 2011; Wu et al., 2011). Earlier, Galeone et al. (2006) provided evidence using multivariate odd ratios that garlic is inversely associated with the risk of several common cancers. Some of its preparations like aged garlic extract and fresh garlic juice hold potential to inhibit cell proliferation and induce apoptosis (Sengupta et al., 2004). Recently, it has been suggested that the DAS, diallyl trisulfide (DADS) and DATS can initiate a cascade of molecular events characteristic of apoptosis of cancerous cells (Chandra-Kuntal and Singh, 2010; Altonsy and Andrews, 2011). The mechanism through which garlic and its functional ingredients imparts anticancer effects needs further elaborations.

Studies suggest that garlic may be useful in preventing the suppression of immune response, associated in increased risk of malignancy. In the nutshell, it can be assumed that garlic is one of the most potential candidates as immune booster, improves the antioxidants status of the body, and protects body from free radicals, inflammation, and cancer insurgence.

## Green Tea

Green tea (*Camellia sinensis*) is one of the most popular beverages consumed worldwide, and its consumption is increasing day by day. The changing trends of consumers are mainly attributed to its perceived antioxidant, anti-inflammatory, and anticarcinogenic properties (Almajano et al., 2011). Catechin (epicatechin, epigallocatechin, epicatechin gallate, and epigallocatechin gallate) are its active ingredients (Baliga and Katiyar, 2006). Among these, epigallocatechin-3-gallate (EGCG) is identified as the principal antioxidant moiety contributing approximately 30% of the total antioxidant capacity (Colic and Pavelic, 2000; Stewart et al., 2005).

Recently, Butt and Sultan (2009) provided molecular and cellular insights of green tea for their antioxidant and anti-inflammatory potential. EGCG has marked effect in modulating production of immunoregulatory cytokines in stimulated dendritic cells and hence acts as suppressor of T cell activation (Rogers et al., 2005). It can also prevent angiogenesis as observed in angiogenic assay in vivo and inhibits the growth of the highly angiogenic Kaposi's sarcoma tumor cells in rodent experimental modeling (Noonan et al., 2007; Kawai et al., 2011). Green tea and EGCG effectively mitigate cellular damage by lowering the inflammatory reaction and reduce lipid peroxidation and formation of NO radicals (Norwood et al., 2006; Tipoe et al., 2007; Bais et al., 2010).

The anticancer perspectives of green tea catechins have been highlighted in the last few decades. It can significantly reduce the risk of cancers as revealed in experimental animal studies, human cell lines, and human clinical studies (Thakur et al., 2012). Overall, it can be used as adjunct in the treatment of bladder, breast, and colon cancer (Sartippour et al., 2002; Singh et al., 2011; Yang et al., 2011). Green tea is also effective in leukemia, liver, biliary tract, and lung cancer (Fujimoto et al., 2002; Choudhury et al., 2011; Wang et al., 2011). Current applications of green tea polyphenols can provide protection against ultra violet-induced skin cancers (Mantena et al., 2005). Some other scientific data reported its beneficial effects in esophageal and prostate cancer too (Gupta et al., 2003). The studies suggested that multiple mechanisms are involved in its chemoprevention properties including induction of apoptosis, cell cycle arrest downregulation of telomerase, inhibition of vascular endothelial growth factor, and suppression of aromatase activity (Sartippour et al., 2002; Butt and Sultan, 2009; Yang et al., 2011). Green tea assumed to induce apoptosis and promote cell growth by altering expression of cell cycle regulatory proteins, altering Bax/Bcl2 function, activating killer caspases, and suppressing nuclear factor kappa B function (Aktas et al., 2004; Khan et al., 2006). Green tea catechins especially EGCG has been shown to block each stage of carcinogenesis by modulating the signal transduction pathways during proliferation, inflammation, apoptosis, and metastasis (Kang et al., 2007; Butt and Sultan, 2009; Al-Hazzani and Alshatwi, 2011; Singh et al., 2011). Some studies do suggest that green consumption effectively improve the overall antioxidant capacities of the body in

human subjects (Ellinger et al., 2011). However, some scientific intervention, cohort studies, and meta-analysis suggested that green tea intake is not completely useful for all populations, e.g., smokers and exsmokers are at risk of lung cancer if they keep consuming green tea. The synergistic and antagonistic effect of green tea catechins with other drugs/natural products needs further attention as some studies suggested that EGCG can help curcumin against cancer insurgence (Manikandan et al., 2011; Xu et al., 2011). The Japanese Public Health Center-based prospective study (JPHC) highlighted the association between green tea consumption and decreased risk of cancer insurgence (Iwasaki and Tsugane, 2011).

Green tea and EGCG are now of clinical importance for the treatment of lifestyle-related disorders and cancer insurgence (Sueoka et al., 2001). Watson et al., 2005 also recommended that green tea and catechins especially EGCG could be useful for enteric immune disorders and T cell driven immunopathologies. The instant discussion is a message for adopting "designer approach" to get maximum health benefit from green tea catechins.

## Ginger

Recent research has rejuvenated the centuries old traditional herbs to cure various ailments by using the modern tools like diet-based therapy or regimen. Ginger (*Zingiber officinale*) is one of the classic examples used for not only culinary preparations but also holds unique therapeutic significance (Baliga et al., 2011; Butt and Sultan, 2011). The bioactive molecules present in ginger include  $\alpha$ -zingiberene,  $\alpha$ -farnesene,  $\beta$ -bisabolene,  $\alpha$ -curcumene, [6]-gingerol and [6]-shogaol, paradol, zingerones, and allied derivatives (Zhan et al., 2008). Owing to the rich phytochemistry, it has been used in Chinese, Ayurvedic medicines (Hung et al., 2009; Jeong et al., 2009).

Ginger and its bioactive molecules are effective in controlling the extent of colorectal, gastric, ovarian, liver, and skin cancers. The molecular targets involved in chemoprevention include inhibition of NF- $\kappa$ B activation, suppresses cIAP1 expression and cell proliferation, arrests cell cycle in G2/M phases, upregulates cytochrome c, Apaf-1, activates PI3K/Akt/I kappaB kinases IKK. It also induces apoptosis, chromatin condensation, and increases caspase-3/7 activation and the number of tumor-infiltrating lymphocytes (Kim et al., 2009; Oyagbemi et al., 2010; Ju et al., 2012).

Ginger is consumed in many cultures as immune boosters (Kannappan et al., 2011), and its anti-inflammatory potential has been highlighted in number of scientific investigations. The mechanisms for its anti-inflammatory properties include inhibition of arachidonic acid-induced platelet aggregation and formation of thromboxane B, upregulation of histone H3 acetylation, and suppressed histone deacetylase (HDAC)1 expression (Shim et al., 2011), inhibition of IL-1, TNF- $\alpha$  and IL-8, downregulation of inflammatory inducible NO synthase (iNOS) and cyclooxygenase 2 (COX-2) gene expression through

inactivation of Nuclear Factor Kappa B (NF- $\kappa$ B) (Rani et al., 2011). The other molecular mechanisms, include extracellular signal-regulated kinases 1 and 2 (ERK1/2), p38 mitogen-activated protein kinase (MAPK), c-Jun N-terminal kinase (JNK), and the inactivation of Nuclear factor- $\kappa$ B (NF- $\kappa$ B) (Zhou et al., 2006; Jung et al., 2009). Likewise, 6-gingerol, 10-gingerol, 8-shogaol, and 10-shogaol can decrease iNOS and TNF- $\alpha$  expression through suppression of I- $\kappa$ B $\alpha$  phosphorylation, Nuclear factor- $\kappa$ B (NF- $\kappa$ B), gene activation of proinflammatory enzymes, COX-2, nuclear activation, and protein kinase C (PKC)- $\alpha$  translocation (Lee et al., 2009; Kim et al., 2010; Li et al., 2011; van Breemen et al., 2012).

Immune system also helps the body by protecting it from hazardous effects of chemicals. The ginger and its bioactive molecules are effective in providing protection against hepatotoxicity induced through bromobenzene, ethanol, and acetaminophen via modulation of enzymes (Mallikarjuna et al., 2008; Atta et al., 2010). 8-gingerol (50, 100 mg/kg) suppressed humoral and cellular immune responses through direct inhibition of sensitized T and B lymphocytes (Ueda et al., 2010; Lu et al., 2011; Nievergelt et al., 2011; Shim et al., 2011). Intraperitoneal injections of ginger extract decreased the eosinophils to the lungs along with diminished levels of IL-4, IL-5, and eotaxin levels (Ahui et al., 2008). Further evidences can also be presented in this regard as [6]-gingerol (25–50 mg/kg) inhibited acetic acid-induced writhing response and formalin-induced licking time. However, higher doses of [6]-gingerol (50–100 mg/kg) are required to inhibit paw edema induced by carrageenin (Young et al., 2005).

The anti-inflammatory potential of ginger is helpful for the management of disorders like respiratory infections (Podlogar and Verspohl, 2012), arthritis (Funk et al., 2009), allergic diseases (Chen et al., 2009), and gout (Sabina et al., 2010). In the nutshell, it can be observed that ginger and its components hold anti-inflammatory activities. They are effective in reducing the extent of chemical toxicity and are of significance importance in treating the inflammatory disorders. Cohort studies and controlled trails should be conducted to warrant its pharmacological applications.

### Purple Coneflower (Echinacea)

*Echinacea* is one of the most important plants having various herbal preparations that purport to improve immune functioning (Brush et al., 2006). The diversity of phytochemicals in *Echinacea* sp. includes betain, sesquiterpenes, caryophyllene, polyacetylene, rosmarinic acid, glycosides, echinacoside, chicoric acid, alkyl amides, and arabinogalactan-proteins (Hwang et al., 2004; Huntley et al., 2005).

Alkyl amides present in *Echinacea* sp. have reported immunomodulatory actions as they suppressed the ability of activated Jurkat T cells (key mediators of antiviral immunity) to produce IL-2 independently (Sasagawa et al., 2006; Woelkart et al., 2006). Arabinogalactan-protein and various other bioactive constituents of this plant were clearly identified as stimula-

tors of both the classical and alternative pathway of complement activation and modulation involving inhibition or stimulation of immunity. These modules would also be interesting targets for drug development to ease the mucosal immune suppression and lessen the duration of upper respiratory tract infections (Alban et al., 2002; Bonifati and Kishore, 2007; Hall et al., 2007).

*Echinacea* preparations activate cellular immunity and stimulate phagocytosis of neutrophils both in vitro and in vivo. Stimulatory effect and immune-modulating action by *Echinacea* juice increase cellular immune parameters, e.g., the percentage of phagocytosing cells, NBT-positive granulocytes and lysozyme level reported by Truchlinski et al. (2006). It effectively stimulates immunocompetence, and the plant extract behaves as hematinic agent, i.e., one can improve the quality of blood by increasing hemoglobin levels and the number of erythrocytes. Consequently, these effects result in improvements of oxygen transport that are further useful in improving the exercise physiology and performance (O'Neill et al., 2002).

*Echinacea* stimulates the immune system against bacterial and viral infections, and its extracts are widely used in prophylaxis and various other therapies (Bany et al., 2003; Gan et al., 2003). In contrary, South and Exon (2001) were on opinion that it holds immunosuppressive activity instead of immunostimulatory activity. However, it is able to stimulate innate immune responses including macrophages and natural killer cells regulated responses. Research carried out by Cundell et al. (2003) suggested that *Echinacea* affects both at mononuclear cell levels and circulating IL-2 levels. The recent advances has advocated the idea that combinatorial therapy, involving specific tumor cell immunization, followed by daily phytotherapy by *E. purpurea*, could lead to life span prolongation greater than immunization alone (Currier and Miller, 2002).

In brief, mode of action of *Echinacea* in enhancing immunity moves around activating and modulating cell-mediated immune responses and, indeed, as hematinic agents. In this context, the use of *E. purpurea* in diet-based therapies has the potential to enhance both humoral immune responses as well as innate immune responses.

### Black Cumin (Nigella sativa)

Seeds of black cumin (*Nigella sativa*) have been used in herbal medicine for the treatment and prevention of a number of diseases, supplement to maintain good health and nutrition (Salem and Hossain, 2000; Sultan et al., 2009; Jafri et al., 2010; Abusnina et al., 2011). In the literature, numbers of evidences unveiled its anti-inflammatory and immunomodulatory effects (Tekeoglu et al., 2006; Salem et al., 2011). *N. sativa* displays its regulatory effects via inflammatory cells (Ozugurlu et al., 2005) by directly influences immune system and improves helper T cell (T4) to suppressor T cell (T8) ratio. It also improves natural killer cell activity. Moreover, due to the presence of antioxidant components, it protects central nervous compartments like brain and medulla spinalis tissues against autoimmune encephalomyelitis (Butt and Sultan, 2010).

The oral administration of black cumin essential oil, a phytochemicals rich fraction, holds significant analgesic effect as reflected from acetic acid-induced writhing, formalin, and light tail flick tests. Moreover, it can inhibit carrageenan-induced paw edema and croton oil-induced edema (Hajhashemi et al., 2004; Ghannadi et al., 2005). The therapeutic potential of black cumin essential oil for curing rheumatism is also validated (Mahmood et al., 2003). Later, Tekeoglu et al. (2006) confirmed that thymoquinone (active ingredient of essential oil) suppressed adjuvant-induced arthritis in rodent experimental modeling. Thymoquinone holds ability to inhibit the leukotrienes (LTs) formation by modulating 5-lipoxygenase and LTC<sub>4</sub> synthase activity (Dijsselbloem et al., 2004; Mansour and Tornhamre, 2004; El Gazzar et al., 2006).

*N. sativa* especially its bioactive component thymoquinone holds anticancer perspectives (Banerjee et al., 2010; Sayed-Ahmed et al., 2010; Arafa et al., 2011). It was reported in one case study that they inhibited carcinogenesis in the post-initiation stage, and inhibition was associated with suppression of cell proliferation in colonic mucosa (Salim and Fukushima, 2003; Gali-Muhtasib et al., 2004). These fractions are effective in reducing the extent of oxidative stress and mammary carcinoma (Norwood et al., 2006; Velho-Pereira et al., 2011; Woo et al., 2011). Likewise, *N. sativa* and thymoquinone reduce the carcinogenic effects of 7, 12 di-methylbenz(a)anthracene and benzo(a)pyrene (B(a)P), respectively (Rooney and Ryan, 2005; Badary et al., 2007).

Another active ingredient of black cumin is  $\alpha$ -hederin that possesses antitumor activity. It is effective against cultured SiHa cells, B16 melanoma cells, Lewis lung carcinoma, hepatocellular carcinoma (HepG2), and murine leukemia P388 cells (Swamy and Huat, 2003; Ng et al., 2011). The mode of action of this active ingredient includes production of reactive oxygen species (ROS) that inhibit DNA, RNA, and protein synthesis (Kumara and Huat, 2001).

Health benefits associated with *N. sativa* supplementation might be due to its antioxidant potential that improves natural defense system of the body. Further studies are urgently required to explore bystander effects of *N. sativa*, and its active ingredients and their possible use as therapeutic agents. The nutritional quality of black cumin can find its prospects of adding up in diet-based strategies to improve the human health (Sultan et al., 2009; Butt and Sultan, 2010).

### *Astragalus*

In Chinese, Astragalus is known as Huang-qi and is written in 2000-year-old Shen Nong Ben Cao Jing and is still considered one of the superior tonics in traditional Chinese medicine (Yip and Kwan, 2006; Xu et al., 2007; Walsh et al., 2011). Astragalus membranaceus extract (AME) has therapeutic potential in regulating human immune functions and antitumor activities and may be applied in clinical practice for immunomodulation and cancer therapy (Wang et al., 2002). The immune boosting

potential of Astragalus includes stimulation of immune cells against stimulus-response action (Brush et al., 2006). Furthermore, AME could promote the proliferation of human peripheral blood mononuclear cell (PBMC), elevates the tumor cell-killing activity of cytotoxic T-lymphocyte, strengthens the tumor cell phagocytosis and cytokines (TNF- $\alpha$  and IL-6) production from peripheral blood adherent monocytes and IgG production from same types of cells (Wang et al., 2002).

Wu et al. (2006) are of the view that auxiliary use of Astragalus injection (AI) is beneficial to the restoration of cellular immunity after serious abdominal trauma (SAT). AI also improves the immune function of congestive heart failure (CHF) patients. Mechanism behind its action involves increased CD4 level and CD4/CD8 ratio while it lowers the levels of IL-2R, IgG, and IgA (Liu et al., 2003). AI supplemented with chemotherapy could inhibit the development of cancer insurgence. It also decreases the adverse consequences of chemotherapy through elevation of immune function that, in turn, improves the quality of life in cancerous patients (Duan and Wang, 2002; Guo et al., 2004).

The aforementioned effects are attributed to the phytochemical profile of Astragalus like Astragalus polysaccharide (APS) and lectins (Yan et al., 2005). APS significantly induces NO production and iNOS transcription through the activation of nuclear factor-kappaB/Rel (Lee and Jeon, 2005). APS promotes lymphocyte proliferation and both humoral and cellular immune responses that would be expected to be new-type of immunopotentiator drugs (Kong et al., 2004). APS and astragalosides (AS) have strong promoting effects on macrophages that result in enhanced secretion of IL-1  $\beta$ , IL-6, and TNF- $\alpha$  (Xu et al., 2007). Recently, Liu et al. (2011) suggested that APS might induce the differentiation of splenic dendritic cells (DCs) to CD11c(high)CD45RB(low) DCs followed by shifting of Th2 to Th1 with enhancement of T lymphocyte immune function. The differentiation of DCs was not associated with the inhibition of IL-10 production in CD11c(low)CD45RB(high) DCs. In addition, APS can induce increased TLR4 expression that can result in upregulation of innate immunity during mucosal bacterial infection (Yin et al., 2010). The same component effectively improves the quality of life in subjects with advanced nonsmall cell lung cancer (Guo et al., 2011).

In herbal medicines, it is common to use herbs in combination in the form of decoction. Such herbal remedies containing Astragalus are also important in improving phagocytic activity of peritoneal macrophage (Ning et al., 2005). The efforts should be directed toward exploration of herbal combinations that could be useful in enhancing the functionality of immune systems. These improvements will not only be helpful in chemotherapies but also immune related disorders.

### *Licorice (Glycyrrhiza glabra)*

Herbal medication in the United States is a popular form of therapy and Licorice, the root extract of *Glycyrrhiza glabra*

is used as a medicine for various diseases (Abebe, 2003). It holds anti-inflammatory as well as antiallergic activities that are mainly attributed to its bioactive constituents, like glycyrrhizin, aglycone, beta-glycyrrhetic acid (Kroes et al., 1997). In this context, Yang et al. (1990) characterized 10 flavonoids that are licochalcone A, isoschaftoside, schaftoside, liquiritin, isoviolanthin, violanthin, ononin, isoliquiritin, 4',7-dihydroxyflavone, and licoflavone A. Medicinal importance of flavonoids present in *Glycyrrhiza* is not known yet, but some of them are reported to play vital roles in proper functionality of immune system.

Glycyrrhizin has anti-inflammatory and anti-allergic properties (Shibata, 2000), and of its mode of action is actually revolve around production of IL-10 and IL-12 in large amount. Glycyrrhizin holds dose-dependent priming effect on lipopolysaccharide (LPS)-induced IL-12 p40 and IL-12 p70 (heterodimer of p40 and p35) protein production by peritoneal macrophages (PM). IL-12, being monocyte/macrophage-derived cytokine it plays a prominent role in the development of T helper type 1 (Th1) cell-mediated immune responses (Dai et al., 2001).

Likewise, the reduction in cellular immunocompetence in the gamma-irradiated mice can be recovered with the treatment of glycyrrhizae and glycyrrhizic acid (active fractions of licorice). These fractions are found to be effective in enhancing the leukocyte count and the blastogenic responses of splenocytes to mitogens (Dorhoi et al., 2006). Glycyrrhizin is well known to have various immunomodulating and biological response-modifier activities. In one case study, SNMC (stronger neominophagen C), active component is glycyrrhizin improved the liver function (Abe et al., 2003).

Some scientific evidences support its use as immunostimulating agent that is involved in proper metabolisms and functioning of various important mediators in adaptive or acquired immunity. The claims mentioned in the earlier section suggested that *Glycyrrhiza glabra* could act as immunostimulating agent (Brush et al., 2006). However, the mechanisms underlying the anti-inflammatory activities of glycyrrhizin are still poorly understood.

### ***St. John's Wort (Hypericum perforatum)***

St. John's wort (*Hypericum perforatum*) is effective in wounds and injuries healing and holds anti-inflammatory and antioxidative activities and indeed acts as antidepressant agent. The characteristic metabolites of St John's wort are the photodynamic active plant pigment hypericin and the phloroglucin-derivative hyperforin. In addition, it contains many polyphenolic compounds, e.g., flavonoids and phenolic acids (Genovese et al., 2006).

Owing to the presence of these functional ingredients, it has high antioxidant potential that can reduced the peroxidative and chlorinating activity of human leukocyte MPO in concentration-dependent manner. The inhibition of MPO activity is further responsible for its anti-inflammatory potential (Pabuccuoglu et al., 2003). Moreover, *Hypericum* extract and its bioactive compo-

nent, hyperforin, inhibit proliferation of T lymphocytes that is helpful in mediating inflammatory skin disorders (Wan et al., 2003). In another study, compounds like hyperguinone and hyperforin have strong potential in reducing oxygen production by polymorphonuclear cells (PMNs) after stimulation with N-formyl-methionyl-leucyl-phenylalanine (Heilmann et al., 2003).

Its potential as antidepressant agent has been highlighted in several studies. Proper functionality of immune system results in faster recovery from state of depression as immunological changes occur in depressed persons including increase in total number of white blood cells and in the numbers and percentages of neutrophils and lymphocytes. Enumeration of lymphocyte subsets also revealed that depression was associated with decreases in the number and percentage of lymphocytes and NK cells (Zorrilla et al., 2001). Immunomodulation is of clinical and practical consideration not only as targets for the treatment of depression but also to reduce the risk of development and progression of infectious diseases. *Hypericum* extract possesses antidepressant activity and is popular herbal supplement used to treat mild to moderate depression. It possesses serotonergic properties such as inhibition of serotonin (5-hydroxytryptamine; 5-HT) reuptake. Serotonergic pharmacotherapy is associated with amelioration of depression as well as increases in natural killer cell activity (Helgason et al., 2000). Postmarketing surveillance conducted by Rudolf and Zeller (2004) that included 4337 depressive patients suffered from mild to moderate depression shows that a single-dose therapy with highly dosed St. John's wort extract reduces severity of depression and significantly improving the quality of life.

St. John's wort no doubt possesses various health benefits, but questions arose about its efficiency. It interferes with certain drugs such as xenobiotics and reduces their efficiency. St. John's wort activates a nuclear receptor called pregnane X receptor (PXR). PXR is a ligand-activated transcription factor that induces a number of xenobiotic-metabolizing enzymes and transporters including cytochrome P450A4 (CYP3A4) in humans. Because CYP3A4 alone metabolizes about 60% of all clinically relevant drugs, induction of CYP3A4 may result in the rapid elimination of these drugs and a consequent reduction in drug efficacy (Choudhuri and Valerio, 2005). Thus, care should be taken for its use and its responsibility of clinician to document use of herbs of botanicals to cure various maladies.

### **CONCLUSIONS**

Use of herbals/botanicals has been gaining wide popularity in recent years in the United States as well as in other parts of the world, but the mechanism of action of most of these herbals/botanicals has not been subjected to thorough scientific investigations. Plants and botanicals hold therapeutic potential in clinical therapy to prevent or cure certain health risks with additional benefit of reduction in prevention cost. Indeed, findings suggested that these plants and their bioactive metabolites

are effective in balancing and proper functionality of immune system through various modules of immune modification like stimulation and suppression. Thus, current scenario demands form scientific research to explore the mode of actions of the selected botanicals/herbs. Nutritionists, physicians, and other health professionals can use such information effectively for the treatment of various ailments in the vulnerable segments. Overall, certain plants and botanicals can be utilized as an additional tool for disease prevention and risk management.

## REFERENCES

- Abe, M., Akbar, F., Hasebe, A., Horiike, N. and Onji, M. (2003). Glycyrrhizin enhances interleukin-10 production by liver dendritic cells in mice with hepatitis. *J. Gastroenterol.* **38**:962–967.
- Abebe, W. (2003). An overview of herbal supplement utilization with particular emphasis on possible interactions with dental drugs and oral manifestations. *J. Dent. Hyg.* **77**:37–46.
- Abusnina, A., Alhosin, M., Keravis, T., Muller, C. D., Fuhrmann, G., Bronner, C. and Lugnier, C. (2011). Down-regulation of cyclic nucleotide phosphodiesterase PDE1A is the key event of p73 and UHRF1 deregulation in thymoquinone-induced acute lymphoblastic leukemia cell apoptosis. *Cell Signal.* **23**:152–160.
- Ahui, M. L., Champy, P., Ramadan, A., Pham Van, L., Araujo, L., Brou André, K., Diem, S., Damotte, D., Kati-Coulibaly, S., Offoumou, M. A., Dy, M., Thieblemont, N. and Herbelin, A. (2008). Ginger prevents Th2-mediated immune responses in a mouse model of airway inflammation. *Int. Immunopharmacol.* **8**:1626–1632.
- Aktas, O., Prozorovski, T., Smorodchenko, A., Savaskan, N. E., Lauster, R., Kloetzel, P. M., Infante-Duarte, C., Brocke, S. and Zipp, F. (2004). Green tea epigallocatechin-3-gallate mediates T cellular NF-kappa B inhibition and exerts neuroprotection in autoimmune encephalomyelitis. *J. Immunol.* **173**:5794–5800.
- Alban, S., Classen, B., Brunner, G. and Blaschek, W. (2002). Differentiation between the complement modulating effects of an arabinogalactan-protein from *Echinacea purpurea* and heparin. *Planta Med.* **68**:1118–1124.
- Al-Hazzani, A. A. and Alshatwi, A. A. (2011). Catechin hydrate inhibits proliferation and mediates apoptosis of SiHa human cervical cancer cells. *Food Chem. Toxicol.* **49**:3281–3286.
- Almajano, M. P., Vila, I. and Gines, S. (2011). Neuroprotective effects of white tea against oxidative stress-induced toxicity in striatal cells. *Neurotox. Res.* **20**:372–378.
- Altonsy, M. O. and Andrews, S. C. (2011). Diallyl disulphide, a beneficial component of garlic oil, causes a redistribution of cell-cycle growth phases, induces apoptosis, and enhances butyrate-induced apoptosis in colorectal adenocarcinoma cells (HT-29). *Nutr. Cancer.* **63**:1104–1113.
- Arafa, el-S. A., Zhu, Q., Shah, Z. I., Wani, G., Barakat, B. M., Racoma, I., El-Mahdy, M. A. and Wani, A. A. (2011). Thymoquinone up-regulates PTEN expression and induces apoptosis in doxorubicin-resistant human breast cancer cells. *Mutat. Res.* **706**:28–35.
- Ares, G., Giménez, A. and Gámbaro, A. (2009). Consumer perceived healthiness and willingness to try functional milk desserts; influence of ingredient, ingredient name and health claim. *Food Qual. Prefer.* **20**:50–56.
- Atta, A. H., Elkoly, T. A., Mouneir, S. M., Kamel, G., Alwabel, N. A. and Zaher, S. (2010). Hepatoprotective effect of methanol extracts of *Zingiber officinale* and *Cichorium intybus*. *Ind. J. Pharm. Sci.* **72**:564–570.
- Badary, O. A., Abd-Ellah, M. F., El-Mahdy, M. A., Salama, S. A. and Hamada, F. M. (2007). Anticlastogenic activity of thymoquinone against benzo(a)pyrene in mice. *Food Chem. Toxicol.* **45**:88–92.
- Bais, H. P., Venkatachalam, L. and Biedrzycki, M. L. (2010). Stimulation or Inhibition: Conflicting evidence for (+/-)-catechin's role as a chemical facilitator and disease protecting agent. *Plant Signal Behav.* **5**:239–246.
- Baliga, M. S., Haniadka, R., Pereira, M. M., D'Souza, J. J., Pallaty, P. L., Bhat, H. P. and Popuri, S. (2011). Update on the chemopreventive effects of ginger and its phytochemicals. *Crit. Rev. Food Sci. Nutr.* **51**:499–523.
- Baliga, M. S. and Katiyar, S. K. (2006). Chemoprevention of photocarcinogenesis by selected dietary botanicals. *J. Photochem. Photobiol. Sci.* **5**:243–53.
- Banerjee, S., Padhye, S., Azmi, A., Wang, Z., Philip, P. A., Kucuk, O., Sarkar, F. H. and Mohammad, R. M. (2010). Review on molecular and therapeutic potential of thymoquinone in cancer. *Nutr. Cancer.* **62**:938–946.
- Bany, J., Zdanowska, D., Zdanowski, R. and Skopinska-Rozewska, E. (2003). The effect of herbal remedy on the development of *Trichinella spiralis* infection in mice. *Polish J. Vet. Sci.* **6**:6–8.
- Bat-Chen, W., Golan, T., Peri, I., Ludmer, Z. and Schwartz, B. (2010). Allicin purified from fresh garlic cloves induces apoptosis in colon cancer cells via Nrf2. *Nutr. Cancer.* **62**:947–957.
- Bhattacharyya, M., Girish G. V., Karmohapatra, S. K., Samad, S. A. and Sinha, A. K. (2007). Systemic production of IFN-alpha by garlic (*Allium sativum*) in humans. *J. Interfer. Cytok. Res.* **27**:377–382.
- Bonifati, D. M. and Kishore, U. (2007). Role of complement in neurodegeneration and neuroinflammation. *Mol. Immunol.* **44**:999–1010.
- Borek, C. (2001). Antioxidant health effects of aged garlic extract. *J. Nutr.* **131**:1010–1015S.
- Bourgeon, S., Raclot, T., Le Maho, Y., Ricquier, D. and Criscuolo, F. (2007). Innate immunity assessed by plasma No measurement is not suppressed during the incubation fast in eiders. *Dev. Comp. Immunol.* **31**:720–728.
- Brush, J., Mendenhall, E., Guggenheim, A., Chan, T., Connelly, E., Soumyanath, A., Buresh, R., Barrett, R. and Zwick, H. (2006). The effect of *Echinacea purpurea*, *Astragalus membranaceus* and *Glycyrrhiza glabra* on CD69 expression and immune cell activation in humans. *Phytother. Res.* **20**:687–695.
- Butt, M. S. and Sultan, M. T. (2009). Green tea: Nature's defense against malignancies. *Cri Rev. Food Sci. Nutr.* **49**:463–473.
- Butt, M. S. and Sultan, M. T. (2010). *Nigella sativa*: Reduces the risk of various maladies. *Cri Rev. Food Sci. Nutr.* **50**:654–665.
- Butt, M. S. and Sultan, M. T. (2011). Ginger and its health claims: Molecular aspects. *Crit. Rev. Food Sci. Nutr.* **51**:383–393.
- Butt, M. S., Nazir, A., Sultan, M. T. and Schroe'n, K. (2008). *Morus alba* L. nature's functional tonic. *Trends Food Sci Technol.* **19**:505–512.
- Butt, M. S., Sultan, M. T., Butt, M. S. and Iqbal, J. (2009). Garlic: Nature's protection against physiological threats. *Cri Rev. Food Sci. Nutr.* **49**:538–55.
- Chanana, V., Majumdar, S. and Rishi, P. (2007). Involvement of caspase-3, lipid peroxidation and TNF- $\alpha$  in causing apoptosis of macrophages by co-ordinately expressed *Salmonella* phenotype under stress conditions. *Mol. Immunol.* **44**:1551–1558.
- Chandra-Kuntal, K. and Singh, S. V. (2010). Diallyl trisulfide inhibits activation of signal transducer and activator of transcription 3 in prostate cancer cells in culture and in vivo. *Cancer Prev. Res. (Phila).* **3**:1473–1483.
- Chandrashekar, P. M., Prashanth, K. V. and Venkatesh, Y. P. (2011). Isolation, structural elucidation and immunomodulatory activity of fructans from aged garlic extract. *Phytochemistry.* **72**:255–264.
- Chandrashekar, P. M. and Venkatesh, Y. P. (2009). Identification of the protein components displaying immunomodulatory activity in aged garlic extract. *J. Ethnopharmacol.* **124**:384–390.
- Chaouat, G., Ledee-Bataille, N. and Dubanchet, S. (2007). Immune cells in uteroplacental tissues throughout pregnancy: A brief review. *Reprod. Biomed. Online.* **14**:256–266.
- Chen, B. H., Wu, P. Y., Chen, K. M., Fu, T. F., Wang, H. M. and Chen, C. Y. (2009). Antiallergic potential on RBL-2H3 cells of some phenolic constituents of *Zingiber officinale* (ginger). *J. Nat. Prod.* **72**:950–953.
- Chen, C. Y., Huang, C. F., Tseng, Y. T. and Kuo, S. Y. (2012). Diallyl disulfide induces Ca(2+) mobilization in human colon cancer cell line SW480. *Arch. Toxicol.* **86**:231–238.
- Choudhuri, S. and Valerio, L. G. Jr. (2005). Usefulness of studies on the molecular mechanism of action of herbals/botanicals: The case of St. John's wort. *J. Biochem. Mol. Toxicol.* **19**:1–11.
- Choudhury, S. R., Balasubramanian, S., Chew, Y. C., Han, B., Marquez, V. E. and Eckert, R. L. (2011). (-)-Epigallocatechin-3-gallate and DZNep reduce



- polycomb protein level via a proteasome-dependent mechanism in skin cancer cells. *Carcinogenesis*. **32**:1525–32.
- Clement, F. and Venkatesh, Y. P. (2010). Dietary garlic (*Allium sativum*) lectins, ASA I and ASA II, are highly stable and immunogenic. *Int. Immunopharmacol.* **10**:1161–1169.
- Colic, M. and Pavelic, K. (2000). Molecular mechanisms of anticancer activity of natural dietetic products. *J. Mol. Med.* **78**:333–336.
- Cooper, M. D. and Alder, M. N. (2006). The evolution of adaptive immune systems. *Cell*. **124**:815–822.
- Cundell, D. R., Matrone, M. A., Ratajczak, P. and Pierce, J. D. Jr. (2003). The effect of aerial parts of Echinacea on the circulating white cell levels and selected immune functions of the aging male Sprague-Dawley rat. *Int. Immunopharmacol.* **3**:1041–1048.
- Currier, N. L. and Miller, S. C. (2002). The effect of immunization with killed tumor cells with/without feeding of Echinacea purpurea in an erythroleukemic mouse model. *J. Alter. Complem. Med.* **8**:49–58.
- Dai, J. H., Iwatani, Y., Ishida, T., Terunuma, H., Kasai, H., Iwakula, Y., Fujiwara H. and Ito, M. (2001). Glycyrrhizin enhances interleukin-12 production in peritoneal macrophages. *Immunology*. **3**:235–243.
- Dijsselbloem, N., Vanden Bergh, W., De Naeyer, A. and Haegeman, G. (2004). Soy isoflavone phyto-pharmaceuticals in interleukin-6 affections. Multi-purpose nutritionaceuticals at the crossroad of hormone replacement anti-cancer and anti-inflammatory therapy. *Biochem. Pharmacol.* **68**:1171–1185.
- Dorhoi, A., Dobrea, V., Zahan, M. and Virag, P. (2006). Modulatory effects of several herbal extracts on avian peripheral blood cell immune responses. *Phytother. Res.* **20**:352–358.
- Duan, P. and Wang, Z. M. (2002). Clinical study on effect of Astragalus in efficacy enhancing and toxicity reducing of chemotherapy in patients of malignant tumor. *Zhongguo Zhong Xi Yi Jie He Za Zhi*. **22**:515–517.
- El Gazzar, M., El Mezayen, R., Nicolls, M. R., Marecki, J. C. and Dreskin, S. C. (2006). Down regulation of leukotriene biosynthesis by thymoquinone attenuates airway inflammation in a mouse model of allergic asthma. *Biochem. Biophys. Acta*. **1760**:1088–1095.
- Ellinger, S., Müller, N., Stehle, P. and Ulrich-Merzenich, G. (2011). Consumption of green tea or green tea products: Is there an evidence for antioxidant effects from controlled interventional studies? *Phytomedicine*. **18**:903–915.
- Fauci, A. S., Mavilio, D. and Kottlilil, S. (2005). NK cells in HIV infection paradigm for protection or targets for ambush. *Nat. Rev. Immunol.* **5**:835–843.
- Fujimoto, N., Sueoka, N., Sueoka, E., Okabe, S., Suganuma, M., Harada, M. and Fujiki, H. (2002). Lung cancer prevention with (-)-epigallocatechin gallate using monitoring by heterogeneous nuclear ribonucleoprotein B1. *Int. J. Oncol.* **20**:1233–1239.
- Funk, J. L., Frye, J. B., Oyarzo, J. N. and Timmermann, B. N. (2009). Comparative effects of two gingerol-containing Zingiber officinale extracts on experimental rheumatoid arthritis. *J. Nat. Prod.* **72**:403–407.
- Galeone, C., Pelucchi, C., Levi, F., Negri, E., Franceschi, S., Talamini, R., Giacosa, A. and La Vecchia, C. (2006). Onion and garlic use and human cancer. *Am. J. Clin. Nutr.* **84**:1027–1032.
- Gali-Muhtasib, H., Diab-Assaf, M., Boltze, C., Al-Hmaira, J., Hartig, R., Roessner, A. and Schneider-Stock, R. (2004). Thymoquinone extracted from black seed triggers apoptotic cell death in human colorectal cancer cells via a p53-dependent mechanism. *Int. J. Oncol.* **25**:857–866.
- Gamboa-Le'on, M. R., Aranda-Gonz'alez, I., Mut-Mart'ın, M., Garc'ia-Miss, M. R. and Dumonteil, E. (2007). In vivo and in vitro control of Leishmania mexicana due to garlic-induced NO production. *Scand. J. Immunol.* **66**:508–514.
- Gan, X. H., Zhang, L., Heber, D. and Bonavida, B. (2003). Mechanism of activation of human peripheral blood NK cells at the single cell level by Echinacea water soluble extracts: Recruitment of lymphocyte-target conjugates and killer cells and activation of programming for lysis. *Int. Immunopharmacol.* **3**:811–824.
- Genovese, T., Mazzon, E., Menegazzi, M., Di Paola, R., Muia, C., Crisafulli, C., Bramanti, P., Suzuki, H. and Cuzzocrea, S. (2006). Neuroprotection and enhanced recovery with Hypericum perforatum extract after experimental spinal cord injury in mice. *Shock*. **25**:608–617.
- Ghannadi, A., Hajhashemi, V. and Jafarabadi, H. (2005). An investigation of the analgesic and anti-inflammatory effects of Nigella sativa seed polyphenols. *J. Med. Food*. **8**:488–493.
- Guo, F. C., Kwakkel, R. P., Williams, B. A., Parmentier, H. K., Li, W. K., Yang, Z. Q. and Versteegen, M. W. (2004). Effects of mushroom and herb polysaccharides on cellular and humoral immune responses of Eimeria tenella-infected chickens. *Poultry Sci.* **83**:1124–1132.
- Guo, L., Bai, S. P., Zhao, L. and Wang, X. H. (2011). Astragalus polysaccharide injection integrated with vinorelbine and cisplatin for patients with advanced non-small cell lung cancer: Effects on quality of life and survival. *Med Oncol.* **29**:1656–1662.
- Gupta, S., Hussain, T. and Mukhtar, H. (2003). Molecular pathway for (-)-epigallocatechin-3-gallate-induced cell cycle arrest and apoptosis of human prostate carcinoma cells. *Arch. Biochem. Biophys.* **410**:177–185.
- Hajhashemi, V., Ghannadi, A. and Jafarabadi, H. (2004). Black cumin seed essential oil as a potent analgesic and anti-inflammatory drug. *Phytother. Res.* **18**:195–199.
- Hall, H., Fahlman, M. M. and Engels, H. J. (2007). Echinacea purpurea and mucosal immunity. *Int. J. Sports Med.* **28**:792–797.
- Heilmann, J., Winkelmann, K. and Sticher, O. (2003). Studies on the antioxidative activity of phloroglucinol derivatives isolated from hypericum species. *Planta Med.* **69**:202–206.
- Helgason, C. M., Frank, J. L., Johnson, D. R., Frank, M. G. and Hendricks, S. E. (2000). The effects of St. John's Wort (Hypericum perforatum) on NK cell activity in vitro. *Immunopharmacology*. **46**:247–251.
- Hung, J. Y., Hsu, Y. L., Li, C. T., Ko, Y. C., Ni, W. C., Huang, M. S. and Kuo, P. L. (2009). [6]-Shogaol, an active constituent of dietary ginger, induces autophagy by inhibiting the AKT/mTOR pathway in human non-small cell lung cancer A549 cells. *J. Agric. Food Chem.* **57**:9809–9816.
- Huntley, A. L., Thompson, C. J. and Ernst, E. (2005). The safety of herbal medicinal products derived from Echinacea species a systematic review. *Drug Safety*. **28**:387–390.
- Hwang, S. A., Dasgupta, A. and Actor, J. K. (2004). Cytokine production by non-adherent mouse splenocyte cultures to Echinacea extracts. *Clinica. Chimica. Acta*. **343**:161–166.
- Iciek, M., Kwiecień, I. and Włodek, L. (2009). Biological properties of garlic and garlic-derived organosulfur compounds. *Environ. Mol. Muta.* **50**:247–265.
- Iwasaki, M. and Tsugane, S. (2011). Risk factors for breast cancer: Epidemiological evidence from Japanese studies. *Cancer Sci.* **102**:1607–1614.
- Jafri, S. H., Glass, J., Shi, R., Zhang, S., Prince, M. and Kleiner-Hancock, H. (2010). Thymoquinone and cisplatin as a therapeutic combination in lung cancer: In vitro and in vivo. *J. Exp. Clin. Cancer Res.* **1**:29: 87.
- Jeong, C. H., Bode, A. M., Pugliese, A., Cho, Y. Y., Kim, H. G., Shim, J. H., Jeon, Y. J., Li, H., Jiang, H. and Dong, Z. (2009). [6]-Gingerol suppresses colon cancer growth by targeting leukotriene a4 hydrolase. *Cancer Res.* **69**:5584–5591.
- Ju, S. A., Park, S. M., Lee, Y. S., Bae, J. H., Yu, R., An, W. G., Suh, J. H. and Kim, B. S. (2012). Administration of 6-gingerol greatly enhances the number of tumor-infiltrating lymphocytes in murine tumors. *Int. J. Cancer*. **130**:2618–2628.
- Jung, H. W., Yoon, C. H., Park, K. M., Han, H. S. and Park, Y. K. (2009). Hexane fraction of Zingiberis Rhizoma Crudus extract inhibits the production of nitric oxide and proinflammatory cytokines in LPS-stimulated BV2 microglial cells via the NF-kappaB pathway. *Food Chem. Toxicol.* **47**:1190–1197.
- Kang, T. H., Lee, J. H., Song, C. K., Han, H. D., Shin, B. C., Pai, S. I., Hung, C. F., Trimble C., Lim J. S., Kim, T. W. and Wu, T.-C. (2007). Epigallocatechin-3-gallate enhances CD8+ T cell-mediated antitumor immunity induced by DNA vaccination. *Cancer Res.* **67**:802–811.
- Kannappan, R., Gupta, S. C., Kim, J. H., Reuter, S. and Aggarwal, B. B. (2011). Neuroprotection by spice-derived nutraceuticals: You are what you eat! *Mol. Neurobiol.* **44**:142–159.
- Kawai, K., Tsuno, N. H., Kitayama, J., Sunami, E., Takahashi, K. and Nagawa, H. (2011). Catechin inhibits adhesion and migration of peripheral blood B cells by blocking CD11b. *Immunopharmacol. Immunotoxicol.* **33**:391–397.

- Khan, N., Afaq, F., Saleem, M., Ahmad, N. and Mukhtar, H. (2006). Targeting multiple signaling pathways by green tea polyphenol (–)-epigallocatechin-3-gallate. *Cancer Res.* **66**:2500–2505.
- Kim, J. Y. and Kwon, O. (2009). Garlic intake and cancer risk: An analysis using the Food and Drug Administration's evidence-based review system for the scientific evaluation of health claims. *Am. J. Clin. Nutr.* **89**:257–264.
- Kim, M., Miyamoto, S., Yasui, Y., Oyama, T., Murakami, A. and Tanaka, T. (2009). Zerumbone, a tropical ginger sesquiterpene, inhibits colon and lung carcinogenesis in mice. *Int. J. Cancer.* **124**:264–271.
- Kim, M. K., Chung, S. W., Kim, D. H., Kim, J. M., Lee, E. K., Kim, J. Y., Ha, Y. M., Kim, Y. H., No, J. K., Chung, H. S., Park, K. Y., Rhee, S. H., Choi, J. S., Yu, B. P., Yokozawa, T., Kim, Y. J. and Chung, H. Y. (2010). Modulation of age-related NF-kappaB activation by dietary zingerone via MAPK pathway. *Exp. Gerontol.* **45**:419–426.
- Kong, X., Hu, Y., Rui, R., Wang, D. and Li, X. (2004). Effects of Chinese herbal medicinal ingredients on peripheral lymphocyte proliferation and serum antibody titer after vaccination in chicken. *Int. Immunopharmacol.* **4**:975–982.
- Kris-Etherton, P. M., Hecker, K. D., Bonanome, A., Coval, S. M., Binkoski, A. E., Hilpert, K. F., Griel, A. E. and Etherton, T. D. (2002). Bioactive compounds in foods: Their role in the prevention of cardiovascular disease and cancer. *Am. J. Med.* **113**:71S–88S.
- Kroes, B. H., Beukeman, C. J., van den Berg, A. J., Wolbink, G. J., van Dijk, H. and Labadie, R. P. (1997). Inhibition of human complement by beta-glycyrrhetic acid. *Immunology.* **90**:115–120.
- Kumara, S. S. and Huat, B. T. (2001). Extraction, isolation and characterization of anti-tumour principle, alpha-hedrin, from the seeds of *Nigella sativa*. *Planta Med.* **67**:29–32.
- Lee, K. Y. and Jeon, Y. J. (2005). Macrophage activation by polysaccharide isolated from *Astragalus membranaceus*. *Int. Immunopharmacol.* **5**:1225–1233.
- Lee, T. -Y., Lee, K. -C., Chen, S. -Y. and Chang, H. -H. (2009). 6-Gingerol inhibits ROS and iNOS through the suppression of PKC- $\alpha$  and NF- $\kappa$ B pathways in lipopolysaccharide-stimulated mouse macrophages. *Biochem. Biophys. Res. Commun.* **382**:134–139.
- Li, S., Yao, X., Liu, H., Li, J. and Fan, B. (2007). Prediction of T-cell epitopes based on least squares support vector machines and amino acid properties. *Analytica Chimica Acta.* **584**:37–42.
- Li, X. H., McGrath, K. C., Nammi, S., Heather, A. K. and Roufogalis, B. D. (2012). Attenuation of liver pro-inflammatory responses by *Zingiber officinale* via inhibition of NF-kappa B activation in high-fat diet-fed rats. *Basic Clin. Pharmacol. Toxicol.* **110**:238–244.
- Liang, D., Qin, Y., Zhao, W., Zhai, X., Guo, Z., Wang, R., Tong, L., Lin, L., Chen, H., Wong, Y. C. and Zhong, Z. (2011). S-allylmercaptocysteine effectively inhibits the proliferation of colorectal cancer cells under *in vitro* and *in vivo* conditions. *Cancer Lett.* **310**:69–76.
- Liu, Q. Y., Yao, Y. M., Zhang, S. W. and Sheng, Z. Y. (2011). *Astragalus* polysaccharides regulate T cell-mediated immunity via CD11c(high)CD45RB(low) DCs *in vitro*. *J. Ethnopharmacol.* **136**:457–464.
- Liu, Z. G., Xiong, Z. M. and Yu, X. Y. (2003). Effect of *astragalus* injection on immune function in patients with congestive heart failure. *Zhongguo Zhong Xi Yi Jie He Za Zhi.* **23**:351–353.
- Lu, J., Guan, S., Shen, X., Qian, W., Huang, G., Deng, X. and Xie, G. (2011). Immunosuppressive activity of 8-gingerol on immune responses in mice. *Molecules.* **16**:2636–2645.
- Mahmood, M. S., Gilani, A. H., Khwaja, A., Rashid, A. and Ashfaq, M. K. (2003). The *in vitro* effect of aqueous extract of *Nigella sativa* seeds on nitric oxide production. *Phytother. Res.* **17**:921–924.
- Mallikarjuna, K., Sahitya-Chetan, P., Sathyavelu-Reddy, K. and Rajendra, W. (2008). Ethanol toxicity: Rehabilitation of hepatic antioxidant defense system with dietary ginger. *Fitoterapia.* **79**:174–178.
- Manikandan, R., Beulaja, M., Arulvasu, C., Sellamuthu, S., Dinesh, D., Prabhu, D., Babu, G., Vaseeharan, B. and Prabhu, N. M. (2011). Synergistic anticancer activity of curcumin and catechin: An *in vitro* study using human cancer cell lines. *Microsc. Res. Tech.*
- Mansour, M. and Tornhamre, S. (2004). Inhibition of 5-lipoxygenase and leukotriene C4 synthase in human blood cells by thymoquinone. *J. Enz. Inhib. Med. Chem.* **19**:431–436.
- Mantena, S. K., Roy, A. M. and Katiyar, S. K. (2005). Epigallocatechin-3-gallate inhibits photocarcinogenesis through inhibition of angiogenic factors and activation of CD8+ T cells in tumors. *Photochem. Photobiol.* **81**:1174–1179.
- Nagaraj, N. S., Anilakumar, K. R. and Singh, O. V. (2010). Diallyl disulfide causes caspase-dependent apoptosis in human cancer cells through a Bax-triggered mitochondrial pathway. *J. Nutr. Biochem.* **21**:405–412.
- Ng, W. K., Yazan, L. S. and Ismail, M. (2011). Thymoquinone from *Nigella sativa* was more potent than cisplatin in eliminating of SiHa cells via apoptosis with down-regulation of Bcl-2 protein. *Toxicol. In Vitro.* **25**:1392–1398.
- Nievergelt, A., Marazzi, J., Schoop, R., Altmann, K. H. and Gertsch, J. (2011). Ginger phenylpropanoids inhibit IL-1{beta} and prostanoid secretion and disrupt arachidonate-phospholipid remodeling by targeting phospholipases A2. *J. Immunol.* **187**:4140–4150.
- Ning, K. J., Ruan, X. C., Lu, J. F. and Cheng, Y. X. (2005). Effects of Huangqi on phagocytic activity of peritoneal macrophage of mice. *Zhongguo Zhong Yao Za Zhi.* **30**:1670–1672.
- Noonan, D. M., Benelli, R. and Albin, A. (2007). Angiogenesis and cancer prevention: A vision. *Recent Res. Cancer Res.* **174**:219–224.
- Norwood, A. A., Tan, M., May, M., Tucci, M. and Benghuzzi, H. (2006). Comparison of potential chemotherapeutic agents, 5-fluorouracil, green tea, and thymoquinone on colon cancer cells. *Biomed. Sci. Instrumen.* **42**:350–356.
- O'Neill, W., McKee, S. and Clarke, A. F. (2002). Immunological and haematinic consequences of feeding a standardised *Echinacea* (*Echinacea angustifolia*) extract to healthy horses. *Equine. Vet. J.* **34**:222–227.
- Oyagbemi, A. A., Saba, A. B. and Azeez, O. I. (2010). Molecular targets of [6]-gingerol: Its potential roles in cancer chemoprevention. *Biofactors.* **36**:169–178.
- Ozugurlu, F., Sahin, S., Idiz, N., Akyol, O., Ilhan, A., Yigitoglu, R. and Isik, B. (2005). The effect of *Nigella sativa* oil against experimental allergic encephalomyelitis via nitric oxide and other oxidative stress parameters. *Cell Mol. Biol.* **51**:337–342.
- Pabuccuoglu, A., Konyalioglu, S., Bas, M. and Meral, G. E. (2003). The *in vitro* effects of *Hypericum* species on human leukocyte myeloperoxidase activity. *J. Ethnopharmacol.* **87**:89–92.
- Podlogar, J. A. and Verspohl, E. J. (2012). Antiinflammatory effects of ginger and some of its components in human bronchial epithelial (BEAS-2B) cells. *Phytother. Res.* **26**:333–336.
- Ramaa, C. S., Shirode, A. R., Mundada, A. S. and Kadam, V. J. (2006). Nutritionaceuticals an emerging era in the treatment and prevention of cardiovascular diseases. *Curr. Pharm. Biotechnol.* **7**:15–23.
- Raman, P., Dewitt, D. L. and Nair, M. G. (2008). Lipid peroxidation and cyclooxygenase enzyme inhibitory activities of acidic aqueous extracts of some dietary supplements. *Phytother. Res.* **22**:204–212.
- Rani, M. P., Padmakumari, K. P., Sankarikutty, B., Cherian, O. L., Nisha, V. M. and Raghu, K. G. (2011). Inhibitory potential of ginger extracts against enzymes linked to type 2 diabetes, inflammation and induced oxidative stress. *Int. J. Food Sci. Nutr.* **62**:106–110.
- Rogers, J., Perkins, I., van Olphen, A., Burdash, N., Klein, T. W. and Friedman, H. (2005). Epigallocatechin gallate modulates cytokine production by bone marrow-derived dendritic cells stimulated with lipopolysaccharide or muramyl dipeptide or infected with *Legionella pneumophila*. *Exp. Biol. Med. (Maywood).* **230**:645–651.
- Roller, M., Clune, Y., Collins, K., Rechkemmer, G. and Watzl, B. (2007). Consumption of prebiotic inulin enriched with oligofructose in combination with the probiotics *Lactobacillus rhamnosus* and *Bifidobacterium lactis* has minor effects on selected immune parameters in polypectomised and colon cancer patients. *Br. J. Nutr.* **97**:676–684.
- Rooney, S. and Ryan, M. F. (2005). Effects of alpha-hederin and thymoquinone, constituents of *Nigella sativa*, on human cancer cell lines. *Anticancer Res.* **25**:2199–2204.
- Rudolf, G. A. and Zeller, K. (2004). Highly dosed St. John's wort extract improves quality of life. *MMW Fortschr. Med.* **146**:77–80.
- Sabina, E. P., Rasool, M., Mathew, L., Ezilrani, P. and Indu, H. (2010). [6]-Shogaol inhibits monosodium urate crystal-induced inflammation - An *in vivo* and *in vitro* study. *Food Chem. Toxicol.* **48**:229–235.

- Salem, M. L., Alenzi, F. Q. and Attia, W. Y. (2011). Thymoquinone, the active ingredient of *Nigella sativa* seeds, enhances survival and activity of antigen-specific CD8-positive T cells in vitro. *Br. J. Biomed. Sci.* **68**:131–137.
- Salem, M. L. and Hossain, M. S. (2000). Protective effect of black seed oil from *Nigella sativa* against murine cytomegalovirus infection. *Int. J. Immunopharmacol.* **22**:729–740.
- Salim, E. I. and Fukushima, S. (2003). Chemopreventive potential of volatile oil from black cumin (*Nigella sativa* L.) seeds against rat colon carcinogenesis. *Nutr. Cancer* **45**:195–192.
- Sartippour, M. R., Shao, Z. M., Heber, D., Beatty, P., Zhang, L., Liu, C., Ellis, L., Liu, W., Go, V. L. and Brooks, M. N. (2002). Green tea inhibits vascular endothelial growth factor (VEGF) induction in human breast cancer cells. *J. Nutr.* **132**:2307–2311.
- Sasagawa, M., Cech, N. B., Gray, D. E., Elmer, G. W. and Wenner, C. A. (2006). Echinacea alkylamides inhibit interleukin-2 production by Jurkat T cells. *Int. Immunopharmacol.* **6**:1214–1221.
- Sayed-Ahmed, M. M., Aleisa, A. M., Al-Rejaie, S. S., Al-Yahya, A. A., Al-Shabanah, O. A., Hafez, M. M. and Nagi, M. N. (2010). Thymoquinone attenuates diethylnitrosamine induction of hepatic carcinogenesis through antioxidant signaling. *Oxid. Med. Cell Longev.* **3**:254–261.
- Schwager, J., Mohajeri, M. H., Fowler, A. and Weber, P. (2008). Challenges in discovering bioactives for the food industry. *Curr. Opin. Biotechnol.* **19**:66–72.
- Sengupta, A., Ghosh, S., Bhattacharjee, S. and Das, S. (2004). Indian food ingredients and cancer prevention - an experimental evaluation of anticarcinogenic effects of garlic in rat colon. *Asian Pac. J. Cancer Prev.* **5**:126–132.
- Shibata, S. (2000). A drug over the millennia: Pharmacognosy, chemistry, and pharmacology of licorice. *Yakugaku Zasshi.* **120**:849–862.
- Shim, S., Kim, S., Choi, D. S., Kwon, Y. B. and Kwon, J. (2011). Anti-inflammatory effects of [6]-shogaol: Potential roles of HDAC inhibition and HSP70 induction. *Food Chem. Toxicol.* **49**:2734–2740.
- Singh, B. N., Shankar, S. and Srivastava, R. K. (2011). Green tea catechin, epigallocatechin-3-gallate (EGCG): Mechanisms, perspectives and clinical applications. *Biochem. Pharmacol.* **82**:1807–1821.
- South, E. H. and Exon, J. H. (2001). Multiple immune functions in rats fed Echinacea extracts. *Immunopharmacol. Immunotoxicol.* **23**:411–421.
- Stewart, A. J., Mullenm, W. and Crozier, A. (2005). On-line high-performance liquid chromatography analysis of the antioxidant activity of phenolic compounds in green and black tea. *Mol. Nutr. Food Res.* **49**:52–60.
- Sueoka, N., Suganuma, M., Sueoka, E., Okabe, S., Matsuyama, S., Imai, K., Nakachi, K. and Fujiki, H. (2001). A new function of green tea: Prevention of lifestyle-related diseases. *Ann. New York Acad. Sci.* **928**:274–280.
- Sultan, M. T., Butt, M. S., Anjum, F. M., Jamil, A., Akhtar, S. and Nasir, M. (2009). Nutritional profile of indigenous cultivar of black cumin seeds and antioxidant potential of its fixed and essential oil. *Pak. J. Bot.* **41**:32–40.
- Swamy, S. M. and Huat, B. T. (2003). Intracellular glutathione depletion and reactive oxygen species generation are important in alpha-hederin-induced apoptosis of P388 cells. *Mol. Cell. Biochem.* **245**:127–139.
- Tekeoglu, I., Dogan, A. and Demiralp, L. (2006). Effects of thymoquinone (volatile oil of black cumin) on rheumatoid arthritis in rat models. *Phytother. Res.* **20**:869–871.
- Thakur, V. S., Gupta, K. and Gupta, S. (2012). The chemopreventive and chemotherapeutic potentials of tea polyphenols. *Curr. Pharm. Biotechnol.* **13**:191–199.
- Timár, K. K., Dallos, A., Kiss, M., Husz, S., Bos, J. D. and Asghar, S. S. (2007). Expression of terminal complement components by human keratinocytes. *Mol. Immunol.* **44**:2578–2586.
- Tipoe, G. L., Leung, T. M., Hung, M. W. and Fung, M. L. (2007). Green tea polyphenols as an anti-oxidant and anti-inflammatory agent for cardiovascular protection. *Cardiovas. Hematol. Disor. Drug-Targets.* **7**:135–144.
- Truchlinski, J., Krauze, M., Cendrowska-Pinkosz, M. and Modzelewska-Banachiewicz, B. (2006). Influence of garlic, synthetic 1,2,4-triazole derivative and herbal preparation echinovit C on selected indices of turkey-hens non-specific immunity. *Polish J. Vet. Sci.* **9**:51–55.
- Ueda, H., Ippoushi, K. and Takeuchi, A. (2010). Repeated oral administration of a squeezed ginger (*Zingiber officinale*) extract augmented the serum corticosterone level and had anti-inflammatory properties. *Biosci. Biotechnol. Biochem.* **74**:2248–2252.
- van Breemen, R. B., Tao, Y. and Li, W. (2011). Cyclooxygenase-2 inhibitors in ginger (*Zingiber officinale*). *Int. J. Food Sci. Nutr.* **62**:106–110.
- Velho-Pereira, R., Kumar, A., Pandey, B. N., Jagtap, A. G. and Mishra, K. P. (2011). Radiosensitization in human breast carcinoma cells by thymoquinone: Role of cell cycle and apoptosis. *Cell Biol. Int.* **35**:1025–1029.
- Walsh, N. P., Gleeson, M., Pyne, D. B., Nieman, D. C., Dhabhar, F. S., Shephard, R. J., Oliver, S. J., Bermon, S. and Kojenien, A. (2011). Position statement. Part two: Maintaining immune health. *Exerc. Immunol. Rev.* **17**:64–103.
- Wan, D., Pei, J., Zhou, M. and Wang, G. (2003). Antidepressant effect of ethanol extracts from three species of sect. *Hypericum* medicinal plants in mice. *Zhong. Yao. Cai.* **26**:187–189.
- Wang, H., Bian, S. and Yang, C. S. (2011). Green tea polyphenol EGCG suppresses lung cancer cell growth through upregulating miR-210 expression caused by stabilizing HIF-1{alpha}. *Carcinogenesis.* **32**:1881–1889.
- Wang, R. T., Shan, B. E. and Li, Q. X. (2002). Extracorporeal experimental study on immuno-modulatory activity of Astragalus membranaceus extract. *Zhongguo Zhong Xi Yi Jie He Za Zhi.* **22**:453–456.
- Wang, X., Jiao, F., Wang, Q. W., Wang, J., Yang, K., Hu, R. R., Liu, H. C., Wang, H. Y. and Wang, Y. S. (2012). Aged black garlic extract induces inhibition of gastric cancer cell growth in vitro and in vivo. *Mol. Med. Report.* **5**:66–72.
- Wang, Z. H., Hsu, C. C. and Yin, M. C. (2009). Antioxidative characteristics of aqueous and ethanol extracts of glossy privet fruit. *Food Chem.* **112**:914–918.
- Watson, J. L., Vicario, M., Wang, A., Moreto, M. and McKay, D. M. (2005). Immune cell activation and subsequent epithelial dysfunction by *Staphylococcus enterotoxin B* is attenuated by the green tea polyphenol (-)-epigallocatechin gallate. *Cell Immunol.* **237**:7–16.
- Wilson, E. A. and Demmig-Adams, B. (2007). Antioxidant, anti-inflammatory, and antimicrobial properties of garlic and onions. *Nutr. Food Sci.* **37**:178–183.
- Woelkart, K., Marth, E., Suter, A., Schoop, R., Raggam, R. B., Koidl, C., Kleinhapfl, B. and Bauer, R. (2006). Bioavailability and pharmacokinetics of Echinacea purpurea preparations and their interaction with the immune system. *Int. J. Clin. Pharmacol. Ther.* **44**:401–408.
- Wojcikowski, K., Myers, S. and Brooks, L. (2007). Effects of garlic oil on platelet aggregation: A double blind placebo controlled crossover study. *Platelets.* **18**:29–34.
- Woo, C. C., Loo, S. Y., Gee, V., Yap, C. W., Sethi, G., Kumar, A. P. and Tan, K. H. (2011). Anticancer activity of thymoquinone in breast cancer cells: Possible involvement of PPAR-γ pathway. *Biochem. Pharmacol.* **82**:464–475.
- Wu, J., Wang, Y. X., Su, W. L., Zhu, W. X., Lu, J. W. and Li, Z. K. (2006). Effect of Astragalus injection on serious abdominal traumatic patients' cellular immunity. *Chin. J. Integ. Med.* **12**:29–31.
- Wu, P. P., Chung, H. W., Liu, K. C., Wu, R. S., Yang, J. S., Tang, N. Y., Lo, C., Hsia, T. C., Yu, C. C., Chueh, F. S., Lin, S. S. and Chung, J. G. (2011). Diallyl sulfide induces cell cycle arrest and apoptosis in HeLa human cervical cancer cells through the p53, caspase- and mitochondria-dependent pathways. *Int. J. Oncol.* **38**:1605–1613.
- Wu, X. J., Hu, Y., Lamy, E. and Mersch-Sundermann, V. (2009). Apoptosis induction in human lung adenocarcinoma cells by oil-soluble allyl sulfides: Triggers, pathways, and modulators. *Environ. Mol. Mut.* **50**:266–275.
- Xu, G., Ren, G., Xu, X., Yuan, H., Wang, Z., Kang, L., Yu, W. and Tian, K. (2011). Combination of curcumin and green tea catechins prevents dimethylhydrazine-induced glossy carcinogenesis. *Food Chem. Toxicol.* **48**:390–395.
- Xu, H. D., You, C. G., Zhang, R. L., Gao, P. and Wang, Z. R. (2007). Effect of Astragalus polysaccharides and astragalosides on the phagocytosis of *Mycobacterium tuberculosis* by macrophages. *J. Inter. Med. Res.* **35**:84–90.
- Yan, Q., Jiang, Z., Yang, S., Deng, W. and Han, L. (2005). A novel homodimeric lectin from *Astragalus mongholicus* with antifungal activity. *Arch. Biochem. Biophys.* **442**:72–81.
- Yang, G., Zheng, W., Xiang, Y. B., Gao, J., Li, H. L., Zhang, X., Gao, Y. T. and Shu, X. O. (2011). Green tea consumption and colorectal cancer risk: A report from the Shanghai Men's Health Study. *Carcinogenesis.* **32**:1684–1688.

- Yang, L., Liu, Y. L. and Lin, S. Q. (1990). HPLC analysis of flavonoids in the root of six *Glycyrrhiza* species. *Yao Xue Xue Bao*. **25**:840–848.
- Yin, X., Chen, L., Liu, Y., Yang, J., Ma, C., Yao, Z., Yang, L., Wei, L. and Li, M. (2010). Enhancement of the innate immune response of bladder epithelial cells by *Astragalus* polysaccharides through upregulation of TLR4 expression. *Biochem. Biophys. Res. Commun.* **397**:232–238.
- Yip, P. Y. and Kwan, H. S. (2006). Molecular identification of *Astragalus membranaceus* at the species and locality levels. *J. Ethnopharmacol.* **106**:222–229.
- Young, H. Y., Luo, Y. L., Cheng, H. Y., Hsieh, W. C., Liao, J. C. and Peng, W. H. (2005). Analgesic and anti-inflammatory activities of [6]-gingerol. *J. Ethnopharmacol.* **96**:207–210.
- Zamani, A., Vahidinia, A. and Ghannad, M. S. (2009). The effect of garlic consumption on Th1/Th2 cytokines in phytohemagglutinin (PHA) activated rat spleen lymphocytes. *Phytother. Res.* **23**:579–581.
- Zhan, K., Wang, C., Xu, K. and Yin, H. (2008). Analysis of volatile and non-volatile compositions in ginger oleoresin by gas chromatography-mass spectrometry. *Se Pu*. **26**:692–696.
- Zhang, T., Li, Y., Wang, Y., Wang, X., Young, M., Young, M., Douglas, S. D. and Ho, W. (2007). Natural killer cell inhibits human immunodeficiency virus replication in chronically infected immune cells. *Antiviral Res.* **73**:132–139.
- Zhou, H., Deng, Y. and Xie, Q. (2006). The modulatory effects of the volatile oil of ginger on the cellular immune response in vitro and in vivo in mice. *J. Ethnopharmacol.* **105**:301–305.
- Zhou, S. F., Xue, C. C., Yu, X. Q. and Wang, G. (2007). Metabolic activation of herbal and dietary constituents and its clinical and toxicological implications: An update. *Curr. Drug Metabol.* **8**:526–553.
- Zorrilla, E. P., Luborsky, L., McKay, J. R., Rosenthal, R., Houldin, A., Tax, A., McCorkle, R., Seligman, D. A. and Schmidt, K. (2001). The relationship of depression and stressors to immunological assays: A meta-analytic review. *Brain Behav. Immun.* **15**:199–26.