



## Electrolyte minerals intake and cardiovascular health

Noushin Mohammadifard, Carolyn Gotay, Karin H. Humphries, Andrew Ignaszewski, Ahmad Esmailzadeh & Nizal Sarrafzadegan

To cite this article: Noushin Mohammadifard, Carolyn Gotay, Karin H. Humphries, Andrew Ignaszewski, Ahmad Esmailzadeh & Nizal Sarrafzadegan (2018): Electrolyte minerals intake and cardiovascular health, Critical Reviews in Food Science and Nutrition, DOI: [10.1080/10408398.2018.1453474](https://doi.org/10.1080/10408398.2018.1453474)

To link to this article: <https://doi.org/10.1080/10408398.2018.1453474>



Accepted author version posted online: 15 Mar 2018.  
Published online: 04 Apr 2018.



Submit your article to this journal [↗](#)



Article views: 90



View Crossmark data [↗](#)



## Electrolyte minerals intake and cardiovascular health

Noushin Mohammadifard<sup>a,b</sup>, Carolyn Gotay<sup>c</sup>, Karin H. Humphries<sup>d</sup>, Andrew Ignaszewski<sup>d</sup>, Ahmad Esmailzadeh<sup>e</sup>, and Nizal Sarrafzadegan<sup>f,g</sup>

<sup>a</sup>Hypertension Research Center, Cardiovascular Research Institute, Isfahan University of Medical Sciences, Isfahan, Iran; <sup>b</sup>Interventional Cardiology Research Center, Cardiovascular Research Institute, Isfahan University of Medical Sciences, Isfahan, Iran; <sup>c</sup>Centre of Excellence in Cancer Prevention, Faculty of Medicine, School of Population and Public Health, The University of British Columbia, Vancouver, Canada; <sup>d</sup>Division of Cardiology, The University of British Columbia, Vancouver, Canada; <sup>e</sup>Department of Community Nutrition, School of Nutritional Sciences and Dietetics, Tehran University of Medical Sciences, Tehran, Iran; <sup>f</sup>Isfahan Cardiovascular Research Center, Cardiovascular Research Institute, Isfahan University of Medical Sciences, Isfahan, Iran; <sup>g</sup>School of Population and Public Health, Faculty of Medicine, The University of British Columbia, Canada

### ABSTRACT

Appropriate intake of micronutrient, such as electrolyte minerals is critical for the well-being of the cardiovascular health system. However, there are some debates regarding the impacts of dietary and/or supplemental intake of these minerals, on the risk of cardiovascular events and associated risk factors. High sodium intake is adversely associated with the risk of hypertension. Although many reports referred to the positive association of Na intake and cardiovascular events and all-cause mortality, however, other studies indicated that low Na intake is related to higher risk of all-cause mortality and HF-related events. By contrast, dietary potassium, magnesium and calcium have an inverse correlation with cardiovascular events and risk factors, especially with blood pressure. There are some controversies about cardiovascular effects and all-cause mortality of high Ca intake, including no effect, preventive or adverse effect with or without vitamin D. Calcium supplementation might be beneficial for prevention of cardiovascular events and all-cause mortality only in individuals with low intake. Moreover, calcium intake showed a J- or U-shaped association with the risk of cardiovascular diseases. Due to the controversies of the effect of electrolyte minerals especially sodium and calcium intake on cardiovascular events, large scale, well-designed long-term randomized clinical trials are required to evaluate the effect of minerals intake on cardiovascular events and all-cause mortality. In this review, we discuss the role of dietary and or supplemental sodium, potassium, magnesium, calcium, in cardiovascular health, as well as their clinical applications, benefits, and risks for the primary prevention of cardiovascular disease, in general population.

### KEYWORDS

Mineral; cardiovascular; sodium; potassium; magnesium; calcium; mortality

## Introduction

Micronutrients consist of vitamins and minerals that are critical for maintaining health and well-being. As inorganic substances, minerals have various functions in the body including regulation of nerve and muscle function and maintenance of acid-base balance and water balance (Otten et al. 2006). Imbalances in electrolytes like sodium (Na), potassium (K), magnesium (Mg) and calcium (Ca) are frequent and potentially hazardous that may lead to many cardiovascular disease (CVD) like hypertension, coronary heart disease, cardiomyopathy, heart failure, and arrhythmias (Aburto et al. 2013, Chrysant and Chrysant 2014, Kolte et al. 2014, Seth et al. 2014).

Ensuring adequate intake of vitamins and minerals, by diet and/or supplementation, is commonly recommended for health promotion (Fortmann et al. 2013). Thus the main reason of micronutrients supplementation is for prevention of CVD and cancer, leading to improvement of associated risk factors, particularly inflammation and oxidative stress (Pham-Huy et al. 2008, Moyer and Force 2014, Mohammadifard et al.). However, recent reviews in humans, have demonstrated that no

consistent evidence implies that vitamins and minerals supplements affect CVD, cancer, or all-cause mortality in healthy individuals, without known nutritional deficiencies. In addition,

few trials have investigated the impacts of micronutrient supplementation on the primary prevention of CVD and cancer. (Fortmann et al. 2013, Moyer and Force 2014). Although many studies and reviews have examined the beneficial effect of vitamins, as a micronutrient on CVD, but only a few reports have comprehensively and critically examined the effect of minerals on CVD health, their benefits and risks, clinical applications and indications and contraindications (Fortmann et al. 2013, Moyer and Force 2014). We reported the effect of trace minerals including iron, copper, zinc and selenium on CVD health and risk recently. (Mohammadifard et al.). Among minerals, Na, K, Mg and Ca, as electrolytes, can be effective on CV health (Aburto et al. 2013, Chrysant and Chrysant 2014, Das De et al. 2015, Kolte et al. 2014, Smyth et al. 2015). Thus, the aim of this review is to evaluate the role of dietary and supplemental intake

of the aforementioned minerals, in all CV events and risk factors.

### Review criteria

A search was performed for original articles published in PubMed, between January 1990 and October 2017. The search terms used were cardiovascular, coronary heart disease, coronary artery disease, sudden cardiac death, cardiomyopathy, heart failure, arrhythmia, atherosclerosis, hypertension, mineral, sodium, potassium, magnesium and calcium individually and in combination. All articles identified were full-text manuscripts published in English. Clinical and population studies were included.

### Sodium intake and cardiovascular disease

#### Physiology and function

Na is the major cation of the extracellular fluid and functions, as an osmotic determinant in regulating extracellular fluid and plasma volume, in order to balance water in the body, as well as regulating muscle and nerve function. The different functions of Na are provided by a transmembrane pumping of  $\text{Na}^+$ - $\text{K}^+$ -ATPase, accomplishing the active electrogenic translocation of  $\text{Na}^+$  and  $\text{K}^+$ , across the plasma membrane of most of the cells. The balance between Na intake, excretion and extracellular fluid regulate Na maintenance in normal status. A variety of regulators, including renin-angiotensin-aldosterone

system (RAAS) via the sympathetic nervous system and carotid baroreceptors and arterial natriuretic peptide, are involved in Na regulation (Coffman 2014). Figure 1 shows the metabolism of Na in the body.

#### Current dietary recommendations

The global mean Na intake was 3950 mg/day in 2010 (powles et al,2013) inspite of several national and international health organizations recommendations like the World Health Organization (WHO), the American Heart Association (AHA) and the National Institute for Health and Clinical Excellences (NICE) in the UK, to reduce sodium intake to less than 2000, 2400 and 1200 mg/day in the general population, respectively (Eckel et al. 2014, Excellence 2010, World Health Organization 2012)., But the Institute of Medicine (IOM) committee reported lack of enogh evidence to support the beneficial effect of lowering sodium intake to the above recommended levels, on health outcomes other than blood pressure (BP) (Committee on the Consequences of Sodium Reduction. 2013).

#### Role in cardiovascular disease

If there is no decline in peripheral resistance, high intake of dietary Na increases extracellular volume and cardiac output, leading to an increase in BP (Schmidlin et al. 2007). Na-induced hypertension may occur by different impaired physiological mechanisms, including renal function, fluid hormone, salt sensitivity, smooth muscle in peripheral vasculature and sympathetic nervous system, as the physiological mechanism of Na-




Figure 1. Metabolism of sodium, potassium and magnesium in the body. The metabolism, regulation and maintenance in the body.

induced increase in BP (Farquhar et al. 2015).  $\text{Na}^+$  associated with chloride has a different effect compared to when it is binding to other anion such as bicarbonate, therefore, since 1904, it has been considered as a major risk factor for CVD and BP, in the form of NaCl (Ambard and Beaujard 1904). The sodium chloride might have more important role in salt-sensitive hypertension than  $\text{Na}^+$ . Diverse documents from monogenic syndromes, dietary and *in vivo* studies reported, a main role for  $\text{Cl}^-$  on BP regulation through renal balance, and  $\text{Cl}^-$  transporters in vascular tissues (McCallum et al. 2015). A potential hypothesis is inhibition of renin release, due to an increase in  $\text{Cl}^-$  level, delivering to the renal tube (Lorenz et al. 1991). Thus, further studies are suggested to examine the effect of Na and Cl on the risk of CVD and hypertension, independently (McCallum et al. 2015).

### Hypertension

A variety of studies in animals and humans, including ecological, observational, clinical trials and meta-analysis of randomized clinical trials (RCTs), have documented the correlation between high Na intake and BP (Whelton 2015). Large scale observational studies such as INTERSALT in 52 centers worldwide, European Prospective Investigation into Cancer in Norfolk (Khaw et al. 2004) and Prospective Urban, Rural Epidemiological Study (PURE) in 18 countries (Mente et al. 2014), as well as some recent meta-analyses of RCTs, demonstrated a positive correlation between Na and BP that was higher in hypertensive than normotensive populations (Graudal et al. 2012, He et al. 2013). Furthermore, the incidence of hypertension declined, after reducing Na intake in the well-designed Trials of Hypertension Prevention (TOPH) II.

### Cardiovascular events and risk factors

There are many controversies regarding the correlation between Na and CVD events. A meta-analysis of 7 RCTs, using at least 6 months sodium restriction, reported no effect of Na intake reduction on CV mortality, morbidity, or all-cause mortality (Taylor et al. 2011). However in other meta-analysis, after excluding the RCT on heart failure (HF) patients, all CV events showed significant decrease of 20% (He and MacGregor 2011). Since no large, long-term RCTs have been performed to examine the effects of Na restriction on CV events (Whelton et al. 2012), most of the evidence has been derived from prospective cohort studies (O'Donnell et al. 2015), which have known methodological limitations (Cobb et al. 2014). Cobb *et al.* reported that diverse studies showed direct, inverse, J-shaped or null associations between Na intake and CV events with 77%, 75%, 100% or 100% systematic error and 31%, 38%, 50% or 0% reverse causality error, respectively (Cobb et al. 2014).

A pooled analysis of four large prospective cohort studies in 49 countries, using diverse methods for Na assessment in various populations, have found a J or U-shaped association between Na intake and CVD mortality and morbidity (O'Donnell et al. 2014b). A meta-analysis of 23 observational studies showed that Na intake less than 3 g/day, increased all-cause mortality and thus argued the recommendation of Na intake reduction, except in hypertensives or subjects with the high Na intake (more than 6 g/day), unless a well done RCT confirm the safety and beneficial effect (Saulnier et al. 2014). It was

assumed that although low Na consumption is expected to improve CV events by lowering BP, but an increase also have been found in CV events and mortality, which might be through RAAS activation, catecholamine and lipid disorders (Saulnier et al. 2014).

The PURE study found that an estimation of Na intake of 3–6 g/day based on single morning spot urine sample was associated with a lower risk of CV events and all-cause mortality compared to lower or higher estimated Na intake (O'Donnell et al. 2014a). However, Cook has criticized the J-shape association of Na intake and CVD mortality in PURE study by indicating that methodological weakness exist, such as the Na assessment method using an overnight Na excretion, reverse causality relevant to recruiting, some patients who reduced their Na intake for therapeutic approaches, using single morning spot urine and non-valid formula, causing under- or over-estimation of Na intake (Cook 2014). The large size of PURE study may not eradicate different biases including selection, confounders or reverse causality (Cook 2014, Cook et al. 2014). Whereas, in TOHP study, Na intake was assessed by an average of three to seven 24-hr urine collections, performed during 1–4 years (Cook et al. 2014) as a most precise method. Recent trial in TOHP study concluded a direct association between Na intake and mortality risk, even at the lowest levels of Na intake, over a 20-year period (Cook 2014). However, it should be notified that the most precise and reliable method for assessment of Na intake, individually is multiple 24-hr urine collection (Lerchl et al. 2015). Therefore, further studies are required in order to examine the effect of Na on CVD events and risk factors, using multiple 24-hr urine collection.

### Endothelial dysfunction

High Na intake may increase the risk of left ventricular (LV) hypertrophy, endothelial dysfunction, vascular dysregulation and remodeling, and renal disease independent of BP (Aaron and Sanders 2013). This aldosterone regulates the activity of epithelial sodium channels in the renal and endothelia, which may damage on these cell to swell, stiffen, and alter its output of nitric oxide (Greaney et al. 2012). The protective barrier of endothelial glycocalyx buffer interferes with high Na intake. It may be due to limiting the sodium pump (i.e. the Na/ K ATPase) and as a mesh of anionic biopolymers covering the endothelium surface, contributes in the progression of stiffness process and thus enhances the permeability of vascular sodium (Oberleithner 2012). Only acute Na loading, can leads to an increase in microvascular Na permeability and induces plasma volume enhancement. It may progress by direct effects of  $\text{Na}^+$  to the endothelium (Rorije et al. 2015). Na restriction improves endothelial dysfunction and elastic artery stiffness, independent of BP (Jablonski et al. 2009).

### Heart failure

Na restriction induces neurohormonal activation and elevates plasma renin activity, which may increase the risk of mortality in HF patients (Verma et al. 2011). Hyponatraemia is considered as a significant determinant of all-cause mortality and HF-related events, such as HF readmissions irrespective of ejection fraction (Rusinaru et al. 2012). Table 1 illustrates

**Table 1.** Mechanism, clinical application, indications and contraindications in CVD prevention.

Mineral	Regulation	Mechanism	Clinical application in CVD	Type of study	Indications	Contraindications
Sodium	Oral uptake	Inducing hypertension by impaired:	Extracellular fluid	Animal	Dehydration	Hypertension
	Kidney excretion	Renal function	Osmotic determinant	Ecological	High sweating	Allergy
	RAAS	Fluid hormone	Muscle function	Cross-sectional		
Potassium	Sympathetic nervous system	Salt sensitivity	Nerve function	Prospective cohort study		
	Carotid baroreceptors	Smooth muscle in peripheral vasculature		RCT		
	Arterial natriuretic peptide					
magnesium	Oral uptake	Sympathetic nervous system	Intracellular osmolality	Meta-analysis	Hypokalemia	Diarrhea
	Kidney excretion	The BP lowering effect is exerted through:	Resting potential cell membrane	Cross-sectional		
	Aldosterone	Natriuresis stimulation		Prospective cohort study	Prolonged illness with diarrhea or vomiting.	Bloating
Calcium	Liver circulation	Endothelial function improvement		RCT		Nausea
		NO release increasing Sodium-potassium pump and plasma membrane potassium channels activity		Meta-analysis		Abdominal painVomiting
	Muscle storage	Reduced sympathetic activity				
magnesium	Oral uptake	Regulating BP by:	Coenzyme in the production and transport	Cross-sectional	Mg deficiency	Heart block
	Kidney excretion	Improving vascular tone	Keep normal heart rhythm	Prospective cohort	Preeclampsia	Myocardial damage
		Vascular smooth muscle contraction	Vascular smooth muscle tone	RCT	BP control	
Calcium	Vitamin D	Endothelial cell function	Glucose and insulin metabolism	Meta-analysis	Encephalopathy	
	Calcitonin	Myocardial excitability			Acute nephritis in children	
	Parathyroid hormone	Normal BP				
Calcium		Bone integrity	Muscle contraction	Cross-sectional	Calcium deficiency	Tumor
		Increase carotid plaque thickness	Vascular tone	Prospective cohort	Osteoporosis	Kidney Stone
		Calcification	Nerve transmission	RCT	Premenstrual syndrome	Kidney Disease
Calcium		BP regulation	Hormonal system	Meta-analysis	Preeclampsia	Hyperparathyroidism
		Insulin secretion and sensitivity	Enzyme-intermediate		Prevention of colon and rectal cancers	Hypocalcaemia
		Obesity				Extreme dehydration
Calcium		Serum lipids				
		Inflammatory agents				
		Anti-thrombotic agent				
Calcium		Vasorelaxation				

RAAS: Renin-angiotensin-aldosterone system; RCT: Randomised clinical trial; BP: Blood pressure; NO: Nitric oxide; GI: Gastrointestinal; ROS: Reactive oxygen species.



regulation, mechanism, clinical application in CVD, type of study, indications and contraindications of the minerals.

### **Potassium intake and cardiovascular disease**

#### **Physiology and function**

K is the major intracellular cation (98%) in the body (Ekmekcioglu et al. 2016). The main functions of K are resting potential of the cell membrane and regulation of water and Na metabolism in the cell by intracellular osmolarity, acid-base balance, normalization of heartbeat, controlling nerve impulses in muscles and, participation in protein biosynthesis and the conversion of blood sugar into glycogen and the activation of a number of enzymes, especially those involved in energy production (Table 1). The total K concentration is regulated by oral uptake, kidney excretion and circulation in the liver and muscle storage (Eaton and Konner 1985a). The kidney retains Na and excretes K, facilitated by aldosterone function (Eaton and Konner 1985b) (Figure 1).

#### **Current dietary recommendations**

Following industrialization, Na intake has been increased, while K intake has been decreased. Thus nowadays dietary guidelines believe that K is a “nutrient of public health concern” (Palmer and Clegg 2016). Different international organizations like the IOM and WHO recommend 90–120 mmol/day (3500–4700 mg/ day) of K for adolescents and adults (Otten et al. 2006, WHO 2012). Global K intake was evaluated as inadequate and less than 70–80 mmol/day (2700–3100 mg/ day) in the INTERSALT study. About 2% of US adult population has adequate consumption of K (Cogswell et al. 2012). K intake among adult population of 10 European countries ranged from 3536 to 4870 mg/ day in men and from 2730 to 3723 mg/ day in women (Welch et al. 2009), respectively. Its intake levels in two Asian populations are 1800 mg/ day in Chinese and 2732 in Iranians respectively (Du et al. 2014, Mohammadifard et al. 2017).

#### **Role in cardiovascular disease**

The crucial role of K in CV system and the importance of K maintenance have become an essential issue in cardioprotective treatment. Abnormal serum levels of K may affect CVD, including hypertension, arrhythmia, fatal and non-fatal stroke and myocardial infarction (MI), as well as sudden cardiac death (Table 1) (Sica et al. 2002).

#### **Hypertension**

In 1928, Addison proposed an inverse relationship between K and BP (Bulpitt 1981), corroborated by the INTERSALT study, demonstrating an inverse correlation between urinary K excretion and BP in various populations. The BP lowering effect of K is exerted through a variety of mechanisms, including natriuresis stimulation, endothelial function improvement, NO release, increasing Na-K pump function and plasma membrane K channel activity, which causes vasodilatation and reduced sympathetic activity, leading to relaxation of arterial muscle (Ekmekcioglu et al. 2016). Dietary and supplemental K intake had a dose-response effect on BP (Geleijnse et al. 2003, Khaw et al. 2004, Mente et al. 2014), but the beneficial effect plateaus,

above 120 mmols/ day of K consumption (Aburto et al. 2013). In addition, increasing K intake was more beneficial in hypertensives, those with high Na intake (Aburto et al. 2013) especially, more than 4 g/ day (Powles et al. 2013). Thus it was concluded that a low Na and high K diet, might be more effective in reducing BP and hypertension risk (Whelton 2014). Whereas some observational studies had inconsistent findings. High intake of Na and low intake of K did not associate with raised BP levels in the normotensive US adult population and serum K level (Sharma et al. 2014). However, among Chinese population, a community-based cohort study, demonstrated a positive correlation between baseline serum K level, but not serum level of Na and incidence of hypertension, after five years (Xi et al. 2015). However, this study had some limitations, including very low sample size and also the serum K level at baseline could not identify K status.

#### **Cardiovascular events and risk factors**

Some prospective cohort studies showed that higher dietary K intake was associated with lower risk of CV mortality, after adjustment for potential confounders, including dietary Na intake (Umesawa et al. 2008), as well as all-cause mortality and ischemic stroke, but not hemorrhage type in non-hypertensive women (Seth et al. 2014). However a recent cohort study showed dietary Na to K ratio directly increased the risk of mortality from all strokes, even haemorrhagic one. CVD and all causes mortality. Two meta-analyses were done on prospective cohort studies. A recent review in diverse populations, demonstrated an inverse association between K intake and stroke risk by an overall 24% reduction in stroke incidence and the highest reduction of 30% and 21% in subjects with K intake of 90–120 mmol/ day and 42 mmol/ day, respectively, but there was no significant correlation with total CVD events and coronary heart disease (CHD) (Aburto et al. 2013, D’Elia et al. 2011). However, it should be considered that for assessment of K intake similar to Na intake in individual, multiple rounds of 24-hr urine collection are needed (Birukov et al. 2016).

#### **Heart failure**

Mild to moderate hypokalemia in patients with HF, LV hypertrophy and cardiac ischemia, leads to cardiac arrhythmias. Hypokalemia is an independent predictor of mortality and sudden cardiac death in HF patients (Dursun and Sahin 2006). Hypokalemia may be a frequent side effect of treatment with diuretics and digoxin, increased activity of neurohormonal or disease progression (Tomat et al. 2011). A pooled analysis of three prospective cohorts of hospitalized HF patients, illustrated that a reduction of >15% in serum K was an independent predictor of all-cause mortality (Salah et al. 2015).

### **Magnesium intake and cardiovascular disease**

#### **Physiology and function**

Mg is an abundant mineral needed for living cells in the human body, playing a critical role as a cofactor in about 300 metabolic reactions (Volpe, 2013). It is involved in several essential physiological, biochemical functions in the body, such as acting as a coenzyme in the production of DNA, RNA, proteins and energy, as well as energy storage (Guasch-Ferre et al. 2014),

mitochondrial membrane stability, maintenance of normal muscle and nerve functions, normal heart rhythm, vascular smooth muscle tone, endothelial cell function, myocardial excitability, normal BP, bone integrity, and glucose and insulin metabolism (Volpe 2013) (Figure 1). Mg regulates BP by improving vascular tone and vascular smooth muscle contraction (Table 1) (Champagne 2008).

#### **Current dietary recommendations**

The Recommended Dietary Allowance (RDA) for Mg in adults is 310–420 mg/d, including 410–420 mg/day and 310–320 mg/day for men and women, respectively (Food and Nutrition Board 1997). Mg deficiency is commonly seen, worldwide. Marginal or subclinical Mg deficiency induced by Mg intakes < 250 mg/day, and serum Mg concentrations  $\leq 0.75$  mmol/l is associated with inflammatory indicators including C-reactive protein, cytokines such as tumor necrosis factor- $\alpha$  and interleukin (IL)-1, IL-6, E-selectin, intracellular adhesion molecule-1, vascular cell adhesion molecule-1, and fibrinogen (Nielsen 2014, WHO 2012).

#### **Role in cardiovascular disease**

Mg is associated with a number of conditions and chronic diseases. Increasing dietary intake and oral supplement of Mg intake can prevent inflammatory and oxidative stress, endothelial dysfunction, increased vascular reactivity, increased vascular tone, hypertension, CVD, stroke, insulin resistance, type 2 diabetes mellitus and obesity (Champagne 2008, Geiger and Wanner 2012, Qu et al. 2013, Fang et al. 2016). Thus the consumption of Mg-rich foods is the best way to maintain a normal serum Mg level, which consequently prevent the associated health problems (Table 1).

#### **Hypertension**

Mg intake and serum concentration levels are correlated with BP and hypertension (Qu et al. 2013, Champagne 2008). A meta-analysis showed that each 10 mmol/day increase in Mg by supplementation, was associated with a 4.3 mm Hg reduction in systolic BP and a 2.3 mm Hg reduction in diastolic BP (Qu et al. 2013). However, there were some controversies on the inverse correlation between dietary Mg intake as well as serum level and the risk of hypertension (Khan et al. 2010, Song et al. 2006, Bain et al. 2015, Grober et al. 2015). One review suggested that the effect of dietary intake of Mg on BP reduction is stronger than supplementation (Champagne 2008). However, no studies have reported adverse effects for Mg supplementation.

#### **Cardiovascular events and risk factors**

An inverse association was reported between low serum Mg and C-reactive protein, Coronary Artery Calcium (CAC), as well as Carotid Intima-Media Thickness (CIMT), both as main predictors of atherosclerosis (Simon et al. 2002, Posadas-Sanchez et al. 2016). Recent RCTs in obese subjects pointed out that Mg supplements of 350 mg/day in 24 weeks improved arterial stiffness (Joris et al. 2016).

There are beneficial effects of both dietary and supplemental intake, as well as serum concentrations of Mg on different CV events (Del Gobbo et al. 2013, Champagne 2008). An increase

of 150 mg/d to 400 mg/d in Mg intake or an increase of 0.1 mEq/L in serum Mg level, caused 9% reduction in CVD risk (Qu et al. 2013). The benefits of increasing Mg intake on CVD risk factors, such as improvement in lipid profiles, diabetes mellitus, obesity, and metabolic syndrome have been shown in several studies (Del Gobbo et al. 2013, Champagne 2008). In addition, rare reports exist on hypermagnesaemia or Mg toxicity in patients with normal renal function. However, parenteral administration of the Mg supplement is contraindicated in patients with heart block or myocardial damage (Waksman and Ajani 2009).

#### **Calcium intake and cardiovascular disease**

##### **Physiology and function**


Ca is the fifth most plentiful mineral in the body. Typically 99% of Ca is retained in the skeleton. In addition to its role in bone structural strength, it is a pivotal component in many physiologic pathways like muscle contraction, vascular tone, blood coagulation, nerve transmission (e.g. acetylcholine from pre-synaptic terminals of nerves), hormonal system involved in digestion, energy and fat metabolism (e.g. insulin from the beta cells in the pancreatic islets) and enzyme-intermediate in  $\alpha$ -amylase, phospholipase and protein-kinase (Vaskonen 2003). The potential roles of Ca on BP modification, are due to Ca regulating hormone, sympathetic nervous system, and electrolyte interactions (Figure 2). Ca deficient diet increases intracellular Ca level which enhances production 1,25-dihydroxyvitamin D<sub>3</sub> and parathyroid hormone (PTH). It causes Ca uptake into vascular smooth muscle cells and consequently increases vascular resistance and BP (Hatton et al. 1995). Ca regulates RAAS activity by inhibiting acute renin secretion and expression. In addition, chronic Ca receptor activation associates with increasing the activity of renin. PTH secretion stimulates due to increasing aldehyde or angiotensin II (Vaidya et al. 2015). Bones provide a pool for these vital functions by remodeling. Vitamin D, calcitonin and PTH regulate Ca homeostasis (Table 1) (Mundy and Guise 1999).

##### **Current dietary recommendations**

The IOM recommended Dietary Reference Intake (DRI) for Ca, as 1000 mg/day for adults, 1200 mg/day for women aged over 51 and men over 71 years to prevent osteoporosis and bone fractures. The Tolerable Upper Intake Levels (ULs) of Ca is 2500 and 2000 mg per day for adults and elderly, respectively (Ross et al. 2011). Ca supplementation is a common recommendation in the management and prevention of osteoporosis in individuals at risk, since insufficient intake of Ca was seen in most regions in the world (Shin and Kim 2015). The dietary Ca intake ranges from 175 to 1233 mg/day among 74 countries worldwide (Balk et al. 2017).

##### **Role in cardiovascular disease**

Dietary Ca has beneficial effect on CV system through some potential mechanisms including improvement of serum lipids, BP, obesity, insulin secretion and sensitivity, inflammatory and anti-thrombotic agents and vasorelaxation (Chrysant and Chrysant 2014, Tomat et al. 2011).



**Figure 2. Metabolism of calcium in the body.** The metabolism, regulation and maintenance in the body.

However, increase in carotid plaque thickness, aortic calcification, fatal and non fatal CV events similar to hypercalcemic patients may occur subsequently by using Ca supplement. Since it may rapidly increase serum Ca for several hours after consumption, and subsequently pyrophosphate bind to Ca-sensing receptor (Reid et al. 2010). Dietary sources of Ca that are ingested slowly during a day, do not consequently increase the serum Ca concentration, since it is regulated by calciotropic hormones (Wang et al. 2012, Lutsey and Michos 2013). Polymorphisms of the Ca-sensing receptor can also lead to altered serum Ca levels and thus can cause vascular events (Reid et al. 2016).

### Cardiovascular events and risk factors

There are some controversies about the effects of high Ca intake on CV events. Two meta-analysis of RCTs have reported that there is no significant association between dietary and supplemental Ca ranged from 500 to 1600 mg/ day (with and without vitamin D) and CHD, MI, stroke, total CVD risks, angina pectoris, acute coronary syndrome and all-cause mortality (Lewis et al. 2015, Wang et al. 2012). Whereas, a meta-analysis of cohort studies (Reid 2013) are popularly known as Women's Health Initiative (WHI), found that 1000 Ca supplemental with and without vitamin D have increased the risk of MI and stroke (Bolland et al. 2011). Other meta-analysis of observational studies demonstrated that total Ca intake is associated with increased risk of CVD mortality in studies with long term follow-up, and decreased risk of all-cause and CVD mortality, in studies with follow-up  $\leq 10$  years. Ca intake inversely related to all-cause mortality risk (Asemi et al. 2015). In a recent cross-sectional study, Ca intake was associated with atrial fibrillation,

which might be one of potential pathways through which, Ca supplementation may cause strokes (Thiele et al. 2015). The adverse effect of Ca supplementation may be due to excess intake, such as 2000–2500 mg/ day (Daly and Ebeling 2010). The Swedish Mammography Cohort found a J-shaped correlation between Ca intake and mortality, since Ca supplementation was positively associated with a mortality rate in women with a dietary Ca intake of  $>1400$  mg/day and it was higher in individuals with Ca intake  $<600$  mg/day (Michaelsson et al. 2013). A meta-analysis by Wang showed U-shaped manner for association CVD mortality and Ca intake and it was not significant with Ca more than 900 mg/ day (Wang et al. 2014). Recently, other meta-analysis by Reid *et al.* found a significant association between serum Ca levels within the normal range and increased risk of CHD, CVD and total mortality as well as plaque, and vascular calcification. The attenuation of this association after adjustment for CVD risk factors including hypertension and dyslipidemia, implies that these risk factors can be the mediators in the correlation of serum Ca and CV events (Reid et al. 2016). The reverse causality is unlikely to be an issue, because serum Ca measurement was performed about 11–30 years before the occurrence of CV events (Reid et al. 2016).

While a recent prospective cohort study performed on 41,514 non-hypertensive elderly populations showed that dietary Ca up to 1346 mg/ day, has a beneficial effect on non-fatal CVD events, strokes and all-cause mortality (Khan et al. 2015), a recent cross-sectional study demonstrated that low serum Ca has a positive relationship with left ventricular systolic dysfunction in CHD patients with and without AMI (Wang et al. 2012).



## Hypertension

Studies on the correlation between Ca and BP go back to the 1980s (McCarron et al. 1982), and it was confirmed after 20 years (McCarron and Reusser 1999). Clinical and experimental studies demonstrated that Ca has some beneficial lowering effect on BP, which was mediated by suppression of renin activity and an improvement in BP rising effect of Na, particularly in salt-sensitive hypertension. In US population, the risk of hypertension was 11–14% vs. 3–6% in those who took Ca less than 300 mg/ day, compared to subjects with 1200 mg/ day of Ca intake (Singh et al. 1987). However, cross-sectional studies showed that serum Ca levels has direct correlation with BP. Evidence proposed acute rising of BP, following an acute increase in Ca levels might be due to the Ca effect on the contraction of vascular smooth muscle (Reid et al. 2016).

## Summary

In summary, there are inconsistent findings regarding the correlation between minerals and CV events, their risk factors and mortality. The incidence of hypertension declined after reducing Na intake in well designed trials. However, there are many controversies, regarding the correlation between Na and CV events. Although, high Na intake is associated with increased risk of CV events and all-cause mortality, several studies indicated that low Na intake and low to normal serum Na level were significantly associated with all-cause mortality and HF-related events, such as HF readmissions. Most evidence support an inverse correlation between dietary K, Mg and Ca with CV events and risk factors, especially BP. Dietary and supplemental K intake have a dose-response effect on BP, but the beneficial effect plateaus above 120 mmols/day of K intake. Higher dietary K intake was associated with lower rates of BP, CV risk and all-cause mortality. A diet that combines low Na and high K might be more effective in reducing BP and hypertension risk. Mg deficiency is associated with a number of conditions and chronic diseases like inflammatory and oxidative stress, endothelial dysfunction, increased vascular reactivity, increased vascular tone, hypertension, CVD, stroke, insulin resistance and type 2 diabetes mellitus. Diverse studies reported the beneficial effects of dietary and supplemental intake of Mg, as well as serum concentrations of Mg on different CV events and risk factors. There is an inverse association between low serum Mg and CAC and CIMT, as main predictors of atherosclerosis. Calcium supplement may rapidly increases serum Ca and subsequently increase carotid plaque thickness, aortic calcification, fatal and non-fatal CV events similar to hypercalcemic patients, but dietary sources of Ca that are ingested slowly during a day, do not increase immediately the serum Ca concentration. There are some controversies about CV effects and all-cause mortality of high Ca intake, including no effect, preventive or adverse with or without vitamin D. A U-shaped correlation between Ca level and CV events has been suggested. The potential adverse effect of Ca intake on CV events has not been supported by most existing evidence. Ca supplementation may be beneficial in individuals with low intake for prevention of fracture and reduction in all-cause

mortality. The adverse effect of Ca supplementation may be due to excess intake, such as 2000–2500 mg/ day. As future directions, it is important to conduct a large, well-designed RCT with long term follow-up, in order to rigorously evaluate the effect of the mineral intake on CV events, risk factors, mortality and all-cause mortality in normal population.

## References

- Aaron, K. J., and P. W. Sanders 2013. Role of dietary salt and potassium intake in cardiovascular health and disease: a review of the evidence. *Mayo Clinic Proceedings* 88:987–95. doi:10.1016/j.mayocp.2013.06.005.
- Aburto, N. J., S. Hanson, H. Gutierrez, L. Hooper, P. Elliott, and F. P. Cappuccio 2013. Effect of increased potassium intake on cardiovascular risk factors and disease: systematic review and meta-analyses. *BMJ* 346: f1378. doi:10.1136/bmj.f1378.
- Ambard, L., and E. Beaujard 1904. Causes de l'hypertension arterielle. *Arch Gen Med* 1:520–33.
- Asemi, Z., P. Saneei, S. S. Sabihi, A. Feizi, and A. Esmailzadeh 2015. Total, dietary, and supplemental calcium intake and mortality from all-causes, cardiovascular disease, and cancer: A meta-analysis of observational studies. *Nutrition, Metabolism and Cardiovascular Diseases* 25:623–34. doi:10.1016/j.numecd.2015.03.008.
- Bain, L. K., P. K. Myint, A. Jennings, M. A. Lentjes, R. N. Luben, K. T. Khaw, N. J. Wareham, and A. A. Welch 2015. The relationship between dietary magnesium intake, stroke and its major risk factors, blood pressure and cholesterol, in the EPIC-Norfolk cohort. *International Journal of Cardiology* 196:108–14. doi:10.1016/j.ijcard.2015.05.166.
- Balk, E. M., G. P. Adam, V. N. Langberg, A. Earley, P. Clark, P. R. Ebeling, A. Mithal, R. Rizzoli, C. A. F. Zerbin, D. D. Pierroz, et al. 2017. Global dietary calcium intake among adults: A systematic review. *Osteoporosis International* 28:3315–24. doi:10.1007/s00198-017-4230-x.
- Birukov, A., N. Rakova, K. Lerchl, R. H. Engberink, B. Johannes, P. Wabel, U. Moissl, M. Rauh, F. C. Luft, and J. Titze 2016. Ultra-long-term human salt balance studies reveal interrelations between sodium, potassium, and chloride intake and excretion. *American Journal of Clinical Nutrition* 104:49–57. doi:10.3945/ajcn.116.132951.
- Bolland, M. J., A. Grey, A. Avenell, G. D. Gamble, and I. R. Reid 2011. Calcium supplements with or without vitamin D and risk of cardiovascular events: reanalysis of the Women's Health Initiative limited access dataset and meta-analysis. *Bmj* 342:d2040. doi:10.1136/bmj.d2040.
- Bulpitt, C. J. 1981. Sodium excess or potassium lack as a cause of hypertension: A discussion paper. *Journal of the Royal Society of Medicine* 74:896–900. doi:10.1177/014107688107401208.
- Champagne, C. M. 2008. Magnesium in hypertension, cardiovascular disease, metabolic syndrome, and other conditions: a review. *Nutrition in Clinical Practice* 23:142–51. doi:10.1177/0884533608314533.
- Chrysant, S. G., and G. S. Chrysant 2014. Controversy regarding the association of high calcium intake and increased risk for cardiovascular disease. *Journal of Clinical Hypertension (Greenwich)* 16:545–50. doi:10.1111/jch.12347.
- Cobb, L. K., C. A. Anderson, P. Elliott, F. B. Hu, K. Liu, J. D. Neaton, P. K. Whelton, M. Woodward, and L. J. Appel 2014. Methodological issues in cohort studies that relate sodium intake to cardiovascular disease outcomes: a science advisory from the American Heart Association. *Circulation* 129:1173–86. doi:10.1161/CIR.0000000000000015.
- Coffman, T. M. 2014. The inextricable role of the kidney in hypertension. *Journal of Clinical Investigation* 124:2341–47. doi:10.1172/JCI72274.
- Cogswell, M. E., Z. Zhang, A. L. Carriquiry, J. P. Gunn, E. V. Kuklina, S. H. Saydah, Q. Yang, and A. J. Moshfegh 2012. Sodium and potassium intakes among US adults: NHANES 2003–2008. *American Journal of Clinical Nutrition* 96:647–57. doi:10.3945/ajcn.112.034413.
- COMMITTEE ON THE CONSEQUENCES OF SODIUM REDUCTION IN, P., FOOD, NUTRITION, B., BOARD ON POPULATION, H., PUBLIC HEALTH, P., and INSTITUTE OF, M. 2013. In: STROM, B. L., A. L. YAKTINE, and M. ORIA. (eds.) *Sodium Intake in Populations: Assessment of Evidence*. Washington (DC): National Academies Press (US)

- Cook, N. R. 2014. Sodium and cardiovascular disease. *New England Journal of Medicine* 371:2134. doi:10.1056/NEJMc1412113.
- Cook, N. R., L. J. Appel, and P. K. Whelton 2014. Lower levels of sodium intake and reduced cardiovascular risk. *Circulation* 129, 981–89. doi:10.1161/CIRCULATIONAHA.113.006032.
- Cooper, S., J. Taggar, S. Lewis, N. Marlow, A. Dickinson, R. Whitmore, and T. Coleman 2014. Effect of nicotine patches in pregnancy on infant and maternal outcomes at 2 years: Follow-up from the randomised, double-blind, placebo-controlled SNAP trial. *The Lancet Respiratory Medicine* 2:728–37. doi:10.1016/S2213-2600(14)70157-2.
- D'elia, L., G. Barba, F. P. Cappuccio, and P. Strazzullo 2011. Potassium intake, stroke, and cardiovascular disease a meta-analysis of prospective studies. *Journal of the American College of Cardiology* 57:1210–19. doi:10.1016/j.jacc.2010.09.070.
- Daly, R. M., and P. R. Ebeling 2010. Is excess calcium harmful to health? *Nutrients* 2:505–22. doi:10.3390/nu2050505.
- DAS DE, S., S. Krishna, and A. Jethwa 2015. Iron status and its association with coronary heart disease: systematic review and meta-analysis of prospective studies. *Atherosclerosis* 238:296–303. doi:10.1016/j.atherosclerosis.2014.12.018.
- Del Gobbo, L. C., F. Imamura, J. H. Wu, M. C. De Oliveira Otto, S. E. Chiuve, and D. Mozaffarian 2013. Circulating and dietary magnesium and risk of cardiovascular disease: a systematic review and meta-analysis of prospective studies. *American Journal of Clinical Nutrition* 98:160–73. doi:10.3945/ajcn.112.053132.
- Du, S., C. Batis, H. Wang, B. Zhang, J. Zhang, and B. M. Popkin 2014. Understanding the patterns and trends of sodium intake, potassium intake, and sodium to potassium ratio and their effect on hypertension in China. *American Journal of Clinical Nutrition* 99:334–43. doi:10.3945/ajcn.113.059121.
- Dursun, I., and M. Sahin 2006. Difficulties in maintaining potassium homeostasis in patients with heart failure. *Clinical Cardiology* 29:388–92. doi:10.1002/clc.4960290904.
- Eaton, S. B., and M. Konner 1985a. Paleolithic nutrition. A consideration of its nature and current implications. *New England Journal of Medicine* 312:283–89. doi:10.1056/NEJM198501313120505.
- Eaton, S. B., and M. Konner 1985b. Paleolithic nutrition. A consideration of its nature and current implications. *New England Journal of Medicine* 312:283–89. doi:10.1056/NEJM198501313120505.
- Eckel, R. H., J. M. Jakicic, J. D. Ard, J. M. De Jesus, N. Houston Miller, V. S. Hubbard, I. M. Lee, A. H. Lichtenstein, C. M. Loria, B. E. Millen, et al. 2014. 2013 AHA/ACC guideline on lifestyle management to reduce cardiovascular risk: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *Circulation* 129:S76–99. doi:10.1161/01.cir.0000437740.48606.d1.
- Ekmekcioglu, C., I. Elmadfa, A. L. Meyer, and T. Moeslinger 2016. The role of dietary potassium in hypertension and diabetes. *Journal of Physiology and Biochemistry* 72:93–106. doi:10.1007/s13105-015-0449-1.
- EXCELLENCE. 2010. *NiFHaC Prevention of cardiovascular disease at the population level: NICE Public Health Guidance* [Online]. UK: National Health Service. Available: [www.nice.org.uk/guidance/PH25](http://www.nice.org.uk/guidance/PH25) [Accessed 3/02/2016 2016].
- Fang, X., K. Wang, D. Han, X. He, J. Wei, L. Zhao, M. U. Imam, Z. Ping, Y. Li, Y. Xu, J. Min, and F. Wang 2016. Dietary magnesium intake and the risk of cardiovascular disease, type 2 diabetes, and all-cause mortality: a dose–response meta-analysis of prospective cohort studies. *BMC Medicine* 14:210. doi:10.1186/s12916-016-0742-z.
- Farquhar, W. B., D. G. Edwards, C. T. Jurkovic, and W. S. Weintraub 2015. Dietary sodium and health: more than just blood pressure. *Journal of the American College of Cardiology* 65:1042–50. doi:10.1016/j.jacc.2014.12.039.
- Food And Nutrition Board, I. O. M. 1997. *Dietary Reference Intakes for calcium, phosphorus, magnesium, vitamin D, and fluoride* Washington, DC, National Academies Press.
- Fortmann, S. P., B. U. Burda, C. A. Senger, J. S. Lin, and E. P. Whitlock 2013. Vitamin and mineral supplements in the primary prevention of cardiovascular disease and cancer: An updated systematic evidence review for the U.S. Preventive Services Task Force. *Ann Intern Med* 159:824–34. doi:10.7326/0003-4819-159-12-201312170-00729.
- Geiger, H., and C. Wanner 2012. Magnesium in disease. *Clinical Kidney Journal* 5:i25–38. doi:10.1093/ndtplus/sfr165.
- Geleijnse, J. M., F. J. Kok, and D. E. Grobbee 2003. Blood pressure response to changes in sodium and potassium intake: a meta-regression analysis of randomised trials. *Journal of Human Hypertension* 17:471–80. doi:10.1038/sj.jhh.1001575.
- Graudal, N. A., T. Hubeck-Graudal, and G. Jurgens 2012. Effects of low-sodium diet vs. high-sodium diet on blood pressure, renin, aldosterone, catecholamines, cholesterol, and triglyceride (Cochrane Review). *American Journal of Hypertension* 25:1–15. doi:10.1038/ajh.2011.210.
- Greaney, J. L., J. J. Dupont, S. L. Lennon-Edwards, P. W. Sanders, D. G. Edwards, and W. B. Farquhar 2012. Dietary sodium loading impairs microvascular function independent of blood pressure in humans: role of oxidative stress. *Journal of Physiology* 590:5519–28. doi:10.1113/jphysiol.2012.236992.
- Grober, U., J. Schmidt, and K. Kisters 2015. Magnesium in Prevention and Therapy. *Nutrients* 7:8199–226. doi:10.3390/nu7095388.
- Guasch-Ferre, M., M. Bullo, R. Estruch, D. Corella, M. A. Martinez-Gonzalez, E. Ros, M. Covas, F. Aros, E. Gomez-Gracia, M. Fiol, et al. 2014. Dietary magnesium intake is inversely associated with mortality in adults at high cardiovascular disease risk. *J Nutr* 144:55–60. doi:10.3945/jn.113.183012.
- Hatton, D. C., Q. Yue, and D. A. McCarron 1995. Mechanisms of calcium's effects on blood pressure. *Seminars in Nephrology* 15:593–602.
- He, F. J., J. Li, and G. A. Macgregor 2013. Effect of longer term modest salt reduction on blood pressure: Cochrane systematic review and meta-analysis of randomised trials. *BMJ* 346:f1325. doi:10.1136/bmj.f1325.
- He, F. J., and G. A. Macgregor 2011. Salt reduction lowers cardiovascular risk: meta-analysis of outcome trials. *Lancet* 378:380–82. doi:10.1016/S0140-6736(11)61174-4.
- Jablonski, K. L., P. E. Gates, G. L. Pierce, and D. R. Seals 2009. Low dietary sodium intake is associated with enhanced vascular endothelial function in middle-aged and older adults with elevated systolic blood pressure. *Therapeutic Advances in Cardiovascular Disease* 3:347–56. doi:10.1177/1753944709345790.
- Joris, P. J., J. Plat, S. J. Bakker, and R. P. Mensink 2016. Long-term magnesium supplementation improves arterial stiffness in overweight and obese adults: results of a randomized, double-blind, placebo-controlled intervention trial. *American Journal of Clinical Nutrition* 103:1260–66. doi:10.3945/ajcn.116.131466.
- Khan, A. M., L. Sullivan, E. McCabe, D. Levy, R. S. Vasan, and T. J. Wang 2010. Lack of association between serum magnesium and the risks of hypertension and cardiovascular disease. *American Heart Journal* 160:715–20. doi:10.1016/j.ahj.2010.06.036.
- Khan, B., C. A. Nowson, R. M. Daly, D. R. English, A. M. Hodge, G. G. Giles, and P. R. Ebeling 2015. Higher Dietary Calcium Intakes Are Associated With Reduced Risks of Fractures, Cardiovascular Events, and Mortality: A Prospective Cohort Study of Older Men and Women. *Journal of Bone and Mineral Research* 30:1758–66. doi:10.1002/jbmr.2515.
- Khaw, K. T., S. Bingham, A. Welch, R. Luben, E. O'Brien, N. Wareham, and N. Day 2004. Blood pressure and urinary sodium in men and women: the Norfolk Cohort of the European Prospective Investigation into Cancer (EPIC-Norfolk). *American Journal of Clinical Nutrition* 80:1397–403. doi:10.1093/ajcn/80.5.1397.
- Kolte, D., K. Vijayaraghavan, S. Khera, D. A. Sica, and W. H. Frishman 2014. Role of magnesium in cardiovascular diseases. *Cardiology in Review* 22:182–92. doi:10.1097/CRD.0000000000000003.
- Lerchl, K., N. Rakova, A. Dahlmann, M. Rauh, U. Goller, M. Basner, D. F. Dinges, L. Beck, A. Agureev, I. Larina, et al. 2015. Agreement between 24-hour salt ingestion and sodium excretion in a controlled environment. *Hypertension* 66:850–57. doi:10.1161/HYPERTENSIONAHA.115.05851.
- Lewis, J. R., S. Radavelli-Bagatini, L. Rejnmark, J. S. Chen, J. M. Simpson, J. M. Lappe, L. Mosekilde, R. L. Prentice, and R. L. Prince 2015. The effects of calcium supplementation on verified coronary heart disease hospitalization and death in postmenopausal women: a collaborative meta-analysis of randomized controlled trials. *Journal of Bone and Mineral Research* 30:165–75. doi:10.1002/jbmr.2311.

- Lorenz, J. N., H. Weihprecht, J. Schnermann, O. Skott, and J. P. Briggs 1991. Renin release from isolated juxtaglomerular apparatus depends on macula densa chloride transport. *American Journal of Physiology* 260:F486–93.
- Lutsey, P. L., and E. D. Michos 2013. Vitamin D, calcium, and atherosclerotic risk: evidence from serum levels and supplementation studies. *Current Atherosclerosis Reports* 15:293. doi:10.1007/s11883-012-0293-5.
- Mccallum, L., S. Lip, and S. Padmanabhan 2015. The hidden hand of chloride in hypertension. *Pflügers Archiv* 467:595–603. doi:10.1007/s00424-015-1690-8.
- Mccarron, D. A., C. D. Morris, and C. Cole 1982. Dietary calcium in human hypertension. *Science* 217:267–69. doi:10.1126/science.7089566.
- Mccarron, D. A., and M. E. Reusser 1999. Finding consensus in the dietary calcium-blood pressure debate. *Journal of the American College of Nutrition* 18:398s–405s. doi:10.1080/07315724.1999.10718904.
- Mente, A., M. J. O'donnell, S. Rangarajan, M. J. McQueen, P. Poirier, A. Wielgosz, H. Morrison, W. Li, X. Wang, C. Di, et al. 2014. Association of urinary sodium and potassium excretion with blood pressure. *New England Journal of Medicine* 371:601–11. doi:10.1056/NEJMoa1311989.
- Michaelsson, K., H. Melhus, E. Warensjö Lemming, A. Wolk, and L. Byberg 2013. Long term calcium intake and rates of all cause and cardiovascular mortality: community based prospective longitudinal cohort study. *BMJ* 346:f228. doi:10.1136/bmj.f228.
- Mohammadifard, N., K. Humphries, C. Gotay, G. Mena-Sánchez, J. Salas-Salvador, A. Esmailzadeh, A. Ignaszewski, and N. Sarrafzadegan 2017. Trace minerals intake: risks and benefits for cardiovascular health. *Critical Reviews in Food Science and Nutrition*, Inpress. doi:10.1080/10408398.2017.1406332.
- Mohammadifard, N., A. Khaledifar, A. Khosravi, F. Nouri, A. Pourmoghadasi, A. Feizi, A. Esmailzadeh, and N. Sarrafzadegan 2017. Dietary sodium and potassium intake and their association with blood pressure in a non-hypertensive Iranian adult population: Isfahan salt study. *Nutr Diet* 74:275–82. doi:10.1111/1747-0080.12304.
- Moyer, V. A., and U. S. P. S.T. Force 2014. Vitamin, mineral, and multivitamin supplements for the primary prevention of cardiovascular disease and cancer: U.S. Preventive services Task Force recommendation statement. *Annals of Internal Medicine* 160:558–64. doi:10.7326/M14-0198.
- Mundy, G. R., and T. A. Guise 1999. Hormonal control of calcium homeostasis. *Clinical Chemistry* 45:1347–52.
- Nielsen, F. H. 2014. Effects of magnesium depletion on inflammation in chronic disease. *Current Opinion in Clinical Nutrition and Metabolic Care* 17:525–30. doi:10.1097/MCO.0000000000000093.
- O'donnell, M., A. Mente, S. Rangarajan, M. J. McQueen, X. Wang, L. Liu, H. Yan, S. F. Lee, P. Mony, A. Devanath, et al. 2014a. Urinary sodium and potassium excretion, mortality, and cardiovascular events. *New England Journal of Medicine* 371:612–23. doi:10.1056/NEJMoa1311889.
- O'donnell, M., A. Mente, and S. Yusuf 2014b. Sodium and cardiovascular disease. *New England Journal of Medicine* 371:2137–38.
- O'donnell, M., A. Mente, and S. Yusuf 2015. Sodium intake and cardiovascular health. *Circulation Research* 116:1046–57. doi:10.1161/CIRCRESAHA.116.303771.
- Oberleithner, H. 2012. Two barriers for sodium in vascular endothelium? *Annals of Medicine* 44 (Suppl 1):S143–8. doi:10.3109/07853890.2011.653397.
- Otten, J. J., J. P. Hellwig, and L. D. E. Meyers 2006. *Dietary Reference Intakes: The Essential Guide to Nutrient Requirements* Washington, DC, National Academies Press.
- Palmer, B. F., and D. J. Clegg 2016. Achieving the Benefits of a High-Potassium, Paleolithic Diet, Without the Toxicity. *Mayo Clinic Proceedings* 91:496–508. doi:10.1016/j.mayocp.2016.01.012.
- Pham-Huy, L. A., H. He, and C. Pham-Huy 2008. Free radicals, antioxidants in disease and health. *International Journal of Biomedical Science* 4:89–96.
- Posadas-Sanchez, R., C. Posadas-Romero, G. Cardoso-Saldana, G. Vargas-Alarcon, M. T. Villarreal-Molina, N. Perez-Hernandez, J. M. Rodriguez-Perez, A. Medina-Urrutia, E. Jorge-Galarza, J. G. Juarez-Rojas, et al. 2016. Serum magnesium is inversely associated with coronary artery calcification in the Genetics of Atherosclerotic Disease (GEA) study. *Nutrition Journal* 15:22. doi:10.1186/s12937-016-0143-3.
- Powles, J., S. Fahimi, R. Micha, S. Khatibzadeh, P. Shi, M. Ezzati, R. E. Engell, S. S. Lim, G. Danaei, D. Mozaffarian, et al. 2013. Global, regional and national sodium intakes in 1990 and 2010: a systematic analysis of 24 h urinary sodium excretion and dietary surveys worldwide. *BMJ Open* 3:e003733. doi:10.1136/bmjopen-2013-003733.
- Qu, X., F. Jin, Y. Hao, H. Li, T. Tang, H. Wang, W. Yan, and K. Dai 2013. Magnesium and the risk of cardiovascular events: a meta-analysis of prospective cohort studies. *PLoS One* 8:e57720. doi:10.1371/journal.pone.0057720.
- Reid, I. R. 2013. Cardiovascular effects of calcium supplements. *Nutrients* 5:2522–29. doi:10.3390/nu5072522.
- Reid, I. R., M. J. Bolland, and A. Grey 2010. Does calcium supplementation increase cardiovascular risk? *Clinical Endocrinology (Oxford)* 73:689–95. doi:10.1111/j.1365-2265.2010.03792.x.
- Reid, I. R., G. D. Gamble, and M. J. Bolland 2016. Circulating calcium concentrations, vascular disease and mortality: a systematic review. *Journal of Internal Medicine* 279:524–40. doi:10.1111/joim.12464.
- Rorije, N., R. O. Engberink, B. Van Den Born, H. Verberne, and L. Vogt 2015. 7D.06: EFFECTS OF AN ACUTE AND CHRONIC SALT LOAD ON MICROVASCULAR PERMEABILITY IN HEALTHY SUBJECTS. *Journal of Hypertension* 33 (Suppl 1):e101. doi:10.1097/01.hjh.0000467623.14967.fc.
- Ross, A. C., J. E. Manson, S. A. Abrams, J. F. Aloia, P. M. Brannon, S. K. Clinton, R. A. Durazo-Arvizu, J. C. Gallagher, R. L. Gallo, G. Jones, et al. 2011. The 2011 Dietary Reference Intakes for Calcium and Vitamin D: what dietetics practitioners need to know. *Journal of The American Dietetic Association* 111:524–27. doi:10.1016/j.jada.2011.01.004.
- Rusinaru, D., C. Tribouilloy, C. Berry, A. M. Richards, G. A. Whalley, N. Earle, K. K. Poppe, M. Guazzi, S. M. Macin, M. Komajda, and R. N. Doughty 2012. Relationship of serum sodium concentration to mortality in a wide spectrum of heart failure patients with preserved and with reduced ejection fraction: an individual patient data meta-analysis(dagger): Meta-Analysis Global Group in Chronic heart failure (MAGGIC). *European Journal of Heart Failure* 14:1139–46. doi:10.1093/eurjhf/hfs099.
- Salah, K., Y. M. Pinto, L. W. Eurlings, M. Metra, S. Stienen, C. Lombardi, J. G. Tijssen, and W. E. Kok 2015. Serum potassium decline during hospitalization for acute decompensated heart failure is a predictor of 6-month mortality, independent of N-terminal pro-B-type natriuretic peptide levels: An individual patient data analysis. *American Heart Journal* 170:531–42 e1. doi:10.1016/j.ahj.2015.06.003.
- Saulnier, P. J., E. Gand, and S. Hadjadj 2014. Sodium and cardiovascular disease. *New England Journal of Medicine* 371:2135–36.
- Schmidlin, O., A. F. Sebastian, and R. C. Morris JR. 2007. What initiates the pressor effect of salt in salt-sensitive humans? Observations in normotensive blacks. *Hypertension* 49:1032–39. doi:10.1161/HYPERTENSIONAHA.106.084640.
- Seth, A., Y. Mossavar-Rahmani, V. Kamensky, B. Silver, K. Lakshminarayan, R. Prentice, L. Van Horn, and S. Wassertheil-Smoller 2014. Potassium intake and risk of stroke in women with hypertension and nonhypertension in the Women's Health Initiative. *Stroke* 45:2874–80. doi:10.1161/STROKEAHA.114.006046.
- Sharma, S., K. Mcfann, M. Chonchol, and J. Kendrick 2014. Dietary sodium and potassium intake is not associated with elevated blood pressure in US adults with no prior history of hypertension. *The Journal of Clinical Hypertension (Greenwich)* 16:418–23. doi:10.1111/jch.12312.
- Shin, C. S., and K. M. Kim 2015. Calcium, Is It Better to Have Less?—Global Health Perspectives. *Journal of Cellular Biochemistry* 116:1513–21. doi:10.1002/jcb.25119.
- Sica, D. A., A. D. Struthers, W. C. Cushman, M. Wood, J. S. Banas JR., and M. Epstein 2002. Importance of potassium in cardiovascular disease. *The Journal of Clinical Hypertension (Greenwich)* 4:198–206. doi:10.1111/j.1524-6175.2002.01728.x.
- Simon, A., J. Gariepy, G. Chironi, J. L. Megnien, and J. Levenson 2002. Intima-media thickness: a new tool for diagnosis and treatment of cardiovascular risk. *Journal of Hypertension* 20:159–69. doi:10.1097/00004872-200202000-00001.



- Singh, R. B., N. K. Singh, P. J. Mehta, and S. S. Rastogi **1987**. Does calcium aggravate and cause hypertension? *Acta Cardiologica* 42:445–67.
- Smyth, A., M. O'donnell, A. Mente, and S. Yusuf **2015**. Dietary sodium and cardiovascular disease. *Current Hypertension Reports* 17:559. doi:10.1007/s11906-015-0559-8.
- Song, Y., H. D. Sesso, J. E. Manson, N. R. Cook, J. E. Buring, and S. Liu **2006**. Dietary magnesium intake and risk of incident hypertension among middle-aged and older US women in a 10-year follow-up study. *American Journal of Cardiology* 98:1616–21. doi:10.1016/j.amjcard.2006.07.040.
- Taylor, R. S., K. E. Ashton, T. Moxham, L. Hooper, and S. Ebrahim **2011**. Reduced dietary salt for the prevention of cardiovascular disease: a meta-analysis of randomized controlled trials (Cochrane review). *American Journal of Hypertension* 24:843–53. doi:10.1038/ajh.2011.115.
- Thiele, I., J. Linseisen, C. Meisinger, S. Schwab, C. Huth, A. Peters, S. Perz, T. Meitinger, F. Kronenberg, C. Lamina, et al. **2015**. Associations between calcium and vitamin D supplement use as well as their serum concentrations and subclinical cardiovascular disease phenotypes. *Atherosclerosis* 241:743–51. doi:10.1016/j.atherosclerosis.2015.06.032.
- Tomat, A. L., L. Costa Mde, and C. T. Arranz **2011**. Zinc restriction during different periods of life: influence in renal and cardiovascular diseases. *Nutrition* 27:392–98. doi:10.1016/j.nut.2010.09.010.
- Umesawa, M., H. Iso, C. Date, A. Yamamoto, H. Toyoshima, Y. Watanabe, S. Kikuchi, A. Koizumi, T. Kondo, Y. Inaba, N. Tanabe, A. Tamakoshi, and J. S. Group **2008**. Relations between dietary sodium and potassium intakes and mortality from cardiovascular disease: the Japan Collaborative Cohort Study for Evaluation of Cancer Risks. *American Journal of Clinical Nutrition* 88:195–202. doi:10.1093/ajcn/88.1.195.
- Vaidya, A., J. M. Brown, and J. S. Williams **2015**. The Renin-Angiotensin-Aldosterone System and Calcium-Regulatory Hormones. *Journal of Human Hypertension* 29:515–21. doi:10.1038/jhh.2014.125.
- Vaskonen, T. **2003**. Dietary minerals and modification of cardiovascular risk factors. *The Journal of Nutritional Biochemistry* 14:492–506. doi:10.1016/S0955-2863(03)00074-3.
- Verma, S., M. Gupta, D. T. Holmes, L. Xu, H. Teoh, S. Gupta, S. Yusuf, and E. M. Lonn **2011**. Plasma renin activity predicts cardiovascular mortality in the Heart Outcomes Prevention Evaluation (HOPE) study. *European Heart Journal* 32:2135–42. doi:10.1093/eurheartj/ehr066.
- Volpe, S. L. **2013**. Magnesium in disease prevention and overall health. *Advances in Nutrition* 4:378s–83s. doi:10.3945/an.112.003483.
- Waksman, R., and A. E. E. D. Ajani **2009**. *Appendix, in Pharmacology in the Catheterization Laboratory* Oxford, UK, Wiley-Blackwell.
- Wang, L., J. E. Manson, and H. D. Sesso **2012**. Calcium intake and risk of cardiovascular disease: a review of prospective studies and randomized clinical trials. *American Journal of Cardiovascular Drugs* 12:105–16. doi:10.2165/11595400-000000000-00000.
- Wang, X., H. Chen, Y. Ouyang, J. Liu, G. Zhao, W. Bao, and M. Yan **2014**. Dietary calcium intake and mortality risk from cardiovascular disease and all causes: a meta-analysis of prospective cohort studies. *BMC Medicine* 12:158. doi:10.1186/s12916-014-0158-6.
- Welch, A. A., H. Fransen, M. Jenab, M. C. Boutron-Ruault, R. Tumino, C. Agnoli, U. Ericson, I. Johansson, P. Ferrari, D. Engeset, et al. **2009**. Variation in intakes of calcium, phosphorus, magnesium, iron and potassium in 10 countries in the European Prospective Investigation into Cancer and Nutrition study. *European Journal of Clinical Nutrition* 63 (Suppl 4):S101–21. doi:10.1038/ejcn.2009.77.
- Whelton, P. K. **2014**. Sodium, potassium, blood pressure, and cardiovascular disease in humans. *Current Hypertension Reports* 16:465. doi:10.1007/s11906-014-0465-5.
- Whelton, P. K. **2015**. Dietary sodium intake: scientific basis for public policy. *Blood Purification* 39:16–20. doi:10.1159/000368975.
- Whelton, P. K., L. J. Appel, R. L. Sacco, C. A. Anderson, E. M. Antman, N. Campbell, S. B. Dunbar, E. D. Frohlich, J. E. Hall, M. Jessup, et al. **2012**. Sodium, blood pressure, and cardiovascular disease: further evidence supporting the American Heart Association sodium reduction recommendations. *Circulation* 126:2880–89. doi:10.1161/CIR.0b013e318279acbf.
- WHO **2012**. *Guideline: Potassium intake for adults and children*. Geneva: World Health Organization (WHO).
- WORLD HEALTH ORGANIZATION **2012**. *Guideline: Sodium Intake for Adults and Children*. Geneva: World Health Organization.
- Xi, L., Y. C. Hao, J. Liu, W. Wang, M. Wang, G. Q. Li, Y. Qi, F. Zhao, W. X. Xie, Y. Li, J. Y. Sun, J. Liu, L. P. Qin, and D. Zhao **2015**. Associations between serum potassium and sodium levels and risk of hypertension: a community-based cohort study. *Journal of Geriatric Cardiology* 12:119–26.