

Critical Reviews in Food Science and Nutrition

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/bfsn20>

Natural Occurrence, Analysis, and Prevention of Mycotoxins in Fruits and their Processed Products

Jinyi Yang^a, Jing Li^a, Yueming Jiang^a, Xuewu Duan^a, Hongxia Qu^a, Bao Yang^a, Feng Chen^b & Dharini Sivakumar^c

^a Key Laboratory of Plant Resource Conservation and Sustainable Utilization, South China Botanical Garden, Chinese Academy of Sciences, People's Republic of China

^b Department of Food Science and Human Nutrition, Clemson University, Clemson, South Carolina, USA

^c Department of Crop Sciences, Tshwane University of Technology, Pretoria (West Campus), Pretoria, South Africa

Accepted author version posted online: 24 Feb 2012. Published online: 04 Nov 2013.

To cite this article: Jinyi Yang, Jing Li, Yueming Jiang, Xuewu Duan, Hongxia Qu, Bao Yang, Feng Chen & Dharini Sivakumar (2014) Natural Occurrence, Analysis, and Prevention of Mycotoxins in Fruits and their Processed Products, Critical Reviews in Food Science and Nutrition, 54:1, 64-83, DOI: [10.1080/10408398.2011.569860](https://doi.org/10.1080/10408398.2011.569860)

To link to this article: <http://dx.doi.org/10.1080/10408398.2011.569860>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

Natural Occurrence, Analysis, and Prevention of Mycotoxins in Fruits and their Processed Products

JINYI YANG,¹ JING LI,¹ YUEMING JIANG,¹ XUEWU DUAN,¹ HONGXIA QU,¹
BAO YANG,¹ FENG CHEN,² and DHARINI SIVAKUMAR³

¹Key Laboratory of Plant Resource Conservation and Sustainable Utilization, South China Botanical Garden, Chinese Academy of Sciences, People's Republic of China

²Department of Food Science and Human Nutrition, Clemson University, Clemson, South Carolina, USA

³Department of Crop Sciences, Tshwane University of Technology, Pretoria (West Campus), Pretoria, South Africa

Mycotoxins are small toxic chemical products formed as the secondary metabolites by fungi that readily contaminate foods with toxins in the field or after harvest. The presence of mycotoxins, such as aflatoxins, ochratoxin A, and patulin, in fruits and their processed products is of high concern for human health due to their properties to induce severe acute and chronic toxicity at low-dose levels. Currently, a broad range of detection techniques used for practical analysis and detection of a wide spectrum of mycotoxins are available. Many analytical methods have been developed for the determination of each group of these mycotoxins in different food matrices, but new methods are still required to achieve higher sensitivity and address other challenges that are posed by these mycotoxins. Effective technologies are needed to reduce or even eliminate the presence of the mycotoxins in fruits and their processed products. Preventive measures aimed at the inhibition of mycotoxin formation in fruits and their processed products are the most effective approach. Detoxification of mycotoxins by different physical, chemical, and biological methods are less effective and sometimes restricted because of concerns of safety, possible losses in nutritional quality of the treated commodities and cost implications. This article reviewed the available information on the major mycotoxins found in foods and feeds, with an emphasis of fruits and their processed products, and the analytical methods used for their determination. Based on the current knowledge, the major strategies to prevent or even eliminate the presence of the mycotoxins in fruits and their processed products were proposed.

Keywords: Metabolites, fungus, control, contamination, detoxification, handling

INTRODUCTION

Mycotoxins are toxic compounds produced by fungi, mostly by saprophytic moulds growing on a variety of foodstuffs (Logrieco, 2010) and they are a group of structurally diverse secondary metabolites (Figure 1) (Turner et al., 2009). However, the reason for the production of mycotoxins is not yet known as they are neither necessary for growth nor the development of the fungi (Fox and Howlett, 2008). Because mycotoxins weaken the receiving host, the fungus may use them as a strategy to better meet the environment for further fungal proliferation. Mycotoxin production can occur depending largely on

the species of fungi. Fungi can infect agricultural crops during crop growth, harvest, storage, or processing. Some molds are capable of producing more than 1 mycotoxin and some mycotoxins are produced by more than 1 fungal species (Sakuda and Kimura, 2010). For example, ochratoxin (OTA) is produced by some species such as *Aspergillus ochraceus* mainly in tropical regions and by *Penicillium verrucosum* a common storage fungus in temperate areas, and in some cases, 1 species can form more than 1 mycotoxin. The mycotoxins include mainly aflatoxins, citrinin, cyclopiazonic acid, fumonisins, moniliformin, ochratoxins, patulin, sterigmatocystin, and trichothecenes. Aflatoxins, ochratoxins, trichothecenes, zearalenone, fumonisins, tremorgenic toxins, and ergot alkaloids are the mycotoxins of the great agro-economic importance. Mycotoxins have adverse effects on humans, animals, and crops that result in illnesses and economic losses (Zain, 2010). The economic impact of mycotoxins include loss of human and animal life, increased health

Address correspondence to Yueming Jiang, Key Laboratory of Plant Resource Conservation and Sustainable Utilization, South China Botanical Garden, Chinese Academy of Sciences, People's Republic of China. E-mail: ymjjiang@scbg.ac.cn




Figure 1 Examples of common mycotoxins demonstrating structural diversity of these compounds. A, citrinin; B, aflatoxin B1; C, patulin; D, deoxynivalenol; and E, fumonsin B1 (Turner et al., 2009).

care and veterinary care costs, reduced livestock production, disposal of contaminated foods and feeds, and investment in research and applications to reduce severity of the mycotoxin problem (Chassy, 2010).

The most commonly found in fruits and their processed products are aflatoxins, ochratoxin A, patulin, and *Alternaria* toxins (Drusch and Ragab, 2003; Barkai-Golan and Paster, 2008; Moss, 2008). These mycotoxins can contaminate food stuffs or human foods. The ingestion of these contaminated materials may be pathogenic in humans as they may lead to serious health problems, such as liver, kidney, or nervous system damage, immunosuppression, biphasic cellular response, and carcinogenesis (Bennett and Klich, 2003; Diesing et al., 2010; Maresca and Fantini, 2010).

Contaminations of mycotoxins were reported in a number of agricultural commodities, foods, feed stuffs as well as bio-

logical fluids at geographically diverse locations (Phillips et al., 1980; Gimeno and Martins, 1983; Heber et al., 2001; CAST, 2003; Marín et al., 2010; Meister, 2004; Soriano and Dragacci, 2004; Dashti and AL-Hamli, 2010; Yuan et al., 2010). Contamination can occur preharvest or postharvest, for example, deoxynivalenol and T-2 toxin produced by *Fusarium* (preharvest), or OTA by *Aspergillus* and *Penicillium* and aflatoxin (AFT) by *Aspergillus* (postharvest) (EMAN, 2003). A lot of mycotoxins have been identified in fruits and their processed products (Drusch and Aumann, 2005; Luttfullah and Hussain, 2010).

The maximum levels have been set for some mycotoxins in food and feed to protect animal and public health, as described in detail by Zain (2010). The worldwide contamination of foods and feed with mycotoxins is a significant problem. Due to the widespread nature of fungi in the natural environment, mycotoxins are considered unavoidable contaminants in food




Figure 2 Chemical structure of aflatoxin B (AFB₁ and AFB₂), aflatoxin G (AFG₁ and AFG₂), and aflatoxin M (AFM₁ and AFM₂) (Zain, 2010).

and feed. On a worldwide basis, at least 99 countries established mycotoxin regulations for food and/or feed in 2003. Although efforts have continued internationally to set guidelines to control mycotoxins, practical measures have not been adequately implemented. Thus, mycotoxin management will meet an international challenge to ensure global food safety (López-García, 2010). It is important to develop rapid, sensitive, and reproducible assays to detect the presence of mycotoxins. At the same time, considerable efforts have been made to reduce or even eliminate the presence of the mycotoxins in foods and feeds. This article reviewed the available information on the major mycotoxins found in foods and feeds, with an emphasis of fruits and their processed products, and the analytical methods used for their determination. Based on the current knowledge, the major strategies to prevent or even eliminate the presence of the mycotoxins in fruits and their processed products were proposed.

TOXICITY OF MYCOTOXINS

Contamination of crops by fungal action has been noted for over 2 millennia. The improvements in food safety in developed countries have eliminated acute human mycotoxicosis, but such outbreaks still occur in rural communities in the developing world where aflatoxins, fumonisins, deoxynivalenol, ochratoxin, and zearalenone present in foods have been involved in the deaths or acute diseases reported (Shephard, 2008; Zain, 2010). The current literature on mycotoxin-related human diseases clearly reveals a linkage between ingesting mycotoxin-contaminated food and illness, especially hepatic, gastrointestinal, carcinogenic, and teratogenic diseases (Fung and Clark, 2004; Shephard, 2008). Based on the effects on human and animal health, aflatoxin, fumonisin, trichothecenes, ochratoxin, zearalenone, and patulin are recognized as the most important

food mycotoxins. The most studied mycotoxins in fruits and their processed products have been AFT, OTA, and patulin. Detailed information on the toxicity of mycotoxins is given in 2 reports reviewed by Turner et al. (2009) and Fernández-Cruza et al. (2010).

MAJOR MYCOTOXINS

Aflatoxins are a group of closely related metabolites produced by *Aspergillus flavus* and *Aspergillus parasiticus*. They are difuranocoumarin derivatives and the main components of this group are aflatoxin B₁, B₂, G₁, and G₂, based on their fluorescence under ultraviolet (UV) light (blue or green) and their relative chromatographic mobility (Figure 2) (Turner et al., 2009). They were first detected and characterized in the 1960s (Asao et al., 1965) and have been found in a variety of agricultural and food products. The LD₅₀ of AFB₁ ranges from 0.3 to 18 mg/kg depending on the animal species and routes of administration. Aflatoxins are classified by the International Agency for Research on Cancer as being carcinogenic to humans (Group 1) (World Health Organization, 1998; Fung and Clark, 2004; Shephard, 2008).

OTA is a phenylalanine-dihydroisocoumarin derivative (Figure 3) (Zain, 2010). The LD₅₀ of OTA ranges from 0.5 mg/kg for dogs to over 50 mg/kg for mice and the toxicological profile of OTA includes teratogenesis, nephrotoxicity, and immunotoxicity (Soleas et al., 2001; Drusch and Ragab, 2003). Animal experiments have implicated cytochrome P450-related reactions and DNA adducts generation as possible mechanisms for the formation of renal tumours.

Patulin (PAT) is an unsaturated heterocyclic lactone. Various acute and chronic effects have been attributed to PAT (Drusch and Ragab, 2003; Fung and Clark, 2004; Sant'Ana et al., 2008).




Figure 3 Chemical structure of ochratoxin A (Zain, 2010).

The LD₅₀ of patulin ranges from 15 to 25 mg/kg and varies with animal species and route of exposure. The acute symptoms include lung and brain edema, liver, spleen, and kidney damage and toxicity to the immune system while the chronic symptoms include genotoxic, neurotoxic, immunotoxic, immunosuppressive, and teratogenic effects. At the cellular level, some examples of these effects by PAT are plasmatic membrane rupture and inhibitions of DNA, RNA, and protein syntheses (Drusch and Ragab, 2003; Fung and Clark, 2004; Sant'Ana et al., 2008; Zhou et al., 2010).

NATURAL OCCURRENCE OF MYCOTOXINS IN FRUITS AND THEIR PROCESSED PRODUCTS

Alternaria fungi are commonly parasitic on plants and may cause spoilage of fruits and vegetables during transport and storage. *Alternaria alternata* produces a number of mycotoxins including the dibenzo- α -pyrones alternariol, alternariol monomethyl ether, and altenuene, altertoxin I and II, and tenuazonic acid—a tetramic acid. Culture extracts of *A. alternata* are mutagenic in various microbial and cell systems, and carcinogenic in rats. Ochratoxin A was originally isolated from *A. ochraceus* in 1965. Other *Aspergillus* species are also capable of producing OTA. *P. verrucosum* is the best known *Penicillium* species to produce OTA. The contamination of foods with OTA in cool climates is usually caused by *P. verrucosum*, whereas the occurrence of OTA in foods in warmer and tropical climates is associated with *A. ochraceus*. Several species of *Penicillium* and *Aspergillus* can also produce PAT. The most important producer of PAT is the apple-rotting fungus *Penicillium expansum*. However, a few literatures are available for the aflatoxins (except in dried figs) and the *Alternaria* toxins in fruits and their products, although the ubiquity and toxicity of the latter are well known. Very recently, some evidence for the presence of fumonisins B1 and B2 in fruits has been reported. Fumonisin B2 has been identified in visibly moldy dried figs (Senyuva and Gilbert, 2008). A high incidence of fumonisin B1 was found in dried figs in Turkey (Karbancioglu-Güler and Heperkan, 2008) while Logrieco et al. (2009) reported fumonisin B2 production by *Aspergillus niger* from grapes and natural occurrence in must. Table 1 gives the occurrence of mycotoxins in fruits and their processed products.

The extent of fungal growth and subsequent possible mycotoxin contamination depends largely on endogenous and exogenous factors (van der Westhuizen et al., 2010). It has been demonstrated that environmental stress conditions such as insect infestation, drought, cultivar susceptibility, mechanical damage, nutritional deficiencies, and unseasonable temperature, rainfall, or humidity can promote mycotoxin production in growing crops (Varga et al., 2010). In fact, changes in farming practices in the past few decades may result in increasing stress on plants and, therefore, enhance fungal invasion and mycotoxin contamination. Various stages of the manufacturing process of apple juice are capable of reducing the amount on the final products on a certain extent, and the incidence of the mycotoxins throughout the world confirms their stabilities (Sant'Ana et al., 2008). The careful selection and proper storage of fruits are the most important implement reducing mycotoxins (Swanson, 1987; Drusch and Ragab, 2003).

THERMAL STABILITY OF MYCOTOXINS IN FRUITS AND THEIR PROCESSED PRODUCTS

Most mycotoxins are chemically stable during storage and processing, even when cooked at quite high temperatures of more than 90°C (Swanson, 1987). Dibenzo- α -pyrones alternariol (AOH) and alternariol monomethyl ether are stable on heating at 100°C in sunflower flour (Combina et al., 1999). Both mycotoxins are very stable in spiked apple juice at room temperature for up to 5 weeks and at 80°C after 20 min. Thus, mycotoxins are difficult to be degraded or decontaminated in fruits and their processed products.

EXTRACTION, SEPARATION, AND/OR DETERMINATION OF MYCOTOXINS

The fact that most mycotoxins are toxic in very low concentrations requires sensitive and reliable methods for their detection. Due to the varied structures and chemical functions of these compounds, it is impossible to use one standard technique to detect all mycotoxins, as each will require a different method (Turner et al., 2009). Practical requirements for high-sensitivity detection and the need for a special laboratory setting create challenges for routine analysis. Depending on the physical and chemical properties of mycotoxins, more procedures have been developed around existing analytical techniques, which offer flexible and broad-based methods of detecting these compounds. The application of simpler, cheaper, and more effective solutions for the detection of mycotoxins is increasingly being required, due to their perceived importance, based around their toxicity and requirements of legislation for limits on amounts in foods. A successful detection method should be robust and sensitive, with a high degree of flexibility, over a wide range of compounds. All techniques should be reproducible to a high level, and the results gained must be relevant and easy to analyze. For fieldwork, the system should also be rapid and portable. Many

Table 1 Occurrence of mycotoxins in fruits and their processed products

Commodities	Toxins	Concentration range	Reference
Apple juice	AF-B ₁ and AF-G ₁	20–30 µg/l	Abdel Sater et al. (2001)
Dried raisins	AFT	Max. 2–550 µg/kg	Saxena and Mehrotra (1990), Trucksess and Scott (2008)
Dried figs	AFT	Max. 10–325 µg/kg	Juan et al. (1991), Drusch and Ragab (2003)
Rotten mandarins	AOH	1000–5200 µg/kg	Logrieco et al. (1990)
Tangerine flavedo	AOH	2.5–17.4 µg/kg	Magnani et al. (2007)
Apple juice	AOH	0.04–2.40 µg/l	Lau et al. (2003)
Red grape juices	AOH	0.03–0.46 µg/l	Scott et al. (2006)
Peaches	OTA	0.21 µg/kg	Engelhardt et al. (1999)
Cherries	OTA	2.71 µg/kg	Engelhardt et al. (1999)
Strawberry	OTA	1.44 µg/kg	Engelhardt et al. (1999)
Apple	OTA	0.41 µg/kg	Engelhardt et al. (1999)
Grape juice	OTA	0.010–5.3 µg/l	Battilani et al. (2006)
Vinegar	OTA	0.22–6.4 µg/l	Battilani et al. (2006)
Raisins	OTA	Max 26–250 µg/kg	Drusch and Ragab (2003), Battilani et al. (2006), Trucksess and Scott (2008)
Dried figs	OTA	<0.12–6900 µg/kg	Drusch and Ragab (2003), Karbancioglu-Güler and Heperkan (2008)
Apples rotten areas	PAT	2–11,3000 µg/kg	Drusch and Ragab (2003)
Apple juice	PAT	0.5–1150 µg/l	Leggott and Shephard (2001), Cheraghali et al. (2005), Murillo-Arbizu et al. (2009)
Apple juice conc	PAT	4.6–467.4 µg/l	Harris et al. (2009)
Apple puree	PAT	3–39 µg/kg	Funes and Resnik (2009)

of the techniques described below have been combined to form protocols, which are most used in laboratories today.

Sample Pretreatment Methods

Accurate determination of a mycotoxin relies on correct extraction and clean-up methods except perhaps enzyme-linked immunosorbent assay (ELISA) which may not require clean-up (Chu, 1992; Şenyuva and Gilbert, 2010). The extraction method used to remove the mycotoxin from a biological matrix is dependent largely on the structure of the mycotoxin. Polar metabolites, such as the fumonisins, require the presence of water, aided by the presence of organic solvents (Shephard, 1998). Hydrophobic mycotoxins such as AFT rely on use of organic solvents (Holcomb et al., 1992). These can be direct extractions, or may be partitioned with other solvents, such as *n*-hexane for partial clean-up, to remove excess components of the biological matrix. The choice of an appropriate extraction solvent is also dependent on the matrix from which the extraction is required, as the inherent components of the mixture can affect it (Wilkes and Sutherland, 1998). The clean-up procedure used in a protocol is the most important step, as the purity of the sample affects the resolution of separation and sensitivity of the results (Scott, 1995). Trace amounts of a target molecule may be masked by interfering compounds, found not only in the matrix but in solvents used in the technique. Glassware should also be free of contamination, such as alkaline detergents, which can form salts with the compounds and then interfere detection or result in false results.

Liquid–liquid Extraction

Liquid–liquid extraction involves exploiting the different solubility of the mycotoxins in 2 immiscible phases, to extract the

target compounds into 1 solvent leaving the rest of the matrix in the other. For example, solvents such as hexane and cyclohexane are commonly used to remove nonpolar contaminants, for example, lipids and cholesterol. The procedure is effective for several mycotoxins and works well in small-scale preparations (Bauer and Gareis, 1987). Holcomb et al. (1996) reported supercritical fluid extraction (SFE) extraction of aflatoxins from corn. However, the method is time consuming and may cause loss of part of targets.

Solid Phase Extraction

The basic principle of solid phase extraction (SPE) technology is a variation of chromatographic techniques based on around small disposable cartridges packed with silica gel, or bonded phases which are called the stationary phase. The sample is loaded in one solvent, generally under reduced pressure, rinsed, where most of the contaminants are removed, and eluted in another solvent (EMAN, 2003; Kataoka et al., 2010). These cartridges have a high capacity for binding small molecules. These cartridges contain different bonding phases, ranging from silica gel, C-18 (octadecylsilane), florisil, phenyl, aminopropyl, ion-exchange materials, both anionic and cationic, to affinity materials such as immunoabsorbents and molecular imprinted polymers (Visconti and Pascale, 1998; Sharma and Marquez, 2001; Visconti et al., 2000; Zamboni et al., 2002; Muñoz et al., 2006; Giraudi et al., 2007; Katerere et al., 2008; Vatinno et al., 2008). Anion exchange bonded phases provide good clean-up of extracts (Sydenham et al., 1992). The SPE systems have many advantages, for example, they use considerably less solvent and are faster in operation. In addition to cleaning sample, they can also be used to preconcentrate the sample providing better detection results. SPE has found widespread use and is an integral part of many extraction and detection protocols.

Table 2 Examples of SPE methods used in the extraction and clean-up of mycotoxin samples

Toxin	SPE phase	Matrix	Reference
Alternariol	C-18-RP	Apple juice	Delgado et al. (1996)
Fumonisin	C-18-RP, SAX, Immunoaffinity	Various	Shephard (1998)
Patulin	Silica gel SPE column	Dried apple rings	Katerere et al. (2008)
Trichothecenes	C-18-RP	Grain	Radová et al. (1998)

Giraudi et al. (2007) described a preparation of a hexapeptide library by combinatorial synthesis and then identified a peptide which showed good affinity with OTA. In another study, a 2-dimensional extraction procedure employed SPE for the extraction of OTA (Maier et al., 2004). Furthermore, detection of OTA using solid phase micro extraction-coupled with liquid chromatography–mass spectrometry–mass spectrometry (SPME–LC–MS/MS) has been reported (Vatinno et al., 2008). High throughput was achieved by simultaneous preparation of up to 96 samples using multifiber SPME (solid phase micro extraction) device and multiwell plates. Katerere et al. (2008) reported the extraction of patulin using SPE and isocratic separation on high-performance liquid chromatography coupled with diode array detector (HPLC-DAD). Using this method, the accuracy and linearity at 10–50 ppb were obtained. Nowadays, SPE is by far the most popular technique used in routine analysis of mycotoxins. Table 2 presents some examples of SPE methods used in the extraction and clean-up of mycotoxin samples. Recently, solid-phase extraction using molecularly imprinted polymers for selective extraction of a mycotoxin has been developed and the method has exhibited a high selectivity and significant capacity of the molecularly imprinted polymers to extract OTA (Ali et al., 2010).

Silica gel is a very popular material, frequently used for SPE. The surface of silica particles is heterogeneous, with a variety of silanol groups which can bind target compounds through multiple electrostatic interactions. The addition of various functional groups can widen the application of this material. Normally, this is achieved by reacting an organosilane with a long chain aliphatic derivative (e.g., C-18 phase). Silica gel can be used directly or after modification for mycotoxin detection. More frequently, however, it is a hydrophobic phase which is used in environmental and food analyses of these toxins, performed in both polar and nonpolar solvents. Leitner et al. (2002) demonstrated the use of C-18 reverse phase materials in the extraction of OTA. They have shown that the use of cartridges made of this material combined with mass spectroscopy offers good results (detection at sub-ppb level) which are comparable to clean-up with OTA-specific immunosorbents.

Ion-exchange materials are often used in SPE to isolate and preconcentrate mycotoxins found in aqueous solutions. They utilize electrostatic interactions formed between the target molecule and charged groups bonded to the silica material. The samples are set to a pH value where both groups are charged, and filtered through the SPE cartridge. The bound material is removed by the addition of a strong ionic solution or by altering pH. Several types exist in both anionic and cationic phases.

Strong anion exchange (SAX) is the favored material for mycotoxin extraction. Pussemier et al. (2006) developed a sensitive protocol for OTA, citrinin (CIT), or deoxynivalenol (DON) detection and quantification using SAX columns in clean-up.

MycosepTM columns are made up of several absorbents specifically selected for recovery of individual mycotoxins, packed into a plastic tube, and used to remove the entire matrix leaving the desired compound in solution on top of the column (Akiyama et al., 2001; Mateo et al., 2002). This method is practical, portable, and quick with no additional rinsing steps required. However, columns are designed per analyte and, therefore, not useful in multitoxin analysis. A MycosepTM column specific for OTA has been developed, which works in tandem with other columns to ensure clean-up from a variety of matrices. With several companies working on development of newer columns for different mycotoxins, it is expected that MycosepTM columns would be used for more applications in future.

Antibody has been intensively investigated over the past few years, and has been shown to be of great value in the mycotoxin analysis (Zimmerli and Dick, 1995; Scudamore and MacDonald, 1998; Visconti and Pascale, 1998; Visconti et al., 2000; Mateo et al., 2002). They offer excellent recovery of analyte because of the specificity gained by using monoclonal or polyclonal antibodies. The main disadvantage in using these materials is high costs involved as each column can only be used once due to denaturation of antibodies. This has been taken into account with further development of sequential injection immunoassay (Garden and Strachan, 2001).

Separation and/or Determination Methods

Recent trends in sample separation include miniaturization, automation, high-throughput performance, and reduction in solvent consumption and operation time. There are several types of chromatographic methods available for mycotoxin analysis (Shephard, 1998; Valenta, 1998; Vaclavik et al., 2010), which will be introduced in more details in the following sections.

Thin Layer Chromatography

Traditionally, the most popular method used for mycotoxins analysis is thin-layer chromatography (TLC), which offers the ability to screen large numbers of samples economically. There are good reviews available on the applications of TLC for mycotoxin analysis (Betina, 1985, 1993; Scott, 1985, 1987;

Table 3 Examples of TLC protocols used for the detection of common mycotoxins

Toxin	Protocol	Mode of Detection	Matrix	Reference
Citrinin	Silica gel precoated G-25 HR TLC plates	Fluorescent	Apples and pears	Gimeno and Martins (1983), Martins et al. (2002)
Citrinin	Silica gel 60 TLC	Fluorescent	Fungi isolated from grapes	Abrunhosa et al. (2001)
Trichothecenes	Comparison with HPLC for detection	UV and fluorescent	Food	Tomlins et al. (1989)
OTA	Silica gel	Fluorescence densitometry	Rice	Abramson et al. (1999a)

Holcomb et al., 1992; Kok, 1994; Lin et al., 1998). The use of TLC analysis for mycotoxins is still popular for both quantitative and semiquantitative purposes (Table 3), due to its high throughput of samples, low operating cost, and ease of identification of target compounds with an aid of UV-V is spectral analysis. In this context, several methods have been developed to obtain the best results with each separate class of mycotoxin and both 1-dimensional and 2-dimensional analyses are used frequently (Gimeno, 1984; Eppley et al., 1986; Lin et al., 1998). However, one of the major requirements for TLC is that there is an inherent need for sample preparation, the type of clean-up protocol dependent on properties and the type of the mycotoxin in study. Silica gel layer seems to be one of the common layers used for TLC while octadecyl bonded ones seem to be the most widely used (Stahr et al., 1985; Edwards and Lillehoj, 1987; Frohlich et al., 1988). Phenyl nonpolar bonded (Bradburn et al., 1990), silanized (Tomlins et al., 1989), and impregnated with organic acid (Colinski and Rabarkiewicz-Szcesna, 1984) were also attempted to separate mycotoxins.

TLC has been reported to be used to assess mould spoilage of *C. sativa* nuts commercially sold in Canadian grocery stores (Overly et al., 2003). Three mycotoxigenic fungi dominating the samples were *P. crustosum*, *P. glabrum/spinulosum*, and *P. discolor*. A 1-dimensional TLC method, combined with immunoaffinity columns (IAC)-based clean-up, was used to determine AF in various food matrices (Stroka et al., 2000). Medina and Schwartz (1992) reported a TLC analysis for detection of estrone, estradiol, diethylstilbestrol, zearalanol, zearalanone, and zearalenol. TLC has been also used for the detection of diphenylindene sulphonyl and esters of TRC mycotoxins (Turner et al., 2009). While TLC is still used continuously, improvement of sample preparation, further development of more sensitive analytical system and improved automation in terms of enhancing its performance could also make TLC more popular tool in future.

High-Performance Liquid Chromatography

Modern analysis of mycotoxins relies heavily on HPLC in 2 aspects: (1) interactions between the HPLC column (adsorbents) and the mycotoxins and (2) detection methods such as UV or fluorescence which relies largely on the presence of a chromophore in the molecules. A number of toxins already have natural fluorescence (e.g., OTA, AFT, and CIT) and can be detected directly by HPLC-fluorescence detector (FD) (Elizalde-Gonzalez et al.,

1998). However, some mycotoxins, such as fumonisins produced by *Fusarium* genus, lack a suitable chromophore, so that their determination requires chemical derivatization (Shephard, 1998; EMAN, 2003). Some derivatizing agents, for example, *o*-phthaldialdehyde and 9-(fluorenylmethyl) chloroformate (Holcomb et al., 1993), can be employed in either pre- or postcolumn (Neely and Emerson, 1990; Kussak et al., 1995; Jimenez et al., 2000; Chiavaro et al., 2001). HPLC has been a standard for mycotoxin detection in food industry (Valenta, 1998). For example, reverse phase (RP)-HPLC coupled to sensitive FD was employed to analyze OTA formation (Hernández et al., 2006), which allowed estimation of OTA in 0.01 ng/ml concentration. In similar study, OTA contamination in dried figs, which was extracted by methanol and orthophosphoric acid followed by clean-up by an IAC, was investigated using HPLC-FD with the limits of detection (LOD) for OTA of 0.12 µg/kg (Karbancioglu-Güler and Heperkan, 2008). HPLC has also been employed for PAT detection in commercial pure apple juices and mixed apple juices marketed in Italy (Spadaro et al., 2007). There have been several other mycotoxins that have been determined using HPLC methods (Josephs et al., 1998; Ambrosino et al., 2004; Xu et al., 2006; González et al., 2008). AFT and cyclopiazonic acid were also detected using HPLC method in Brazilian peanuts (González et al., 2008). The main benefit of using HPLC along with the high quality of separation and low LOD provides the possibility to combine multiple detection systems (fluorescent, UV, and diode array) with this technology, allowing for multiple detections of compounds from one sample. For instance, an HPLC-FD has been used for simultaneous detection of *Fusarium* mycotoxins [zearalanone (ZON), nivalenol, and DON] in cereals (Tanaka et al., 1985). ZON in food has been determined using clean-up by either conventional SPE or IAC and detected by HPLC-MS with an atmospheric pressure chemical ionization interface (Tanaka et al., 1985; Rosenberg et al., 1998). Application of APCI-MS detection increased the sensitivity by a factor of ca. 50 in comparison with HPLC-FD. Due to the selectivity of MS detection, it also was possible to quantitatively determine ZON both in raw extracts without clean-up. Leitner et al. (2002) presented an excellent comparison of MS and FD for OTA analysis. Furthermore, a LC-MS/MS method for multimycotoxin determination was developed for 19 analytes and was applied for a semiquantitative screening of 87 moldy food samples from private households, including bread, fruits, vegetables, cheeses, nuts, and jam (Sulyok et al., 2010). Table 4 lists some examples of protocols using HPLC for common mycotoxin analyses. MS became the industry standard detection method allowing highly

Table 4 Examples of protocols using HPLC for common mycotoxins

Toxin	Protocol	Detection	Matrix	Reference
Aflatoxins	UV-HPLC	Spectrofluorimetric	Brazilian peanut kernels	González et al. (2008), Afsah-Hejri et al. (2010)
Aflatoxin B1 and OTA	RP- HPLC	Fluorescent	Gapes	Khoury et al. (2008)
Aflatoxin B1, citrinin, and OTA	Normal HPLC	Fluorescent	Various samples	Nguyen et al. (2007), Brera et al. (2010)
Aflatoxins, the aflatoxin metabolite M1, ochratoxin a	Ultra HPLC-MS	Orthogonal Z-spray-electrospray interface.	Food	Beltrán et al. (2010)
Beauvericin	HPLC followed by confirmation by LC-MS	Diode array detector	Maize	Ambrosino et al., (2004)
Citrinin	Normal-HPLC	Fluorescent	Corn	Abramson et al. (1999b)
Citrinin	Normal-HPLC	Fluorescent	Cereals	Meister (2004)
Citrinin	Many chromatographic techniques compared	Various detection methods	Various samples	Xu et al. (2006)
Ergosterol	HPLC compared with ELISA and TLC	Various detection methods	Various samples	Janardhana et al. (1999)
Fumisin	(HPLC-ELSD)	Evaporative laser scattering detection	Corn samples	Wang et al. (2008)
Patulin	Microemulsion electrokinetic chromatography for quantification	UV and fluorescent	Commercial apple juice	Murillo-Arbizu et al. (2008)
Patulin	Normal HPLC	UV diode array detector	Fruit juice	Spadaro et al. (2007), Moukas et al. (2008)
Penitrem A, chaetoglobosin A and C, emodin, and OTA	Normal HPLC	Diode array detection	Extracts prepared from naturally infected nut tissue	Overy et al. (2003)
Trichothecene	HPLC analysis after SFE	Diode array detection or GC with electron capture detection	Wheat	Josephs et al. (1998)
OTA	Normal HPLC	Fluorescent	Dried figs	Karbancioglu-Güler and Heperkan, (2008)
OTA	Reversed-phase coupled to HPLC	Fluorescent	Grape	Hernández et al. (2006)
Zearalenone	Atmospheric-pressure chemical ionization mass spectrometry	Fluorescent	Food	Rosenberg et al. (1998)
Deoxynivalenol, T-2	Direct analysis without derivatization	GC-MS	Cereals	Onji et al. (1998)
Deoxynivalenol and OTA	GC-MS combined with electronic nose	GC-MS	Barley grains	Olsson et al. (2002)
Trichothecenes	Trichothecenes in fungal cultures using GC-tandem MS	GC-MS	Fungal cultures	Nielsen and Thrane (2001)
Sterigmatocystin	GC-MS along with other methods for comparison such as LC-MS, LC-MS/MS, and LC-UV methods for STE determination	UV detection	Food samples	Tanaka et al. (2007)

accurate and specific detection of mycotoxins. However, the use of MS as an analytical tool will also add the high cost of equipment, complex laboratory requirements, and limitations in the type of the solvents used in extraction and separation.

Gas Chromatography

Gas chromatography (GC) is regularly used to identify and quantify the presence of mycotoxins in food samples, by which many protocols have been developed for these samples. Normally, the system is linked to MS, flame ionization detector or Fourier transform infrared spectroscopy detection techniques (Onji et al., 1998). However, most mycotoxins are not volatile and, therefore, have to be chemically derivatized prior to GC analysis (Scott, 1995). So far, several techniques have been

developed for the derivatization of mycotoxins. For example, chemical reactions such as silylation or polyfluoroacylation are employed in order to obtain a volatile material (Scott, 1995). Several examples of OTA detection utilizing derivatization have been reported (Jiao et al., 1992). This approach gave a LOD of 0.1 µg/kg. In general, the use of GC detection is not expected for commercial protocols due to existence of cheaper and faster alternatives such as HPLC.

Capillary Electrophoresis

The individual detection of closely related mycotoxins requires sophisticated separation technique. The effective separation of components can be based upon charge- and mass-dependent migration in an electrical field. The fast separations

can be accomplished by capillary electrophoresis in aqueous buffer solutions, excluding the need for organic solvents. The combination of CE with sensitive fluorescence-based detection methods have been described for AFT (Pena et al., 2002) and fumonisin (Wilkes and Sutherland, 1998). Maragos and Appell (2007) and Murillo-Arbizu et al. (2010) reported the determinations of zearalenone in maize and patulin in apple juice using capillary electrophoresis, respectively, with the high selectivity, satisfactory recovery, and low-limit detection.

OTHER DETERMINATION METHODS

Colorimetric Technique

Some mycotoxins such as citrinin have a conjugated, planar structure that gives natural fluorescence, which makes it feasible for qualitative and quantitative determination by using a fluorometer. Rapid semiquantitative fluorimetric assay was used for citrinin testing in peanuts (Trantham and Wilson, 1984). Under an acidic environment condition, the weak native fluorescence of citrinin could be remarkably enhanced. Due to diverse influencing factors of the colorimetric technique, relatively low sensitivity restricted the commercial analysis of citrinin, but this analysis procedure is rapid and bears a low cost for analyzing a large number of samples. In addition, this technique allows the high-throughput analysis of a large number of industrial samples. Yang et al. (2010) reported an aptasensor for OTA using unmodified gold nanoparticles indicator. The method exhibited a linear range for a wide OTA concentration from 20 to 625 nM, with the detection limit of 20 nM. However, the colorimetric detection method requires a tedious sample preparation of extraction and clean-up. Besides, there is loss of mycotoxins during sample treatment, or due to instable chromatographic behavior or relatively low sensitivity and recovery.

Bioassay Technique

Bioassays have become increasingly useful for mycotoxin detection (Watson and Lindsay, 1982; Yates, 1986) as an alternative of chemical analysis. Bioassay provided a rapid means for screening samples and allowed the analyst to make an informed decision. Cheli et al. (2008) reported the detection of mycotoxin-producing fungi and mycotoxin toxicity cell-based bioassays as rapid tools for evaluating food and feed mycotoxin contamination. Mitterbauer et al. (2003) reported the bioassay of strain YZRM7 allowed qualitative detection and quantification of total estrogenic activity in cereal extracts without requiring additional clean-up steps. Its high sensitivity makes this assay suitable for low-cost monitoring of contamination of maize and small grain cereals with estrogenic *Fusarium* mycotoxins.

Immunoassay Technique

Immunochemical methods provided a convenient and sensitive alternative for detecting many mycotoxins due to their

relatively low cost and easy application (Morgan, 1989; Chu, 1991; Goryacheva et al., 2007; Guan et al., 2011). Commercially available ELISA kits for detection of mycotoxins are normally based on a competitive assay format that uses either a primary antibody specific for the target molecule or a conjugate of an enzyme and the required target (Morgan, 1989; Pestka et al., 1995; Stanker et al., 2008). The simultaneous detection of multiple mycotoxins (aflatoxins B1, zearalenone, and HT-2) by the giant magnetoresistive-based immunoassay was also developed, with a detection limit of 50 pg/ml (Mak et al., 2010). A commercial ELISA kit was used to determine OTA concentrations in commonly consumed food items in Sri Lanka (Wanigasuriya et al., 2008). Table 5 presents some examples of protocols and detection limits using ELISA for the most common mycotoxins. An indirect competitive ELISA using rabbit antisera was firstly performed to detect citrinin in buffer solutions at 1–13 ng/ml (0.05–0.65 ng per assay) (Abramson et al., 1995). A direct competitive ELISA was established also, with citrinin coupled to horseradish peroxidase as the labeled antigen (Abramson et al., 1999a). More recently, a competitive ELISA for determination of citrinin concentration in dietary supplements was developed (Heber et al., 2001). Direct ELISA is quick since only 1 antibody is used and cross reactivity of secondary antibody in some cases is completely eliminated, but they suffer from the fact that immunoreactivity of the primary antibody may be reduced as a result of labeling and signal amplification becomes difficult. Conversely, indirect ELISA has several advantages, in the sense that a wide range of labeled secondary antibodies are commercially available. Indirect ELISA is generally more sensitive because each primary antibody contains several epitopes that can be bound by the labeled secondary antibody thereby enhancing signal amplification. However, the cross-reactivity may occur with the secondary antibody, resulting in nonspecific signal. Thus, the results of ELISA require to be confirmed further by other analytical methods such as sensitive HPLC. The antibodies developed for ELISA were also used for self-made immunoaffinity columns for HPLC extract clean-up (Vrabcheva et al., 1999). In addition, based on specific mycotoxin-monoclonal as well as polyclonal antibodies, a strip tests for mycotoxin screening of food and feed was developed (Cvak et al., 2010; Xu et al., 2010).

Biosensor Technique

While highly specific and sensitive, traditional immunochemical assays such as enzyme-linked immunosorbent assays are adopted, the rapid and easy use of biosensors has been developed. Paniel et al. (2010) reported the development of an electrochemical immunosensor for the detection of aflatoxin M₁ while a screen-printed carbon electrode is chosen as the transducer. Actis et al. (2010) used the ion nanogating technology as a label-free biosensor capable of identifying DNA and proteins and the sensor exhibited the ultrasensitive detection of HT-2 toxin with a detection limit of 100 fg/ml, which can compare the

Table 5 Examples of protocols and detection limits using ELISA for the most common mycotoxins

Toxin	Protocol	Matrix	Reference
Aflatoxin	Diverse microplate-based hybridoma screening methods for the generation of hapten-(aflatoxin-)-specific McAbs	Commercial mycotoxin samples	Cervino et al. (2008)
Aflatoxins	Sequential injection immunoassay	Spiked food	Betina (1984), Ye et al. (2010)
Aflatoxins	Noncommercial ELISA	Peanuts and maize	Smart et al. (1990)
Deoxynivalenol	Noncommercial ELISA	Kenyan wheat samples	Molloy et al. (2003)
Deoxynivalenol	A commercial ELISA Kit, sol-gel immunoaffinity chromatography	Wheat samples	Torres et al. (1998), Klinglmayr et al. (2010)
Deoxynivalenol and Zearalenone	Noncommercial ELISA	Food samples	Cavaliere et al. (2005)
Dihydroergosine	Competitive ELISA	Field contaminated maize	Kiihn et al. (1995)
Ergosterol	Monoclonal antibody based indirect competitive antibody based ELISA	Grains and mixed animal feed	Janardhana et al. (1999)
OTA	Commercial ELISA 96 well format	Maize grains	Matrella et al. (2006)
OTA	CE with laser-induced fluorescence	Various products	Corneli and Maragos (1998)
OTA	Comparison of CE and ELISA	Food samples	Köller et al. (2006)
OTA	Noncommercial ELISA	Food samples	Pronyk et al. (2006)

conventional assay techniques. Furthermore, Khan et al. (2010) demonstrated mycotoxin detection using antibody immobilized conducting polymer supported electrochemically polymerized acacia gum while Yang et al. (2010) reported the aptamer-based colorimetric biosensor to detect Ochratoxin A using unmodified gold nanoparticles indicator. In addition, horseradish peroxidase has been successfully immobilized in a polypyrrole matrix onto disposable screen-printed carbon electrodes for the selective detection of ochratoxin A, with the capability of 0.1 ng/ml (Alonso-Lomillo et al., 2010). Thus, the biosensor technique exhibits great potential for rapid and simple monitoring of food contamination.

Molecular Technique

Unprecedented biotechnological advances in the past decade have delivered powerful transcriptomics methods that provide new technique for mycotoxin detection. Interestingly, the real-time PCR-based system has been developed for the detection and quantification of the most common *Fusarium* species with the similar profiles of mycotoxins they can potentially produce, and such a system enables simultaneous identification of different mycotoxin producer groups in samples (Stakheev et al., 2010). Furthermore, the fingerprinting method based on metabolomics, with respect to the identification of emerging contaminants at

all stages of the food-production chain, can be developed for risk assessment (Lancova et al., 2010).

PREVENTION OR DECONTAMINATION OF MYCOTOXINS

Prevention or decontamination of mycotoxins from fungi in food is an important issue. Although spoiled products must be removed, inoculum sources such as agricultural residues or dirty farming materials have to be minimized to reduced or avoid contamination (FAO/WHO/UNEP, 1999a; Codex Alimentarius Commission, 2003; Amézqueta et al., 2009). The use of chemical compounds is a very effective strategy to prevent mycotoxin production (FAO/WHO/UNEP, 1999b). For example, in the case of grapes, some fungicides have been found to be effective against fungal colonization and OTA production (Varga and Kozakiewicz, 2006). However, fungicides must be applied with special care since some of them, such as carbendazim, have been found to reduce fungal flora but to stimulate OTA production (Medina et al., 2007b). Application of other fungicide azoxystrobin can inhibited mycotoxin production in a carbendazim-resistant strain of *Fusarium sporotrichioides* (D'Mello et al., 2001). Table 6 lists some examples to prevent mycotoxins produced by fungi.

Table 6 Examples of prevention from mycotoxins

Toxin	Protocol	Matrix	Reference
OTA	Gamma radiation	Black pepper	Jalili et al. (2010)
Aflatoxins	Gamma radiation	Black pepper	Jalili et al. (2010)
Aflatoxins	Ozone	Agricultural products	Denvir et al. (1999)
Ochratoxin A	Temperature and relative humidity	Grapes	Bellí et al. (2007b)
Ochratoxin A	Preharvest fungicides	Dehydrating grapes	Valero et al. (2007b)
Ochratoxin A	Adsorbent	Food stuff	Daković et al. (2003)
Citrinin	High hydrostatic pressure	Black table olives	Tokuşoğlu et al. (2010)
OTA	<i>A. nigri</i>	Dehydrating grapes	Valero et al. (2007b)
Patulin	<i>R. glutinis</i> strain LS11	Apple	Castoria et al. (2005)

Preharvest Strategies

Cultural practices including crop rotation, tillage, planting date, and management of irrigation and fertilization have limited effects on infection and subsequent mycotoxins accumulation, whereas environmental conditions greatly affect crops, and fungi and their mycotoxins (Pose et al., 2010; Tirado et al., 2010). Kokkonen et al. (2010) investigated the influence of environmental conditions on the toxin production by *Fusarium langsethiae*, *Fusarium avenaceum*, and *Fusarium tricinctum*. To avoid fungal invasion, plant moisture, and drought stress must be reduced. Some of the strategies involved are appropriate irrigation, mineral nutrition and fertilization (FAO/WHO/UNEP, 1999b). Other good agricultural practices (GAP) are crop rotation, culture, and harvest in the appropriate seasons and conditions (space each plant for optimal growth, avoid extreme drought periods) and land deep plough (Codex Alimentarius Commission, 2003; Elmholt, 2003; Aldred and Magan, 2004). At present, new and promising preharvest biotechnology strategies are being explored (FAO/WHO/UNEP, 1999b). These new approaches include the design and production of new plants that reduce the incidence of fungal infection, restrict the growth of toxigenic fungi, prevent toxin accumulation, or are resistant to the action of insects.

Strategies during Harvest

The correct performance and cleanliness of farming equipment is recommended (Codex Alimentarius Commission, 2003). Mature fruits must be collected in sacks or containers free of contamination. Those overripe, damaged, or fallen onto the soil must be discarded and eliminated even off the field, as they are likely to have high OTA levels or to harbor OTA-producing molds which could be rapidly propagated (Bucheli and Taniwaki, 2002; Paulino de Moraes and Luchese, 2003; Pérez de Obanos et al., 2005; Belli et al., 2007a). Blend with poor quality products or with other commodities is also unadvisable (López-García et al., 2008).

Postharvest Handling Strategies

Prevention through preharvest management is the effective method for controlling mycotoxin contamination. In the postharvest phase, the major topic is to prevent contamination from mycotoxins (FAO/WHO/UNEP, 1999c). Postharvest handling strategies to prevent mycotoxins from fungi include physical, chemical, or microbiological treatments, storage, and processing by destroying, modifying or absorbing these mycotoxins (FAO/WHO/UNEP, 1999a; Morales et al., 2010). The ideal methods would be easy to use and economical, and would not generate toxic compounds or alter other fruit quality parameters such as nutrient content (EMAN, 2003).

Physical Methods

A freezing (-20°C) process, and UV and gamma treatments are able to diminish the production of mold conidia. Nevertheless, only gamma radiation can destroy the mycotoxin (Deberghes et al., 1993, 1995; Aziz et al., 2004; Bhat et al., 2010), but the gamma rays even at 60 kGy were not effective in completely destroying of ochratoxin and aflatoxins present in black pepper (Jalili et al., 2010). Application of combined infrared radiation heating with ultraviolet irradiation reduced surface decontamination of fig fruit (Hamanaka et al., 2010). However, thermal treatments do not completely eliminate OTA (Boudra et al., 1995). Another proposed method is ozonization. The development of electrochemical techniques has allowed the ozone use for OTA removal up to undetectable levels in foodstuffs such as grains, nuts, or vegetables (McKenzie et al., 1997; Denvir et al., 1999).

Chemical Methods

Postharvest treatment with fungicides can reduce production of mycotoxins by fungi. In grapes, it has been demonstrated that some fungicides, pesticides, and insecticides have positive effects on mycotoxin detoxification (Varga and Kozakiewicz, 2006; Reddy et al., 2007; Belli et al., 2007b; Valero et al., 2007b; Lehocski-Krsjak et al., 2010). Some fungicides such as itraconazole and amphotericin B have been shown to effectively control the aflatoxin-producing *Aspergillus* species (Ni and Streett, 2005). Application of alternative fungicide azoxystrobin can inhibit mycotoxin production in a carbendazim-resistant strain of *F. sporotrichioides* (D'Mello et al., 2001). Overall, use of fungicides is being discouraged due to economic reasons and growing concern for environment- and food-safety issues. Other chemicals such as ethanol inhibited food spoilage fungi and, thus, reduced mycotoxin production (Dao and Dantigny, 2010).

For OTA detoxification with chemical compounds, many adsorbent materials such as activated charcoal, cholestyramine, sodium and calcium aluminum silicates (mainly zeolites), bentonite, or wood fragments have been tested (Bauer, 1994; Leong et al., 2006; Varga and Kozakiewicz, 2006; Péteri et al., 2007; Ringot et al., 2007; Savino et al., 2007). However, their activities in vivo were not as high as expected except for the activated charcoal (Scott, 1996; Huwig et al., 2001). A new insoluble vegetal fiber has been developed in order to adsorb the OTA present in liquid food products (Tangni et al., 2005). Currently, the most promising adsorbent materials are modified zeolites as they have shown good results in foodstuff decontamination (Schall et al., 2002; Daković et al., 2003; Tomasevic-Canovic et al., 2003; Daković et al., 2005). It is noted that there is a risk that nonspecific adsorption agents using enterosorbents may prevent uptake of micronutrients from the food.

As OTA is known to cause cell membrane damage through increased lipid peroxidation, the protective properties of antioxidant substances are being investigated. These properties of these

compounds can act as superoxide anion scavengers, thereby protecting cell membrane from the mycotoxin-induced damage (Atroshi et al., 2000; Atroshi et al., 2002; Ruiz, 2010). A new strategy under study is the use of antioxidants such as vanillic acid or 4-hydroxybenzoic acid (Palumbo et al., 2007) and essential oils extracted from plants such as *Thymus vulgaris*, *Aframomum danielli*, *Ocimum gratissimum* L., or *Trachyspermum ammi* (L.) Sprague ex Turill (Aroyeun and Adegoke, 2007; Nguefack et al., 2007; Prakash et al., 2010; Velazhahan et al., 2010), which can inhibit mold growth and OTA synthesis. Furthermore, the degradation of mycotoxins can be reduced with an ethanolic solution and intermittent ultrasounds (Lindner, 1992). Alkaline hydrogen peroxide, sodium hydroxide, and monomethylamine or ammonium with calcium hydroxide treatments are also effective methods against OTA in this matrix.

Biological Methods

Significant efforts have been made in establishing various biocontrol strategies such as development of atoxigenic biocontrol fungi that can out-compete their closely related and toxigenic strains in field environments, thus reducing the levels of mycotoxins in the crops. Bejaoui et al. (2004) reported ochratoxin A removal in synthetic and natural grape juices by selected oenological *Saccharomyces* strains while Castoria et al. (2005) evaluated the effect of a yeast *Rhodotorula glutinis* strain LS11 on patulin accumulation in stored apples. In the case of products affected by *A. carbonarius*, natamycin appears to be an effective substance for controlling fungal growth and OTA production (Medina et al., 2007a). Application of nontoxigenic biocompetitive agents as nontoxigenic *A. niger* strains is also a potentially useful strategy (Valero et al., 2007a). Carboxypeptidase A is an enzyme capable of destroying OTA (Deberghes et al., 1995) and the use of atoxigenic *A. niger* strains as carboxypeptidase sources has been suggested (Varga et al., 2000). Other enzymes that can be obtained from *A. niger* strains and can efficiently degrade OTA are lipases and metalloenzyme (Stander et al., 2000; Abrunhosa et al., 2006; Abrunhosa and Venancio, 2007). A carboxypeptidase present in *Phaffia rhodozyma* can also degrade OTA up to 90% (Péteri et al., 2007). Moreover, certain bacteria belonging to *Streptococcus*, *Bifidobacterium*, *Lactobacillus*, *Butyrivibrio*, *Phenyllobacterium*, *Pleurotus*, *Saccharomyces*, *Bacillus*, and *Acinetobacter* genera (Varga et al., 2000; Varga et al., 2005; Fuchs et al., 2008) and certain fungi belonging to *Aspergillus* (*A. fumigatus*, *A. niger*, *A. carbonarius*, *A. japonicus*, *A. versicolor*, *A. wentii*, and *A. ochraceus*), *Alternaria*, *Botrytis*, *Cladosporium*, *Phaffia*, *Penicillium*, and *Rhizopus* (*R. stolonifer* and *R. oryzae*) genera (Abrunhosa et al., 2002; Varga et al., 2005; Péteri et al., 2007) are able to degrade OTA in vitro up to more than 95%. Recently, Heintz et al. (2010) have reported the degradation of fumonisin B1 by the consecutive action of 2 bacterial enzymes from *Sphingopyxis* sp. MTA144. Moreover, some of them have shown detoxifying properties in vivo assays (Fuchs et al., 2008). For example, detoxification of aflatoxin B1

and patulin by *Enterococcus faecium* strains was reported by Topcu et al. (2010). In addition, the established the *Monascus* genetic system showed high efficiency and specificity, which provides us a potential approach to manipulate and improve industrial *Monascus* strains to eliminate the citrinin production (Jia et al., 2010).

Fruit Storage and Processing

Storage environmental conditions greatly affect fungi and their mycotoxins (van der Westhuizen et al., 2010). Proper storage is necessary to prevent biological activity through adequate drying to less than 10% moisture, elimination of insect activity that can increase moisture content through condensation of moisture resulting from respiration, low temperatures, and inert atmospheres. A storage temperature of 6°C or below could be considered as safe for tomato fruits and high moisture enhanced production of *Alternaria* toxins (Pose et al., 2010). Application of the controlled atmospheres containing 80% carbon dioxide and 20% oxygen reduced significantly the growth and mycotoxin production by *Mucor plumbeus*, *Fusarium oxysporum*, *Byssoschlamys fulva*, *Byssoschlamys nivea*, *Penicillium commune*, *Penicillium roqueforti*, *A. flavus*, *Eurotium chevalieri*, and *Xeromyces bisporus* (Taniwaki et al., 2010). Furthermore, the modified atmosphere packaging conditions inhibited growth and reduced aflatoxin production of *A. flavus* (Ellis et al., 1993). In addition, the sealed or vacuum conditions can prevent mycotoxin occurrence of red chili pepper during storage (Duman, 2010). Thus, fruit storage is needed to optimize the conditions of temperatures, humidity, gas atmosphere, and packaging.

During vine fruits production, efficient drying and turning over are required because, in these stages, moisture and sugar content stimulates *Aspergillus* and OTA synthesis (Magan and Aldred, 2005). It has also been proposed to eliminate the products colonized by molds from the manufacturing process. For example, high hydrostatic pressure as a nonthermal technology is an effective tool for microbiologically safe and can reduce the citrinin production by filamentous fungi such as *Penicillium citrinum*, *Penicillium*, *Aspergillus*, and *Monascus* (Tokuşoğlu et al., 2010). For grape case, fruit washing, cooking, and pressing can also promote the OTA content drop (Delage et al., 2003; Ratola et al., 2005).

HACCP System

Hazard Analysis and Critical Control Point (HACCP) is a quality system that identifies, evaluates, and controls hazards which are significant for food safety. It is a structured, systematic approach for the control of food safety throughout the commodity system, from the plough to the plate (Martínez-Rodríguez and Carrascosa, 2009). It requires a good understanding of the relationship between cause and effect and it is a key element in total quality management. HACCP builds on the foundations

of well-established quality management systems such as good manufacturing practice, good hygienic practice, GAP, and good storage practice (FAO, 2001). By applying the HACCP concept to the management of the likelihood of an adverse health effect as a result of exposure to mycotoxins, an adequate, wholesome, and safe food supply would be maintained. In order to design and develop effective HACCP-based integrated mycotoxin management programs, a given country would have to consider factors such as climate, farming systems, preharvest and postharvest technologies, public health significance of the contaminant, producer/processor compliance, the availability of analytical resources, and even the economy. In an ideal integrated management system that incorporates the HACCP concept, mycotoxins would be minimized at every stage of production, harvesting, processing, and distribution (FAO/WHO/UNEP, 1999c).

CONCLUSION AND PROSPECTS

The presence of mycotoxins such as aflatoxins, ochratoxin A, and patulin in fruits and their processed products such as juices, wines, or cider is of high concern for human health due to their properties to induce severe acute and chronic toxicity at low-dose levels (Garcia et al., 2009; Chassy, 2010). There are increasing reports on different and less-obvious sources of alimentary exposure, in addition to the conventional studied and worldwide consumed fruit matrices for these mycotoxins, that is, apple for patulin, grape for ochratoxin, and fig for aflatoxins (Sakuda and Kimura, 2010). Particularly, the occurrence of mycotoxins in juices is of high concern because children are one of the main consumers and because juice consumption is greater than that of wine. Consequently, improved monitoring programs should be encouraged. The cooccurrence of these different mycotoxins in the same matrix is another point that requires more studies from a toxicological and occurrence point of view.

A broad range of detection techniques used for practical analysis and detection of a wide spectrum of mycotoxins are available (Turner et al., 2009). Many analytical methods have been developed for the determination of each group of these mycotoxins in different matrices. Though there have been several recent successes in detection of mycotoxins, new methods are still required to achieve higher sensitivity and address other challenges that are posed by these mycotoxins. The application of MS in conjunction with other tools for decreasing limits of detection has been of increased interest in the recent times. Future trends would focus on rapid assays and tools that would measure multiple toxins from a single matrix and the development of rapid screening methods is also advisable in order to increase the number of monitored samples.

Apart from the regulatory controls, some major strategies have been adopted to decrease or even eliminate the presence of the mycotoxins in foods: prevention of mycotoxin contamination during the preharvest and postharvest periods, detoxification of mycotoxins present in foods, and inhibition of mycotoxin absorption in the gastrointestinal tract. Preventive mea-

sures aimed at the inhibition of mycotoxin formation in agricultural products are the most effective approach for avoiding consumer exposure. Good farm management, appropriate culture methods, use of insecticides, fungicides and biological control, irrigation, and cultivar selection ensure plants less vulnerable to fungal infection. Postharvest contamination can be avoided by controlling moisture, temperature, and microbiological, insect and animal pests. However, detoxification of mycotoxins by different physical, chemical, and biological methods are less effective and sometimes restricted because of concerns of safety, possible losses in nutritional quality of the treated commodities and cost implications. Some of the most promising interventions studied to date involve the use of microorganisms to reduce absorption of mycotoxins from consumed foods in the gastrointestinal tract, but further studies are needed. In addition, efforts have been done to study the critical points of the mycotoxin presence in the manufacturing chain of affected commodities. To effectively eliminate mycotoxins, an integrated management system should be adopted at every stage of production, harvesting, processing, and distribution.

As fungus-derived mycotoxins are unavoidable, new approaches should be developed using biotechnology. Potential biochemical and genetic resistance markers have been identified in crops in different parts of the world which are being utilized as selectable markers in breeding for resistance to aflatoxin contamination. Prototypes of genetically engineered crops have been developed which contain genes for resistance to the phytotoxic effects of certain trichothecenes, thereby helping reduce fungal virulence or contain genes encoding fungal growth inhibitors for reducing fungal infection. Breeding approaches allowed in the past the creation of new varieties with lower toxin levels (Lehoczki-Krsjak et al., 2010). With the rapid development of molecular biology, gene technology could further improve crop varieties to better prevent contamination of mycotoxins by fungi in fruits and their processed products.

ACKNOWLEDGMENTS

This work was supported by the National Natural Science Foundation of China (Grant No. 30928017), the CAS/SAFEA international partnership program for creative research teams, and the 5th China-South African Joint Research Programme (grant no. CS05-L08) by the Ministry of Science and Technology, China.

ABBREVIATIONS

AFB-1	= Aflatoxin B1;
AFG-1	= Aflatoxin G1;
AFT	= Aflatoxin;
AOH	= Dibenzo- α -pyrones alternariol;
ALT	= Altenuene;

APCI	= Atmospheric pressure chemical ionization;	OTA	= Ochratoxin;
ATX-I	= Alvertoxin I;	PAT	= Patulin; ppb, parts per billion; ppm, parts per million;
ATX-II	= Alvertoxin II;	RNA	= Ribonucleic acid;
C-18	= Octadecylsilane polymer resin;	RP	= Reverse phase;
CE	= Capillary electrophoresis;	RP-HPLC-ESI-MS	= Reversed-phase HPLC-electrospray ionization-mass spectrometry;
CI-ELISA	= Competitive indirect enzyme-linked immunosorbent assay;	SAX	= Strong anion exchange;
CIT	= Citrinin;	SFE	= Supercritical fluid extraction;
CPZ	= Cyclopiazonic acid;	SIIA	= Sequential injection immunoassay;
DAD	= Diode array detection;	SPE	= Solid phase extraction;
DNA	= Deoxyribonucleic acid;	SPME	= Solid phase microextraction;
DON	= Deoxynivalenol;	SPME-LC-MS/MS	= Solid phase microextraction coupled with liquid chromatography-mass spectrometry-mass spectrometry;
ECD	= Electron capture detector;	SPR	= Surface plasmon resonance;
ELISA	= Enzyme-linked immunosorbent assay;	TLC	= Thin-layer chromatography;
FB1	= Fuminosin B1;	TRC	= Trichothecenes;
FB2	= Fuminosin B2;	WHO	= World Health Organization;
FI	= Flame ionization;	ZEA	= Zearalone;
FID	= Flame ionization detector;	ZON	= Zearalanone
FD	= Fluorescence detector;		
FTIR	= Fourier transform infrared spectroscopy;		
GC	= Gas chromatography;		
GC-ECD	= Gas chromatography coupled with electron capture detector;		
GC-MS	= Gas chromatography-mass spectrometry;		
HACCP	= Hazard analysis and critical control point;		
HPLC	= High-performance liquid chromatography;		
HPLC-DAD	= High-performance liquid chromatography coupled with diode array detector; HPLC-ELSD, HPLC coupled with an evaporative laser scattering detector;		
HPLC-FD	= High-performance liquid chromatography coupled with fluorescence detector; HT-2, HT-2 toxin (15-Acetoxy-3 α ,4 β -dihydroxy-8 α -(3-methylbutyryloxy)-12,13-epoxytrichothec-9-ene);		
IAC	= Immuno affinity columns;		
IARC	= International Agency for Research on Cancer;		
LC-MS	= Liquid chromatography-mass spectrometry;		
LLC	= Liquid-liquid chromatography;		
LLE	= Liquid-liquid extraction;		
LOD	= Limits of detection;		
LOQ	= Limits of quantification;		
MIP	= Molecularly imprinted polymers;		
MS	= Mass spectrometry;		
MW	= Molecular weight;		
NICI	= Negative ion chemical ionization;		
NIV	= Nivalenol;		
OPA	= o-phthaldialdehyde;		

REFERENCES

- Abdel Sater, M. A., Zohri, A. A., and Ismail, M. A. (2001). Natural contamination of some Egyptian. *J. Food Sci. Technol.* **38**:407-411.
- Abrunhosa, L., Santos, L., and Venancio, A. (2006). Degradation of ochratoxin A by proteases and by a crude enzyme of *Aspergillus niger*. *Food Biotechnol.* **20**:231-242.
- Abrunhosa, L., Serra, R., and Venancio, A. (2002). Biodegradation of ochratoxin A by fungi isolated from grapes. *J. Agric. Food Chem.* **50**:7493-7496.
- Abrunhosa L., Paterson, R. R. M., Kozakiewicz, Z., Lima, N., and A. Venancio, A. (2001). Mycotoxin production from fungi isolated from grapes. *Lett. Appl. Microbiol.* **32**:40-242.
- Abramson, D., Usleber, E., and Martlbauer, E. (1995). An indirect enzyme immunoassay for the mycotoxin citrinin. *Appl. Environ. Microbiol.* **61**:2007-2009.
- Abramson, D., Usleber, E., and Märtilbauer, E. (1999a). Rapid determination of citrinin in corn by fluorescence liquid chromatography and enzyme immunoassay. *J. AOAC Int.* **82**:1353-1356.
- Abramson, D., Hulasare, R., White, N. D. G., Jayas, D. S., and Marquardt, R. (1999b). Mycotoxin formation in hullless barley during granary storage at 15 and 19% moisture content. *J. Stored Prod. Res.* **35**:297-305.
- Abrunhosa, L. and Venancio, E. A. (2007). Isolation and purification of an enzyme hydrolyzing ochratoxin A from *Aspergillus niger*. *Biotechnol. Lett.* **29**:1909-1914.
- Actis, P., Jejelowo, O., and Pourmand, N. (2010). Ultrasensitive mycotoxin detection by STING sensors. *Biosens. Bioelectron.* **26**:333-337.
- Afsah-Hejri, L., Jinap, S., Arzandeh, S., and Mirhosseini, H. (2011). Optimization of HPLC conditions for quantitative analysis of aflatoxins in contaminated peanut. *Food Control.* **22**:381-388.
- Akiyama, H., Goda, Y., Tanaka, T., and Toyoda, M. (2001). Determination of aflatoxins B1, B2, G1 and G2 in spices using a multifunctional column clean-up. *J. Chromatogr. A.* **932**:153-157.
- Aldred, D. and Magan, N. (2004). Prevention strategies for trichothecenes. *Toxicol. Lett.* **153**:165-171.
- Al-Hazmi, N. A. (2010). Determination of Patulin and Ochratoxin A using HPLC in apple juice samples in Saudi Arabia. *Saudi J. Biol. Sci.* **17**:353-359.
- Ali, W. H., Derrien, D., Alix, F., Pérolle, C., Lépine, O., Bayoudh, S., Chapuis-Hugon, F., and Pichon, V. (2010). Solid-phase extraction using molecularly

- imprinted polymers for selective extraction of a mycotoxin in cereals. *J. Chromatogr. A*. **1217**:6668–6673.
- Alonso-Lomillo, M. A., Domínguez-Renedo, O., Ferreira-Gonçalves, L., and Arcos-Martínez, M. J. (2010). Sensitive enzyme-biosensor based on screen-printed electrodes for Ochratoxin A. *Biosens. Bioelectron.* **25**:1333–1337.
- Ambrosino, P., Galvano, F., Fogliano, V., Logrieco, A., Fresa, R., and Ritieni, A. (2004). Supercritical fluid extraction of Beauvericin from maize. *Talanta*. **62**:523–530.
- Amézqueta, S., González-Peñas, E., Murillo-Arbizu, M., and Cerain, A. L. D. (2009). Ochratoxin A decontamination: A review. *Food Control*. **20**:326–333.
- Aroyeun, S. O. and Adegoke, G. O. (2007). Reduction of ochratoxin A (OTA) in spiked cocoa powder and beverage using aqueous extracts and essential oils of *Aframomum danielli*. *Afr. J. Biotechnol.* **6**:612–616.
- Asao, T., Buchi, G., Abdel Kader, M. M., Chang, S. B., Wick, E. L., and Wogan, G. N. (1965). Structures of aflatoxins B and G1. *J. Amer. Chem. Soc.* **87**:882–886.
- Atroschi, F., Rizzo, A., Westermarck, T., and Ali-Vehmas, T. (2002). Antioxidant nutrients and mycotoxins. *Toxicol.* **180**:151–167.
- Atroschi, F., Biese, I., Saloniemi, H., Ali-Vehmas, T., Saari, S., Rizzo, A., and Veijalainen, P. (2000). Significance of apoptosis and its relationship to antioxidants after ochratoxin A administration in mice. *J. Pharm. Pharm. Sci.* **3**:281–291.
- Aziz, N. H., Moussa, L. A. A., and Far, F. M. E. (2004). Reduction of fungi and mycotoxins formation in seeds by gamma-radiation. *J. Food Safety*. **24**:109–127.
- Bailly, J. D., Querin, A., Le Bars-Bailly, S., Benard, G., and Guerre, P. (2002). Citrinin production and stability in cheese. *J. Food Prot.* **65**:1317–1321.
- Barkai-Golan, R. and Paster, N. (2008). *Mycotoxins in Fruits and Vegetables*. Academic Press, New York.
- Food and Agriculture Organization of the United Nations (FAO). (2003). World-wide regulation for mycotoxins in food and feed in 2003. 120.
- Battilani, P., Magan N., and Logrieco, A. (2006). European research on ochratoxin A in grapes and wine. *Int. J. Food Microbiol.* **111**(Supplement 1):S2–S4.
- Bauer, J. (1994). Möglichkeiten zur entgiftung mykotoxinhaltiger futtersmittel. *Monatshefte für Veterinärmedizin*. **49**:175–181.
- Bauer, J. and M. Gareis, M. (1987). Ochratoxin-A in the food-chain. *J. Vet. Med. B* **34**:613–627.
- Bejaoui, H., Mathieu, F., Taillandier, P., and Lebrihi, A. (2004). Ochratoxin A removal in synthetic and natural grape juices by selected oenological *Saccharomyces* strains. *J. Appl. Microbiol.* **97**:1038–1044.
- Belli, N., Marín, S., Argiles, E., Ramos, A. J., and Sanchís, V. (2007a). Effect of chemical treatments on ochratoxigenic fungi and common mycobiota of grapes (*Vitis vinifera*). *J. Food Prot.* **70**:157–163.
- Bellí, N., Marín, S., Coronas, I., Sanchís, V., and Ramos, A. J. (2007b). Skin damage, high temperature and relative humidity as detrimental factors for *Aspergillus carbonarius* infection and ochratoxin A production in grapes. *Food Control*. **18**:1343–1349.
- Beltrán, E., Ibáñez, M., Sancho, J. V., Cortés, M. A., Yúsá, V., and Hernández, F. (2010). UHPLC-MS/MS highly sensitive determination of aflatoxins, the aflatoxin metabolite M1 and ochratoxin A in baby food and milk. *Food Chemistry*. in press.
- Bennett, J. W. and Klich, M. (2011). Mycotoxins. *Clin. Microbiol. Rev.* **66**:737–744.
- Betina, V. (1984). *Mycotoxins: Production, Isolation, Separation, and Purification*. Elsevier, Amsterdam, The Netherlands.
- Betina, V. (1985). Thin-layer chromatography of mycotoxins. *J. Chromatogr.* **334**:211–276.
- Betina, V. (1993). Chapter 7: Thin-layer chromatography of mycotoxins. *J. Chromatogr. Libr.* **54**:141–251.
- Bhat, R., Sridhar, K. R., and Karim, A. A. (2010). Microbial quality evaluation and effective decontamination of nutraceutically valued lotus seeds by electron beams and gamma irradiation. *Radiat. Phys. Chem.* **79**:976–981.
- Boudra, H., Le Bars, P., and Le Bars, J. (1995). Thermostability of ochratoxin A in wheat under two moisture conditions. *Appl. Environ. Microbiol.* **61**:1156–1158.
- Bradburn, N., Coker, R. D., and Jewers, K. (1990). Development of rapid methods for the analysis of aflatoxin in agricultural products. 7. A comparative study of phenyl bonded-phase, CB and romer cleanup procedures for determining aflatoxin levels in maize by bidirectional HPTLC. *Chromatographia*. **29**:177–181.
- Brera, C., Debegnach, F., De Santis, B., Pannunzi, E., Berdini, C., Prantera, E., Gregori, E., and Miraglia, M. (2011). Simultaneous determination of aflatoxins and ochratoxin A in baby foods and paprika by HPLC with fluorescence detection: A single-laboratory validation study. *Talanta*. **83**:1442–1446.
- Bucheli, P. and Taniwaki, M. H. (2002). Review. Research on the origin, and on the impact of post-harvest handling and manufacturing on the presence of ochratoxin A in coffee. *Food Addit. Contam.* **19**:655–665.
- CAST. (2003). *Mycotoxin: Risks in Plant, Animal, and Human Systems*. Council of Agricultural Science and Technology, Task force report No. 139. Ames, Iowa, USA.
- Castoria, R., Morena, V., Caputo, L., Panfili, G., De Curtis, F., and De Cicco, V. (2005). Effect of the biocontrol yeast *Rhodotorula glutinis* strain LS11 on patulin accumulation in stored apples. *Phytopathol.* **95**:1271–1278.
- Cavaliere, C., Ascenzo, G. D., Foglia, P., Pastorini, E., Samperi, R., and Lagana, A. (2005). Determination of type B trichothecenes and macrocyclic lactone mycotoxins in field contaminated maize. *Food Chem.* **92**:559–568.
- Cervino, C., Weber, E., Knopp, D., and Niessner, R. (2008). Comparison of hybridoma screening methods for the efficient detection of high-affinity hapten-specific monoclonal antibodies. *J. Immunol. Methods*. **329**:184–193.
- Chassy, B. M. (2010). Food safety risks and consumer health. *New Biotechnol.* **27**:534–544.
- Cheli, F., Pinotti, L., Campagnoli, A., Fusi, E., Rebucci, R., and Baldi, A. (2008). Mycotoxin analysis, mycotoxin-producing fungi assays and mycotoxin toxicity bioassays in food mycotoxin monitoring and surveillance. *Ital. J. Food Sci.* **20**:447–462.
- Cheraghali, A. M., Mohammadi, H. R., Amirahmadi, M., Yazdanpanah, H., Abouhossain, G., Zamanian, F., Khansari, M. G., and Afshar, M. (2005). Incidence of patulin contamination in apple juice produced in Iran. *Food Control*. **16**:165–167.
- Chiavaro E., Dall'Asta, C., Galaverna, G., Biancardi, A., Gambarelli, E., and Dossena, A. (2001). New reversed-phase liquid chromatographic method to detect aflatoxins in food and feed with cyclodextrins as fluorescence enhancers added to the eluent. *J. Chromatogr. A*. **937**:31–40.
- Chu, F. S. (1991). Current immunochemical methods for mycotoxin analysis. In: *Immunoassays for Trace Chemical Analysis: Monitoring Toxic Chemicals in Humans, Food and the Environment*, pp. 140–157. Vandelaan, M., Stanker, L. H., Watkins, B. E., and Roberts, D. W., Eds., American Chemical Society, Washington, DC.
- Chu, F. S. (1992). Mycotoxins in ecological systems. In: *Handbook of Applied Mycology*, pp. 87–136. Bhatnagar, D., Lillehoj, E. B., and Arora, D. K., Eds., Marcel Dekker, New York.
- Castoria, R., Morena, V., Caputo, L., Panfili, G., De Curtis, F., and De Cicco, V. (2005). Effect of the biocontrol yeast *Rhodotorula glutinis* strain LS11 on patulin accumulation in stored apples. *Phytopathol.* **95**:1271–1278.
- Codex Alimentarius Commission. (2003). Report of the 34th session of the codex committee on food additives and contaminants.
- Colinski, P. and Rabarkiewicz-Szcesna, J. (1984). Chemical confirmatory tests for ochratoxin A, citrinin, penicillic acid, sterigmatocystin, and zearalenone performed directly on thin-layer chromatographic plates. *J. Assoc. Off. Anal. Chem.* **67**:1108–1110.
- Combina, M., Dalcero, A., Varsavsky, E., Torres, A., Etcheverry, M., Rodriguez, M., and Gonzalez, Q. H. (1999). Effect of heat treatments on stability of alternariol, alternariol monomethyl ether and tenuazonic acid in sunflower flour. *Mycotoxin Res.* **15**:33–38.
- Corneli, S. and Maragos, C. M. (1998). Capillary electrophoresis with laser-induced fluorescence: Method for the mycotoxin ochratoxin A. *J. Agric. Food Chem.* **46**:3162–3165.
- Cvak, B., Molinelli, A., and Krska, R. (2010). Development of immunodiagnostic strip tests for mycotoxin screening of food and feed. *Toxicol. Lett.* **196**(Supplement 1):S337–S338.

- Daković A., Tomašević-Čanović, M., Rottinghaus, G. E., Dondur, V., and Mašić, A. (2003). Adsorption of ochratoxin A on octadecyldimethyl benzyl ammonium exchanged-clinoptilolite-heulandite tuff. *Colloid Surf. B-Biointerfaces*. **30**:157–165.
- Daković, A., Tomašević-Čanović, M., Dondur, V., Rottinghaus, G. E., Medaković, V., and Zarić, S. (2005). Adsorption of mycotoxins by organozeolites. *Colloid Surf. B-Biointerfaces*. **46**:20–25.
- Dao, T. and Dantigny, P. (2011). Control of food spoilage fungi by ethanol. *Food Control*. **22**:360–368.
- Dashti, B. H. R. and AL-Hamli, S. F. (2010). Investigation of mycotoxins in selected food commodities in Kuwait: A case study. *J. Amer. Diet. Assoc.* **110**(Supplement 1):A92.
- Deberghes, P., Deffieux, G., Gharbi, A., Betbeder, A. M., Boisard, F., Blanc, R., Delaby, J. F., and Creppy, E. E. (1993). Detoxification de l'ochratoxine A par des moyens physiques, chimiques et enzymatiques. *Human Ochratoxicosis Pathol.* **231**:75–82.
- Deberghes, P., Betbeder, A. M., Boisard, F., Blanc, R., Delaby, J. F., Krivobok, S., Steiman, R., Seigle-Murandi, F., and Creppy, E. E. (1995). Detoxification of ochratoxin A, a food contaminant: Prevention of growth of *Aspergillus ochraceus* and its production of ochratoxin A. *Mycotoxin Res.* **11**:37–47.
- Delage, N., d'Harlingue, A., Ceccaldi, B. C., and Bompeix, G. (2003). Occurrence of mycotoxins in fruit juices and wine. *Food Control*. **14**:225–227.
- Delgado, T., Cordovés, C. G., and Scott, P. M. (1996). Determination of alternariol and alternariol methyl ether in apple juice using solid-phase extraction and high-performance liquid chromatography. *J. Chromatogr. A*. **731**:109–114.
- Denvir A. J., McKenzie, K. S., Rogers, T. D., Miller, D. R., Hitchens, G. D., and Andrews, C. C. (1999). Ozone treatment of agricultural products such as grains, nuts or vegetables to destroy mycotoxins and aflatoxins. Patent number WO9939723-A2.
- Diesing, A. K., Nossol, C., Panther, P., Walk, N., Post, A., Kluess, J., Kreutzmann, P., Dänike, S., Rothkötter, H. J., and Kahlert, S. (2011). Mycotoxin deoxynivalenol (DON) mediates biphasic cellular response in intestinal porcine epithelial cell lines IPEC-1 and IPEC-J2. *Toxicol. Lett.* **200**:8–18.
- D'Mello, J. P. F., Macdonald, A. M. C. and Rinna, R. (2001). Effects of azoxystrobin on mycotoxin production in a carbendazim-resistant strain of *Fusarium sporotrichioides*. *Phytoparasitica*. **29**:431–440.
- Drusch, S. and Aumann, J. (2005). Mycotoxins in fruits: Microbiology, occurrence, and changes during fruit processing. *Adv. Food Nutr. Res.* **50**:33–78.
- Drusch, S. and Ragab, W. (2003). Mycotoxins in fruits, fruit juices and dried fruits. *J. Food Prot.* **66**:1514–1527.
- Duman, A. D. (2010). Storage of red chili pepper under hermetically sealed or vacuum conditions for preservation of its quality and prevention of mycotoxin occurrence. *J. Stored Prod.s Res.* **46**:155–160.
- Edwards, J. V. and Lillehoj, E. B. (1987). Isolation and liquid-chromatographic determination of the cyclic peptide mycotoxin cyclosporine A from rice. *J. Assoc. Off. Anal. Chem.* **70**:126–129.
- Elizalde-Gonzalez, M. P., Mattusch, J., and Wennrich, R. (1998). Stability and determination of aflatoxins by high-performance liquid chromatography with amperometric detection. *J. Chromatogr. A*. **828**:439–444.
- Ellis, W. O., Smith, J. P., Simpson, B. K., Khanizadeh, S., and Oldham, J. H. (1993). Control of growth and aflatoxin production of *Aspergillus flavus* under modified atmosphere packaging (MAP) conditions. *Food Microbiol.* **10**:9–21.
- Elmholt, S. (2003). Ecology of the ochratoxin A producing *Penicillium verrucosum*: Occurrence in field soil and grain with especial attention to farming system and on-farm drying practices. *Biol. Agric. Hort.* **20**:311–337.
- EMAN. (2003). European Mycotoxin Awareness Network co-ordinated by Leatherhead Food Research Association, UK.
- Engelhardt, G., Ruhland, M., and Wallnofer, P. R. (1999). Occurrence of ochratoxin A in moldy vegetables and fruits analysed after removal of rotten tissue parts. *Adv. Food Sci.* **21**:88–92.
- Eppley, R. M., Truckess, M. W., Nesheim, S., Thorpe, C. W., and Pohland, A. E. (1986). Thin-layer chromatographic method for determination of deoxynivalenol in wheat—collaborative study. *J. Assoc. Off. Anal. Chem.* **69**:37–40.
- FAO. (2001). Manual on the application of the HACCP system in mycotoxin prevention and control. *FAO Food and Nutrition Paper*, pp. 1–124. Available at: <http://ftp.fao.org/docrep/fao/005/y1390e/y1390e00.pdf>.
- FAO. (2006). Reducing OTA in coffee. Available at: <http://www.coffee-ota.org/>.
- FAO/WHO/UNEP. (1999a). Minimising risks posed by mycotoxins utilising the HACCP Concept. *Third Joint FAO/WHO/UNEP International Conference on Mycotoxins*, pp. 1–13. Available at: <ftp://ftp.fao.org/es/esn/food/myco8b.pdf>.
- FAO/WHO/UNEP. (1999b). Mycotoxin prevention and decontamination, corn—a case study. *Third Joint FAO/WHO/UNEP International Conference on Mycotoxins*, pp. 1–11. Available at: <ftp://ftp.fao.org/es/esn/food/myco6b.pdf>.
- FAO/WHO/UNEP. (1999c). Mycotoxin prevention and decontamination, HACCP and its mycotoxin control potential: An evaluation of ochratoxin A in coffee production. *Third Joint FAO/WHO/UNEP International Conference on Mycotoxins*, pp. 1–13. Available at: <ftp://ftp.fao.org/es/esn/food/myco6c.pdf>.
- Fernández-Cruza, M. L., Mansilla, M. L., and Tadeo, J. L. (2010). Mycotoxins in fruits and their processed products: Analysis, occurrence and health implications. *J. Adv. Res.* **1**:113–122.
- Fox, E. M. and Howlett, B. J. (2008). Secondary metabolism: Regulation and role in fungal biology. *Curr. Opin. Microbiol.* **11**:481–7.
- Frohlich, A. A., Marquardt, R. R., and Bernatsky, A. (1988). Quantization of ochratoxin A – use of reverse phase thin-layer chromatography for sample cleanup followed by liquid-chromatography or direct fluorescence measurement. *J. Assoc. Off. Anal. Chem.* **71**:949–953.
- Fuchs, S., Sontag, G., Stidl, R., Ehrlich, V., Kundi, M., and Knasmüller, S. (2008). Detoxification of patulin and ochratoxin A, two abundant mycotoxins, by lactic acid bacteria. *Food Chem. Toxicol.* **46**:1398–1407.
- Funes, G. J. and Resnik, S. L. (2009). Determination of patulin in solid and semisolid apple and pear products marketed in Argentina. *Food Control*. **20**:277–280.
- Fung, F. and Clark, R. F. (2004). Health effects of mycotoxins: A toxicological overview. *J. Toxicol. – Clin. Toxicol.* **42**:217–234.
- Garcia, D., Ramos, A. J., Sanchis, V., and Marín, S. (2009). Predicting mycotoxins in foods: A review. *Food Microbiol.* **26**:757–769.
- Garden, S. R. and Strachan, N. J. C. (2001). Novel colorimetric immunoassay for the detection of aflatoxin B1. *Anal. Chim. Acta*. **444**:187–191.
- Gimeno, A. (1984). Determination of citrinin in corn and barley on thin-layer chromatographic plates impregnated with glycolic acid. *J. Assoc. Off. Anal. Chem.* **67**:194–196.
- Gimeno, A. and Martins, M. L. (1983). Rapid thin layer chromatographic determination of patulin, citrinin, and aflatoxin in apples and pears, and their juices and jams. *J. Assoc. Off. Anal. Chem.* **66**:85–91.
- Giraudi, G., Anfossi, L., Baggiani, C., Giovannoli, C., and Tozzi, C. (2007). Solid-phase extraction of ochratoxin A from wine based on a binding hexapeptide prepared by combinatorial synthesis. *J. Chromatogr. A*. **1175**:174–180.
- Gonçalves, E., Nogueira, J. H. C., Fonseca, H., Felicio, J. D., Pino, F. A., and Corrêa, B. (2008). Mycobiota and mycotoxins in Brazilian peanut kernels from sowing to harvest. *Int. Food Microbiol.* **123**:184–190.
- Goryacheva, I. Y., De Saeger, S., Eremin, S. A., and Van Peteghem, C. (2007). Immunochemical methods for rapid mycotoxin detection: Evolution from single to multiple analyte screening: A review. *Food Addit. Contam.* **10**:1169–1183.
- Guan, D., Li, P. W., Zhang, Q., Zhang, W., Zhang, D. H., and Jiang, J. (2011). An ultra-sensitive monoclonal antibody-based competitive enzyme immunoassay for aflatoxin M1 in milk and infant milk products. *Food Chem.* **125**:1359–1364.
- Hamanaka, D., Norimura, N., Baba, N., Mano, K., Kakiuchi, M., Tanaka, F., and Uchino, T. (2011). Surface decontamination of fig fruit by combination of infrared radiation heating with ultraviolet irradiation. *Food Control*. **22**:375–380.
- Harris, K. L., Bobe, G., and Bourquin, L. D. (2009). Patulin surveillance in apple cider and juice marketed in Michigan. *J. Food Prot.* **72**:1255–1261.
- Heber, D., Lembertas, A., Lu, Q. Y., Bowerman, S., and Go, V. L. W. (2001). An analysis of nine proprietary Chinese red yeast rice dietary supplements: Implications of variability in chemical profile and contents. *J. Altern. Complement. Med.* **7**:133–139.
- Heinl, S., Hartinger, D., Thamhesl, M., Vekiru, E., Krska, R., Schatzmayr, G., Moll, W. D., and Grabherr, R. (2010). Degradation of fumonisin B1 by the consecutive action of two bacterial enzymes. *J. Biotechnol.* **145**:120–129.

- Hernández, M. A., Garcia-Moreno, M. V., Duran, E., Guillen, D., and Barroso, C. G. (2006). Validation of two analytical methods for the determination of ochratoxin A by reversed-phased high-performance liquid chromatography coupled to fluorescence detection in musts and sweet wines from Andalusia. *Anal. Chim. Acta*. **566**:117–121.
- Holcomb, M., Thompson, H. C., and Hankins, L. J. (1993). Analysis of fumonisin B1 in rodent feed by gradient elution HPLC using precolumn derivatization with fmoc and fluorescence detection. *J. Agric. Food Chem.* **41**:764–767.
- Holcomb, M., Thompson, Jr. H. C., and Cooper, W. H. (1996). SFE extraction of aflatoxins (B-1, B-2, G(1), and G(2)) from corn and analysis by HPLC. *J. Supercrit. Fluids*. **9**:118–121.
- Holcomb, M., Wilson, D. M., Trucksess, M. W., and Thompson, Jr. H. C. (1992). Determination of aflatoxins in food-products by chromatography. *J. Chromatogr.* **624**:341–352.
- Huwig, A., Freimund, S., Käppeli, O., and Dutler, H. (2001). Mycotoxin detoxication of animal feed by different adsorbents. *Toxicol. Lett.* **122**:179–188.
- Jalili, M., Jinap, S., and Noranizan, A. (2010). Effect of gamma radiation on reduction of mycotoxins in black pepper. *Food Control*. **21**:1388–1393.
- Janardhana, G. R., Raveesha, K. A., and Shetty, H. S. (1999). Mycotoxin contamination of maize grains grown in Karnataka (India). *Food Chem. Toxicol.* **37**:863–868.
- Jia, X. Q., Xu, Z. N., Zhou, L. P., and Sung, C. K. (2010). Elimination of the mycotoxin citrinin production in the industrial important strain *Monascus purpureus* SM001. *Metab. Eng.* **12**:1–7.
- Jiao, Y. Y., Blaas, W., Rühl, C., and Weber, R. (1992). Identification of ochratoxin-A in food samples by chemical derivatization and gas-chromatography mass-spectrometry. *J. Chromatogr. A*. **595**:364–367.
- Jimenez, M., Mateo, J. J., and Mateo, R. (2000). Determination of type A trichothecenes by high-performance liquid chromatography with coumarin-3-carbonyl chloride derivatisation and fluorescence detection. *J. Chromatogr. A*. **870**:473–481.
- Josephs, R. D., Krska, R., Grasserbauer, M., and Broekaert, J. A. C. (1998). Determination of trichothecene mycotoxins in wheat by use of supercritical fluid extraction and high-performance liquid chromatography with diode array detection or gas chromatography with electron capture detection. *J. Chromatogr. A*. **795**:297–304.
- Juan, C., Zinedine, A., Moltó, J. C., Idrissi, L. and Mañes, J. (2008). Aflatoxins levels in dried fruits and nuts from Rabat-Salé area, Morocco. *Food Control*. **19**:849–853.
- Karbancioglu-Guler, F. and Heperkan, D. (2009). Natural occurrence of fumonisin B1 in dried figs as an unexpected hazard. *Food Chem. Toxicol.* **47**:289–292.
- Karbancioglu-Güler, F. and Heperkan, D. (2008). Natural occurrence of ochratoxin A in dried figs. *Anal. Chim. Acta*. **617**:32–36.
- Kataoka, H., Ishizaki, A., Nonaka, Y. and Saito, K. (2010). Developments and applications of capillary microextraction techniques: A review. *Anal. Chim. Acta*. **655**:8–29.
- Katerere, D., Stockenström, S., and Shephard, G. S. (2008). HPLC-DAD method for the determination of patulin in dried apple rings. *Food Control*. **19**:389–392.
- Khan, R., Dey, N. C., Hazarika, A. K., Saini, K. K., and Dhayal, M. (2011). Mycotoxin detection on antibody immobilized conducting polymer supported electrochemically polymerized acacia gum. *Anal. Biochem.* **410**:185–190.
- Khoury, A. E. L., Rizk, T., Lteif, R., Azouri, H., Delia, M. L., and Lebrhi, A. (2008). Fungal contamination and aflatoxin B1 and ochratoxin A in Lebanese wine-grapes and musts. *Food Chem. Toxicol.* **46**:2244–2250.
- Kiihn, I., Valenta, H., and Rohr, K. (1995). Determination of ochratoxin-A in bile of swine by high-performance liquid-chromatography. *J. Chromatogr. B*. **668**:333–337.
- Klinglmayr, C., Nöbauer, K., Razzazi-Fazeli, E., and Cichna-Markl, M. (2010). Determination of deoxynivalenol in organic and conventional food and feed by sol-gel immunoaffinity chromatography and HPLC-UV detection. *J. Chromatogr. B*. **878**:187–193.
- Kok, W. T. (1994). Derivatization reactions for the determination of aflatoxins by liquid-chromatography with fluorescence detection. *J. Chromatogr. B*. **659**:127–137.
- Kokkonen, M., Ojala, L., Parikka, P. and Jestoi, M. (2010). Mycotoxin production of selected *Fusarium* species at different culture conditions. *Int. J. Food Microbiol.* **143**:17–25.
- Köller, G., Wichmann, G., Rolle-Kampczyk, U., Popp, P., and Herbarth, O. (2006). Comparison of ELISA and capillary electrophoresis with laser-induced fluorescence detection in the analysis of Ochratoxin A in low volumes of human blood serum. *J. Chromatogr. B*. **840**:94–98.
- Kussak, A., Andersson, B., Andersson, K., and Andersson, J. (1995). Immunoaffinity column cleanup for the high-performance liquid-chromatographic determination of aflatoxins B-1, B-2, G1, G2, M1 and Q1 in urine. *J. Chromatogr. B*. **672**:253–259.
- Lancova, K., Dip, R., Antignac, J. P., Bizec, B. L., Elliott, C., and Naegeli, H. (2011). Detection of hazardous food contaminants by transcriptomics fingerprinting. *TrAC Trends Anal. Chem.* **30**:181–191.
- Lau, B. P. Y., Scott, P. M., Lewis, D. A., Kanhere, S. R., Cleroux, C., and Roscoe, V. A. (2003). Liquid chromatography-mass spectrometry and liquid chromatography-tandem mass spectrometry of the *Alternaria* mycotoxins alternariol and alternariol monomethyl ether in fruit juices and beverages. *J. Chromatogr. A*. **998**:119–131.
- Leggott, N. L. and Shephard, G. S. (2001). Shephard. Patulin in South African commercial apple products. *Food Control*. **12**:73–76.
- Lehoczki-Krsjak, S., Szabo-Hever, A., Toth, B., Kotai, C., Bartok, T., Varga, M., Farady, L., and Mesterhazy, A. (2010). Prevention of *Fusarium* mycotoxin contamination by breeding and fungicide application to wheat. *Food Addit. Contam.* **27**:616–628.
- Leitner, A., Zöllner, P., Paolillo, A., and Stroka, J. (2002). Papadopoulos-Bouraoui, S. Jaborek, E. Anklam and W. Lindner, Comparison of methods for the determination of ochratoxin A in wine. *Anal. Chim. Acta*. **453**:33–41.
- Leong, S. L. L., Hocking, A. D., and Scott, E. S. (2006). The effect of juice clarification, static or rotary fermentation and fining on ochratoxin A in wine. *Aust. J. Grape Wine Res.* **12**:245–252.
- Lin, L., Zhang, J., Wang, P., Wang, Y., and Chen, J. (1998). Thin-layer chromatography of mycotoxins and comparison with other chromatographic methods. *J. Chromatogr. A*. **815**:3–20.
- Lindner, W. (1992). Decontamination and detoxication of cereals contaminated by mycotoxins. Patent number WO9216116.
- Logrieco, A. F. (2010). Mycotoxicological risks of main European food/feed chains. *Toxicol. Lett.* **196**(Supplement 1):S24.
- Logrieco, A., Ferracane, R., Haidukowsky, M., Cozzi, G., Visconti, A., and Ritieni, A. (2009). Fumonisin B2 production by *Aspergillus niger* from grapes and natural occurrence in must. *Food Addit. Contam. A*. **26**:1495–1500.
- Logrieco, A., Visconti, A., and Bottalico, A. (1990). Mandarin fruit rot caused by *Alternaria alternata* and associated mycotoxins. *Plant Dis.* **74**:415–417.
- López-García, R. (2010). Mycotoxin Management: An International Challenge. Elsevier, New York.
- López-García, R., Mallmann, C. A., and Pineiro, M. (2008). Design and implementation of an integrated management system for ochratoxin A in the coffee production chain. *Food Addit. Contam.* **25**:231–240.
- Luttfallah, G. and Hussain, A. (2011). Studies on contamination level of aflatoxins in some dried fruits and nuts of Pakistan. *Food Control*. **22**:426–429.
- Magan, N. and Aldred, D. (2005). Conditions of formation of ochratoxin A in drying, transport and in different commodities. *Food Addit. Contam.* **22**:10–16.
- Magnani, R. F., De Souza, G. D., and Rodrigues Filho, E. (2007). Analysis of alternariol and alternariol monomethyl ether on flavedo and albedo tissues of tangerines (*Citrus reticulata*) with symptoms of *Alternaria* brown spot. *J. Agric. Food Chem.* **55**:4980–4986.
- Maier, N. M., Buttinger, G., Welhartzki, S., Gavioli, E., and Lindner, W. (2004). Molecularly imprinted polymer-assisted sample clean-up of ochratoxin A from red wine: Merits and limitations. *J. Chromatogr. B*. **804**:103–111.
- Mak, A. C., Osterfeld, S. J., Yu, H., Wang, S. X., Davis, R. W., Jejelowo, O. A., and Pourmand, N. (2010). Sensitive giant magnetoresistive-based immunoassay for multiplex mycotoxin detection. *Biosens. Bioelectron.* **25**:1635–1639.
- Maragos, C. M. and Appell, M. (2007). Capillary electrophoresis of the mycotoxin zearalenone using cyclodextrin-enhanced fluorescence. *J. Chromatogr. A*. **1143**:252–257.

- Maresca, M. and Fantini, J. (2010). Some food-associated mycotoxins as potential risk factors in humans predisposed to chronic intestinal inflammatory diseases. *Toxicon*. **56**:282–294.
- Marín, S., Mateo, E. M., Sanchis, V., Valle-Algarra, F. M., Ramos, A. J., and Jiménez, M. (2010). Patulin contamination in fruit derivatives, including baby food, from the Spanish market. *Food Chem.* **124**:563–568.
- Martínez-Rodríguez, A. J., and Carrascosa, A. V. (2009). HACCP to control microbial safety hazards during winemaking: Ochratoxin A. *Food Control*. **20**:469–475.
- Martins, M. L., Gimeno, H. M., Martins, H. M. and Bernardo, F. (2002). Co-occurrence of patulin and citrinin in Portuguese apples with rotten spots. *Food Addit. Contam.* **19**:568–574.
- Mateo, J. J., Mateo, R., Hinojo, M. J., Llorens, A. and Jimenez, M. (2002). Liquid chromatographic determination of toxigenic secondary metabolites produced by *Fusarium* strains. *J. Chromatogr. A*. **955**:245–256.
- Matrella, R., Monaci, L., Milillo, M. A., Palmisano, F., and Tantillo, M. G. (2006). Ochratoxin A determination in paired kidneys and muscle samples from swines slaughtered in southern Italy. *Food Control*. **17**:114–117.
- McKenzie, K. S., Sarr, A. B., Mayura, K., Bailey, R. H., Miller, D. R., Rogers, T. D., Norred, W. P., Voss, K. A., Plattner, R. D., Kubena, L. F., and Phillips, T. D. (1997). Oxidative degradation and detoxification of mycotoxins using a novel source of ozone. *Food Chem. Toxicol.* **35**:807–820.
- Medina, A. and Magan, N. (2011). Temperature and water activity effects on production of T-2 and HT-2 by *Fusarium langsethiae* strains from north European countries. *Food Microbiol.* **28**:392–398.
- Medina, M. B. and Schwartz, D. P. (1992). Thin-layer chromatographic detection of zeranol and estradiol in fortified plasma and tissue-extracts with fast Corinth V. *J. Chromatogr. A*. **581**:119–128.
- Medina, A., Jiménez, M., Mateo, R., and Magan, N. (2007a). Efficacy of natamycin for control of growth and ochratoxin A production by *Aspergillus carbonarius* strains under different environmental conditions. *J. Appl. Microbiol.* **103**:2234–2239.
- Medina, A., Mateo, R., Valle-Algarra, F. M., Mateo E. M., and Jiménez, M. (2007b). Effect of carbendazim and physicochemical factors on the growth and ochratoxin A production of *Aspergillus carbonarius* isolated from grapes. *Int. J. Food Microbiol.* **119**:230–235.
- Meneely, J. P., Ricci, F., van Egmond, H. P., and Elliott, C. T. (2010). Current methods of analysis for the determination of trichothecene mycotoxins in food. *TrAC Trends Anal. Chem.* **30**:192–203.
- Meister, U. (2004). New method of citrinin determination by HPLC after polyamide column clean-up. *Eur. Food Res. Technol.* **218**:394–399.
- Mitterbauer, R., Weindorfer, H., Safaie, N., Krska, R., Lemmens, M., Ruckenhauer, P., Kuchler, K., and Adam, G. (2003). A sensitive and inexpensive yeast bioassay for the mycotoxin zearalenone and other compounds with estrogenic activity. *Appl. Environ. Microbiol.* **69**:805–811.
- Molloy, J. B., Moore, C. J., Bruyeres, A. G., Murray, S. A., and Blane, B. J. (2003). Determination of dihydroergosine in sorghum ergot using an immunoassay. *J. Agric. Food Chem.* **51**:3916–3919.
- Morales, H., Marín, S., Ramos, A. J. and Sanchis, V. (2010). Influence of post-harvest technologies applied during cold storage of apples in *Penicillium expansum* growth and patulin accumulation: A review. *Food Control*. **21**:953–962.
- Morgan, M. R. A. (1989). Mycotoxin immunoassays (with special reference to ELISAs). *Tetrahedron*. **45**:2237–2249.
- Moss, M. O. (2008). Fungi, quality and safety issues in fresh fruits and vegetables. *J. Appl. Microbiol.* **104**:1239–1243.
- Moukas, A., Panagiotopoulou, V., and Markaki, P. (2008). Determination of patulin in fruit juices using HPLC-DAD and GC-MSD techniques. *Food Chem.* **109**:860–867.
- Muñoz, K., Vega, M., Rios, G., Muñoz, S., and Madariaga, R. (2006). Preliminary study of Ochratoxin A in human plasma in agricultural zones of Chile and its relation to food consumption. *Food Chem. Toxicol.* **14**:1884–1889.
- Murillo-Arbizu, M., Penas, E. G., Hansen, S. H., Amézqueta, S., and Ostergaard, J. (2008). Development and validation of a microemulsion electrokinetic chromatography method for patulin quantification in commercial apple juice. *Food Chem. Toxicol.* **46**:2251–2257.
- Murillo-Arbizu, M., Gonzalez-Penas, E., and Amezceta, S. (2010). Comparison between capillary electrophoresis and high performance liquid chromatography for the study of the occurrence of patulin in apple juice intended for infants. *Food Chem. Toxicol.* **48**:2429–2434.
- Murillo-Arbizu, M., Amézqueta, S., González Peñas, E., and de Cerain, A. L. (2009). Occurrence of patulin and its dietary intake through apple juice consumption by the Spanish population. *Food Chem.* **113**:420–423.
- Neely, F. L. and Emerson, C. S. (1990). Determination of termination of sterigmatocystin in fermentation broths by reversed-phase high-performance liquid-chromatography using postcolumn fluorescence enhancement. *J. Chromatogr.* **523**:305–311.
- Nguefack, J., Nguikwie, S. K., Fotio, D., Dongmo, B., Amvam Zollo, P. H., Leth, V., Nkengfack, A. E., and Poll, L. (2007). Fungicidal potential of essential oils and fractions from *Cymbopogon citratus*, *Ocimum gratissimum* and *Thymus vulgaris* to control *Alternaria padwickii* and *Bipolaris oryzae*, two seed-borne fungi of rice (*Oryza Sativa* L.). *J. Essent. Oil Res.* **19**:581–587.
- Nguyen, H., Campi, E., Jackson, M., and Patti, A. F. (2007). Effect of oxidative deterioration on flavour and aroma components of lemon oil. *Food Chem.* **105**:42–47.
- Ni, X. and Streett, D. A. (2005). Modulation of water activity on fungicide effect on *Aspergillus niger* growth in Sabouraud dextrose agar medium. *Lett. Appl. Microbiol.* **41**:428–433.
- Nielsen, K. F. and Thrane, U. (2001). Fast methods for screening of trichothecenes in fungal cultures using gas chromatography-tandem mass spectrometry. *J. Chromatogr. A*. **929**:75–87.
- Olsson, J., Borjesson, T., Lundstedt, T. and Schnurer, J. (2002). Detection and quantification of ochratoxin A and deoxynivalenol in barley grains by GC-MS and electronic nose. *Int. J. Food Microbiol.* **72**:203–214.
- Onji, Y., Aoki, Y., Tani, N., Umebayashi, K., Kitada, Y., and Dohi, Y. (1998). Direct analysis of several *Fusarium* mycotoxins in cereals by capillary gas chromatography mass spectrometry. *J. Chromatogr. A*. **815**:59–65.
- Overy, D. P., Seifert, K. A., Savard, M. E. and Frisvad, J. C. (2003). Spoilage fungi and their mycotoxins in commercially marketed chestnuts. *Int. J. Food Microbiol.* **88**:69–77.
- Ozay, G. and Alperden, I. (1991). Aflatoxin and ochratoxin – a contamination of dried figs (*Ficus carina* L) from the 1988 crop. *Mycotoxin Res.* **7**:85–91.
- Palumbo, J. D., O’Keeffe, T. L., and Mahoney, N. E. (2007). Inhibition of ochratoxin A production and growth of *Aspergillus* species by phenolic antioxidant compounds. *Mycopathol.* **164**:241–248.
- Paniel, N., Radoi, A., and Marty, J. L. (2010). Development of an electrochemical biosensor for the detection of aflatoxin M-1 in milk. *Sensors*. **10**:9439–9448.
- Park, D. L., Njapau, H., and Boutrif, E. (1999). Minimizing risks posed by mycotoxins utilizing the HACCP concept. *FAO* **23**:49–55.
- Paulino de Moraes, M. H. and Luchese, R. H. (2003). Ochratoxin A on green coffee: Influence of harvest and drying processing procedures. *J. Agric. Food Chem.* **51**:5824–5828.
- Pena, R., Alcaraz, M. C., Arce, L., Rios, A., and Valcárcel, M. (2002). Screening of aflatoxins in feed samples using a flow system coupled to capillary electrophoresis. *J. Chromatogr. A*. **967**:303–314.
- Pérez de Obanos, A., González-Peñas, E., and López de Cerain, A. (2005). Influence of roasting and brew preparation on the ochratoxin A content in coffee infusion. *Food Addit. Contam.* **22**:463–471.
- Pestka, J. J., Islam, Z., and Amuzie, C. J. (2008). Immunochemical assessment of deoxynivalenol tissue distribution following oral exposure in the mouse. *Toxicol. Lett.* **178**:83–87.
- Pestka, J. J., Abouzied, M. N., and Sutikno, M. (1995). Immunological assays for mycotoxin detection. *Food Technol.* **49**:120–128.
- Péteri, Z., Téren, J., Vágvölgyi, C., and Varga, J. (2007). Ochratoxin degradation and adsorption caused by astaxanthin-producing yeasts. *Food Microbiol.* **24**:205–210.
- Phillips, R. D., Hayes, A. W., and Berndt, W. O. (1980). High-performance liquid chromatographic analysis of the mycotoxin citrinin and its application to biological fluids. *J. Chromatogr.* **190**:419–427.

- Pose, G., Patriarca, A., Kyanko, V., Pardo, A., and Fernández Pinto, V. (2010). Water activity and temperature effects on mycotoxin production by *Alternaria alternata* on a synthetic tomato medium. *Int. J. Food Microbiol.* **142**:348–353.
- Prakash, B., Shukla, R., Singh, P., Mishra, P. K., Dubey, N. K., and Kharwar, R. N. (2011). Efficacy of chemically characterized *Ocimum gratissimum* L. essential oil as an antioxidant and a safe plant based antimicrobial against fungal and aflatoxin B1 contamination of spices. *Food Res. Int.* **44**:385–390.
- Pronyk, C., Cenkowski, S., and Abramson, D. (2006). Superheated steam reduction of deoxynivalenol in naturally contaminated wheat kernels. *Food Contam.* **17**:789–796.
- Pussemier, L., Pierard, J. Y., Anselme, M., Tangni, E. K., Motte, J. C., and Larondelle, Y. (2006). Development and application of analytical methods for the determination of mycotoxins in organic and conventional wheat. *Food Addit. Contam.* **23**:1208–1218.
- Radová, Z., Holadová, K., and Hajslová, J. (1998). Comparison of two clean-up principles for determination of trichothecenes in grain extract. *J. Chromatogr. A*. **829**:259–267.
- Ratola, N., Abade, E., Simoes, T., Venâncio, A., and Alves, A. (2005). Evolution of ochratoxin A content from must to wine in port wine microvinification. *Anal. Bioanal. Chem.* **382**:405–409.
- Reddy, K. R. N., Reddy, C. S., and Muralidharan, K. (2007). Exploration of ochratoxin A contamination and its management in rice. *Amer. J. Plant Physiol.* **2**:206–213.
- Ringot, D., Lerzy, B., Chaplain, K., Bonhoure, J. P., Auclair, E., and Larondelle, Y. (2007). *In vitro* biosorption of ochratoxin A on the yeast industry by-products: Comparison of isotherm models. *Biores. Technol.* **98**:1812–1821.
- Rosenberg, E., Krska, R., Wissiack, R., Kmetov, V., Josephs, R., Razzazi, E., and Grasserbauer, M. (1998). High-performance liquid chromatography atmospheric-pressure chemical ionization mass spectrometry as a new tool for the determination of the mycotoxin zearalenone in food and feed. *J. Chromatogr. A*. **819**:277–288.
- Ruiz, M. J. (2010). Protective effect of antioxidants against mycotoxins induced cytotoxicity in cells. *Toxicol. Lett.* **196**(Supplement 1):S24.
- Sakuda, S. and Kimura, M. (2010). Toxins of Microorganisms. *Compr. Nat. Prod. II: Chem. Biol.* **4**:411–455.
- Sant'Ana, A. D., Rosenthal, A., and de Massaguer, P. R. (2008). The fate of patulin in apple juice processing: A review. *Food Res. Int.* **41**:441–453.
- Savino, M., Limosani, P., and García-Moruno, E. (2007). Reduction of ochratoxin A contamination in red wines by oak wood fragments. *Amer. J. Enol. Viticult.* **58**:97–101.
- Saxena, J. and Mehrotra, B. S. (1990). The occurrence of mycotoxins in some dry fruits retail marketed in Nainital district of India. *Acta Aliment.* **19**:221–224.
- Schall, N., Simmler-Hübenthal, H., and Süd-Chemie, A. G. (2002). Use of activated layered silicates for the adsorption of mycotoxins. Patent number WO0240150.
- Scott, P. M. (1996). Effects of processing and detoxification treatments on ochratoxin A: Introduction. *Food Addit. Contam.* **13**(Supplement):19–21.
- Scott, P. M. (1985). Report on mycotoxins. *J. Assoc. Off. Anal. Chem.* **68**:242–248.
- Scott, P. M. (1995). Mycotoxin methodology. *Food Addit. Contam.* **12**:395–403.
- Scott, P. M. (1987). Mycotoxins. *J. Assoc. Off. Anal. Chem.* **70**:276–281.
- Scott, P. M., Lawrence, G. A., and Lau, B. P. Y. (2006). Analysis of wines, grape juices and cranberry juices for *Alternaria* toxins. *Mycotoxin Res.* **22**:142–147.
- Scudamore, K. A. and MacDonald, S. J. (1998). A collaborative study of an HPLC method for determination of ochratoxin A in wheat using immunoaffinity column clean-up. *Food Addit. Contam.* **15**:401–410.
- Senyuva, H. Z. and Gilbert, J. (2008). Identification of fumonisin B2, HT-2 toxin, patulin and zearalenone in dried figs by liquid chromatography-time-of-flight mass spectrometry and liquid chromatography-mass spectrometry. *J. Food Prot.* **71**:1500–1504.
- Senyuva, H. Z. and Gilbert, J. (2010). Immunoaffinity column clean-up techniques in food analysis: A review. *J. Chromatogr. B*. **878**:115–132.
- Sharma, M. and Marquez, C. (2001). Determination of aflatoxins in domestic pet foods (dog and cat) using immunoaffinity column and HPLC. *Anim. Feed Sci. Technol.* **93**:109–114.
- Shephard, G. S. (1998). Chromatographic determination of the fumonisin mycotoxins. *J. Chromatogr. A*. **815**:31–39.
- Shephard, G. S. (2008). Impact of mycotoxins on human health in developing countries. *Food Addit. Contam.* **25**:146–151.
- Smart, M. G., Shotwell, O. L., and Caldwell, R. W. (1990). Pathogenesis in *Aspergillus* ear rot of maize — aflatoxin B1 levels in grains around wound-inoculation sites. *Phytopathol.* **80**:1283–1286.
- Spadaro, D., Ciavarella, A., Frati, S., Garibaldi, A., and Gullino, M. L. (2007). Incidence and level of patulin contamination in pure and mixed apple juices marketed in Italy. *Food Control.* **18**:1098–1102.
- Soleas, G. J., Yan, J., and Goldberg, D. M. (2001). Assay of ochratoxin A in wine and beer by high-pressure liquid chromatography photodiode array and gas chromatography mass selective detection. *J. Agric. Food Chem.* **49**:2733–2740.
- Soriano, J. M. and Dragacci, S. (2004). Occurrence of fumonisins in foods. *Food Res. Int.* **37**:985–1000.
- Stakheev, A. A., Ryazantsev, Y. D., Gagkaeva, T. Y., and Zavriev, S. K. (2011). PCR detection of *Fusarium* fungi with similar profiles of the produced mycotoxins. *Food Control.* **22**:462–468.
- Stahr, H. M., Domoto, M., Hyde, W., and Martin, P. J. (1985). Volatilization of trichothecene mycotoxins for analysis. *Microchem. J.* **32**:266–271.
- Stander, M. A., Bornscheuer, U. T., Henke, E. and Steyn, P. S. (2000). Screening of commercial hydrolases for the degradation of ochratoxin A. *J. Agric. Food Chem.* **48**:5736–5739.
- Stanker, L. H., Merrill, P., Scotcher, M. C., and Cheng, L. W. (2008). Development and partial characterization of high-affinity monoclonal antibodies for botulinum toxin type A and their use in analysis of milk by sandwich ELISA. *J. Immunol. Methods.* **336**:1–8.
- Stroka, J., Otterdijk, R. V., and Anklam, E. (2000). Development of a simplified densitometer for the determination of aflatoxins by thin-layer chromatography. *J. Chromatogr. A*. **904**:251–256.
- Sulyok, M., Krska, R., and Schuhmacher, R. (2010). Application of an LC-MS/MS based multi-mycotoxin method for the semi-quantitative determination of mycotoxins occurring in different types of food infected by moulds. *Food Chem.* **119**:408–416.
- Swanson, B. G. (1987). Mycotoxins on fruits and vegetables. *Acta Hort.* **207**:49–61.
- Sydenham, E. W., Shephard, G. S., and Thiel, P. G. (1992). Liquid-chromatographic determination of fumonisin B1, fumonisin B2, and fumonisin B3 in foods and feeds. *J. AOAC Int.* **75**:313–318.
- Tangni, E. K., Simonis, J., Larondelle, Y., and De Meeus, D. A. L. (2005). Decontaminating and detoxifying liquid food media, especially beer, using insoluble vegetable fibers to adsorb mycotoxins. Patent number WO2005007794-A1.
- Tanaka, T., Hasegawa, A., Matsuki, Y., Lee, U. S. and Ueno, Y. (1985). Rapid and Sensitive determination of zearalenone in cereals by high-performance liquid-chromatography with fluorescence detection. *J. Chromatogr. A*. **328**:271–278.
- Tanaka, K., Sago, Y., Zheng, Y., Nakagawa, H., and Kushihiro, M. (2007). Mycotoxins in rice. *Int. J. Food Microbiol.* **119**:59–66.
- Taniwaki, M. H., Hocking, A. D., Pitt, J. I., and Fleet, G. H. (2010). Growth and mycotoxin production by fungi in atmospheres containing 80% carbon dioxide and 20% oxygen. *Int. J. Food Microbiol.* **143**:218–225.
- Tirado, M. C., Clarke, R., Jaykus, L. A., McQuatters-Gollop, A., and Frank, J. M. (2010). Climate change and food safety: A review. *Food Res. Int.* **43**:1745–1765.
- Tokuşoğlu, O., Alpas, H., and Bozoğlu, F. (2010). High hydrostatic pressure effects on mold flora, citrinin mycotoxin, hydroxytyrosol, oleuropein phenolics and antioxidant activity of black table olives. *Innov. Food Sci. Emerg. Technol.* **11**:250–258.
- Tomasevic-Canovic, M., Dakovic, A., Rottinghaus, G. E., Matijašević, S., and Duricic, M. (2003). Surfactant modified zeolites-new efficient adsorbents for mycotoxins. *Microporous Mesoporous Mat.* **61**:173–180.
- Tomlins, K. I., Jewers, K., Coker, R. C., and Nagler, M. J. (1989). Development of rapid methods for the aflatoxin in agricultural products. 2. A bi-directional hplc development method for the detection of low-levels of aflatoxin in maize extracts. *Chromatographia.* **27**:49–52.

- Topcu, A., Bulat, T., Wishah, R., and Boyacı, I. H. (2010). Detoxification of aflatoxin B1 and patulin by *Enterococcus faecium* strains. *Int. J. Food Microbiol.* **139**:202–205.
- Torres, M. R., Sanchis, V., and Ramos, A. J. (1998). Occurrence of fumonisins in Spanish beers analyzed by an enzyme-linked immunosorbent assay method. *Int. J. Food Microbiol.* **39**:139–143.
- Trantham, A. L. and Wilson, D. M. (1984). Fluorometric screening method for citrinin in corn, barley, and peanuts. *J. Assoc. Off. Anal. Chem.* **67**:37–38.
- Trucksess, M. W. and Scott, P. M. (2008). Mycotoxins in botanicals and dried fruits: A review. *Food Addit. Contam.* **25**:181–192.
- Turner, N. W., Subrahmanyam, S., and Piletsky, S. A. (2009). Analytical methods for determination of mycotoxins: A review. *Anal. Chim. Acta.* **632**:168–180.
- Udagawa, S. I. (2005). Fungal spoilage of foods and its risk assessment. *Jpn. J. Med. Mycol.* **46**:11–15.
- Vaclavik, L., Zachariasova, M., Hrbek, V. and Hajslova, J. (2010). Analysis of multiple mycotoxins in cereals under ambient conditions using direct analysis in real time (DART) ionization coupled to high resolution mass spectrometry. *Talanta.* **82**:1950–1957.
- Valenta, H. J. (1998). Chromatographic methods for the determination of ochratoxin A in animal and human tissues and fluids. *J. Chromatogr. A.* **815**:75–92.
- Valero, A., Olivan, S., Marin, S., Sanchis, V., and Ramos, A. J. (2007a). Effect of intra and interspecific interaction on OTA production by *A. niger* in grapes during dehydration. *Food Microbiol.* **24**:254–259.
- Valero, A., Marín, S., Ramos, A. J., and Sanchis, V. (2007b). Effect of pre-harvest fungicides and interacting fungi on *Aspergillus carbonarius* growth and ochratoxin A synthesis in dehydrating grapes. *Lett. Appl. Microbiol.* **45**:194–199.
- van der Westhuizen, L., Shephard, G. S., Rheeder, J. P., Burger, H. M., Gelderblom, W. C. A., Wild, C. P., and Gong, Y. Y. (2011). Optimising sorting and washing of home-grown maize to reduce fumonisin contamination under laboratory-controlled conditions. *Food Control.* **22**:396–400.
- Varga, J. and Kozakiewicz, Z. (2006). Ochratoxin A in grapes and grape-derived products. *Trends Food Sci. Technol.* **17**:72–81.
- Varga, J., Peteri, Z., Tabori, K., Teren, J. and Vagvolgyi, C. (2005). Degradation of ochratoxin A and other mycotoxins by *Rhizopus* isolates. *Int. J. Food Microbiol.* **99**:321–328.
- Varga, J., Rigó, K. and Téren, J. (2000). Degradation of ochratoxin A by *Aspergillus* species. *Int. J. Food Microbiol.* **59**:1–7.
- Varga, J., Kocsubé, S., Suri, K., Szigeti, G., Szekeres, A., Varga, M., Tóth, B. and Bartók, T. (2010). Fumonisin contamination and fumonisin producing black *Aspergilli* in dried vine fruits of different origin. *Int. J. Food Microbiol.* **143**:143–149.
- Vatinno, R., Vuckovic, D., Zamboni, C. G., and Pawliszyn, J. (2008). Automated high-throughput method using solid-phase microextraction-liquid chromatography-tandem mass spectrometry for the determination of ochratoxin A in human urine. *J. Chromatogr. A.* **1201**:215–221.
- Velazhahan, R., Vijayanandraj, S., Vijayasamundeeswari, A., Paranidharan, V., Samiyappan, R., Iwamoto, T., Friebe, B., and Muthukrishnan, S. (2010). Detoxification of aflatoxins by seed extracts of the medicinal plant, *Trachyspermum ammi* (L.) Sprague ex Turrill – Structural analysis and biological toxicity of degradation product of aflatoxin G1. *Food Control.* **21**:719–725.
- Visconti, A. and Pascale, M. (1998). Determination of zearalenone in corn by means of immunoaffinity clean-up and high-performance liquid chromatography with fluorescence detection. *J. Chromatogr. A.* **818**:133–140.
- Visconti, A., Pascale, M., and Centonze, G. (2000). Determination of ochratoxin A in domestic and imported beers in Italy by immunoaffinity clean-up and liquid chromatography. *J. Chromatogr. A.* **888**:321–326.
- Vrabcheva, T., Usleber, E., Dietrich, R., and Martlbauer, E. (1999). Ochratoxin A and citrinin in cereals from Bulgarian villages with a history of Balkan endemic nephropathy. In: Proceedings of 21st Mycotoxin workshop, pp. 257–263. Jena, Germany: Federal Institute of Health Protection and Veterinary Medicine.
- Wang, Y., Chai, T., Lu, G., Quan, C., Duan, H., Yao, M., Zucker, B. A., and Schlenker, G. (2008). Simultaneous detection of airborne Aflatoxin, Ochratoxin and Zearalenone in a poultry house by immunoaffinity clean-up and high-performance liquid chromatography. *Environ. Res.* **107**:139–144.
- Wanigasuriya, K. P., Peiris, H., Ilerperuma, N., John, R. J. P., and Wickremasinghe, R. (2008). Could ochratoxin A in food commodities be the cause of chronic kidney disease in Sri Lanka? *Trans. Roy. Soc. Trop. Med. Hyg.* **102**:726–728.
- Watson, D. H. and Lindsay, D. G. (1982). A critical review of biological methods for the detection of fungal toxins in foods and feedstuffs. *J. Sci. Food Agric.* **33**:59–67.
- Wilkes, J. G. and Sutherland, J. B. (1998). Sample preparation and high-resolution separation of mycotoxins possessing carboxyl groups. *J. Chromatogr. B.* **717**:135–156.
- World Health Organization (WHO). (1998). In: Safety Evaluation of Certain Food Additives and Contaminants, Aflatoxins, Series 40, pp. 359–469. Geneva.
- Xu, B. J., Jia, X. Q., Gu, L. J., and Sung, C. K. (2006). Review on the qualitative and quantitative analysis of the mycotoxin citrinin. *Food Control.* **17**:271–285.
- Xu, Y., Huang, Z. B., He, Q. H., Deng, S. Z., Li, L. S., and Li, Y. P. (2010). Development of an immunochromatographic strip test for the rapid detection of deoxynivalenol in wheat and maize. *Food Chem.* **119**:834–839.
- Yang, C., Wang, Y., Marty, J. L., and Yang, X. R. (2011). Aptamer-based colorimetric biosensing of Ochratoxin A using unmodified gold nanoparticles indicator. *Biosens. Bioelectron.* **26**:2724–2722.
- Yates, I. E. (1986). Bioassay systems and their use in diagnosis of mycotoxicoses. In: Diagnosis of Mycotoxicoses, pp. 333–381. Richards, J. L. and Thurston, J. R. Eds., Kluwer Academic Press, Dordrecht.
- Ye, Y., Zhou, Y. X., Mo, Z. Z., Cheng, W., Yang, S., Wang, X. H., and Chen, F. S. (2010). Rapid detection of aflatoxin B1 on membrane by dot-immunogold filtration assay. *Talanta.* **81**:792–798.
- Young, J. C. and Games, D. E. (1994). Analysis of fusarium mycotoxins by gas-chromatography fourier-transform infrared-spectroscopy. *J. Chromatogr. A.* **663**:211–218.
- Yuan, Y., Zhuang, H., Zhang, T., and Liu, J. (2011). Patulin content in apple products marketed in Northeast China. *Food Control.* **21**:1488–1491.
- Zain, M. E. (2011). Impact of mycotoxins on humans and animals. *J. Saudi Chem. Soc.* **15**:129–144.
- Zamboni, C. G., Monaci, L., and Aresta, A. (2002). Solid-phase microextraction-high performance liquid chromatography and diode array detection for the determination of mycophenolic acid in cheese. *Food Chem.* **78**:249–254.
- Zhou, S. M., Jiang, L. P., Geng, C. Y., Cao, J., and Zhong, L. F. (2010). Patulin-induced oxidative DNA damage and p53 modulation in HepG2 cells. *Toxicol.* **55**:390–395.
- Zimmerli, B. and Dick, R. (1995). Determination of ochratoxin A at the ppt level in human blood, serum, milk and some foodstuffs by high-performance liquid-chromatography with enhanced fluorescence detection and immunoaffinity column cleanup – methodology and swiss data. *J. Chromatogr. B.* **666**:85–99.