



Bioactive peptides from selected latin american food crops – A nutraceutical and molecular approach

Domancar Orona-Tamayo, María Elena Valverde & Octavio Paredes-López

To cite this article: Domancar Orona-Tamayo, María Elena Valverde & Octavio Paredes-López (2018): Bioactive peptides from selected latin american food crops – A nutraceutical and molecular approach, Critical Reviews in Food Science and Nutrition, DOI: [10.1080/10408398.2018.1434480](https://doi.org/10.1080/10408398.2018.1434480)

To link to this article: <https://doi.org/10.1080/10408398.2018.1434480>



Accepted author version posted online: 01 Feb 2018.
Published online: 26 Feb 2018.



Submit your article to this journal [↗](#)



Article views: 101



View related articles [↗](#)



View Crossmark data [↗](#)



Bioactive peptides from selected latin american food crops – A nutraceutical and molecular approach

Domancar Orona-Tamayo, María Elena Valverde, and Octavio Paredes-López

Centro de Investigación y de Estudios Avanzados de Instituto Politécnico Nacional. Km. 9.6 Libramiento Norte Carretera Irapuato-León, Irapuato, Guanajuato, México, CP

ABSTRACT

This review reported an updated survey on the molecular functional properties of bioactive peptides derived from different Latin American ancient grains such as Maize, common Bean, Amaranth, Quinoa and Chia seeds. Seed storage proteins encode in their sequences diverse peptides associated with a wide range of beneficial effects on the human health and the most studied are antihypertensive, anti-cholesterolemic, antioxidant, anti-inflammatory, anticancer, antimicrobial and immunomodulatory properties. Additionally, in the last decades molecular properties have been also used for their characterization to understand their activities and it makes them highly attractive to be incorporated into food formulations and to complement or replace some conventional cereal grains. Due to the nutraceutical effects, today, these seeds are one of the main gastronomic trends of consumption worldwide due to their nutritional benefits and are part of the shopping lists of many people, among them vegetarians, vegans, celiacs or lovers of raw food. These seeds are a legacy of pre-Columbian civilizations reason why in our time they are considered as “Superfoods of the Gods”, “The pre-hispanic superfoods of the future” and “The new golden seeds of the XXI century”.

KEYWORDS

Nutraceutical peptides; Maize; Amaranth; Quinoa; Common bean; Chia

1. Introduction

Nowadays, different scientific areas like chemistry, biology, medicine and pharmacology have focused their research on foods and food products, mainly due to their health benefits. Bioactive substances are known as “components of foods that modulate metabolic processes, provide benefits to health and a positive impact on the function of the body”. The role of a healthy diet for the prevention of diseases is well accepted today, and the border between food and medicine is very small (Pravst 2012; Minkiewicz et al. 2015).

Proteins are important components of foods, and are essential for the maintenance of the body; they are also a source of bioactive peptides which exert biological functions to promote health and prevent some diseases. Specific amino acids sequences encrypted in the protein have health effects, but it is still unclear if they are more potent than the whole proteins. These short amino acid sequences are found in a wide range of proteins and varied foods. They can play an important role as a fragment of the whole protein (epitopes responsible for interactions between proteins and antibodies) or they may be useful after the release by proteolytic enzymes. Peptides with two or three amino acids are very useful because they pass easily through the gastrointestinal tract to the blood. It is possible that each protein is a source of biological active dipeptides and tripeptides, and thus it is possible that all existing proteins are precursors of peptides with certain biological activity. The study of the proteome, or total set of proteins in a food in particular, may be of great importance upon providing

adequate food for the mankind (Minkiewicz et al. 2008; Möller et al. 2008; Minkiewicz et al. 2015; Wang et al. 2015; Cicero, Fogacci, and Colletti 2017).

Bioactive peptides have been known for several years and their identification in plant and animal sources are increasing as result of new scientific methods. These compounds have high nutraceutical potential with great benefits for human health; i.e, on blood pressure and lipid metabolism; as well as anticancer, immunomodulatory, analgesic, antimicrobial, antioxidants, anti-thrombosis, anti-atherosclerotic, opioid and anti-inflammatory; some of them have more than one activity. They are also useful to improve the absorption of minerals (Hartmann and Meisel 2007; Cicero et al. 2011a; Udenigwe and Aluko 2012; Aluko 2015; Cicero and Colletti 2015; Maestri, Marmiroli, and Marmiroli 2016; Bouglé and Bouhallab 2017).

Bioactive peptides are product of the hydrolysis by gastrointestinal digestive enzymes (pepsin, trypsin, and chymotrypsin), or by *in vitro* producers with enzymes, temperature or pH. They usually are composed by two and up to 30 amino acids, with some exceptions, such as Lunasin (peptide derived from cereals and soy proteins) that has 43 amino acids. Other characteristics of the bioactive peptides are the presence of hydrophobic amino acids in their sequences, a positive charge and the resistance to digestion by proteases and peptidases, and a proline C terminal. Small peptides with a dipeptide of proline-proline at their C terminal are the most resistant to degradation by proteases and peptidases of stomach, pancreas or intestine.

Large peptides (with 6 or more) are usually active outside the intestinal epithelium (Galvez et al. 2001; Korhonen and Pihlanto 2003; Panchaud, Affolter, and Kussmann 2012; Perez Espitia et al. 2012; Maestri, Marmiroli, and Marmiroli 2016).

By silico, *in vitro*, *ex vivo* and *in vivo* models have been reported over 3566 different potential bioactive peptides and distinct biological activities; this information has been compiled into a peptide database (BIOPEP database) (Minkiewicz et al. 2008). Bioactive peptides are identified by mass spectrometry, and used for comparative proteomics. At present, there are several databases to analyze food peptides (Table 1) (Dave et al. 2016; Maestri, Marmiroli, and Marmiroli 2016). On the other hand, Gupta et al. (2013) built a Web site to predict the toxicity of peptides

called ToxinPred. This site has been very useful for scientists because it allows to determine the appropriate function of the peptide in human health.

2. Functional properties of bioactive peptides

Bioactive peptides have a wide range of benefits for the human body, mainly due to the composition of amino acids present in each peptide and the different reactivity; the most studied are antihypertensive, anti-cholesterolemic, antioxidant, anti-inflammatory, anticancer, antimicrobial and immunomodulatory properties. Beside, some peptides have multimodal activities and can exert more than one beneficial effect (Meisel 2004; Hartmann and Meisel 2007; Santiago-López et al. 2016).

Table 1. Bioactive peptides database.

Name of Database	Description of database	Link
ACEpepDB	Peptides inhibiting Angiotensin I converting enzyme	http://www.re3data.org/repository/r3d100010972
AHTPDB	Antihypertensive peptides	http://crdd.osdd.net/raghava/ahtpdb/
AllergenOnline Celiac Disease	Database of celiac-toxic peptides and proteins	http://www.allergenonline.org/celiachome.shtml
AMPer	Antimicrobial peptidess	https://omictools.com/amper-tool
ANTISTAPHYBASE	Antimicrobial peptidess	http://www.antistaphybase.com
ArachnoServer	Toxic peptides and proteins from spider venoms	http://www.arachnoserver.org/mainMenu.html
ATDB	Animal Toxin Database	http://bactpepdb.rpbs.univ-paris-diderot.fr/cgi-bin/links.pl
APD	Antimicrobial and anticancer peptides	http://aps.unmc.edu/AP/main.php
AVPdb	Antiviral peptides	http://crdd.osdd.net/servers/avpdb/
BactPepDB	Predicted bacterial peptides	http://bactpepdb.rpbs.univ-paris-diderot.fr/cgi-bin/home.pl
BACTIBASE	Antibacterial peptides	http://bactibase.hammamilab.org/main.php
BAGEL2	Antibacterial peptides	http://bagel2.molgenrug.nl
BAGEL3	Database of antibacterial peptides (bacteriocins)	http://bagel.molgenrug.nl
BaAMPs	Antimicrobial peptidess against microbial biofilms	http://www.baamps.it
Brainpeps	Blood-Brain peptides	http://brainpeps.ugent.be
BIOPEP	Biologically active peptides	http://www.uwm.edu.pl/biochemia/index.php/pl/biopep
CAMP	Antimicrobial peptides and proteins	http://www.camp.bicnirrh.res.in
CPPSite	Cellular peptides	http://crdd.osdd.net/raghava/cppsite/
CancerPP	Anticancer peptides	http://crdd.osdd.net/raghava/cancerppd/
Cybase	Cyclic proteins and peptides with various bioactivities	http://www.cybase.org.au
DADP	Anuran defense peptides	http://split4.pmfst.hr/dadp/
DAMPD	Antimicrobial peptides.	http://apps.sanbi.ac.za/dampd/
Defensins knowledgebase	Database of defensins from antimicrobial peptides	http://defensins.bii.a-star.edu.sg
Effective DB	Database of predicted secreted bacterial peptides	http://effectors.org
EROP-Moscow	Regulatory oligopeptides	http://erop.inbi.ras.ru
HIPdb	HIV inhibiting peptides	http://crdd.osdd.net/servers/hipdb/
MHCBN	Major Histocompatibility Complex (MHC) Binding, Non-binding peptides and T-cell epitopes	http://crdd.osdd.net/raghava/mhcbn/
HMRBase	Hormones and peptides	http://crdd.osdd.net/raghava/hmrbase/
NeuroPep	Neuropeptides	http://isyslab.info/NeuroPep/
Norine	Nonribosomally synthesized bioactive peptides.	http://bioinfo.lifl.fr/norine/
ParaPep	Antiparasitic peptides	http://crdd.osdd.net/raghava/parapep/
PepBank	Peptides shorter than 20 residues	http://pepbank.mgh.harvard.edu
PepBind	Protein-peptide interactions	https://test5.bicpu.edu.in
Peptaibols	Antimicrobial peptides	http://peptaibol.cryst.bbk.ac.uk/home.shtml
PeptideAtlas	Compendium of peptides from multi-organisms	http://www.peptideatlas.org
PeptideDB	Compendium of peptides from multi-organisms	http://www.peptides.be
PeptideLocator	Identified bioactive peptides within protein sequences	http://bioware.ucd.ie/~compass/biowareweb/Serv_er_pages/biopred.php
PhytAMP	Antimicrobial Plant peptides	http://phytamp.hammamilab.org/main.php
Quorumpeps	Resource of quorum sensing signalling peptides	http://quorumpeps.ugent.be
SMS	A database to study the structural plasticity of short peptide fragments in non-redundant proteins.	http://cluster.physics.iisc.ernet.in/sms/
SPdb	A signal peptide database containing signal sequences of archae, prokaryotes and eukaryotes	http://proline.bic.nus.edu.sg/spdb/
SwePep	A database of biologically active peptides including sophisticated computing tools for peptide mass spectra interpretation.	http://www.swepep.org
YADAMP	Database of Antimicrobial Peptides	http://yadamp.unisa.it
THPdb	Therapeutic peptides	http://crdd.osdd.net/raghava/thpdb/
ToxinPred TumorHoPe	<i>In silico</i> method to predict and design toxic/non-toxic peptides Tumor-recognizing peptides	http://crdd.osdd.net/raghava/toxinpred/ http://crdd.osdd.net/raghava/tumorhope/

2.1. Anti-hypertensive

Hypertension is the major risk factor for the myocardial infarction, heart failure, atherosclerosis and kidney disease. The use of nutraceuticals in the treatment or prevention of this condition is of great impact to reduce the side effects of the antihypertensive drugs (Sirtori, Arnoldi, and Cicero 2015; Borghi and Cicero 2017).

The main mechanism to regulate blood pressure is the renin-angiotensin aldosterone system (ACE), which plays an important role in the control and regulation of blood pressure and salt balance. Antihypertensive drugs, like Captopril, Lisinopril and Enalapril, inhibit the renin-angiotensin system but produce some side effects (sleep apnea, dry cough, angioedema, etc). Some bioactive peptides show competitive and/or non-competitive ACE inhibition, which is responsible of the conversion of angiotensin I in angiotensin II. Angiotensin-I catalyzes the conversion to the vasoconstrictor angiotensin II, which increases the peripheral vascular resistance, inducing hypertensive action. ACE also determines the cleavage and inactivation of bradykinin, a vasodilator peptide. The increased of protein intake has been associated with a reduced risk of increased blood pressure and coronary heart diseases; food-derived peptides have been identified as potential antihypertensive agents. Peptides with ACE inhibitory activity are small and have hydrophobic amino acids (W, Y, F, and P) (Table 2) in C-terminal which determine the affinity to the active site of the enzyme. In addition, the effectiveness of these peptides depends on their resistance to degradation, to the absorption into the blood stream and to amino acid sequence. Many peptides have been identified with this activity in protein of plants, animals and microorganisms, and the beneficial effect on blood pressure and vascular system has been confirmed in animal and human models (Foltz et al. 2007; Aluko 2015; Mora and Hayes 2015; Cicero et al. 2016; Bhat, Kumar, and Bhat 2017; Bouglé and Bouhallab 2017).

The richest source of protein and bioactive peptides is milk, which have been identified as antihypertensive agents; IPP and VPP are the best-characterized in human studies (Table 2). Cicero et al. (2011b) reported that in Asian patients the most efficient dose of peptides to decrease blood pressure ranged between 3.07 mg/d (1.60 mg of VPP and 1.47 mg of IPP) and 52.5 mg/d (30 mg of VPP and 22.5 mg of IPP), and was independent of age, by addition, the systolic and diastolic pressure decreased around 5 and 2 mm Hg, respectively, without negative impact on health; however, the European Food Safety Authorities (EFSA) rejected this health benefit (Jauhiainen et al. 2005; Sano et al. 2005; Panel 2012; Bhat, Kumar, and Bhat 2017). This kind of antihypertensive peptides can be found now on the market. Serum proteins and casein are bioactive peptides with significant antihypertensive effects in normal-tense/pre-retinopathy obese subjects. Numerous studies in humans and animals show that milk, whey and casein peptides are able to inhibit ACE and may positively affect blood pressure. Peptides sequences from some animals proteins are shown in Table 2 (Bhat, Kumar, and Bhat 2015; Nongonierma and FitzGerald 2015).

The search for ACE inhibitory peptides has been increasing in the last years; Nakashima et al. (2002) identified two in

Table 2. Bioactive peptides from various animal proteins with antihypertensive/ACE-inhibition activity.

Protein source	Peptide/sequence
Casein	FP
	YP
	IPP
	IPA
	GKP
	LKP
	VPP
	FFVAP
	LKPNM
	VYPFPG
	YKVPQL
	EMPFPK
	TTMPLW
	FALPQY
	YQEPVL
	KKYKVPQ
	AVYPYQR
	FALPQYLK
	YPVEPFTE
	RPKHPIKHQ
	GPVRGPFPIIV
	FFVAPFPEVFGK
Whey proteins	YP
	LLF
	IPA
	WLAHK
	ALPMHIRY
Porcine skeletal muscle Myosin	IWWNP
	TNP
	ITTNP
	MNPPK
	KRVITY
	KRQKYDI
	RMLGQTPT
	VKKVLGNP
Gelatin (bovine skin)	GPL
	GPV
Chicken	IKW
	LKP
Egg	GFHyGTHyGLHyGF
	LW
	YPI
	SALAM
	RADHP
	RADHPF
	RADHPFL
	ERKIKVYL
	FRADHPFL
	RADHPFLT
Fish	FFGRCVSP
	YAEERYPIL
Beef	VY
	VLAQYK
	LAQYWK

The amino acid nomenclature: A: Alanine; C: Cysteine; D: Aspartic acid; E: Glutamic acid; F: Phenylalanine; G: Glycine; H: Histidine; I: Isoleucine; K: Lysine; L: Leucine; M: Methionine; N: Asparagine; P: Proline; Q: Glutamine; R: Arginine; S: Serine; T: Threonine; V: Valine; W: tryptophan; Y: Tyrosine; B: either of D or N; Z: either of E or Q.

porcine skeletal muscle, MNPPK and IWWNP, and Katayama et al. (2003) found RMLGQTPT. In porcine myosin, Katayama et al. (2003; 2007) identified VKKVLGNP, KRVITY and KRQKIDI, and Arihara et al. (2001) described ITTNP and NNPPK. Jang and Lee (2005) identified VLAQYK in beef. Kawasaki et al. (2000) observed a reduction of blood pressure in patients with a daily ingestion of 3 mg of sardine dipeptide VY, 9 and 5 mm Hg in systolic and diastolic pressure, respectively.

Saiga et al. (2003) reported the strongest ACE inhibitory activity with GFHYGTHYGLHYGF in chicken breast meat, and Cheng et al. (2008) (Cheng et al. 2008) in chicken leg bones. Also, it has been reported that ovokinin, an octapeptide (FRAHPFL) isolated from ovalbumin, significantly lowered systolic blood pressure in spontaneously hypertensive rats. Some analogs of this peptide also exhibited antihypertensive properties; with the replacement of some amino acids in the peptide enhanced 100-fold antihypertensive activity (Matoba et al. 2001; Yamada et al. 2002). Miguel et al. (2005) reported RADHPFL, YAEERYPIL and IVF peptides in crude egg white with potent ACE-inhibitory and antihypertensive activities. RVPSL peptide from egg white showed the highest level of ACE inhibitory activity, and QIGLF potential ACE inhibition. Hydrolysate of egg yolk also have ACE inhibitory activity, peptides with 1 kDa or less reduced the hypertension in spontaneously hypertensive rats after 12 weeks (Yoshii et al. 2001; Liu et al. 2010; Yu et al. 2011).

Peptides with antihypertensive activity have been detected in marine species (LKP, IKP, LRP, MVGSAPGVL, LGPLGHQ and AHIII). Also, plants provide antihypertensive peptides, i.e., peptides extracted from oats and barley (IVY and INP) showed strong inhibitory action on ACE. It is important to distinguish between the effect of plant proteins and other components, like isoflavones, phenols compounds and carotenoids on blood pressure levels (Motoi and Kodama 2003; Altorf-van der Kuil et al. 2010; Malaguti et al. 2014; Cheung, Ng, and Wong 2015; Gangopadhyay et al. 2015).

In general, peptides derived from any type of food have significant antihypertensive effect in humans, which has been confirmed by clinical trials in normotensive and pre-hypertensive patients (Cicero et al. 2016).

2.2. Anti-cholesterolemic

The bioactive peptides with clinical evidence of cholesterol lowering effect are found in soybean, lupin and milk proteins; a decrease in total cholesterol (2%), LDL (3%), HDL (3%) and triacylglycerol (4%) has been reported (Lammi et al. 2014; Tokede et al. 2015; Butteiger et al. 2016). Lunasin (a peptide of 43 amino acid) is found in soybean, barley, rye, and wheat and has shown a cholesterol-lowering activity and increasing production of LDL receptors; however, further studies are necessary to assess its efficacy and safety in clinical practice. Protein conglutin, the protein from lupin, lowered effectively LDL-cholesterol in a rat model, and increases LDL receptors in a hepatoma G2 cell line. Besides, peptides from cowpea inhibited cholesterol synthesis (Hernandez-Ledesma, Hsieh, and Ben 2009a; Hernandez-Ledesma, Hsieh, and O De Lumen 2013; Lule et al. 2015; Marques et al. 2015; Sirtori, Arnoldi, and Cicero 2015).

The main mechanism to reduce the serum cholesterol levels is the inhibition of the hydroxymethylglutaryl-CoA (HMG-CoA) reductase and the increase in fecal excretion of bile salts. β -lactotensin is a peptide of milk that also lowers cholesterol; it significantly reduced cholesterol in mice at a dose of 100 mg/kg. It probably acts on dopamine D1 and neurotensin NTS2 receptor, which increase the synthesis of bile acids. In summary, the mechanisms of action of the lipid-lowering peptides

are mainly by inhibition of the production of endogenous cholesterol and the fecal excretion of exogenous cholesterol (Yamauchi, Ohinata, and Yoshikawa 2003; Lammi et al. 2014; Cicero et al. 2016).

2.3. Antioxidant

Reactive oxygen species (ROS) and reactive nitrogen species (RNS) are free radicals produced by endogenous oxidation-reduction (REDOX) reactions, i.e., through normal respiration in aerobic organisms. ROS/RNS are useful to facilitate biological processes such as gene expression, cell proliferation, angiogenesis, programmed cell death, senescence and defense against infections. Under normal conditions, the body antioxidant defense systems can remove reactive species; however, in certain circumstances the endogenous defense system fails, resulting in oxidative stress. When accumulated ROS/RNS exceeds the threshold of the body, cellular damage is induced, which results in macromolecular damage such as atherosclerosis, arthritis, hypertension, cancer, diabetes, neurodegenerative disorders, heart disease, stroke, and aging (Castro and Freeman 2001; Hancock, Desikan, and Neill 2001; Evans et al. 2003; Johansen et al. 2005; Valko et al. 2007; Dowling and Simmons 2009; Scherz-Shouval and Elazar 2011; Liu et al. 2016).

Once the redox balance is interrupted, it is necessary to reinforce the intake of antioxidants from food sources. Vitamin C, herbal extracts and artificial antioxidants have been marketed for a long time as natural antioxidants. However, amino acids and peptides could also provide antioxidant effect to protect cells against oxidative damage. Peptides with antioxidant capacity are safe, healthy, stable and without dangerous immunological reactions (Shahidi 2000; Sarmadi and Ismail 2010).

Antioxidants peptides comprise 4 to 16 amino acids and approximately 400 – 2000 Da molecular weight. Low molecular weight peptides (<3 kDa) present better antioxidant potential than high ones. Small molecules can easily cross the gastrointestinal barrier and enter to peripheral tissue; they can be digested by gastrointestinal proteases, and their antioxidant capacity increases because they can be absorbed more easily (Rubas and Grass 1991; Roberts et al. 1999; Wang et al. 2008; Liu et al. 2010).

Peptides with strong antioxidant activities have been found in crude albumin proteins. The egg white peptide TAQQRT-PIL, with ACE-inhibitory activity, also exhibited a high radical-scavenging activity (Davalos et al. 2004; Bhat, Kumar, and Bhat 2017).

2.4. Anti-inflammatory

Inflammation is an important defense mechanism of the body to protect it for infection and injury. The body cells respond to different stimulus by producing inflammatory mediators (eicosanoids, vasoactive amines, vasoactive peptides, lipid mediators, cytokines, chemokines and proteolytic enzymes), which interact with all cells to release inflammatory response. Inflammation is related to a wide variety of diseases, including asthma, ulcerative colitis, Crohn's disease, cancer, type 2 diabetes, cardiovascular diseases (atherosclerosis), and central nervous

system-related disorders (Parkinson's disease and cognitive impairment) (Laveti et al. 2013).

Pharmacological therapies against chronic inflammatory diseases often produce side effects; hence, scientists suggest that the intake of food compounds than can modulate chronic inflammatory conditions. *In vitro* cell and *in vivo* animal studies indicate that peptides from food proteins are safe, without any deleterious side effects and are well recognized by their enormous potential as anti-inflammatory agents (Kovacs-Nolan et al. 2012; Nakamura et al. 2013; Kobayashi et al. 2015; Minihihane et al. 2015).

The anti-inflammatory capacity of bioactive peptides is not completely understood. It could be related to the modulation of transcription factors and the inhibition of expression of pro-inflammatory cytokines and chemokines. These peptides are found in different food sources such as bovine milk, egg, fish, and soybean (Cicero et al. 2016; Majumder, Mine, and Wu 2016). Holmer-Jensen et al. (2011) reported a reduction in postprandial inflammation in obese and non-diabetic subjects using milk proteins. Majumder et al. (2013) found a tripeptide from ovotransferrin, albumen of eggs, with anti-inflammatory activity. Peptides derived from soybean, bean and milk have also shown anti-inflammatory effects on intestinal inflammation. Lunasin, a soybean peptide, has also shown anti-inflammatory properties. The most studied peptides with this activity are IPP and VPP, from different sources (Cam and de Mejia 2012; Majumder, Mine, and Wu 2016).

2.5. Anti-carcinogenic

Cancer occurs when some cells in the body ignore growth inhibitory signals, avoid cell death, replicate without limit, sustain angiogenesis and invade tissues, at the end the system fails to eliminate cancer cells. It is one of the major causes of mortality around the world. Radiation and chemotherapy are the principal methods for cancer management; however, both therapies are highly toxic and with some severe side effects. Nowadays, the development of novel tumor-targeted to eliminate tumor cells, in a specific and effective way, with low toxicity, is considered of great importance (Kang et al. 2012; Ortiz-Martinez, Winkler, and García-Lara 2014). On the other hand, the relationship between the risk increasing of colorectal cancer and eating habits, physical activity and obesity is well established (Wolin et al. 2010). Currently, the pharmaceutical industry is developing novel peptides that are more precise to kill cancer cells. Small peptides possess favorable advantages like high affinity, strong specificity, low toxicity, distinct tissue distribution patterns, and good solubility properties, which could ensure high uptake into target tissues and rapid clearance from the blood and non-target tissues. The explorations and modifications of anticancer peptides will play important roles in cancer prevention and treatment (Shadidi and Sioud 2003; Bhutia and Maiti 2008; Chen and Conti 2010; Lee, Xie, and Chen 2010; Segura-Campos et al. 2011; Wu et al. 2014).

Bioactive peptides derived from plants, animals, milk, eggs, marine organisms and microorganism, could be utilized against some cancer types due to their effect on one or more carcinogenesis molecular pathways. Bioactive peptides present different mechanism of action, i.e., apoptosis, blockage of

intermediate tumor generation and regulation of immune system, inhibition of cell migration and tumor angiogenesis, antioxidant activity, inhibition of cell proliferation, tubulin structure disorganization and cytotoxicity (Schweizer 2009; Thundimadathil 2012; Tyagi et al. 2014; Blanco-Míguez et al. 2016).

In vitro studies on lunasin have confirmed anti-neoplastic effects in breast, skin, colon, prostate, leukemia and lymphoma cell lines. Besides, the pentapeptide EGRPA of rice bran, at a dose of 600–700 $\mu\text{g/mL}$, inhibited the 84% of the growth of colon cancer cells (Caco-2 cell line and human colorectal adeno-carcinoma, HCT-116), 80% in breast cancer (MCF-7 and MDA-MB-231) and 84% in liver cancer (HepG-2 cells) (Hsieh, Hernández-Ledesma, and De Lumen 2010; Kannan et al. 2010; Diaz and de Mejia 2011; Chang et al. 2014; Hernández-Ledesma and Hsieh 2017).

Bowman-Birk inhibitor (BBI), a bioactive peptide of legume seeds, is a source of protease inhibitors and has preventive effect against prostate, breast and colon cancer, and could be an effective suppressor of carcinogenesis. The use of BBI in patients with benign prostatic hyperplasia was approved by the Food and Drug Administration (FDA); patients show tolerance for a prolonged period of time (Armstrong et al. 2000; Malkowicz et al. 2001; Park, Jeong, and De Lumen 2005).

Lectins, plant derived peptide, are able to recognize specific carbohydrates produced by malignant cells or tissues. Lectins show antiproliferative and cytotoxic effects by inhibiting colony formation on cervical and human colon carcinoma cell lines. They also showed, in a mouse model, an immunomodulatory effect that delayed the development of colon cancer and improved the survival of mice with leukemia cells (Ma et al. 2008; Seifert et al. 2008; Valadez-Vega et al. 2011).

Lactoferrin, a whey protein, inhibits the growth of nasopharyngeal carcinoma and breast (MDA-MB-231) cancer cells. These effects were confirmed with an *in vivo* model, in addition this protein can inhibit spontaneous growth of melanoma cells and suppresses metastasis of liver and lung lymphoma. Lactoferrin acts against different types of cancer cell lines, by inducing apoptosis, modulating gene expression, preventing angiogenesis and disorderly growth of the cells, and stopping the cell cycle. Another whey protein, α -lactalbumin, activated apoptosis *in vitro* chemoprevention. In milk two peptides derived from casein β -casomorphin 5 showed the capacity to arrests the cell cycle in breast cancer cells and induce apoptosis in HT-29 and AZ-97 (Hatzoglou et al. 1996; Yoo et al. 1997; Damiens et al. 1999; Varadhachary et al. 2004; Fast et al. 2005; de Mejia and Dia 2010; Perego et al. 2012; Sun et al. 2012; Deng et al. 2013).

Some egg-derived peptides like hydrolysates from egg yolk protein, lysozyme and ovomucin, have shown anticancer properties. *In vivo* studies demonstrate that hydrolysates from egg yolk protein inhibit the proliferation of tumor cells in colorectal cancer; lysozyme reduces the formation of lung metastases of B16 melanoma and ovomucin inhibits tumor growth (Watanabe et al. 1998; Azuma et al. 2000).

Some bioactive peptides derived from different marine species, induce antiproliferative activity against different cancer types. *In vitro* assays showed positive effect of tuna dark muscle peptides on human breast cancer cells; also besides, peptides

from squid gelatin had cytotoxic activity on human breast carcinoma (MCF-7) and glioma cell lines, and peptide reniochalstatin E from the sponge *Reniochalina stalagmites* showed cytotoxic activity toward several tumor cell lines. On the other hand, peptides called BEPT II, BEPT II-1, Dolastatins and Pardaixin induced apoptosis toward different types of cancer (Bai, Petit, and Hamel 1990; Anderson et al. 1996; Alemán et al. 2011b; Hsu, Li-Chan, and Jao 2011; Ma et al. 2013; Ting et al. 2014; Zhan et al. 2014). Additionally, *in vivo* studies confirmed that the polypeptide P2 of *Arca subcrenata* has an antiproliferative action on HeLa and HT-29 cells in S-180 tumor-bearing mice, and peptides from oyster (*Crassostrea gigas*) showed an antiproliferative effect against sarcoma-S180 in mice (Wang et al. 2010; Hu et al. 2012).

Bioactive peptides possess outstanding characteristics to produce anti-tumor drugs, like high selectivity, wide range of targets and low toxicity; however, they present some problems, such a poor stability, low membrane permeability and susceptibility to proteolytic digestion. In order to avoid these difficulties, some modifications have been developed, i.e., amino acid substitution, structural fusion of functional peptides and conjugation with chemotherapeutic drugs (Vlieghe et al. 2010; Zhang, Eden, and Chen 2012; Craik et al. 2013; Wu et al. 2014).

2.6. Anti-microbial activity and immunomodulatory effect

The immune system represents the first line of defense against infections, and its optimum performance is vital to provide protection. Innate immunity is mainly represented by different cells, including dendritic, macrophages and natural killer (NK); some immune stimulants increase the activation of these cells (Akira, Uematsu, and Takeuchi 2006).

Over the years, anti-microbial plant derived peptides have become interesting tools for the production of new antibiotics for the treatment of different infections. This kind of peptides are obtained from various sources and are the first line of defense in animals. These peptides contained different amino acids positively charged such as arginine, lysine, histidine, and an excess of glutamic and aspartic acids negatively charged. They usually disintegrate the cell membrane by altering its permeability (Kitts and Weiler 2003; Otvos Jr 2016).

Lactoferrin, protein fraction from human and bovine milk, tuna and sardine meat, barley, and pumpkin, is produced in Japan. It is considered a potent anti-microbial and also is used in the prevention of infections and the improvement of bowel flora of milk-fed infants and in other health conditions (Bellamy et al. 1992; Nayak et al. 1999).

Some plants produce anti-microbial peptides involved in defense against infection or natural invaders. Thionins are basic peptides identified in wheat, barley, rye and oats. They are homologous polypeptides with 43–46 amino acid, rich in cysteine and lysine residues. Thionins have anti-microbial activity against plant pathogenic fungi and bacteria (García-Olmedo et al. 1998; Kitts and Weiler 2003). Small peptides from maize kernels showed anti-microbial properties; they were similar to thionins and one was purified, MBP-1, with 33 residues and 4127 Da (Duvick et al. 1992).

Immunomodulatory peptides do not target the pathogen directly; however, they can enhance host defense mechanisms

by activating immune cell functions, like lymphocyte proliferation, natural killer cell activity, antibody synthesis and cytokine regulation; besides, they might reduce allergic reactions in atopic humans and enhance mucosal immunity in the gastrointestinal tract (Korhonen and Pihlanto 2003; Fitzgerald and Murray 2006; McCann et al. 2006; Hartmann and Meisel 2007; Mookherjee and Hancock 2007). For example, some casein peptides stimulated phagocytes and the production of IgG in lymphocytes, as well as the proliferation of T lymphocytes. On the other hand, fish peptides presented immunomodulatory effects in animals, and oyster hydrolysates increased the activity of natural killer cells. Peptides obtained from rice and soybean proteins activated the defense systems, and stimulated anion superoxide (ROS), which activated non-specific immune defense systems (Cheung et al. 1980; Lahov and Regelson 1996; Hata, Higashiyama, and Otani 1998; Kitts and Weiler 2003; Yang et al. 2009; Wang et al. 2010; Cheung, Ng, and Wong 2015).

Hypothetically, immunopeptides interact on the opioid receptors that regulate the peripheral immune system. In addition, peptides with arginine, tryptophan and phosphoserine in the N or C terminal residues exert high reactivity in immune cells. In addition, glutamine residues involved in the recognition of opioid receptors are located on the surface of immune cells (Elitsur and Luk 1991; Mookherjee and Hancock 2007; Haque, Chand, and Kapila 2008; Santiago-López et al. 2016).

3. Maize

Maize (*Zea mays* L.) is one of the most important grains originated in the New World. It was the main food for different pre-Columbian societies in America, and was domesticated approximately 6000–10,000 years ago. In the Mayan culture, maize was as important that was used in spiritual and religious rituals (Piperno and Flannery 2001; Matsuoka et al. 2002) (<http://maize.teacherfriendlyguide.org/index.php/what-is-maize/cultural-uses-of-maize>). Maize is native to Central Mexico and several studies indicate that annual teosinte (*Zea mays* ssp. *parviglumis*) was its predecessor in at least two separate domestication sites in Mexico. Isoenzymes, molecular and cytological data, indicate that wild maize is native to the Balsas River Valley in Michoacan and Guerrero, Mexico. However, another theory argues that it is native to Tehuacan, Puebla, Mexico (Piperno and Flannery 2001; Matsuoka et al. 2002; Wu et al. 2008; Somerville et al. 2016; Orona-Tamayo and Paredes-López 2017a; Orona-Tamayo, Valverde, and Paredes-López 2017b).

3.1. Nutritional composition

Maize is one of the world's three main cereal crops and is consumed by millions of people specially from the developing countries. The main parts of the maize kernel are the endosperm and germ; both tissues contain most of the starch (62%) and oil (3.8%). The protein content is 6 to 12% and almost 75 to 80% is in the endosperm, the rest is found in the germ and bran. The proportion of endosperm protein fractions is 3 to 12% for albumins, 3 to 10% for globulins, 38 to 60% for prolamins or zeins, and 25 to 40% for glutelins. Zeins are well balanced with glutamine (20%), leucine (20%), alanine (14%),

proline (9%) and serine (7%); but lack of two important essential amino acids, lysine and tryptophan. In 1960, efforts were made to increase the nutritional quality of maize protein. The high proportion of zeins is related to the poor quality of protein; hence research studies have focused on the reduction of this fraction in order to increase other fractions with more lysine and tryptophan content (Dierks-Ventling 1981; Larkins et al. 1984; Vasal 2000; Shukla and Cheryan 2001; Rascón-Cruz et al. 2004).

3.2. Bioactive peptides

Zein peptide LPP shows strong ACE inhibitory activities, and have applications in the medical field for lowering blood pressure (Table 3) (García et al. 2013; Corradini et al. 2014). Maruyama et al. (1989) reported that the native sequence VHLPPP from γ -zein had stronger antihypertensive effect in spontaneously hypertensive rats than synthesized peptides. Similarly, Miyoshi et al. (1991) found *in vitro* that antihypertensive tripeptides isolated from α -zein hydrolysates with thermolysin (LGP, LSP and LRP) showed similar ACE inhibitory activities; however, the LRP peptide showed the strongest inhibition effect on blood pressure in spontaneously hypertensive rats. (Puchalska, Marina, and García 2012; 2013) found significant differences in the content of these three Zein peptides in diverse maize lines (Table 3). Ren et al. (2014) found that sweeping frequency ultrasound (SFU) treatment of zein increased hydrolysis, and the hydrolysates presented a higher inhibition effect on ACE. It shows that the use of this methodology promotes the release of ACE-inhibitory peptides from zein by altering the secondary structure and loosening the protein.

Kong and Xiong (2006) reported that hydrolyzed zein is an excellent antioxidant and also possesses a strong chelating activity against Cu^{2+} , as well as strong reducing power. Although intact zein displayed antioxidant effect, it was far less potent than the hydrolyzed zein peptides. Tang et al. (2009) separated zein hydrolysates by gel filtration, ultrafiltration, and reversed-phase HPLC, and concluded that peptides of <1 kDa exhibited a strong DPPH, ABTS and O_2^- scavenging activity. They also purified and identified two peptides (YA, and LMCH) exhibiting high antioxidant reactivity.

Several small peptides with antimicrobial properties have been isolated from maize kernels. Duvick et al. (1992) found a 4.1 kDa peptide with antimicrobial properties, named MBP-1, similar to thionins. MBP-1 inhibits spore germination and the

hyphal elongation of several plant pathogenic fungi (*Fusarium montiliforme* and *F. graminearum*) and maize phyto bacteria such as *Clavivacter michiganense*. Wound infection is a common problem in hospitals and it is typically caused by the antibiotic-resistant *Staphylococcus aureus*, which is the major pathogen for skin and soft tissues infections. Salouti et al. (2016) investigated the antibacterial effect of MBP-1 peptide in combination with silver nanoparticles on infected wounds caused by *S. aureus* in animal models, and found that this combination increased wound healing rates and decreased colonization of *S. aureus*. Sousa et al. (2016) identified antibacterial activity in MBP-1 and concluded that by replacing cysteine and tryptophan can abolish this activity against gram negative and positive bacteria.

3.3. Genetic engineering to improve the functional properties of peptides

In maize seeds, different studies have been performed to elucidate genes encoding storage proteins. These genes are under strict control and are expressed in a tissue specific manner. The zein protein fraction is loaded with a high amount of bioactive peptides, and is a great blank to transform different plants (Yamamoto, Ejiri, and Mizuno 2003; Sarmadi and Ismail 2010). From the middle of the 1960s, great efforts were performed to improve the nutritional quality of the maize endosperm proteins with the mutant lines of opaque-2 and floury-2. These mutants alter the amino acids compositions of maize endosperm proteins, two-fold increase in levels of lysine and tryptophan; and other amino acids such as histidine, arginine, aspartic acid, and glycine are also increased. These mutations affect some agronomic performance including yield and consumer aspects of the kernel. Subsequently, several other mutant genes in maize have been discovered such as opaque-7, opaque-6 and floury-3; these mutants reduce the zein content and increase lysine and tryptophan levels due to synthesis of non-zein proteins such as glutelins, globulins and albumins. These studies were focused to improve the nutritional quality resulting in the Quality Protein Maize, hybrids that gained at the beginning certain popularity in development countries (Huang et al. 2004).

In the 80s, the high potential to produce transgenic plants with the expression of chimeric genes transferred into plants cells via *A. tumefaciens*. This technology led to researchers to obtain transgenic plants with a high level of essential amino

Table 3. Bioactive peptides from maize (*Zea mays*).

Bioactive function	Protein source	Bioactive Peptide/Peptide sequence	Reference
Antihypertensive	Zein (Prolamin)	LPP	Corradini et al. 2014; García et al. 2013
	γ -Zein (Prolamin)	VHLPPP	
	α -Zein (Prolamin)	FNQ, FSQ, LAY, LNPA, LSPA, AF, LPN, FYQQ, YLEPA, FY, LSPY, LNSPAY, LLP, LF, FLP, LVP, FLPP, LQQ, YSQQQQ, FNQ, ASY, AY, AYPQQ, LSPQSY, LSLP	Maruyama et al. 1989
	α -Zein (Prolamin)	LQP, LSP, LRP	Yano, Suzuki, and Funatsu 1996
Antioxidant	Zein	YA, LMCH	Miyoshi et al. 1991; Puchalska, Marina, and García 2012; Puchalska et al., 2013
Antimicrobial	Total protein	RSGRGECRRQCLRRHEGQPWETQECMRRCRRR	Tang et al. 2009
			Duvick et al. 1992; Sousa et al. 2016

See Table 2 for the amino acid nomenclature used in this table.

acids with the introduction and expression of maize zein gen. Matzke et al. (1984), reported the first evidence for transcription of a genomic clone of maize zein introduced with T-DNA into sunflower cells; they found that the zein mRNA made in sunflower can be translated *in vitro* to yield an immunoprecipitable protein with expected molecular weight as native zein in maize. Moreover, transformed tobacco plants were developed containing maize gene encoding a zein protein using a modified Ti plasmid vector; zein proteins were detected in roots, leaves and endosperm tissues. In addition, constructed chimeric genes containing a maize α -zein coding sequence and a β -phaseolin gene promoter and were inserted into genome of tobacco plants via *A. tumefaciens* transformation. α -zein mRNA in tobacco seeds varied between 1.0 and 2.5% of total mRNA; the α -zein protein was immunologically detected in seed extracts of transgenic plants (Scherthaner, Matzke, and Matzke 1988; Ohtani et al. 1991). These genetic efforts may enrich some plant products by increasing the concentration of selected bioactive peptides.

4. Common bean

Phaseolus vulgaris species or common beans are native to Mexico and Guatemala, and were associated with the development of the pre-hispanic cultures and today they play an important role in the nutrition of an important part of the world's population. They represent more than 50% of the legumes that are used for human consumption worldwide, which is due to its outstanding nutritional content sensorial properties and health effects. Beans are a major source of highly valuable plant protein, dietary fiber and micronutrients and are associated to lowering the risk of non-transmissible sicknesses, including diabetes and cardiovascular diseases, lowering blood glucose and lipid levels, decreasing cholesterol and triglycerides, as well ascertain benefits against cancer, and the treatment of obesity and metabolic syndrome. Some legume seeds also have anti-tumor, antiviral and antifungal activities (de Mejia et al. 2005; Wang, Rao, and Ye 2009; Lam and Ng 2011; Campos-Vega et al. 2012; Olmedilla-Alonso et al. 2013; Romero-Arenas et al. 2013; Rubiales and Mikic 2015). On the other hand, common beans have a key role in the diversification and intensification of agriculture by contributing to improve the environment due to their biological nitrogen fixation, effects on the soil, and control of weeds (Rubiales and Mikic 2015).

4.1. Nutritional composition

Common beans are rich in carbohydrates (60%) and proteins (19–27%), as well as a good source of vitamin B complex such as niacin, riboflavin, folic acid and thiamine; they also comprise iron, copper, zinc, phosphorus, potassium, magnesium and calcium. Besides, they have high fiber content and are an excellent source of polyunsaturated fatty acids. Phaseolin (30.2 to 53.5% of the total protein) is the principal storage protein and the most abundant protein in common beans; it has high resistance to proteolysis due to its globular and glycosylated structure (Montoya et al. 2010). *P. vulgaris* proteins are excellent source

of bioactive peptides, which provide antioxidant, antihypertensive, anticancer and antiinflammatory activities. On the other hand, common beans have phenolic acids, flavan-3-ols, flavonols, flavanones and anthocyanidins that show antioxidant, antiinflammatory, antihypertensive and antiatherosclerotic properties (Lin et al. 2008; del Rio et al. 2013; Romero-Arenas et al. 2013; Luna-Vital et al. 2015).

4.2. Bioactive peptides

Common bean protein is a low cost source for bioactive peptides and a great alternative as nutraceutical product or functional ingredient. Garcia-Mora et al. (2016) reported several ACE-inhibitory and antioxidant peptides in Pinto beans through BIOPEP Database; and concluded that the small peptides (<3 kDa) were the main contributors in these activities as well as specific amino acids in the peptide sequence; however, some peptides had more than one potential biological activity as a result of the amino acid sequences. Tagliazucchi et al. (2015) identified ACE-inhibitory peptides after gastrointestinal digestion of Pinto bean; they were di- or tri- peptides and the majority had a hydrophobic amino acid, (iso) leucine or phenylalanine, or proline at the C-terminal position which is crucial for the ACE-inhibitory activity. (Mojica and González de Mejia 2015) and Luna-Vital et al. (2015) concluded that ACE inhibition is the most frequent biological function of bean peptides (87.5%); they also reported DPP-IV inhibition in more than half of sequences analyzed (74 peptides), important in type-2 diabetes treatment and antioxidant capacity (36 sequences). On the other hand, Mojica and González de Mejia (2016) established a systematic evaluation and characterization for anti-diabetic peptides from black bean protein and found that peptides EGLELLLLLAG, AKSPLF and FEELN inhibited DPP-IV more efficiently than the control, and TTGGKGGK had higher inhibitory potential on α -glucosidase compared to the control acarbose. (Oseguera-Toledo, De Mejia, and Amaya-Llano 2015) also reported, peptides that inhibit *in vitro* α -amylase and α -glucosidase. Mojica, Luna-Vital, and Mejía (2017) found that phaseolin remained almost unhydrolyzed after digestion with pepsin/pancreatin; they also found four peptides: KKSSG, KTYGL, GGGLHK and CPGNK with potential to inhibit DPP-IV, ACE and α -glucosidase enzymes. Besides, they concluded that the biological activities are related to small amino acid sequences and demonstrated significant antioxidant, antidiabetic and antihypertensive properties (Table 4).

Wang and Ng (2007) found a 7.3 kDa peptide from *P. vulgaris* with potent antiproliferative activity for leukaemia cell line L1210 and lymphoma cell line MBL2. Luna-Vital et al. (2015) reported five peptides (GLTSK, LSGNK, GEGSGA, MPACGSS and MTEYY) in non-digestible fractions of common bean with antiproliferative activity on human colon cancer cells, that were produced by modifying molecules involved in cell cycle arrest or apoptosis. On the other hand, Lam and Ng (2011) reported that hemagglutinin from *P. vulgaris* protein, suppressed the proliferation of breast cancer MCF-7 cells (IC₅₀ of 0.2 μ M), and induced cell cycle arrest in G2/M phase, phosphatidyl serine externalization, mitochondrial membrane depolarization and apoptosis by activating the death receptor-mediated pathway.

Table 4. Bioactive peptides from common beans (*Phaseolus vulgaris*).

Bioactive function	Bioactive peptide sequence	Reference
Antihypertensive	KKSSG	Mojica, Luna-Vital, and Mejía 2017
	KTYGL	
	GGGLHK	
	CPGNK	
	LSFNT	
Antioxidant	KMARPV	Mojica, Luna-Vital, and Mejía 2017
	GHVPP	
	TACKD	
	GGGLHK	
	KTYGL	
Antidiabetic	MPHLK	Mojica and De Mejia 2015
	EGLELLLLLAG	
	AKSPLF	
	FEELN	
	TTGGKGGK	
Anticancer	KKSSG	Luna-Vital et al., 2015
	KTYGL	
	GGGLHK	
	CPGNK	
	VKFMT	
	GLTSK	
	LSGNK	
	GEGSGA	
	MPACGSS	
	MTEEY	

See Table 2 for the amino acid nomenclature used in this table.

4.3. Genetic engineering to improve the functional properties of peptides

The global economic importance of bean is due to mainly their high amount of storage proteins. In contrast to other legume seeds, common beans are composed of high amounts of globulins or phaseolin (7S fraction). Phaseolin contains different bioactive peptides encrypted in its amino acid sequence, and is coded by a multigene family; their expression is temporally regulated during embryogenesis and is localized in the embryonic tissues of common bean; from the 80s this protein fraction has been under several studies to improve the nutritional properties in different plants (Slightom et al. 1985; Chappell and Chrispeels 1986; Montoya et al. 2010; Carrasco-Castilla et al. 2012). Murai et al. (1983) transformed sunflower tissue cell with sequences coding phaseolin protein from common bean, consequently the protein was detected in transformed cells with different biochemical tools.

Similarly, Bustos et al. (1991) transformed tobacco plants with four phaseolin genes, the specific phaseolin mRNA was similar to those found in plants transformed with a bean genomic DNA sequence that encodes an identical β -phaseolin subunit. In addition, Bagga et al. (1992) introduced a chimeric β -phaseolin gene into alfalfa plants to study the pattern of accumulation of this protein in different parts of the transgenic plant including the mature seeds. They found high concentrations of phaseolin transcripts, however mature seeds only presented significant accumulation of the phaseolin protein. Zheng et al. (1995) expressed phaseolin in the endosperm of transgenic rice plants; promoter fragments of the rice glutelin gene were fused transcriptionally to either genomic or cDNA of phaseolin gene. These studies demonstrated the expression of a phaseolin gene derived from common beans transferred to improve the nutritional properties to distinct taxonomically plant species.

Many legume seeds are deficient in one or more essential amino acids, and improving the sulphur-containing amino acids has been a challenge for different research groups (Montoya et al. 2010). It is well known that common beans have a low content of sulphur-containing amino acids (Evans and Bandemer 1967); there is a great interest by plant breeders in increasing its methionine content. Protein engineering has offered alternatives for manipulating the amino acid content in the phaseolin. In this sense, (Dyer, Nelson, and Murai 1993) utilized molecular mechanic calculations to evaluate several strategies for improving the nutritional seed quality; these authors used this structure to simulate modifications aimed at increasing the methionine content of phaseolin. As a result of this simulation, methionine content was increased by the replacement of hydrophobic amino acids with methionine in the central structure of the phaseolin protein. Experimentally, researches have attempted to correct the methionine-deficiency by the introduction of a transgene coding for a methionine-rich storage albumin (2S albumin) from Brazil nut via biolistic methods that derived in bean transgenic lines (Aragão et al. 1992; Araújo et al. 1996; Araújo et al. 1999). Araújo et al. (1992) reported the transient expression of the Brazil nut 2S-albumin gene in cells of the bean embryonic that derived in transgenic bean plants containing and expressing the 2S-albumin gene (Aragão et al. 1996); in addition, the same authors found that in two of the five transgenic plants the methionine content was significantly increased by around 14–23% over the values found in non-transformed plants (Aragão et al. 1999).

The knowledge acquired about two extensively studied storage proteins, γ -zein of *Z. mays* and phaseolin of *P. vulgaris*, has been used to create a fusion protein called zeolin which has the solubility and assembly properties of wild-type γ -zein (Bellucci et al. 2007). Hoffman et al. (1987) used the maize zein gene under regulation of common bean β -phaseolin gene at flanking regions; *A. tumefaciens*-mediated transformation was used to insert the chimeric phaseolin-zein gene into tobacco genome, which gave as a result transgenic plants synthesizing zein in roots, hypocotyls, and cotyledons of germinating transgenic tobacco seeds. Similarly, Mainieri et al. (2004) produced a chimeric protein with the entire phaseolin coding sequence gene fused to the N-terminal region of γ -zein (including the tandem repeated and the Pro-rich domains). While γ -zein and phaseolin usually accumulate in the ER and in storage vacuoles, respectively; in transgenic tobacco leaves, zeolin is retained in the ER where its accumulation in protein bodies can reach 3.5% of the total leaf protein content. Similarly, zeolin was introduced to transform alfalfa plants, and the chimeric protein was expressed and was stably accumulated in alfalfa leaves because it forms endoplasmic reticulum-located protein bodies in cells (Bellucci et al. 2007). The authors conclude that the γ -zein portion is sufficient to induce the formation of protein bodies when fused to another protein as well. Because the storage proteins of cereals and legumes nutritionally complement each other, zeolin can be used as a starting point to produce nutritionally balanced and highly stable chimeric storage proteins (Mainieri et al. 2004).

5. Amaranth

Amaranth is a dicotyledonous plant. It grows fast in soil low in nutrients, and is resistant or tolerant to extreme conditions,

changes of temperature and radiation, which is due to its high osmotic adjustment capacity (Montoya-Rodríguez et al. 2015; Orona-Tamayo and Paredes-López 2017a). Amaranth plants are also widespread in temperate and tropical areas, and there are three main amaranth species native to the New World: *Amaranthus hypocondriacus* from the northwestern and central area of Mexico, *A. caudatus* from the Andes, and *A. cruentus* from the southern Mexico and the central region of Guatemala. Aztec civilizations planted *A. hypocondriacus* as one of their basic foods, Maya and Inca civilizations used *A. cruentus* and *A. caudatus*, respectively. These civilizations used its grains and leaves in different form; i.e., drinks, sauces that were mixed with maize to prepare tortilla, medicinal treatments and also in religious practices. It is worth to mention that amaranth was prohibited by Spaniards conquerors who were scornful of its use in native religious ceremonies (Kauffman 1992; Mallory et al. 2008; Mlakar et al. 2010; Pavlik 2011; Rastogi and Shukla 2013; Orona-Tamayo and Paredes-López 2017a).

5.1. Nutritional composition

The recent agronomic uses of amaranth began in 1970 when some researchers found that it was a promising plant; they increased the interest on this crop as potential staple food due to its high-quality proteins and outstanding nutritional compounds (Tucker 1986; Paredes-López 1994). The proximate composition of amaranth seeds depends to certain extent on the specie and culture place. The seed contains a high amount of digestive proteins; total protein ranges from 13.2 to 18.4% comprising a good balance of essential amino acids, close to the optimum required in the human diet. Amaranth contains important amount of crude fiber with ranges from 2.2 to 5.8%; the lipid level ranges between 1.9 and 13.0%, and it is composed by different fatty acids being the most abundant linoleic (36.7 to 55.9%), followed by oleic (18.7 to 38.9%) and palmitic (19.1 to 23.4%) acids. The main carbohydrate is starch (57–62.0%), and sucrose, glucose, raffinose, fructose, stachyose, maltose, and inositol are found in lower concentration. The grain is a good source of minerals such as Ca, Fe, Mn, Mg, K, P and Na. Also contains B complex vitamins. In addition, amaranth has other important bioactive compounds that include diverse metabolites such as phenolic acids, flavonoids, anthocyanins, tannins, and phytosterols (Becker et al. 1981; Segura-Nieto, Barba De La Rosa, and Paredes-López 1994; FAO 1985; Pasko et al. 2008; Repo-Carrasco-Valencia et al. 2009; Alvarez-Jubete, Arendt, and Gallagher 2010a; Ferreira and Areas 2010).

Amaranth contains proteins of high quality. Globulins are the main fraction at the storage proteins, and the most abundant are the 11S and 7S globulins. The 11S globulin, known as amarantin, has similar function as legumins, a molecular weight of 300–400 kDa and six subunits of 50–60 kDa. 7S globulins comprise four subunits (66, 52, 28 and 16 kDa) which integrate a polypeptide of 150–200 kDa. Amaranth globulins show a high concentration of lysine (5.7%) and high levels of sulfur amino acids (5.1%). 11S globulins contain important amounts of methionine (1.7%) and tryptophan (1.6%), high levels of lysine (4.3%), histidine (2.6%), phenylalanine (6.9%), valine (6.3%) and isoleucine (5.0%); while 7S globulin shows low levels of some essential amino acids, such as tryptophan

(0.4%) and methionine (0.5%), lysine (3.4%), histidine (1.5%), phenylalanine (4.9%), valine (5.5%), and isoleucine (6.2%) (Paredes-López, Mora-Escobedo, and Ordorica-Falomir 1988; Bressani and Garcia-Vela 1990; Barba de la Rosa et al. 1992; Segura-Nieto, Barba De La Rosa, and Paredes-López 1994; Vasco-Méndez and Paredes-López 1994; Barba de La Rosa et al. 1996; Marcone 1999a; Marcone 1999b; Tovar-Pérez et al. 2009; Quiroga et al. 2010)

5.2. Bioactive peptides

Most of the amaranth proteins encrypts mainly peptides with antihypertensive effect (Table 5). Montoya-Rodríguez et al. (2015) reported that 15 proteins mostly presented ACE-inhibitory peptides, and Fritz et al. (2011) evaluated *in vitro* and *in vivo* ACE-inhibition in peptides from *A. mantegazzium* and *A. hypocondriacus*. Montoya-Rodríguez et al. (2015) reported that 50% of peptides from amarantin have ACE inhibitory activity. Besides, Silva-Sánchez et al. (2008) found di- and tripeptides in globulins and glutelins with a strong inhibitory effect against ACE. Tovar-Pérez et al. (2009) identified important ACE-inhibitory effects in peptides from albumin (40%) and globulin (35%) fractions from *A. hypocondriacus*. Moreover, Vecchi and Añón (2009) studied tetrapeptides from 11S globulin of *A. hypocondriacus* with ACE-inhibitory activities (Table 5).

Luna-Suárez et al. (2010) modified the acidic subunit of the 11S globulin from amaranth through the insertion of four peptides with antihypertensive functions. These peptides showed a high inhibition effect against the ACE, eight-fold more than the non-modified 11S globulin. In our laboratory, these peptides were overexpressed in *E. coli* strains to obtain a highly production of antihypertensive peptides (Arano-Varela, Dominguez-Dominguez, and Paredes-López 2012; Castro-Martínez, Luna-Suárez, and Paredes-López 2012; Morales-Camacho, Dominguez-Dominguez, and Paredes-López 2013). Later, these antihypertensive peptides were evaluated by Medina-Godoy et al. (2013) in spontaneously hypertensive rats, and they found considerable reduction of arterial pressure, similar to the control group treated with commercial captopril. Amaranth 7S globulin also encrypts antihypertensive peptides with high ACE-inhibitory capacity (Quiroga et al. 2012). (Barba de la Rosa et al. 2010) reported that peptides from amaranth glutelins possess various amino acids with high ACE-inhibitory effects, comparable to captopril.

Qureshi Lehmann, and Peterson (1996) assessed the influence of amaranth in the cholesterologenesis in chickens of 6 weeks of age and found that it reduced total and LDL-cholesterol; also, they concluded that cholesterol-7- α -hydroxylase enzyme and 3-hydroxy-3-methylglutaryl coenzyme A (HMG-CoA) decreased. Similarly, Plate and Arêas (2002) reported a reduction in LDL and total cholesterol in hypercholesterolemic rabbits supplemented with extruded amaranth; they suggested that amaranth consumption may be an alternative option to prevent coronary diseases. However, these studies attributed the anti-cholesterolemic effects to different compounds, such as fiber and the bioactive peptides encrypted in the protein sequences. On the other hand, Escudero et al. (2006) reported that Wistar rats fed with concentrates of amaranth protein

Table 5. Bioactive peptides from Amaranth (*Amaranthus hypochondriacus*).

Bioactive function	Protein source	Bioactive Peptide/Peptide sequence	Reference
Antihypertensive	Total protein (hydrolysates)	YL, RW, RR, KL, LF, EG, GT, HK, RP, HP, PG, GG, GL, LG, GA, AG, GT, HG, FG, PR, GP, PRY, DG, GP,	Montoya-Rodríguez et al. 2014
	Albumin	AG, GL, RY, LQ, LA, EV, VE, RW, LY, EG LG, GH, TE, VG, GG, GS, TG, VF, IF, TF, KG, NG, GK, GV, FG, GY, YG, LN, AH, MF, GT, KF, PR, EA, AA, FR, KR, SG, TQ, AI, EI, IE, RL GKP, LF, FP, YL, RF, HY, VV ALEP, VIKP VY	Montoya-Rodríguez et al. 2015
	Amarantin (11S globulin)		Silva-Sánchez et al., 2008 Vecchi and Anon 2009 Luna-Suárez et al. 2010; Arano-Varela, Dominguez-Dominguez, and Paredes-López 2012; Castro-Martínez et al., 2012; Morales-Camacho, Dominguez-Dominguez, and Paredes-López 2013; Medina-Godoy et al. 2013 Montoya-Rodríguez et al. 2015
7S Globulin		SG, GF, HP, PT, NG, MG, GE, EG, GR, FR, QG, GL, TE, EV, AG, GV, RR, IE, PH, HG, AP, IY, GI, TG, GM, IP, PG, GS, GG, KF, FG, GD, HL, IF, EA, AA, KG, LY, LN, AH, MY, GQ, VF, VP, AI, AF, EW, MF, RL, IR, IKP, GKP, IEP, ALEP, DA, GK, QG, LQ, RL, 2VF, PR, PL, AP, GF, FR, VG, AG, 2 GE, MG, GD, EG, EA, 2LQ, LN, PQ, EW, HP LVL, HY, FP, PR, 4LF, GPL, 2GP, PL, VK, 2AF, 2LA, KR, VP, 3FR, GL, GH, GR, 2FG, GK, GT, GG, 2EG, NG, 2PG, VR, GHF, NY, NF, 2LQ, LN, EK VLP, LNP, 2YL, GPL, PL, AF, AP, 3GA, 3AG, 2FG, GS, 2 GV, 2GT, QG, 4SG, GHF, RR, AR, PH, 2VF, GY, AY, AA, VG, GE, QG, EA, NG, NF, YK, EV 2LY, LVL, RF, HY, GY, LNP, YL, 2LF, VK, 2AF, AP, 2LA, FR, GA, 3GL, GR, FG, GS, GT, 2SG, 2LG, 2EG, NY, 2 SF, KL, LVE, VE, LQ, LN, TQ, EK, KE, PH LP, RL, PR, LSP, LF, FFL, VP, GF, AG, SY, LN, PQ, TF, EL, FA, LP, VLP, VF, HY, GY, AF, LA, KR, FR, VG, 2 NF VF, FP, GF, GL, FG, LN, HY, PR, LF, GP, AF, LA, KR, PG, NF AP, AIP, ALPP, AVP, AY, FNQ, FP, FQP, FY, GGY, GKP, GRP, GY, HIR, HY, IKP, ILP, IR, IRA, IV, LAA, LAMA, LAY, LF, LLP, LNP, LPP, LQP, LQQ, LRP, LSP, LVL, LVR, LW, LY, MF, MY, PLP, PQR, PR, PRY, RF, RL, RY, TAP, VAA, VAP, VAY, VF, VLP, VPP, VPR, VSP, VW, VY, VYP, YG, YGGY, YL, YP AY, FP, FY, GY, HY, IR, IY, LF, LW, LY, ME, MY, PR, RF, RL, RY, VF, VW, VY, YG, YL, YP AAP, AIP, AVP, FNQ, FQP, GGY, GKP, GRP, HIR, IKP, ILP IRA, LAA, LAY, LLP, LNP, LPP, LQP, LQQ, LRP, LSP, LVL, LVR, PLP, PQR, PRY, TAP, VAA, VAP, VAY, VLP, VPP, CRP, VSP, VYP, ALPP, LAMA, YGGY PYY, RY, WY, RW, PWW, PWR, PW, PWY, WYS, PW, WY LK, TY, RW, LH, KP, EL, LHV, LVR IR, EL, TY, HL, YL, LH, KP, YV, YL, PHG, PEL, RHL, LHV AWEE, REQ, GSR, YLAGK, PQQEH, IYIEQ, NGITGM, TEWDS, NEQ HH, HL, LH, LHH VVV, GQ SSDIKE	(Quiroga et al. 2012)
	Glutelin		Silva-Sánchez et al. 2008
Antioxidant	Total protein (hydrolysates)		Barba de la Rosa et al. 2010
	Albumin		Montoya-Rodríguez et al. 2014
	Amarantin (11S globulin)		Montoya-Rodríguez et al. 2015
Anticancer Immunomodulatory	Glutelin		Orsini-Delgado et al. 2016
	Amarantin (11S globulin)		Silva-Sánchez et al. 2008
	Total protein (hydrolysates)		Montoya-Rodríguez et al. 2015 Moronta et al. 2016a; Moronta et al. 2016b
Glutelin		EAE, GFL, KRP, TKPR, YG, YGG	Silva-Sánchez et al. 2008

(continued on next page)

Table 5. (Continued).

Bioactive function	Protein source	Bioactive Peptide/Peptide sequence	Reference
Hypocholesterolemic	Total protein (hydrolysates)	GGV, IVG, VGVL	Soares et al. 2015
Antimicrobial	Total protein	VGECVRGRCPSGMCCSQFGYCGKPKYCG (Ac-AMPT1)	Broekaert et al. 1992
	Total Protein	VGECVRGRCPSGMCCSQFGYCGKPKYCGR (Ac-AMP2)	Lipkin et al. 2005
	Total protein	AGECVQGRCPSGMCCSQFGYCGRGPYCGR (Ar-AMP)	Rivillas-Acevedo and Soriano-García 2007a; Rivillas-Acevedo and Soriano-García 2007b
	Albumin	CPSGMCCSQFGYCGK (AyAMP)	Montoya-Rodríguez et al. 2015
Antidiabetic	Albumin	KK	Montoya-Rodríguez et al. 2014
	Total protein (hydrolysates)	AP, GP, FP, GP	Montoya-Rodríguez et al. 2015
	Albumin	MA, LP, LL, LA, FA, VA, FP	Montoya-Rodríguez et al. 2014
	Amarantin (11S globulin)	HA, MA, LL, LP, AP, FA, MP, PA, VA, FP, LA, PP, KA, VV, VP,	Montoya-Rodríguez et al. 2015
	Glutelin	AP, FA, FP, GP, GPR, GQ, HA, IPI, KA, LA, LL, LP, MA, MP, PA, PP, PPLP, PPPA, VA, VP, VPL, VV	Montoya-Rodríguez et al. 2015 Silva-Sánchez et al. 2008

See Table 2 for the amino acid nomenclature used in this table.

concentrated presented a reduction of serum LDL protein and triglycerides compared to control group; the liver also showed a reduction of total cholesterol and triglycerides, and the fatty acid synthase decreased in its activity. Mendonça et al. (2009) showed that hamster fed with amaranth protein isolates reduced total plasma cholesterol and non-HDL-cholesterol levels. Soares et al. (2015) identified peptides with HMG-CoA enzyme activity, a key enzyme in the biosynthesis of cholesterol. They reported GGV, IVG, and VGVV peptides inhibiting the activity of HMG-Co reductase, suggesting a possible hypocholesterolemic effect.

The physiological effect of the antioxidant peptides from amaranth seed proteins has been proved in different studies (Guzmán-Maldonado and Paredes-López 1998). Silva-Sánchez et al. (2008) reported the antioxidant potential of four peptides rich in histidine encrypted in the amaranth 11S globulin. Tironi and Añón (2010) found peptides with high antioxidant capacity from *A. mantegazzianus* in all protein fractions. (Orsini-Delgado, Tironi, and Añón 2011) probed *in vitro* the antioxidant effect of amaranth peptides derived from multi-enzymatic peptidases. Later on, the same group investigated the antioxidant potential of peptides released by gastrointestinal digestion and identified a high increment in the scavenging activity against different radicals, such as peroxy, hydroxyl and peroxynitrites (Orsini-Delgado et al. 2015). Also, Sabbione et al. (2016) obtained antioxidant peptides derived from the activation of endogenous amaranth proteases, and they found more antioxidant activity by the hydrolysates than the protein isolate from the amaranth seed flour. Many studies have reported the benefits of the peptides derived from amaranth proteins; however, amaranth hydrolysates provide a complex combination of peptides with different physical properties (Le Maux, Nongonierma, and Fitzgerald 2015).

In the last years, bioinformatics and protein sequencing have simplified the analysis of the effect of peptides and facilitated subsequent experimentation (Moruz and Käll 2016). In this sense, Montoya-Rodríguez et al. (2014) characterized antioxidant peptides from amaranth hydrolysates, and identified different antioxidant amino acids, such as lysine, proline, histidine, glycine and threonine. In Montoya-Rodríguez et al. 2015 determined, by bioinformatic studies, that the amaranth 11S globulin encrypted diverse di- and tripeptides with antioxidant activities. Besides, Orsini-Delgado et al. (2016) identified four peptides from the acidic subunit of the 11S globulin with high antioxidant capacities.

Amaranth contains proteins and peptides with anti-tumoral and anti-proliferative effects (Martínez-Cruz, Cabrera-Chávez, and Paredes-López 2014). For example, Yu et al. (2001) isolated a lectin from *A. mantegazzianus* and concluded that amaranth lectin can have a marked effect on the inhibition of malignant cancer cells. Barrio and Añón (2010) worked with *A. mantegazzianus* protein hydrolysate and found that it produced morphological changes and caused a rearrangement of the cytoskeleton in UMR106 cell line, inhibited cell adhesion and induced apoptosis and necrosis. Lunasin is also a promising peptide involved in prevention of different types of cancers. Lunasin was discovered in soybean and then was found in amaranth (Hernandez-Ledesma, Hsieh, and Ben 2009b; Hsieh, Hernández-Ledesma, and De Lumen 2010). Maldonado-Cervantes

et al. (2010) reported that amaranth lunasin is a highly efficient peptide to prevent cancer, mainly because it is capable of enter into the nucleus of the cell and inhibits the transformation of NIH-3T3 spotlights cancerous cells.

Peptides from extruded amaranth inhibited inflammatory process in human and mouse macrophages by preventing the inactivation of NF- κ B signaling (Montoya-Rodríguez et al. 2014). Similarly, Moronta et al. (2016a) determined that amaranth protein hydrolysates have immunomodulatory effects on epithelial cells through the NF- κ B signaling pathways; also, they identified the peptide SSEDIKE, which has the ability to attenuate the activation of epithelial cells. The same authors reported that SSEDIKE peptide inhibits the allergic reaction in mice with food allergy, prevents IgE secretion, and intestinal inflammation control preventing the activation of NF- κ B (Moronta et al. 2016b).

Peptides with antimicrobial effect in the plasma membrane of microorganism are found in amaranth grains (Broekaert et al. 1997). Broekaert et al. (1992) isolated two antimicrobial peptides (Ac-AMP1 and Ac-AMP2) from *A. caudatus*, which bind to chitin in an irreversible way and inhibit the growth of different plant pathogenic fungi and show some activities against Gram-positive bacteria. The antimicrobial peptide Ar-AMP was isolated from *A. retroflexus* seeds, its sequence belongs to the hevein-like family of antimicrobial peptides with six cysteine residues; also, this peptide effectively inhibited the growth of different fungi (Lipkin et al. 2005). An antifungal peptide Ay-AMP was isolated from *A. hypochondriacus* seeds which has the capacity to inhibit the growth of *Candida albicans*, *Trichoderma* sp., *Fusarium solani*, *Penicillium chrysogenum*, *Geotrichum candidum*, *Aspergillus candidus*, *Aspergillus ochraceus*, and *Alternaria alternata* (Rivillas-Acevedo and Soriano-García 2007a; Rivillas-Acevedo and Soriano-García 2007b). In addition, Pribylova et al. (2008) identified that the gene AMP2 in *A. caudatus*, *A. albus*, *A. cruentus*, *A. blitum*, *A. hybridus*, *A. hypochondriacus*, *A. retroflexus* and *A. tricolor*.

Amaranth peptides have the potential to diminish some key enzymes associated to type 2 diabetes, such as α -amylase and DPPIV. Conforti et al. (2005) reported the *in vitro* inhibition of α -amylase from two amaranth varieties that involve the digestion of starch and reduction of glucose absorption; however, they did not find if this effect was caused by peptides. (Velarde-Salcedo et al. 2013) reported the inhibitory effect against DPPIV enzyme by different peptides encrypted in amaranth globulin and glutelin. These amaranth peptides could be used as a novel therapy to prevent and diabetes treatment.

5.3. Genetic engineering to improve the functional properties of peptides

Amaranth protein globulin 11S or amarantin contained an array in their amino acid sequences of different potential bioactive peptides that confer important health benefits such as ACE, DPPIV and antioxidant inhibition effects (Montoya-Rodríguez et al. 2015). Amarantin gene is an excellent candidate to be expressed in diverse crops to improve their nutritional value as staple foods with a higher amount of bioactive peptides with different functional activities. Since the middle of 90's, this protein was subjected to different molecular studies to

understand its processing and deposition in the amaranth grain (Barba De La Rosa et al. 1996). The importance of amaranth globulin 11S has led researchers in the last few decades to isolate, modify, overexpressed and purified amarantin gene and protein in *E. coli* cells. In this context, Osuna-Castro et al. (2000) overexpressed amarantin in *E. coli* and after purification found *in vitro* a proper refolding of this protein; in addition, the protein expressed and purified from *E. coli*, exhibited the same properties than the native amarantin from the seed (Barba de La Rosa et al. 1996). (Medina-Godoy et al. 2004) improved the potency of purification of the amarantin. They cloned and expressed a modified cDNA of histidine-tagged 11S globulin from *A. hypocondriacus* in *E. coli* bacteria; the resulting amarantin expressed was purified in one step with high yields. Amarantin protein is susceptible to be modified in its acidic subunit with the potential as a functional and nutraceutical molecule. This subunit contains four hypervariable regions of five regions detected in the 11S seed globulin (Tandang-Silvas et al. 2012). In that sense, Luna-Suárez et al. (2008) modified the third variable region of the acidic subunit of a His-tagged 11S amarantin and expressed in *E. coli*. The hydrolyzed and modified amarantin, showed a higher inhibitory activity against ACE than the native seed amarantin (Luna-Suárez et al. 2010). Similarly, (Arano-Varela, Dominguez-Dominguez, and Paredes-López 2012) expressed amarantin and evaluated different parameters in a bioreactor to obtain a high concentration of the recombinant acidic subunit, and demonstrated that the aeration rate and temperature were the most important variable to increase the protein yield. (Castro-Martínez, Luna-Suárez, and Paredes-López 2012) increased in a reactor around three-fold the protein yield expressed as a modified amarantin using different parameters such as temperature, agitation speed, and oxygen conditions.

Later, (Morales-Camacho, Dominguez-Dominguez, and Paredes-López 2013) expressed the acidic subunit of the amarantin manipulating parameters in a bioreactor when they used two minimal media supplemented with the inducer isopropyl- β -D-thiogalactopyranoside or lactose, and found a high yield of the recombinant acidic subunit in a short time of induction. Similarly, Morales-Camacho et al. (2016) modified the C-terminal in the fourth variable region of acidic subunit of amarantin by inserting four VY antihypertensive peptides and expressed in *E. coli*. They found that modification of the variable region improves the thermal stability of amarantin acidic subunit. Another system was used to modified and expressed amarantin gene. In this context Medina-Godoy et al. (2006) used *Pichia pastoris* to insert a peptide that was tagged to retrieve the amarantin and overexpress and enhance the protein.

Amarantin gen is a candidate to be expressed in different important crops in order to improve the bioactive peptides and nutritional value of staple foods. Valdez-Ortiz et al. (2005) cloned the amarantin gene and transformed tobacco plants; they found that the protein retrieved its biochemical and physiological properties of the native amarantin. Rascón-Cruz et al. (2004) isolated and used the 11S globulin cDNA to transform maize plants; they analyzed and evaluated the insertion and expression of the amarantin gene in transformed plants. When the amarantin was accumulated in the maize seed, their content

of essential amino acids profile such as Lys and Trp increased in relatives non-transformed plants. Similarly, Germán-Báez et al. (2014) used acidic subunit of amarantin from *A. hypocondriacus* and transformed tomato fruits; they found an increase of total protein content and higher concentrations of essential amino acids such as Val, Tyr, Ile and Leu. By the way, albumin gene that coded an important protein in amaranth, was used to transform potato tuber plants with the aim of improving nutritional quality of tuber proteins; the introduction of this gene imparted an increase in total protein and in essential amino acids as well as in phytoesters (Chakraborty, Chakraborty, and Datta 2000; Shekhar et al. 2016).

6. Chia

Chia (*Salvia hispanica* L.) is native to Central Mexico up to Northern Guatemala, and its use as a food source has been dated as far back as 3,500 BC. The *Salvia* genus is the largest of the Lamiaceae family, with nearly 1000 species scrubs. The word “Chia” is derivate from Nahuatl (Mexican indigenous language) and it means oily. It is an annual plant that grows around 1 m and has oppositely arranged leaves with small white and purple hermaphrodite flowers. This plant was cultivated for its taste and was also used in folk medicine. Later, between 1500 and 900 BC Aztecs, Mayas and Incas used the seeds to prepare medicine, foods and paintings. In pre-Columbian cultures, it was the main staple crop after beans and maize, and was an integral part of diet of these civilizations, providing to these civilization important compounds that made their diet even superior to today's one. In addition, chia seeds are also a source of energy and are promoted as health food product (Ayerza, 2009). Nowadays the plant is considered as the super-food due of the 21st century to its nutraceutical potential (Capitani et al. 2012; Jamboonsri et al. 2012; Muñoz et al. 2013; Ullah et al. 2015; Orona-Tamayo, Valverde, and Paredes-López 2017b).

People has increased recently the consumption of chia due to its health benefits properties and nutritional value. It can now be found at markets in different brands and preparations like raw seeds, flour, food supplement, oil capsules, in yoghurts, drinks, energy bars, cereal, peanut butter and bakery products, and also animal feed formulations; mainly for horses, birds and rabbits. Raw chia seeds and seed flour are becoming popular and they may be added to muesli, breakfast cereals and fruit, nut and different seed mixes, shakes, desserts or homemade bakery products (Reyes-Caudillo, Tecante, and Valdivia-López 2008; Peperkamp 2014; Valdivia-López and Tecante 2015; Orona-Tamayo, Valverde, and Paredes-López 2017b).

6.1. Nutritional composition

The nutritional content of chia seeds includes 25–40% fat, 15–24% protein, and 26–41% carbohydrates and several minerals. The fat fraction has a high content of essential polyunsaturated fatty acids which comprise 55–64% linolenic acid (ω –3) and 19–20% linoleic acid (ω –6). ω –3 fatty acids from chia (α -linolenic, eicosapentanoic and docosahexanoic acid) and ω –6 (linoleic and arachidonic acids) are essential polyunsaturated fatty acids (PUFAs). Other important nutritional aspects of the

seeds are the content of total dietetic fiber, 42% of total dry weight (Gómez-Favela et al. 2017), and the high-quality amino acids. Also, chia seeds contain high levels of phenolic compounds with important antioxidant capacity. The seeds contain gallic, caffeic, chlorogenic, ferulic and rosmarinic acids; other important phenols such as myricetin, quercetin, and kaempferol, as well as different isoflavones like daidzin, glycitin, genistin, glycitein, and genistein (Muñoz et al. 2013; Martínez-Cruz and Paredes-López 2014). The presence of these nutraceutical components and the healthier life style of the population have increased the popularity of this plant, and led to main producers of chia such as Bolivia and Paraguay. Mexico, Argentina and Australia are emerging producer as a result of the increasing demand of this seed.

The protein content of chia seeds (15–24%) is higher than other cereals like maize (14%), wheat (14%), oats (15.3%), barley (9.2%) and rice (8.5%). Besides, it comprises high concentration of essential amino acids (histidine, isoleucine, methionine, phenylalanine, threonine and tryptophan) and is an excellent source of sulfur amino acids (cysteine and methionine) as well as arginine, aspartic and glutamic acids. Globulins are the major protein fraction which constitute around 55% of the total storage proteins, followed by 18% albumins, 15% glutelin and 12% prolamin (Sandoval-Oliveros and Paredes-López 2013; Orona-Tamayo et al. 2015; Gómez-Favela et al. 2017; Orona-Tamayo, Valverde, and Paredes-López 2017b).

6.2. Bioactive peptides

The high amount of proteins in chia encrypts in their sequences biologically active peptides with different nutraceutical

properties, including antihypertensive, antioxidant, antidiabetic and other important functions (Figure 1). Salazar-Vega et al. (2012) and Segura-Campos et al. (2013b) showed that peptides obtained from extensive hydrolysis inhibited more efficiently ACE activity than peptides from shorter hydrolysis. Segura-Campos et al. (2013a) reported that peptides from the globulin fraction, subjected to an extensive enzymatic hydrolysis, had high ACE-inhibitory activity and this was increased in peptides with low-molecular weight (<1 kDa; 69.3% of ACE-inhibition). Recently, Orona-Tamayo et al. (2015) found that peptides from albumin and globulin fractions exert more inhibition effect against ACE-enzyme than peptides from the other proteins fractions, similar results were found by Segura-Campos et al. (2011). The biological activity of chia proteins hydrolysates with ACE inhibition is related to the amino acid sequence of the peptides encrypted in the storage proteins, and the activity is influenced by the C-terminal. Chia seeds contain a high concentration of hydrophobic amino acids that include proline, isoleucine and phenylalanine, which have high ACE-inhibitory activity (Murray and FitzGerald 2007; Alemán et al. 2011a; Segura-Campos et al. 2013a; Orona-Tamayo, Valverde, and Paredes-López 2017b). Therefore, peptides with ACE-inhibitory activity from chia seeds could be a natural treatment for hypertensive patients. Diabetic individuals with cardiovascular risk, supplemented with 37 g/day of ground chia seeds, showed reduction in systolic blood pressure (6.3 mm Hg) compared to control patients (Vuksan et al. 2007). However, Nieman et al. (2009) evaluated the hypertensive effect of 50 g/day of whole or ground chia for 12 weeks and no differences in blood pressure were found. The same authors supplemented overweight women with 25 g/day of whole or ground chia for 10 weeks




Figure 1. Bioactive peptides from chia (*Salvia hispanica*) globulin fraction.

and reported that the ingestion of chia did not influence cardiovascular risk factors (Nieman et al. 2012). On the other hand, Toscano et al. (2014) evaluated, in under treatment and non-treatment hypertensive patients, the incorporation of 35 g/day of chia seed flour into their diet for 12 weeks, and found a reduction of blood pressure (5.2 mm Hg) in treated patients, while the control group showed no change, concluding that chia consumption is able to decrease the blood pressure in hypertensive patients.

Orona-Tamayo et al. (2015) showed that peptides from albumin and globulin fractions presented high antioxidant capacity against DPPH and ABTS with lower IC₅₀, and peptides from glutelin and prolamin fractions showed lower antiradical activity against these oxidants. They concluded that albumin and globulins encrypted important peptides with a high antioxidant capacity. These authors determined that peptides from prolamin showed a higher capacity to chelate Fe⁺² ion than peptides from globulin and glutelin fractions. In living systems, the presence of transient metal ion, such as Fe²⁺ and Cu²⁺ involve the formation of ROS (Canabady-Rochelle et al. 2018). These results indicate that peptides from globulin and prolamin are the most powerful chelating components in chia seeds.

6.3. Genetic engineering to improve the functional properties of peptides

Nowadays genetic, molecular biology and genomic information on bioactive peptides from chia is missing, therefore there is an opportunity to use different mutagenesis and other advanced molecular tools to facilitate rapid genetic improvement of proteins and peptides in this plant. In that sense, Jamboonsri et al. (2012) chemically mutagenized chia seeds with ethyl methanesulfonate to obtain seeds that generated chia early flowering lines able to flower under a short photoperiod; under field conditions these lines flowered at a photoperiod of around 14 h, and the plants produce seeds in a range of environments in temperate areas.

New generation tools can help us to reveal different gene function related to important metabolic networks. The high polyunsaturated fatty acids content in chia seeds makes this crop very attractive as food and for different industrial uses. From a commercial point view, it is desirable that seeds contain high levels of triacylglycerols enriched with beneficial fatty acids (Brown et al. 2012). Different researches have focused their attention the identification of genes that encoding enzymes related to fatty acids and triacylglycerol biosynthesis (Ruiz-Lopez et al. 2015). In this context, Xue et al. (2017) isolated from RACE method two full-length Fatty Acid Desaturases (*ShFAD2-1* and *ShFAD2-2*) genes that encoded key enzymes for linoleic acid and α -linolenic acid biosynthesis in Chia. Both *ShFAD2-1* and *ShFAD2-2* proteins possess strong transmembrane helices, three histidine motifs and a C-terminal ER-located signal, and their heterologous expression in *Saccharomyces cerevisiae* indicated that both genes encoded a bio-functional delta-12 oleate desaturase. *ShFAD2-2* was mainly expressed in flowers and early-stage seeds while *ShFAD2-1* expression was almost constitutive in different organs. This study can provide insights into molecular mechanism of high α -Linolenic acid (ALA) traits of *S. hispanica*. Sreedhar et al.

(2015) analyzed a global transcriptome in chia seed development, and found genes involved in the lipid metabolism and oil accumulation in chia seeds. However, chia is a diploid plant with only 12 chromosomes (Estilai, Hashemi, and Truman 1990), which facilitates rapid genetic improvement and the plant breeding (Araki and Ishii 2015) by new molecular generation technologies. This genomic information may be useful to improve the quality and quantity of proteins, peptides and oil from chia seeds.

7. Quinoa

Quinoa (*Chenopodium quinoa*) is a flowering plant from the Amaranthaceae family. It was cultivated 7,000 years ago by the Incas in the Titicaca Lake and then was spread to Ecuador and Chile. Actually it is mainly cultivated in Bolivia, Chile, Peru, Ecuador, and Colombia; as well as in few other countries such as England, Sweden, Denmark, the Netherlands, Italy, Kenya and France (Bhargava, Shukla, and Ohri 2006; Bhargava, Shukla, and Ohri 2007; Fuentes et al. 2009; Iqbal 2015).

Quinoa has large genetic variability and great capacity to adapt to different agro-ecological edaphic conditions. It shows tolerance to frost, salinity and drought; also has the potential to grow on marginal soils, it may also be tolerant to disease, pests, and it can grow at pH between 4.8 and 9.5. Five ecotypes have been identified: *salares* (salt flats), highlands, inter-Andean valleys, *yungas* and coastal-lowlands (Jacobsen, Mujica, and Jensen 2003; Repo-Carrasco, Espinoza, and Jacobsen 2003; Iqbal 2015; Aloisi et al. 2016; Rizzello et al. 2017).

This grain is considered an excellent functional food. It is part of FAO strategy to encourage the cultivation of traditional crops as means of contributing to food security for the 21st century; and it is considered by NASA as an emerging crop with excellent nutritional properties, mainly due to its protein and unique amino acid composition. Regarding economics and sustainability, sweet quinoa varieties are the most promising because they provide more high-quality protein than the bitter crops. It is considered a pseudocereal and the whole grain may be used to fermented to produce beer, or in bread, soups, biscuits, drinks, etc.; also, the whole plant is used as green forage to feed cattle, pigs and poultry (Schlick and Bubenheim 1993; Jacobsen, Mujica, and Jensen 2003; Jancurová, Minarovicová, and Dandar 2009; Vega-Gálvez et al. 2010; Iqbal 2015).

7.1. Nutritional composition

The nutritional composition of quinoa varies among ecotypes due to genetic variability and environmental conditions (Repo-Carrasco, Espinoza, and Jacobsen 2003). Its protein content (13.8 to 16.5%) is higher than barley (11.0%), rice (8.8%), maize (10.5%), rye (11.6%) and sorghum (12.4%); but, the importance relies on the high level of essential amino acids, which is also higher than common cereals. Quinoa has a well-balanced amino acid profile with high amounts of lysine and methionine; it contains 1.4 times more lysine than soybean, twice more than maize, 2.3 more than wheat. Its proteins do not contain gluten, which is a promising characteristic for celiac patients (Kozioł 1992; Aubrecht and Biacs 2001; Mujica-Sanchez et al. 2001; Drzewiecki et al. 2003; Comai et al. 2007; Alves, Rocha, and

Gomes 2008; James 2009; Jancurová, Minarovicová, and Dandár 2009; Vega-Gálvez et al. 2010; Stikic et al. 2012; Escuredo et al. 2014). Storage proteins of quinoa are mainly albumins (2S) and globulins (11S) (35 and 37%, respectively); among the 11S, globulins A and B are the most abundant fractions, and their amino acid sequence was almost identical (Brinegar and Goundan 1993; Brinegar, Sine, and Nwokocha 1996; Gorinstein et al. 2002; Wright et al. 2002; Watanabe et al. 2003; Gorinstein et al. 2005; Capriotti et al. 2015). On the other hand, Aloisi et al. (2016) determined that salinity altered the seed proteome and amino acid profiles, which, increasing the concentration of bioactive molecules and oxidant activity of protein extracts.

Quinoa contains important levels of vitamins (A, B2, E) and minerals (Ca, Fe, Cu, Mg, Zn); lipids between 2.0 to 9.5% are rich in linoleic (49 to 56%) and α -linolenic (9 to 12%) acids, which correspond to 88% approximated of the total fatty acids (Koziot 1990; Koziol 1992; Ando et al. 2002; Repo-Carrasco, Espinoza, and Jacobsen 2003; Ryan et al. 2007; Borges et al. 2010).

The seeds also comprise important bioactive compounds, such as vitamin B2 and E, carotene, tocopherols and phenolic compounds. It is good source of flavonoids, quercetin and kaempferol glycosides. Phenolic compounds, vitamins, proteins and fat may vary between quinoa ecotypes; which are related to the environmental conditions, farming techniques and genetic factors; antioxidant and antimicrobial activities also vary among genotypes and environment (Repo-Carrasco, Espinoza, and Jacobsen 2003; Nsimba, Kikuzaki, and Konishi 2008; Alvarez-Jubete et al. 2010b; Miranda et al. 2011; Miranda et al. 2013; Miranda et al. 2014; Iqbal 2015).

7.2. Bioactive peptides

There is scarce information reported in literature about bioactive peptides of quinoa proteins. That is why it is necessary to carry out more studies to demonstrate the nutraceutical potential of the proteins of this pseudocereal.

Fortunately, Rizzello et al. (2017) reported that five bioactive peptides from quinoa (5 to 9 amino acid residues) contain important antioxidant activity. They were found in the NCBI nr database as encrypted in different quinoa proteins. IVLVQEG and TLFRPEN are encrypted in a quinoa globulin B. VGFGI and FTLLIN released from salt sensitive protein, and LENS GDKKY encrypted in a maturase K protein.

Genetic engineering to improve the functional properties of peptides

As indicated before, there is almost no information based on genetic engineering of this important seed. Fortunately, very recently Zou et al. (2017) sequenced a genome of stress tolerance of quinoa and found that it is associated with the expansion of genes involved in ion and nutrient transport, abscisic acid (ABA) homeostasis and signaling, and enhancing basal-level ABA response. Also, Jarvis et al. (2017) sequenced a chromosome-scale reference genome for quinoa which facilitated the identification of the transcription factor likely to control the production of anti-nutritional triterpenoid saponins found in this crop.

8. Safety of bioactive peptides from seed

Seeds contain high concentration of storage proteins with important levels of different amino acids, which have diverse functional biopeptides. There is plenty information on the benefits of seeds peptides; moreover, the research has emphasized the peptides efficacy, and now it is necessary to analyze their safety, allergenicity and potential toxicity (Aluko and Monu 2003; Yamamoto, Ejiri, and Mizuno 2003; Luna-Vital et al. 2015; Montoya-Rodríguez et al. 2015; Nongonierma et al. 2015; Orona-Tamayo et al. 2015; Valverde et al. 2016; Orona-Tamayo and Paredes-López 2017a; Orona-Tamayo, Valverde, and Paredes-López 2017b).

Toxic sequences of peptide contain high concentrations of proline, histidine and asparagine, as well as cysteine residues. These peptides contain dominant amino acids such as valine, arginine, threonine, glutamine, methionine, leucine, lysine, isoleucine, phenylalanine, and alanine. Peptides with ACE, renin and DPPIV inhibitory effects, contain residues of phenylalanine, leucine, arginine, tryptophan, isoleucine and high concentration of proline residues which suggests that these peptides can be potentially toxic (Gupta et al. 2013; Lafarga and Hayes 2017).

Gupta et al. (2013) developed *in silico* models to predict toxicity of peptides and proteins, named as “ToxinPred” (<http://www.imtech.res.in/raghava/toxinpred/>). They use various properties of the peptides to predict toxicity, minimal mutations in peptides to increase or decrease toxicity, and toxic regions of proteins. It is worth to mention that the safety of the bioactive peptides, and their hydrolysates used as food ingredients for therapeutic proposes, should be also evaluated *in vitro* and *in vivo*. However, no side effects have been reported in consumers using bioactive peptides from food.

9. Conclusions

Nutraceuticals or functional foods are any product derived from food source with extra health benefits, in addition to the basic value of the food. They may be considered non-specific biological therapies used to promote general well-being, control symptoms and prevent malignant process. In recent years, the relationship between diet and health has been more clear; however, it was known since the time of Hippocrates, who said “let your food be your medicine”. Moreover, further clinical human studies should be conducted to verify the health benefits of bioactive peptides. The increase rate in the advancement proteomics, bioinformatics, peptide libraries, and modification strategies, will support the use of bioactive peptides as novel drugs in future clinical applications.

On the other hand, the development of healthier foods with bioactive peptides is a great scientific and technological challenge. Nowadays, it is possible to find them in the market, especially in countries like Japan. These products are accepted because peptides possess favorable pharmacokinetic profiles, tissue distribution patterns, rapid clearance from the blood, non-target tissues, low toxicity, immunogenicity and good solubility properties.

Some techniques have been widely used to improve the nutritional quality of the protein of maize, common bean,

amaranth, chia and quinoa; however, only in amaranth has tried to raise the profile of the bioactive peptides, but, indirectly have increased in each case that became transformation with the gene that encodes for any of the mentioned seed storage protein.

Acknowledgments

We thank Talia Hernandez-Perez for her kind and helpful assistance. This project was partially supported by Consejo Nacional de Ciencia y Tecnología (CONACyT-Mexico).

References

- Akira, S., S. Uematsu, and O. Takeuchi. 2006. Pathogen recognition and innate immunity. *Cell* 124 (4):783–801. doi:10.1016/j.cell.2006.02.015.
- Alemán, A., B. Giménez, E. Pérez-Santín, M. C. Gómez-Guillén, and P. Montero. 2011a. Contribution of Leu and Hyp residues to antioxidant and ace-inhibitory activities of peptide sequences isolated from squid gelatin hydrolysate. *Food Chemistry* 125 (2):334–41. doi:10.1016/j.foodchem.2010.08.058.
- Alemán, A., E. Pérez-Santín, S. Bordenave-Juchereau, I. Arnaudín, M. Gómez-Guillén, and P. Montero. 2011b. Squid gelatin hydrolysates with antihypertensive, anticancer and antioxidant activity. *Food Research International* 44 (4):1044–51. doi:10.1016/j.foodres.2011.03.010.
- Aloisi, I., L. Parrotta, K. B. Ruiz, C. Landi, L. Bini, G. Cai, S. Biondi, and D. Duca, S.. 2016. New insight into quinoa seed quality under salinity: Changes in proteomic and amino acid profiles, phenolic content, and antioxidant activity of protein extracts. *Frontiers in Plant Science* 7:656. doi:10.3389/fpls.2016.00656.
- Altorf-Van Der Kuil, W., M. F. Engberink, E. J. Brink, M. A. Van Baak, S. J. Bakker, G. Navis, P. Van't Veer, and J. M. Geleijnse. 2010. Dietary protein and blood pressure: A systematic review. *Plos One* 5 (8): e12102. doi:10.1371/journal.pone.0012102.
- Aluko, R., and E. Monu. 2003. Functional and bioactive properties of quinoa seed protein hydrolysates. *Journal of Food Science* 68 (4):1254–58. doi:10.1111/j.1365-2621.2003.tb09635.x.
- Aluko, R. E.. 2015. Antihypertensive peptides from food proteins. *Annual Review of Food Science and Technology* 6:235–62. doi:10.1146/annurev-food-022814-015520.
- Alvarez-Jubete, L., E. K. Arendt, and E. Gallagher. 2010a. Nutritive value of pseudocereals and their increasing use as functional gluten-free ingredients. *Trends in Food Science & Technology* 21 (2):106–13. doi:10.1016/j.tifs.2009.10.014.
- Alvarez-Jubete, L., H. Wijngaard, E. Arendt, and E. Gallagher. 2010b. Polyphenol composition and in vitro antioxidant activity of amaranth, quinoa buckwheat and wheat as affected by sprouting and baking. *Food Chemistry* 119 (2):770–78. doi:10.1016/j.foodchem.2009.07.032.
- Alves, L. F., M. S. Rocha, and C. C. F. Gomes. 2008. Avaliação da qualidade protéica da quinua real (*Chenopodium quinoa* willd) através de métodos biológicos. *e-Scientia* 1 (1):1–16.
- Anderson, H. J., J. E. Coleman, R. J. Andersen, and M. Roberge. 1996. Cytotoxic peptides hemiasterlin, hemiasterlin a and hemiasterlin b induce mitotic arrest and abnormal spindle formation. *Cancer Chemotherapy and Pharmacology* 39 (3):223–26. doi:10.1007/s002800050564.
- Ando, H., Y.-C. Chen, H. Tang, M. Shimizu, K. Watanabe, and T. Mitsunaga. 2002. Food components in fractions of quinoa seed. *Food Science and Technology Research* 8 (1):80–84. doi:10.3136/fstr.8.80.
- Aragão, F., L. Barros, A. Brasileiro, S. Ribeiro, F. Smith, J. Sanford, J. Faria, and E. Rech. 1996. Inheritance of foreign genes in transgenic bean (*Phaseolus vulgaris* L.) co-transformed via particle bombardment. *Theoretical and Applied Genetics* 93 (1–2):142–50. doi:10.1007/BF00225739.
- Aragão, F. J., M. F. G. De Sa, E. R. Almeida, E. S. Gander, and E. L. Rech. 1992. Particle bombardment-mediated transient expression of a brazil nut methionine-rich albumin in bean (*Phaseolus vulgaris* L.). *Plant Molecular Biology* 20 (2):357–59. doi:10.1007/BF00014508.
- Aragão, F. J. L., L. M. G. Barros, M. De Sousa, M. Grossi De Sá, E. Almeida, E. Gander, and E. Rech. 1999. Expression of a methionine-rich storage albumin from the brazil nut (*Bertholletia excelsa* hbk, *lecyrthidaceae*) in transgenic bean plants (*Phaseolus vulgaris* L., Fabaceae). *Genetics and Molecular Biology* 22 (3):445–49. doi:10.1590/S1415-47571999000300026.
- Araki, M., and T. Ishii. 2015. Towards social acceptance of plant breeding by genome editing. *Trends in Plant Science* 20 (3):145–49. doi:10.1016/j.tplants.2015.01.010.
- Arano-Varela, H., J. Dominguez-Dominguez, and O. Paredes-López. 2012. Effect of environmental conditions on the expression levels of a recombinant 11s amaranth globulin in *Escherichia coli*. *Recent Patents on Biotechnology* 6 (1):23–31. doi:10.2174/187220812799789181.
- Arihara, K., Y. Nakashima, T. Mukai, S. Ishikawa, and M. Itoh. 2001. Peptide inhibitors for angiotensin I-converting enzyme from enzymatic hydrolysates of porcine skeletal muscle proteins. *Meat Science* 57 (3):319–24. doi:10.1016/S0309-1740(00)00108-X.
- Armstrong, W. B., A. R. Kennedy, X. S. Wan, J. Atiba, C. E. McLaren, and F. L. Meyskens. 2000. Single-dose administration of Bowman-Birk inhibitor concentrate in patients with oral leukoplakia. *Cancer Epidemiology, Biomarkers & Prevention* 9 (1):43–47.
- Aubrecht, E., and P. Biacs. 2001. Characterization of buckwheat grain proteins and its products. *Acta Aliment* 30 (1):71–80. doi:10.1556/AAlim.30.2001.1.8.
- Ayerza, R. 2009. The seed's protein and oil content, fatty acid composition, and growing cycle length of a single genotype of chia (*Salvia hispanica* L.) as affected by environmental factors. *Journal of Oleo Science* 58 (7):347–54. doi:10.5650/jos.58.347.
- Azuma, N., H. Suda, H. Iwasaki, N. Yamagata, T. Saeki, R. Kanamoto, and K. Iwami. 2000. Antitumorogenic effects of several food proteins in a rat model with colon cancer and their reverse correlation with plasma bile acid concentration. *Journal of Nutritional Science and Vitaminology* 46 (2):91–96. doi:10.3177/jnsv.46.91.
- Bagga, S., D. Sutton, J. D. Kemp, and C. Sengupta-Gopalan. 1992. Constitutive expression of the β -phaseolin gene in different tissues of transgenic alfalfa does not ensure phaseolin accumulation in non-seed tissue. *Plant Molecular Biology* 19 (6):951–58. doi:10.1007/BF00040527.
- Bai, R., G. R. Petit, and E. Hamel. 1990. Dolastatin 10, a powerful cytostatic peptide derived from a marine animal: Inhibition of tubulin polymerization mediated through the vinca alkaloid binding domain. *Biochemical Pharmacology* 39 (12):1941–49. doi:10.1016/0006-2952(90)90613-P.
- Barba De La Rosa, A. P., A. Barba Montoya, P. Martínez-Cuevas, B. Hernández-Ledesma, M. F. León-Galván, A. De León-Rodríguez, and C. González. 2010. Tryptic amaranth glutelin digests induce endothelial nitric oxide production through inhibition of ace: Antihypertensive role of amaranth peptides. *Nitric Oxide* 23 (2):106–11. doi:10.1016/j.niox.2010.04.006.
- Barba De La Rosa, A. P., J. Gueguen, O. Paredes-Lopez, and G. Viroben. 1992. Fractionation procedures, electrophoretic characterization, and amino acid composition of amaranth seed proteins. *Journal of Agricultural and Food Chemistry* 40 (6):931–36. doi:10.1021/jf00018a002.
- Barba De La Rosa, A. P., A. Herrera-Estrella, S. Utsumi, and O. Paredes-López. 1996. Molecular characterization, cloning and structural analysis of a cDNA encoding an amaranth globulin. *Journal of Plant Physiology* 149 (5):527–32. doi:10.1016/S0176-1617(96)80329-4.
- Barrio, D. A., and M. C. Añón. 2010. Potential antitumor properties of a protein isolate obtained from the seeds of *Amaranthus mantegazzianus*. *European Journal of Nutrition* 49 (2):73–82. doi:10.1007/s00394-009-0051-9.
- Bellamy, W., M. Takase, K. Yamauchi, H. Wakabayashi, K. Kawase, and M. Tomita. 1992. Identification of the bactericidal domain of lactoferrin. *Biochim. Biophys. Acta Prot. Struct. Mol. Enzymol* 1121 (1–2):130–36. doi:10.1016/0167-4838(92)90346-F.
- Bellucci, M., F. De Marchis, I. Nicoletti, and S. Arcioni. 2007. Zeolin is a recombinant storage protein with different solubility and stability properties according to its localization in the endoplasmic reticulum or in the chloroplast. *Journal of Biotechnology* 131 (2):97–105. doi:10.1016/j.jbiotec.2007.06.004.
- Becker, R., E. Wheeler, K. Lorenz, A. Stafford, O. Grosjean, A. Betschart, and R. Saunders. 1981. A compositional study of Amaranth Grain.

- Journal of Food Science* 46 (4):1175–80. doi:10.1111/j.1365-2621.1981.tb03018.x.
- Bhargava, A., S. Shukla, and D. Ohri. 2006. *Chenopodium quinoa*—an indian perspective. *Industrial Crops and Products* 23 (1):73–87. doi:10.1016/j.indcrop.2005.04.002.
- Bhargava, A., S. Shukla, and D. Ohri. 2007. Genetic variability and interrelationship among various morphological and quality traits in quinoa (*Chenopodium quinoa* willd.). *Field Crops Research* 101 (1):104–16. doi:10.1016/j.fcr.2006.10.001.
- Bhat, Z., S. Kumar, and H. F. Bhat. 2015. Bioactive peptides of animal origin: A review. *Journal of Food Science and Technology* 52 (9):5377–92. doi:10.1007/s13197-015-1731-5.
- Bhat, Z. F., S. Kumar, and H. F. Bhat. 2017. Antihypertensive peptides of animal origin: A review. *Critical Reviews in Food Science and Nutrition* 57 (3):566–78. doi:10.1080/10408398.2014.898241.
- Bhutia, S. K., and T. K. Maiti. 2008. Targeting tumors with peptides from natural sources. *Trends in Biotechnology* 26 (4):210–17. doi:10.1016/j.tibtech.2008.01.002.
- Blanco-Míguez, A., A. Gutiérrez-Jácome, M. Pérez-Pérez, G. Pérez-Rodríguez, S. Catalán-García, F. Fdez-Riverola, A. Lourenço, and B. Sánchez. 2016. From amino acid sequence to bioactivity: Scientific evidence on antitumor peptides. *Protein Science* 25:1084–95. doi:10.1002/pro.2927.
- Borges, J., R. Bonomo, C. Paula, L. Oliveira, and M. Cesário. 2010. Physico-chemical and nutritional characteristics and uses of quinoa (*Chenopodium quinoa* willd.). *Temas Agrários* 15 (1):9–23.
- Borghi, C., and A. F. Cicero. 2017. Nutraceuticals with a clinically detectable blood pressure—lowering effect: A review of available randomized clinical trials and their meta-analyses. *British Journal of Clinical Pharmacology* 83 (1):163–71. doi:10.1111/bcp.12902.
- Bouglé, D., and S. Bouhallab. 2017. Dietary bioactive peptides: Human studies. *Critical Reviews in Food Science and Nutrition* 57 (2):335–43. doi:10.1080/10408398.2013.873766.
- Bressani, R., and L. A. Garcia-Vela. 1990. Protein fractions in amaranth grain and their chemical characterization. *Journal of Agricultural and Food Chemistry* 38 (5):1205–09. doi:10.1021/jf00095a010.
- Brinegar, C., and S. Goundan. 1993. Isolation and characterization of chenopodin, the 11s seed storage protein of quinoa (*Chenopodium quinoa*). *Journal of Agricultural and Food Chemistry* 41 (2):182–85. doi:10.1021/jf00026a006.
- Brinegar, C., B. Sine, and L. Nwokocha. 1996. High-cysteine 2s seed storage proteins from quinoa (*Chenopodium quinoa*). *Journal of Agricultural and Food Chemistry* 44 (7):1621–23. doi:10.1021/jf950830+.
- Broekaert, W. F., B. P. Cammue, M. F. De Bolle, K. Thevissen, G. W. De Samblanx, R. W. Osborn, and K. Nielson. 1997. Antimicrobial peptides from plants. *Critical Reviews in Plant Sciences* 16 (3):297–323. doi:10.1080/07352689709701952.
- Broekaert, W. F., W. Marien, F. R. Terras, M. F. De Bolle, P. Proost, J. Van Damme, L. Dillen, M. Claeys, and S. B. Rees. 1992. Antimicrobial peptides from *Amaranthus caudatus* seeds with sequence homology to the cysteine/glycine-rich domain of chitin-binding proteins. *Biochemistry* 31 (17):4308–14. doi:10.1021/bi00132a023.
- Brown, A. P., J. Kroon, D. Swarbreck, M. Febrer, T. R. Larson, I. A. Graham, M. Caccamo, and A. R. Slabas. 2012. Tissue-specific whole transcriptome sequencing in castor, directed at understanding triacylglycerol lipid biosynthetic pathways. *Plos One*. 7 (2):e30100. doi:10.1371/journal.pone.0030100.
- Bustos, M. M., F. A. Kalkan, K. A. Vandenbosch, and T. C. Hall. 1991. Differential accumulation of four phaseolin glycoforms in transgenic tobacco. *Plant Molecular Biology* 16 (3):381–395. doi:10.1007/BF00023990.
- Butteiger, D. N., A. A. Hibberd, N. J. McGraw, N. Napawan, J. M. Hall-Porter, and E. S. Krul. 2016. Soy protein compared with milk protein in a western diet increases gut microbial diversity and reduces serum lipids in golden syrian hamsters. *Journal of Nutrition* 146 (4):697–705. doi:10.3945/jn.115.224196.
- Cam, A., and E. G. De Mejia. 2012. Rgd-peptide lunasin inhibits Akt-mediated NF- κ B activation in human macrophages through interaction with the α V β 3 integrin. *Molecular Nutrition & Food Research* 56 (10):1569–81. doi:10.1002/mnfr.201200301.
- Campos-Vega, R., T. García-Gasca, R. N. Guevara-Gonzalez, M. Ramos-Gomez, B. D. Oomah, and G. Loarca-Piña. 2012. Human gut flora-fermented nondigestible fraction from cooked bean (*Phaseolus vulgaris* L.) modifies protein expression associated with apoptosis, cell cycle arrest, and proliferation in human adenocarcinoma colon cancer cells. *Journal of Agricultural and Food Chemistry* 60 (51):12443–450. doi:10.1021/jf303940r.
- Canabady-Rochelle, L. L., K. Selmececi, S. Collin, A. Pasc, L. Muhr, and S. Boschi-Muller. 2018. Spr screening of metal chelating peptides in a hydrolysate for their antioxidant properties. *Food Chemistry* 239:478–85. doi:10.1016/j.foodchem.2017.06.116.
- Capitani, M., V. Spotorno, S. Nolasco, and M. Tomás. 2012. Physicochemical and functional characterization of by-products from chia (*Salvia hispanica* L.) seeds of argentina. *LWT- Food Science and Technology* 45 (1):94–102. doi:10.1016/j.lwt.2011.07.012.
- Capriotti, A. L., C. Cavaliere, S. Piovesana, S. Stampachiachiere, S. Ventura, R. Zenezini Chiozzi, and A. Laganà. 2015. Characterization of quinoa seed proteome combining different protein precipitation techniques: Improvement of knowledge of nonmodel plant proteomics. *Journal of Separation Science* 38 (6):1017–25. doi:10.1002/jssc.201401319.
- Carrasco-Castilla, J., A. J. Hernández-Álvarez, C. Jiménez-Martínez, C. Jacinto-Hernández, M. Alaiz, J. Girón-Calle, J. Vioque, and G. Dávila-Ortiz. 2012. Antioxidant and metal chelating activities of peptide fractions from phaseolin and bean protein hydrolysates. *Food Chemistry* 135 (3), 1789–95. doi:10.1016/j.foodchem.2012.06.016.
- Castro, L., and B. A. Freeman. 2001. Reactive oxygen species in human health and disease. *Nutrition (Burbank, Los Angeles County, Calif.)* 17 (2):161–65. doi:10.1016/S0899-9007(00)00570-0.
- Castro-Martínez, C., S. Luna-Suárez, and O. Paredes-López. 2012. Overexpression of a modified protein from amaranth seed in *escherichia coli* and effect of environmental conditions on the protein expression. *Journal of Biotechnology* 158 (1–2):59–67. doi:10.1016/j.jbiotec.2011.12.012.
- Chakraborty, S., N. Chakraborty, and A. Datta. 2000. Increased nutritive value of transgenic potato by expressing a nonallergenic seed albumin gene from *Amaranthus hypochondriacus*. *Proceedings of the National Academy of Sciences of the United States of America*. 97 (7):3724–29. doi:10.1073/pnas.97.7.3724.
- Chappell, J., and M. J. Chrispeels. 1986. Transcriptional and posttranscriptional control of phaseolin and phytohemagglutinin gene expression in developing cotyledons of *Phaseolus vulgaris*. *Plant Physiology* 81 (1):50–54. doi:10.1104/pp.81.1.50.
- Chang, H.-C., D. Lewis, C.-Y. Tung, L. Han, S. M. Henriquez, L. Voiles, I. P. Lupov, D. Pelloso, A. L. Sinn, and K. E. Pollok. 2014. Soy peptide lunasin in cytokine immunotherapy for lymphoma. *Cancer Immunology, Immunotherapy* 63 (3):283–95. doi:10.1007/s00262-013-1513-8.
- Chen, K., and P. S. Conti. 2010. Target-specific delivery of peptide-based probes for pet imaging. *Advanced Drug Delivery Reviews* 62 (11):1005–22. doi:10.1016/j.addr.2010.09.004.
- Cheng, F. Y., Y. T. Liu, T. C. Wan, L. C. Lin, and R. Sakata. 2008. The development of angiotensin i-converting enzyme inhibitor derived from chicken bone protein. *Animal Science Journal* 79 (1):122–28. doi:10.1111/j.1740-0929.2007.00507.x.
- Cheung, H.-S., F.-L. Wang, M. A. Ondetti, E. F. Sabo, and D. W. Cushman. 1980. Binding of peptide substrates and inhibitors of angiotensin-converting enzyme. Importance of the cooh-terminal dipeptide sequence. *Journal of Biological Chemistry* 255 (2):401–07.
- Cheung, R. C. F., T. B. Ng, and J. H. Wong. 2015. Marine peptides: Bioactivities and applications. *Marine Drugs* 13 (7):4006–43. doi:10.3390/md13074006.
- Cicero, A., B. Gerocarni, L. Laghi, and C. Borghi. 2011a. Blood pressure lowering effect of lactotripeptides assumed as functional foods: A meta-analysis of current available clinical trials. *Journal of Human Hypertension* 25 (7):425–36. doi:10.1038/jhh.2010.85.
- Cicero, A., B. Gerocarni, L. Laghi, and C. Borghi. 2011b. Blood pressure lowering effect of lactotripeptides assumed as functional foods: A meta-analysis of current available clinical trials. *Journal of Human Hypertension* 25 (7):425. doi:10.1038/jhh.2010.85.
- Cicero, A., M. Rosticci, S. D'addato, F. Fogacci, R. Urso, and C. Borghi. 2016. Effects of a combined blood pressure and lipid-lowering

- nutraceutical on mildly hypertensive hypercholesterolemic subjects: A pilot, double-blind, placebo-controlled. *Journal of Human Hypertension* 34:e312. doi:10.1097/01.hjh.0000492253.96038.c3.
- Cicero, A. F., and A. Colletti. 2015. Nutraceuticals and blood pressure control: Results from clinical trials and meta-analyses. *High Blood Pressure & Cardiovascular Prevention* 22 (3):203–13. doi:10.1007/s40292-015-0081-8.
- Cicero, A. F., F. Fogacci, and A. Colletti. 2017. Potential role of bioactive peptides in prevention and treatment of chronic diseases: A narrative review. *British Journal of Pharmacology* 174 (11):1378–94. doi:10.1111/bph.13608.
- Comai, S., A. Bertazzo, L. Bailoni, M. Zancato, C. V. Costa, and G. Allegri. 2007. The content of proteic and nonproteic (free and protein-bound) tryptophan in quinoa and cereal flours. *Food Chemistry* 100 (4):1350–55. doi:10.1016/j.foodchem.2005.10.072.
- Conforti, F., G. Statti, M. R. Loizzo, G. Sacchetti, F. Poli, and F. Menichini. 2005. In vitro antioxidant effect and inhibition of α -amylase of two varieties of *Amaranthus caudatus* seeds. *Biological & Pharmaceutical Bulletin* 28 (6):1098–102. doi:10.1248/bpb.28.1098.
- Corradini, E., P. S. Curti, A. B. Meniqueti, A. F. Martins, A. F. Rubira, and E. C. Muniz. 2014. Recent advances in food-packing, pharmaceutical and biomedical applications of zein and zein-based materials. *International Journal of Molecular Sciences* 15 (12):22438–470. doi:10.3390/ijms151222438.
- Craik, D. J., D. P. Fairlie, S. Liras, and D. Price. 2013. The future of peptide-based drugs. *Chemical Biology and Drug Design* 81 (1):136–47. doi:10.1111/cbdd.12055.
- Damiens, E., I. El Yazidi, J. Mazurier, I. Duthille, G. Spik, and Y. Boily-Marer. 1999. Lactoferrin inhibits G1 cyclin-dependent kinases during growth arrest of human breast carcinoma cells. *Journal of Cellular Biochemistry* 74 (3):486–98. doi:10.1002/(SICI)1097-4644(19990901)74:3<486::AID-JCB16>3.0.CO;2-6.
- Davalos, A., M. Miguel, B. Bartolome, and R. Lopez-Fandino. 2004. Antioxidant activity of peptides derived from egg white proteins by enzymatic hydrolysis. *Journal of Food Protection* 67 (9):1939–44. doi:10.4315/0362-028X-67.9.1939.
- Dave, L. A., M. Hayes, C. A. Montoya, S. M. Rutherford, and P. J. Moughan. 2016. Human gut endogenous proteins as a potential source of angiotensin-I-converting enzyme (ACE-I)-, renin inhibitory and antioxidant peptides. *Peptides* 76:30–44. doi:10.1016/j.peptides.2015.11.003.
- De Mejia, E. G., and V. P. Dia. 2010. The role of nutraceutical proteins and peptides in apoptosis, angiogenesis, and metastasis of cancer cells. *Cancer and Metastasis Reviews* 29 (3):511–28. doi:10.1007/s10555-010-9241-4.
- De Mejia, E. G., M. D. C. Valadez-Vega, R. Reynoso-Camacho, and G. Loarca-Pina. 2005. Tannins, trypsin inhibitors and lectin cytotoxicity in tepary (*Phaseolus acutifolius*) and common (*Phaseolus vulgaris*) beans. *Plant Foods for Human Nutrition* 60 (3):137–45. doi:10.1007/s11130-005-6842-0.
- Del Rio, D., A. Rodriguez-Mateos, J. P. Spencer, M. Tognolini, G. Borges, and A. Crozier. 2013. Dietary (poly) phenolics in human health: Structures, bioavailability, and evidence of protective effects against chronic diseases. *Antioxidants & Redox Signaling* 18 (14):1818–92. doi:10.1089/ars.2012.4581.
- Deng, M., W. Zhang, H. Tang, Q. Ye, Q. Liao, Y. Zhou, M. Wu, W. Xiong, Y. Zheng, and X. Guo. 2013. Lactotransferrin acts as a tumor suppressor in nasopharyngeal carcinoma by repressing akt through multiple mechanisms. *Oncogene* 32 (36):4273–83. doi:10.1038/onc.2012.434.
- Dia, V. P., and E. G. De Mejia. 2011. Lunasin potentiates the effect of oxaliplatin preventing outgrowth of colon cancer metastasis, binds to $\alpha_5\beta_1$ integrin and suppresses FAK/ERK/NF- κ B signaling. *Cancer Letters* 313 (2):167–80. doi:10.1016/j.canlet.2011.09.002.
- Dierks-Ventling, C. 1981. Storage proteins in *Zea mays* (L.): Interrelationship of albumins, globulins and zeins in the Opaque-2 mutation. *European Journal of Biochemistry Febs* 120 (1):177–82. doi:10.1111/j.1432-1033.1981.tb05686.x.
- Dowling, D. K., and L. W. Simmons. 2009. Reactive oxygen species as universal constraints in life-history evolution. *Proceedings of the Royal Society of London. Series B. Biological Sciences* 2008:1791.
- Drzewiecki, J., E. Delgado-Licon, R. Haruenkit, E. Pawelzik, O. Martin-Beloso, Y.-S. Park, S.-T. Jung, S. Trakhtenberg, and S. Gorinstein. 2003. Identification and differences of total proteins and their soluble fractions in some pseudocereals based on electrophoretic patterns. *Journal of Agricultural and Food Chemistry* 51 (26):7798–804. doi:10.1021/jf030322x.
- Duvick, J., T. Rood, A. G. Rao, and D. R. Marshak. 1992. Purification and characterization of a novel antimicrobial peptide from maize (*Zea mays* L.) kernels. *Journal of Biological Chemistry* 267 (26):18814–820.
- Dyer, J. M., J. W. Nelson, and N. Murai. 1993. Strategies for selecting mutation sites for methionine enhancement in the bean seed storage protein phaseolin. *Journal of Protein Chemistry* 12 (5):545–60. doi:10.1007/BF01025119.
- Elitsur, Y., and G. Luk. 1991. Beta-casomorphin (BCM) and human colonic lamina propria lymphocyte proliferation. *Clinical and Experimental Immunology* 85 (3):493–97. doi:10.1111/j.1365-2249.1991.tb05755.x.
- Escudero, N., F. Zirulnik, N. Gomez, S. Mucciarelli, and M. Gimenez. 2006. Influence of a protein concentrate from *Amaranthus cruentus* seeds on lipid metabolism. *Experimental Biology and Medicine* 231 (1):50–59. doi:10.1177/153537020623100106.
- Escuredo, O., M. I. G. Martín, G. W. Moncada, S. Fischer, and J. M. H. Hierro. 2014. Amino acid profile of the quinoa (*Chenopodium quinoa* willd.) using near infrared spectroscopy and chemometric techniques. *Journal of Cereal Science* 60 (1):67–74. doi:10.1016/j.jcs.2014.01.016.
- Estilai, A., A. Hashemi, and K. Truman. 1990. Chromosome number and meiotic behavior of cultivated chia, *Salvia hispanica* (Lamiaceae). *Hortscience* 25 (12):1646–47.
- Evans, J. L., I. D. Goldfine, B. A. Maddux, and G. M. Grodsky. 2003. Are oxidative stress-activated signaling pathways mediators of insulin resistance and β -cell dysfunction?. *Diabetes* 52 (1):1–8. doi:10.2337/diabetes.52.1.1.
- Evans, R. J., and S. L. Bandemer. 1967. Nutritive value of legume seed proteins. *Journal of Agricultural and Food Chemistry* 15:439–43. doi:10.1021/jf60151a011.
- FAO. 1985. *Energy and protein requirements*. World Health Organization, Rome.
- Fast, J., A.-K. Mossberg, H. Nilsson, C. Svanborg, M. Akke, and S. Linse. 2005. Compact oleic acid in hamlet. *FEBS Letters* 579 (27):6095–100. doi:10.1016/j.febslet.2005.08.089.
- Ferreira, T. A., and J. A. G. Areas. 2010. Calcium bioavailability of raw and extruded amaranth grains. *Food Science and Technology* 30 (2):532–38. doi:10.1590/S0101-20612010000200037.
- Fitzgerald, R. J., and B. A. Murray. 2006. Bioactive peptides and lactic fermentations. *International Journal of Dairy Technology* 59 (2):118–25. doi:10.1111/j.1471-0307.2006.00250.x.
- Foltz, M., E. E. Meynen, V. Bianco, C. Van Platerink, T. M. Koning, and J. Klok. 2007. Angiotensin converting enzyme inhibitory peptides from a lactotripeptide-enriched milk beverage are absorbed intact into the circulation. *Journal of Nutrition* 137 (4):953–58. doi:10.1093/jn/137.4.953.
- Fritz, M., B. Vecchi, G. Rinaldi, and M. C. Añón. 2011. Amaranth seed protein hydrolysates have in vivo and in vitro antihypertensive activity. *Food Chemistry* 126 (3):878–84. doi:10.1016/j.foodchem.2010.11.065.
- Fuentes, F., E. Martinez, P. Hinrichsen, E. Jellen, and P. Maughan. 2009. Assessment of genetic diversity patterns in chilean quinoa (*Chenopodium quinoa* willd.) germplasm using multiplex fluorescent microsatellite markers. *Conservation Genetics* 10 (2):369–77. doi:10.1007/s10592-008-9604-3.
- Galvez, A. F., N. Chen, J. Macasieb, and O. Ben. 2001. Chemopreventive property of a soybean peptide (lunasin) that binds to deacetylated histones and inhibits acetylation. *Cancer Research* 61 (20):7473–78.
- Gangopadhyay, N., M. B. Hossain, D. K. Rai, and N. P. Brunton. 2015. A review of extraction and analysis of bioactives in oat and barley and scope for use of novel food processing technologies. *Molecules (Basel, Switzerland)* 20 (6):10884–909. doi:10.3390/molecules200610884.
- García, M., P. Puchalska, C. Esteve, and M. Marina. 2013. Vegetable foods: A cheap source of proteins and peptides with antihypertensive, antioxidant, and other less occurrence bioactivities. *Talanta* 106:328–49. doi:10.1016/j.talanta.2012.12.041.

- García-Mora, P., E. Peñás, J. Frias, H. Zieliński, W. Wiczowski, D. Zielińska, and C. Martínez-Villaluenga. 2016. High-pressure-assisted enzymatic release of peptides and phenolics increases angiotensin converting enzyme I inhibitory and antioxidant activities of pinto bean hydrolysates. *Journal of Agricultural and Food Chemistry* 64 (8):1730–40. doi:10.1021/acs.jafc.5b06080.
- García-Olmedo, F., A. Molina, J. M. Alamillo, and P. Rodríguez-Palenzuela. 1998. Plant defense peptides. *Peptide Science* 47 (6):479–91. doi:10.1002/(SICI)1097-0282(1998)47:6<479::AID-BIP6>3.0.CO;2-K.
- Germán-Báez, L. J., A. Cruz-Mendivil, S. Medina-Godoy, J. Milán-Carrillo, C. Reyes-Moreno, and A. Valdez-Ortiz. 2014. Expression of an engineered acidic-subunit 11S globulin of amaranth carrying the antihypertensive peptides VY, in transgenic tomato fruits. *Plant Cell, Tissue and Organ Culture* 118 (2):305–12. doi:10.1007/s11240-014-0483-7.
- Gómez-Favela, M. A., R. Gutiérrez-Dorado, E. O. Cuevas-Rodríguez, V. A. Canizalez-Román, C. del Rosario León-Sicaire, J. Milán-Carrillo, and C. Reyes-Moreno. 2017. Improvement of chia seeds with antioxidant activity, GABA, essential amino acids, and dietary fiber by controlled germination bioprocess. *Plant Foods for Human Nutrition* 72 (4):345–52. doi:10.1007/s11130-017-0631-4.
- Gorinstein, S., J. Drzewiecki, E. Delgado-Licon, E. Pawelzik, A. L. M. Ayala, O. J. Medina, R. Haruenkit, and S. Trakhtenberg. 2005. Relationship between dicotyledone-amaranth, quinoa, fagopyrum, soybean and monocots-sorghum and rice based on protein analyses and their use as substitution of each other. *European Food Research and Technology* 221 (1):69–77. doi:10.1007/s00217-005-1208-2.
- Gorinstein, S., E. Pawelzik, E. Delgado-Licon, R. Haruenkit, M. Weisz, and S. Trakhtenberg. 2002. Characterisation of pseudocereal and cereal proteins by protein and amino acid analyses. *Journal of the Science of Food and Agriculture* 82:886–91. doi:10.1002/jsfa.1120.
- Gupta, S., P. Kapoor, K. Chaudhary, A. Gautam, R. Kumar, G. P. Raghava, and O. S. D. D. Consortium. 2013. *In silico* approach for predicting toxicity of peptides and proteins. *Plos One* 8 (9):e73957. doi:10.1371/journal.pone.0073957.
- Guzmán-Maldonado, S. H., and O. Paredes-López. 1998. Functional products of plants indigenous to latin america: Amaranth, quinoa, common beans, and botanicals. In *Functional foods: Biochemical and processing aspects*, ed. G. Mazza, 293–328. Lancaster, Pennsylvania, U.S.A: Technomic Publishing Company.
- Hancock, J., R. Desikan, and S. Neill. 2001. *Role of reactive oxygen species in cell signalling pathways*. Portland: Portland Press Limited.
- Haque, E., R. Chand, and S. Kapila. 2008. Biofunctional properties of bioactive peptides of milk origin. *Food Reviews International* 25 (1):28–43. doi:10.1080/87559120802458198.
- Hartmann, R., and H. Meisel. 2007. Food-derived peptides with biological activity: From research to food applications. *Current Opinion in Biotechnology* 18 (2):163–69. doi:10.1016/j.copbio.2007.01.013.
- Hata, I., S. Higashiyama, and H. Otani. 1998. Identification of a phosphopeptide in bovine α s1-casein digest as a factor influencing proliferation and immunoglobulin production in lymphocyte cultures. *Journal of Dairy Research* 65 (4):569–78. doi:10.1017/S0022029998003136.
- Hatzoglou, A., E. Bakogeorgou, C. Hatzoglou, P.-M. Martin, and E. Castanas. 1996. Antiproliferative and receptor binding properties of α - and β -casomorphins in the T47D human breast cancer cell line. *European Journal of Pharmacology* 310 (2–3):217–23. doi:10.1016/0014-2999(96)00339-1.
- Hernandez-Ledesma, B., C. C. Hsieh, and B. O De Lumen. 2013. Chemopreventive properties of peptide lunasin: A review. *Protein and Peptide Letters* 20 (4):424–32.
- Hernandez-Ledesma, B., and C.-C. Hsieh. 2017. Chemopreventive role of food-derived proteins and peptides: A review. *Critical Reviews in Food Science and Nutrition* 57 (11):2358–76. doi:10.1080/10408398.2015.1057632.
- Hernandez-Ledesma, B., C.-C. Hsieh, and O. Ben. 2009a. Lunasin, a novel seed peptide for cancer prevention. *Peptides* 30 (2):426–30. doi:10.1016/j.peptides.2008.11.002.
- Hernandez-Ledesma, B., C.-C. Hsieh, and O. Ben. 2009b. Antioxidant and anti-inflammatory properties of cancer preventive peptide lunasin in raw 264.7 macrophages. *Biochemical and Biophysical Research Communications* 390 (3):803–08. doi:10.1016/j.bbrc.2009.10.053.
- Hoffman, L. M., D. D. Donaldson, R. Bookland, K. Rashka, and E. M. Herman. 1987. Synthesis and protein body deposition of maize 15-kD zein in transgenic tobacco seeds. *Embo Journal* 6 (11):3213.
- Holmer-Jensen, J., T. Karhu, L. S. Mortensen, S. B. Pedersen, K.-H. Herzig, and K. Hermansen. 2011. Differential effects of dietary protein sources on postprandial low-grade inflammation after a single high fat meal in obese non-diabetic subjects. *Nutrition Journal* 10 (1):115. doi:10.1186/1475-2891-10-115.
- Hsieh, C., B. Hernández-Ledesma, and B. O. De Lumen. 2010. Soybean peptide lunasin suppresses in vitro and in vivo 7, 12-Dimethylbenz [a] anthracene-induced tumorigenesis. *Journal of Food Science* 75 (9):311–16. doi:10.1111/j.1750-3841.2010.01861.x.
- Hsu, K.-C., E. C. Li-Chan, and C.-L. Jao. 2011. Antiproliferative activity of peptides prepared from enzymatic hydrolysates of tuna dark muscle on human breast cancer cell line MCF-7. *Food Chemistry* 126 (2):617–22. doi:10.1016/j.foodchem.2010.11.066.
- Hu, X., L. Song, L. Huang, Q. Zheng, and R. Yu. 2012. Antitumor effect of a polypeptide fraction from arca subcrenata in vitro and in vivo. *Marine Drugs* 10 (12):2782–94. doi:10.3390/md10122782.
- Huang, S., W. R. Adams, Q. Zhou, K. P. Malloy, D. A. Voyles, J. Anthony, A. L. Kriz, and M. H. Luethy. 2004. Improving nutritional quality of maize proteins by expressing sense and antisense zein genes. *Journal of Agricultural and Food Chemistry* 52 (7):1958–64. doi:10.1021/jf0342223.
- Iqbal, M. A. 2015. An assessment of quinoa (*Chenopodium quinoa* willd.) potential as a grain crop on marginal lands in pakistan. *American-Eurasian Journal Of Agricultural & Environmental Sciences* 15 (1):16–23.
- Jacobsen, S.-E., A. Mujica, and C. Jensen. 2003. The resistance of quinoa (*Chenopodium quinoa* willd.) to adverse abiotic factors. *Food Reviews International* 19 (1–2):99–109. doi:10.1081/FRI-120018872.
- Jamboonsri, W., T. D. Phillips, R. L. Geneve, J. P. Cahill, and D. F. Hildebrand. 2012. Extending the range of an ancient crop, *Salvia hispanica* L.—a new ω 3 source. *Genetic Resources and Crop Evolution* 59 (2):171–78. doi:10.1007/s10722-011-9673-x.
- James, L. E. A. 2009. Quinoa (*Chenopodium quinoa* willd.): Composition, chemistry, nutritional, and functional properties. *Advances in Food and Nutrition Research* 58:1–31. doi:10.1016/S1043-4526(09)58001-1.
- Jancurová, M., L. Minarovicová, and A. Dandar. 2009. Quinoa—a review. *Czech Journal of Food Science* 27 (2):71–79.
- Jang, A., and M. Lee. 2005. Purification and identification of angiotensin converting enzyme inhibitory peptides from beef hydrolysates. *Meat Science* 69 (4):653–61. doi:10.1016/j.meatsci.2004.10.014.
- Jarvis, D. E., Y. S. Ho, D. J. Lightfoot, S. M. Schmöckel, B. Li, T. J. Borm, H. Ohyanagi, K. Mineta, C. T. Michell, and N. Saber. 2017. The genome of *Chenopodium quinoa*. *Nature* 542 (7641):307–12. doi:10.1038/nature21370.
- Jauhainen, T., H. Vapaatalo, T. Poussa, S. Kyrönpalo, M. Rasmussen, and R. Korpela. 2005. Lactobacillus helveticus fermented milk lowers blood pressure in hypertensive subjects in 24-h ambulatory blood pressure measurement. *American Journal of Hypertension* 18 (12):1600–605. doi:10.1016/j.amjhyper.2005.06.006.
- Johansen, J. S., A. K. Harris, D. J. Rychly, and A. Ergul. 2005. Oxidative stress and the use of antioxidants in diabetes: Linking basic science to clinical practice. *Cardiovascular Diabetology* 4 (1):5. doi:10.1186/1475-2840-4-5.
- Kang, T. H., C.-P. Mao, L. He, Y.-C. Tsai, K. Liu, V. La, T.-C. Wu, and C.-F. Hung. 2012. Tumor-targeted delivery of IL-2 by NKG2D leads to accumulation of antigen-specific CD8⁺ T cells in the tumor loci and enhanced anti-tumor effects. *Plos One* 7 (4):e35141. doi:10.1371/journal.pone.0035141.
- Kannan, A., N. S. Hettiarachchi, J. O. Lay, and R. Liyanage. 2010. Human cancer cell proliferation inhibition by a pentapeptide isolated and characterized from rice bran. *Peptides* 31 (9):1629–34. doi:10.1016/j.peptides.2010.05.018.
- Katayama, K., H. E. Anggraeni, T. Mori, A. M. Ahmmed, S. Kawahara, M. Sugiyama, T. Nakayama, M. Maruyama, and M. Muguruma. 2007. Porcine skeletal muscle troponin is a good source of peptides with angiotensin-I converting enzyme inhibitory activity and antihypertensive effects in spontaneously hypertensive rats. *Journal of Agricultural and Food Chemistry* 56 (2):355–60. doi:10.1021/jf071408j.

- Katayama, K., M. Tomatsu, H. Fuchu, M. Sugiyama, S. Kawahara, K. Yamauchi, Y. Kawamura, and M. Muguruma. 2003. Purification and characterization of an angiotensin I-converting enzyme inhibitory peptide derived from porcine troponin C. *Animal Science Journal* 74 (1):53–58. doi:10.1046/j.1344-3941.2003.00086.x.
- Kauffman, C. S. 1992. Realizing the potential of grain amaranth. *Food Reviews International* 8 (1):5–21. Available from: <https://doi.org/10.1080/87559129209540927>. doi:10.1080/87559129209540927.
- Kawasaki, T., E. Seki, K. Osajima, M. Yoshida, K. Asada, T. Matsui, and Y. Osajima. 2000. Antihypertensive effect of Valyl-Tyrosine, a short chain peptide derived from sardine muscle hydrolyzate, on mild hypertensive subjects. *Journal of Human Hypertension* 14 (8):519.
- Kitts, D. D., and K. Weiler. 2003. Bioactive proteins and peptides from food sources. Applications of bioprocesses used in isolation and recovery. *Current Pharmaceutical Design* 9 (16):1309–23.
- Kobayashi, Y., J. Kovacs-Nolan, T. Matsui, and Y. Mine. 2015. The anti-atherosclerotic dipeptide, Trp-His, reduces intestinal inflammation through the blockade of L-type Ca^{2+} channels. *Journal of Agricultural and Food Chemistry* 63 (26):6041–50.
- Kong, B., and Y. L. Xiong. 2006. Antioxidant activity of zein hydrolysates in a liposome system and the possible mode of action. *Journal of Agricultural and Food Chemistry* 54 (16):6059–68.
- Korhonen, H., and A. Pihlanto. 2003. Food-derived bioactive peptides—opportunities for designing future foods. *Current Pharmaceutical Design* 9 (16):1297–308.
- Kovacs-Nolan, J., H. Zhang, M. Ibuki, T. Nakamori, K. Yoshiura, P. V. Turner, T. Matsui, and Y. Mine. 2012. The pept1-transportable soy tripeptide vpy reduces intestinal inflammation. *Biochimica et Biophysica Acta (BBA)* 1820 (11):1753–63.
- Kozioł, M. 1992. Chemical composition and nutritional evaluation of quinoa (*Chenopodium quinoa* willd.). *Journal of Food Composition and Analysis* 5 (1):35–68.
- Kozioł, M. 1990. Composición química. In *Quinoa hacia su cultivo comercial*, 137–59. Quito: Latinreco SA.
- Lafarga, T., and M. Hayes. 2017. Bioactive protein hydrolysates in the functional food ingredient industry: Overcoming current challenges. *Food Reviews International* 33 (3):217–46.
- Lahov, E., and W. Regelson. 1996. Antibacterial and immunostimulating casein-derived substances from milk: Casacidin, isracidin peptides. *Food and Chemical Toxicology* 34 (1):131–45.
- Lam, S. K., and T. B. Ng. 2011. Apoptosis of human breast cancer cells induced by hemagglutinin from *Phaseolus vulgaris* Cv. Legumi secchi. *Food Chemistry* 126 (2):595–602.
- Lammi, C., C. Zanoni, G. M. Scigliuolo, A. D'amato, and A. Arnoldi. 2014. Lupin peptides lower low-density lipoprotein (ldl) cholesterol through an up-regulation of the LDL receptor/sterol regulatory element binding protein 2 (SREBP2) pathway at HEPG2 cell line. *Journal of Agricultural and Food Chemistry* 62 (29):7151–59.
- Larkins, B. A., K. Pedersen, M. D. Marks, and D. R. Wilson. 1984. The zein proteins of maize endosperm. *Trends in Biochemical Sciences* 9 (7):306–08.
- Laveti, D., M. Kumar, R. Hemalatha, R. Sistla, V. Gm Naidu, V. Talla, V. Verma, N. Kaur, and R. Nagpal. 2013. Anti-inflammatory treatments for chronic diseases: A review. *Inflammation & Allergy - Drug Targets* 12 (5):349–61.
- Le Maux, S., A. B. Nongonierma, and R. J. Fitzgerald. 2015. Improved short peptide identification using HILIC-MS/MS: Retention time prediction model based on the impact of amino acid position in the peptide sequence. *Food Chemistry* 173:847–54.
- Lee, S., J. Xie, and X. Chen. 2010. Peptides and peptide hormones for molecular imaging and disease diagnosis. *Chemical Reviews* 110 (5):3087–111.
- Lin, L.-Z., J. M. Harnly, M. S. Pastor-Corrales, and D. L. Luthria. 2008. The polyphenolic profiles of common bean (*Phaseolus vulgaris* L.). *Food Chemistry* 107 (1):399–410.
- Lipkin, A., V. Anisimova, A. Nikonorova, A. Babakov, E. Krause, M. Bienert, E. Grishin, and T. Egorov. 2005. An antimicrobial peptide Ar-AMP from amaranth (*Amaranthus retroflexus* L.) seeds. *Phytochemistry* 66 (20):2426–31.
- Liu, Q., B. Kong, Y. L. Xiong, and X. Xia. 2010. Antioxidant activity and functional properties of porcine plasma protein hydrolysate as influenced by the degree of hydrolysis. *Food Chemistry* 118 (2):403–10.
- Liu, R., L. Xing, Q. Fu, G.-H. Zhou, and W.-G. Zhang. 2016. A review of antioxidant peptides derived from meat muscle and by-products. *Antioxidants* 5 (3):32.
- Lule, V. K., S. Garg, S. D. Pophaly, and S. K. Tomar. 2015. Potential health benefits of lunasin: A multifaceted soy-derived bioactive peptide. *Journal of Food Science* 80 (3):R485–94.
- Luna-Suárez, S., S. Medina-Godoy, A. Cruz-Hernández, and O. Paredes-López. 2008. Expression and characterization of the acidic subunit from 11S amaranth seed protein. *Biotechnology Journal* 3 (2):209–19.
- Luna-Suárez, S., S. Medina-Godoy, A. Cruz-Hernández, and O. Paredes-López. 2010. Modification of the amaranth 11S globulin storage protein to produce an inhibitory peptide of the angiotensin I converting enzyme, and its expression in *Escherichia coli*. *Journal of Biotechnology* 148:240–47.
- Luna-Vital, D. A., L. Mojica, E. G. De Mejía, S. Mendoza, and G. Loarca-Piña. 2015. Biological potential of protein hydrolysates and peptides from common bean (*Phaseolus vulgaris* L.): A review. *Food Research International* 76:39–50.
- Ma, J., F. Huang, H. Lin, and X. Wang. 2013. Isolation and purification of a peptide from bullacta exarata and its impact of apoptosis on prostate cancer cell. *Marine Drugs* 11 (1):266–73.
- Ma, Y.-H., W.-Z. Cheng, F. Gong, A.-L. Ma, Q.-W. Yu, J.-Y. Zhang, C.-Y. Hu, X.-H. Chen, and D.-Q. Zhang. 2008. Active chinese mistletoe lectin-55 enhances colon cancer surveillance through regulating innate and adaptive immune responses. *World Journal of Gastroenterology: Wjg* 14 (34):5274.
- Maestri, E., M. Marmiroli, and N. Marmiroli. 2016. Bioactive peptides in plant-derived foodstuffs. *Journal of Proteomics* 147:140–155.
- Mainieri, D., M. Rossi, M. Archinti, M. Bellucci, F. De Marchis, S. Vavasori, A. Pompa, S. Arcioni, and A. Vitale. 2004. Zeolin. A new recombinant storage protein constructed using maize γ -zein and bean phaseolin. *Plant Physiology* 136 (3):3447–3456.
- Majumder, K., S. Chakrabarti, J. S. Morton, S. Panahi, S. Kaufman, S. T. Davidge, and J. Wu. 2013. Egg-derived tri-peptide IRW exerts antihypertensive effects in spontaneously hypertensive rats. *Plos One* 8 (11):e82829.
- Majumder, K., Y. Mine, and J. Wu. 2016. The potential of food protein-derived anti-inflammatory peptides against various chronic inflammatory diseases. *Journal of the Science of Food and Agriculture* 96 (7):2303–11.
- Malaguti, M., G. Dinelli, E. Leoncini, V. Bregola, S. Bosi, A. F. Cicero, and S. Hrelia. 2014. Bioactive peptides in cereals and legumes: Agronomical, biochemical and clinical aspects. *International Journal of Molecular Sciences* 15 (11):21120–35.
- Maldonado-Cervantes, E., H. J. Jeong, F. León-Galván, A. Barrera-Pacheco, A. De León-Rodríguez, E. G. De Mejía, B. O. De Lumen, and A. P. Barba De La Rosa. 2010. Amaranth lunasin-like peptide internalizes into the cell nucleus and inhibits chemical carcinogen-induced transformation of NIH-3T3 cells. *Peptides* 31 (9):1635–42.
- Malkowicz, S. B., W. G. McKenna, D. J. Vaughn, X. S. Wan, K. J. Probert, K. Rockwell, S. H. Marks, A. J. Wein, and A. R. Kennedy. 2001. Effects of Bowman-Birk inhibitor concentrate (BBIC) in patients with benign prostatic hyperplasia. *The Prostate* 48 (1):16–28.
- Mallory, M., R. Hall, A. McNabb, D. Pratt, E. Jellen, and P. Maughan. 2008. Development and characterization of microsatellite markers for the grain amaranths. *Crop Science* 48:1098–106.
- Marcone, M. 1999a. Possible nutritional implications of varietal influence on the 7S/11S seed globulin ratios in amaranth. *Plant Foods for Human Nutrition* 54 (4):375–80.
- Marcone, M. F. 1999b. Biochemical and biophysical properties of plant storage proteins: A current understanding with emphasis on 11S seed globulins. *Food Research International* 32 (2):79–92.
- Marques, M. R., R. Freitas, A. C. C. Carlos, É. S. Siguemoto, G. G. Fontanari, and J. Arêas. 2015. Peptides from cowpea present antioxidant activity, inhibit cholesterol synthesis and its solubilisation into micelles. *Food Chemistry* 168:288–93.
- Martínez-Cruz, O., F. Cabrera-Chávez, and O. Paredes-López. 2014. Biochemical characteristics, and nutraceutical and technological uses of amaranth globulins. In *Globulins: Biochemistry, production and role in*

- immunity, ed. S. D. Milford, 41–70. New York: Nova Science Publishers.
- Martínez-Cruz, O., and O. Paredes-López. 2014. Phytochemical profile and nutraceutical potential of chia seeds (*Salvia hispanica* L.) by ultra high performance liquid chromatography. *Journal of Chromatography A* 1346:43–48.
- Maruyama, S., S. Miyoshi, T. Kaneko, and H. Tanaka. 1989. Angiotensin I-converting enzyme inhibitory activities of synthetic peptides related to the tandem repeated sequence of a maize endosperm protein. *Agricultural and Biological Chemistry* 53 (4):1077–81.
- Matoba, N., Y. Yamada, H. Usui, R. Nakagiri, and M. Yoshikawa. 2001. Designing potent derivatives of ovokinin (2–7), an anti-hypertensive peptide derived from ovalbumin. *Bioscience, Biotechnology, and Biochemistry* 65 (3):736–39.
- Matsuoka, Y., Y. Vigouroux, M. M. Goodman, J. Sanchez, E. Buckler, and J. Doebley. 2002. A single domestication for maize shown by multilocus microsatellite genotyping. *Proceedings of the National Academy of Sciences of the United States of America* 99 (9):6080–84.
- Matzke, M., M. Susani, A. Binns, E. Lewis, I. Rubenstein, and A. Matzke. 1984. Transcription of a zein gene introduced into sunflower using a ti plasmid vector. *Embo Journal* 3 (7):1525.
- Mccann, K., B. Shiell, W. Michalski, A. Lee, J. Wan, H. Roginski, and M. Coventry. 2006. Isolation and characterisation of a novel antibacterial peptide from bovine α s1-casein. *International Dairy Journal* 16 (4):316–23.
- Medina-Godoy, S., N. C. Nielsen, and O. Paredes-López. 2004. Expression and characterization of a His-tagged 11S seed globulin from *Amaranthus hypochondriacus* in *Escherichia coli*. *Biotechnology Progress* 20 (6):1749–56.
- Medina-Godoy, S., S. K. Rodriguez-Yanez, N. A. Bobadilla, R. Perez-Villalva, R. Valdez-Ortiz, E. Hong, S. Luna-Suarez, O. Paredes-López, and A. Valdez-Ortiz. 2013. Antihypertensive activity of AMC3, an engineered 11S amaranth globulin expressed in *Ascherichia coli*, in spontaneously hypertensive rats. *Journal of Functional Foods* 5 (3):1441–49.
- Medina-Godoy, S., A. Valdez-Ortiz, M. E. Valverde, and O. Paredes-López. 2006. Endoplasmic reticulum-retention c-terminal sequence enhances production of an 11s seed globulin from *Amaranthus hypochondriacus* in *Pichia pastoris*. *Biotechnology Journal* 1 (10):1085–92.
- Meisel, H. 2004. Multifunctional peptides encrypted in milk proteins. *Bio-factors* 21 (1–4):55–61.
- Mendonça, S., P. H. Saldiva, R. J. Cruz, and J. A. Arêas. 2009. Amaranth protein presents cholesterol-lowering effect. *Food Chemistry* 116 (3):738–42.
- Miguel, M., R. López-Fandino, M. Ramos, and A. Aleixandre. 2005. Short-term effect of egg-white hydrolysate products on the arterial blood pressure of hypertensive rats. *British Journal of Nutrition* 94 (5):731–37.
- Minihane, A. M., S. Vinoy, W. R. Russell, A. Baka, H. M. Roche, K. M. Tuohy, J. L. Teeling, E. E. Blaak, M. Fenech, and D. Vauzour. 2015. Low-grade inflammation, diet composition and health: Current research evidence and its translation. *British Journal of Nutrition* 114 (7):999–1012.
- Minkiewicz, P., M. Darewicz, A. Iwaniak, J. Sokołowska, P. Starowicz, J. Bucholska, and M. Hryniewicz. 2015. Common amino acid subsequences in a universal proteome—relevance for food science. *International Journal of Molecular Sciences* 16 (9):20748–73.
- Minkiewicz, P., J. Dziuba, A. Iwaniak, M. Dziuba, and M. Darewicz. 2008. Biopep database and other programs for processing bioactive peptide sequences. *Journal of Aoac International* 91 (4):965–80.
- Miranda, M., J. Delatorre-Herrera, A. Vega-Gálvez, E. Jorquera, I. Quispe-Fuentes, and E. A. Martínez. 2014. Antimicrobial potential and phytochemical content of six diverse sources of quinoa seeds (*Chenopodium quinoa* willd.). *Journal of Agricultural Science* 5 (11):1015.
- Miranda, M., A. Vega-Gálvez, E. A. Martínez, J. López, R. Marín, M. Aranda, and F. Fuentes. 2013. Influence of contrasting environments on seed composition of two quinoa genotypes: Nutritional and functional properties. *Chilean Journal of Agricultural Research* 73 (2):108–16.
- Miranda, M., A. Vega-Gálvez, E. Uribe, J. López, E. Martínez, M. J. Rodríguez, I. Quispe, and K. Di Scala. 2011. Physico-chemical analysis, antioxidant capacity and vitamins of six ecotypes of Chilean quinoa (*Chenopodium quinoa* willd.). *Procedia Food Science* 1:1439–46.
- Miyoshi, S., H. Ishikawa, T. Kaneko, F. Fukui, H. Tanaka, and S. Maruyama. 1991. Structures and activity of angiotensin-converting enzyme inhibitors in an α -zein hydrolysate. *Agricultural and Biological Chemistry* 55 (5):1313–18.
- Mrakar, S. G., M. Turinek, M. Jakop, M. Bavec, and F. Bavec. 2010. Grain amaranth as an alternative and perspective crop in temperate climate. *Journal of Geographical Sciences* 5 (1):135–45.
- Mojica, L., and E. G. De Mejía. 2015. Characterization and comparison of protein and peptide profiles and their biological activities of improved common bean cultivars (*Phaseolus vulgaris* L.) from Mexico and Brazil. *Plant Foods for Human Nutrition* 70 (2):105–12.
- Mojica, L., and E. G. De Mejía. 2016. Optimization of enzymatic production of anti-diabetic peptides from black bean (*Phaseolus vulgaris* L.) proteins, their characterization and biological potential. *Food & Function* 7 (2):713–27.
- Mojica, L., D. A. Luna-Vital, and G. D. Mejía, E. 2017. Characterization of peptides from common bean protein isolates and their potential to inhibit markers of type-2 diabetes, hypertension and oxidative stress. *Journal of the Science of Food and Agriculture* 97 (8):2401–10.
- Möller, N. P., K. E. Scholz-Ahrens, N. Roos, and J. Schrezenmeir. 2008. Bioactive peptides and proteins from foods: Indication for health effects. *European Journal of Nutrition* 47 (4):171–82.
- Montoya, C. A., J.-P. Lallès, S. Beebe, and P. Leterme. 2010. Phaseolin diversity as a possible strategy to improve the nutritional value of common beans (*Phaseolus vulgaris*). *Food Research International* 43 (2):443–49.
- Montoya-Rodríguez, A., M. A. Gómez-Favela, C. Reyes-Moreno, J. Milán-Carrillo, and E. González De Mejía. 2015. Identification of bioactive peptide sequences from amaranth (*Amaranthus hypochondriacus*) seed proteins and their potential role in the prevention of chronic diseases. *Comprehensive Reviews in Food Science and Food Safety* 14 (12):139–58.
- Montoya-Rodríguez, A., E. G. Mejía, V. P. Dia, C. Reyes-Moreno, and J. Milán-Carrillo. 2014. Extrusion improved the anti-inflammatory effect of amaranth (*Amaranthus hypochondriacus*) hydrolysates in lps-induced human THP-1 macrophage-like and mouse raw 264.7 macrophages by preventing activation of NF- κ B signaling. *Molecular Nutrition & Food Research* 58 (5):1028–41.
- Mookherjee, N., and R. Hancock. 2007. Cationic host defence peptides: Innate immune regulatory peptides as a novel approach for treating infections. *Cellular and Molecular Life Sciences* 64 (7–8):922.
- Mora, L., and M. Hayes. 2015. Cardioprotective cryptides derived from fish and other food sources: Generation, application, and future markets. *Journal of Agricultural and Food Chemistry* 63 (5):1319–31.
- Morales-Camacho, J., J. Dominguez-Dominguez, and O. Paredes-López. 2013. Overexpression of a modified amaranth protein in *Escherichia coli* with minimal media and lactose as inducer. *Recent Patents on Biotechnology* 7 (1):61–70.
- Morales-Camacho, J. I., O. Paredes-López, E. Espinosa-Hernández, D. A. F. Velasco, and S. Luna-Suárez. 2016. Expression, purification and thermal stability evaluation of an engineered amaranth protein expressed in *Escherichia coli*. *Electronic Journal of Biotechnology* 22:44–51.
- Moronta, J., P. L. Smaldini, G. H. Docena, and M. C. Añon. 2016a. Peptides of amaranth were targeted as containing sequences with potential anti-inflammatory properties. *Journal of Functional Foods* 21:463–73.
- Moronta, J., P. L. Smaldini, C. A. Fossati, M. C. Añon, and G. H. Docena. 2016b. The anti-inflammatory SSEDIKE peptide from amaranth seeds modulates IgE-mediated food allergy. *Journal of Functional Foods* 25:579–87.
- Moruz, L., and L. Käll. 2016. Peptide retention time prediction. *Mass Spectrometry Reviews* 35:615–23. DOI 10.1002/mas.21488
- Motoi, H., and T. Kodama. 2003. Isolation and characterization of angiotensin I-converting enzyme inhibitory peptides from wheat gliadin hydrolysate. *Molecular Nutrition & Food Research* 47 (5):354–58.
- Mujica-Sanchez, A., S. Jacobsen, J. Izquierdo, and J. Marathe. 2001. *Quinoa (Chenopodium quinoa willd.): Ancestral cultivo andino, alimento del presente y del futuro*. Santiago: FAO.
- Muñoz, L. A., A. Cobos, O. Diaz, and J. M. Aguilera. 2013. Chia seed (*Salvia hispanica*): An ancient grain and a new functional food. *Food Reviews International* 29 (4):394–408. Available from: <https://doi.org/10.1080/87559129.2013.818014>.

- Murai, N., D. W. Sutton, M. G. Murray, J. L. Slightom, D. J. Merlo, N. A. Reichert, C. Sengupta-Gopalan, C. A. Stock, R. Baker, and J. D. Kemp. 1983. Phaseolin gene from bean is expressed after transfer to sunflower via tumor-inducing plasmid vectors. *Science* 222 (4623):476–82.
- Murray, B., and R. Fitzgerald. 2007. Angiotensin converting enzyme inhibitory peptides derived from food proteins: Biochemistry, bioactivity and production. *Current Pharmaceutical Design* 13 (8):773–91.
- Nakamura, T., T. Hirota, K. Mizushima, K. Ohki, Y. Naito, N. Yamamoto, and T. Yoshikawa. 2013. Milk-derived peptides, val-pro-pro and Ile-Pro-Pro, attenuate atherosclerosis development in apolipoprotein E-deficient mice: A preliminary study. *Journal of Medicinal Food* 16 (5):396–403.
- Nakashima, Y., K. Arihara, A. Sasaki, H. Mio, S. Ishikawa, and M. Itoh. 2002. Antihypertensive activities of peptides derived from porcine skeletal muscle myosin in spontaneously hypertensive rats. *Journal of Food Science* 67 (1):434–37.
- Nayak, S., P. Pattnaik, S. Arora, and J. Sindhu. 1999. Functional peptides from milk proteins. *Indian Dairyman* 51:29–34.
- Nieman, D. C., E. J. Cayea, M. D. Austin, D. A. Henson, S. R. Mcanulty, and F. Jin. 2009. Chia seed does not promote weight loss or alter disease risk factors in overweight adults. *Nutrition Research* 29 (6):414–18.
- Nieman, D. C., N. Gillitt, F. Jin, D. A. Henson, K. Kennerly, R. A. Shanely, B. Ore, M. Su, and S. Schwartz. 2012. Chia seed supplementation and disease risk factors in overweight women: A metabolomics investigation. *Journal of Alternative and Complementary Medicine* 18 (7):700–08.
- Nongonierma, A. B., and R. J. Fitzgerald. 2015. Bioactive properties of milk proteins in humans: A review. *Peptides* 73:20–34.
- Nongonierma, A. B., S. Le Maux, C. Dubrulle, C. Barre, and R. J. Fitzgerald. 2015. Quinoa (*Chenopodium quinoa* willd.) protein hydrolysates with in-vitro Dipeptidyl Peptidase IV (DPP-IV) inhibitory and antioxidant properties. *Journal of Cereal Science* 65:112–18. Available from: <http://www.sciencedirect.com/science/article/pii/S0733521015300357>.
- Nsimba, R. Y., H. Kikuzaki, and Y. Konishi. 2008. Antioxidant activity of various extracts and fractions of *Chenopodium quinoa* and *Amaranthus* spp. Seeds. *Food Chemistry*. 106 (2):760–66.
- Ohtani, T., G. Galili, J. C. Wallace, G. A. Thompson, and B. A. Larkins. 1991. Normal and lysine-containing zeins are unstable in transgenic tobacco seeds. *Plant Molecular Biology* 16 (1):117–28.
- Olmedilla-Alonso, B., M. M. Pedrosa, C. Cuadrado, M. Brito, and C. Asensio-Vegas. 2013. Composition of two spanish common dry beans (*Phaseolus vulgaris*), 'almonga' and 'curruquilla', and their postprandial effect in type 2 diabetics. *Journal of the Science of Food and Agriculture* 93 (5):1076–82.
- Orona-Tamayo, D., and O. Paredes-López. 2017a. Amaranth—sustainable crop for the 21st century: Food properties and nutraceutical for improving human health. In *Sustainable protein sources*, ed. S. Nadathur, 1st ed., 239–56. San Diego: Elsevier-Academic Press.
- Orona-Tamayo, D., M. E. Valverde, and O. Paredes-López. 2017b. Chia—the new golden seed for the 21st century: Nutraceutical properties and technological uses. In *Sustainable protein sources*, ed. S. Nadathur, 1st ed., 265–81. San Diego: Elsevier-Academic Press.
- Orona-Tamayo, D., M. E. Valverde, B. Nieto-Rendón, and O. Paredes-López. 2015. Inhibitory activity of chia (*Salvia hispanica* L.) protein fractions against angiotensin I-converting enzyme and antioxidant capacity. *LWT- Food Science and Technology* 64 (1):236–42.
- Orsini-Delgado, M. C., V. A. Tironi, and M. C. Añón. 2011. Antioxidant activity of amaranth protein or their hydrolysates under simulated gastrointestinal digestion. *LWT- Food Science and Technology* 44 (8):1752–60.
- Orsini-Delgado, M. C., A. Nardo, M. Pavlovic, H. Rogniaux, M. C. Añón, and V. A. Tironi. 2016. Identification and characterization of antioxidant peptides obtained by gastrointestinal digestion of amaranth proteins. *Food Chemistry* 197:1160–67.
- Orsini-Delgado, M. C. O., M. Galleano, M. C. Añón, and V. A. Tironi. 2015. Amaranth peptides from simulated gastrointestinal digestion: Antioxidant activity against reactive species. *Plant Foods for Human Nutrition* 70 (1):27–34.
- Ortiz-Martinez, M., R. Winkler, and S. García-Lara. 2014. Preventive and therapeutic potential of peptides from cereals against cancer. *Journal of Proteomics* 111:165–83.
- Oseguera-Toledo, M. E., E. G. De Mejia, and S. L. Amaya-Llano. 2015. Hard-to-cook bean (*Phaseolus vulgaris* L.) proteins hydrolyzed by alcalase and bromelain produced bioactive peptide fractions that inhibit targets of type-2 diabetes and oxidative stress. *Food Research International* 76:839–51.
- Osuna-Castro, J. A., Q. Rascón-Cruz, J. Napier, R. J. Fido, P. R. Shewry, and O. Paredes-López. 2000. Overexpression, purification, and in vitro refolding of the 11S globulin from amaranth seed in *Escherichia coli*. *Journal of Agricultural and Food Chemistry* 48 (11):5249–55.
- Otvos, L. Jr. 2016. Immunomodulatory effects of anti-microbial peptides. *Acta Microbiologica Et Immunologica Hungarica* 63 (3):257–77.
- Panchaud, A., M. Affolter, and M. Kussmann. 2012. Mass spectrometry for nutritional peptidomics: How to analyze food bioactives and their health effects. *Journal of Proteomics* 75 (12):3546–59.
- Panel, E. N. 2012. Scientific opinion on the substantiation of health claims related to Isoleucine-Proline-Proline (IPP) and Valine-Proline-Proline (VPP) and maintenance of normal blood pressure. *EFSA Journal* 10 (6):2715.
- Paredes-López, O. 1994. *Amaranth: Biology, chemistry and technology*. 1st ed. Boca Raton: CRC Press.
- Paredes-López, O., R. Mora-Escobedo, and C. Ordorica-Falomir. 1988. Isolation of amaranth proteins. *LWT* 21:59–61.
- Park, J. H., H. J. Jeong, and B. O. De Lumen. 2005. Contents and bioactivities of lunasin, Bowman-Birk inhibitor, and isoflavones in soybean seed. *Journal of Agricultural and Food Chemistry* 53 (20):7686–90.
- Pasko, P., M. Sajewicz, S. Gorinstein, and Z. Zachwieja. 2008. Analysis of selected phenolic acids and flavonoids in *Amaranthus cruentus* and *Chenopodium quinoa* seeds and sprouts by hplc. *Acta Chromatographica* 20 (4):661–72.
- Pavlik, V. 2011. The revival of amaranth as a third-millennium food. *Neuro Endocrinology Letters* 33:3–4.
- Peperkamp, M. 2014. Chia from bolivia: A modern super seed in a classic pork cycle. In *M. O. F. Affairs*, ed., 1–15. Netherland: CBI Marked Intelligence. <http://www.cbi.eu/market-information/grains-pulses/chia/>.
- Perego, S., S. Cosentino, A. Fiorilli, G. Tettamanti, and A. Ferraretto. 2012. Casein phosphopeptides modulate proliferation and apoptosis in HT-29 cell line through their interaction with voltage-operated l-type calcium channels. *Journal of Nutritional Biochemistry* 23 (7):808–16.
- Perez Espitia, P. J., N. De Fátima Ferreira Soares, J. S. Dos Reis Coimbra, N. J. De Andrade, R. Souza Cruz, A. Medeiros, and E. Antonio. 2012. Bioactive peptides: Synthesis, properties, and applications in the packaging and preservation of food. *Comprehensive Reviews in Food Science and Food Safety* 11 (2):187–204.
- Piperno, D. R., and K. V. Flannery. 2001. The earliest archaeological maize (*Zea mays* L.) from highland Mexico: New accelerator mass spectrometry dates and their implications. *Proceedings of the National Academy of Sciences of the United States of America* 98 (4):2101–03.
- Plate, A. Y., and J. Arêas. 2002. Cholesterol-lowering effect of extruded amaranth (*Amaranthus caudatus* L.) in hypercholesterolemic rabbits. *Food Chemistry* 76 (1):1–6.
- Pravst, I. 2012. Functional foods in Europe: A focus on health claims. In *Scientific, health and social aspects of the food industry*, ed. B. Valdez, Chapter 9:165–08. InTech.
- Pribylova, R., P. Kralik, B. Pisarikova, and I. Pavlik. 2008. Detection of the antimicrobial peptide gene in different *Amaranthus* species. *Biologia* 63 (2):217–20.
- Puchalska, P., M. L. Marina, and M. C. García. 2012. Development of a reversed-phase high-performance liquid chromatography analytical methodology for the determination of antihypertensive peptides in maize crops. *Journal of Chromatography A* 1234:64–71.
- Puchalska, P., M. L. Marina, and M. C. García. 2013. Development of a high-performance liquid chromatography-electrospray ionization-quadrupole-time-of-flight-mass spectrometry methodology for the determination of three highly antihypertensive peptides in maize crops. *Journal of Chromatography A* 1285:69–77.

- Quiroga, A., E. N. Martínez, H. Rognaux, A. Geairon, and M. C. Añón. 2010. Amaranth (*Amaranthus hypochondriacus*) vicilin subunit structure. *Journal of Agricultural and Food Chemistry* 58 (24):12957–63.
- Quiroga, A. V., P. Aphalo, J. L. Ventureira, E. N. Martínez, and M. C. Añón. 2012. Physicochemical, functional and angiotensin converting enzyme inhibitory properties of amaranth (*Amaranthus hypochondriacus*) 7S globulin. *Journal of the Science of Food and Agriculture* 92 (2):397–403.
- Qureshi, A. A., J. W. Lehmann, and D. M. Peterson. 1996. Amaranth and its oil inhibit cholesterol biosynthesis in 6-week-old female chickens. *Journal of Nutrition* 126 (8):1972–78.
- Rascón-Cruz, Q., S. Sinagawa-García, J. Osuna-Castro, N. Bohorova, and O. Paredes-López. 2004. Accumulation, assembly, and digestibility of amarantin expressed in transgenic tropical maize. *Theoretical and Applied Genetics* 108 (2):335–42.
- Rastogi, A., and S. Shukla. 2013. Amaranth: A new millennium crop of nutraceutical values. *Critical Reviews in Food Science and Nutrition* 53 (2):109–25.
- Ren, X., H. Ma, S. Mao, and H. Zhou. 2014. Effects of sweeping frequency ultrasound treatment on enzymatic preparations of ACE-inhibitory peptides from zein. *European Food Research and Technology* 238 (3):435–42.
- Repo-Carrasco, R., C. Espinoza, and S.-E. Jacobsen. 2003. Nutritional value and use of the andean crops quinoa (*Chenopodium quinoa*) and kañiwa (*Chenopodium pallidicaule*). *Food Reviews International* 19 (1–2):179–89.
- Repo-Carrasco-Valencia, R., J. Peña, H. Kallio, and S. Salminen. 2009. Dietary fiber and other functional components in two varieties of crude and extruded kiwicha (*Amaranthus caudatus*). *Journal of Cereal Science* 49 (2):219–24.
- Reyes-Caudillo, E., A. Tecante, and M. Valdivia-López. 2008. Dietary fibre content and antioxidant activity of phenolic compounds present in mexican chia (*Salvia hispanica* L.) seeds. *Food Chemistry* 107 (2):656–63.
- Rivillas-Acevedo, L., and M. Soriano-García. 2007a. Antifungal activity of a protean extract from *Amaranthus hypochondriacus* seeds. *Journal of the Mexican Chemical Society* 51 (3):136–40.
- Rivillas-Acevedo, L. A., and M. Soriano-García. 2007b. Isolation and biochemical characterization of an antifungal peptide from *Amaranthus hypochondriacus* seeds. *Journal of Agricultural and Food Chemistry* 55 (2):10156–61.
- Rizzello, C. G., A. Lorusso, V. Russo, D. Pinto, B. Marzani, and M. Gobetti. 2017. Improving the antioxidant properties of quinoa flour through fermentation with selected autochthonous lactic acid bacteria. *International Journal of Food Microbiology* 241:252–61.
- Roberts, P. R., J. Burney, K. W. Black, and G. P. Zaloga. 1999. Effect of chain length on absorption of biologically active peptides from the gastrointestinal tract. *Digestion* 60 (4):332–37.
- Romero-Arenas, O., M. Damián-Huato, J. Rivera-Tapia, A. Báez-Simón, M. Huerta-Lara, and E. Cabrera-Huerta. 2013. The nutritional value of beans (*Phaseolus vulgaris* L.) and its importance for feeding of rural communities in puebla, mexico. Introduction: Legumes in sustainable agriculture. *Critical Reviews in Plant Sciences* 34:2–3. Taylor & Francis.
- Rubas, W., and G. M. Grass. 1991. Gastrointestinal lymphatic absorption of peptides and proteins. *Advanced Drug Delivery Reviews* 7 (1):15–69.
- Rubiales, D., and A. Mikic. 2015. *Introduction: Legumes in sustainable agriculture*. Taylor & Francis.
- Ruiz-Lopez, N., S. Usher, O. V. Sayanova, J. A. Napier, and R. P. Haslam. 2015. Modifying the lipid content and composition of plant seeds: Engineering the production of LC-PUFA. *Applied Microbiology and Biotechnology* 99 (1):143–54.
- Ryan, E., K. Galvin, T. O'connor, A. Maguire, and N. O'brien. 2007. Phytosterol, squalene, tocopherol content and fatty acid profile of selected seeds, grains, and legumes. *Plant Foods for Human Nutrition* 62 (3):85–91.
- Sabbione, A. C., S. M. Ibañez, E. N. Martínez, M. C. Añón, and A. A. Scilingo. 2016. Antithrombotic and antioxidant activity of amaranth hydrolysate obtained by activation of an andogenous protease. *Plant Foods for Human Nutrition* 71 (2):174–82.
- Saiga, A., T. Okumura, T. Makihara, S. Katsuta, T. Shimizu, R. Yamada, and T. Nishimura. 2003. Angiotensin I-converting enzyme inhibitory peptides in a hydrolyzed chicken breast muscle extract. *Journal of Agricultural and Food Chemistry* 51 (6):1741–45.
- Salazar-Vega, I. M., M. R. Segura-Campos, L. Chel-Guerrero, and D. Betancur-Ancona. 2012. Antihypertensive and antioxidant effects of functional foods containing chia (*Salvia hispanica*) protein hydrolysates. In *Agricultural and Biological Sciences Scientific, Health and Social Aspects of the food industry*, ed. B. Valdez, 381–98. InTech.
- Salouti, M., F. Mirzaei, R. Shapouri, and A. Ahangari. 2016. Synergistic antibacterial activity of plant peptide MBP-1 and silver nanoparticles combination on healing of infected wound due to *Staphylococcus aureus*. *Jundishapur Journal of Microbiology* 9 (1):e27997.
- Sandoval-Oliveros, M. R., and O. Paredes-López. 2013. Isolation and characterization of proteins from chia seeds (*Salvia hispanica* L.). *Journal of Agricultural and Food Chemistry* 61 (1):193–201.
- Sano, J., K. Ohki, T. Higuchi, K. Aihara, S. Mizuno, O. Kajimoto, S. Nakagawa, Y. Kajimoto, and Y. Nakamura. 2005. Effect of casein hydrolysate, prepared with protease derived from *Aspergillus oryzae*, on subjects with high-normal blood pressure or mild hypertension. *Journal of Medicinal Food* 8 (4):423–30.
- Santiago-López, L., A. Hernández-Mendoza, B. Vallejo-Cordoba, V. Mata-Haro, and A. F. González-Córdova. 2016. Food-derived immunomodulatory peptides. *Journal of the Science of Food and Agriculture* 96 (11):3631–41.
- Sarmadi, B., and A. Ismail. 2010. Antioxidative peptides from food proteins: A review. *Peptides* 31 (10):1949–56.
- Scherthaner, J., M. Matzke, and A. Matzke. 1988. Endosperm-specific activity of a zein gene promoter in transgenic tobacco plants. *Embo Journal* 7 (5):1249.
- Scherz-Shouval, R., and Z. Elazar. 2011. Regulation of autophagy by ros: Physiology and pathology. *Trends in Biochemical Sciences* 36 (1):30–38.
- Schlick, G., and D. L. Bubenheim. 1993. Quinoa: An emerging new crop with potential for cells. *NASA Technical Paper*. 3422(https://www.researchgate.net/profile/David_Bubenheim/publication/24326431_Quinoa_An_emerging_new_crop_with_potential_for_CELSS/links/53fc41390cf2364ccc047708.pdf).
- Schweizer, F. 2009. Cationic amphiphilic peptides with cancer-selective toxicity. *European Journal of Pharmacology* 625 (1):190–94.
- Segura-Campos, M., L. Chel-Guerrero, D. Betancur-Ancona, and V. M. Hernandez-Escalante. 2011. Bioavailability of bioactive peptides. *Food Reviews International* 27 (3):213–26.
- Segura-Campos, M. R., F. Peralta González, L. Chel Guerrero, and D. Betancur Ancona. 2013a. Angiotensin I-converting enzyme inhibitory peptides of chia (*Salvia hispanica*) produced by enzymatic hydrolysis. *International Journal of Food Science* 8:ArticleID 158482, 8 p.
- Segura-Campos, M. R., I. M. Salazar-Vega, L. A. Chel-Guerrero, and D. A. Betancur-Ancona. 2013b. Biological potential of chia (*Salvia hispanica* L.) protein hydrolysates and their incorporation into functional foods. *Journal of Food Science and Technology* 50 (2):723–31.
- Segura-Nieto, M., A. Barba De La Rosa, and O. Paredes-López. 1994. Biochemistry of amaranth proteins. In *Biology, Chemistry and Technology*, ed. O. Paredes-López, 75–106. Boca Raton, FL: CRC Press.
- Seifert, G., P. Jesse, A. Laengler, T. Reindl, M. Lüth, S. Lobitz, G. Henze, A. Prokop, and H. N. Lode. 2008. Molecular mechanisms of mistletoe plant extract-induced apoptosis in acute lymphoblastic leukemia in vivo and in vitro. *Cancer Letters* 264 (2):218–28.
- Shadidi, M., and M. Sioud. 2003. Selective targeting of cancer cells using synthetic peptides. *Drug Resistance Updates* 6 (6):363–71.
- Shahidi, F. 2000. Antioxidants in food and food antioxidants. *Molecular Nutrition & Food Research* 44 (3):158–163.
- Shekhar, S., L. Agrawal, D. Mishra, A. K. Buragohain, M. Unnikrishnan, C. Mohan, S. Chakraborty, and N. Chakraborty. 2016. Ectopic expression of amaranth seed storage albumin modulates photoassimilate transport and nutrient acquisition in sweetpotato. *Scientific Reports* 6, Article Number:25384. doi: 10.1038/srep25384.
- Shukla, R., and M. Cheryan. 2001. Zein: The industrial protein from corn. *Industrial Crops and Products* 13 (3):171–92.

- Silva-Sánchez, C., A. B. De La Rosa, M. León-Galván, B. De Lumen, A. De Leon-Rodríguez, and E. G. De Mejía. 2008. Bioactive peptides in amaranth (*Amaranthus hypochondriacus*) seed. *Journal of Agricultural and Food Chemistry* 56 (4):1233–40.
- Sirtori, C. R., A. Arnoldi, and A. F. Cicero. 2015. Nutraceuticals for blood pressure control. *Annals of Medicine* 47 (6):447–56.
- Slightom, J. L., R. F. Drong, C. Klassy, and L. M. Hoffman. 1985. Nucleotide sequences from pbaseolln cDNA clones: the major storage proteins from *Phaseolus vulgaris* are encoded by two unique gene families. *Nucleic Acids Research* 13 (18):6483–98.
- Soares, R. A. M., S. Mendonça, L. De Castro, A. Menezes, and J. Arêas. 2015. Major peptides from amaranth (*Amaranthus cruentus*) protein inhibit HMG-CoA reductase activity. *International Journal of Molecular Sciences* 16 (2):4150–60.
- Somerville, A. D., N. Sugiyama, L. R. Manzanilla, and M. J. Schoeninger. 2016. Animal management at the ancient metropolis of Teotihuacan, Mexico: Stable isotope analysis of leporid (Cottontail and Jackrabbit) bone mineral. *Plos One* 11 (8):e0159982.
- Sousa, D. A., W. F. Porto, M. Z. Silva, T. R. Da Silva, and O. L. Franco. 2016. Influence of cysteine and tryptophan substitution on DNA-binding activity on maize α -hairpinin antimicrobial peptide. *Molecules (Basel, Switzerland)* 21 (8):1062.
- Sreedhar, R., P. Kumari, S. D. Rupwate, R. Rajasekharan, and M. Srinivasan. 2015. Exploring triacylglycerol biosynthetic pathway in developing seeds of chia (*Salvia hispanica* L.): A transcriptomic approach. *Plos One* 10 (4):e0123580.
- Stikic, R., D. Glamoclija, M. Demin, B. Vucelic-Radovic, Z. Jovanovic, D. Milojkovic-Opsenica, S.-E. Jacobsen, and M. Milovanovic. 2012. Agromonomical and nutritional evaluation of quinoa seeds (*Chenopodium quinoa* willd.) as an ingredient in bread formulations. *Journal of Cereal Science* 55 (2):132–38.
- Sun, X., R. Jiang, A. Przepiorski, S. Reddy, K. P. Palmano, and G. W. Krisanssen. 2012. “Iron-saturated” bovine lactoferrin improves the chemotherapeutic effects of tamoxifen in the treatment of basal-like breast cancer in mice. *Bmc Cancer [Electronic Resource]* 12 (1):591.
- Tagliazucchi, D., S. Martini, A. Bellesia, and A. Conte. 2015. Identification of ace-inhibitory peptides from *Phaseolus vulgaris* after in vitro gastrointestinal digestion. *International Journal of Food Sciences and Nutrition* 66 (7):774–82.
- Tandang-Silvas, M. R., C. S. Cabanos, L. D. Carrasco Peña, A. P. B. De La Rosa, J. A. Osuna-Castro, S. Utsumi, B. Mikami, and N. Maruyama. 2012. Crystal structure of a major seed storage protein, 11S proglubulin, from *Amaranthus hypochondriacus*: Insight into its physico-chemical properties. *Food Chemistry* 135 (2):819–26.
- Tang, X., Z. He, Y. Dai, Y. L. Xiong, M. Xie, and J. Chen. 2009. Peptide fractionation and free radical scavenging activity of zein hydrolysate. *Journal of Agricultural and Food Chemistry* 58 (1):587–93.
- Thundimadathil, J. 2012. Cancer treatment using peptides: Current therapies and future prospects. *Journal of Amino Acids* 2012 (2012), Article ID 967347, 13 p.
- Ting, C.-H., H.-N. Huang, T.-C. Huang, C.-J. Wu, and J.-Y. Chen. 2014. The mechanisms by which pardaxin, a natural cationic antimicrobial peptide, targets the endoplasmic reticulum and induces c-fos. *Biomaterials* 35 (11):3627–40.
- Tironi, V. A., and M. C. Añón. 2010. Amaranth proteins as a source of antioxidant peptides: Effect of proteolysis. *Food Research International* 43 (1):315–22.
- Tokede, O. A., T. A. Onabanjo, A. Yansane, J. M. Gaziano, and L. Djoussé. 2015. Soya products and serum lipids: A meta-analysis of randomised controlled trials. *British Journal of Nutrition* 114 (6):831–43.
- Toscano, L. T., C. S. O. Da Silva, L. T. Toscano, A. E. M. De Almeida, A. Da Cruz Santos, and A. S. Silva. 2014. Chia flour supplementation reduces blood pressure in hypertensive subjects. *Plant Foods for Human Nutrition* 69 (4):392–98.
- Tovar-Pérez, E., I. Guerrero-Legarreta, A. Farrés-González, and J. Soriano-Santos. 2009. Angiotensin I-converting enzyme-inhibitory peptide fractions from albumin 1 and globulin as obtained of amaranth grain. *Food Chemistry* 116 (2):437–44.
- Tucker, J. 1986. Amaranth – the once and future crop. *Bioscience* 36:9–13.
- Tyagi, A., A. Tuknait, P. Anand, S. Gupta, M. Sharma, D. Mathur, A. Joshi, S. Singh, A. Gautam, and G. P. Raghava. 2014. CancerPPD: A database of anticancer peptides and proteins. *Nucleic Acids Research* 43 (D1):D837–43.
- Udenigwe, C. C., and R. E. Aluko. 2012. Food protein-derived bioactive peptides: Production, processing, and potential health benefits. *Journal of Food Science* 77 (1):R11–24.
- Ullah, R., M. Nadeem, A. Khaliq, M. Imran, S. Mehmood, A. Javid, and J. Hussain. 2015. Nutritional and therapeutic perspectives of chia (*Salvia hispanica* L.): A review. *Journal of Food Science and Technology* 53 (4):1750–58. doi:10.1007/s13197-015-1967-0.
- Valadez-Vega, C., G. Alvarez-Manilla, L. Riverón-Negrete, A. García-Carrancá, J. A. Morales-González, C. Zuñiga-Pérez, E. Madrigal-Santillán, J. Esquivel-Soto, C. Esquivel-Chirino, and R. Villagómez-Ibarra. 2011. Detection of cytotoxic activity of lectin on human colon adenocarcinoma (SW480) and epithelial cervical carcinoma (C33-A). *Molecules (Basel, Switzerland)* 16 (3):2107–18.
- Valdez-Ortiz, A., Q. Rascón-Cruz, S. Medina-Godoy, S. R. Sinagawa-García, M. E. Valverde-González, and O. Paredes-López. 2005. One-step purification and structural characterization of a recombinant His-tag 11S globulin expressed in transgenic tobacco. *Journal of Biotechnology* 115 (4):413–23.
- Valdivia-López, M. A., and A. Tecante. 2015. Chia (*Salvia hispanica*): A review of native mexican seed and its nutritional and functional properties. In *Adv Food Nutr Res*, ed. H. Jeyakumar, 54–75. Elsevier.
- Valko, M., D. Leibfritz, J. Moncol, M. T. Cronin, M. Mazur, and J. Telser. 2007. Free radicals and antioxidants in normal physiological functions and human disease. *International Journal of Biochemistry & Cell Biology* 39 (1):44–84.
- Valverde, M. E., D. Orona-Tamayo, B. Nieto-Rendón, and O. Paredes-López. 2016. Antioxidant and antihypertensive potential of protein fractions from flour and milk substitutes from canary seeds (*Phalaris canariensis* L.). *Plant Foods for Human Nutrition* 72 (1):20–25.
- Varadhachary, A., J. S. Wolf, K. Petrak, B. W. O’malley, M. Spadaro, C. Curcio, G. Forni, and F. Pericle. 2004. Oral lactoferrin inhibits growth of established tumors and potentiates conventional chemotherapy. *International Journal of Cancer* 111 (3):398–403.
- Vasal, S. 2000. The quality protein maize story. *Food and Nutrition Bulletin* 21 (4):445–50.
- Vasco-Méndez, N. L., and O. Paredes-López. 1994. Antigenic homology between amaranth glutelins and other storage proteins. *Journal of Food Biochemistry* 18 (4):227–38.
- Vecchi, B., and M. Añón. 2009. Ace inhibitory tetrapeptides from *Amaranthus hypochondriacus* 11S globulin. *Phytochemistry* 70 (7):864–70.
- Vega-Gálvez, A., M. Miranda, J. Vergara, E. Uribe, L. Puente, and E. A. Martínez. 2010. Nutrition facts and functional potential of quinoa (*Chenopodium quinoa* willd.), an ancient andean grain: A review. *Journal of the Science of Food and Agriculture* 90 (15):2541–47.
- Velarde-Salcedo, A. J., A. Barrera-Pacheco, S. Lara-González, G. M. Montero-Morán, A. Díaz-Gois, E. G. De Mejia, and A. P. B. De La Rosa. 2013. In vitro inhibition of Dipeptidyl Peptidase IV by peptides derived from the hydrolysis of amaranth (*Amaranthus hypochondriacus* L.) proteins. *Food Chemistry* 136 (2):758–64.
- Vlieghe, P., V. Lisowski, J. Martinez, and M. Khrestchatsky. 2010. Synthetic therapeutic peptides: Science and market. *Drug Discovery Today* 15 (1):40–56.
- Vuksan, V., D. Whitham, J. L. Sievenpiper, A. L. Jenkins, A. L. Rogovik, R. P. Bazinet, E. Vidgen, and A. Hanna. 2007. Supplementation of conventional therapy with the novel grain salba (*salvia hispanica* L.) improves major and emerging cardiovascular risk factors in type 2 diabetes: Results of a randomized controlled trial. *Diabetes Care* 30 (11):2804–10.
- Wang, H., and T. Ng. 2007. Isolation and characterization of an antifungal peptide with antiproliferative activity from seeds of *Phaseolus vulgaris* Cv: ‘Spotted bean’. *Applied Microbiology and Biotechnology* 74 (1):125–30.
- Wang, J. Z., H. Zhang, M. Zhang, W. T. Yao, X. Y. Mao, and F. Z. Ren. 2008. Antioxidant activity of hydrolysates and peptide fractions of porcine plasma albumin and globulin. *Journal of Food Biochemistry* 32 (6):693–707.
- Wang, L., Q. Wang, J. Qian, Q. Liang, Z. Wang, J. Xu, S. He, and H. Ma. 2015. Bioavailability and bioavailable forms of collagen after oral

- administration to rats. *Journal of Agricultural and Food Chemistry* 63 (14):3752–56.
- Wang, S., P. Rao, and X. Ye. 2009. Isolation and biochemical characterization of a novel leguminous defense peptide with antifungal and antiproliferative potency. *Applied Microbiology and Biotechnology* 82 (1):79–86.
- Wang, Y.-K., H.-L. He, G.-F. Wang, H. Wu, B.-C. Zhou, X.-L. Chen, and Y.-Z. Zhang. 2010. Oyster (*Crassostrea gigas*) hydrolysates produced on a plant scale have antitumor activity and immunostimulating effects in BALB/C mice. *Marine Drugs* 8 (2):255–68.
- Watanabe, K., A. Ibuki, Y.-C. Chen, Y. Kawamura, and T. Mitsunaga. 2003. Composition of quinoa protein fractions. *The Japanese Society for Food Science and Technology* 50 (11):546–49.
- Watanabe, K., Y. Tsuge, M. Shimoyamada, N. Ogama, and T. Ebina. 1998. Antitumor effects of pronase-treated fragments, glycopeptides, from ovomucin in hen egg white in a double grafted tumor system. *Journal of Agricultural and Food Chemistry* 46 (8):3033–38.
- Wolin, K. Y., A. V. Patel, P. T. Campbell, E. J. Jacobs, M. L. McCullough, G. A. Colditz, and S. M. Gapstur. 2010. Change in physical activity and colon cancer incidence and mortality. *Cancer Epidemiology, Biomarkers & Prevention* 19 (12):3000–04.
- Wright, K., O. Pike, D. Fairbanks, and C. Huber. 2002. Composition of atriplex hortensis, sweet and bitter chenopodium quinoa seeds. *Journal of Food Science* 67 (4):1383–85.
- Wu, D., Y. Gao, Y. Qi, L. Chen, Y. Ma, and Y. Li. 2014. Peptide-based cancer therapy: Opportunity and challenge. *Cancer Letters* 351 (1):13–22.
- Wu, W., Q. Su, X. Xia, Y. Wang, Y. Luan, and L. An. 2008. The suaeda liaotungensis kitag betaine aldehyde dehydrogenase gene improves salt tolerance of transgenic maize mediated with minimum linear length of DNA fragment. *Euphytica* 159:17–25.
- Xue, Y., N. Yin, B. Chen, F. Liao, A. N. Win, J. Jiang, R. Wang, X. Jin, N. Lin, and Y. Chai. 2017. Molecular cloning and expression analysis of two fad2 genes from chia (*Salvia hispanica*). *Acta Physiologiae Plantarum* 39 (4):95.
- Yamada, Y., N. Matoba, H. Usui, K. Onishi, and M. Yoshikawa. 2002. Design of a highly potent anti-hypertensive peptide based on ovokin (2–7). *Bioscience, Biotechnology, and Biochemistry* 66 (6):1213–17.
- Yamamoto, N., M. Ejiri, and S. Mizuno. 2003. Biogenic peptides and their potential use. *Current Pharmaceutical Design* 9 (16):1345–55.
- Yamauchi, R., K. Ohinata, and M. Yoshikawa. 2003. B-lactotensin and neurotensin rapidly reduce serum cholesterol via NT 2 receptor. *Peptides* 24 (12):1955–61.
- Yang, R., Z. Zhang, X. Pei, X. Han, J. Wang, L. Wang, Z. Long, X. Shen, and Y. Li. 2009. Immunomodulatory effects of marine oligopeptide preparation from chum salmon (*Oncorhynchus keta*) in mice. *Food Chemistry* 113 (2):464–70.
- Yano, S., K. Suzuki, and G. Funatsu. 1996. Isolation from α -zein of thermolysin peptides with angiotensin I-converting enzyme inhibitory activity. *Bioscience, Biotechnology, and Biochemistry* 60 (4):661–63.
- Yoo, Y. C., S. Watanabe, R. Watanabe, K. Hata, K. I. Shimazaki, and I. Azuma. 1997. Bovine lactoferrin and lactoferricin, a peptide derived from bovine lactoferrin, inhibit tumor metastasis in mice. *Cancer Science* 88 (2):184–90.
- Yoshii, H., N. Tachi, R. Ohba, O. Sakamura, H. Takeyama, and T. Itani. 2001. Antihypertensive effect of ace inhibitory oligopeptides from chicken egg yolks. *Comparative Biochemistry and Physiology Part C: Pharmacology, Toxicology and Endocrinology* 128 (1):27–33.
- Yu, L., J. D. Milton, D. G. Fernig, and J. M. Rhodes. 2001. Opposite effects on human colon cancer cell proliferation of two dietary thomsen-friedenreich antigen-binding lectins. *Journal of Cellular Physiology* 186 (2):282–87.
- Yu, Z., W. Zhao, J. Liu, J. Lu, and F. Chen. 2011. QIGLF, a novel angiotensin I-converting enzyme-inhibitory peptide from egg white protein. *Journal of the Science of Food and Agriculture* 91 (5):921–26.
- Zhan, K.-X., W.-H. Jiao, F. Yang, J. Li, S.-P. Wang, Y.-S. Li, B.-N. Han, and H.-W. Lin. 2014. Reniochalistatins A–E, cyclic peptides from the marine sponge reniochalina stalagmitis. *Journal of Natural Products* 77 (12):2678–84.
- Zhang, X.-X., H. S. Eden, and X. Chen. 2012. Peptides in cancer nanomedicine: Drug carriers, targeting ligands and protease substrates. *Journal of Controlled Release* 159 (1):2–13.
- Zheng, Z., K. Sumi, K. Tanaka, and N. Murai. 1995. The bean seed storage protein [beta]-phaseolin is synthesized, processed, and accumulated in the vacuolar type-II protein bodies of transgenic rice endosperm. *Plant Physiology* 109 (3):777–86.
- Zou, C., A. Chen, L. Xiao, H. M. Muller, P. Ache, G. Haberer, M. Zhang, W. Jia, P. Deng, and R. Huang. 2017. A high-quality genome assembly of quinoa provides insights into the molecular basis of salt bladder-based salinity tolerance and the exceptional nutritional value. *Cell Research* 27 (11):1327.