



An Overview of Advanced Technologies for Selection of Probiotics and Their Expediency: A Review

Ruby Yadav & Pratyosh Shukla

To cite this article: Ruby Yadav & Pratyosh Shukla (2015): An Overview of Advanced Technologies for Selection of Probiotics and Their Expediency: A Review, Critical Reviews in Food Science and Nutrition, DOI: [10.1080/10408398.2015.1108957](https://doi.org/10.1080/10408398.2015.1108957)

To link to this article: <http://dx.doi.org/10.1080/10408398.2015.1108957>



Accepted author version posted online: 27 Oct 2015.



Submit your article to this journal [↗](#)



View related articles [↗](#)



View Crossmark data [↗](#)

An overview of advanced technologies for selection of probiotics and their expediency:**A review**

Ruby Yadav, Pratyosh Shukla*

Enzyme Technology and Protein Bioinformatics Laboratory, Department of Microbiology,

Maharshi Dayanand University, Rohtak-124001, Haryana, INDIA

*Corresponding author: pratyosh.shukla@gmail.com

Abstract

It has become easy to identify and select an appropriate microorganism with the advancement in various molecular biology and analytical techniques. The majority of the novel techniques is being implemented for the identification and characterization of microorganisms used for probiotic application. Standard microbial techniques such as biochemical testing and culture techniques routinely used for probiotic microbes screening, identification and selection. However, these standard techniques may not give complete information on the microbes that can be used for probiotic production. Furthermore, alternative molecular and analytical techniques such as 16S and 23S ribosomal DNA sequencing, RNA analysis by reverse transcriptase (RT-PCR), fluorescent in situ hybridization (FISH), quantitative analysis by real time PCR (RT-PCR or qPCR) and fluorescent activated cell sorting (FACS) are potentially used to confirm and select all types of lactic acid microorganism. All these approaches can be employed in the screening and selection of appropriate lactic acid bacteria which can be potentially used for the production

of human use probiotics in large scale fermentation. This review mainly focuses on various tools and techniques used for effective screening and selection of a better candidate bacterium for probiotic applications.

Keywords Probiotics. Quantitative analysis. Fluorescent in situ hybridization (FISH). Real time PCR (RT-PCR). Fluorescent activated cell sorting (FACS)

1. Introduction:

Probiotic microbial food supplements are live microflora used to facilitate advantageous effect on individual health by advancing the microbiological flora in the intestine. Most commonly used bacteria which have excellent probiotic properties are from human normal microflora and they belong to genera *Lactobacilli* and *Bifidobacteria*. These bacteria exert beneficial effects in the human gastrointestinal tract, such as, reduction of serum cholesterol, alleviation of lactose intolerance, anti-infection, antimicrobial, antioxidant, anti-allergic, antimutagenic, anticarcinogenic properties, blood lipid reduction and immune system stimulation (Mortazavian et al., 2008; Lee et al., 2014). Probiotic bacteria may produce a variety of inhibitory substances such as organic acids and bacteriocins, which can inhibit the pathogenic bacteria present in the human intestine. Organic acids lower the body pH and affect the growth of pathogens. Probiotics, prebiotics and synbiotics (combination of probiotics and prebiotics) are considered as food products, not drugs (Pundir et al., 2013). There has been a rapid rise in use of probiotic product in dairy applications. The effect of probiotics against allergies, diarrhoea, malnutrition, antibiotic side effects etc. provides the scientific basis for determining a dynamic future for dairy probiotics (Reid et al., 2015). The importance of probiotic lactic acid bacteria and *Bifidobacteria* in the prevention and treatment of inflammatory bowel and other associated diseases have been studied. It has been well established that probiotics act as an important prophylactic or therapeutic strategy for many mucosal and non mucosal immune-related conditions, such as celiac disease, metabolic syndrome, and diabetes (Ivanov et al., 2012). Lactic acid bacteria and *bifidobacteria*, have been characterized as probiotics that can modify the gut microflora and may be advantageous

for prevention and treatment of Inflammatory Bowel Disease (Lara et al., 2015). Probiotics can also induce some physiological effects even in healthy person. For example, in healthy adults four week consumption of *Lactobacillus paracasei* subsp. *paracasei* LC01 resulted in reduced faecal *Escherichia coli* and ammonia, and increases in *Lactobacillus*, *Bifidobacterium*, and *Roseburia intestinalis* and acetic and butyric acid (Zhang et al., 2013). The safety, functionality and quality must be fulfilled by a probiotic strain, if used as human food. The phenotypic methods have been used as an identification tool for many years to characterize a probiotic strain. These phenotypic methods include morphology examination, gram staining, biochemical methods, carbohydrate fermentation pattern, growth at different salt, pH and temperature, however temperature sometime may lead to misidentification (Margolles et al., 2009). We understand that, a successful and effective probiotic product can only be regarded as useful, if a suitable bacteria used as the product is properly identified, characterized and qualified for desired use. So, identification of probiotic strains is a chief concern for industrial, technological, and probiotic applications (Collado et al., 2006). Currently available molecular techniques such as 16S-23S rRNA sequencing, DGGE, PFGE, ARDRA contributes in a greater extent for identification, classification and characterization of these bacteria. Most commonly used identification techniques now a day's used for genotyping of probiotic bacteria are PFGE, AFLP, RAPD-PCR and Ribotyping. These identification techniques further enable identify and classify the bacteria at the strain level. Probiotic products especially for human consumption requires properly identified and well characterized microbial strains so that it does not lead to any erroneous effect to the

customers. In this review advanced as well as common techniques used recent past for identification and enumeration of probiotic bacteria are discussed.

2. History and definition of probiotics

The word probiotics consist of Latin word 'pro' means for and Greek word 'bios' means life (Lalitha, 2011). The idea of the beneficial effect of microorganisms on human health was proposed by Noble Prize winner Eli Metchnikoff. Metchnikoff has shown that *L. bulgaricus* is able to abolish pathogenic bacteria from intestinal microflora and result in health benefits (Metchnikoff, 1907). He suggested that the long life of Bulgarian peasants is correlated to high consumption of the fermented milk products as the lactic acid bacteria present in these products replaces the harmful intestinal bacteria (Singh et al., 2011). But the actual concept of probiotics was introduced by Lilly and Stillwell as after their work probiotics characterized as beneficial microorganisms that promotes the growth of other bacteria (Lilly et al., 1965). Parker defined probiotic microorganisms and their substances to facilitate the intestinal microflora (Parker, 1974; Goyal et al., 2013). Probiotics have been defined by Fuller as live microbial food supplements which beneficially affects the host by improving its intestinal microflora (Fuller, 1989). WHO and FAO defined probiotics as live microorganisms, its administration in adequate amount has an advantageous effect for human health. To assess the efficacy of probiotics in humans, it is necessary to have an understanding of the probiotic strains, therefore novel methods are needed for selection and characterization of these strains.

3. Selection criteria

For the selection of a microbial strain that to be used as probiotic should have certain characteristics includes (Fig. 1) Safety, production/manufacturing, viability during storage, resistance to gastric acid, bile tolerant, adherence to gut epithelial cells, colonization to the gastrointestinal tract (GIT), production of antimicrobial substances, ability to stimulate the immune response, and ability to influence metabolic activities such as vitamin production of a host (Savodago et al., 2006). The initial screening and selection of probiotics also includes testing of genotypic and phenotypic stability, plasmid stability, carbohydrate utilization patterns, antibiotic resistance patterns, ability to inhibit known pathogens, and immunogenicity. A probiotic strain should survive, proliferate and colonize in GIT, it should neither be pathogenic nor trigger allergic response to the host. Furthermore, probiotics should be easily culturable on a larger scale and must be able to resist technological manipulations such as heating and low oxygen conditions during packaging. As human stomach is highly acidic due to the presence of HCL, hence acidic environment may act as a barrier for probiotics. Therefore, the probiotic microflora must be able to survive gastric as well as the bile in the upper digestive tract before their entry into the small intestine (Lee et al., 2014). The probiotic species *L. casei* and *L. acidophilus* survives well in the acidic conditions of artificial gastric juice at pH 3.0 at 37⁰C while *L. delbrueckii* ssp. *bulgaricus* does not (Eric et al., 2015). Microencapsulation is the most noteworthy promising and competent technology these days used for the safeguarding of probiotics against unfavorable environmental conditions like oxygen, low pH, gastric juice, bile salts, salinity, heat and cold shocks (Tripathi et al., 2014). Most common encapsulating materials are alginate, Chitosan, gums (locust bean, gellan gum, xanthan gum, etc.), gelatin, carrageenan, whey protein, starch. The

encapsulating materials significantly improved the viability of probiotic microorganisms during storage, processing, and passage in the gastrointestinal tract (Riaz et al., 2013). When exposed to simulated gastrointestinal juice, survivability of probiotic strains *Lactobacillus acidophilus* PTCC1643 and *Lactobacillus rhamnosus* PTCC1637 trapped in alginate coating, was considerably higher than their uncoated forms (Mokarram et al., 2009). A bakery product, pan bread was prepared by using sodium alginate and whey protein-encapsulated and air-dried *L. rhamnosus* GG (Soukoulis et al., 2014).

4. Health benefits

Interaction between probiotics and cells of the immune system are essential for maintaining mucosal tissue homeostasis and innate immunity. The gastrointestinal tract of human contains bacterial species which acts as a barrier against potential pathogens and help in regulating the immune response of the body. Human gut microflora comprises mainly Gram positive Firmicutes (60–80%) and Gram-negative Bacteroidetes (20–40%). *Lactobacillus* and *Bifidobacterium* are the predominant genres in the human intestine, both have a long history of safe use and are considered as GRAS (Backhed, 2012). *Lactobacillus* is predominant in the small intestine and *Bifidobacterium* in the large intestine. However, species belonging to the genera *Lactococcus*, *Enterococcus* (*Enterococcus faecium*), *Streptococcus* (*Streptococcus salivarius*), yeasts (e.g. *Saccharomyces cerevisiae* and *Saccharomyces boulardii*) and filamentous fungi (e.g. *Aspergillus oryzae*) are also used as probiotics due to their health promoting effects. Some commonly used probiotic strains and their commercial applications is given in Table 1.

Gut microflora differ among individuals and can vary according to genotype, age, health condition, diet, and exposure to antibiotics to host (Henao et al., 2013). The presence of bacterial species in the gastrointestinal tract is influenced by many factors such as diet, intestinal pH, bacterial adhesion, secretion of mucin and bacterial antagonism etc. It has been established by Neish, 2009 that changes in any of these factors can affect the environment that supports the gut microbiota, thus altering the number of microorganisms available to maintain intestinal homeostasis. This whole event cascade leads to excessive energy production and calorie extraction, with negative effects on metabolism and obesity which is a major cause of liver disease (Neish, 2009). Probiotics have been shown to have favorable effects when used to treat several liver diseases by reducing the production of bacterial toxins and by modulating autoimmune responses, intestinal permeability, and the inflammatory response. The mechanisms through which probiotic bacteria alter the progression of liver diseases are thought to relate to modifications in the composition of the natural intestinal microbiota and its activity. For example, supplementation with *Lactobacillus paracasei* decreases the number of *Enterobacteriaceae* and *Enterococcus spp*, and promotes the growth of species of *Lactobacillus*, *Bifidobacterium* and *Bacteroides* (Norberto et al., 2015). Bacteriocins produced by Probiotics, can slow down the growth of pathogenic bacteria. Organic acids produced can penetrate the cell membrane and inhibit nutrients transport and ATPase activity (Levy et al., 2006). Altered microbiota, termed dysbiosis, could lead to altered immune functions and increased risk of diseases, gastrointestinal disorders and triggers many inflammatory bowel diseases. Some evidences of probiotic bacteria are improving antibiotic-associated diarrhoea, acute infectious diarrhoea, irritable bowel

syndrome, pouchitis, ulcerative colitis, necrotizing enterocolitis and Crohn's disease (Chong, 2014). Probiotics have been validated to improve antioxidant condition, enhance immune resistance against viral infections and tumours and also boost GIT barrier functionality (Toucheffeu et al., 2014; Vieira et al., 2013). Probiotics have been used as adjuvant treatment for many diseases that affect the gastrointestinal tract, such as inflammatory bowel disease, cancer, *Helicobacter pylori* infection, vaginosis, allergy, and lactose intolerance (Hill et al., 2014; McFarland, 2014; Narbona et al., 2014). It has been reported that allergic Rhinitis which is triggered by grass pollen might be mitigated using *L. paracasei* subsp Paracasei LP-33 (Costa et al., 2014). Further, a recent study suggested that IRT5 probiotics (a mixture of 5 probiotics) could suppress various inflammatory disorders, including encephalomyelitis, a T-cell mediated inflammatory autoimmune disease of the central nervous system (Kwon et al., 2013). It is illustrious that probiotics restores the gut homeostasis by involving regeneration of the intestine, which repairs infection associated damage (Vieira et al., 2013). Furthermore, the improved significance of probiotics in cancer prophylaxis and therapy has been widely studied (Patel et al., 2013). Bifidobacteria have been reported to produce a protective coat of exopolysaccharides that could help bacteria to survive against high acidic condition and high bile concentration of the gastrointestinal tract without a detrimental effect on their viability (Prasanna et al., 2014). Probiotics provide various health benefits (Fig. 2) which includes balancing intestinal microflora, improvement of the immune system, provide protection against gastrointestinal pathogens, reduction of serum cholesterol level, prevention of food allergy, protection against colon and urinary bladder cancer. Probiotics help in modulation of different signalling pathways of innate immunity through the Toll like

receptors. Different strains of probiotic *Lactobacilli* can generate different intensity of immune responses (Maldonado et al., 2015).

5. Techniques used for Strain Identification

5.1 Screening by Invitro testing

In vitro testing methods are very useful to gain knowledge on the mechanism of probiotic strain. The Invitro assay is recommended for the testing of probiotics as it is used for human trials. The Potential probiotic microflora screened using in vitro methods have shown resistance to gastric acids (Casey et al., 2004), bile salt tolerance (Hoque et al., 2010) antimicrobial activity against pathogenic bacteria, bile salt hydrolase activity (Franz et al., 2001), adherence to human epithelial cells and resistance to 0.4% phenol (Kumar et al., 2012).

5.2 Molecular techniques for identification

5.2.1 Species specific identification techniques

Various molecular methods have been developed for identification and categorization of probiotic bacteria summarized in table 1. These techniques can be further divided into nucleic acid based and complete cell activity based identification. However, most frequently used tool is nucleic acid based because of its specificity and high throughput potential which includes ribotyping, RAPD, AFLP and PFGE. These methods have been extensively used for

identification and genotyping of bifidobacteria and LAB (lactic acid bacteria). Some of the most common and relevant methods of identifications are described below.

5.2.1.1 Amplified ribosomal DNA restriction analysis (ARDRA)

ARDRA is an excellent tool for identification and differentiation of bacteria which are isolated from composite environment and related groups of bacteria. The identification of strains at the species level is first described in 1992 (Vanechoutte et al., 1992). It involves enzymatic amplification followed by digestion with restriction enzyme patterns, the restriction pattern obtained for new strain is compared with patterns obtained from the reference strain (Donelli et al., 2013). It is a good molecular tool for identification of human *bifidobacterium* species. A study showed that the *Bifidobacterium* species present in the intestinal tract of human can be differentiated using a six restriction pattern combination from 511bp to 525 bp 16S ribosomal fragment with *Taq1*, *Sau3A1*, *Rsa1*, *Alu1*, *Sau961*, *Nci1* enzymes, respectively (Venema et al., 2003). ARDRA has been shown as the most appropriate method for a quick discrimination between different species of *Propionibacteria* used as probiotics (Dherbecourt et al., 2006). In addition, ARDRA has been successfully used for identification of *Lactobacilli*, *Bifidobacteria* and *Streptococci* species. (Collado et al., 2007). Recent studies on 28 faecal samples of *B. longum* suggested ARDRA as an appropriate method of distinguishing between different strains (Srutkova et al., 2011).

5.2.1.2 The 16S and 23S rRNA sequencing

Phenotypic identification of bacteria has been improved with the introduction of 16S/23S-5S rRNA sequencing methods (Collins et al., 1991). Some regions of 16S and 23S rRNA are

conserved throughout all bacteria, hence these sequences can be used as a source of species specific bacterial identification (Klijn et al., 1991). The primers encoding consensus sequences of 16S and 23S rRNA are used for PCR amplification of the genomic DNA of identifying probiotic bacteria and further the PCR products obtained can be analysed on agarose gel electrophoresis (Bested et al., 2013). The intergenic spacer region of 16S& 23S rRNA, generally used for the identification of *Lactobacillus* genus (Han et al., 2005). Rapid identification of *Lactobacilli* can be done by using colony PCR method, crude cell lysate and species specific primers targeting 16SrRNA (Luo et al., 2012).

5.3 Strain specific identification techniques

5.3.1 Ribotyping

Ribotyping is used for identification and classification of bacteria based on differences in the rRNA make-up which provides high discrimination of bacteria at the species and subspecies level. DNA extraction from bacterial colony, enzymatic digestion with appropriate restriction endonuclease followed by agarose gel electrophoresis analysis, transferred to hybridization membrane followed by incubation with a specific enzyme linked DNA probe. After washing, the enzyme substrate produces chromogenic signals, DNA fragments are visualized by adding on membrane made up of nylon, the pattern observed is then used for detection of strain. This technique has been used to differentiate *lactobacilli* (Rodtong et al., 1993). It has been also applied to characterize *Lactobacillus* and *Bifidobacterium* strains (Zhong et al., 1998; Giraffa et al., 2000). A current taxonomic study has been paying attention on typing of *Lactobacillus* spp. occurring in dental caries (Svec et al., 2010).

5.3.2 Random amplified polymorphic DNA (RAPD)

RAPD have been broadly reported as simple, quick, economical technique for genetic typing of diverse strains of bifidobacteria and lactic acid bacteria. A large number of samples can be analyzed within a short period of time (Satokari et al., 2003). It is a PCR based method that makes use of arbitrary primers or short primers which anneals to multiple random target sequences and directed to the formation of fingerprints of DNA bands differentiated between bacterial species (Fujimoto et al., 2008). This method has been successfully applied to distinguish between various species of lactobacilli at interspecies as well as intraspecific level. It allows to distinguish between bifidobacteria (Vincent et al., 1998) and *Lactobacillus acidophilous* group and related strains (Gancheva et al., 1999). Many reports on RAPD assay are available used to differentiate at strain and genus among different probiotic lactobacilli species (Du plesses et al., 1995; Roy et al., 2000) and bifidobacteria (Schillinger, 1999; Pal et al., 2012). It is also studied that RAPD method has a very low degree of discrimination among *L. fermentum* and *L. gasseri* in comparison with PFGE methods, (Gosiewski et al., 2012). However, RAPD has been used as a suitable technique to validate lactobacilli results received by Pulse field gel electrophoresis method (Herbal et al., 2013).

5.3.3 Pulse field gel electrophoresis (PFGE) and amplified fragment length polymorphism (AFLP)

PFGE allow the separation of large sized DNA fragments (larger than 10 kb). DNA fragments are subjected to periodic change in polarity of the electric field. The fingerprints of DNA obtained depend on the restriction enzyme specificity as well as the genome sequence

of the bacteria. It is reported that this method has been used for discrimination among significant probiotic bacterial strains such as *B. animalis* and *B. longum*, (Roy et al., 1996), *L. casei* and *L. rhamnosus* (Tynkkynen et al., 1999). PFGE has been used to categorize different lactobacilli strains in various probiotic products for human use (Coeuret et al., 2004). It has been also used in diversity and dynamics study of lactobacilli isolates (Henri et al., 2008). In a recent study PFGE has been used as an identification tool to evaluate vaginal epithelial cell colonization by a mixture of orally administrated different *Lactobacilli* (Strus et al., 2012). Furthermore, few studies on AFLP suggest that this technique is a hybrid of RFLP and RAPD techniques. It has better reproducibility, greater resolution and sensitivity at genome level as compared to RAPD-PCR. AFLP has become one of the most amplified techniques for bacterial taxonomic studies (Janssen et al., 1996). AFLP have been used as strain specific identification technique for probiotic strain Lactobacilli (Sisto et al., 2009) and *Bifidobacteria longum subsp.* (Makino et al., 2011).

5.3.4 PCR-Denaturing gradient gel electrophoresis (DGGE)

This method provides an easy evaluation of microorganisms used in probiotic products as it does not require prior culturing of cells or individual strains separation. The separation of similar length PCR products in denaturing gradient gels is based on sequence differences (Herbel et al., 2013). DGGE a molecular fingerprinting method allows separation of up to 500 bp DNA fragment. The melting point is an important factor in this method as the days DNA migrating through gel stops at a specific melting temperature which results in separation (Logan et al., 2003). Various studies reveal that the bacterial population

identification in the fermentation of Italian sausages (Cocolin et al., 2001), traditional sourdough (Randazzo et al., 2005) and wine making (Renouf et al., 2006) is done using this technique. It has been used also used for the identification of *Lactobacillus* species in the gastrointestinal microflora of mice (Walter et al., 2000). Study of intestinal persistence of orally administrated probiotic *Lactobacillus* sp. strain in healthy adults has been also done using PCR-Denaturing gradient gel electrophoresis (Hutt et al., 2011). To assure results this technique should be used in combination with 16S rRNA sequencing of the V3 region (Liu et al., 2012).

5.3.5 FACS (Fluorescent activated cell sorting) method

FACS is a high throughput tool allows sorting of heterogenous biological cell mixtures based on the fluorescent characteristics and light scattering. It allows large number cellular examination at a given time (i.e. 200- 2000 cells/second). There are numerous diverse parameters that can be linked to various cellular characteristics (Tracy et al., 2010). Different fluorescent probes can be used for the examination of physiological characterization of a live cell, these include membrane integrity and membrane potential, pH of cytoplasm, intracellular enzyme activity etc. These characteristics provide viability measure of bacteria (Chen et al., 2012). Hence FACS can be potentially used for high throughput cell sorting which could minimize the time required for determination of size of cell, abundance and cellular metabolic activity of probiotic strains (Davis, 2014).

5.3.6 Nucleic acid-based enumeration methods (PCR methods)

PCR is a method of obtaining genomic DNA fingerprints of bifidobacteria (Roy et al., 1996). It uses primers complementary to intersperse repetitive consensus sequences that can amplify sequences between the repetitive elements (Olive et al., 1999). Numerous studies have used three different primers which are BOX, ERIC and REP (Gomez et al., 2000). Primers ERIC and BOX have been used for classification and differentiation of Bifidobacteria (Ventura et al., 2002). RT-PCR uses reverse transcriptase enzyme to transcribe RNA into its complementary DNA. mRNA is short lived, extremely labile molecule in bacteria, hence transcription of bacterial mRNA should provide a useful marker of viability (Davis, 2014). Hellyer and Nadeau (2004) used RT-SDA (Reverse transcriptase-strand displacement amplification) as a marker for bacterial viability of bacteria. qPCR is culture independent, rapid amplification method that uses fluorescent dyes for the amplification and detection of DNA and RNA. Intensity of fluorescent signal produced by fluorescent dyes is proportional to the amount of DNA/RNA produced per cycle of PCR. This technique allows discrimination of different species and quantification of microbial populations by measuring the abundance of the target sequence in the DNA samples which are extracted from various food products (Postellac et al., 2011). It is evident from the literature that commonly used qPCR techniques use TaqMan labelled primers, molecular beacons and SYBR Green. TaqMan and molecular beacons are probe based techniques on reporter-quencher system (Bustin et al., 2012) whereas fluorescent dyes e.g. SYBR Green has a binding affinity to dsDNA and it is a quantitative approach for enumeration of bacterial communities (Miller et al., 2012; Castoldi et al., 2013). Interestingly, qPCR is also used to detect minor bacterial populations in the dominant population of bacteria and using this method even the non

cultivable bacterial species can be detected and quantified (Postellac et al., 2011). Recently a more understanding on these aspects was provided and a map has been published for obtaining more reliable results using MIQE guidelines (Taylor, 2013).

5.3.7 Fluorescent in situ hybridization (FISH)

Culture-independent detection methods have been used to enumerate probiotic strains based on viability. These techniques quantify viable cells either by using dyes to differentiate live and dead cells, by direct observation, measure the membrane integrity or characterize metabolic activity of cell such as nucleic acids synthesis. Probiotic bacteria can be directly visualized by microscopic techniques, but to enumerate viable bacteria requires differentiation of live and dead bacteria. FISH involves direct imaging and visual enumeration of bacteria. The technique enables rapid and accurate way to identify and quantify bacterial species by using epifluorescence, confocal microscopy, flow cytometry techniques (Zwirgmaier, 2005). This technique combines the versatility of microscopic and DNA/rRNA hybridization methods. The technique is based on hybridization of a rRNA-targeted fluorescent probe (a small stretch of oligonucleotide) to a specific region of 16S/23S rRNA ribosomal sequences of bacteria. Cells marked with fluorescent probe can be differentiated by using specific light filters (Lahtinen et al., 2006). It can be accomplished in a few hours, which allows fast in situ analysis (Juste et al., 2008). Now a days, an alternative to DNA probes is peptide nucleic acid (PNA) probes which have emerged as a rapid microbial detection method (Stender et al., 2002). PNA probes recognize and bind to specific region of 16S and 23S rRNA with higher thermal stability. The Lac663 probe bound

exclusively to *Lactobacillus* strains and capable of directly quantifying *Lactobacillus* spp. at concentrations which considered as effective on human health. The advantage of this technique is possible to localize and observe target cells within their native environment by the help of this technique. The technique also used in aquaculture systems for monitoring of pathogen and probiotics. Bacterial identification in intestinal tract of tilapia species have been studied using this technique. Also, it can also help to control the release of pathogenic bacteria to the environment results from pond water exchange, which could be a serious health problem for wildlife. The results of this study evidently verified that the FISH is a potent tool used in aquaculture (Alessandro et al., 2015). A new Histo-FISH method which is based on specific fluorochrome labeled probes, able to detect *Lactobacillus* spp. and *Bifidobacterium* spp. within biofilms on the mucosal surface of the GIT embedded in paraffin in histological slices. This is also appropriate method for visualization of bacterial populations in gastrointestinal tract (John et al., 2016). The Histo-FISH method permits us to analyse bacterial colonization as well as biofilm formation in stomach and caecum of BALB/c and germ-free mice (Madar et al., 2015). It has been also used in food industry for specific lactic acid bacteria detection for the production of hard cooked cheeses (Bottari et al., 2010). In addition it is also noteworthy to mention that FISH methods in combination with qPCR can also be used for enumeration of probiotic strains into a varied range of cheese based products.

Conclusion

This review describes the updated information towards use of various tools and techniques for identification and selection of probiotic microorganisms. The advanced tools and techniques used in this prominent area of research are also recapitulated in this review. Both conventional as well as advanced alternative techniques are reviewed which offers various options for selecting an appropriate microbial strain towards testing its potential for use of probiotic isolate in a variety of applications. These approaches are summarized in figure 3 which gives an overall idea of using such techniques in this field. Although the cost of advanced tools are high compared to traditional techniques, but nevertheless, these tools offer various advantages over traditional techniques. The high throughput screening tools such as FACS and Q-PCR offers fast and reliable identification of probiotics which in turns reduces the cost of experiments with high success rate. The cell based bioassay methods could predominantly use to check the activity as well as side effects of identified bacterial strains. The advanced techniques such as 16S and 23S RNA analysis, FACS and q-PCR can be potentially used for the identification as well as a selection of such microorganisms for industrial as well as academic applications.

Acknowledgements

The authors duly acknowledge the University Research Scholarship (URS) to Ruby Yadav (Ref. 10-RUR-8058) by M.D. University, Rohtak, India. The authors also acknowledge the support from SERB, Department of Science and Technology, Government of India (DST Fast Track Grant. No. SR/FT/LS-31/2012).

References

Backhed, F., Ding, H., Wang, T., Hooper, L. V., Koh, G. Y., Nagy, A., Semenkovich, C. F. and Gordon, J. I. (2004). The gut microbiota as an environmental factor that regulates fat storage. *Proc Natl Acad Sci USA* **101**: 15718–15723.

Banan-Mwine, E. D. and Lee, B. H. (2015). New perspectives on probiotics in health and disease. *Food sci. human well.* **15**:1–29.

Bested, A. C., Logan, A. C. and Selhub, E. M. (2013). Intestinal microbiota, probiotics and mental health: from Metchnikoff to modern advances: Part I - autointoxication revisited. *Gut. Pathog.* **5**:1–16.

Bustin, S. A., Zaccara, S. and Nolan, T. (2012). An introduction to the real-time polymerase chain reaction. In: Filion MN (ed) *Quantitative real-time PCR in applied microbiology*. Caister Academic Press, UK 3–25.

Bottari, B., Santarelli, M., Neviani, E., Gatti, M., 2010. Natural whey starter for Parmigiano Reggiano: culture-independent approach. *J. Appl. Microbiol.* **108**:1676–1684.

Caglar, E., Kargul, B. and Tanboga, I. (2005). Bacteriotherapy and probiotics role on oral health. *Oral Dis.* **11**:131–137.

Caglar, E., Topcuoglu, N., Ozbey, H., Sandalli, N. and Kulekci, G. (2015). Early colonization of *Lactobacillus reuteri* after exposure to Probiotics. *J. Clin. Pediatr. Dent.* **39**:326–330.

Casey, P., Casey, G. D., Gardiner, G. E., Tangney, M., Stanton, C., Ross, R. P., Hill, C. and Fitzgerald, G. F. (2004). Isolation and characterization of anti-Salmonella lactic acid bacteria from the porcine gastrointestinal tract. *Lett. Appl. Microbiol.* **39**:431–438.

Castoldi M et al (2013) Expression profiling of MicroRNAs by quantitative real-time PCR. In: Nolan T, Bustin SA (ed) PCR technology: current innovations. CRC Press, Boca Raton, Taylor & Francis Group, pp 601–608

Chen, S., Cao, Y., Ferguson, L. R., Shu, Q. and Garg, S. (2012). Flow cytometric assessment of the protectants for enhanced in vitro survival of probiotic lactic acid bacteria through simulated human gastro-intestinal stresses. *Appl. Microbiol. Biotechnol.* **95**:345–356.

Chong, E. S. L. (2014). A potential role of probiotics in colorectal cancer prevention: review of possible mechanisms of action. *World J. Microbiol. Biotechnol.* **30**: 351–374.

Claassen, E. (2014). Cost-benefit relation of diet and probiotics in iatrogenic bowel irregularity (IBI). *Front. Pharmacol. | Pharmaceutical Medicine and Outcomes Research* doi: 10.3389/fphar.2014.00014.

Cocolin, L., Manzano, M., Cantoni, C. and Comi, C. (2001). Denaturing gradient gel electrophoresis analysis of the 16S rRNA gene V1 region to monitor dynamic changes in the bacterial population during fermentation of Italian sausages. *Appl. Environ. Microbiol.* **67**:5113–5121.

Coeuret, V., Gueguen, M. and Vernoux, J. P. (2004). Numbers and strains of lactobacilli in some probiotic products. *Int. J. Food Microbiol.* **97**:147–156.

Collado, C. M. and Hernandez, M. (2007). Identification and differentiation of *Lactobacillus*, *Streptococcus* and *Bifidobacterium* species in fermented milk products with bifidobacteria. *Microbiol. Res.* **16**:86–92.

Collado, M., Moreno, Y., Cobo, J. and Hernández, M. (2006). Microbiological evaluation and molecular characterization of bifidobacteria strains in commercial fermented milks. *Eur. Food Res. Tech.* **222**:112–117.

Collins, M. D., Rodrigues, U., Ash, C., Aguirre, M., Farrow, J. A. E. and Martinez-Murcia, A. (1991). Phylogenetic analysis of the genus *Lactobacillus* and related lactic acid bacteria as determined by reverse transcriptase sequencing of 16S rRNA. *FEMS. Microbiol. Lett.* **77**:5–12.

Costa, D. J., Marteau, P., Amouyal, M., Poulsen, L. K., Hamelmann, E., Cazaubiel, M., Housez, B., Leuillet, S., Stavnsbjerg, M., Molimard, P., Courau, S. and Bousquet, J. (2014). Efficacy and safety of the probiotic *Lactobacillus paracasei* LP-33 in allergic rhinitis: A double-blind, randomized, placebo-controlled trial (GA2LEN Study). *Eur. J. Clin. Nutr.* **68**:602–607.

Davis, C. (2014). Enumeration of probiotic strains: Review of culture-dependent and alternative techniques to quantify viable bacteria. *J. Microbiol. Methods* **103**:9–17.

Dherbecourt, J., Thierry, A., Madec, M. N. and Lortal, S. (2006). Comparison of amplified ribosomal DNA restriction analysis, peptidoglycan hydrolase and biochemical profiles for rapid dairy *Propionibacteria* species identification. *Res. Microbiol.* **157**:905–913.

Donelli, G., Vuotto, C. and Mastromarino, P. (2013). Phenotyping and genotyping are both essential to identify and classify a probiotic microorganism. *Microb. Ecol. Health Dis.* **24**:1–8.

Duca, A. D., Cesar E. D., Abreu, P. C. (2015). Bacterial community of pond's water, sediment and in the guts of tilapia (*Oreochromis niloticus*) juveniles characterized by fluorescent in situ hybridization technique. *Aqua. Res.* **46**:707–715.

Franz, C., Specht, I., Haberer, P. and Holzapfel, W. H. (2001). Bile salt hydrolase activity of enterococci isolated from food: screening and quantitative determination. *J. Food Prot.* **64**:725–729.

Fujimoto, J., Matsuki, T., Sasamoto, M., Tomii, Y. and Watanabe, K. (2008). Identification and quantification of *Lactobacillus casei* strain Shirota in human feces with strain-specific primers derived from randomly amplified polymorphic DNA. *Int. J. Food Microbiol.* **126**:210–215.

Fuller, R. (1989). Probiotics in man and animals. *J. Appl. Bacteriol.* **66**:365–378.

Galdeano, C. M., Ivanna, N. N., Esteban, C., Alejandra, M. B. and Gabriela, P. (2015). Role of Probiotics and Functional Foods in Health: Gut Immune Stimulation by Two Probiotic Strains and a Potential Probiotic Yoghurt. *Endocr. Metab. Immune Disord. Drug Targets.* **15**:37–45.

Gancheva, A., Pot, B., Vanhonacker, K., Hoste, B. and Kersters, K. (1999). A polyphasic approach towards the identification of strains belonging to *Lactobacillus acidophilus* and related species. *Syst. Appl. Microbiol.* **22**:573–585.

Giraffa, G., Gatti, M., Rossetti, L., Senini, L. and Neviani, E. (2000). Molecular diversity within *Lactobacillus helveticus* as revealed by genotypic characterization. *Appl. Environ. Microbiol.* **66**:1259–1265.

Gomez, Z. A., Urraza, D. P., Antoni DG (2000) Characterization of Bifidobacterium strains using box primers. *Anaerobe.* **6**:169–177.

Gosiewski, T., Chmielarczyk, A., Strus, M., Brzychczy, W. M. and Heczko, P. B. (2012). The application of genetics methods to differentiation of three *Lactobacillus* species of human origin. *Ann. Microbiol.* **62**:1437–1445.

Gotteland, M., Andrews, M., Toledo, M., Muñoz, L., Caceres, P., Anziani, A., Wittig, E., Speisky, H. and Salazar, G. (2008). Modulation of *Helicobacter pylori* colonization with cranberry juice and *Lactobacillus johnsonii* La1 in children. *Nutr.* **24**:421–426.

Gungor, O. E., Kirzioglu, Z. and Kivanc, M. (2015). Probiotics: can they be used to improve oral health? *Benef. Microbes.* 1–10.

Goyal, S., Raj, T., Banerjee, C., Imam, J. and Shukla, P. (2013). Isolation and characterization of probiotic microorganisms from curd and chili sauce. *Int J probiotics prebiotics.* **8**:91–94.

Hajela, N., Ramakrishna, B. S., Nair, B. G., Abraham, P., Gopalan, S. and Ganguly, N. K. (2015). Gut microbiome, gut function, and probiotics: Implications for health. *Ind. J. Gastroenterol.* **34**:93–107.

Han, K. S., Kim, Y., Choi, S., Oh, S., Park, S., Kim S. H. and Whang, K. Y. (2005). Rapid identification of *Lactobacillus acidophilus* by restriction analysis of the 16S-23S rRNA intergenic spacer region and flanking 23S rRNA gene. *Biotechnol. Lett.* **27**:1183–1188.

Hellyer, T. J. and Nadeau, J. G. (2004). Strand displacement amplification: a versatile tool for molecular diagnostics. *Expert Rev. Mol. Diagn.* **4**:251–261.

Henao, M. J., Elinav, E., Thaiss, C. A., Licona, L. P. and Flavell, R. A. (2013). Role of the intestinal microbiome in liver disease. *J. Autoimmu.* **46**:66–73.

Henri, D. S., Desmasures, N. and Gueguen, M. (2008). Diversity and dynamics of lactobacilli populations during ripening of RDO Camembert cheese. *Can. J. Microbiol.* **54**:218–228.

Herbel, S. R., Vanjen, W., Wieler, L. H. and Guenther, S. (2013). Timely approaches to identify probiotic species of the genus *Lactobacillus*. *Gut. Pathogens.* **5**:1–13.

Hill, C., Guarner, F., Reid, G., Gibson, G. R., Merenstein, D. J., Pot, B., Morelli, L., Canani, R. B., Flint, H. J., Salminen, S., Calder, P. C. and Sanders, M. E. (2014). Expert consensus document. The International Scientific Association for Probiotics and Prebiotics consensus statement on the scope and appropriate use of the term probiotic. *Nature Reviews. Gastroenterol. Hepatol.* **11**:506–514.

- Hoque, M. Z., Akter, F., Hossain, K. M., Rahman, M. S. M., Billah, M. M. and Islam, K. M. D. (2010). Isolation, Identification and Analysis of Probiotic Properties of *Lactobacillus Spp.* From Selective Regional Yoghurts. *World J. Dairy & Food Sci.* **5**:39–46.
- Hutt, P., Koll, P., Stsepetova, J., Alvarez, B., Mandar, R., Krogh, A. K., Marcotte, H., Hammarstrom, L. and Mikelsaar, M. (2011). Safety and persistence of orally administered human *Lactobacillus sp.* strains in healthy adults. *Benef. Microbes.* **2**:79–90.
- Ivanov, I. I., Honda, K. (2012). Intestinal commensal microbes as immune modulators. *Cell Host Microbe.* **12**:496–508.
- Jain, M., Gupta, K. and Jain, P. (2014). Significance of Probiotics and Prebiotics in Health and Nutrition. *Malaya J. Biosci.* **1**:181–195.
- Janssen, P., Coopman, R., Huys, G., Swings, J., Bleeker, M., Vos, P., Zabeau, M. and Kersters, K. (1996). Evaluation of the DNA fingerprinting method AFLP as a new tool in bacterial taxonomy. *Microbiol.* **142**:1881–1893.
- John, B. G., Barbara, A. W., Michael, J. G., Deirdre, M. (2016). Visualization of microbe dietary remnant interactions in digesta from pigs, by fluorescence *in situ* hybridization and staining methods; effects of a dietary arabinoxylan-rich wheat fraction. *Food Hydrocoll.* **52**:952–962.
- Jothika, M., Vanajassun, P. P. and Someshwar, B. (2015). Effectiveness of probiotic, chlorhexidine and fluoride mouthwash against *Streptococcus mutans* – Randomized, single-blind, *in vivo* study. *J. Int. Soc. Prev. Community. Dent.* **5**:44–48.

Juste, A., Thommad, B., Lievens, B., (2008). Recent advances in molecular techniques to study microbial communities in food-associated matrices and processes. *Food Microbiol.* **25**:745–761.

Klijn, N., Weerkamp, A. H. and De Vos, W. M. (1991). Identification of mesophilic lactic acid bacteria by using polymerase chain reaction-amplified variable regions of 16S rRNA and specific DNA probes. *Appl. Environ. Microbiol.* **57**:3390–3393.

Kumar, M., Ghosh, M. and Ganguli, A. (2012). Mitogenic response and probiotic characteristics of lactic acid bacteria isolated from indigenously pickled vegetables and fermented beverages. *World J. Microbiol. Biotechnol.* **28**:703–711.

Kwon, H. K., Kim, G. C., Kim, Y., Hwang, W., Jash, A., Sahoo, A., Kim, J. E., Nam, J. H. and Im, S. H. (2013). Amelioration of experimental autoimmune encephalomyelitis by probiotic mixture is mediated by a shift in T helper cell immune response. *Clin. Immunol.* **146**:217–227.

Lahtinen, S. J., Gueumonde, M., Ouwehand, A. C., Reinikainen, J. P., Salminen, S. J. (2006). Comparison of four methods to enumerate probiotic bifidobacteria in a fermented food product. *Food Microbiol.* **23**:571–577.

Lalitha, T. A. (2011). Probiotics and oral health. *J. Oral Res. Rev.* **3**:20–26.

Lara, M. J. S., Llorente, C. G., Diaz, J. P., Gil, A. (2015). The Role of Probiotic Lactic Acid Bacteria and Bifidobacteria in the Prevention and Treatment of Inflammatory Bowel Disease

and Other Related Diseases: A Systematic Review of Randomized Human Clinical Trials. *BioMed. Res. Int.* 1–15.

Lee, N. K., Kim, S. Y., Han, K. J., Eom, S.J. and Paik, H. D. (2014). Probiotic potential of Lactobacillus strains with anti-allergic effects from kimchi for yogurt starters. *LWT - Food Sci. Tech.* **58**:130–134.

Lee, I. (2014). Critical pathogenic steps to high risk Helicobacter pylori gastritis and gastric carcinogenesis. *World J. gastroenterol.* **20**:6412–6419.

Levy, C. and Lindor, K. D. (2006). Primary sclerosing cholangitis: Epidemiology, natural history, and prognosis. *Semin. Liver Dis.* **26**:22–30.

Lilly, D. M. and Stillwell, R. H. (1965). Probiotics: Growth-Promoting Factors Produced by Microorganisms. *Sci.* **147**:747–748.

Liu, W., Baq, Q., Jirimutu, Qing, M., Siriguleng, Chen, X., Sun, T., Li, M., Zhang, J., Yu, J., Bilige, M., Sun, T. and Zhang, H. (2012). Isolation and identification of lactic acid bacteria from Tarag in Eastern Inner Mongolia of China by 16S rRNA sequences and DGGE analysis. *Microbiol. Res.* **167**:110–115.

Logan, A. C., Rao, A. V. and Irani, D. (2003). Chronic fatigue syndrome: lactic acid bacteria may be of therapeutic value. *Med. Hypotheses.* **60**:915–923.

Luo, Y., Ma, B. C., Zou, L. K., Cheng, J. G., Cai, Y. H., Kang, J. P., Li, B., Gao, X. H., Wang, P. and Xiao, J. J. (2012). Identification and characterization of lactic acid bacteria from forest musk deer feces. *Afr. J. Microbiol. Res.* **6**:5871–5881.

Madar, M., Slizova, M., Czerwinski, J., Hrcokova, G., Mudronova, D., Gancarcikova, S., Popper, M., Pistl, J., Soltys, J., Nemcova, R. (2015). Histo-FISH protocol to detect bacterial compositions and biofilms formation *in vivo*. *Benef. Microbes*. DOI: <http://dx.doi.org/10.3920/BM2015.0016>. 1876–2883.

Makino, H., Kushiro, A., Ishikawa, E., Muylaert, D., Kubota, H. and Sakai, T. (2011). Transmission of intestinal *Bifidobacterium longum subsp. longum* strains from mother to infant, determined by multilocus sequencing typing and amplified fragment length polymorphism. *Appl. Environ. Microbiol.* **77**: 678–893.

Maldonado, G. C., Lemme-Dumit, J. M., Thieblemont, N., Carmuega, E., Weill, R. and Perdigon, G. (2015). Stimulation of Innate Immune Cells Induced by Probiotics: Participation of TollLike Receptors. *J. Clin. Cell Immunol.* **6**:1–9.

Margolles, A., Mayo, B. and Ruas, M. P. (2009). Screening, identification, and characterization of Lactobacillus and Bifidobacterium strains. **In:** Handbook of probiotics and prebiotics, pp. 45–58. Nomoto, K., Salminen, S., Lee, Y. K., Eds., Hoboken, Wiley, New Jersey.

Martin, R., Miquel, S., Ulmer, J., Langella, P. and Bermudez-Humaran, L. G. (2014). Gut ecosystem: how microbes help us. *Benef. Microbes*. **5**:219–233.

McFarland, L. V. (2014). Use of probiotics to correct dysbiosis of normal microbiota following disease or disruptive events: A systematic review. *BMJ Open*. **4**:1–18.

Metchnikoff, E. (1907). The Prolongation of Life. Optimistic Studies London. 16–183.

- Mokarram, R. R., Mortazavi, S. A., Najafi, M. B. H. and Shahidi, F. (2009). The influence of multi stage alginate coating on survivability of potential probiotic bacteria in simulated gastric and intestinal juice. *Food Res. Int.* **42**:1040–1045.
- Mortazavian, A. M., Ehsani, M. R., Azizi, A., Razavi, S. H., Mousavi, S. M., Sohrabvandi, S. and Reinheimer, J. A. (2008). Viability of calcium-alginate-microencapsulated probiotic bacteria in Iranian yogurt drink (Doogh) during refrigerated storage and under simulated gastrointestinal conditions. *Aust. J. Dairy Tech.* **63**:24–29.
- Miller, D. M., Dudley, E. G. and Roberts, R. F. (2012). Technical note: development of a quantitative PCR method for monitoring strain dynamics during yogurt manufacture. *J. Dairy Sci.* **95**:4868–4872.
- Narbona, L. E., Uberos, F. J., Armada, M. M. I., Couce, P. M. L., Rodriguez, M. G. and Saenz, de P. M. (2014). Nutrition and Metabolism Group of the Spanish Neonatology Society: Recommendations and evidence for dietary supplementation with probiotics in very low birth weight infants. *An. Pediatr.* **81**:1–8.
- Neish, A. S. (2009). Microbes in gastrointestinal health and disease. *Gastroenterol.* **136**:65–80.
- Norberto, C., Tapia, C., Rodriguez, L. G., Jeong, M. S., Ramirez, Y. L., Becerra, V. B., Eva, J. H., Juan, L., Flores, R., Arrese, M., Sanchez, N. M. and Uribe, M. (2015). Current evidence on the use of probiotics in liver diseases. *J. Func. Food.* **17**:137–151.
- Olive, D. M. and Bean, P. (1999). Principles and applications of methods for DNA-based typing of microbial organisms. *J. Clin. Microbiol.* **37**:1661–1669.

Pal, K., Szen, O., Kiss, A. and Naar, Z. (2012). Comparison and evaluation of molecular methods used for identification and discrimination of lactic acid bacteria. *J. Sci. Food Agric.* **92**:1931–1936.

Panigrahi, P. (2014). Probiotics and prebiotics in neonatal necrotizing enterocolitis: New opportunities for translational research. *Pathophys.* **21**:35–46.

Parker, R. (1974). Probiotics, the other half of the antibiotic story. *Anim. Nutr. Health.* **28**:240–255.

Patel, S. (2015). Emerging trends in nutraceutical applications of whey protein and its derivatives. *J. Food Sci. Technol.* 1–12.

Patel, S. and Goyal, A. (2013). Evolving roles of probiotics in cancerprophylaxis and therapy. *Antimicrob. Proteins.* **5**:59–66.

Patel, S., Shukla, R. and Goyal, A. (2015). Probiotics in valorization of innate immunity across various animal models. *J. Func. Foods.* **14**:549–561.

Paturi, G., Butts, C. A., Hewitt, K. L. B., Hedderley, D., Stoklosinski, H. and Ansell, J. (2015). Differential effects of probiotics, prebiotics, and synbiotics on gut microbiota and gene expression in rats. *J. Func. Foods.* **13**:204–213.

Pierre, W. and Denis, R. L. (2005). Review of molecular methods for identification, characterization and detection of Bifidobacteria. *INRA, EDP Sci.* **85**:23–32.

Postellec, F., Falentin, H., Pavan, S., Combrisson, J. and Sohier, D. (2011). Recent advances in quantitative PCR (qPCR) applications in food microbiology. *Food Microbiol.* **28**:848–861.

Prasanna, P. H. P., Grandison, A. S. and Charalampopoulos, D. (2014). Bifidobacteria in milk products: An overview of physiological and biochemical properties, exopolysaccharide production, selection criteria of milk products and health benefits. *Food Res. Int.* **55**:247–262.

Pundir, R. K., Rana, S., Kashyap, N. and Kaur, A. (2013). Probiotic potential of lactic acid bacteria isolated from food samples: an in vitro study. *J. App. Pharm. Sci.* **3**:85-93.

Raghuwanshi, S., Misra, S., Sharma, R. and Bisen, P. S. (2015). Indian perspective for probiotics: A review. *Ind. J. Dairy Sci.* **68**:195–205.

Randazzo, C. L., Heilig, H., Restuccia, C., Giudici, P. and Caggia, C. (2005). Bacterial population in traditional sourdough evaluated by molecular methods. *J. Appl. Microbiol.* **99**:251–258.

Rebolledo, M., Rojas, E. and Salgado, F. (2013). Effect of two probiotics containing *Lactobacillus rhamnosus* and *Lactobacillus johnsonii* variety on the in vitro growth of *Streptococcus mutans*. *Int. J. odontostomatol.* **7**:415–419.

Reid, G. (2015). The growth potential for dairy probiotics. *Int. Dairy J.* **49**:16–22.

Renouf, V., Miot, S. C. and Lonvaud, F. A. (2006). Lactic acid bacteria evolution during winemaking: Use of *rpoB* gene as a target for PCR-DGGE analysis. *Food Microbiol.* **23**:136–145.

Riaz, Q. U. and Masud, T. (2013). Recent trends and applications of encapsulating materials for probiotic stability. *Crit. Rev. Food Sci. Nutr.* **53**:231–244.

Rodtong, S. and Tannock, G. W. (1993). Differentiation of *Lactobacillus* strains by ribotyping. *Appl. Environ. Microbiol.* **59**:3480–3484.

Roy, D., Ward, P. and Champagne, G. (1996). Differentiation of bifidobacteria by use of pulsed-field gel electrophoresis and polymerase chain reaction. *Int. J. Food Microbiol.* **29**:11–29.

Sadana, G., Sikri, C. and Sadana, S. (2015). Role of probiotics in general and oral health: a review. *Ind. J. Compr. Dent. Care.* **5**:571–57.

Satokari, R. M. et al (2003). Molecular approaches for the detection and identification of bifidobacteria and lactobacilli in the human gastrointestinal tract. *Syst. Appl. Microbiol.* **26**:572–584.

Savadago, A., Ouattara, C. A. T., Bassole, I. H. N. and Traore, S. A. (2006). Bacteriocins and lactic acid bacteria—a mini review. *Afr. J. biotec.* **5**:678–683.

Saxer, S., Schwenninger, S. M. and Lacroix, C. (2013). Characterization of the microflora of industrial Mexican cheeses produced without added chemical preservatives. *LWT - Food Sci. Technol.* **10**:0023–6438.

Schillinger, U. (1999). Isolation and identification of lactobacilli from novel-type probiotic and mild yoghurts and their stability during refrigerated storage. *Int. J. Food Microbiol.* **47**:79–87.

Schulz, C., Koch, N., Schutte, K., Pieper, D. H. and Malfertheiner, P. (2015).

H. pylori and its modulation of gastrointestinal microbiota. *J. Dig. Dis.* **16**:109–117.

Sharma, P., Tomar, S. K., Goswami, P., Sangwan, V. and Singh, R. (2014). Antibiotic resistance among commercially available probiotics. *Food Res. Int.* **57**:176–195.

Shehata, I. (2012). Molecular identification of probiotics *Lactobacillus* strain isolates by: Amplified Ribosomal DNA Restriction Analysis (ARDRA). *Sci. J. Microbiol.* 1–8.

Siepert, B., Reinhardt, N., Kreuzer, S., Bondzio, A., Twardziok, S. and Brockmann, G. (2014). *Enterococcus faecium* NCIMB 10415 supplementation affects intestinal immune-associated gene expression in post-weaning piglets. *Vet. Immunol. Immunopathol.* **157**: 65–77.

Singh, A. K., Pandey, S. K., Saha, G. and Gattupalli, N. K. (2015). Pyrroloquinoline quinone (PQQ) producing *Escherichia coli* Nissle 1917 (EcN) alleviates age associated oxidative stress and hyperlipidemia, and improves mitochondrial function in ageing rats. *Exp. Gerontol.* **66**:1–9.

Singh, K., Kallali, B., Kumar, A. and Thaker, V. (2011). Probiotics: A review. *Asian. Pac. J. Trop. Biomed.* **1**:287–290.

Sisto, A., De Bellis, P., Visconti, A., Morelli L. and Lavermicocca, P. (2009). Development of a PCR assay for the strain-specific identification of probiotic strain *Lactobacillus paracasei*. *Int. J. Food Microbiol.* **136**:59–65.

Soukoulis, C., Yonekura, L., Gan, H. H., Behboudi, J. S., Parmenter, C. and Fisk, I. (2014). Probiotic edible films as a new strategy for developing functional bakery products: The case of pan bread. *Food Hydrocoll.* **39**:231–242.

Srutkova, D., Spanova, A., Spano, M., Drab, V., Schwarzer, M. and Kozakova, H. (2011). Efficiency of PCR-based methods in discriminating *Bifidobacterium longum* ssp. *longum* and *Bifidobacterium longum* ssp. *infantis* strains of human origin. *J. Microbiol. Methods.* **87**:10–16.

Stefan, K. and Rossitza, V. D. (2014). Bacterial vaginosis - local *Lactobacillus casei* var *rhannosus* doderlein monotherapy. *Int. J. Probiotics Prebiotics*. **9**:129–133.

Stender, H., Fiandaca, M., Hyldig-Nielsen, J., Coull, J., (2002). PNA for rapid microbiology. *J. Microbiol.Methods*. **48**:1–17.

Strus, M., Chmielarczyk, A., Kochan, P., Adamski, P., Chelmicki, Z. and Chelmicki, A. (2012). Studies on the effects of probiotic *Lactobacillus* mixture given orally on vaginal and rectal colonization and on parameters of vaginal health in women with intermediate vaginal flora. *Eur. J. Obstet. Gynecol. Reprod. Biol.* **163**:210–215.

Svec, P., Kukletova, M. and Sedlacek, I. (2010). Comparative evaluation of automated ribotyping and RAPD-PCR for typing of *Lactobacillus* spp. occurring in dental caries. *A Van Leeuw*. **98**:85–92.

Taylor, S. (2013). A map to more reliable RT-qPCR results using MIQE guidelines. *Am. Lab.* **18**:20–21.

Tian, Q., Wang, T. T., Tang, X., Han, M. Z., Leng, X. J. and Mao, X. Y. (2015.) Developing a potential prebiotic of yogurt: growth of *Bifidobacterium* and yogurt cultures with addition of glycomacropeptide hydrolysate. *Int. J. Food Sci. Tech.* **50**:120–127.

Toucheffeu, Y., Montassier, E., Nieman, K., Gastinne, T., Potel, G., Brule, V. S., Le, V. F. and de La, C. M. F. (2014). Systematic review: The role of the gut microbiota in chemotherapy or radiation induced gastrointestinal mucositis - Current evidence and potential clinical applications. *Aliment. Pharmacol. Ther.* **40**:409–421.

Tracy, B. P., Gaida, S. M. and Papoutsakis, E. T. (2010). Flow cytometry for bacteria: enabling metabolic engineering, synthetic biology and the elucidation of complex phenotypes. *Curr. Opin. Biotechnol.* **21**:85–99.

Tripathi, M. K. and Giri, S. K. (2014). Probiotic functional foods: Survival of probiotics during processing and storage. *J. Func. Food.* **9**:225–241.

Tynkkynen, S., Satokari, R., Saarela, M., Mattila-Sandholm, T. and Saxelin, M. (1999). Comparison of ribotyping, randomly amplified polymorphic DNA Analysis, and pulsed-field gel electrophoresis in typing of *Lactobacillus rhamnosus* and *L. casei* strains. *Appl. Environ. Microbiol.* **65**:3908–3914.

Vaneechoutte, M., Rossau, R., De Vos, P., Gillis, M., Janssens, D. and Paepe, N. (1992). Rapid identification of bacteria of the Comamonadaceae with amplified ribosomal DNA-restriction analysis (ARDRA). *FEMS. Microbiol. Lett.* **72**:227–233.

Venema, K. and Maathuis, A. J. H. (2003). A PCR-based method for identification of bifidobacteria from the human alimentary tract at the species level. *FEMS. Microbiol. Lett.* **224**:143–149.

Ventura, M. and Zink, R. (2002). Rapid identification, differentiation, and proposed new taxonomic classification of *Bifidobacterium lactis*. *Appl. Environ. Microbiol.* **68**:6429–6434.

Vieira, A. T., Teixeira, M. M. and Martins, F. S. (2013). The role of probiotics and prebiotics in inducing gut immunity. *Front. Immunol.* **4**(445): 1–12.

Vincent, D., Roy, D., Mondou, F. and Dery, C. (1998). Characterization of bifidobacteria by random DNA amplification. *Int. J. Food Microbiol.* **43**:185–193.

Walter, J., Tannock, G. W., Tilsala-Timisjarvi, A., Rodtong, A., Loach, D. M. and Munro. (2000). Detection and identification of gastrointestinal *Lactobacillus* species by using denaturing gradient gel electrophoresis and species-specific PCR primers. *Appl. Environ. Microbiol.* **66**:297–303.

West, N. P., Horn, P. L., Pyne, D. B., Gebiski, V. J., Lahtinen, S. J., Fricker, P. A. and Cripps, A.W. (2014). Probiotic supplementation for respiratory and gastrointestinal illness symptoms in healthy physically active individuals. *Clin. Nutr.* **33**:581–587.

Zhang, H., Sun, J., Liu, X., Hong, C., Zhu, Y., Liu, A., et al. (2013). *Lactobacillus paracasei* subsp. *paracasei* LC01 positively modulates intestinal microflora in healthy young adults. *J. Microbiol.* **51**:777–782.

Zhong, W., Millsap, K., Bialkowska-Hobrzanska, H. and Reid, G. (1998). Differentiation of *Lactobacillus* species by molecular typing. *Appl. Environ. Microbiol.* **64**:2418–2423.

Zwirgmaier, K. (2005). Fluorescent in situ hybridization (FISH) – The next generation. *FEMS. Microbiol. Lett.* **246**:151–158.

Table 1 Commercially available probiotic products, their strain designations and commercial applications

Probiotic strain	Product (company)	Applications and proven Health benefits	References
<i>L. acidophilus</i> NCFMs	HOWARU® Premium Probiotics (Danisco A/S)	Beverages, confectionery, dairy, dietary supplements and frozen desserts, anti allergic	West et al. 2014; Patel et al. 2015
<i>L. acidophilus</i> La-14	Flora FIT® (Danisco A/S)	Food and beverages, skin health	Caglar et al. 2015; Sadana et al. 2015
<i>L. acidophilus</i> LA5	Biorich® (Chr. Hansen A/S)	Starters for yoghurt manufacture, anti- diarrhoea, immune boosting, anti-infection	Reid 2015; Sadana et al. 2015
<i>L. casei</i> Shirota	Yakult® (Yakult Honsha Co. Ltd)	Probiotic beverage, intestinal flora reposition, improve digestion	Raghuwanshi et al. 2015; Hajela et al. 2015
<i>L. casei</i> DN114001	Actimel® (Danone)	Protection against pathogens	Martin et al. 2014; Galdeano et al. 2015

<i>Lactobacillus casei</i> <i>rhamnosus</i> Lcr 35	Ginophilus® (Probionov)	Lowers vaginal pH, prevents harmful pathogenic bacteria from colonizing and proliferating	Stefan et al. 2014; Paturi et al. 2015
<i>Lactobacillus</i> <i>johnsonii</i> La1	SVELTY® Gastro Protect (Nestle)	Controls <i>H. pylori</i> infection and stomach discomfort, immune boosting	Gotteland et al. 2008; Rebolledo et al. 2014; Schulz et al. 2015
<i>Lactobacillus</i> <i>plantarum</i> 299v	LP299V® (Probi AB)	Anti-diarrhoea, immune boosting	Jain et al. 2014; Patel et al. 2015
<i>Lactobacillus</i> <i>rhamnosus</i> GG	Actifit®, Gefilus®, LGG®, Vifit® (Valio)	Antiallergy, anti-diarrhoea, immune boosting, oral health	Caglar et al. 2005; Rebolledo et al. 2014; Schulz et al. 2015
<i>L. paracasei</i> CRL431	L. casei 431® (Chr. Hansen A/S)	Immune boosting, anti- infection	Galdeano et al. 2015; Paturi et al. 2015
<i>L. paracasei</i> GMNL-33	GenMont biotech	Anti allergic	Patel et al. 2015 Schulz et al. 2015
<i>L. reuteri</i> ATCC 55730 (Protectis)	BioGaia (Biogaia)	Antiallergy, anti-diarrhoea, oral health, immune	Paturi et al. 2015

		boosting, anti-infection	
<i>B. animalis</i> ssp. lactis Bb12	Chr. Hansen	Reduced salivary counts of <i>S. mutans</i>	Caglar et al. 2005; Jothika et al. 2015
<i>B. animalis</i> DN173010	DanActive fermented milk (Danone)	Digestive health improvement, Inhibits growth of <i>S. mutans</i>	Caglar et al. 2005; Gungor et al. 2015
<i>B. lactis</i> HN019 (DR10)	Fonetera	Immune boosting, anti-infection	Gungor et al. 2015; Paturi et al. 2015
<i>B. longum</i> BB536	Morinaga Milk Industry Co. Ltd.	Anti allergy	Caglar et al. 2015; Schulz et al. 2015
<i>B. infantis</i> 35624	Align (Procter y Gamble)	Maintain the digestive balance fortifying the digestive system with healthy bacteria	Raghuwanshi et al. 2015; Tian et al. 2015
<i>Enterococcus faecium</i> SF68	Bioflorin (Cerbios – Pharma)	Prevention and treatment of intestinal disorders, anti-diarrhoea	Hajela et al. 2015; Siepert et al. 2014
<i>Streptococcus salivarius</i> K12	BLIS technologies	immune boosting, anti-infection, oral health	Gungor et al. 2015; Patel et al. 2015; Paturi et al.

			2015
<i>Streptococcus oralis</i> KJ3 <i>Streptococcus uberis</i> KJ2 <i>Streptococcus rattus</i> JH145	ProBiora3 (Oragenics Inc.)	Low-acid-producing oral inhabitants quickly colonize the oral cavity to inhibit the growth of the pathogenic streptococci	Caglar et al. 2005; Gungor et al. 2015; Jothika et al. 2015
<i>Escherichia coli</i> Nissle 1917	Mutaflor (Ardeypharm)	Prevention and treatment of intestinal disorders	Siepert et al. 2014; Singh et al. 2015
<i>Escherichia coli</i>	URO VAXOM® (Apsen)	Immunotherapy, prevention recurrent infections of the lower urinary tract	Stefan et al. 2014; Paturi et al. 2015; Singh et al. 2015
<i>Saccaromyces</i> <i>boulardi</i>	Florastor (Biocodex)	Immune boosting	Patel et al. 2015; Paturi et al. 2015

Table 2 Genotypic methods and their applications

Method	Strain/species specific typing	Applications	References
ARDR A	Species specific	Identification and discrimination between closely related bacterial species	Srutkova et al. 2011; Shehata et al. 2012; Gianfranco et al. 2013
16S/23SrRN A sequencing	Species specific	Identification based upon 16S/23S conserved region	Luo et al. 2012; Bested et al. 2013
Ribotyping	Strain specific	Identification and classification based upon differences in rRNA	Svec et al. 2010
RAPD	Strain specific	Identification and distinguish between different bacterial species based upon arbitrary primers	Pal et al. 2012; Gosiewski et al. 2012
PFGE	Strain specific	Identification and	Coeuret et al.

		differentiation between different bacterial strains	2004; Henri et al. 2008; Strus et al. 2012
PCR- DGGE	Strain specific	Strain level identification	Hutt et al. 2011; Liu et al. 2012; Herbel et al. 2013
AFLP	Strain specific	Taxonomic studies of bacteria	Sisto et al. 2009; Makino et al. 2011




Fig. 1 Criteria for selection of probiotics




Fig. 2 Health benefits of probiotic bacteria




Fig. 3 A combinatory approach for selection of probiotics