



Fermented and malted millet products in Africa: Expedition from traditional/ethnic foods to industrial value added products

J.A. Adebiyi, A.O. Obadina, O.A. Adebo & E. Kayitesi

To cite this article: J.A. Adebiyi, A.O. Obadina, O.A. Adebo & E. Kayitesi (2016): Fermented and malted millet products in Africa: Expedition from traditional/ethnic foods to industrial value added products, Critical Reviews in Food Science and Nutrition, DOI: [10.1080/10408398.2016.1188056](https://doi.org/10.1080/10408398.2016.1188056)

To link to this article: <http://dx.doi.org/10.1080/10408398.2016.1188056>



Accepted author version posted online: 31 May 2016.
Published online: 31 May 2016.



[Submit your article to this journal](#)



Article views: 10



[View related articles](#)



[View Crossmark data](#)

Fermented and malted millet products in Africa: Expedition from traditional/ethnic foods to industrial value added products

Adebiyi J.A.^{1,*}, Obadina A.O.^{1,2}, Adebo, O.A.¹, Kayitesi E.¹

¹Department of Biotechnology and Food Technology, Faculty of Science, University of Johannesburg, P. O. Box 17011, Doornfontein 2028, Johannesburg, South Africa.

²Department of Food Science and Technology, College of Food Sciences and Human Ecology, Federal University of Agriculture, Abeokuta, Nigeria.

***Corresponding author:** janet.adeyinka@yahoo.com; Tel: +27843485939

Abstract

With the prevalent food insecurity in Africa, there is a growing need to utilize the available crops to develop nutritious, affordable and palatable food for the populace. Millet is critical in this role, relative to its abundance in the continent and good nutritional composition. For ages, fermentation and malting have been traditionally used to transform millet into variety of produce. A paradigm shift has however occurred over the years, giving birth to new commercially available products. This review thus appraises and gives an overview of traditional and modern fermented and malted products. Although, millet has been diversified to several products, its major food uses are still restrained to traditional consumers and largely remains underutilized. Considering the potential embedded in this grain, it is important to explore this crop through the application of appropriate modern fermentation and malting technologies. This will ensure the availability of ready to eat (RTE) and ready to use (RTU) food products and to a large extent address the incessant food security challenges plaguing Africa.

Keywords

Millet, fermentation, malting, food products, processing, Africa.

1.0 Introduction

Cereals are important sources of the world's food supply and their role in human diet throughout the world is extremely vital. Cereal grains generally provide significant amount of nutrients and non-nutrients in the diet of populations all over the world, including developed and developing nations (Joshi *et al.*, 2008). In the light of this, the major cereals contributing to diet in the world are rice, barley, maize, wheat, sorghum, oat, rye and millet (Mridula and Sharma, 2015).

From this lot, are millets, commonly referred to as small-seeded grains, belonging to the *Poaceae* (*Gramineae*) family (Zhu, 2014). They are regarded as minor cereals (Shobana *et al.*, 2013) and are majorly known for their starchy edible seeds (Kothari *et al.*, 2005). Millets are important source of nutrients and an indispensable food for millions of people in developing countries such as Africa (Filli *et al.*, 2010; Amadou *et al.*, 2013; Mridula and Sharma, 2015). According to Joshi *et al.* (2008), millet is an important ecological food security crop known for its drought resistance and nutritional quality and can be an immediate subsistence food for a nutrient-scarce populace. This group of cereal crop has significant potential in widening the genetic diversity in the food basket and ensuring improved food and nutrition security (Mal *et al.*, 2010).

For centuries ago and at present, millet have been processed to transform them into other food products to increase their palatability, aesthetic value, acceptability and sensory attributes. Fermentation and malting are few of the most used processing techniques utilized for the diversification and transformation of millet. According to Lei and Michaelsen (2006), these groups of foods continue to constitute 20-40% of the food supply worldwide. Particularly in

ancient times and the situation in most developing countries where proper storage is unavailable, these processing techniques are widely utilized and of crucial importance. According to Onweluzo and Nwabugwu (2009) fermentation process prolongs the shelf-life of foods, increases protein content and digestibility, relative nutritive value, availability of minerals, functional properties and decreases the concentration of phytic acid while enhancing sensory properties and imparting antimicrobial properties on the food product (Jay *et al.*, 2005; Inyang and Zakari, 2008). Alternatively, malting is an easily adaptable technology which causes increased activities of hydrolytic enzymes leading to an increase in essential amino acids contents, total sugars, B-group vitamins, with a subsequent decrease in dry matter, starch and anti-nutrients (Traore *et al.*, 2004; Coulibaly *et al.*, 2012). Accordingly, both techniques maximize the levels of some of the utilizable nutrients and lower the dietary bulk in cereals (Akinhanmi *et al.*, 2008).

Historically in Africa, millets are processed by using two basic food processing techniques; fermentation and malting to obtain different products. However with the trend and drive in industrialization and improvement of these processing techniques, millet has been transformed to give birth to new, contemporary value-added products, further creating a huge potential and opportunity for the evolution of other ones. This review thus presents an appraisal and expedition of fermented and malted millet products from a traditional and rural food product to the commercially available contemporary, modern and industrialized food product. This review also draws attention to its prospects which would further increase its utilization, diversification and consequent consumption.

2.0 Millets

Millet is a generic term referring to an extraneous group of forage grasses (Table 1) recognized for producing small-sized grains (Muthamilarasan *et al.*, 2016). It is a word derived from the French word “*mille*” which means thousand, literarily indicating that a handful of millet has thousands of grains (Taylor and Emmambux, 2008). They belong to the grass family *Poaceae* (*Gramineae*) and are believed to be cultivated since establishment of human civilization (Shobana *et al.*, 2013). Species of this cereal crop have been said to originate from China, Asia and Africa (Oelke *et al.*, 1990). Relative to this, they are sometime referred to as the first cereal crop (Lu *et al.*, 2005). In terms of world production, it is the sixth most important cereal crop after wheat, maize, rice, barley and sorghum with an annual production of about 29 million tonnes in 2013 (FAOSTAT, 2015).

This crop is unique compared to other cereals like maize and rice, because of its superior nutritional qualities especially valuable amino acid, methionine, higher protein, quality macro and micronutrients (Shobana and Malleshi, 2007; Saleh *et al.*, 2013). Millets are also good sources of nutraceutical and functional food ingredients in health promotion, mainly due to their antioxidant, antimicrobial, anti-inflammatory, antiviral, anticancer, antiplatelet aggregation, and cataractogenesis inhibitory activities (Chandrasekara and Shahidi, 2011; Taylor and Duodu, 2015; Muthamilarasan *et al.*, 2016).

Though millets are usually consumed by people from the lower economic strata, they are nutritionally rich (Taylor *et al.*, 2010a; Saleh *et al.*, 2013). Millet has been termed as “nutri-cereals”, because they are rich in vitamins and sulphur containing amino acids (Table 2) (Sangeetha and Devi, 2012). According to Leder (2004), millet contains 22-28% albumins and

globulins, 28-32% glutelin and glutelin-like proteins, while its prolamin content ranges from 22-35%. These nutrients are reportedly lacking in diets of the poor who live mostly on starchy staple meals (Srivastava and Sharma, 2012). Millets are also rich sources of phytochemicals, polyphenols, soluble and insoluble dietary fibers, minerals, and antioxidants (Chethan and Malleshi 2007; Taylor and Duodu, 2015; Muthamilarasan *et al.*, 2016). Its fatty acids are highly poly unsaturated and contain relatively high proportions of non-starchy polysaccharides (Hegde and Chandra 2005; Muthamilarasan *et al.*, 2016). Millets are also known to slowly release sugars and consequently have a low glycemic index (Bala-Ravi, 2004). It is also gluten free, which makes it an excellent choice for people suffering from celiac diseases caused due to gluten intolerance (Taylor *et al.*, 2006).

3.0 Processing of millet through fermentation and malting

Processing of foods is an essential step in transforming foods to improve sensorial, nutritive value and enhancing bioavailability of nutrients (Liu *et al.*, 2012). Other desired physical and chemical changes include modification of the macro- and micro- structure of the product and starch digestibility (Concy, 2009; Amadou *et al.*, 2014; Dharmaraj *et al.*, 2014). Processing of millet has also been reported to decrease levels of antinutrients (Hotz and Gibson 2007; Taylor and Duodu, 2015). While several techniques have been utilized for processing millets, fermentation and malting are the oldest and most common forms of processing techniques (Pugalenth and Vadivel, 2005; Singh *et al.*, 2015). This can be attributed to their relative simplicity, cheapness and availability. Notwithstanding, several attempts towards improved techniques for these processes have been made and are existing. These are highlighted in the proceeding sections of this review.

3.1 Fermentation

Fermentation is one of the oldest and widely used methods adopted for the processing of millets. It is a process dependent on the biological activity of certain microorganisms [lactic acid bacteria (LAB) and yeasts], leading to the production of a wide range of metabolites (Ross *et al.*, 2002). It has been regarded as an economical technique of processing food and its relevance is evident in the myriad of available products in African.

Simply defined, it is a process performed to transform substrates into new products through the action of microorganisms. Biochemical changes that occur during this process lead to the modification of the substrate and production of volatiles (Singh *et al.*, 2015). The fermentation process lead to the activation of enzymes, adjustment of pH and performance enhancement of certain enzymes including proteases, amylases and hemicellulases (Jay *et al.*, 2005). These changes result in the beneficial properties associated with fermentation including preservation through alcoholic, lactic acid, alkaline, acetic acid fermentation and the impartation of antimicrobial properties (Jay *et al.*, 2005). Fermentation also ensures the hygienic safety of the food by suppressing the growth and survival of undesirable microflora in food (Ross *et al.*, 2002). According to Osman (2004) and Eltayeb *et al.* (2007), this process also leads to detoxification, decrease in trypsin and amylase inhibitory activity, phytic acid and tannins. Nutritional enrichment by increasing protein content, mineral extractability and palatability have also been reported (El Hag *et al.*, 2002; Ali *et al.*, 2003; Badau *et al.*, 2005 Osman, 2011; Ranasalva and Visvanathan, 2014). Fermentation further results in a decrease in cooking time (Jay *et al.*, 2005; Sasikumar, 2014) and enrichment of sensorial qualities through the

development of diverse flavors, aromas and texture (Jay *et al.*, 2005; Kohajdova and Karovicova, 2007).

3.1.1 Natural or spontaneous fermentation

Majority of the fermented foods from Africa are being produced through natural or spontaneous fermentation. This process is mostly managed by women in households or at a commercial small scale level (Figure 1). Depending on the type of product, such fermentation could be only lactic acid fermentation or both lactic acid and alcoholic fermentation for the production of alcoholic beverages (Kohajdova and Karovicova, 2007). Under such conditions, millet grains are winnowed to remove debris, stones and other foreign materials. These are rarely cleaned and soaked into water for days (depending on the period required for the desired product). Spontaneously, fermentation results as a consequence of the competitive activities of different microorganisms and strains best adapted and with higher growth rates usually dominate (Aka *et al.*, 2014). Investigations into microbial populations responsible for this type of fermentation have been investigated over the years and a dominance of LAB have been reported (Galati *et al.*, 2014). Nevertheless, different undesirable species of bacteria, yeasts and fungi have been detected and implicated for contributing to spoilage and pathogenic traits (Humblot and Guyot, 2009; Aka *et al.*, 2014; Galati *et al.*, 2014).

Similar to natural or spontaneous fermentation is back-slopping, commonly practiced traditionally, in low-income households and small scale commercial enterprises. This involves the introduction of a small portion of a previous successful fermentation batch (Aka *et al.*, 2014)

which serve as a source of starter cultures and guarantees the successful transfer of the responsible LAB responsible for fermentation.

3.1.2 Controlled fermentation

As earlier stated, undesirable microorganisms may develop during the spontaneous fermentation of foods and upon reaching a maximum level, undesirable microflora of fermentation could be detected in fermentation process (Corsetti and Settanni, 2007). Microorganisms implicated in such conditions include *Fusarium*, *Aspergillus*, *Acetivibrio*, *Candida*, *Enterobacter*, *Aerobacter* and *Bacillus* (Galati *et al.*, 2014). The presence and occurrence of these organisms represents a food safety concern as some are pathogenic while some of the fungal species produce some toxigenic mycotoxins (Abriouel *et al.*, 2006; Humblot and Guyot, 2009; Njobeh *et al.*, 2010; Oguntuyinbo *et al.*, 2011; Galati *et al.*, 2014; Adebo *et al.*, 2015).

Over the years, several efforts have been directed towards the identification and selection of strains for a better fermentation process, referred to as controlled fermentation (Galati *et al.*, 2014). This had involved monitoring of dominant fermentation microorganisms and the analysis of the respective microbial biodiversity of fermented millet foods (Table 3). Controlled fermentation have since evolved into the selection of strains and starter cultures based on their specific biotechnological traits with a goal of obtaining final products of desired characteristics (Figure 1). While LAB is known for its rapid acidification, yeasts are selected for their higher alcoholic fermentation rate (Galati *et al.*, 2014; Singh *et al.*, 2015). Using the selected precise strain, the fermentation process is done under an optimal condition required by the strain, to yield a desired and safer product. Although concerted efforts from research entities, industries

and the government have yielded commercially available strains, challenges including financial implications, social and religious beliefs of African inhabitants have frustrated a holistic acceptance of this approach.

3.2 Malting

Similar to fermentation, malting of millet is a traditional practice in Africa, where the malt is subsequently used in lactic acid and alcoholic fermented beverages and infant food production (Adekunle, 2012). It could be simply defined as the germination of grain in moist air under controlled conditions to promote the development of hydrolytic enzymes which were inactive in the raw grain (Taylor and Dewar, 2001; Ayernor and Ocloo, 2007). These hydrolytic enzymes structurally modify and solubilize the grain structure, causes the breakdown of carbohydrates, lipids and proteins to simple sugars, lipids and amino acids respectively (Adekunle, 2012; Patil, 2013; Swami *et al.*, 2013). Malting also activates proteases which degrades protein and improves its bioavailability (Taylor *et al.*, 1985; Pyler and Thomas, 2000), decreases phytic acid and increases the extractability of calcium, iron and zinc in millets (Badau *et al.*, 2005; Lestienne *et al.*, 2005). This process has also been reported to enhance flavor and bioactivity of grains (Pyler and Thomas, 2000; Heinio *et al.*, 2001). In addition to starch degradation, there is a conversion of the proteinaceous matrix in the millet grain into polypeptides and amino acids by proteolytic enzymes (Umaru and Genwa, 2013).

The malting process involves three main operations, which are soaking, germination, and drying/kilning. The cleaned grains are stepped in water for 24 hours (Figure 2) after which they are drained. The wet grains are then spread on trays, floor or jute bags (Figure 2) for

germination, and water has to be added intermittently to facilitate the germination process. During the germination process, the grains are raked and turned to keep grains loose and aerated (Taylor and Emmambux, 2008). The germinated grains are then dried under controlled temperature to obtain malted grains which can be further processed to other food products discussed in the later part of this review. Due to the warm and damp conditions of grains, during the malting process, growth of molds and fungi are encouraged. However in rare cases, some species of these fungi produce mycotoxins which have been implicated in different poisonous outbreaks (Njobeh *et al.*, 2010).

Different researchers have suggested some propositions for a modern day malting process (Dewar *et al.*, 1997; Francakova *et al.*, 2012; Frere *et al.*, 2013). These include steeping grains with caustic soda or chloride salt, keeping germination time possibly shorter, using turners to prevent clumping of the germinating grains and preventing germinative stress. Further innovative and controlled malting systems, to provide high quality malted products which will enhance utilization of millet have also been industrialized (Figure 3).

4.0 Traditional fermented and malted millet-based food products

Africa is home to a vast and large number of fermented and millet food products (Table 4), most of which are traditionally processed and delivered to its intended consumers. Notable examples of these traditional products are given below;

4.1 Gruels, porridges and dumplings

Ogi is a traditional gruel or porridge prepared from fermented millet, sorghum or maize (Blandino *et al.*, 2003). It is a staple food for the inhabitants of West Africa and a meal/weaning

food for infants. *Ogi* (Figure 4) is traditionally prepared by soaking the grains in water for 24-72 hours, wet milling and sieving to remove the bran. The filtrate is further fermented for 2-3 days to yield *ogi*, a sour starchy sediment (Teniola and Odunfa, 2001). Traditionally, this product is often marketed as a wet cake wrapped in leaves or transparent plastic bags. Similarly to this is *ben saalga* (Figure 4), a semi-solid millet based gruel commonly eaten by children (Guyot *et al.*, 2003). It is a popular food in Burkina Faso (Tou *et al.*, 2007) and its processing has been described to involve soaking, milling, sieving, settling (during which fermentation occurs) followed by a cooking process (Tou *et al.*, 2006).

Ghanaian *koko* is a millet porridge daily consumed as an in-between meal or snack (Muller, 1970; Lei and Jakobsen, 2004). Production is done by steeping pearl millet overnight, wet milling the grain (with spices), addition of water (to make slurry), sieving, fermentation and sedimentation (Lei and Jakobsen, 2004). This is followed by boiling of the top water and addition of the sedimented bottom layer to obtain the desired consistency (Muller, 1970; Lei and Jakobsen, 2004; Lei and Michaelsen, 2006). Likewise, *fura* (*furah*) (Figure 4) is a traditional semi-solid dumpling prepared by steam cooking of fermented moist flour balls of pearl millet, followed by pounding in a mortar and rolling (Owusu-Kwarteng *et al.*, 2010). It is a staple food in Nigeria, Ghana and Burkina Faso and produced by blending millet with water and spices, compressing them into dough ball and subsequent cooking (Jideani *et al.*, 2002). From Botswana and Southern Africa is *bogobe* (*ting*), made from fermented maize, sorghum or pearl millet. *Dégué* (*tchobal*) is also a spontaneous, fermented and gelatinized pearl millet product, consumed by children and adults from Burkina Faso (Abriouel *et al.*, 2006). After dehulling, the grains are grinded with water and steam cooked to produce gelatinized balls which are further fermented

for 24 h and afterwards consumed with dairy products and juice (Abriouel *et al.*, 2006; Hama *et al.*, 2009). From Benin is *gowe*, a sweetish gelatinized, fermented and malted, cooked paste made from sorghum, millet and/or maize flour (Adinsi *et al.*, 2014). It is usually consumed as a porridge or as a beverage, after dilution with water and addition of sugar, ice and sometimes milk.

4.2 Alcoholic and non-alcoholic beverages

Alcoholic and non-alcoholic beverages produced from millet are more prevalent compared to other types of products. Pundits have argued that this can be attributed to survival instincts of the first African settlers. Known as *malwa*, *bantu*, *pombe*, opaque or *kaffir* beer are alcoholic beverages made from malted or fermented millet grains, although other grains such as maize and sorghum are commonly used (Figure 5). They are consumed in different parts of Central, East and West Africa and plays an important role in the culture and other social activities of the inhabitants of this region. *Bjalwa bja leotsa* and *ndlovo* are alcoholic traditional beers of the Pedi people and Zimbabweans respectively (Quin, 1959; Taylor and Emmambux, 2008). Similar to these other aforementioned alcoholic beverages, is *pito* (Figure 5) a light brown alcoholic beverage prepared from fermented millet or sorghum and consumed in Nigeria, Ghana and other parts of West Africa (Ajiboye *et al.*, 2014). It is a nutritious source of instant energy for Nigerian tribes and used for cultural purposes (Tamang, 2010). Accordingly, majority of these traditional millet beers are mostly consumed in their active state of fermentation. In Egypt *boza* (*buza*, *bouza*) is commonly brewed from pearl millet (Arici and Dagliogu, 2002). It is thick, pale yellow and has a characteristic acid-alcoholic aroma. Also in Ethiopia, a traditional beer and spirit has been prepared from finger millet is commonly known as *tella* and *katikalla* respectively (Bultosa

and Taylor, 2004). In Benin and Burkina Faso, a similar traditional millet beer called *tchouk*, *dolo* or *chapalo* has also been prepared. *Merrisa* is a Sudanese traditional alcoholic beverage manufactured using malted sorghum or millet (Dirar, 1992; Lyumugabe *et al.*, 2012). Its production has been described as a complex procedure and probably the most complicated brewing process of all traditional millet-based African alcoholic beverages (Lyumugabe *et al.*, 2012). The process has been extensively described and the resulting drink is reported to have a pH of 4 and alcohol level of 5% (v/v) (Dirar, 1976; Lyumugabe *et al.*, 2012).

Kunu-zaki (*kunu*) (Figure 6) is a non-alcoholic fermented beverage commonly consumed in Nigeria, especially the Northern part (Blandino *et al.*, 2003). It is of high social and economic importance and can be prepared from maize, sorghum or pearl millet (Efiuvwevwere and Akona, 1995). Grains are washed, dried and coarsely ground. The resulting flour is mixed with hot water to form a paste which is spontaneously fermented for days to obtain the product. It is also spiced with ginger to improve sensorial qualities. *Oshikundu*, a similar fermented product has been prepared in Namibia from fermented-cooked pearl millet and sorghum malt flour (Taylor, 2004), while *togwa* is from maize meal and finger millet malt (Oi and Kitabake, 2003). In South-Western Uganda is *bushera* (*obushera*), a fermented and malted product of millet or sorghum, consumed by both young and old people in (Mukisa *et al.*, 2010). It is regarded as the most common traditional fermented beverage and a thirst quencher, energy drink and utilized for social purposes (Muyanga *et al.*, 2003). To prepare *bushera*, sorghum or millet malt flour is mixed with boiling water and left to cool, after which the malted millet/sorghum or a previously fermented batch is added to ferment for another 1-6 days (Muyanga *et al.*, 2003).

Mageu (mahewu) (Figure 6) is one of the most popular cereal based beverage in Southern Africa and some other Arabian Gulf countries (Bvochora *et al.*, 1999; McMaster *et al.*, 2005). It is a refreshing drink and also a weaning beverage for infants. Although it is commonly prepared with maize flour, some ethnic groups use sorghum and millet instead. Its preparation starts with cooking maize meal in water until a thin porridge is obtained, cooling and addition of sorghum or millet malt after the cooking stage for fermentation (Simatende *et al.*, 2015). Another prominent non-alcoholic beverage in East Africa (notably Uganda) is *uji*. It can be made from sorghum, maize or finger millet and could be fermented and unfermented (Onyango *et al.*, 2004; Kohajdova, 2014). The fermented one is prepared by a spontaneous or back slop fermentation process after which it is boiled, sweetened and consumed hot (Nout, 2009; Kohajdova, 2010; Nyanzi and Jooste, 2012).

4.3 Flatbreads and pancakes

Injera (Figure 7), is a leavened pancake with a honey-comb like texture (Belton and Taylor, 2004) and undergoes a mixed LAB and yeast fermentation (Gashe *et al.*, 1982). It is a common staple food in Ethiopia, Eritrea and Somalia. It is mostly made from *teff*, mixed with water and fermented for 24-48 hours. Part of the dough is gelatinized, remixed and baked. Gelatinization of the dough is to make the dough viscous and allow the *injera* rise and hold gas. Similar to *injera* is *kisra* (Figure 7), which is relatively thin, with neither holes nor a spongy texture (Badi *et al.*, 1989).

5.0 Value added and modern fermented and malted food products from millet

According to Jayne *et al.* (2014), five of the top ten fastest growing economies in the world are located in sub-Saharan Africa. This rapid economic transformation ongoing in the continent has spawned efforts channeled towards increasing industrialization and urbanization. This economic growth has thus led to the increased demand for convenience, RTU and RTE foods. By default, considerable and new opportunities towards the development of industries to meet this demand have risen.

Traditionally available raw materials and products are the first point of call to local and potential foreign investors and millet have positioned itself among other cereals as a viable resource. Additionally, demand for gluten free products from consumers with celiac disease has created a wider prospect for processed millet. As defined by Schuppan and Zimmer (2013), celiac disease is an inflammatory disease of the small intestine which arises in response to the consumption of gluten by genetically or medically intolerant persons. The primary and possibly best treatment of this allergy is through the consumption of a gluten-free diet (Rubi-Tapia *et al.*, 2013), which the consumption of millet based products readily provides. Hence, this section appraises and mentions some notable and significant fermented and malted millet based products which have transited from being a traditional home based product to a modern day commercially available merchandize, on the shelf in shops and stores.

One West African product that has undergone technological transformation over the years is *ogi* and *koko*. By the application of appropriate cereal science and technology, this produce have been transformed from the moist starchy sediment to a dehydrated packaged product (Figure 8). Further investigation into value-added potential of *ogi* have also led to the fortification of this

product with vitamins and compositing with soy flour to produce soy-*ogi* (Figure 8). Responding to the needs of alcohol consuming celiacs, a brewing company, formulated and created a gluten free ale from millet and sorghum (Figure 8). *Mbege* ale is made from millet, sorghum and banana. Likewise is the success story of the Southern African opaque beer that has been industrialized, packaged and commercialized as *chibuku shake shake* (Figure 9). It remains a popular beer especially in Botswana, Zambia and Zimbabwe. In Nigeria is *kunun tsamiya*, an instant beverage modified from *kunun-zaki* and in Uganda is *bushera* (Figure 10).

Consequent to the birth of these commercialized fermented millet foods, many African countries where such products are manufactured have benefited from the industrialization of their indigenous fermented foods through value addition to millet and from sale of the final processed products. Further to this is self-empowerment and the daily profits which comes along with the sale of these products.

6.0 Future prospects

Millet stands as a cereal crop which has being potentially unexplored. They are however viable raw materials for the development of food manufacturing industries, in contrast to other cereal crops which are not locally grown in Africa. While malting and fermentation remains a cheap and economical way of transforming millets into value added products, the unavailability of RTE products have however hindered the acceptability and usage of these products. With a growing population coupled with the challenges of mal- and under-nutrition, food shortage and food insecurity especially in the developing world, millet is vital in addressing this. Although, numerous millet based products are available, there is a huge potential to develop diverse food

products by using traditional and contemporary methods. Value addition to millets will thus offer variety, quality and convenience foods. With the growing demand for gluten free, functional and nutraceutical food products, millet products can be established as a substitute. Furthermore, these foods could be potential sources of beneficial bioactive compounds e.g. polyphenols which exerts health promoting properties and will address increased rates of diet related diseases.

7.0 Conclusion

While different studies have demonstrated the utilization of millet for food products, recommendations from such studies have been poorly accepted and rarely adopted by the appropriate authorities, thereby reducing wide range of acceptability. With the increasing need for the world to feed its growing population, it is imperative to explore food crops such as millets that are grown locally and consumed by low income households in developing nations. Moving forward, the agricultural productivity of this crop must be substantially improved to reduce their costs, which in turn make them more beneficial and attractive as compared to other cereals. Application of industrial processing technologies using modern equipment and optimized conditions will enhance the production of high quality and diversified food products at a commercial scale. This will further improve the productivity of millet and substantially contribute and drive economic development and food security in Africa.

Acknowledgement

This piece of work was financially supported via the University Research Committee (URC) Fellowship of the University of Johannesburg and the National Research Foundation (NRF) Fellowship granted to the main author (Adebiyi, J. A).

References

- Abriouel, H., Ben Omar, N., Lopez, R. L., Martinez-Canamero, M., Keleke, S., and Galvez, A. (2006). Culture-independent analysis of the microbial composition of the African traditional fermented foods *poto poto* and *dégué* by using three different DNA extraction methods. *Int. J. Food Microbiol.* **111**: 228–233.
- Adebo, O. A., Njobeh, P. B., Gbashi, S., Nwinyi, O. C., and Mavumengwana, V. (2015). Review on microbial degradation of aflatoxins. *Crit. Rev. Food Sci. Nutr.* In Press <http://dx.doi.org/10.1080/10408398.2015.1106440> (Accessed 25 December 2015).
- Adekunle, A. A. Agricultural Innovation in Sub-Saharan Africa: Experiences from Multiple Stakeholder Approaches. Forum for Agricultural Research in Africa, Ghana. 2012.
- Adinsi, L., Vieira-Dalode, G., Akissoe, N. H., Anihouvi, V., Mestres, C., Jacobs, A., Dlamini, N., Pallet, D., and Hounhouigan, J. D. (2014). Processing and Quality Attributes of *Gowe*: a Malted and Fermented Cereal-Based Beverage from Benin. *Food Chain* **4**(2): 171-183.
- Ajiboye, T. O., Iliasu, G. A., Adeleye, A. O., Abdussalam, F. A., Akinpelu, S. A., Ogunbode, S. M., Jimoh, S. O., and Oloyede, O. B. (2014). Nutritional and antioxidant dispositions of sorghum/millet beverages indigenous to Nigeria. *Food Sci. Nutr.* **2**(5): 597-604.
- Aka, S., Konan, G., Fokou, G., Dje, K. M., and Bassirou, B. (2014). Review on African traditional cereal beverages. *Am. J. Res. Comm.* **2**(5): 103-153.
- Ali, M. A. M., El Tinay, A. H., Abdalla, A. H. (2003). Effect of fermentation on the in vitro protein digestibility of pearl millet. *Food Chem.* **80**(1): 51-54.

- Akinhanmi, F. T., Arogundade, A. L., Tiamiyi, O. M., and Oloruntoba, E. (2008). Protein fractions of legumes and cereals consumed in Nigeria. *Int. J. Agric. Sci. Environ. Technol.* **7**(1): 54-62.
- Amadou, I., Mahamadou, E. G., and Guo-Wei, L. (2013). Millets: nutritional composition, some health benefits and processing – a review. *Emir. J. Food Agric.* **25**(7): 501-508.
- Amadou, I., Gounga, M. E., Shi, Y. H., and Le, G.W. (2014). Fermentation and heat moisture treatment induced changes on the physicochemical properties of foxtail millet (*Setaria italica*) flour. *Food Bioprod. Process.* **92**: 38-45.
- Arici, M., and O. Daglioglu, (2002). *Boza*: A lactic acid fermented cereal beverage as a traditional Turkish food. *Food Rev. Int.* **18**: 39–48.
- Ayernor, G. S., and Ocloo, F. C. K. (2007). Physico-chemical changes and dia-static activity associated with germinating paddy rice (PSB. Rc 34). *African J. Food Sci.* **1**: 37–41.
- Badau, M., Nkama, H. I., and Jideani, A. I. (2005). Phytic acid content and hydrochloric acid extractability of minerals in pearl millet as affected by germination time and cultivar. *J. Food Chem.* **92**: 425-435.
- Badi, S. M., Bureng, P. L., and Monowar, L. Y. (1989). Commercial production: A breakthrough in *kisra* technology. **In**: *ICC 4th Quadrennial Symposium on Sorghum and Millets, Lausanne, Switzerland*. Dendy, D. A. V. Ed., International Association for Cereal Science and Technology, Vienna.
- Bala-Ravi, S. (2004). Neglected millets that save the poor from starvation. Low external input sustain. *Agric. India* **6**(1): 34-36.

- Blandino, A., Al-Aseeri, M. E., Pandiella, S. S., Cantero D., and Webb, C. (2003). Cereal-based fermented foods and beverages. *Food Res. Int.* **36**(6): 527-543.
- Bultosa, G., and Taylor, J. R. N. (2004). Teff. **In:** Encyclopedia of Grain Science, Wrigley, C., Corke, H., and Walker, C. E, Eds., Elsevier, Amsterdam.
- Bvochora J. M., Reed J. D., Read J. S., and Zvauya, R. (1999). Effect of fermentation processes on proanthocyanidins in sorghum during preparation of *mahewu*, a non-alcoholic beverage. *Process Biochem.* **35**: 21-25.
- Chandrasekara, A., and Shahidi, F. (2011). Bioactivities and antiradical properties of millet grains and hulls. *J. Agr. Food Chem.* **59**: 9563–9571.
- Chethan, S., Malleshi, N. G. (2007). Finger millet polyphenols: Optimization of extraction and the effect of pH on their stability. *Food Chem.* **105**(2): 862–70.
- Chipo, C. (2013). Mahewu, a traditional African beverage. The African pot nutrition. <http://theafricanpotnutrition.com/2013/10/01/mahewu-a-traditional-african-beverage/> (Accessed 13 January 2016).
- Concy, A. J. (2009). Effect of roasting on the nutritional quality of finger millet. An MSc dissertation submitted to the school of graduate studies, Makerere University, Uganda.
- Corsetti, A., and Settanni, L. (2007). *Lactobacilli* in sourdough fermentation: A review. *Food Res. Int.* **40**: 539-558.
- Coulibaly, A., Kouakou, B., and Chen, J. (2012). Extruded adult breakfast based on milled and soybean: Nutritional and functional qualities, source of glycemic low food. *J. Nutr. Food Sci.* **2**(7): 1-9.

- Dewar, J., Taylor, J. R. N., and Berjak, P. (1997). Determination of improved steeping conditions for sorghum malting. *J. Cereal Sci.* **26**: 129-136.
- Dharmaraj, U., Parameswara, P., and Somashekar, R. (2014). Effect of processing on the microstructure of finger millet by x-ray diffraction and scanning electron microscopy. *J. Food Sci. Tech.* **51**(3): 494-502.
- Dirar, H. A. (1976). The art and science of Merissa fermentation. Sudan Notes and Records **57**: 115-129.
- Dirar, H. A. (1992). Sudan's fermented food heritage. **In**: Applications of Biotechnology to Fermented Foods: Report of an Ad Hoc Panel of the Board on Science and Technology for International Development. National Academies Press, Washington D C, USA.
- Efiuvwevwere B. J., and Akona, O. (1995). The microbiology of 'kunun-zaki', a cereal beverage from Northern Nigeria, during the fermentation (production) process. *World J. Microbiol. Biotech.* **11**(5): 491-493.
- Egwim, E., Musa, A., Abubakar, Y., and Mainuna, B. (2013). Nigerian indigenous fermented foods: processes and prospects. **In**: Mycotoxin and Food Safety in Developing Countries. Makun, H. A., Ed., InTech, Croatia.
- El Hag, M. E., El Tinay, A. H., and Yousif, N. E. (2002). Effect of fermentation and dehulling on starch, total polyphenols, phytic acid and invitro digestibility of pearl millet. *Food Chem.* **77**: 193-196.
- Eltayeb, M. M., Hassan, A. B., Sulieman, M. A., and Babiker, E. E. (2007). Effect of processing followed by fermentation on antinutritional factors content of pearl millet (*Pennisetum glaucum* L.) cultivars. *Pak. J. Nutr.* **6**: 463-467.

- FAOSTAT (Food and Agriculture Organization Statistics) (2015). 2013 data for major commodities. <http://faostat.fao.org/site/567/DesktopDefault.aspx?PageID=567#ancor> (Accessed 24 November 2015).
- Filli, K. B., and Nkama, I. (2007). Hydration properties of extruded *fura* from millet and legumes. *British Food J.* **109**: 68-80.
- Filli, K. B., Nkama, I., Abubakar, U. M., and Jideani, V. A. (2010). Influence of extrusion variables on some functional properties of extruded millet-soybean for the manufacture of *fura*: a Nigerian traditional food. *Afr. J. Food Sci.* **4**: 342-352.
- Francakova, H., Liskova, M., Bojnanska, T., and Marecek, J. (2012). Germination index as an indicator of malting potential. *Czech J. Food Sci.* **30**(4): 377-384.
- Franz, C. M. A. P., Huch, M., Mathara, J. M., Abriouel, H., Benomar, N., Reid, G., Galvez, A., and Holzapfel, W. H. (2014). African fermented foods and probiotics. *Int. J. Food Microbiol.* **190**: 84-96.
- Frere, S., Phillips, P. J., and Yin, X. S. (2013). Method for improving yield in malting process. Patent no: WO2011127372. <http://www.google.com/patents/WO2011127372A1?cl=en> (Accessed 14 January 2016).
- Gadaga, T. H., Mutakumira, A. N., Narvhus, J. A., and Ferusu, S. B. (1999). A review of traditional fermented foods and beverages in Zimbabwe. *Int. J. Food Microbiol.* **53**: 1–11.
- Galati, A., Oguntotinbo, F. A., Moschetti, G., Crescimanno, M., and Settanni, L. (2014). The cereal market and the role of fermentation in cereal-based food production in Africa. *Food Rev. Int.* **30**: 317-337.

- Gashe, B. A., Girma, M., and Bisrat, A. (1982). Teff fermentation. I. The role of microorganisms in fermentation and their effect on the nitrogen content of teff. *SINET: Eth. J. Sci.* **5**: 69–76.
- Geervani, P., and Eggum, O. (1989). Nutrient composition and protein composition of minor millets. *Plant Food Hum. Nutr.* **39**: 201–208.
- Gopalan, C., Rama Sastri, B. V., and Balasubramanian, S. C. (2009). Nutritive value of Indian foods. Hyderabad, India: National institute of Nutrition.
- Guyot, J. P., Mouquet, C., Tou, E., Counil, E., Traore, A. S., and Treche, S. (2003). Study of the processing of pearl millet (*Pennisetum glaucum*) into *ben-saalga*, a fermented gruel from Burkina Faso. 2nd International Workshop on Food based approaches for a healthy nutrition. http://horizon.documentation.ird.fr/exl-doc/pleins_textes/divers11-08/010036797.pdf (Accessed 15 January 2016).
- Hama, F., Savadogo, A., Quattara, C. A. T., and Traore, A. S. (2009). Biochemical, microbial and processing study of *dégué* a fermented food from pearl millet dough from Burkina Faso. *Pak. J. Nutr.* **8**: 759-764.
- Hegde, P. S., and Chandra, T. S. (2005). ESR spectroscopic study reveals higher free radical quenching potential in *kodo* millet (*Paspalum Scrobiculatum*) compared to other millets. *Food Chem.* **92**: 177–82.
- Heinio, R. L., Oksman-Caldentey, K. M., Latva-Kala, K., Lehtinen, P., and Poutanen, K. (2001). Effects of drying treatment conditions on sensory profile of germinated oat. *Cereal Chem.* **78**: 707–714.

- Hotz, C. and Gibson, R. S. (2007). Traditional food-processing and preparation practices to enhance the bioavailability of micronutrients in plant-based diets. *J. Nutr.* **37**: 1097–100.
- Humblot, C., and Guyot, J. P. (2009). Pyrosequencing of tagged 16S rRNA gene amplicons for rapid deciphering of the microbiomes of fermented foods such as pearl millet slurries. *Appl. Environ. Microbiol.* **75**: 4354–4361.
- Icard-Verniere, C., Ouattara, L., Avallone, S., Joseph, H., Kayode, P., Amoussa, W., and Fatoumata, B. H. (2010). Traditional recipes of millet-, sorghum- and maize-based dishes and related sauces frequently consumed by young children in Burkina Faso and Benin. https://www.google.co.za/url?sa=t&rct=j&q=&esrc=s&source=web&cd=2&cad=rja&uact=8&ved=0ahUKEwiBio7v3avKAhXIAxoKHfJWDZcQFgghMAE&url=http%3A%2F%2Fwww.instapa.org%2Fdownloadattachment%2F2198%2F1687%2FInstapa_booklet_EN_webversion.pdf&usg=AFQjCNEJ4TsT_k69MH_MWnbUbwAXx-DqTA&sig2=-zIImHHXxcPSx9ahEMMYuw (Accessed 15 January 2016).
- Ilango, S., and Antony, U. (2013). Assessment of the microbiological quality of *koozh* a fermented millet beverage. *Afr. J. Microbiol. Res.* **8**(3): 308-312.
- Inyang, C. U., and Zakari, U. M. (2008). Effect of germination and fermentation of pearl millet on proximate, chemical and sensory properties of instant “*fura*”- a Nigerian cereal food. *Pak. J. Nutr.* **7**: 9-12.
- Jay, J. M., Loessner, M. J., and Golden, D. A. (2005). Modern Food Microbiology, 7th ed. Springer, India.

- Jayne, T. S., Meyer, F., and Traub, L. N. (2014). Africa's evolving food systems: drivers of change and the scope for influencing them. <http://pubs.iied.org/pdfs/14637IIED.pdf> (Accessed 13 January 2016).
- Jideani, V. A., Nkama, I., Agbo, E. B., and Jideani, I. A. (2002). Modelling of visual shelf life of millet (*Pennisetum glaucum*) dough (*fura*) as affected by time- temperature and product packaging parameters. *Cereal Chem.* **79**(5): 710 – 714.
- Joshi, A., Rawat, K., and Bhawana, K. (2008). Millet as “religious offering” for nutritional, ecological, and economical security. *Comp. Rev. Food Sci. Food Safety.* **7**: 369–372.
- Kohajdova, Z., and Karovicova, J. (2007). Fermentation of cereals for specific purpose. *J. Food Nutr. Res.* **46**(2): 51-57.
- Kohajdova, Z. (2010). Fermented cereal products. **In:** Comprehensive Food Fermentation Biotechnology. Pandey, A., Soccol, C. R., Gnansounou, E., Larroche, C., Nigam, P. S., and Dussap, C. G., Eds., Asiatech Publishers Inc., New Dehli.
- Kohajdova, Z. (2014). Fermented cereal products. **In:** Microorganisms and Fermentation of Traditional Foods. Ramesh, R. C., and Didier, M., Eds., CRC Press, London.
- Kothari, S. L., Kumar, S., Vishnoi, R. K., Kothari, A., and Watanabe, K. N. (2005). Applications of biotechnology for improvement of millet crops: Review of progress and future prospects. *Plant Biotech.* **22**(2): 81-88.
- Leder, I. (2004). Sorghum and millets. **In:** Cultivated Plants, Primary as Food Sources - Encyclopedia of Life Support Systems, Fuleky, G, Ed., EOLSS Publishers, Oxford, UK.

- Lei, V., and Jakobsen, M. (2004). Microbiological characterization and probiotic potential of koko and koko sour water, African spontaneously fermented millet porridge and drink. *J. Appl. Microbiol.* **96**(2): 384-397.
- Lei, V., Frilis, H., and Michaelsen, K. F. (2006). Spontaneously fermented millet product as a natural probiotic treatment for diarrhea in young children: An intervention study in Northern Ghana. *Int. J. Food Microbiol.* **110**(1): 246-253.
- Lestienne, I., Besancon, P., Caporiccio, B., Lullien-Pellerin, V. and Treche, S. (2005). Iron and zinc in vitro availability in pearl millet flours (*Pennisetum glaucum*) with varying phytate, tannin, and fiber contents. *J. Agric. Food Chem.* **53**: 3240–3247.
- Liu, J., Tang, X., Zhang, Y., and Zhao, W. (2012). Determination of the volatile composition in brown millet, milled millet and millet bran by gas chromatography/mass spectrometry. *Mol.* **17**: 2271–82.
- Lu, H., Yang X., Ye, M., Liu, K. B., Xia, Z., and Ren, X. (2005). Millet noodles in late Neolithic China. *Nature* **437**: 13–14.
- Lyumugabe, F., Gros, J., Nzungize, J., Bajyana, E., and Thonart, P. (2012). Characteristics of African traditional beers brewed with sorghum malt: a review. *Biotechnol. Agron. Soc. Environ.* **16**: 509-530.
- Mal, B., Padulosi, S., and Ravi, S. B. (2010). Minor millets in South Asia: Learnings from IFAD-NUS Project in India and Nepal. Maccaresse, Rome, Italy: Bioversity Intl., and Chennai, India: M.S. Swaminathan Research Foundation: 1-185.
- Marsh, A. J., Hill, C., Ross, R. P., and Cotter, P. D. (2014). Fermented beverages with health-promoting potential: Past and future perspectives. *Trends Food Sci. Tech.* **38**: 113-124.

- McMaster, L. D., Kokott, S. A., Reid, S. J., and Abratt, V. R. (2005). Use of traditional African fermented beverages as delivery vehicles for *Bifidobacterium lactis* DSM 10140. *Int. J. Food Microbiol.* **102**(2): 231-237.
- Michael and Doria's Travel Tales. (2007). Geek Corps Mali. http://thembsterstravels.blogspot.co.za/2007_01_01_archive.html (Accessed 13 January 2016).
- Mridula, D., and Sharma, M. (2015). Development of non-dairy probiotic drink utilizing sprouted cereals, legume and soymilk. *LWT - Food Sci. Technol.* **62**: 482-487.
- Mukisa, I. M., Nsiimire, D. G., Byaruhanga, Y. B., Muyanja, C. M. B. K., Langsrud, T., and Narvhus, J. A. (2010). Obushera: Descriptive sensory profiling and consumer acceptability. *J. Sens. Stud.* **25**: 190-214.
- Muller, H. G. (1970). Traditional cereal processing in Nigeria and Ghana. *Ghana J. Agric. Sci.* **3**: 187-195.
- Muthamilarasan, M., Dhaka, A., Yadav, R., and Prasad, M. (2016). Exploration of millet models for developing nutrient rich graminaceous crops. *Plant Sci.* **242**: 89-97.
- Muyanga, C. M. B. K., Narvhus, J. A., Treimo, J., and Langsrud, T. (2003). Isolation, characterization and identification of lactic acid bacteria from *Bushera*: a Ugandan traditional fermented beverage. *Int. J. Food Microbiol.* **80**: 201-210.
- Nkama, I., and Malleshi, N. G. (1998). Production and nutritional quality of traditional Masa form mixture of rice, pearl millet, cowpea and groundnut. *Food Nutr. Bull.* **19**: 336-373.
- Nout, M. J. R. (2009). Rich nutrition for the poorest-Cereal fermentations in Africa and Asia. *Food Microbiol.* **26**: 685-692.

- Njobeh, P. B., Dutton, M. F., and Makun, H. A. (2010). Mycotoxins and human health: significance, prevention and control. **In:** Smart Biomolecules in Medicine. Mishra, A. K., Tiwari, A., and Mishra, S. B., Eds., VBRI Press, India.
- Nyanzi, R., and Jooste, P. J. (2012). Cereal based functional foods. **In:** Probiotics. Rigobelo, E.C., Ed., Intech, Rijeka.
- Obadina, A. O., Oyewole, O. B., and Awojobi, T. M. (2008). Effect of steeping time of milled grains on the quality of *kunnu-zaki* (A Nigerian beverage). *Afr. J. Food Sci.* **2**: 33-36.
- Oelke, E. A., Oplinger, E. S., Putnam, D. H., Durgan, B. R., Doll, J. D., and Undersander, D. J. (1990). Millets. <https://hort.purdue.edu/newcrop/afcm/millet.html> (Accessed 13 January 2016).
- Oguntoyinbo, F. A., Turlomousis, P., Gasson, M., and Narbad, A. (2011). Analysis of bacterial communities of traditional fermented West African cereal foods using culture independent methods. *Int. J. Food Microbiol.* **145**: 205–210.
- Oi, Y., and Kitabatake, N. (2003). Chemical composition of East African traditional beverage, *togwa*. *J. Agric. Food Chem.* **51**: 7024–7028.
- Okonkwo, S. L., and Ogbunike, R. U. (2011). Application of chemistry in the development of Africa using local resources: Production of malt drink from millet. *Res. J. Chem. Sci.* **1**(9): 87-89.
- Onweluzo, J. C., and Nwabugwu, C. C. (2009). Fermentation of millet (*Pennisetum americanum*) and pigeon pea (*Cajanus cajan*) seeds for flour production: Effects on composition and selected functional properties. *Pak. J. Nutr.* **8**: 737-744.

- Onyango, C., Henle, T., Hofmann, T., and Bley, T. (2004). Production of high energy density fermented *uji* using a commercial alpha amylase or by single-screw extrusion. *Lebensm. Wiss. Technol.* **37**: 401–407.
- Osman, M. A. (2004). Changes, in sorghum enzyme inhibitors, phytic acid, tannins, and invitro protein digestibility occurring during *khamir* (local bread) fermentation. *Food Chem.* **88**: 129-134.
- Osman, M. A. (2011). Effect of traditional fermentation process on the nutrient and antinutrient contents of pearl millet during preparation of *Lohoh*. *J. Saudi Soc. Agric. Sci.* **10**(1), 1-6.
- Owusu-Kwarteng, J., Tano-Debrah, K., Glover, R. L. K., and Akabanda, F. (2010). Process characteristics and microbiology of *fura* produced in Ghana. *Nat. Sci.* **8**(8): 41-51.
- Patil, S. S. (2013). Consumer acceptance and marketing of *ragi* malt. *Indian J. Appl. Res.* **3**(12): 322-323.
- Pugalenthi, M., and Vadivel, V. (2005). Nutritional evaluation and the effect of process sing methods on antinutritional factors of sword bean (*Canavalia gladiata*). *J. Food Sci. Technol.* **42**: 510–516
- Pyler, R. E., and Thomas, D. A. (2000). Malted cereals: their production and use. **In**: Handbook of Cereal Science and Technology, Kulp, K., and Ponte, J. G, Eds., Marcel Dekker, New York.
- Quin, P. J. (1959). Foods and the Feeding Habits of the *Pedi*. Johannesburg: Witwatersrand University Press.
- Ranasalva, N., and Visvanathan, R. (2014). Development of cookies and bread from cooked and fermented pearl millet flour. *Afr. J. Food Sci.* **8**(6): 330-336.

- Ross, R. P., Morgan, S., and Hill, C. (2002). Preservation and fermentation: past, present and future. *Int. J. Food Microbiol.* **79**: 3-16.
- Rubio-Tapia, A., Hill, I. D., Kelly, C. P., Calderwood, A. H., and Murray, J. A. (2013). Diagnosis and management of celiac disease. *Am. J. Gastroenterol.* **108**: 656-676.
- Saleh, A. S. M., Zhang, Q., Chen, J., and Shen, Q. (2013). Millet Grains: Nutritional Quality, Processing, and Potential Health Benefits. *Compr. Rev. Food Sci. Food Safety.* **12**: 281-295.
- Sangeetha, N., and Devi, P. M. (2012). Effect of dehydration on the quality characteristics of extruded pasta using millet milk powder. *J. Nutr. Food Sci.* **2**(10): 1-5.
- Sasikumar, R. (2014). Fermentation technologies in food production. **In**: Progress in Biotechnology for Food Applications. Lai, W. F., Ed., OMICS Group, California, USA.
- Schuppan, D., and Zimmer, K. P. (2013). The diagnosis and treatment of celiac disease. *Dtsch Arztebl. Int.* **110**: 835-846.
- Shobana, S., and Malleshi N. G. (2007). Preparation and functional properties of decorticated finger millet (*Eleusine coracana*). *J. Food Eng.* **79**: 529-538.
- Shobana, S., Krishnaswamy, K., Sudha, V., Malleshi, N. G., Anjana, R. M., Palaniappan, L., and Mohan, V. (2013). Finger millet (*ragi*, *Eleusine coracana* L): a review of its nutritional properties, processing, and plausible health benefits. *Adv. Food Nutr. Res.* **69**: 1-39.
- Simatende, P., Gadaga, T. H., Nkambule, S. J., and Siwela, M. (2015). Methods of preparation of Swazi traditional fermented foods. *J. Ethn. Foods* **2**: 119-125.
- Singh, A. K., Rehal, J., Kaur, A., and Jyot, A. (2015). Enhancement of attributes of cereals by germination and fermentation: A review. *Crit. Rev. Food Sci. Nutr.* **55**: 1575-1589.

- Srivastava, K., Sharma, A. K. (2012). Nutraceutical importance of finger millet (*Eleusine coracana*) for improved human health. The European Journal of Plant Science and Biotechnology, 1-4.
- Swami, S. B., Thakor, N. J., and Gurav, H. S. (2013). Effect of soaking and malting on finger millet (*Eleusine Coracana*) grain. *Agric. Eng. Int.: CGIR J.* **15**(1): 194-200.
- Tamang, J. P. (2010). Dietary cultures and antiquity of fermented foods and beverages. In: Fermented foods and beverages of the world. Hui, Y. H, Özgül Evranuz, E., and Samuel, D., Eds., CRC Press, Boca Raton.
- Taylor, J. R. N., Noveille, L., and Liebenberg, N. W. (1985). Protein body degradation in the starchy endosperm of germinating sorghum. *J. Exp. Biol.* **36**: 1287–1295.
- Taylor, J. R. N., and Dewar, J. (2001). Developments in sorghum food technologies. *Adv. Food Nutr. Res.* **43**: 217–264.
- Taylor, J. R. N. (2004). Foods and nonalcoholic beverages. **In:** Encyclopedia of Grain Science. Wrigley, C., Corke, H., and Walker, C. E., Eds., Elsevier, Amsterdam.
- Taylor, J. R. N., Schober, T. J., and Bean, R. S. (2006). Novel food and non-food uses for sorghum and millets. *J. Cereal Sci.* **44**: 252-271.
- Taylor, J. R. N., and Emmambux, M. N. (2008). Gluten-free cereal products and beverages. **In:** Gluten-Free Foods and Beverages from Millets. Arendt, E. K., and Bello, F. D., Eds., Elsevier, Amsterdam.
- Taylor, J. R. N., Barrion, S. C., and Rooney, L. W. (2010a). Pearl millet - new developments in ancient food chain. *Cereal Foods World* **55**: 16–19.

- Taylor, J. R. N. (2010b). Sorghum malting: Introduction, objectives and scientific principles. International Sorghum and Millet Collaborative Research Support Program (INTSORMIL CRSP) Presentations, paper 21. <http://digitalcommons.unl.edu/cgi/viewcontent.cgi?article=1020&context=intsormilpresent> (Accessed 11 January 2016).
- Taylor, J. R. N., and Duodu, K. G. (2015). Effects of processing sorghum and millets on their phenolic phytochemicals and the implications of this to the health-enhancing properties of sorghum and millet food and beverage products. *J. Sci. Food Agric.* **95**: 225–237.
- Teniola, O. D., and Odunfa, S. A. (2001). The effects of processing methods on the levels of lysine, methionine and the general acceptability of *ogi* processed using starter cultures. *Int. J. Food Microbiol.* **63**: 1-9.
- Tou, E. H., Guyot, J. P., Mouquet-Rivier, C., Rochette, I., Counil, E., Traore, A. S., and Treche, A. S. (2006). Study through surveys and fermentation kinetics of the traditional processing of pearl millet (*Pennisetum glaucum*) into “*ben-saalga*”, a millet-based fermented gruel from Burkina Faso used as complementary food. *Int. J. Food Microbiol.* **106**: 52–60.
- Tou, E. H., Moquet-River, C., Picq, C., Traore, A. S., Treche, S., and Guyot, J. P. (2007). Improving the nutritional quality of *ben-saalga*, a traditional fermented millet-based gruel, co-fermenting millet with groundnut and modifying the processing method. *LWT-Food Sci. Technol.* **40**: 1561-1569.
- Traoré, T., Mouquet, C., Icard-Vernière, C., Traoré, A., and Trèche, S. (2004). Changes in nutrient composition, phytate and cyanide contents and α -amylase activity during cereal

malting in small production units in Ouagadougou (Burkina Faso). *Food Chem.* **88**: 105-114.

Umaru, A. A., and Genwa, E. I. (2013). Proximate analysis and hydrolysis of malted millet (*Panicum miliaceum*) blended with maize and rice flour. *Acad. J. Interdiscp. Stud.* **2**(12): 2281-3993.

Uno, D. (2005). Farmer's selection of local and improved pearl millet varieties in Ovambo land, Northern Namibia. *Afri. Study Monograph.* **30**: 107-117.

Zhu, F. (2014). Structure, physicochemical properties, and uses of millet starch. *Food Res. Int.* **64**: 200–211.

Table 1: Common millet species and botanical names

Species	Common name
<i>Brachiaria ramose</i>	Browntop millet
<i>Coix lachryma jobi</i>	Job's tears
<i>Digitaria exilis</i>	Hungry rice
<i>Digitaria ibura</i>	Fonio
<i>Echinochloa colona</i>	Jungle rice
<i>Echinochloa decompositum</i>	Australian millet
<i>Echinochloa frumentacea</i>	Japanese barnyard millet
<i>Eleusine coracana</i>	Finger millet
<i>Eragrostis teff</i>	Teff
<i>Panicum miliaceum</i>	Proso millet
<i>Panicum miliare</i>	Little millet
<i>Paspalum notatum</i>	Bahia grass
<i>Paspalum scrobiculatum</i>	Kodo millet
<i>Pennisetum glaucum</i>	Pearl millet
<i>Setaria italic</i>	Foxtail millet
<i>Urochloa deflexa</i>	Guinea millet

(Adapted from: Kothari *et al.*, 2005)

Table 2: Nutritional composition of millet varieties

	Finger millet	Proso millet	Foxtail millet	Little millet	Kodo millet	Barnyard millet
Proximates						
Moisture (g)	13.1	11.9	11.2	11.5	12.8	11.9
Protein (g)	7.3	12.5	12.3	7.7	8.3	6.2
Fat (g)	1.3	1.1	4.3	4.7	1.4	2.2
Minerals	2.7	1.9	3.3	1.5	2.6	4.4
Dietary fiber (g)	11.5	-	2.4	2.53	2.47	1.98
Carbohydrates (g)	72.0	70.4	60.9	67	65.9	65.5
Energy (kcal)	328	341	331	341	309	307
Vitamins						
Carotene (µg)	42	0	32	0	0	0
Thiamine (mg)	0.42	0.20	0.59	0.30	0.33	0.33
Riboflavin (mg)	0.19	0.18	0.11	0.09	0.09	0.10
Niacin (mg)	1.1	2.3	3.2	3.2	2.0	4.2
Choline (mg)	-	748	-	-	-	-
Minerals and trace elements						
Calcium (mg)	344	14	31	17	27	20
Phosphorus (mg)	283	206	290	220	188	280

	Finger millet	Proso millet	Foxtail millet	Little millet	Kodo millet	Barnyard millet
Iron (mg)	3.9	0.8	2.8	9.3	0.5	5.0
Magnesium (mg)	137	153	81	133	147	82
Sodium (mg)	11	8.2	4.6	8.1	4.6	-
Potassium (mg)	408	113	250	129	144	-
Copper (mg)	0.47	1.60	1.40	1.00	1.60	0.60
Manganese (mg)	5.49	0.60	0.60	0.68	1.10	0.96
Molybdenum (mg)	0.102	-	0.70	0.016	-	-
Zinc (mg)	2.3	1.4	2.4	3.7	0.7	3.0
Chromium (mg)	0.028	0.020	0.030	0.180	0.020	0.090
Sulfur (mg)	160	157	171	149	136	-
Chlorine (mg)	44	19	37	13	11	-
Essential amino acids						
Arginine (mg/g N)	300	290	220	250	270	-
Histidine (mg/g N)	130	110	130	120	120	-
Lysine (mg/g N)	220	190	140	110	150	-
Tryptophan (mg/g N)	100	50	60	60	50	-
Phenyl alanine (mg/g N)	310	310	420	330	430	-
Tyrosine (mg/g N)	220	-	-	-	-	-
Methionine (mg/g N)	210	160	180	180	180	-

	Finger millet	Proso millet	Foxtail millet	Little millet	<i>Kodo</i> millet	Barnyard millet
Cystine (mg/g N)	140	-	100	90	110	-
Threonine (mg/g N)	240	150	190	190	200	-
Leucine (mg/g N)	690	760	1040	760	650	-
Isoleucine (mg/g N)	400	410	480	370	360	-
Valine (mg/g N)	480	410	430	350	410	-

(Adapted from Geervani and Eggum, 1989; Gopalan *et al.*, 2009)

Table 3: African based millet fermented foods and their respective microflora

Product	Microflora	Product type
<i>Amgba</i>	LAB, yeasts	Alcoholic beverage
<i>Bel-saalga</i>	LAB	Gruel
<i>Bogbe</i>	<i>Lactobacilli</i> and yeast	Porridge
<i>Busa</i>	<i>Lactobacilli</i> , <i>Saccharomyces</i>	Liquid drink
<i>Bushera</i>	LAB, yeasts	Non-alcoholic beverage
<i>Dégué</i>	<i>Lactobacilli</i>	Dumpling
<i>Doro</i>	LAB, <i>S. cerevisiae</i>	Alcoholic beverage
<i>Injera</i>	<i>Candida guilliermondii</i>	Pancake
<i>Kenkey</i>	<i>Lactobacilli</i> and yeast	Porridge
<i>Kisra</i>	<i>Lactobacilli</i>	Pancake
<i>Kunu-zaki</i>	LAB, yeasts	Gruel or beverage
<i>Mahewu</i>	<i>C. krusei</i> , <i>S. cerevisiae</i> , <i>Lactobacilli</i>	Liquid drink
<i>Merissa</i>	<i>Saccharomyces</i>	Alcoholic drink
<i>Ogi</i>	Bacteria, yeasts	Gruel
<i>Tchouhoutou</i>	<i>Lactobacilli</i> , yeasts	Alcoholic beverage
<i>Togwa</i>	LAB, yeasts	Gruel or beverage
<i>Uji</i>	LAB	Porridge

Table 4: Some fermented and malted millet products from Africa

Products	Country	Reference
<i>Ben saalga</i>	Burkina Faso	Franz <i>et al.</i> (2014)
<i>Bushera</i>	Uganda	Marsh <i>et al.</i> (2014)
<i>Dégué</i>	Burkina Faso	Franz <i>et al.</i> (2014)
<i>Doro</i>	Zimbabwe	Gadaga <i>et al.</i> (1999)
<i>Dolo (chapalo)</i>	Burkina Faso	Anon
<i>Fura</i>	Nigeria	Jideani <i>et al.</i> (2002)
<i>Gowe</i>	Benin	Adinsi <i>et al.</i> (2014)
<i>Injera</i>	Ethiopia, Eritrea	Gashe <i>et al.</i> (1982)
<i>Kisra</i>	Sudan	Gashe <i>et al.</i> (1982)
<i>Koko</i>	Nigeria, Ghana	Marsh <i>et al.</i> (2014)
<i>Koozh</i>	Nigeria	Ilango and Antony (2013)
<i>Kunu-zaki</i>	Nigeria	Obadina <i>et al.</i> (2008)
<i>Mahewu</i>	South Africa	Marsh <i>et al.</i> (2014)
Malt drink	Nigeria	Okonkwo and Ogbuneke (2011)
<i>Masa (waina)</i>	Nigeria, Chad, Burkina Faso	Egwim <i>et al.</i> (2013)
<i>Masvusvu</i>	Zimbabwe	Amadou <i>et al.</i> (2013)
<i>Merissa</i>	Sudan	Dirar (1992)
<i>Ndlovo</i>	Zimbabwe	Taylor and Emmanbux (2008)
<i>Ogi</i>	Nigeria	Teniola and Odunfa (2001)

<i>Ontaku (oshikundu)</i>	Namibia	Uno (2005)
<i>Tchouk</i>	Benin	Anon
<i>Ting</i>	Botswana, South Africa	Quin (1959)
<i>Togwa</i>	Tanzania	Marsh <i>et al.</i> (2014)
<i>Uji</i>	East Africa	Onyango <i>et al.</i> (2004)




Figure 1: Natural/Spontaneous fermentation process for millet (A), Controlled fermentation process (B)




Figure 2: Traditional millet malting process; soaking (sprouting) in pots (A), germination on jute bags (B) (Adapted from Michael and Doria's Travel Tales, 2007).



Figure 3: A modern industrialized malting equipment (Adapted from Taylor, 2010b).



Figure 4: Millet based gruel, dumplings and porridges; *Ogi* (A), *ben saalga* (B) (Adapted from Icard-Verniere *et al.*, 2010), *fura* (C).



Figure 5: Traditional millet based alcoholic beverages; *pombe* from Mozambique (A), opaque beer from Zimbabwe (B), millet beer from Cameroon (C), *pito* from Nigeria (D).



Figure 6: Traditional millet based beverages; *kunu* from Nigeria (A), *bushera* from Uganda (B), *mahewu* from Southern Africa (C) (Chipo, 2013).



Figure 7: Millet based flatbreads/pancakes; *injera* being sold in an Ethiopian market (A), a Sudanese woman making *kisra* (B).



Figure 8: Modern millet products; packaged *ogi* (A) and (B), packaged *koko* (C) and (D).



Figure 9: Modern millet based alcohol; gluten free ale (A), *mbege* (B), *chibuku shake shake* (C).



Figure 10: Millet based beverages; instant *kunun* (A), *bushera* (B).