



Exploration of collagen recovered from animal by-products as a precursor of bioactive peptides: Successes and challenges

Yu Fu, Margrethe Therkildsen, Rotimi E. Aluko & René Lametsch

To cite this article: Yu Fu, Margrethe Therkildsen, Rotimi E. Aluko & René Lametsch (2018): Exploration of collagen recovered from animal by-products as a precursor of bioactive peptides: Successes and challenges, Critical Reviews in Food Science and Nutrition, DOI: [10.1080/10408398.2018.1436038](https://doi.org/10.1080/10408398.2018.1436038)

To link to this article: <https://doi.org/10.1080/10408398.2018.1436038>



Accepted author version posted online: 02 Feb 2018.



Submit your article to this journal [↗](#)



View related articles [↗](#)



View Crossmark data [↗](#)

Publisher: Taylor & Francis

Journal: *Critical Reviews in Food Science and Nutrition*

DOI: <https://doi.org/10.1080/10408398.2018.1436038>

Exploration of collagen recovered from animal by-products as a precursor of bioactive peptides: Successes and challenges

Yu Fu^{a*}, Margrethe Therkildsen^b, Rotimi E. Aluko^c, René Lametsch^a

Author affiliations:

^aDepartment of Food Science, Faculty of Science, University of Copenhagen, Rolighedsvej 26, 1958 Frederiksberg C, Denmark

^bDepartment of Food Science, Aarhus University, Blichers Allé 20, 8830 Tjele, Denmark

^cDepartment of Human Nutritional Sciences, University of Manitoba, Winnipeg, MB, Winnipeg, R3T 2N2, Canada

Corresponding author:

Yu Fu, Ph.D.

Department of Food Science, Faculty of Science, University of Copenhagen, Rolighedsvej 26, 1958 Frederiksberg C, Denmark

E-mail: yu@food.ku.dk

Phone: +45 35334298

Abstract

A large amount of food-grade animal by-products is annually produced during industrial processing and they are normally utilized as animal feed or other low-value purposes. These by-products are good sources of valuable proteins, including collagen or gelatin. The revalorization of collagen may lead to development of a high benefit-to-cost ratio. In this review, the major approaches for generation of collagen peptides with a wide variety of bioactivities were summarized, including antihypertensive, antioxidant and antidiabetic activities, and beneficial effects on bone, joint and skin health. The biological potentials of collagen peptides and their bioavailability were reviewed. Moreover, the unique advantages of collagen peptides over other therapeutic peptides were highlighted. In addition, the current challenges for development of collagen peptides as functional food ingredients were also discussed. This article discusses the opportunity to utilize collagen peptides as high value-added bio-functional ingredients in the food industry.

Keywords: Collagen; Gelatin; Bioactive peptides; Bioavailability; Functional foods

Abbreviations:

ACE: Angiotensin I-converting enzyme

DPP-IV: Dipeptidyl peptidase-IV

GIP: Glucose-dependent insulintropic polypeptide

GLP-1: Glucagon-like peptide-1

IC₅₀: Half-maximal inhibitory concentration

ROS: Reactive oxygen species

QSAR: Quantitative structure-activity relationship

T2D: Type 2 diabetes

Introduction

Industrial processing of land and aquatic animal produces tons of by-products. In the European Union, it is estimated that over 20 million tons of animal by-products are generated annually (Henchion et al., 2016; Martínez-Alvarez et al., 2015; Pal and Suresh, 2016). Nevertheless, these by-products are normally utilized for production of low-value products, including fertilizers, animal feed, pet food, etc (Alao et al., 2017). Therefore, alternative utilization of animal by-products for high-value products not only reduces the environmental impact but also provides new food ingredients and increases the potential revenue for food industry (Henchion et al., 2016). Mammalian skins, bones, connective tissue, and fish bone, fins and scales as main components of animal by-products can serve as promising sources of collagen and gelatin (Alemán and Martínez-Alvarez, 2013). Gelatin can be obtained by partial hydrolysis of collagen. In this regard, conversion of collagen/gelatin into high value-added food ingredients, e.g. collagen hydrolysates or peptides with potential bioactivities, has gained a great attention, which leads to a high benefit-to-cost ratio (Toldrá et al., 2016).

Bioactive peptides are defined as certain peptide fragments with 2-20 amino acid residues that can exert a positive impact on human health (Kitts and Weiler, 2003). They are inactive within the sequences of their parent proteins but exhibit the beneficial health-promoting effects once they are released. Enzymatic hydrolysis of collagen or gelatin can release encrypted bioactive peptide sequences with various physiological functions (Gómez-Guillén et al., 2011). Collagen peptides can be released from their precursors during food processing, gastric digestion and enzymatic hydrolysis through exogenous and endogenous proteases (Fu et al., 2017b; Udenigwe, 2014). The past decade has witnessed a growing number of studies on collagen peptides with antihypertensive, antioxidant, antidiabetic, bone, joint and skin health-promoting effects (Alemán and Martínez-Alvarez, 2013; Gómez-Guillén et al., 2011). Furthermore, a growing body of evidence supports the excellent bioavailability of collagen peptides (Liu et al., 2015; Zague, 2008). Although some review

papers focus on the functional and bioactive properties of collagen/gelatin peptides (Gómez-Guillén et al., 2011; Pal & Suresh, 2016), the major approaches for generating collagen peptides with various bioactivities are classified and the biological potentials of collagen peptides and their underlying mechanisms are highlighted in the present review. More importantly, unique advantages of collagen peptides and the current challenges for exploitation of collagen peptides-based functional food ingredients are summarized. The present work contributes to development of collagen peptides from animal by-products as functional food ingredients and provides some future directions and strategies to overcome these challenges.

Structure and physicochemical characteristics of collagen

Collagen, a major structural protein, can be found in animal processing by-products, including mammalian skin, bones and connective tissue, and also fish skin, bones and scales (Ferraro et al., 2016; Pal and Suresh, 2016). Collagen accounts for approximately 25-35% of all vertebrate body proteins (Banerjee and Shanthi, 2016). So far 29 different types (type I–IXXX) of collagen have been identified (Liu et al., 2015), among which type I collagen is the most abundant. Collagen molecule mainly consists of three almost identical α -chains, namely collagen triple helix. This triple helix is composed of the continuous repetitive motif, Gly-X-Y, where X is proline (Pro) and Y is hydroxyproline (Hyp). These two imino acids are specific amino acids of collagen, which contributes to the helical structure of each chain. Compared with mammalian collagen, a lower amount of imino acids (proline and hydroxyproline) is present in marine collagen, which leads to lower melting point and weaker gel strength (Karim and Bhat, 2008, 2009).

Collagen can be extracted from land and marine animal by-products through chemical or biochemical pretreatment, followed by extraction process by warm water (above 40 °C) (Gómez-Guillén et al., 2011). This process can cleave the covalent and hydrogen bonds, leading to destabilizing the triple-helix and further conversion to soluble gelatin (Gómez-Guillén et al., 2011).

Dependent on the pH, temperature and extraction time used, the extracted gelatin products vary in the functional properties. However, gelatin as a denatured product of collagen shares similar structures with collagen (Schrieber and Gareis, 2007). In general, there are two types of gelatin, namely type A and type B. Type A gelatin is derived from the acid-extracted raw material, while type B gelatin is obtained through an alkaline extraction process. Even though mammalian gelatins have been extensively used in different fields, the utilization of gelatin derived from marine sources is increasingly attractive, as it is not involved in the risk of diseases, such as bovine spongiform encephalopathy and also meet certain religious requirements (Karim and Bhat, 2009).

Collagen peptides prepared from hydrolyzed collagen and low-molecular-weight gelatin products (0.3 - 8 kDa) are water-soluble, readily digestible and easily absorbed by the body (Sibilla et al., 2015). These characteristics coupled with the unique Gly-X-Y motif confer collagen peptides with versatile functionalities and various applications as food ingredients and supplements in the form of powders, drinks as well as liquid sprays (Hayes, 2011; Venkatesan et al., 2017). The structural features and functionalities of collagen and collagen peptides have been reviewed by Gómez-Guillén et al. (2011), as summarized in Fig. 1.

Generation of collagen-derived bioactive peptides

In recent years, there has been an increasing number of investigations on bioactive collagen peptides derived from different animal by-products. Several approaches for generation of collagen peptides with different bioactivities have been employed, such as *in vitro* and *in silico* hydrolysis. In addition, the integrated approach (peptidomics combined with *in silico* prediction) can also be used to identify novel collagen peptides with diverse bioactivities. The major research approaches on collagen-derived bioactive peptides are summarized in Fig. 2.

Discovery of collagen peptides by the in vitro approach

In vitro enzymatic hydrolysis of collagen or gelatin is one of the most common and empirical approach to release bioactive collagen peptides with the aid of different commercial proteases or their combinations, such as Alcalase[®], papain, pepsin, trypsin, chymotrypsin and other food-grade proteases (Fu et al., 2015; Fu and Zhao, 2013; Gómez-Guillén et al., 2011; Zhang et al., 2013). The classic *in vitro* method is primarily initiated by collagen or gelatin extraction from animal by-products, followed by enzymatic hydrolysis to produce collagen hydrolysates. Then, depending on different bioactivities, *in vitro* bioassays are conducted to screen the most active hydrolysates. The subsequent fractionation and purification by a range of chromatography technologies are implemented to screen the most active peptide fractions (Udenigwe and Aluko, 2012). Afterwards, the most active peptide fractions can be further analyzed by LC-MS/MS to identify amino acid sequences, which enables their bioefficacy validation using synthetic peptides. In recent years, *in vitro* classic approach has been successfully employed to isolate and identify novel bioactive collagen peptides (Chen et al., 2017b; Choonpicharn et al., 2016; Guo et al., 2015). Nevertheless, several drawbacks of this approach include the time-consuming procedure (especially, isolation and purification process) and the relatively low yield of the target peptides. In addition, some potent peptides may be excluded during the process of fractionation and purification due to the synergistic or antagonistic effects of other peptides within the whole hydrolysate (Udenigwe, 2014).

In silico approach

In order to overcome the drawbacks and challenges caused by the *in vitro* approach, *in silico* approach has been proposed and used to release bioactive peptide sequences from collagen (Fu et al., 2016b; Lafarga et al., 2014; Yu et al., 2017). Compared to empirical experiments, it is much more time-saving and economical to screen and identify novel bioactive collagen peptides. Several bioactive peptides have been identified with the aid of *in silico* analysis from a variety of protein sources, especially meat or meat by-products (Lafarga et al., 2014), cereal proteins (Udenigwe et

al., 2013) and milk proteins (Nongonierma and FitzGerald, 2014) and potato proteins (Fu et al., 2016a). There are several frequently-used tools of *in silico* analysis, including BIOPEP “enzyme action” (<http://www.uwm.edu.pl/biochemia/index.php/en/biopep>), ExPASy PeptideCutter (http://web.expasy.org/peptide_cutter), MEROPS (<https://www.ebi.ac.uk/merops/>) and MS-Digest (<http://prospector.ucsf.edu/prospector/cgi-bin/msform.cgi?form=msdigest>, Protein Prospector). Typically, this approach is associated with hydrolysis of a protein sequence obtained from a protein database using the known enzymatic cleavage sites. In addition, the QSAR (quantitative structure-activity relationship) model, a regression model that correlates structural characteristics of molecules with their bioactivities, can be employed to predict activities of unidentified collagen peptides. The peptide sequence with the highest predicted potency can be further chemically synthesized in order to validate their actual *in vitro* potencies. Furthermore, identification of collagen peptide structures with certain bioactivity contributes to structure-function relationship studies and use of specific protease or protease cocktail to generate specific hydrolysates (Udenigwe, 2014). Molecular docking, a computational simulation method that can help predict binding sites between receptor (enzyme) and ligand (inhibitor) and assess binding energy based on the scoring system (Pripp and Ardö, 2007), has been applied to better understand specific binding of collagen peptides to enzymes, e.g. Angiotensin I-converting enzyme (ACE) (Fu et al., 2017a; Fu et al., 2016c).

However, variable results between *in vitro* and *in silico* hydrolysis have been observed. Fu et al. (2016b) reported the discordance in ACE-inhibitory peptides from a papain digest of bovine collagen between *in silico* and *in vitro* protein digestions. Similarly, pepsin-catalyzed hydrolysis of potato proteins was found to be different by *in vitro* hydrolysis compared to *in silico* hydrolysis according to certain enzymatic cleavage sites (Rajendran et al., 2016). The differences between *in vitro* and *in silico* predictions may be due to different reasons. *In silico* hydrolysis assumes that all

cleavable peptide bonds would be accessible to the enzyme and be readily hydrolyzed (Udenigwe, 2014). Thus, it is not guaranteed that virtually-generated peptides can be experimentally released, as the cleavage of peptide bonds during enzymatic hydrolysis is dependent on their accessibility as well as the enzyme activity, which is influenced by the external experimental hydrolysis conditions. Also, in terms of the complicated interaction of protease-proteins, the complex spatial structure of collagen might hinder enzyme access to cleavage sites of native proteins, leading to a mismatch between predicted peptides and experimentally released peptides (Fu et al., 2016b; Nongonierma and FitzGerald, 2017). Furthermore, post-translational modifications (PTMs) and amino acid modifications (e.g. high content of Hyp in collagen) of peptides during food storage and processing is not considered during *in silico* hydrolysis (Rajendran et al., 2016). Overall, the *in silico* approach can serve as an alternative approach for exploration of possible bioactive collagen peptides, but the limitations should not be ignored.

Integrated approaches

Given that the limitations of the methods discussed above, the integrated approach may be promising to discover bioactive peptides from collagen. The integrated approach involves screening the optimal enzyme based on *in silico* prediction. The subsequent *in vitro* hydrolysis using the optimal protease is implemented to release collagen peptides. The peptide profile of hydrolysates can be characterized through peptidomics (LC-MS/MS) analysis (Sasaki et al., 2010). Thereafter, the identified sequences can be further predicted for their *in vitro* potential of bioactivities by an *in silico* method. The PeptideRanker server (http://bioware.ucd.ie/~compass/biowareweb/Server_pages/peptideranker.php) can rank peptide sets and assign scores ranging from 0 to 1 based on structure-function relationships (Mooney et al., 2012). The closer the predicted score is to 1, the higher possibility that the peptide is bioactive (Mooney et al., 2012). Eventually, *in vitro* bioactivities of collagen peptides can be further

confirmed by synthetic peptides. However, it should be noted that the relevant knowledge on the structure-function pattern is necessary to predict meaningful information (Udenigwe, 2014), as PeptideRanker cannot predict the particular type or potency of bioactivity of the peptides. More important, experimental hydrolysis is essential to apply the integrated approach (Udenigwe, 2014).

Health beneficial potentials of collagen peptides

In recent years, an increasing number of research findings have indicated a variety of bioactivities for collagen hydrolysates or peptides (Fig. 3), such as antihypertensive, antioxidant and antidiabetic activities in addition to beneficial effects on bone, joint and skin health. Furthermore, the bioactivities of some collagen peptides have been confirmed by *in vivo* animal models and human clinical trials.

Antihypertensive and ACE-inhibitory activity

Hypertension, a serious global health issue, represents one of the major risk factors for developing cardiovascular diseases (Bhat et al., 2015; Organization, 2013). Angiotensin-I converting enzyme (ACE) plays a pivotal physiological role in regulation of blood pressure and electrolyte homeostasis (Perazella and Setaro, 2003). Effective inhibition of ACE has been considered the main target for treatment of hypertension (Udenigwe and Mohan, 2014). However, long-term administration of synthetic ACE inhibitors, such as Captopril, Lisinopril and Enalapril, has been reported to have adverse side effects (Flather et al., 2000). Thus, there has been an increasing interest in development of the natural antihypertensive and ACE-inhibitory peptides as a safer alternative for lowering blood pressure. It has been reported that collagen is a good precursor of ACE-inhibitory peptides (Minkiewicz et al., 2011). A number of identified collagen peptides with ACE-inhibitory activities are mainly derived from fish skins (Lee et al., 2011; Ngo et al., 2015; Ngo et al., 2014), squid gelatin (Alemán et al., 2011), chum salmon gelatin (Lee et al., 2014), Pacific cod gelatin (Himaya et al., 2012), bovine connective tissue (Fu et al., 2016c) and jellyfish collagen (Zhuang et

al., 2010). Some identified collagen peptides with ACE-inhibitory and antihypertensive activities are listed in Table 1.

It is widely documented that the ACE-inhibitory properties of collagen peptides are closely related to their unique amino acid compositions (Aluko, 2015b; Martínez-Alvarez et al., 2015). The ACE-inhibitory activities of collagen peptides can be strongly influenced by the occurrence of hydrophobic (aromatic or branched side chains) amino acid residues, such as Pro, Phe, and Val, at the C-terminal positions (Aluko, 2015a, b). Therefore, the high proportion of Pro residues within the sequence of ACE-inhibitory peptides, to a large extent, contributes to higher potency (Gómez-Guillén et al., 2011; Martínez-Alvarez et al., 2015). In addition, the identified collagen peptides tend to be short and of low-molecular-weight, which potentiates higher activities (Herregods et al., 2011), as small peptides are more efficiently embedded into the active sites of ACE in order to impair its activity (Iwaniak et al., 2014).

Currently, several collagen peptides with ACE-inhibitory activities have been confirmed for their *in vivo* antihypertensive effects. Oral administration of collagen hydrolysates derived from squid skin gelatin, salmon skin gelatin and jellyfish collagen can successfully lower blood pressure in spontaneously hypertensive rats (SHR) (Lee et al., 2014; Lin et al., 2012; Zhuang et al., 2010). Ingestion of a collagen-derived dipeptide Gly-Pro at 500 mg/kg can also induce an antihypertensive effect in SHR after 24 h (Ichimura et al., 2009). Zhuang et al. (2012) evaluated antihypertensive effect of long-term oral administration (100 mg/kg for 30 days) of jellyfish collagen peptides on renovascular hypertension rats (RVH) and found that both systolic blood pressure and diastolic blood pressure of RVH were significantly decreased by 34.8 and 52.6 mmHg. Lee et al. (2014) suggested the antihypertensive effect in SHR of Gly-Leu-Pro (20 mg/kg), an ACE-inhibitory peptide from chum salmon skin gelatin, by decreasing 31 mm Hg systolic blood pressure after 6 h. However, further study on the *in vivo* antihypertensive effects of collagen peptides based on animal

and human clinical trials is still needed for further development of new antihypertensive collagen peptides.

Antioxidant activity

Oxidative stress is an imbalance between excessive production and quenching of reactive oxygen species (ROS). Oxidative stress is implicated in a diversity of diseases, such as aging, hypertension, neurodegenerative diseases, inflammation, gastric ulcers and diabetes mellitus (Fu and Zhao, 2015; Valko et al., 2006). Nowadays, due to safety concerns and consumer preferences, there is an increasing interest in the utilization of natural antioxidants (Sarmadi and Ismail, 2010). In this regard, a wide array of peptides that possess antioxidant activity have been found from various collagenous sources and verified in different oxidative systems (Gómez-Guillén et al., 2011). In recent years, the collagen-derived antioxidant peptides have been identified from blue shark skin gelatin (Weng et al., 2014), Amur sturgeon skin gelatin (Nikoo et al., 2014), grass carp skin (Cai et al., 2015), duck skin gelatin (Lee et al., 2012), croceine croaker scales (Wang et al., 2013), cod skin gelatin (Ngo et al., 2011) and tilapia skin gelatin (Zhang et al., 2012). The identified collagen peptide sequences with antioxidant activities are listed in Table 2.

To date, the exact mechanism responsible for the antioxidant activity of collagen peptide has not been fully elucidated. The antioxidant capacities of collagen peptides have been reported to involve free radical scavenging, singlet oxygen quenching or chelation of metal ion (Kitts and Weiler, 2003). Chen et al. (1998) suggested that the total antioxidant effect of peptides is probably associated with the synergistic effects of the above mechanisms. It is demonstrated that the sequence, composition, peptide conformation, positioning of amino acids, hydrophobicity and molecular weight exert some impacts on antioxidant activities (Alemán and Martínez-Alvarez, 2013; Martínez-Alvarez et al., 2015). The radical-scavenging activity of collagen peptides is attributed to the presence of certain amino acids within their sequences, which can donate protons to

electron-deficient radicals (Samaranayaka and Li-Chan, 2011). The presence of Tyr, Phe, Trp, His, Met and Pro in the peptide sequences has been suggested to enhance chelating and free radical scavenging properties (Banerjee and Shanthi, 2016). Recently, collagen peptides isolated from cod and skate skin gelatin have been demonstrated to enhance cell viability and protect living cells against free radical-mediated oxidative damage (Ngo et al., 2014; Ngo et al., 2011). Two collagen peptides, Thr-Cys-Ser-Pro and Thr-Gly-Gly-Gly-Asn-Val derived from Pacific cod skin can protect H₂O₂-mediated DNA damage and ROS production in RAW 264.7 cells in a dose-dependent manner. At the low peptide concentration (50 µg/mL), DNA damage was attenuated more than 90% and ROS (over 60%) was largely scavenged after 3h (Ngo et al., 2011). Similarly, 10 µM of skate skin gelatin peptides (Met-Val-Gly-Ser-Ala-Pro-Gly-Val-Leu and Leu-Gly-Pro-Leu-Gly-His-Gln) can scavenge intracellular ROS in EA.hy926 cells, evaluated by flow cytometry and light microscope analysis (Ngo, Ryu and Kim 2014). Meanwhile, the mRNA and protein expression levels of antioxidant enzymes (superoxide dismutase and glutathione) were up-regulated with the treatment of Met-Val-Gly-Ser-Ala-Pro-Gly-Val-Leu and Leu-Gly-Pro-Leu-Gly-His-Gln in a dose-dependent manner (10-100 µM) (Ngo, Ryu and Kim 2014). Therefore, the underlying mechanism responsible for antioxidant activities of collagen peptides in cell models may be due to the ability to enhance the expression of antioxidant enzymes in cultured cells by treatment with collagen peptides (Cheung et al., 2015; Fu and Zhao, 2015). In addition, collagen peptides have been reported to retard lipid peroxidation, which is promising for future application in food preservation for the enhanced shelf life and oxidative stability (Sohaib et al., 2016).

Dipeptidyl peptidase-IV (DPP-IV) inhibitory activity

Type 2 diabetes (T2D), a chronic metabolic disorder, is increasing at an alarming rate worldwide. It is characterized by hyperglycemia resulting from inadequate insulin secretion, resistance to insulin action and pancreatic beta-cell dysfunction (Jao et al., 2015; Patil et al., 2015). For T2D

management, one of the latest strategies is associated with inhibition of dipeptidyl peptidase-IV (DPP-IV) activity (Lacroix and Li-Chan, 2016; Nongonierma and FitzGerald, 2016). DPP-IV inhibitors can play a key role in regulation of glucose level through retarding degradation of glucose-dependent insulintropic polypeptide (GIP) and glucagon-like peptide-1 (GLP-1), further leading to insulin secretion (Power et al., 2014). Enzymatic hydrolysis of dietary proteins has been reported to release DPP-IV inhibitory peptides (Nongonierma & FitzGerald, 2015). Among the different sources of proteins, collagen is predicted to be one of the optimal potential precursors of DPP-IV inhibitory peptides based on *in silico* analysis (Lacroix and Li-Chan, 2012). In the past years, a range of collagen-derived DPP-IV inhibitory peptides has been discovered. Table 3 shows a summary of sequences and potencies of DPP-IV inhibitory peptides from different collagenous sources, such as bovine collagen (Hatanaka et al., 2014, Lafarga et al., 2014), Atlantic salmon gelatin (Li-Chan et al., 2012), barbel fish gelatin (Sila et al., 2015), deer collagen (Jin et al., 2015), porcine skin gelatin (Huang et al., 2014) and silver carp collagen (Zhang et al., 2016). Interestingly, a common feature of the identified collagen-derived DPP-IV inhibitory peptides is that most of them share the similar repeating motif, Gly-X-Y sequence.

The structure-activity relationship and the underlying mechanisms responsible for collagen peptide-induced DPP-IV inhibition have not been fully elucidated. Nevertheless, some structural characteristics of DPP-IV inhibitory peptides have been observed in recent studies. Typically, several potent DPP-IV inhibitory peptides contain 2-7 amino acid residues with Pro or Ala as the penultimate amino acid located at the N-terminal (Power et al., 2014). Furthermore, it has been demonstrated that DPP-IV can preferably cleave Pro, Ala, Ser and Hyp in protein sequence as the second N-terminal residue (Davy et al., 2000), so collagen hydrolysates with high proportions of these peptide motifs, such as Xaa-Pro and Xaa-Ala, may have the potential to be DPP-IV inhibitors. This fact is recently confirmed based on *in silico* and *in vitro* analyses (Nongonierma and

FitzGerald, 2013; Nongonierma and FitzGerald, 2014; Wang et al., 2017). In addition to amino acid composition, the peptide sequence can also exert an impact on DPP-IV inhibitory activity. A practical example is that DPP-IV inhibitory dipeptides Ile-Pro and Trp-Val exhibit no inhibitory activity once the peptide sequences are reversed (Hatanaka et al., 2014; Nongonierma and FitzGerald, 2017; Power et al., 2014).

Although *in vitro* DPP-IV inhibitory activity of collagen peptide is widely reported, a limited amount of *in vivo* studies using diabetic animals or humans have been performed to evaluate their bioefficacy. Zhu et al. (2010) claimed that supplement of marine collagen hydrolysates (13 g/day for 3 months) in Chinese patients with T2D may benefit glucose and lipid metabolism, insulin sensitivity and renal function in Chinese patients with T2D. Recently, Hsieh et al. (2015) reported that oral administration of Atlantic salmon skin gelatin hydrolysate at the dose of 300 mg/day for 5 weeks could inhibit GLP-1 degradation by DPP-IV, leading to the enhanced insulin secretion and improved glycemic control in streptozotocin-induced diabetic rats. Similarly, oral administration of porcine skin gelatin hydrolysate (300 mg per day for 42 days) can also improve the glucose tolerance in streptozotocin-induced diabetic rats through inhibition of plasma DPP-IV activity by 63.4% (Huang et al., 2014). However, the molecular mechanisms responsible for DPP-IV inhibition by collagen peptides remains to be unraveled. Besides, clinical trials are needed to validate and confirm the efficacy and bioavailability of collagen peptides.

Bone and joint health

With the increasingly aging population, osteoporosis and osteoarthritis have become major age-related musculoskeletal diseases (Dequeker et al., 2003). Osteoporosis is a systemic skeletal disease characterized by low bone mass, deterioration of skeletal microarchitectural and increased risk of bone fragility and fracture (Inderjeeth and Poland, 2010). Likewise, osteoarthritis is the most prevalent type of joint disease with chronic musculoskeletal pain and disability mainly due to the

imbalance between synthesis and degradation of the articular cartilage (Bello and Oesser, 2006).

Over the years, a large and growing body of evidence indicates that collagen hydrolysates/peptides play a positive role in management of osteoporosis, joint disorders and osteoarthritis (Bello and Oesser, 2006; Daneault et al., 2017).

A number of *in vitro* studies have confirmed that collagen peptides can stimulate proliferation and differentiation of osteoblasts and improve calcium absorption (Fu and Zhao, 2013; Guillerminet et al., 2012). Fu and Zhao (2013) demonstrated that salmon skin gelatin hydrolysates (0.1 mg/mL for 48h) were capable of inducing osteoblast proliferation, accelerating cell cycle progression and inhibiting cell apoptosis (from 6.9% to 4.5%) in human hFOB1.19 cells. In parallel, collagen hydrolysates (0.1-6.0 mg/mL for 7 days) have been confirmed to stimulate osteoblast proliferation and differentiation (Daneault et al., 2017; Kim et al., 2014; Liu et al., 2014). Moreover, it has been shown that collagen peptides (0.1 mg/mL for 7 days) can significantly up-regulated expression of the bone-related genes in osteoblasts, including type I collagen, alkaline phosphatase, osteocalcin and osteopontin (Han et al., 2009; Kim et al., 2014; Liu et al., 2014; Yamada et al., 2013). However, the related mechanisms underlying the beneficial actions on osteoblasts are not completely clarified. Currently, there are potential reasons that help shed lights upon underlying beneficial effects. Firstly, collagen peptides can activate some related receptors on the cell membrane and exert an anabolic effect on bone cells (Daneault et al., 2017). It has been documented that collagen hydrolysates can induce the expression of bone-related genes in osteoblasts (Daneault et al., 2017). Furthermore, due to its richness in Pro and Hyp, collagen hydrolysate may exhibit the beneficial effects on osteoblasts through inducing insulin-like growth factor 1 (IGF1) secretion and activating the calcium sensing receptor (Conigrave et al., 2008; Daneault et al., 2017; Dawson-Hughes et al., 2007). In addition, it is claimed that the source and

molecular weight of collagen hydrolysates may be also key factors for mediating beneficial actions on bone metabolism (Daneault et al., 2017).

Daneault et al. (2017) recently summarized some evidence from different animal models based on bone growth, bone loss and bone healing models and implied that collagen hydrolysates/peptides can help maintain a balanced bone turnover in different physiological conditions (growth, bone loss and healing) via promoting bone formation and development. To date, a limited number of clinical studies have been implemented to evaluate beneficial effects of collagen hydrolysates on bone metabolism. Supplement of calcium-collagen chelate has been shown to improve bone mass and prevent excessive bone loss and turnover (Elam et al., 2015; Hooshmand et al., 2013). A 4-month randomized double-blind trial executed by Martin-Bautista and colleagues revealed that the daily intake of collagen hydrolysates (250 mL hydrolyzed gelatin products) can exert a positive effect on bone remodeling by significantly increasing the levels of IGF1 and bone alkaline phosphatase (Martin-Bautista et al., 2011). Nevertheless, future standardized clinical trials remain to be conducted to strengthen this scientific evidence.

Scientific evidence suggests that collagen hydrolysates exert a positive therapeutic effect on osteoarthritis (Bello and Cesser, 2006; Kumar et al., 2015). *In vitro* cell studies have revealed that collagen hydrolysates/peptides stimulate synthesis of proteoglycans and collagen in human chondrocytes (Cesser and Seifert, 2003; Porfírio and Fanaro, 2016; Schadow et al., 2017). Meanwhile, the clinical studies have demonstrated that intake of collagen peptides (5 g/day for 12 weeks) can lead to the relief of joint pain (Bruyere et al., 2012; Zdzieblik et al., 2017) and stimulate regeneration of type II collagen and the biosynthesis of proteoglycans in cartilage tissue osteoarthritis patients (McAlindon et al., 2011). Trč and Bohmová (2011) claimed that ingestion of collagen hydrolysate (1.5g/day) for 24 weeks resulted in improvement of joint conditions. These

findings provide a theoretical basis for exploitation of collagen hydrolysate as a regenerative agent for osteoarthritis patients.

Skin health

Skin characteristics are subject to deterioration by chronological aging, dermatological disorders and environmental conditions, which can be exacerbated by photo-aging or other undesirable lifestyle issues (Sibilla et al., 2015; Zague, 2008; Zague et al., 2011). An overview of the beneficial impacts of collagen peptides on the skin by Sibilla et al. (2015) indicates that collagen peptides can stimulate proliferation and motility of fibroblasts (Chai et al., 2010), increase the density and diameter of collagen fibers (Matsuda et al., 2006), elevate production of hyaluronic acid (Matsuda et al., 2006), promote the expression of type I and IV collagen and protect against photoaging (Sibilla et al., 2015; Sun et al., 2013). A number of controlled clinical trials have evidenced the bioefficacy and benefits of collagen peptides on skin properties, including hydration (Choi et al., 2013), skin elasticity (Proksch et al., 2014) and reduction of wrinkles (Schwartz and Park, 2012).

Other potential health benefits

Apart from the above-mentioned bioactivities, collagen peptides have been reported to exhibit some other physiological activities, including renin-inhibitory activity, metal chelating capacity, immunostimulatory activity, lipid-lowering activity, neuroprotective effect and wound-healing activity (Pal and Suresh, 2016). In addition to ACE inhibition, several recent investigations have revealed the inhibitory activity of renin by collagen peptides (Fu et al., 2017b; Lafarga et al., 2014), which is an emerging target to better lower the blood pressure (Udenigwe and Mohan, 2014).

Collagen peptides have been demonstrated to display the excellent metal chelating capacity, which can improve the absorption of mineral ions in the human body, including calcium (Guo et al., 2015), copper (Zhuang et al., 2009) and ferrous (Nakchum and Kim, 2016) ions. Zhang et al. (2011) and Wang et al. (2015a) reported that oral administration of chum salmon collagen peptides could

accelerate wound healing. Niu et al. (2016) indicated that cod skin collagen peptides exerted a protective action on the acetic acid-induced gastric ulcer in rats. It has been reported that collagen hydrolysates show immunostimulatory activity in RAW 264.7 macrophage cells (Sae-leaw et al., 2016), Caco-2 cells (Chen et al., 2017a) and C57BL/6.KOR-ApoE^{shl} mice (Zhang et al., 2010). In addition, the beneficial effects of on lipid metabolism have been reported. Oral administration of collagen hydrolysates (0.4 g for 2h) can significantly lower the levels of total lipids and triglycerides in plasma (Saito et al., 2009). Minaguchi et al. (2012) claimed that Pro-Hyp (200 ng/mL), collagen-derived dipeptide significantly reduced the size of lipid droplets in mouse 3T3-L1 adipocytes after 14 days. Besides, the neuroprotective effect of collagen peptides (3.0 g/kg for 90 days) was reported to benefit learning and memory in the rat brain via attenuating oxidative damage and acetylcholinesterase activity in the brain (Pei et al., 2010; Xu et al., 2015). According to Lima et al. (2015), collagen hydrolysates (5 mg/mL) exhibited antibacterial activity against Gram-positive bacteria and Gram-negative bacteria. The low-molecular-weight of collagen peptides may facilitate interactions with bacterial membranes due to better structure acquisition of amino acid residues (Lima et al., 2015). It has been documented that Pro-rich peptides are potential antimicrobial peptides (Reddy et al., 2004), which is compatible with collagen structure with abundant Pro in its repeating Gly-X-Y motif. The present interest in the collagen peptides can guide exploration of new bioactivities in future research activities.

Unique advantages of collagen peptides over other therapeutic peptides

Bioavailability of collagen peptides and underlying mechanisms

A growing amount of recent studies have indicated that collagen peptides, especially those with C-terminal Pro or Hyp residue, can be transported across the intestinal epithelial monolayer, enter the bloodstream and exhibit their bioactive properties (Alemán and Martínez-Alvarez, 2013). An increasing number of collagen-derived peptides has been identified in the blood after oral ingestion

of collagen/gelatin hydrolysates, among which Pro-Hyp and Hyp-Gly account for a major percentage (Ohara et al., 2007; Shigemura et al., 2011; Shigemura et al., 2014). The presence of collagen-derived oligopeptides with higher concentrations in the blood suggests a high resistance to digestion by the enzymes in plasma and gastrointestinal tract and blood protease (Taga et al., 2014, 2016; Wang et al., 2015b). The unique amino acid composition and structure confer collagen peptides with excellent stability and less protease cleavage sites. The high Pro level exists in collagen peptides (10-30% abundance) may increase their stability towards digestive enzymes and intestinal peptidases in comparison to other therapeutic peptides (Banerjee and Shanthi, 2016). This is because the presence of a Pro residue adjacent to the cleavage site prevents proteases from cleaving the normally susceptible peptide bond (Liu et al., 2015). Examples of detection of collagen peptides in the plasma of animals and humans after the ingestion of collagen/gelatin hydrolysates are listed in Table 4.

In vitro, animal experiments and human clinical trials have highlighted the beneficial effects of collagen peptides, especially Hyp-containing peptides (Matsuda et al., 2006; Ngo et al., 2015; Tsuruoka et al., 2007; Wu et al., 2004). Collagen peptides can stimulate the growth of fibroblasts in the skin as well as the synthesis of hyaluronic acid (Ohara et al., 2010; Shigemura et al., 2009) and improve skin barrier dysfunction (Shimizu et al., 2015). Pro-Hyp and Pro-Hyp-Gly exhibit chemotactic activities in human fibroblast, peripheral blood neutrophils and monocytes (Zague, 2008). Furthermore, oral intake of collagen peptides (2.5 g/day for 8 weeks) with beneficial effects on dermal matrix synthesis have been confirmed by human clinical trials (Proksch et al., 2014) and animal experiments (Zague et al., 2011). The reported evidence has indicated that collagen peptides, as a biological messenger, can trigger the synthesis of new collagen and modulate fibroblast cells reorganization in the extracellular matrix (Wang et al., 2015b). Iwai et al. (2009) examined oral ingestion of chicken collagen hydrolysates and detected a range of Hyp-containing collagen

peptides in human blood with ACE-inhibitory activity, among which Ala-Hyp had the strongest ACE-inhibitory activity ($IC_{50}=177\text{ }\mu\text{M}$). Several collagen peptides identified in the plasma, such as Gly-Pro-Y and X-Hyp-Gly types, may have significant antidiabetic effects *in vivo* (Taga et al., 2016).

The transepithelial transport mechanisms of collagen peptides have been extensively investigated. Aito-Inoue et al. (2007) investigated the mechanism for the transcellular transport of Gly-Pro-Hyp based on porcine brush border membrane vesicles. Gly-Pro-Hyp was further degraded into Gly and Pro-Hyp in the apical side, while the intact Gly-Pro-Hyp was not identified on the basolateral side, leading to failure of transport across the intestinal epithelial apical membrane. By contrast, two collagen peptides (Val-Gly-Pro-Val and Gly-Pro-Arg-Gly-Phe) have been reported to show stability against epithelial cell peptidases and transported across Caco-2 cell monolayers via paracellular pathway (Fu et al., 2016c). Similarly, low-molecular-weight (<2 kDa) collagen hydrolysates can be efficiently transported across the Caco-2 cell monolayer (Feng and Betti, 2017). The difference might be due to collagen type and source as well as the variations in substrate specificities peptidases of animal species on the intestinal epithelial apical membrane (Liu et al., 2015). In addition, Watanabe-Kamiyama et al. (2010) investigated ingestion of low-molecular-weight (approximately 800 Da) collagen hydrolysates containing a high amount of Gly-Pro-Hyp (288 mg for 0.5-6h) and reported that Gly-Pro-Hyp could be absorbed into the bloodstream through the intestinal brush border membrane in an intact form using Wistar rats as the animal model. These above-mentioned results may provide some evidence for the excellent bioavailability of collagen peptides.

Sustainable sources and biocompatibility of collagen peptides

The utilization of inexpensive protein sources, especially animal by-products as the raw materials to prepare bioactive peptides cannot only reduce production costs but also by-products. Collagen is

one of the most abundant proteins in the world that is widely available and inexpensive to recover from animal by-products (Banerjee and Shanthi, 2016). Currently, utilization of collagen as high value-added ingredients via enzyme technology has been the top trend in the meat industry, leading to a high benefit-to-cost ratio and deriving value from slaughterhouse by-products (Lafarga and Hayes, 2014). The resultant collagen peptides have an enormous commercial potential as food ingredients or nutraceuticals. In addition, collagen peptides are biocompatible and safe due to their unique biological and structural characteristics. Regardless of different collagen types, they all share the nearly identical sequence and structure, which guarantee weak *in vivo* immunogenicity (Banerjee and Shanthi, 2016; Lee et al., 2001).

Neutral taste of collagen peptides

Peptides have a wide array of tastes, including sweet, bitter, umami, sour or salty taste (Temussi, 2012). However, bitter peptides are frequently released during enzymatic protein hydrolysis (Li-Chan, 2015), which influences consumer acceptability and impedes the further development of peptide-based food products (Udenigwe, 2014). Humans can distinguish the bitter taste of peptides mainly through taste receptor cells of taste buds on the surface of the tongue (Maehashi and Huang, 2009). The bitter taste is suggested to be mediated by taste receptor type 2 (T2Rs), a large family of G-protein-coupled receptors. The interaction of bitter peptides with T2Rs through binding and stimulating units can trigger the signaling cascades (Lafarga and Hayes, 2016; Maehashi and Huang, 2009).

Bitter peptides have been widely identified in various protein sources, including milk protein (Singh et al., 2005), soy protein (Cho et al., 2004), wheat gluten protein (Liu et al., 2016) and bovine hemoglobin (Aubes-Dufau et al., 1995). To our knowledge, however, it has not been reported that collagen hydrolysates possess bitter tastes. Moreover, hydrolyzed collagen contains a high content of glycine, which also contributes to neutral taste. During the industrial production of collagen

hydrolysates, release of bitter peptides is low when compared to the amount formed from other hydrolyzed proteins (Schrieber and Gareis, 2007). Several approaches have been employed to predict the peptide bitterness. The 'Q rule', which was first proposed by Ney (Ney, 1979), can help estimate the bitterness of peptides based on amino acid composition, where Q value is calculated from the average hydrophobicity of peptide sequences. According to the Ney's Q rule, the peptides (under 6 kDa) with Q values more than 1400 cal/mol are estimated to be bitter. However, the average hydrophobicity (Q) of collagen is 1280 cal/mol (Ney, 1979). This value is much lower than casein (1605 cal/mol) and soy protein (1540 cal/mol). The low Q value implies that collagen hydrolysates/peptides tend to be less bitter and with neutral taste, which promotes its potential as a functional ingredient in the food industry.

The main hurdles for commercialization of collagen peptides

The past decade has witnessed an increasing number of studies on bioactive collagen peptides, which contributes to development of functional foods with health-promoting functions to combat the growing global burden of chronic diseases. Even with the unique advantages of collagen peptides as compared to other therapeutic peptides, conversion to functional foods has confronted several challenges, such as inadequate evidence of clinical bioefficacy, industrial-scale production as well as food processing and storage.

Inadequate clinical trials

Even though collagen peptides with various bioactivities have been confirmed on the basis of *in vitro* or animal models, adequate human clinical studies are scanty to substantiate bioefficacy and support future health claims. A number of nutritional intervention studies on the detection of Hyp-containing peptides in human plasma or tissues after ingestion of collagen or gelatin hydrolysates have been reported (Shigemura et al., 2014; Sugihara et al., 2012; Taga et al., 2014; Zague, 2008). Given that the bioavailability of bioactive peptides in humans is relatively very low (Foltz et al.,

2010; Nongonierma and FitzGerald, 2017), the amount of the collagen peptides that should be ingested to exhibit an *in vivo* effect remains to be explored. Thus, utilization of dose-response approaches in humans may allow assessment of their *in vivo* potency. Recently, Shigemura et al. (2014) reported ingestion of three different doses (30.8, 153.8 and 384.6 mg/kg for 15-360 min) of cod skin gelatin hydrolysates by humans and demonstrated a dose-response pattern between the amount of hydrolysate ingested and the level of Hyp and Hyp-containing peptides in plasma. There was no absorption limit of Hyp and Hyp-containing peptide, suggesting that higher doses would lead to elevated plasma level. In contrast, a well-known health claim on milk-derived antihypertensive tripeptides (Ile-Pro-Pro and Val-Pro-Pro) was rejected by the European Food Safety Authority (EFSA), suggesting lack of cause and effect relationship for consumption of Ile-Pro-Pro and Val-Pro-Pro at the proposed dosage (EFSA Panel on Dietetic Products and Allergies, 2011). Although several clinical studies of collagen peptides have been executed to demonstrate efficacy and benefits on skin properties (Sibilla et al., 2015), more clinical trials are still needed to confirm other *in vivo* bioactive properties of collagen peptides.

Industrial scale production

Novel strategies, such as *in silico* and the integrated approaches can predict and optimize generation of bioactive peptides with targeted activities. However, a major challenge is industrial-scale production of pure collagen peptides. Chemical synthesis of peptides involves the consumption of large amounts of solvents and leads to the generation of waste, which is environmentally unfriendly. Recombinant expression of peptides may serve as an alternative way for the production of longer peptides (more than 25 amino acids in length) (Lafarga and Hayes, 2016; Rodríguez et al., 2014). Unfortunately, this method is not suitable for the generation of smaller peptides with short amino acid sequences. Therefore, industrial-scale production of crude protein hydrolysates containing bioactive peptide sequences is a more practically feasible strategy, given that the peptides retain

their bioactivities within the hydrolysates (Agyei and Danquah, 2011; Udenigwe, 2014). From a viewpoint of industrial-scale production, the use of immobilized enzymes and membrane separation techniques (e.g. ultrafiltration) can offer several benefits, including higher efficiency and lower operating costs for the large-scale production. Integration of complementary processes to establish economical and efficient industrial-scale processes for fractionation and continuous production of bioactive peptides has been proposed (Wu et al., 2013). For example, electrodialysis with ultrafiltration membrane technology based on the separation of molecules according to the charge and molecular mass may be compatible with large-scale production of peptides (Doyen et al., 2012). Therefore, the exploration of novel methodologies to generate bioactive collagen peptides in an inexpensive and efficient way is imperative.

Food processing and storage

The stability of collagen peptides during food processing and storage is crucial for their application as functional ingredients, as peptides are prone to suffer from chemical modifications of the backbone or side chains. These chemical reactions include disulfide bond formation, dehydration, glycation and aromatic ring oxidation (Udenigwe and Fogliano, 2017), giving rise to changes in structure and bioactivity of peptides (López-Fandiño et al., 2006).

During food processing and storage, interactions between peptides and food matrix may bring about some physicochemical reactions, such as hydrophobic interactions, disulfide interactions and Maillard reaction (Rao et al., 2016). Moreover, peptide aggregates formed by covalent bond can be widely detected in different powder-based protein hydrolysates (Rao et al., 2016), leading to impaired quality of products. Maillard reaction has been frequently observed during the storage and thermal food processing in the presence of reducing sugar (Jiang et al., 2013; Rao et al., 2016).

These processes can exert some deleterious impacts on the sensory profile, solubility, color and bioactivity of the peptide-based functional food products (Udenigwe, 2014). However, it is worth

noting that the altered bioactivities of peptide-based Maillard reaction products can be partly attributed to the formation of new compounds (Udenigwe & Fogliano, 2017). Therefore, the unwanted Maillard reaction during processing and storage of peptide-rich products could be minimized through application of sugar substitute (e.g. sugar alcohol). Recently, collagen peptides have been demonstrated to retain potent ACE-inhibitory activity after exposure to different processing temperatures (20-100 °C) and pH values (2-10) (Fu et al., 2015). Unfortunately, the information on the food matrix-collagen peptides interaction and its impact on availability is extremely limited.

In addition, oxidation of amino acids via endogenous production of oxidative compounds and exogenous oxidizing agents during processing and storage may also lead to the changes in molecular structure and bioactivity (López-Fandiño et al., 2006). For instance, Met is easily oxidized to methionine sulfone and homocysteic acid in the presence of peroxides, which makes the oxidized products biologically unavailable (López-Fandiño et al., 2006). Therefore, the appropriate packaging materials and conditions should be used to prevent oxidative damage in peptide-based products.

Conclusion

Over the years, there has been an increasing interest in exploration of collagen peptides as bioactive ingredients. This review indicates that collagen from animal by-products can serve as an excellent precursor to release bioactive peptides. Collagen peptides with various bioactivities and health benefits have a promising potential as functional food ingredients. The integration of interdisciplinary approaches may result in improved generation and identification of novel bioactive collagen peptides. The sustainable sources of collagen, good bioavailability and neutral taste render collagen peptides with exclusive health benefits over other therapeutic peptides. Further advances in effective large-scale production and recovery processes are crucial, and the standardized clinical

trials need to be executed to evaluate and validate the bioefficacy of collagen peptides based on human intervention studies. Finally, to fully explore the benefits of collagen peptides, their interactions with the food matrix during food storage and processing need to be considered as part of holistic approach for functional food development.

Funding

This work was supported by Danish Ministry of Food, Agriculture and Fisheries under GUDP grant (No. 34009-16-1085).

References

- Agyei, D., and Danquah, M. K. (2011). Industrial-scale manufacturing of pharmaceutical-grade bioactive peptides. *Biotechnol Adv.* **29**: 272-277.
- Aito-Inoue, M., Lackeyram, D., Fan, M. Z., Sato, K., and Mine, Y. (2007). Transport of a tripeptide, Gly-Pro-Hyp, across the porcine intestinal brush - border membrane. *J Peptide Sci.* **13**: 468-474.
- Alao, B. O., Falowo, A. B., Chulayo, A., and Muchenje, V. (2017). The Potential of Animal By-Products in Food Systems: Production, Prospects and Challenges. *Sustainability.* **9**: 1089.
- Alemán, A., Giménez, B., Pérez-Santin, E., Gómez-Guillén, M. C., and Montero, P. (2011). Contribution of Leu and Hyp residues to antioxidant and ACE-inhibitory activities of peptide sequences isolated from squid gelatin hydrolysate. *Food Chem.* **125**: 334-341.
- Alemán, A., and Martínez-Alvarez, O. (2013). Marine collagen as a source of bioactive molecules: A review. *Nat Prod J.* **3**: 105-114.
- Aluko, R. E. (2015a). Antihypertensive peptides from food proteins. *Annu Rev Food Sci Technol.* **6**: 235-262.
- Aluko, R. E. (2015b). Structure and function of plant protein-derived antihypertensive peptides. *Curr Opin Food Sci.* **4**: 44-50.

- Aubes-Dufau, I., Seris, J.-L., and Combes, D. (1995). Production of peptic hemoglobin hydrolyzates: Bitterness demonstration and characterization. *J Agric Food Chem.* **43**: 1982-1988.
- Banerjee, P., and Shanthi, C. (2016). Cryptic Peptides from Collagen: A critical review. *Protein Peptide Lett.* **23**: 664-672.
- Bello, A. E., and Oesser, S. (2006). Collagen hydrolysate for the treatment of osteoarthritis and other joint disorders: a review of the literature. *Curr Med Res Opin.* **22**: 2221-2232.
- Bhat, Z. F., Kumar, S., and Bhat, H. F. (2015). Bioactive peptides of animal origin: a review. *J Food Sci Technol.* **52**: 5377-5392.
- Bruyere, O., Zegels, B., Leonori, L., Rabenda, V., Janssen, A., Bourges, C., and Reginster, J. Y. (2012). Effect of collagen hydrolysate in articular pain: a 6-month randomized, double-blind, placebo controlled study. *Complement Ther Med.* **20**: 124-130.
- Cai, L., Wu, X., Zhang, Y., Li, X., Ma, S., and Li, J. (2015). Purification and characterization of three antioxidant peptides from protein hydrolysate of grass carp (*Ctenopharyngodon idella*) skin. *J Funct Foods.* **16**: 234-242.
- Chai, H.-J., Li, J.-H., Huang, H.-N., Li, T.-L., Chan, Y.-L., Shiau, C.-Y., and Wu, C.-J. (2010). Effects of sizes and conformations of fish-scale collagen peptides on facial skin qualities and transdermal penetration efficiency. *J BioMed Res.* **2010**: 757301.
- Chen, H.-M., Muramoto, K., Yamauchi, F., Fujimoto, K., and Nokiara, K. (1998). Antioxidative properties of histidine-containing peptides designed from peptide fragments found in the digests of a soybean protein. *J Agric Food Chem.* **46**: 49-53.
- Chen, Q., Chen, O., Martins, I. M., Hou, H., Zhao, X., Blumberg, J. B., and Li, B. (2017a). Collagen peptides ameliorate intestinal epithelial barrier dysfunction in immunostimulatory Caco-2 cell monolayers via enhancing tight junctions. *Food Funct.* **8**: 1144-1151.

- Chen, Q., Guo, L., Du, F., Chen, T., Hou, H., and Li, B. (2017b). The chelating peptide (GPAGPHGPPG) derived from Alaska pollock skin enhances calcium, zinc and iron transport in Caco-2 cells. *Int Food Sci Tech.* **52**: 1283-1290.
- Cheung, R. C., Ng, T. B., and Wong, J. H. (2015). Marine Peptides: Bioactivities and Applications. *Mar Drugs.* **13**: 4006-4043.
- Cho, M. J., Unklesbay, N., Hsieh, F.-H., and Clarke, A. D. (2004). Hydrophobicity of bitter peptides from soy protein hydrolysates. *J Agric Food Chem.* **52**: 5895-5901.
- Choi, J. W., Kwon, S. H., Huh, C. H., Park, K. C., and Youn, S. W. (2013). The influences of skin visco-elasticity, hydration level and aging on the formation of wrinkles: a comprehensive and objective approach. *Skin Res Technol.* **19**: e349-e355.
- Choonpicharn, S., Tateing, S., Jaturasitha, S., Rakariyatham, N., Suree, N., and Niamsup, H. (2016). Identification of bioactive peptide from *Oreochromis niloticus* skin gelatin. *J Food Sci Technol.* **53**: 1222-1229.
- Conigrave, A., Brown, E., and Rizzoli, R. (2008). Dietary protein and bone health: roles of amino acid-sensing receptors in the control of calcium metabolism and bone homeostasis. *Annu. Rev. Nutr.* **28**: 131-155.
- Daneault, A., Prawitt, J., Fabien Soule, V., Coxam, V., and Wittrant, Y. (2017). Biological effect of hydrolyzed collagen on bone metabolism. *Crit Rev Food Sci Nutr.* **57**: 1922-1937.
- Davy, A., Thomsen, K. K., Juliano, M. A., Alves, L. C., Svendsen, I., and Simpson, D. J. (2000). Purification and characterization of barley dipeptidyl peptidase IV. *Plant Physiol.* **122**: 425-432.
- Dawson-Hughes, B., Harris, S., Rasmussen, H., and Dallal, G. (2007). Comparative effects of oral aromatic and branched-chain amino acids on urine calcium excretion in humans. *Osteoporosis Int.* **18**: 955-961.

- Dequeker, J., Aerssens, J., and Luyten, F. P. (2003). Osteoarthritis and osteoporosis: clinical and research evidence of inverse relationship. *Aging Clin Exp Res.* **15**: 426-439.
- Doyen, A., Saucier, L., Beaulieu, L., Pouliot, Y., and Bazinet, L. (2012). Electro separation of an antibacterial peptide fraction from snow crab by-products hydrolysate by electrodialysis with ultrafiltration membranes. *Food Chem.* **132**: 1177-1184.
- EFSA Panel on Dietetic Products, Nutrition and Allergies (2011). Scientific Opinion on the substantiation of a health claim related to isoleucyl-prolyl-proline (IPP) and valyl-prolyl-proline (VPP) and maintenance of normal blood pressure pursuant to Article 13(5) of Regulation (EC) No 1924/2006. *EFSA J.* **9**: 1-18.
- Elam, M. L., Johnson, S. A., Hooshmand, S., Feresin, R. G., Payton, M. E., Gu, J., and Arjmandi, B. H. (2015). A calcium-collagen chelate dietary supplement attenuates bone loss in postmenopausal women with osteopenia: a randomized controlled trial. *J Med Food.* **18**: 324-331.
- Feng, M., and Betti, M. (2017). Transepithelial transport efficiency of bovine collagen hydrolysates in a human Caco-2 cell line model. *Food Chem.* **224**: 242-250.
- Ferraro, V., Anton, M., and Santé-Lhoutellier, V. (2016). The “sisters” α -helices of collagen, elastin and keratin recovered from animal by-products: Functionality, bioactivity and trends of application. *Trends Food Sci Technol.* **51**: 65-75.
- Flather, M. D., Yusuf, S., Køber, L., Pfeffer, M., Hall, A., Murray, G., Torp-Pedersen, C., Ball, S., Pogue, J., and Moyé, L. (2000). Long-term ACE-inhibitor therapy in patients with heart failure or left-ventricular dysfunction: a systematic overview of data from individual patients. *Lancet.* **355**: 1575-1581.
- Foltz, M., van der Pijl, P. C., and Duchateau, G. S. (2010). Current in vitro testing of bioactive peptides is not valuable. *J Nutr.* **140**: 117-118.

- Fu, Y., Alashi, A. M., Young, J. F., Therkildsen, M., and Aluko, R. E. (2017a). Enzyme inhibition kinetics and molecular interactions of patatin peptides with angiotensin I-converting enzyme and renin. *Int J Biol Macromol.* **101**: 207-213.
- Fu, Y., Young, J. F., and Therkildsen, M. (2017b). Bioactive peptides in beef: Endogenous generation through postmortem aging. *Meat Sci.* **123**: 134-142.
- Fu, Y., Wu, W., Zhu, M., and Xiao, Z. (2016a). In silico assessment of the potential of patatin as a precursor of bioactive peptides. *J Food Biochem.* **40**: 366-370.
- Fu, Y., Young, J. F., Løkke, M. M., Lametsch, R., Aluko, R. E., and Therkildsen, M. (2016b). Revalorisation of bovine collagen as a potential precursor of angiotensin I-converting enzyme (ACE) inhibitory peptides based on in silico and in vitro protein digestions. *J Funct Foods.* **24**: 196-206.
- Fu, Y., Young, J. F., Rasmussen, M. K., Dalsgaard, T. K., Lametsch, R., Aluko, R. E., and Therkildsen, M. (2016c). Angiotensin I-converting enzyme-inhibitory peptides from bovine collagen: insights into inhibitory mechanism and transepithelial transport. *Food Res Int.* **89**: 373-381.
- Fu, Y., Young, J. F., Dalsgaard, T. K., and Therkildsen, M. (2015). Separation of angiotensin I - converting enzyme inhibitory peptides from bovine connective tissue and their stability towards temperature, pH and digestive enzymes. *Int J Food Sci Tech.* **50**: 1234-1243.
- Fu, Y., and Zhao, X.-H. (2015). Utilization of chum salmon (*Oncorhynchus keta*) skin gelatin hydrolysates to attenuate hydrogen peroxide-induced oxidative injury in rat hepatocyte BRL cell model. *J Aquat Food Prod T.* **24**: 648-660.
- Fu, Y., and Zhao, X.-H. (2013). In vitro responses of hFOB1. 19 cells towards chum salmon (*Oncorhynchus keta*) skin gelatin hydrolysates in cell proliferation, cycle progression and apoptosis. *J Funct Foods.* **5**: 279-288.

- Gómez-Guillén, M. C., Giménez, B., López-Caballero, M. E., and Montero, M. P. (2011). Functional and bioactive properties of collagen and gelatin from alternative sources: A review. *Food Hydrocolloid*. **25**: 1813-1827.
- Guillerminet, F., Fabien-Soulé, V., Even, P., Tomé, D., Benhamou, C.-L., Roux, C., and Blais, A. (2012). Hydrolyzed collagen improves bone status and prevents bone loss in ovariectomized C3H/HeN mice. *Osteoporosis Int*. **23**: 1909-1919.
- Guo, L., Harnedy, P. A., O'Keeffe, M. B., Zhang, L., Li, B., Hou, H., and FitzGerald, R. J. (2015). Fractionation and identification of Alaska pollock skin collagen-derived mineral chelating peptides. *Food Chem*. **173**: 536-542.
- Han, X., Xu, Y., Wang, J., Pei, X., Yang, R., Li, N., and Li, Y. (2009). Effects of cod bone gelatin on bone metabolism and bone microarchitecture in ovariectomized rats. *Bone*. **44**: 942-947.
- Hatanaka, T., Kawakami, K., and Uraji, M. (2014). Inhibitory effect of collagen-derived tripeptides on dipeptidylpeptidase-IV activity. *J Enzyme Inhib Med Chem*. **29**: 823-828.
- Hayes, M. (2011). Marine bioactive compounds: Sources, characterization and applications. Springer Science & Business Media, New York.
- Henchion, M., McCarthy, M., and O'Callaghan, J. (2016). Transforming beef by-products into valuable ingredients: Which spell/recipe to use? *Front Nutr*. **3**: 53.
- Herregods, G., Van Camp, J., Morel, N., Ghesquiere, B., Gevaert, K., Vercruysse, L., Dierckx, S., Quanten, E., and Smagghe, G. (2011). Angiotensin I-converting enzyme inhibitory activity of gelatin hydrolysates and identification of bioactive peptides. *J Agric Food Chem*. **59**: 552-558.
- Himaya, S. W., Ryu, B., Ngo, D. H., and Kim, S. K. (2012). Peptide isolated from Japanese flounder skin gelatin protects against cellular oxidative damage. *J Agric Food Chem*. **60**: 9112-9119.

- Hooshmand, S., Elam, M., Browne, J., Campbell, S., and Payton, M. (2013). Evidence for bone reversal properties of a calcium-collagen chelate, a novel dietary supplement. *J Food Nutr Disor.* **1**: 2.
- Hsieh, C. H., Wang, T. Y., Hung, C. C., Chen, M. C., and Hsu, K. C. (2015). Improvement of glycemic control in streptozotocin-induced diabetic rats by Atlantic salmon skin gelatin hydrolysate as the dipeptidyl-peptidase IV inhibitor. *Food Funct.* **6**: 1887-1892.
- Huang, S.-L., Hung, C.-C., Jao, C.-L., Tung, Y.-S., and Hsu, K.-C. (2014). Porcine skin gelatin hydrolysate as a dipeptidyl peptidase IV inhibitor improves glycemic control in streptozotocin-induced diabetic rats. *J Funct Foods.* **11**: 235-242.
- Ichimura, T., Yamanaka, A., Otsuka, T., Yamashita, E., and Maruyama, S. (2009). Antihypertensive effect of enzymatic hydrolysate of collagen and Gly-Pro in spontaneously hypertensive rats. *Biosci Biotechnol Biochem.* **73**: 2317-2319.
- Inderjeeth, C. A., and Poland, K. E. (2010). Management of osteoporosis in older people. *J Pharm Pract Res.* **40**: 229-234.
- Iwai, K., Zhang, Y., Kouguchi, T., Saiga-Egusa, A., Shimizu, M., Ohmori, T., Takahata, Y., and Morimatsu, F. (2009). Blood concentration of food-derived peptides following oral intake of chicken collagen hydrolysate and its angiotensin-converting enzyme inhibitory activity in healthy volunteers. *J Jpn Soc Food Sci.* **56**: 326-330.
- Iwaniak, A., Minkiewicz, P., and Darewicz, M. (2014). Food-originating ACE inhibitors, including antihypertensive peptides, as preventive food components in blood pressure reduction. *Compr Rev Food Sci F.* **13**: 114-134.
- Jao, C. L., Hung, C. C., Tung, Y. S., Lin, P. Y., Chen, M. C., and Hsu, K. C. (2015). The development of bioactive peptides from dietary proteins as a dipeptidyl peptidase IV inhibitor for the management of type 2 diabetes. *Biomed (Taipei).* **5**: 14.

- Jiang, Z., Wang, L., Wu, W., and Wang, Y. (2013). Biological activities and physicochemical properties of Maillard reaction products in sugar–bovine casein peptide model systems. *Food Chem.* **141**: 3837-3845.
- Jin, Y., Yan, J., Yu, Y., and Qi, Y. (2015). Screening and identification of DPP-IV inhibitory peptides from deer skin hydrolysates by an integrated approach of LC–MS/MS and in silico analysis. *J Funct Foods.* **18**: 344-357.
- Karim, A. A., and Bhat, R. (2008). Gelatin alternatives for the food industry. recent developments, challenges and prospects. *Trends Food Sci Techn.* **19**: 644-656.
- Karim, A. A., and Bhat, R. (2009). Fish gelatin: properties, challenges, and prospects as an alternative to mammalian gelatins. *Food Hydrocolloid.* **23**: 563-576.
- Kim, H. K., Kim, M.-G., and Leem, K.-H. (2014). Collagen hydrolysates increased osteogenic gene expressions via a MAPK signaling pathway in MG-63 human osteoblasts. *Food Funct.* **5**: 573-578.
- Kitts, D. D., and Weiler, K. (2003). Bioactive proteins and peptides from food sources. Applications of bioprocesses used in isolation and recovery. *Curr Pharm Design.* **9**: 1309-1323.
- Kumar, S., Sugihara, F., Suzuki, K., Inoue, N., and Venkateswarathirukumara, S. (2015). A double-blind, placebo-controlled, randomised, clinical study on the effectiveness of collagen peptide on osteoarthritis. *J Sci Food Agric.* **95**: 702-707.
- Lacroix, I. M. E., and Li-Chan, E. C. Y. (2012). Evaluation of the potential of dietary proteins as precursors of dipeptidyl peptidase (DPP)-IV inhibitors by an in silico approach. *J Funct Foods.* **4**: 403-422.
- Lacroix, I. M. E., and Li-Chan, E. C. Y. (2016). Food-derived dipeptidyl-peptidase IV inhibitors as a potential approach for glycemic regulation – Current knowledge and future research considerations. *Trends in Food Sci Tech.* **54**: 1-16.

- Lafarga, T., and Hayes, M. (2014). Bioactive peptides from meat muscle and by-products: generation, functionality and application as functional ingredients. *Meat Sci.* **98**: 227-239.
- Lafarga, T., and Hayes, M. (2016). Bioactive protein hydrolysates in the functional food ingredient industry: Overcoming current challenges. *Food Rev Int.* **33**: 217-246.
- Lafarga, T., O'Connor, P., and Hayes, M. (2014). Identification of novel dipeptidyl peptidase-IV and angiotensin-I-converting enzyme inhibitory peptides from meat proteins using in silico analysis. *Peptides.* **59**: 53-62.
- Lee, C. H., Singla, A., and Lee, Y. (2001). Biomedical applications of collagen. *Int J Pharm.* **221**: 1-22.
- Lee, J. K., Jeon, J.-K., and Byun, H.-G. (2011). Effect of angiotensin I converting enzyme inhibitory peptide purified from skate skin hydrolysate. *Food Chem.* **125**: 495-499.
- Lee, J. K., Jeon, J.-K., and Byun, H.-G. (2014). Antihypertensive effect of novel angiotensin I converting enzyme inhibitory peptide from chum salmon (*Oncorhynchus keta*) skin in spontaneously hypertensive rats. *J Funct Foods.* **7**: 381-389.
- Lee, S.-J., Kim, Y.-S., Hwang, J.-W., Kim, E.-K., Moon, S.-H., Jeon, B.-T., Jeon, Y.-J., Kim, J. M., and Park, P.-J. (2012). Purification and characterization of a novel antioxidative peptide from duck skin by-products that protects liver against oxidative damage. *Food Res Int.* **49**: 285-295.
- Li-Chan, E. C., Hunag, S. L., Jao, C. L., Ho, K. P., and Hsu, K. C. (2012). Peptides derived from atlantic salmon skin gelatin as dipeptidyl-peptidase IV inhibitors. *J Agric Food Chem.* **60**: 973-978.
- Li-Chan, E. C. Y. (2015). Bioactive peptides and protein hydrolysates: research trends and challenges for application as nutraceuticals and functional food ingredients. *Curr Opin Food Sci.* **1**: 28-37.

- Lima, C. A., Campos, J. F., Filho, J. L., Converti, A., da Cunha, M. G., and Porto, A. L. (2015). Antimicrobial and radical scavenging properties of bovine collagen hydrolysates produced by *Penicillium aurantiogriseum* URM 4622 collagenase. *J Food Sci Technol*. **52**: 4459-4466.
- Lin, L., Lv, S., and Li, B. (2012). Angiotensin-I-converting enzyme (ACE)-inhibitory and antihypertensive properties of squid skin gelatin hydrolysates. *Food Chem*. **131**: 225-230.
- Liu, B.-Y., Zhu, K.-X., Guo, X.-N., Peng, W., and Zhou, H.-M. (2016). Changes in the enzyme-induced release of bitter peptides from wheat gluten hydrolysates. *RSC Adv*. **6**: 102249-102257.
- Liu, D., Nikoo, M., Boran, G., Zhou, P., and Regenstein, J. M. (2015). Collagen and gelatin. *Annu Rev Food Sci Technol*. **6**: 527-557.
- Liu, J., Zhang, B., Song, S., Ma, M., Si, S., Wang, Y., Xu, B., Feng, K., Wu, J., and Guo, Y. (2014). Bovine collagen peptides compounds promote the proliferation and differentiation of MC3T3-E1 pre-osteoblasts. *PloS one*. **9**: e99920.
- López-Fandiño, R., Otte, J., and Van Camp, J. (2006). Physiological, chemical and technological aspects of milk-protein-derived peptides with antihypertensive and ACE-inhibitory activity. *Int Dairy J*. **16**: 1277-1293.
- Maehashi, K., and Huang, L. (2009). Bitter peptides and bitter taste receptors. *Cell Mol Life Sci*. **66**: 1661-1671.
- Martin-Bautista, E., Martin-Matillas, M., Martin-Lagos, J. A., Miranda-Leon, M. T., Muñoz-Torres, M., Ruiz-Requena, E., Rivero, M., Quer, J., Puigdueta, I., and Campoy, C. (2011). A nutritional intervention study with hydrolyzed collagen in pre-pubertal Spanish children: influence on bone modeling biomarkers. *J Pediatr Endocr Met*. **24**: 147-153.

- Martínez-Alvarez, O., Chamorro, S., and Brenes, A. (2015). Protein hydrolysates from animal processing by-products as a source of bioactive molecules with interest in animal feeding: a review. *Food Res Int.* **73**: 204-212.
- Matsuda, N., Koyama, Y.-i., Hosaka, Y., Ueda, H., Watanabe, T., Araya, T., Irie, S., and Takehana, K. (2006). Effects of ingestion of collagen peptide on collagen fibrils and glycosaminoglycans in the dermis. *J Nutr Sci Vitaminol.* **52**: 211-215.
- McAlindon, T. E., Nuite, M., Krishnan, N., Ruthazer, R., Price, L. L., Burstein, D., Griffith, J., and Flechsenhar, K. (2011). Change in knee osteoarthritis cartilage detected by delayed gadolinium enhanced magnetic resonance imaging following treatment with collagen hydrolysate: a pilot randomized controlled trial. *Osteoarthritis Cartilage.* **19**: 399-405.
- Minaguchi, J., Tometsuka, C., Koyama, Y.-i., Kusubata, M., Nagayasu, A., Sawaya, S., Shiga, T., Shima, H., Hara, T., and Takehana, K. (2012). Effects of collagen-derived oligopeptide prolylhydroxyproline on differentiation of mouse 3T3-L1 preadipocytes. *Food Sci Technol Res.* **18**: 593-599.
- Minkiewicz, P., Dziuba, J., and Michalska, J. (2011). Bovine meat proteins as potential precursors of biologically active peptides--a computational study based on the BIOPEP database. *Food Sci Technol Int.* **17**: 39-45.
- Mooney, C., Haslam, N. J., Pollastri, G., and Shields, D. C. (2012). Towards the improved discovery and design of functional peptides: common features of diverse classes permit generalized prediction of bioactivity. *PloS one.* **7**: e45012.
- Nakchum, L., and Kim, S. M. (2016). Preparation of squid skin collagen hydrolysate as an antihyaluronidase, antityrosinase, and antioxidant agent. *Prep Biochem Biotech.* **46**: 123-130.

- Ney, K. H. (1979). Bitterness of peptides: amino acid composition and chain length. *ACS Symp Ser.* **115**:149–173.
- Ngo, D. H., Kang, K. H., Ryu, B., Vo, T. S., Jung, W. K., Byun, H. G., and Kim, S. K. (2015). Angiotensin-I converting enzyme inhibitory peptides from antihypertensive skate (*Okamejei kenojei*) skin gelatin hydrolysate in spontaneously hypertensive rats. *Food Chem.* **174**: 37-43.
- Ngo, D. H., Ryu, B., and Kim, S. K. (2014). Active peptides from skate (*Okamejei kenojei*) skin gelatin diminish angiotensin-I converting enzyme activity and intracellular free radical-mediated oxidation. *Food Chem.* **143**: 246-255.
- Ngo, D. H., Ryu, B., Vo, T. S., Himaya, S. W., Wijesekara, I., and Kim, S. K. (2011). Free radical scavenging and angiotensin-I converting enzyme inhibitory peptides from Pacific cod (*Gadus macrocephalus*) skin gelatin. *Int J Biol Macromol.* **49**: 1110-1116.
- Nikoo, M., Benjakul, S., Ehsani, A., Li, J., Wu, F., Yang, N., Xu, B., Jin, Z., and Xu, X. (2014). Antioxidant and cryoprotective effects of a tetrapeptide isolated from Amur sturgeon skin gelatin. *J Funct Foods.* **7**: 609-620.
- Niu, H., Wang, Z., Hou, H., Zhang, Z., and Li, B. (2016). Protective effect of cod (*Gadus macrocephalus*) skin collagen peptides on acetic acid-induced gastric ulcer in rats. *J Food Sci.* **81**: H1807-H1815.
- Nongonierma, A. B., and FitzGerald, R. J. (2013). Dipeptidyl peptidase IV inhibitory and antioxidative properties of milk protein-derived dipeptides and hydrolysates. *Peptides.* **39**: 157-163.
- Nongonierma, A. B., and FitzGerald, R. J. (2014). An in silico model to predict the potential of dietary proteins as sources of dipeptidyl peptidase IV (DPP-IV) inhibitory peptides. *Food Chem.* **165**: 489-498.

- Nongonierma, A. B., and FitzGerald, R. J. (2016). Prospects for the management of type 2 diabetes using food protein-derived peptides with dipeptidyl peptidase IV (DPP-IV) inhibitory activity. *Curr Opin Food Sci.* **8**: 19-24.
- Nongonierma, A. B., and FitzGerald, R. J. (2017). Strategies for the discovery and identification of food protein-derived biologically active peptides. *Trends Food Sci Tech.* DOI:10.1016/j.tifs.2017.03.003
- Oesser, S., and Seifert, J. (2003). Stimulation of type II collagen biosynthesis and secretion in bovine chondrocytes cultured with degraded collagen. *Cell Tissue Res.* **311**: 393-399.
- Ohara, H., Ichikawa, S., Matsumoto, H., Akiyama, M., Fujimoto, N., Kobayashi, T., and Tajima, S. (2010). Collagen - derived dipeptide, proline-hydroxyproline, stimulates cell proliferation and hyaluronic acid synthesis in cultured human dermal fibroblasts. *J Dermatol.* **37**: 330-338.
- Ohara, H., Matsumoto, H., Ito, K., Iwai, K., and Sato, K. (2007). Comparison of quantity and structures of hydroxyproline-containing peptides in human blood after oral ingestion of gelatin hydrolysates from different sources. *J Agric Food Chem.* **55**: 1532-1535.
- World Health Organization. (2013). A global brief on hypertension: silent killer, global public health crisis: World Health Day 2013.
- Pal, G. K., and Suresh, P. V. (2016). Sustainable valorisation of seafood by-products: Recovery of collagen and development of collagen-based novel functional food ingredients. *Innov Food Sci Emerg.* **37**: 201-215.
- Patil, P., Mandal, S., Tomar, S. K., and Anand, S. (2015). Food protein-derived bioactive peptides in management of type 2 diabetes. *Eur J Nutr.* **54**: 863-880.

- Pei, X., Yang, R., Zhang, Z., Gao, L., Wang, J., Xu, Y., Zhao, M., Han, X., Liu, Z., and Li, Y. (2010). Marine collagen peptide isolated from Chum Salmon (*Oncorhynchus keta*) skin facilitates learning and memory in aged C57BL/6J mice. *Food Chem.* **118**: 333-340.
- Perazella, M. A., and Setaro, J. F. (2003). Renin-angiotensin-aldosterone system: fundamental aspects and clinical implications in renal and cardiovascular disorders. *J Nucl Cardiol.* **10**: 184-196.
- Porfírio, E., and Fanaro, G. B. (2016). Collagen supplementation as a complementary therapy for the prevention and treatment of osteoporosis and osteoarthritis: a systematic review. *Rev Bras de Geriatr Gerontol.* **19**: 153-164.
- Power, O., Nongonierma, A. B., Jakeman, P., and FitzGerald, R. J. (2014). Food protein hydrolysates as a source of dipeptidyl peptidase IV inhibitory peptides for the management of type 2 diabetes. *Proc Nutr Soc.* **73**: 34-46.
- Prupp, A. H., and Ardö, Y. (2007). Modelling relationship between angiotensin-(I)-converting enzyme inhibition and the bitter taste of peptides. *Food Chem.* **102**: 880-888.
- Proksch, E., Segger, D., Degwert, J., Schunck, M., Zague, V., and Oesser, S. (2014). Oral supplementation of specific collagen peptides has beneficial effects on human skin physiology: a double-blind, placebo-controlled study. *Skin Pharmacol Phys.* **27**: 47-55.
- Rao, Q., Klaassen Kamdar, A., and Labuza, T. P. (2016). Storage Stability of Food Protein Hydrolysates—A Review. *Crit Rev Food Sci Nutr.* **56**: 1169-1192.
- Rajendran, S. R. C. K., Mason, B., and Udenigwe, C. C. (2016). Peptidomics of peptic digest of selected potato tuber proteins: Post-translational modifications and limited cleavage specificity. *J Agric Food Chem.* **64**: 2432-2437.
- Reddy, K. V. R., Yedery, R. D., and Aranha, C. (2004). Antimicrobial peptides: premises and promises. *Int J Antimicrob Ag.* **24**: 536-547.

- Rodríguez, V., Asenjo, J. A., and Andrews, B. A. (2014). Design and implementation of a high yield production system for recombinant expression of peptides. *Microb Cell Fact.* **13**: 65.
- Sae-leaw, T., O'Callaghan, Y. C., Benjakul, S., and O'Brien, N. M. (2016). Antioxidant, immunomodulatory and antiproliferative effects of gelatin hydrolysates from seabass (*Lates calcarifer*) skins. *Int J Food Sci Tech.* **51**: 1545-1551.
- Saito, M., Kiyose, C., Higuchi, T., Uchida, N., and Suzuki, H. (2009). Effect of collagen hydrolysates from salmon and trout skins on the lipid profile in rats. *J Agric Food Chem.* **57**: 10477-10482.
- Samaranayaka, A. G., and Li-Chan, E. C. (2011). Food-derived peptidic antioxidants: A review of their production, assessment, and potential applications. *J Funct Foods.* **3**: 229-254.
- Sarmadi, B. H., and Ismail, A. (2010). Antioxidative peptides from food proteins: a review. *Peptides.* **31**: 1949-1956.
- Sasaki, K., Takahashi, N., Satoh, M., Yamasaki, M., and Minamino, N. (2010). A peptidomics strategy for discovering endogenous bioactive peptides. *J Proteome Res.* **9**: 5047-5052.
- Schadow, S., Simons, V. S., Lochnit, G., Kordelle, J., Gazova, Z., Siebert, H. C., and Steinmeyer, J. (2017). Metabolic response of human osteoarthritic cartilage to biochemically characterized collagen hydrolysates. *Int J Mol Sci.* **18**: 207.
- Schrieber, R., and Gareis, H. (2007). Gelatine handbook: theory and industrial practice. John Wiley & Sons. Weinheim.
- Schwartz, S. R., and Park, J. (2012). Ingestion of BioCell Collagen[®], a novel hydrolyzed chicken sternal cartilage extract; enhanced blood microcirculation and reduced facial aging signs. *Clin Interv Aging.* **7**: 267.
- Shigemura, Y., Akaba, S., Kawashima, E., Park, E. Y., Nakamura, Y., and Sato, K. (2011). Identification of a novel food-derived collagen peptide, hydroxyprolyl-glycine, in human

peripheral blood by pre-column derivatisation with phenyl isothiocyanate. *Food Chem.* **129**: 1019-1024.

Shigemura, Y., Iwai, K., Morimatsu, F., Iwamoto, T., Mori, T., Oda, C., Taira, T., Park, E. Y., Nakamura, Y., and Sato, K. (2009). Effect of prolyl-hydroxyproline (Pro-Hyp), a food-derived collagen peptide in human blood, on growth of fibroblasts from mouse skin. *J Agric Food Chem.* **57**: 444-449.

Shigemura, Y., Kubomura, D., Sato, Y., and Sato, K. (2014). Dose-dependent changes in the levels of free and peptide forms of hydroxyproline in human plasma after collagen hydrolysate ingestion. *Food Chem.* **159**: 328-332.

Shimizu, J., Asami, N., Kataoka, A., Sugihara, F., Inoue, N., Kimira, Y., Wada, M., and Mano, H. (2015). Oral collagen-derived dipeptides, prolyl-hydroxyproline and hydroxyprolyl-glycine, ameliorate skin barrier dysfunction and alter gene expression profiles in the skin. *Biochem Biophys Res Commun.* **456**: 626-630.

Sibilla, S., Godfrey, M., Brewer, S., Budh-Raja, A., and Genovese, L. (2015). An overview of the beneficial effects of hydrolysed collagen as a nutraceutical on skin properties: Scientific background and clinical studies. *Open Nutraceuticals J.* **8**: 29-42.

Sila, A., Martinez-Alvarez, O., Haddar, A., Gomez-Guillen, M. C., Nasri, M., Montero, M. P., and Bougatef, A. (2015). Recovery, viscoelastic and functional properties of Barbel skin gelatine: investigation of anti-DPP-IV and anti-prolyl endopeptidase activities of generated gelatine polypeptides. *Food Chem.* **168**: 478-486.

Singh, T. K., Young, N. D., Drake, M., and Cadwallader, K. R. (2005). Production and sensory characterization of a bitter peptide from β -casein. *J Agric Food Chem.* **53**: 1185-1189.

- Sohaib, M., Anjum, F. M., Sahar, A., Arshad, M. S., Rahman, U. U., Imran, A., and Hussain, S. (2016). Antioxidant proteins and peptides to enhance the oxidative stability of meat and meat products: A comprehensive review. *Int J Food Prop*: 1-13.
- Sugihara, F., Inoue, N., Kuwamori, M., and Taniguchi, M. (2012). Quantification of hydroxyprolyl-glycine (Hyp-Gly) in human blood after ingestion of collagen hydrolysate. *J Biosci Bioeng*. **113**: 202-203.
- Sun, L., Zhang, Y., and Zhuang, Y. (2013). Antiphotaging effect and purification of an antioxidant peptide from tilapia (*Oreochromis niloticus*) gelatin peptides. *J Funct Foods*. **5**: 154-162.
- Taga, Y., Kusubata, M., Ogawa-Goto, K., and Hattori, S. (2014). Highly accurate quantification of hydroxyproline-containing peptides in blood using a protease digest of stable isotope-labeled collagen. *J Agric Food Chem*. **62**: 12096-12102.
- Taga, Y., Kusubata, M., Ogawa-Goto, K., and Hattori, S. (2016). Efficient Absorption of X-Hydroxyproline (Hyp)-Gly after Oral Administration of a Novel Gelatin Hydrolysate Prepared Using Ginger Protease. *J Agric Food Chem*. **64**: 2962-2970.
- Temussi, P. A. (2012). The good taste of peptides. *J Pept Sci*. **18**: 73-82.
- Toldrá, F., Mora, L., and Reig, M. (2016). New insights into meat by-product utilization. *Meat Sci*. **120**: 54-59.
- Trč, T., and Bohmová, J. (2011). Efficacy and tolerance of enzymatic hydrolysed collagen (EHC) vs. glucosamine sulphate (GS) in the treatment of knee osteoarthritis (KOA). *Int Orthop*. **35**: 341-348.
- Tsuruoka, N., Yamato, R., Sakai, Y., Yoshitake, Y., and Yonekura, H. (2007). Promotion by collagen tripeptide of type I collagen gene expression in human osteoblastic cells and fracture healing of rat femur. *Biosci Biotechnol Biochem*. **71**: 2680-2687.

- Udenigwe, C. C. (2014). Bioinformatics approaches, prospects and challenges of food bioactive peptide research. *Trends Food Sci Tech.* **36**: 137-143.
- Udenigwe, C. C., and Aluko, R. E. (2012). Food protein-derived bioactive peptides: production, processing, and potential health benefits. *J Food Sci.* **77**: R11-24.
- Udenigwe, C. C., and Fogliano, V. (2017). Food matrix interaction and bioavailability of bioactive peptides: Two faces of the same coin? *J Funct Foods.* **35**: 9-12.
- Udenigwe, C. C., Gong, M., and Wu, S. (2013). In silico analysis of the large and small subunits of cereal RuBisCO as precursors of cryptic bioactive peptides. *Process Biochem.* **48**: 1794-1799.
- Udenigwe, C. C., and Mohan, A. (2014). Mechanisms of food protein-derived antihypertensive peptides other than ACE inhibition. *J Funct Foods.* **8**: 45-52.
- Valko, M., Rhodes, C., Moncol, J., Izakovic, M., and Mazur, M. (2006). Free radicals, metals and antioxidants in oxidative stress-induced cancer. *Chemico-biological interactions.* **160**: 1-40.
- Venkatesan, J., Anil, S., Kim, S. K., and Shim, M. S. (2017). Marine Fish Proteins and Peptides for Cosmeceuticals: A Review. *Mar Drugs.* **15**: 143.
- Wang, B., Wang, Y.-M., Chi, C.-F., Luo, H.-Y., Deng, S.-G., and Ma, J.-Y. (2013). Isolation and characterization of collagen and antioxidant collagen peptides from scales of croceine croaker (*Pseudosciaena crocea*). *Mar Drugs.* **11**: 4641-4661.
- Wang, J., Xu, M., Liang, R., Zhao, M., Zhang, Z., and Li, Y. (2015a). Oral administration of marine collagen peptides prepared from chum salmon (*Oncorhynchus keta*) improves wound healing following cesarean section in rats. *Food Nutr Res.* **59**: 26411.
- Wang, L., Wang, Q., Liang, Q., He, Y., Wang, Z., He, S., Xu, J., and Ma, H. (2015b). Determination of bioavailability and identification of collagen peptide in blood after oral ingestion of gelatin. *J Sci Food Agric.* **95**: 2712-2717.

- Wang, T. Y., Hsieh, C. H., Hung, C. C., Jao, C. L., Lin, P. Y., Hsieh, Y. L., and Hsu, K. C. (2017). A study to evaluate the potential of an in silico approach for predicting dipeptidyl peptidase-IV inhibitory activity in vitro of protein hydrolysates. *Food Chem.* **234**: 431-438.
- Watanabe-Kamiyama, M., Shimizu, M., Kamiyama, S., Taguchi, Y., Sone, H., Morimatsu, F., Shirakawa, H., Furukawa, Y., and Komai, M. (2010). Absorption and effectiveness of orally administered low molecular weight collagen hydrolysate in rats. *J Agric Food Chem.* **58**: 835-841.
- Weng, W., Tang, L., Wang, B., Chen, J., Su, W., Osako, K., and Tanaka, M. (2014). Antioxidant properties of fractions isolated from blue shark (*Prionace glauca*) skin gelatin hydrolysates. *J Funct Foods.* **11**: 342-351.
- Wu, J., Fujioka, M., Sugimoto, K., Mu, G., and Ishimi, Y. (2004). Assessment of effectiveness of oral administration of collagen peptide on bone metabolism in growing and mature rats. *J Bone Miner Metab.* **22**: 547-553.
- Wu, S., Qi, W., Li, T., Lu, D., Su, R., and He, Z. (2013). Simultaneous production of multi-functional peptides by pancreatic hydrolysis of bovine casein in an enzymatic membrane reactor via combinational chromatography. *Food Chem.* **141**: 2944-2951.
- Xu, L., Dong, W., Zhao, J., and Xu, Y. (2015). Effect of Marine collagen peptides on physiological and neurobehavioral development of male rats with perinatal asphyxia. *Mar Drugs.* **13**: 3653-3671.
- Yamada, S., Nagaoka, H., Terajima, M., Tsuda, N., Hayashi, Y., and Yamauchi, M. (2013). Effects of fish collagen peptides on collagen post-translational modifications and mineralization in an osteoblastic cell culture system. *Dent Mater J.* **32**: 88-95.
- Yazaki, M., Ito, Y., Yamada, M., Goulas, S., Teramoto, S., Nakaya, M. A., Ohno, S., and Yamaguchi, K. (2017). Oral ingestion of collagen hydrolysate leads to the transportation of

highly concentrated Gly-Pro-Hyp and its hydrolyzed form of Pro-Hyp into the bloodstream and skin. *J Agric Food Chem.* **65**: 2315-2322.

Yu, Y., Jin, Y., Wang, F., Yan, J., Qi, Y., and Ye, M. (2017). Protein digestomic analysis reveals the bioactivity of deer antler velvet in simulated gastrointestinal digestion. *Food Res Int.* **96**: 182-190.

Zague, V. (2008). A new view concerning the effects of collagen hydrolysate intake on skin properties. *Arch Dermatol Res.* **300**: 479-483.

Zague, V., de Freitas, V., Rosa, M. d. C., de Castro, G. Á., Jaeger, R. G., and Machado-Santelli, G. M. (2011). Collagen hydrolysate intake increases skin collagen expression and suppresses matrix metalloproteinase 2 activity. *J Med Food.* **14**: 618-624.

Zdzieblik, D., Oesser, S., Gollhofer, A., and Konig, D. (2017). Improvement of activity-related knee joint discomfort following supplementation of specific collagen peptides. *Appl Physiol Nutr Metab.* **42**: 588-595.

Zhang, Y., Chen, R., Chen, X., Zeng, Z., Ma, H., and Chen, S. (2016). Dipeptidyl Peptidase IV-Inhibitory Peptides Derived from Silver Carp (*Hypophthalmichthys molitrix* Val.) Proteins. *J Agric Food Chem.* **64**: 831-839.

Zhang, Y., Duan, X., and Zhuang, Y. (2012). Purification and characterization of novel antioxidant peptides from enzymatic hydrolysates of tilapia (*Oreochromis niloticus*) skin gelatin. *Peptides.* **38**: 13-21.

Zhang, Y., Kouguchi, T., Shimizu, K., Sato, M., Takahata, Y., and Morimatsu, F. (2010). Chicken collagen hydrolysate reduces proinflammatory cytokine production in C57BL/6. KOR-ApoEshl mice. *J Nutr Sci Vitaminol.* **56**: 208-210.

- Zhang, Y., Olsen, K., Grossi, A., and Otte, J. (2013). Effect of pretreatment on enzymatic hydrolysis of bovine collagen and formation of ACE-inhibitory peptides. *Food Chem.* **141**: 2343-2354.
- Zhang, Z., Wang, J., Ding, Y., Dai, X., and Li, Y. (2011). Oral administration of marine collagen peptides from Chum Salmon skin enhances cutaneous wound healing and angiogenesis in rats. *J Sci Food Agric.* **91**: 2173-2179.
- Zhu, C. F., Li, G. Z., Peng, H. B., Zhang, F., Chen, Y., and Li, Y. (2010). Treatment with marine collagen peptides modulates glucose and lipid metabolism in Chinese patients with type 2 diabetes mellitus. *Appl Physiol Nutr Metab.* **35**: 797-804.
- Zhuang, Y., Zhao, X., and Li, B. (2009). Optimization of antioxidant activity by response surface methodology in hydrolysates of jellyfish (*Rhopilema esculentum*) umbrella collagen. *J Zhejiang Univ Sci B.* **10**: 572-579.
- Zhuang, Y., Sun, L., and Li, B. (2010). Production of the angiotensin-I-converting enzyme (ACE)-inhibitory peptide from hydrolysates of jellyfish (*Rhopilema esculentum*) collagen. *Food and Bioprocess Tech.* **5**: 1622-1629.
- Zhuang, Y., Sun, L., Zhang, Y., and Liu, G. (2012). Antihypertensive effect of long-term oral administration of jellyfish (*Rhopilema esculentum*) collagen peptides on renovascular hypertension. *Mar Drugs.* **10**: 417-426.

Figure captions:




Fig. 1 Structural features and functionalities of collagen peptides.




Fig. 2 The approaches for the discovery and production of bioactive collagen peptides.




Fig. 3 Biological activities of collagen peptides.

Table 1 Antihypertensive/ACE inhibitory peptides different sources of collagen.

Source	Enzyme used	Sequence	In vitro/in vivo bioefficacy	Reference
Squid tunic gelatin	Alcalase	GPPGPPGFPGPPGPS	IC ₅₀ =90.03 μM	(Alemán et al. 2011)
Skate skin	α-Chymotrypsin	PGPLGLTGP QLGFLGPR	IC ₅₀ = 95 μM IC ₅₀ = 148 μM	(Lee et al. 2011)
Nile Tilapia gelatin	Alcalase	DPALATEPDMPF	IC ₅₀ = 62.2 μM	(Vo et al. 2011)
Pacific cod gelatin	Pepsin, Trypsin & α-Chymotrypsin	LLMLDNDLPP	IC ₅₀ = 35.7 μM	(Himaya et al. 2012)
Jellyfish collagen	Alcalase	N/A	IC ₅₀ = 43 μg/mL Decreased blood pressure in RVH, 100 mg/kg	(Zhuang et al. 2012)
Squid collagen	Pepsin & Pancreatin	GRGSVPAPGP	IC ₅₀ =47.78 μM	(Alemán et al. 2013)
Mackerel gelatin	Pepsin	FGN	N/A	(Khiari et al. 2013)
Chum salmon skin	Trypsin	GLPLNLP	IC ₅₀ = 18.7 μM Decreased blood pressure in SHR (20 mg/kg)	(Lee et al. 2014)

Skate gelatin	Alcalase	MVGSAPGVL LGPLGHQ	IC ₅₀ = 3.09 μ M IC ₅₀ = 4.22 μ M	(Ngo et al. 2014) (Ngo et al. 2015)
Bovine collagen	Alcalase Papain	VGPV GPRGF	IC ₅₀ = 405.12 μ M IC ₅₀ = 200.92 μ M	(Fu et al. 2016a) (Fu et al. 2016b)

Note: P with boldface denotes Hyp (hydroxylated proline). N/A: not available. SHR, spontaneously hypertensive rats. RHV, renovascular hypertensive (RVH) rat.

Table 2 Antioxidant peptides identified from different sources of collagen.

Source	Enzyme used	Sequence	Antioxidant activity tests	IC ₅₀ (μ M)	Reference
Pacific cod gelatin	Papain	TCSP TGGGNV	DPPH and hydroxyl radical scavenging, DNA damage and ROS	Scavenging activities of DPPH and hydroxyl radicals, 72% and 56% at 1 mg/mL	(Ngo et al. 2011)
Skate skin gelatin	Alcalase, Flavourzyme, Neutrase and Protamex	MVGSAPGVL LGPLGHQ	ROS, glutathione and superoxide dismutase	N/A	(Ngo et al. 2014)
Scales of croceine coaker	Trypsin & Pepsin	GFRGTIGLVG GPAGPAG GFPSG	Hydroxyl, DPPH, superoxide and ABTS radical scavenging	Hydroxyl radical (0.293, 0.240, and 0.107 mg/mL), DPPH radical (1.271, 0.675, and 0.283 mg/mL), superoxide radical (0.463, 0.099, and 0.151 mg/mL) and ABTS radical (0.421, 0.309, and 0.210 mg/mL).	(Wang et al. 2013)
Alaska pollock skin collagen	Trypsin & Flavourzyme	GPAGPHGPPG	Ca, Fe and Cu chelating activity	11.52 \pm 2.23, 1.71 \pm 0.17 and 0.43 \pm 0.02 μ mol/ μ mol	(Guo et al. 2015)
Duck skin gelatin	Collagenase & pepsin	HTVQCMFQ	Hydroxyl, DPPH and alkyl radical scavenging	32.6, 22.7, 55.1, and 49.8 μ g/mL	(Lee et al. 2012)

			ROS		
Spanish dry-cured ham	-	GLAGA	DPPH radical scavenging and ferric-reducing antioxidant power	0.5 units of absorbance at 1 mg/mL	(Escudero et al. 2013)
Amur sturgeon skin gelatin	Alcalase	PAGT	DPPH, ABTS and hydroxyl radical scavenging	5.38, 0.008 and 0.89 mg/mL	(Nikoo et al. 2014)
Blue shark skin gelatin	Protamex	-	DPPH and hydroxyl radical scavenging	0.57 and 2.04 mg/mL	(Weng et al. 2014)
Tilapia skin gelatin	Properase E	LSGYGP	Hydroxyl radical scavenging	22.47 µg/mL.	(Sun et al. 2013)
Collagen from Nile tilapia	Pepsin, trypsin & chymotrypsin	GPM	DPPH radical	25.64 µg/mL	(Liang et al. 2011)
Tilapia skin gelatin	Properase E	EGL YGDEY	Hydroxyl radical scavenging	4.61 and 6.45 µg/mL	(Zhang et al. 2012)
Grass carp skin	Alcalase	PYSFK GFGPEL VGGAP	DPPH, hydroxyl, and ABTS radical scavenging	DPPH radical (2.459, 3.634 and 6.063 mM), hydroxyl radical (3.563, 2.606 and 4.241 mM), ABTS radical (0.281, 0.530 and 0.960 mM)	(Cai et al. 2015)

Table 3 DPP-IV inhibitory peptides from different sources of collagen.

Source	Enzyme used	Sequence	IC ₅₀ (µM)	Reference
Bovine collagen	Papain, Ficin or Bromelain	PPG	2252.68	(Lafarga et al., 2014)
Atlantic salmon gelatin	Alcalase, Bromelain or Flavourzyme	GPAE GPGA	49.6 41.9	(Li-Chan et al., 2012)
Barbel fish gelatin	Esperase, Savinase, Alcalase,	-	-	(Sila et al., 2015)

	Trypsin, Izyme G, Protamex, Neutrase, Peptidase			
Deer collagen	Pepsin, Pepsin/Alcalase, Pepsin/Trypsin	GPGSPGGPL GPVGXAGPPGK GPM(O)GPXGVK GPVGPSGPXGK GPAGPXGVXGL	1638.3 83.3 226.9 93.7 318.1	(Jin et al., 2015)
Halibut, hake, tilapia, milkfish	Flavourzyme	SPGSSGPQGFTG GPVGPAGNPGANGLN PPGPTGPRGQPGNIGF IPGDPGPPGPPGP LPGERGRPGAPGP GPKGDRGLPGPPGRDGM	101.6 81.3 146.7 65.4 76.8 89.6	(Wang et al., 2015)
Pork, cattle, fish, chicken feet	Streptomyces or Collagenase	GAX GPA GPX	>20,000 5030 2510	(Hatanaka et al., 2012)
Porcine skin gelatin	Alcalase	GPX GPAG	45.3 41.1	(Hsu et al., 2013)
Silver Carp collagen	Neutrase	APGPAGP LPIIDI	229.14 105.44	(Zhang et al., 2016)
Alaska pollock skin collagen	Pepsin, Corolase PP & Trpsin	-	-	(Guo et al., 2014)

Table 4 Summary of animal and human intervention studies on the identification and quantification of collagen peptides *in vivo*.

Source	Peptides	Locations	Maximal concentration	Bioactivity	Reference
Commercial fish scale gelatin hydrolysate	Hyp containing di- and tri- peptides	Plasma of adults	60.65 nmol/mL Pro-Hyp	Fibroblast proliferation	(Ichikawa et al., 2010)
Commercial fish scale gelatin	Pro-Hyp and Hyp-Gly	Plasma of adults	120 nmol/mL	Fibroblast proliferation and improved skin	(Shigemura et al., 2011) (Shimizu et

hydrolysate				barrier dysfunction	al., 2015) (Haratake et al., 2015)
Porcine skin gelatin hydrolysate	Hyp-Gly	Plasma of adults	4.2 nmol/mL	Beneficial effects on bone tissue	(Sugihara et al., 2012)
Commercial fish scale gelatin hydrolysate	Hyp and Hyp containing di- and tri-peptides	Plasma of adults	0.663 nmol/mL Pro-Hyp-Gly 163 nmol/mL Pro-Hyp	Fibroblast proliferation	(Taga et al., 2014)
Bovine skin gelatin hydrolysates	Tripeptides (X-Hyp-Gly)	Plasma of rats	0.78 nmol/mL Ala-Hyp-Gly	Chondroprotective effect and DPP-IV inhibitory activity	(Taga et al., 2016)
Porcine skin Collagen tripeptides	Gly-X-Y tripeptide	Plasma of rats	130 nmol/mL Gly-Pro-Hyp	Interaction with platelets and the central nervous system	(Yamamoto et al., 2015)
Chicken collagen hydrolysate	Ala-Hyp	Plasma of adults	2.27 nmol/mL	ACE-inhibitory activity	(Iwai et al. 2009)
Porcine skin collagen hydrolysates	Tripeptides (Gly-X-Y)	Plasma of adults	Ala-Hyp (26.01 nmol/mL) Gly-Pro-Hyp (21.13 nmol/mL) Pro-Hyp (16.84 nmol/mL)	Diverse activities	(Yazaki et al., 2017)