



Biotechnological potential of microbial inulinases: Recent perspective

Hemant Kumar Rawat, Hemant Soni, Helen Treichel & Naveen Kango

To cite this article: Hemant Kumar Rawat, Hemant Soni, Helen Treichel & Naveen Kango (2016): Biotechnological potential of microbial inulinases: Recent perspective, Critical Reviews in Food Science and Nutrition, DOI: [10.1080/10408398.2016.1147419](https://doi.org/10.1080/10408398.2016.1147419)

To link to this article: <http://dx.doi.org/10.1080/10408398.2016.1147419>



Accepted author version posted online: 10 Mar 2016.



Submit your article to this journal [↗](#)



View related articles [↗](#)



View Crossmark data [↗](#)

Title page**Biotechnological potential of microbial inulinases: Recent perspective**

Hemant Kumar Rawat^a, Hemant Soni^a, Helen Treichel^b and Naveen Kango^{a*}

^aDepartment of Applied Microbiology and Biotechnology,
Dr. Harisingh Gour University, Sagar, (M.P.), 470003, India

^bUniversidade Federal da Fronteira Sul-Campus de Erechim,
Av. Dom Joao Hoffmann, 313, Erechim, RS 99700-000, Brazil

* Correspondence

E-mail addresses: **nkango@gmail.com** (N. Kango), **helentreichel@gmail.com** (H. Treichel)

Abstract

Among microbial enzymes, inulinases or fructofuranosylhydrolases have received considerable attention in past decade and as a result a variety of applications based on enzymatic hydrolysis of inulin have been documented. Inulinases are employed for generation of fructose and inulooligosaccharides (IOS) in a single step reaction with specificity. The high fructose syrup can be biotransformed into value added products such as ethanol, single cell protein while IOS are indicated in nutraceutical industry as prebiotic. Myriad microorganisms produce inulinases and a number of exo and endoinulinases have been characterized and expressed in heterologous hosts. Initially predominated by *Aspergilli*, *Penicillia* and some yeasts (*Kluyveromyces* spp.), the

list of prominent inulinase producers has gradually expanded and now includes extremophilic prokaryotes and marine derived microorganisms producing robust inulinases. The present article summarizes important developments about microbial inulinases and their applications made in the last decade.

Keywords Inulin, exoinulinase, endoinulinase, fructose, fructooligosaccharide, immobilization

1. Introduction

Microbial inulinases are an important class of industrial enzymes, which are usually inducible and extracellular. Microbial inulinases belong to the members of glycosylhydrolases (GH 32 family). Inulinases hydrolyzing β -(2 \rightarrow 1) fructosyl linkage are produced by many microorganisms (bacteria, fungi and yeast). Exoinulinase (EC 3.8.1.80; β -D-fructan fructohydrolase or β -D-fructosidase) liberates the terminal fructose unit of inulin linked by hydrolyzing β -fructofuranosidic bonds. High fructose syrup (upto 95%) generated by inulinase action can be utilized as a fermentable substrate. This process is preferable over starch or cellulose degradation as these require enzyme consortia for generation of feedstock glucose. The single step process involving one enzyme encoded in a single gene is therefore much attractive over use of enzyme consortia and has better prospects in terms of cloning and heterologous expression (Chi et al., 2011; Kango and Jain, 2011; Liu et al., 2013). Endoinulinase (EC 3.2.1.7; β -D-fructan fructanohydrolase) hydrolyzes the internal bonds of inulin and liberates inulooligosaccharides (IOS) including inulotriose, inulotetraose and inulopentaose as major products. Fructose is a low-calorie healthy sweetener and a readily fermentable feedstock sugar. IOS have similar structure and functions of fructooligosaccharides (FOS) which have been well known as functional food and nutraceutical (Ganaie et al., 2014; Tsurumaki et al., 2015; Singh et al., 2016).

Inulin, the second most abundant storage polysaccharide after starch, can be exploited as a substrate for the production of ultra-high fructose syrup (UHFS), IOS, bioethanol, single cell oil (SCO), citric acid, lipids, single cell protein (SCP), 2, 3-butanediol, lactic acid and other

important chemicals (Chi et al., 2009; Zhang et al., 2010a; Cui et al., 2011; Kuhn et al., 2012; Wang et al., 2012 and 2014; Rawat et al., 2015b). Most of the industrial processes using inulin as raw substrate employ microbial inulinases preceding its biotransformation. Among fungi, members of *Aspergillus*, *Penicillium* and *Kluyveromyces* spp. are reported as prolific inulinase producers (Liu et al., 2013; Rawat et al., 2015a). Some bacteria such as *Xanthomonas* and *Arthrobacter* are also known sources of microbial inulinases (Chen et al., 2009; Vijayaraghavan et al., 2009; Zhou et al., 2014a).

With recent explorations, the realm of microbial inulinase producers has expanded to extremophilic prokaryotes and marine derived yeasts (Chi et al., 2009). Heat, salt and cold active inulinases sourced from extremophilic microorganisms can be appropriately utilized in developing industrial bioprocesses. The present review covers the more recent developments on microbial inulinases in the last ten years. It discusses their properties, improved production *via*. parametric optimization, heterologous expression and biotechnological applications.

2. Microbial sources of inulinase

Inulin hydrolyzing enzymes are reported to occur in diverse and phylogenetically different microorganisms including bacteria, filamentous fungi, yeasts and actinomycetes (Kango and Jain, 2011). More recently, the focus has been to identify novel inulinases exhibiting tolerance of extreme conditions to suit industrial applications. Halotolerant and cold active exoinulinase from *Arthrobacter* sp. MN8 was active at low temperatures (0 to 20°C) (Zhou et al., 2014a). In another study, *Arthrobacter* sp. S37 endoinulinase active over cold and acidic pH has been cloned in *Yarrowia lipolytica* (Li et al., 2012). Similarly, a novel exoinulinase tolerant to detergent, salt and protease has been reported from *Sphingomonas* sp. JB 13 isolated from

phosphate rock site (Zhou et al., 2015). *Lactobacillus paracasei* B41 isolated from Bulgarian traditional beverage ‘boza’ producing inulinase (232 U/ml) has been demonstrated to hydrolyze chicory flour inulin and convert it into lactic acid (Petrova et al., 2015). *Lactobacillus casei* produced distinct zone of clearing on inulin agar plate and inulinase gene (lev H1) has been deciphered (Kuzuwa et al., 2012). Thermophilic soil isolate, *Bacillus smithii* T7 produced 135.2 U/ml thermostable endoinulinase releasing fructose and oligosaccharides with optimum activity at 70°C (Gao et al., 2007). Exclusive endoinulinase production has been reported from *Bacillus safensis* AS-08. Some inulinase producing bacteria *Xanthomonas oryzae*, *Pseudomonas* sp., *Bacillus licheniformis* and some actinomycetes *Nocardiopsis* sp. DN-K15, *Streptomyces* sp. etc. are recent additions (Naidoo et al., 2009; Lu et al., 2013; Singh and Singh, 2014; Li et al., 2014).

Among filamentous fungi, members of *Aspergillus* and *Penicillium* are reported as prolific producers of inulin hydrolyzing enzymes (Kango and Jain, 2011). Several strains of *A. niger*, a prominent producer, sourced from a variety of ecological niches have been demonstrated to produce high titers of inulinase when subjected to growth on media containing inulin-rich vegetal extracts (Housseiny, 2014; Rawat et al., 2015a). Screening of 200 fungal strains isolated from Jerusalem artichoke fields resulted in identification of *Penicillium subrubescens* FBCC 1632 that efficiently produced inulinase (Mansouri et al., 2013). Such explorations have lead to the availability of a number of inulinase genes and present opportunity of developing industrial strains (Chi et al., 2011; Chen et al., 2013; Flores-Gallegos et al., 2015).

Recently, inulinases from *Aspergillus niger* F0602, *Aspergillus ficuum* JNSP5-06 and *Penicillium janthinellum* have been expressed in yeast, *Pichia pastoris* (Chen et al., 2013; He et al., 2014; Chen et al., 2015). A fungal strain, *Aspergillus tubingensis* CR16 isolated from chicory rhizospheric soil produced 257 U/gds yield of inulinase in SSF (Trivedi et al., 2012). *Penicillium* sp. NFCC 2768 has been isolated from dahlia rhizospheric soil and explored for inulinase (3.83 U/ml) production (Rawat et al., 2015b). *A. niger* AUMC 9375 originating from soil samples of Cairo city Egypt, has been reported to be a prolific inulinase (440 U/ml) producer (Housseiny, 2014). Inulinase production is reported from *Aspergillus terreus*, *Aspergillus versicolor*, *Aspergillus parasiticus* and *Penicillium brevicompactum* originating from marine-derived samples collected from Ismalia, Egypt (Abd El Aty et al., 2014). Some other fungi *Thermomyces lanuginosus*, *Fusarium oxysporum*, *Aspergillus awamori*, *Aspergillus fumigatus* are also reported as inulinase producers (Nguyen et al., 2013; Chen et al., 2015; Rawat et al., 2015a).

Diploid yeast, *Kluyveromyces marxianus*, is a known producer of exoinulinase while some of the marine derived yeasts such as *Meyerozyma guilliermondii*, *Cryptococcus aureus*, *Zygosaccharomyces cerevisiae* are recent additions (Kango and Jain, 2011; Chi et al., 2011; Liu et al., 2014). Among these, *Pichia guilliermondii* sourced from sea water from China, produced 60 U/ml exoinulinase. Increase in NaCl concentration (up-to 2%) and addition of sea water to the culture medium enhanced inulinase production by two-folds (Gong et al., 2007). Novel yeast strains *Zygosaccharomyces*, *Gordonia*, *Hanseniaspora*, *Torulospora*, *Saccharomyces*, *Metschnikowia*, *Lachancea* were screened for inulinase production, and *Zygosaccharomyces bailii* (Talf1) was identified as prominent inulinase

(8.67 U/ml) producer when Jerusalem artichoke juice was used as substrate (Paixao et al., 2013).

3. Production and optimization of inulinases

Cost-effectiveness of an enzymatic bioprocess is a prime requisite for realizing its industrial application. Use of low-cost inulin-rich plant extracts, strategic statistical optimization, over-expression of inulinase and high cell density fermentations are being explored in order to achieve this objective. *B. safensis* AS-08 produced 28 U/ml of endoinulinase in medium containing raw dahlia tuber extract for extracellular inulinase production (Singh and Singh, 2014). *L. paracasei* DSM 23505, a prolific exoinulinase producer strain has been used for saccharification of chicory flour (136 g/L) and its biotransformation into lactic acid (123.7 g/L) (Petrova et al., 2015). *Nocadiopsis* sp. DN-K15, novel alkali-tolerant and thermostable actinomycete strain produced 25 U/ml of inulinase using dahlia tuber extract (Lu et al., 2013).

Several reports suggest use of low-value particulate inulin-rich substrates in solid state fermentation (SSF) for inulinase production by filamentous fungi (**Table 1**). *A. niger* AUMC 9335 produced 300 U/gds of endoinulinase on sunflower tubers and lettuce roots mixed in 6:1 (w/w) ratio (Housseiny, 2014). Trivedi *et al.* (2012) have used wheat bran and corn steep liquor for production of inulinase (257 U/gds) by newly isolated *A. tubingensis* CR16 in SSF. Inulinase extraction from fermented bran has also been optimized leading to enhanced yield upto 193 U/gds (Chen et al., 2011).

Several inulin-rich substrates have been used for the production of microbial inulinases *viz.* dandelion, dahlia, asparagus and Jerusalem artichoke (Mansouri et al., 2013; Paixao et al.,

2013; Rawat et al., 2015a). Kango (2008) used dandelion tap root extract and yeast extract used for inulinase production from *A. niger* NK-126 and obtained 55 U/ml in 96 h. *Penicillium citrinum* isolated from Mexican desert produced 18.7 U/L inulinase at 30°C after 48 h (Adriana et al., 2012).

K. marxianus NRRLY-7571 produced 586 U/gds on sugarcane bagasse in fed-batch mode (Astolfi et al., 2011). *K. marxianus* has been demonstrated to produce inulinase on dahlia tuber extract (1.51 U/ml) and asparagus root extract (1.49 U/ml) (Jain et al., 2012; Rawat et al., 2015a). Induced mutant, *Geotrichum candidum* MEU-5fc-6 showed 50-fold enhancement in exoinulinase titers (71.8 U/ml) when grown on sucrose (5 g/L) in 48 h in a fermentor (Mughal et al., 2009). Sheng et al. (2007) isolated marine yeast *Cryptococcus aureus* G7a producing 85 U/ml exoinulinase in shake flask after 42 h fermentation.

Optimization of medium components is very important in the development any bioprocess and helps in understanding interaction of parameters and identifying optimal range for higher yield. Several workers have reported effective optimization of inulinase production using fungal, bacterial and yeast strains and a variety of substrates such as Jerusalem artichoke, sugarcane bagasse, molasses in submerged or solid state cultivation. A comprehensive list of microorganisms and optimized experimental variables is presented in **Table 2**. Novel inulinases from *M. guilliermondii* and *Kluyveromyces cicerisporus* were expressed in *Saccharomyces cerevisiae* and *P. pastoris* X33 and produced 43.8 and 45.2 U/ml, respectively (Liu et al., 2014; Ma et al., 2015). High titers of inulinase (167.4 U/ml) were produced by alginate immobilized *P. guilliermondii* M-30 on Jerusalem artichoke tuber extract in 48 h (Zhao et al., 2011).

4. Heterologous expression and molecular regulation of inulinase

Heterologous expression in yeast and bacterial systems for efficient production of exo and endoinulinase sourced from bacterial and fungal strains has been a matter of intense research. Yeasts play important host for heterologous expression of inulinase and can grow to very high cell densities with established easy recovery methods (Chi et al., 2011; Liu et al., 2013; Zhou et al., 2014b). Characteristics of inulinase genes expressed in heterologous hosts are presented in **Table 3**. Most commonly used yeasts for heterologous expression of inulinase are *P. pastoris*, *Y. lipolytica*, *S. cerevisiae*, *Kluyveromyces lactis* (Zhang et al., 2010b; Yuan et al., 2013; Liu et al., 2014). Simultaneous inulin hydrolysis and fructose bioconversion to ethanol by engineered yeast presents an opportunity of consolidated bioprocessing (CBP) of inulin (Yuan et al., 2012). Endoinulinase (*inuA*) gene of *A. niger* was cloned and expressed in *S. cerevisiae* JZIC-J producing 3.1 U/ml inulinase and 55.3 g/L ethanol (Yuan et al., 2013). Chen et al. (2013) have expressed exoinulinase gene (exon I and exon II) of *A. ficuum* JNSP5-06 in *Escherichia coli*. The molecular weight of recombinant inulinase was 63 kDa with optimum temperature and pH 55°C and 5.0, respectively. Over expression of *P. guilliermondii* *Inu1* (pPlcZaA) gene in *P. pastoris* X-33 resulted in high level of inulinase titers (286.8 U/ml) in 120 h (Zhang et al., 2009a). Exoinulinase gene from *P. janthinellum* B01 was cloned in *P. pastoris* and high level of expression 272.8 U/ml was achieved which was 11 fold higher. The deduced amino acid sequence indicated highly conserved N-terminal region GH32 and 82% identity with *Talaromyces stipitatus* exoinulinase but was divergent from other microbial exoinulinases (Wang et al., 2011). Thermostable exoinulinase (T_{opt} 55°C) from *K. cicerisporus*

was expressed in *P. pastoris* X33. Very high degree of sequence identity was noticed with *K. marxianus* and *C. aureus* exoinulinases (92-99%).

Studies on regulation of inulinase synthesis have been mostly limited to fungal and yeasts expression systems. Several such reports have established that synthesis of inulinase is subject to repression by glucose and fructose present in the medium. The transcriptional repressor Mig1 coded by *MIG1* gene, a C₂H₂ zinc finger protein, binds to the promoter of genes repressed by glucose (Zhou et al., 2014c). Exoinulinases have conserved motifs WMNDPNGL, RDP, ECP, SVEVF, FS and Q while those in endoinulinases are reported to be WMNEPNGL, RDP, EVP, SVEVF, FT and Q (Liu et al., 2013; Cao et al., 2013). The alignment studies with other exoinulinases showed WMNDPNG, RDP and ECP as three highly conserved motifs (Ma et al., 2015). The expression of *A. niger* 12 *inuE* gene is repressed by fructose and glucose (Yuan et al., 2006). Deletion of *MIG1* gene responsible for glucose repression resulted in enhanced exoinulinase production from *Saccharomycopsis fibuliegra* (Liu et al., 2011). *K. marxianus* strain with deleted *MIG1* gene produced 5.7 U/mg inulinase which was two-fold higher as compared to wild type (Liu et al., 2013). Mig1 binding site, however, is reported to be absent in *A. niger* endoinulinase gene. Thus the gene expression is independent from carbon sources (Liu et al., 2013). In another filamentous fungus *Penicillium* sp. TN 88 the expression of *inuC* and *inuD* was enhanced by 42 and 3260-fold when grown on inulin containing media, while sucrose and fructose showed no induction and glucose showed strong catabolite repression (Moriyma et al., 2006). Inulinolytic genes of *A. niger* have been shown to be co-regulated by inulin and sucrose through a positively acting transcription factor InuR. This factor is a member of fungal specific transcription factor of Zn(II)₂Cys₆. Expression of

inulinolytic genes (*inuE*, *inuA*) and sucrolytic gene (*sucA*) was found to be dependent on InuR (Yuan et al., 2008). Recently, *K. marxianus* derepressed mutant (KM-69) with disrupted *MIG1* gene produced 101.7 U/ml inulinase, while the native strain KM-0 was sensitive to glucose repression (Zhou et al., 2014c).

5. Extremophilic inulinases

Although inulinase production from mesophilic counterparts has been reported by many workers, studies on extremophilic inulinase producing microorganisms are relatively less. However, several reports on cold-active and thermophilic inulinases indicate the recent spurt of interest in this area (**Table 4**). Zhou et al. (2014a) reported cold-adaptive *Arthrobacter* sp. MN8 bacterium isolated from lead-zinc rich soil and exploited for its ability for exoinulinase production in generation of fructose syrup at low temperature. Thermostable *Bacillus smithii* T7 strain has been reported to produce 135.2 U/ml thermostable endoinulinase in media containing inulin. The strain was stable over pH range of 4.0 to 8.0 and half life of inulinase 9 h at 70°C (Gao et al., 2009). A novel detergent, salt and protease tolerant exoinulinase from *Sphingomonas* sp. JB13 (rInuAJB13) was optimally active at 55°C and pH 5.0 and stable at 70°C (Zhou et al., 2015).

6. Biotechnological potential of inulinases

Inulin is one of the abundant storage polysaccharides that await exploitation. It can be used as a source of fructose and fructooligosaccharides which can further be utilized as feedstock for production of ethanol, gluconic acid, sorbitol and other important products (Chi et al., 2011; Kango and Jain, 2011).

6.1 Production of fructose and fructooligosaccharides

Fructose is a generally recognized as safe (GRAS) sweetener, sweeter than sucrose (up to 1.5 times), with lower calories and functional properties that enhance flavor, color and product stability. Furthermore, fructose metabolism bypasses the known metabolic pathway of glucose and therefore does not require insulin. High fructose syrup (HFS) produced by the action of inulinase can be beneficially used for diabetic patients. Thus it is widely used in many food, pharmaceutical and beverage preparations instead of sucrose (Ettalibi and Baratti, 2008; Barclay et al., 2010).

Fructose can be produced by one-step enzymatic hydrolysis of inulin. Exoinulinases from different microorganisms are used for production of ultra high fructose syrup (UHFS) from inulin and inulin-containing materials (Mansouri et al., 2013). Yeast inulinases are generally reported to be exo-acting enzymes and only fructose is released from inulin by their action. Such exoinulinases from marine yeasts *C. aureus* G7a (Zhang et al., 2009a) and *P. guilliermondii* strain (Zhang et al., 2009b) have been purified and applied for fructose generation. *A. niger* inulinase produced 37.5 g/L fructose from 100 g/L inulin in 20 h while the *Candida guilliermondii* TISTR 5844 inulinase produced 35.3 g/L fructose in 25 h under similar conditions (Sirisansaneeyakul et al., 2007). *Asparagus racemosus* raw inulin extract and pure inulin have been used to obtain fructose yields of 39.2 and 40.2 g/L, respectively in 4 h, using *K. marxianus* YS-1 exoinulinase (Singh et al., 2006). Exoinulinase from *A. ficuum* produced 98 mg/L fructose from inulin (g/L) at 50°C (Mutanda et al., 2009). Optimized kinetic parameters ($K_m = 24.2$ g/L/h and $V_{max} = 0.108$ g/L min) were proposed for complete inulin hydrolysis and fructose production from chicory using inulinase at 40°C, isolated from *A. ficuum* (Ricca et al., 2009). 40°C

A. niger inulinase was immobilized on Concanavalin-A attached cryogel (poly ethylene glycol dimethacrylate) for the production of high fructose syrup and the conversion efficiency with pure inulin was found to be 0.23 mg/mL after 5 min hydrolysis (Altunbas et al., 2013). Some other inulinase immobilization and fructose generation studies are described in **Table 5**. Zhao et al. (2011) reported direct conversion of inulin and Jerusalem artichoke extract into single cell oil (SCO) mediated by calcium alginate immobilized inulinase of *Rhodotorula mucilaginosa* TJY15a. The bioprocess yielded 56.6% (w/w) oil from inulin rich tuber extract with SCO density reaching 12.8 g/L in 48 h in a 2 l fermenter.

The immobilized endoinulinase of *A. niger* on chitosan was used for continuous production of oligofructose syrup generation from Jerusalem artichoke juice. End product analysis suggested that higher DP (DP8 and more) oligosaccharides were hydrolyzed into syrup containing DP3-DP7 (Nguyen et al., 2011). Inulin was hydrolyzed by the endoinulinases to produce IOS such as inulotriose and inulotetraose and other inulooligosaccharides (Ronkart et al., 2007; Mutanda et al., 2014). Fructo-oligosaccharides produced enzymatically by the action of two different enzymes viz. (i) transfructosylation of sucrose by fructosyl transferase (FTase) and (ii) hydrolysis of inulin by endoinulinase where it randomly cleaves β -2,1 linkages of inulin to yield inulo-oligosaccharides (Kim et al., 2008; Silva et al., 2013; Ganaie et al., 2014; Bali et al., 2015). Oligofructose have numerous applications in food industries like confectionary, milk desserts, chocolate, ice cream and sauces (Kuntz et al., 2013). The use of IOS in animal nutrition has also attracted considerable recent interests, primarily because they significantly change colonic bacterial populations and fermentation end-products (Kelly, 2009). Endoinulinase has been characterized from several species of filamentous fungi and bacteria (Kango and Jain, 2011; Li et

al., 2012). Inulobiose (F2), inulotriose (F3), inulotetraose (F4) and IOS were obtained by the action of *A. niger* NK-126 fungal inulinase (Kango, 2008). Some other *A. niger* strains are also reported to produce endoinulinases (Kango and Jain, 2011). Rawat et al. (2015a) screened 25 fungal strains, in which *A. fumigatus*, *P. citrinum* and *Penicillium rugulosum* produced mixture of endo and exo-inulinase. Combined action of endo and exoinulinase from *Penicillium* sp. NFCC 2768 and produced a mixture of oligosaccharides, GF2, GF3, and GF4 along with glucose and fructose. An overview of biotechnological applications of microbial inulinases for achieving various objectives are given in **Table 6**.

6.2 Production of ethanol and single cell oil (SCO)

Ethanol is one of the most employed biofuel either as a fuel or as a gasoline enhancer. Selective hydrolysis of inulin using exoinulinase produces fructose that can be selectively converted into ethanol (Chi et al., 2011). Mostly *S. cerevisiae* and *K. marxianus* are used for the production of bio-ethanol from feedstock sugars (Abe et al., 2009; Yuan et al., 2012). Different raw materials used for bioethanol production include sugarcane, starch and lignocelluloses. However, these materials require different pre-treatments and sometimes suffer several disadvantages such as competition for food-grade starch and sucrose, interfering lignin, availability of lignocelluloses etc. (Schmitt et al., 2012). Inulin, produced by large scale cultivation of Jerusalem artichoke or chicory can be utilized for generation of fructose feedstock. Fructose, a readily fermentable substrate can be utilized and converted into ethanol (Chi et al., 2009; Hu et al., 2012).

Kluyveromyces sp., although reported to be a prolific producer of inulinase, cannot tolerate high concentrations of ethanol during production compared to *S. cerevisiae* (Lane and Morrissey,

2010). *K. marxianus* exoinulinase gene was expressed in *S. cerevisiae* and the recombinant yeast was grown on Jerusalem artichoke tubers extract. The conversion efficiency of the inulin type sugar to ethanol was 70% (Lim et al., 2010; Yuan et al., 2013). 106.5% mg/ml of ethanol was produced from 300.0 g/L inulin within 120 h by using endoinulinase obtained from *Arthrobacter* sp. (Li et al., 2013). Liu et al. (2014) reported bioethanol production (126.3 mg/mL) from inulin (300 g/L) by using *M. guilliermondii* exoinulinase expressed in *Saccharomyces* sp. W0 (Table 6).

Fresh Jerusalem artichoke contains nearly 20% (w/w) carbohydrate of which 70-90% (w/w) is inulin. About 50-60 g/kg of fresh weight of Jerusalem artichoke is inulin type fructan and the crop yield is estimated to 5.4 ton/ha (Li and Chan-Halbrecht, 2009). Immobilized cells of inulinase producers *Pichia guilliermondii* M-30 were co-cultured with oleaginous yeast *R. muciliginosa* on Jerusalem artichoke tuber extracts. The oleaginous yeast utilized sugar liberated from Jerusalem artichoke tubers and 48.59% (w/w) oil was accumulated (Zhao et al., 2010b; Wu et al., 2011; Wang et al., 2014). *R. mucilaginosa* TJY15a yeast culture was grown on Jerusalem artichoke juice for single cell oil production and accumulated maximum 55.4% oil (Zhao et al., 2011). Wang et al. (2014) used *R. toruloides* 2F5 yeast strain for direct conversion of inulin into lipid (62.14% w/w) at flask level while 70.36% (w/w) lipid production was reported in optimized conditions in a 2 l fermenter. Lipid content obtained after fermentation was also used for generation of biodiesel (Zhao et al., 2010c).

7. Conclusion and future perspectives

Owing to prospects in food and nutraceutical industries, inulin hydrolysis and bioconversion using microbial enzymes has received attention by several workers. Newer raw

materials with high inulin content have been demonstrated to be useful in generation of bioethanol and single cell oil. Heterologous expression of novel inulinases has paved way for consolidated bioprocessing of inulin by ethanol tolerant yeast expressing high titers of exoinulinase. Robust inulinases sourced from extremophiles will allow newer applications particularly at low temperature or in high salt foods. Although a number of exoinulinases have been characterized from yeasts and overexpressed in suitable hosts, fewer endoinulinases and their lesser understood catalytic mechanism presents a greater challenge. Availability of inulin as raw-material from regular and horticultural crops and developing a self-sustainable bioprocess either for generation of high fructose syrup or oligosaccharides is a prime concern that needs to be worked out.

Critical analysis of current literature shows that microbial inulinase is one of the most promising group of enzymes to be employed in food industries. This review showed that many researchers worldwide direct their activities to the screening of new inulinase-producing microorganisms and, subsequently, on the optimization of the medium composition and operational variables. All these efforts are justified by the relevance of inulinase applications.

Acknowledgement

Authors are thankful to Madhya Pradesh Biotechnology Council (M.P.B.C.), Bhopal, project- (PA-23/656) India, for financial assistance in form of major research project to NK.

References

- Abd El Aty, A. A., Wehaidy, H. R., and Mostafa, F. A. (2014). Optimization of inulinase production from low cost substrates using Plackett-Burman and Taguchi methods. *Carb. Pol.* **102**: 261-268.
- Abe, H., Fujita, Y., Takaoka, Y., Kurita, E., Yano, S., Tanaka, N., and Nakayama, K. I. (2009). Ethanol tolerant *Saccharomyces cerevisiae* strains isolated under selective conditions by over-expression of a proofreading deficient DNA polymerase δ . *J. Biosci. Bioeng.* **108**: 199-204.
- Adriana, C., Flores, G., Jesus, M. C., Cristobal, N. A., and Raul, R. H. (2012). Inulinase production by a Mexican semi desert xerophilic *Penicillium citrinum* strain under submerged culture. *J. Food Sci. Technol.* **4**: 46-50.
- Altunbas, C., Uygun, M., Uygun, A. D., Akgol, S., and Denizli, A. (2013). Immobilization of inulinase on Concanavalin A attached super macroporous cryogel for production of high fructose syrup. *Appl. Biochem. Biotechnol.* **170**: 1909-1921
- Astolfi, V., Joris, J., Verlindo, R., Oliveira, J. V., Maugeri, F., Mazutti, M. A., Oliveira, D. D., and Treichel, H. (2011). Operation of a fixed bed bioreactor in batch and fed batch modes for production of inulinase by solid state fermentation. *Bioche. Eng. J.* **58**:39-49.
- Ayyachamy, M., Khelawan, K., Pillay, D., Permaul, K., and Singh, S. (2007). Production of inulinase by *Xanthomonas campestris* pv. *phaseoli* using onion (*Allium cepa*) and garlic (*Allium sativum*) peels in solid state cultivation. *Lett. Appl. Microbiol.* **45**: 439-444.

- Bali, V., Panesar, P. S., Bera, M. B., and Penesar, R. (2015). Fructooligosaccharides: Production, purification and potential applications. *Crit. Rev. Food Sci. Nutri.* **55**: 1475-1490.
- Barclay, T., Ginic-Markovic, M., Cooper, P., and Petrovsky, N. (2010). Inulin: A versatile polysaccharide with multiple pharmaceutical and food chemical uses. *J. Excipie. Food Chemi.* **1**: 27–50.
- Bonciu, C.N., Constantin O., and Bahrim, G. (2012). Increase in extracellular inulinase production for a new *Rhizoctonia* sp. strain by using buckwheat (*Fagopyrum esculentum*) flour as a single carbon source. *Lett. Appl. Microbiol.* **55**: 195-201.
- Cao, T. S., Wang, G. Y., Chi, Z., Wang, Z. P., Chi, Z. M. (2013). Cloning, characterization and heterologous expression of the INU1 gene from *Cryptococcus aureus* HYA. *Gene.* **516**: 255-262.
- Chen, H. Q., Chen, X. M., Li, Y., Jin, Z. Y., Xu, X. M., Zhao, J. W., Chen, T. X., and Xie, Z. J. (2009). Purification and characterization of exo- and endoinulinase from *Aspergillus ficuum* JNSP5-06. *Food Chem.* **115**: 1206-1212.
- Chen, H., Chen, X., Chen T., Xu, X., and Jin, Z. (2011). Extraction optimization of inulinase obtained by solid state fermentation of *Aspergillus ficuum* JNSP5-06. *Carb. Pol.* **85**: 446-451.
- Chen, M., Lei, X., Chen, C., Zhang, S., Xie, J., and Wei, D. (2015). Cloning, overexpression and characterization of a highly active endoinulinase gene from *Aspergillus fumigatus* Cl1 for production of inulo-oligosaccharides. *Appl. Biochem. Biotechnol.* **175**:1153-1167.

- Chen, X. M., Xu, X. M., Jin, Z.Y. and Chen, H. Q. (2013). Expression of an exoinulinase gene from *Aspergillus ficuum* in *Escherichia coli* and its characterization. *Carb. Pol.* **92**: 1984-1990.
- Chen, X.M., Xu, X.M., Jin, Z.Y., and Chen, H.Q. (2012). Expression of an endoinulinase gene from *Aspergillus ficuum* JNSP 5-06 in *Escherichia coli* and its characterization. *Carb. Pol.* **88**: 748-753.
- Chesini, M., Neila, L. P., Parra, D. F. D. L., Rojas, N. L., Esquivel, J. C. C., Cavalitto, S. F., Ghiringhelli, P. D., and Hours, R. A. (2013). *Aspergillus kawachii* produces an inulinase in culture with yacon (*Smallanthus sonchifolius*) as substrate. *Electro. J. Biotechnol.* **16**: DOI: 10.2225/vol16-issue3-fulltext-13.
- Chi, Z. M., Chi, Z., Liu, G. L., and Yue, L. (2009). Inulinase expressing microorganisms and application of inulinases. *Appl. Microbiol. Biotechnol.* **82**: 211-220.
- Chi, Z., Zhang, T., Cao, T. S., Liu, X. Y., Cui, W., and Zhao, C. H. (2011). Biotechnological potential of inulin for bioprocess. *Bioresour. Technol.* **102**: 4295-4303.
- Cui, W., Wang, Q., Zhang, F., Zhang, S.C., Chi, Z.M., and Madzak, C. (2011). Direct conversion of inulin into single cell protein by the engineered *Yarrowia lipolytica* carrying inulinase gene. *Proc. Biochem.* **46**:1442-1448.
- Dilipkumar, M., Rajamohan, N., and Rajsimman, M. (2013). Inulinase production in a packed bed reactor by solid state fermentation. *Carb. Pol.* **96**: 196-199.
- Dilipkumar, M., Rajsimman, M., and Rajamohan, N. (2010). Optimization of inulinase production using copra waste by *Kluyveromyces marxianus* var. *marxianus*. *Chem. Indus. Chem. Eng. Quarterly.* **16**: 319-327.

- Dilipkumar, M., Rajsimman, M., and Rajamohan, N. (2011). Optimization of media component in inulinase production using garlic by *Penicillium rugulosum*. *J. Advan. Scien. Res.* **2**: 24-33.
- Dilipkumar, M., Rajsimman, M., and Rajamohan, N. (2014a). Optimization, kinetics, and modeling of inulinase production by *K. marxianus* var. *marxianus*. *Prep. Biochem. Biotechnol.* **44**: 291-309.
- Dilipkumar, M., Rajsimman, M., and Rajamohan, N. (2014b). Utilization of copra waste for the solid state fermentative production of inulinase in batch and packed bed reactor. *Carb. Pol.* **102**: 662-668.
- Dinarvand, M., Rezaee, M., Masomian, M., Jazayeri, S. D., Zareian, M., Abbasi, S., and Ariff, A. B. (2013). Effect of C/N ratio and media optimization through response surface methodology on simultaneous productions of intra- and extracellular inulinase and invertase from *Aspergillus niger* ATCC 20611. *Biomed Resea. Internat.* doi.org/10.1155/2013/508968.
- Ettalibi, M., and Baratti, J. C. (2008). Sucrose hydrolysis by thermostable immobilized inulinases from *Aspergillus ficuum*. *Enz. Microbial Technol.* **28**: 473-478.
- Fernandez-Arrojo, L., Santos-Moriano, P., Rodriguez-Colinas, B., Ballesteros, A. O., and Plou, F. J (2015). Micro-scale procedure for enzyme immobilization screening and operational stability assays. *Biotechnol. Lett.* **37**: 1593-600.
- Flores-Gallegos, A. C., Morlett-Chavez, J. A., Aguilar, C. N., Riutort, M., Rodriguez-Herrera R. (2015). Gene encoding inulinase isolated from *Penicillium citrinum* ESS and its molecular phylogeny. *Appl. Biochem. Biotechnol.* **175**: 1358-1370.

- Ganaie, M.A., Rawat, H.K., Wani, O.A., Gupta, U.S., Kango, N. (2014). Immobilization of fructosyltransferase by chitosan and alginate for efficient production of fructooligosaccharides. *Proc. Biochem.* **49**: 840-844.
- Gao, J., Xu, H., Li, Q. J., Feng, X.H., and Li, S. (2010). Optimization of medium for one-step fermentation of inulin extract from Jerusalem artichoke tubers using *Paenibacillus polymyxa* ZJ-9 to produce R, R-2, 3-butanediol. *Bioresour. Technol.* **101**: 7076–7082.
- Gao, J., Yuan, W., Li, Y., Xiang, R., Hou, S., Zhong, S., and Bai, F. (2015). Transcriptional analysis of *Kluyveromyces marxianus* for ethanol production from inulin using consolidated bioprocessing technology. *Biotechnol. Biofuels.* **14**: 98-115.
- Gao, L., Chi, Z., Sheng, J., Wang, L., Li, J., and Gong, F. (2007). Inulinase-producing marine yeasts: evaluation of their diversity and inulin hydrolysis by their crude enzymes. *Microb. Ecol.* **54**: 722-729.
- Gao, W., Bao, Y., Liu, Y., Zhang, X., Wang, J., and An, L. (2009). Characterization of thermostable endoinulinase from a new strain *Bacillus smithii* T7. *Appl. Biochem. Biotechnol.* **157**: 498-506.
- Ge, X.Y., Qian, H., and Zhang, W.G. (2009). Improvement of L-lactic acid production from Jerusalem artichoke tubers by mixed culture of *Aspergillus niger* and *Lactobacillus* sp. *Biores. Technol.* **100**: 1872-1874.
- Golunski, S., Astolfi, V., Carniel, N., Oliveira D.de, Marco Di Luccio, Mazutti. M. A. and Treichel, H. (2011). Ethanol precipitation and ultrafiltration of inulinases from *Kluyveromyces marxianus*. *Sepa. Purifi. Technol.* **78**: 261-265.

- Gong, F., Shen, J., Chi, Z., and Li, J. (2007). Inulinase production by a marine yeast *Pichia guilliermondii* and inulin hydrolysis by the crude inulinase. *J. Indu. Microbiol. Biotechnol.* **34**: 179-185.
- He, M., Wu, D., Wu, J., and Chen, J. (2014). Enhanced expression of endoinulinase from *Aspergillus niger* by codon optimization in *Pichia pastoris* and its application in inulo-oligosaccharide production. *J. Indus. Microbiol. Biotechnol.* **41**: 105-114.
- Housseiny, M. M. (2014). Production of an endoinulinase from *Aspergillus niger* AUMC 9375, by solid state fermentation of agricultural waste, with purification and characterization of the free and immobilized enzyme. *J. Microbiol.* **52**: 389-398.
- Hu, N., Yuan, B., Sun, J., Wang, S.A., and Li, F.L. (2012). Thermotolerant *Kluyveromyces marxianus* and *Saccharomyces cerevisiae* strains representing potentials for bioethanol production from Jerusalem artichoke by consolidated bioprocessing. *Appl. Microbiol. Biotechnol.* **95**: 1359-1368.
- Jain, S. C., Jain, P. C. and Kango, N. (2012). Production of inulinase from *Kluyveromyces marxianus* using dahlia tuber extract. *Braz. J. Microbiol.* **43**: 62-69.
- Kango, N. (2008). Production of inulinase using tap roots of dandelion (*Taraxacum officinale*) by *Aspergillus niger*. *J. Food Eng.* **85**: 473-478.
- Kango, N., and Jain, S. C. (2011). Production and properties of microbial inulinases: Recent advances. *Food Biotechnol.* **25**:165-212.
- Kelly, G. (2009). Inulin-type prebiotics- A review: Part 2. *Alternative Medi. Rev.* **14**: 36-55.

- Kim, H. C., Kim, H. J., Choi, W. B. and Nam, S. W. (2008). Inulooligosaccharide production from inulin by *Saccharomyces cerevisiae* strain displaying cell-surface endoinulinase. *J. Microbiol. Biotechnol.* **16**: 360-367.
- Kobayashi, T., Uchimura, K., Deguchi, S., and Horikoshi, K. (2012). Cloning and sequencing of inulinase and β -fructofuranosidase genes of a deep-sea *Microbulbifer* species and properties of recombinant enzymes. *Appl Environ. Microbiol.* **78**: 2493-2495.
- Kuhn, G. D. O., Rosa, C. D., Silva, M. F., Treichel, H., Oliveira, D. D., and Oliveira, J. V (2012). Synthesis of fructooligosaccharides from *Aspergillus niger* commercial inulinase immobilized in montmorillonite pretreated in pressurized propane and LPG. *Appl. Biochem. Biotechnol.* **169**: 750-760.
- Kuntz, M. G. F., Fiates, G. M. R., and Teixeira, E. (2013). Characteristics of prebiotic food products containing inulin. *British. Food. J.* **115**: 235-251.
- Kuzuwa, S., Yokoi, K. J., Kondo, M., Kimoto, H., Yamakawa, A., Taketo, A., and Kodaira, K. L. (2012). Properties of the inulinase gene *levH1* of *Lactobacillus casei* IAM 1045; cloning, mutational and biochemical characterization. *Gene.* **495**: 154-162.
- Lane, M. M., and Morrissey, J. P. (2010). *Kluyveromyces marxianus*: a yeast emerging from its sister's shadow. *Fungal Biol.* **24**: 17-26.
- Li, M., Liu, G. L., and Chi, Z. M. (2010a). Single cell oil production from hydrolysates of cassava starch by marine derived yeast *Rhodotorula mucilaginosa* TJY15a. *Biomass Bioene.* **4**: 101-107.

- Li, D., Dai, J. Y. and Xiu, Z. L. (2010b). A novel strategy for integrated utilization of Jerusalem artichoke stalk and tuber for production of 2, 3-butanediol by *Klebsiella pneumonia*. *Bioresour. Technol.* **101**: 8342-8347.
- Li, L., Chen, C., Li, K., Wang, Y., Gao, C., Ma, C., and Xu, P. (2014). Efficient simultaneous saccharification and fermentation of inulin to 2, 3-butanediol by thermophilic *Bacillus licheniformis* ATCC 14580. *Appl. Environ. Microbiol.* **80**: 6458-6464.
- Li, S. Z., Chan-Halbrendt, S. Z. (2009). Ethanol production in (the) People's Republic of China: potential and technologies. *Appl. Energ.* **86**: 162-169.
- Li, Y, Liu, G. L., and Chi Z. M. (2013). Ethanol production from inulin and unsterilized meal of Jerusalem artichoke tubers by *Saccharomyces* sp. W0 expressing the endoinulinase gene from *Arthrobacter* sp. *Bioresour. Technol.* **147**: 254-259.
- Li, Y., Liu, G-L., Wang, K., Chi, Z-M. and Madzak, C. (2012). Overexpression of the endoinulinase gene from *Arthrobacter* sp. S37 in *Yarrowia lipolytica* and characterization of the recombinant endoinulinase. *J. Mol. Catal. B: Enz.* **74**: 109-115.
- Lim, S. H., Ryu, J. M., Lee, H., Jeon, J. H., Sok, D. E., and Choi, E. S. (2010). Ethanol fermentation from Jerusalem artichoke powder using *Saccharomyces cerevisiae* KCCM50549 without pretreatment for inulin hydrolysis. *Bioresour. Technol.* **102**: 2109-2111.
- Liu, G. L., Chi, Z., and Chi, Z. M. (2013). Molecular characterization and expression of microbial inulinase genes. *Crit. Rev. Microbiol.* **39**: 152-165.

- Liu, G. L., Fu, G.Y., Chi, Z. and Chi, Z. M. (2014). Enhanced expression of the codon-optimized *exo*-inulinase gene from the yeast *Meyerozyma guilliermondii* in *Saccharomyces* sp. W0 and bioethanol production from inulin. *Appl. Microbiol. Biotechnol.* **98**: 9129-9138.
- Liu, Y.X., Chi, Z., Liu, GL., Wang, F., Madzak C., and Chi Z. M. (2010). Inulin hydrolysis and citric acid production from inulin using the surface engineered *Yarrowia lipolytica* displaying inulinase. *Metabol. Eng.* **12**: 469-476.
- Liu, G. L., Wang, D. S., Wang, L. F., Zhao, S. F., Chi, Z.M. (2011). Mig1 is involved in mycelia formation and expression of the genes encoding extracellular enzymes in *Saccharomycopsis fibuligera* A11. *Fungal Genet. Biol.* **48**: 904-913.
- Lu, W. D., Li, A. X., and Guo, Q. L. (2013). Production of novel alkalitolarant and thermostable inulinase from marine actinomycete *Nocardiopsis* sp. DN-K15 and inulin hydrolysis by the enzyme. *Ann. Microbiol.* **64**: 441-449.
- Ma, J. Y., Cao, H. L., Tan, H. D., Hu, X. J., Liu, W. J., Du, Y. G., and Yin, H. (2015). Cloning, expression, characterization, and mutagenesis of a thermostable exoinulinase from *Kluyveromyces cicerisporus*. *Appl. Biochem. Biotechnol.* **178**: 144-158.
- Makino, Y., Treichel, H., Mazutti, M. A., Maugeria, F., and Rodrigues, M. I. (2009). Inulinase bio-production using agroindustrial residues: screening of microorganisms and process parameters optimization. *J. Chem. Technol. Biotechnol.* **84**: 1056-1062.
- Mansouri, M., Houbraken, J., Samson, R. A., Frisvad, J. C., Christensen, M., Tuthill, D. E., Koutaniemi, S., Hatakka, A., and Lankinen, P. (2013). *Penicillium subrubescens*, a new species efficiently producing inulinase. *Antonie van Leeuwenhoek.* **103**: 1343-1357.

- Mazutti, M.A., Corazza, M.L., Maugeri, F., Rodrigues, M.I., Oliveira, J.V., Treichel, H., and Corazza, F.C. (2010a). Hybrid modeling of inulinase bioproduction process. *J. Chem. Tech. Biotech.* **85**: 512-519.
- Mazutti, M. A., Skrowonski, A., Boni, G., Zabet, G. L., Silva, M. F., de Oliveira, D., Luccio, M. D., Filho, F. M., Rodriguez, M. I., and Treichel, H. (2010b). Partial characterization of inulinases obtained by submerged and solid-state fermentation using agroindustrial residues as substrates: a comparative study. *Appl. Biochem. Biotechnol.* **160**: 682-693.
- Moriyama, S., Muguruma, M., Ohta, K. (2006). Quantitative expression analysis of inulinase gene cluster of *Penicillium* sp. strain TN-88. *J. Biosci. Bioeng.* **101**: 277-279.
- Mughal, M. S., Ali, S., Ashiq, M., and Talish, A. S. (2009). Kinetics of an extracellular exo-inulinase production from a 5-fluorocytosine resistant mutant of *Geotrichum candidum* using two-factorial design. *Bioresour. Technol.* **100**: 3657-3662.
- Mutanda, T., Mokoena, M. P., Olaniran, a. O., Wilhelmi, B. S., and Whiteley, C. G. (2014). Microbial enzymatic production and applications of short-chain fructooligosaccharides and inulooligosaccharides: Recent advances and current perspectives. *J. Indus. Microbiol. Biotechnol.* **41**: 893-906.
- Mutanda, T., Wilhelmi, B. S., and Whiteley, C. G. (2008). Response surface methodology: synthesis of inulooligosaccharides with an endoinulinase from *Aspergillus niger*. *Enzy. Microbial. Technology.* **43**: 363-368.
- Mutanda, T., Wilhelmi, B., and Whiteley, C. G. (2009). Controlled production of fructose by an exoinulinase from *Aspergillus ficuum*. *Appl. Biochem. Biotechnol.* **159**: 65-77.

- Naidoo, K., Ayyachamy, M., Permaul, K., and Singh, S. (2009). Enhanced fructooligosaccharides and inulinase production by *Xanthomonas campestris* pv. *phaseoli* KM 24 mutant. *Bioproc. Biosys. Eng.* **32**: 689-695.
- Nguyen, Q. D., Rezessy-Szabo, J. M., Czukor, B., and Hoschke, A. (2011). Continuous production of oligofructose syrup from Jerusalem artichoke juice by immobilized endoinulinase. *Proc. Biochem.* **46**: 298-303.
- Nguyen, Q. D., Sujtő, N. M., Bujna, E., Hoschke Á., and Rezessy-Szabó J. M. (2013). Effects of medium composition and process parameters on the production of extracellular inulinase by *Thermomyces lanuginosus*. *Food Technol. Biotechnol.* **51**: 36-44.
- Paixao, S.M., Teixeira, P.D., Silva, T.P., Teixeira, A.V., and Alves L. (2013). Screening of novel yeast inulinases and further application to bioprocesses. *New Biotechnol* **30**: 598-606.
- Petrova, P., Velikova, P., Popova, L., and Petrov, K. (2015). Direct conversion chicory flour into L (+)-lactic acid by the highly effective inulinase producer *Lactobacillus paracasei* DSM 23505. *Bioresour. Technol.* **186**: 329-333.
- Rawat, H. K., Ganaie, M. A., and Kango, N. (2015a). Production of inulinase, fructosyltransferase and sucrase from fungi on low-value inulin-rich substrates and their use in generation of fructose and fructooligosaccharides. *Antonie van Leeuwenhoek.* **107**: 799-811.
- Rawat, H. K., Jain, S. C., and Kango, N. (2015b). Production and properties of inulinase from *Penicillium* sp. NFCC 2768 grown on inulin containing vegetal infusions. *Biocata. Biotrans.* **33**: 61-68.

- Ricca, E., Calabro, V., Curcio, S., and Lorio, G. (2009). Fructose production by chicory inulin enzymatic hydrolysis: a kinetic study and reaction mechanism. *Proc. Biochem.* **44**: 466-470.
- Ronkart, S. N., Blecker, C. S., Fourmanoir, H., Fougnes, C., Deroanne, C., Van Herck, J.-C., et al. (2007). Isolation and identification of inulooligosaccharides resulting from inulin hydrolysis. *Analy. Chim. Acta.* **604**: 81-87.
- Saber, W. I. A., and El-Naggar, N. E. (2009). Optimization of fermentation conditions for the biosynthesis of inulinase by the new source: *Aspergillus tamari* and hydrolysis of some inulin containing agro waste. *Biotechnol.* **8**: 425-433.
- Schmitt, E., Bura, R., Gustafson, R., Cooper, J. and Vajzovic, A. (2012). Converting lignocellulosic solid waste into ethanol for the State of Washington: an investigation of treatment technologies and environmental impacts. *Bioresour. Technol.* **104**: 400-409.
- Squarezi, C., Longo, C., Ceni, G., Boni, G., Silva, M. F., Luccio, M., Mazutti, M. A., Maugeri, F., Rodrigues, M. I., and Treichel, H. (2009). Inulinase production by agroindustrial residues: optimization of pretreatment of substrates and production medium. *Food Bioproc. Technol.* **2**: 409-414.
- Sheng, J., Chi, Z., Li, J., Gao, L., and Gong, F. (2007). Inulinase production by the marine yeast *Cryptococcus aureus* G7a and inulin hydrolysis by crude inulinase. *Proc. Biochem.* **42**: 805-811.
- Singh, R. S., Singh, R. P., Kennedy J. F. (2016). Recent insights in enzymatic synthesis of fructooligosaccharides from inulin *Inter J. Biol. Macromol.*
doi:10.1016/j.ijbiomac.2016.01.026

- Silva, M. F., Rigo, D., Mossi, V., Golunski, S., Kuhn, G. D. O., Di Luccio, M., et al. (2013). Enzymatic synthesis of fructooligosaccharides by inulinases from *Aspergillus niger* and *Kluyveromyces marxianus* NRRL Y-7571 in aqueous-organic medium. *Food Chem.* **138**: 148-153.
- Singh, R. S., and Bhermi, H. K. (2008). Production of extracellular exoinulinase from *Kluyveromyces marxianus* YS-1 using root tubers of *Asparagus officinalis*. *Bioresour. Technol.* **99**: 7418-7423.
- Singh, R. S., and Singh, R. P. (2014). Response surface optimization of endoinulinase production from a cost effective substrate by *Bacillus safensis* AS-08 for hydrolysis of inulin. *Biocat. Agricul. Biotechnol.* **3**: 365-372.
- Singh, R. S., Dhaliwal, R., and Puri, M. (2006). Production of inulinase from *Kluyveromyces marxianus* YS-1 using root extract of *Asparagus racemosus*. *Proc. Biochem.* **41**: 1703-1707.
- Sirisansaneeyakul, S., Worawuthiyanan, N., Vanichsriratana, W., Srinophakun, P. and Chisti, Y. (2007). Production of fructose from inulin using mixed inulinases from *Aspergillus niger* and *Candida guilliermondii*. *Worl. J. Microbiol. Biochnol.* **23**: 543-552.
- Songpim, M., Vaithanomsat, P., Vanichsriratana, W., and Sirisansaneeyakul, S. (2011). Enhancement of inulinase and invertase production from a newly isolated *Candida guilliermondii* TISTR 5844. *Kasetsart J.* **45**: 675-685.
- Treichel, H., Mazutti, M. A., Maugeri, F., and Rodrigues, M. I. (2009). Use of a sequential strategy of experimental design to optimize the inulinase production in a batch bioreactor. *J. Ind. Microbiol. Biotechnol.* **36**: 895-900.

- Trivedi, S., Divecha, J., and Shah, A. (2012). Optimization of inulinase production by newly isolated *Aspergillus tubingensis* CR16 using low cost substrates. *Carb. Pol.* **90**: 483-490.
- Trytek, M., Fiedurek, J., Podkoscielna, B., Gawdzik, B., and Skowronek M. (2015). An efficient method for the immobilization of inulinase using new types of polymers containing epoxy groups. *J. Indus. Microbiol. Biotechnol.* **42**: 985-996.
- Tsurumaki, M., Kotake, M., Iwasaki, M., Saito, M., Tanaka, K., Aw, W., Fukudu, S., and Tomita, M. (2015). The application of omics technologies in the functional evaluation of inulin and inulin-containing prebiotics dietary supplementation. *Nutr. Diabetes.* **30**: doi: 10.1038/nutd.2015.35
- Vijayaraghavan, K., Yamini, D., Ambika, V., and Sowdamini, N. S. (2009). Trends in inulinase production- a review. *Criti. Rev. Biotechnol.* **29**: 67-77.
- Waleckx, E., Mateos-Diaz, J.C., Gschaedler, A., Colonna-Ceccaldi, B., Brin, N., Garcia-Quezada, G., Villanueva-Rodriguez, S., Monsan, P. (2011). Use of inulinases to improve fermentable carbohydrate recovery during tequila production. *Food Chem.* **124**: 1533-1542.
- Wang, G.Y., Chi, Z., Song, B., Wang, Z.P., Chi, Z.M. (2012). High level lipid production by a novel inulinase-producing yeast *Pichia guilliermondii* Pcl22. *Bioresour. Technol.* **124**: 77-82.
- Wang, L., Huang, Y., Long, X., Meng, X., and Liu, Z. (2011). Cloning of exoinulinase gene from *Penicillium janthinellum* strain B01 and its high-level expression in *Pichia pastoris*. *J. Appl. Microbiol.* **111**: 1371-1380.

- Wang, Z. P., Fu, W. J., Xu, H. M., and Chi, Z. M. (2014). Direct conversion of inulin into cell lipid by an inulinase-producing yeast *Rhodospiridium toruloides* 2F5. *Bioresour. Technol.* **161**: 131-136.
- Wu, S., Zhao, X., Shen, H., Wang, Q., and Zhao, Z. K. (2011). Microbial lipid production by *Rhodotorula toruloides* under sulfate limited conditions. *Bioresour. Technol.* **102**: 1803-1807.
- Yu, J., Jiang, J., Fang, Z., Li, Y., Lv, H., and Liu, J. (2010). Enhanced expression of heterologous inulinase in *Kluyveromyces lactis* by disruption of hap1 gene. *Biotechnol. Lett.* **32**: 507-512.
- Yu, X., Guo, N., Chi, Z., Gong, F., Sheng, J., and Chi, Z. (2009). Inulinase overproduction by a mutant of the marine yeast *Pichia guilliermondii* using surface response methodology and inulin hydrolysis. *Biochem. Eng. J.* **43**: 266-271.
- Yuan, B., Wang, S. A., and Li, F. L. (2013). Improved ethanol fermentation by heterologous endoinulinase and inherent invertase from inulin by *Saccharomyces cerevisiae*. *Bioresour. Technol.* **139**: 402-405.
- Yuan, X. L., Roubos, J. A., van den Hondel, C.A.M.J.J., Ram, A.F.J. (2008). Identification of InuR, a new Zn(II)₂Cys₆ transcriptional activator involved in the regulation of inulinolytic genes in *Aspergillus niger*. *Mol. Genet. Genom.* **279**: 11-26.
- Yuan, X. L., Goosen, C., Kools, H., van der Maarel, M. J., van den Hondel, C. A., Dijkhuizen, L., Ram, A. F. (2006). Database mining and transcriptional analysis of genes encoding inulin-modifying enzymes of *Aspergillus niger*. *Microbiol.* **152**: 3061-3073.

- Yuan, W. J., Chang, B. L., Ren, J. G., Liu, J. P., Bai, F. W., and Li, Y. Y. (2012). Consolidated bioprocessing strategy for ethanol production from Jerusalem artichoke tubers by *Kluyveromyces marxianus* under high gravity conditions. *J. Appl. Microbiol.* **112**: 38-44.
- Zhang, S., Yang, F., Wang, Q., Hua, Y., and Zhao, Z. K. (2012). High-level secretory expression and characterization of the recombinant *Kluyveromyces marxianus* inulinase. *Proc. Biochem.* **47**: 151-155.
- Zhang T, Chi Z, Chi Z, Parrou JL, Gong F. (2010a). Expression of the inulinase gene from the marine-derived *Pichia guilliermondii* in *Saccharomyces sp.* W0 and ethanol production from inulin. *Microb. Biotechnol.* **3**: 576-582.
- Zhang, T., Chi, Z., Zhao, C. H., Chi, Z. M., and Gong, F. (2010b). Bioethanol production from hydrolysates of inulin and the tuber meal of Jerusalem artichoke by *Saccharomyces sp.* W0. *Bioresour. Technol.* **10**: 8166-8170.
- Zhang, T., Chi, Z. M., Sheng, J. (2009a). A highly thermosensitive and permeable mutant of the marine yeast *Cryptococcus aureus* G7a potentially useful for single-cell protein production and its nutritive components. *Mar. Biotechnol.* **11**: 280-286.
- Zhang, T., Gong, F., Chi, Z., Liu, G., Chi, Z., Sheng, J., Li, J., and Wang, X. (2009b). Cloning and characterization of the inulinase gene from a marine yeast *Pichia guilliermondii* and its expression in *Pichia pastoris*. *Antonie van Leeuwenhoek.* **95**: 13-22.
- Zhao, C. H., Chi, Z. Zhang, F., Guo, F. J., Li, M., Song, W. B., and Chi, Z. M. (2011). Direct conversion of inulin and extract of tubers of Jerusalem artichoke into single cell oil by co-cultures of *Rhodotorula mucilaginosa* TJY15a and immobilized inulinase-producing yeast cells. *Bioresour. Technol.* **102**: 6128-6133.

- Zhao, C. H., Cui, W., Liu, X. Y., Chi, Z. M., and Madzak, C. (2010a). Expression of inulinase gene in the oleaginous yeast *Yarrowia lipolytica* and single cell oil production from inulin containing materials. *Metab. Eng.* **12**: 510-517.
- Zhao, C. H., Zhang, T., Li, M., and Chi, Z. M. (2010b). Single cell oil production from hydrolysates of inulin and extracts of tubers of Jerusalem artichoke by *Rhodotorula mucilaginosa* TJY15a. *Proc. Biochem.* **45**: 1121-1126.
- Zhao, X., Wu, S., Hu, C., Wang, Q., Hua, Y., and Zhao, Z. K. (2010c). Lipid production from Jerusalem artichoke by *Rhodotorula toruloides* Y4. *J. Indus. Microbiol. Biotechnol.* **37**: 581-585.
- Zhou, J., Lu, Q., Peng, M., Zhang, R., Mo, M., Tang, X., Li J, Xu, B., Ding, J., and Huang, Z. (2014a). Cold active and NaCl-tolerant exoinulinase from a cold adapted *Arthrobacter* sp. MN8 and its potential for use in the production of fructose at low temperatures. *Biosci Bioeng.* **119**: 267-274.
- Zhou, J. P., Gao, Y. J., Zhang, R., Mo, M. H., Tang, X. H., Li, J. J., Xu, B., Ding, J. M., and Huang, Z. X. (2014b). A novel low temperature-active exo-inulinase identified based on molecular activity strategy from *Sphingobacterium* sp. GN25 isolated from feces of *Grus nigricollis*. *Proc. Biochem.* **49**: 1656-1663.
- Zhou, H. X., Xin, F. H., Chi, Z., Liu, G.L., Chi, Z. M. (2014c). Inulinase production by the yeast *Kluyveromyces marxianus* with the disrupted MIG1 gene and the over-expressed inulinase gene. *Proc. Biochem.* **49**: 1867-1874.

Zhou, J., Peng, M., Zhang, R., Li, J., Tang, X., Xu, B., Ding, J., Gao, Y., Ren, J., and Huang, Z. (2015). Characterization of *Sphingomonas* sp. JB13 exo-inlinase: a novel detergent-, salt-, and protease-tolerant exo-inulinase. *Extremo*. **19**: 9129-9138.

Table 1 Production of inulinase by microorganisms

SN	Microorganisms	Carbon source ^a	Nitrogen source ^b	Temp (°C)	pH	RPM	Time (h)	Activity (U/ml)	Reference
1	<i>Aspergillus niger</i>	DTE	YE	28	-	150	72	12.2	Rawat et
	GNCC 2655	DRE						4.07	al., 2015a
	<i>A. niger</i> GNCC	ARE						8.21	
2	<i>Penicillium</i> sp.	DTE	YE	30	5.0	150	120	3.87	Rawat et
3	<i>A. niger</i> AUMC	ST, LR	YE	30	5.0	-	168	290-	Housseiny,
4	<i>Penicillium subrubescense</i>	JA	-	28	6.0	180	120	7.7	Mansouri et al., 2013
5	<i>Aspergillus</i>	Yacon	-	30	-	250	68	80	Chesini et
6	<i>A. niger</i> ATCC	Suc	ME	30	6.5	150	96	319	Dinarvand
7	<i>Kluyveromyces</i>	DTE	YE	28	4.0	120	96	17.9	Jain et al.,
8	<i>K. marxianus</i>	SCB,	SB, CSL	30	-	SSF	24	436.7*	Mazutti et
	NRRL Y-7571	SCM							al., 2010a
9	<i>K. marxianus</i>	SCM	YE,	36	5.5	150	72	722*	Makino et
	NRRL Y-7571		CSL						al., 2009
10	<i>K. marxianus</i>	SCM	YE,	-	-	300	-	1317*	Treichel et
	NRRL Y-7571		CSL						al., 2009
11	<i>K. marxianus</i>	SCB	YE,	-	-	-	-	1139*	Squarezi et
			CSL						al., 2009
12	<i>Aspergillus tamaraii</i> AR-IN9	DTE, JA	CSL	35	5.5	-	72	71.97	Saber and Naggar, 2009

	<i>Xanthomonas</i>	Suc	Tryptone	37	7.0	150	18	22.09	Naidoo et
	<i>Bacillus smithi</i>	Inulin	YE,	50	7.0	200	72	135.2	Gao et al.,
	<i>A. niger</i> NK-126	DRE	YE	30	-	150	96	54	Kango,
	<i>K. marxianus</i> YS-1	ARE	BE	30	6.5	200	60	50.2	Singh and Bharmi, 2008
	<i>Cryptococcus</i> <i>aureus</i>	Inulin	YE	28	5.0	170	42	85	Sheng et al., 2007
	<i>Debaryomyces</i> <i>hansenii</i>	Inulin	YE	28	5.0	170	72	52.5	Gao et al., 2007
	<i>X. campestris</i>	Inulin	YE	37	7.0	150	120	17.4	Ayyachamy et al., 2007
	<i>Yarrowia</i> <i>lipolytica</i>	Inulin	YE	28	5.0	170	72	52.3	Gao et al., 2007

^aARE- Asparagus root extract, DTE-Dahlia tuber extract, DRE-Dandelion root extract, JA- Jerusalem artichoke, LT- Lettuce root, SCB- Sugarcane bagasse, SCM- Sugarcane molasses, Suc- Sucrose, ST- Sunflower tuber,; ^bYE- Yeast extract, BE- Beef extract, SB- Soyabean bran, ME- Meat extract, CSL- Corn steep liquor, ADP- Ammonium dihydrogen phosphate, AS- Ammonium sulphate; Production of inulinase using SSF *U/gds

Table 2 Optimization of process parameters for inulinase production

S	Organism	Optimized experimental variables	Substrate	Type of Fermentation	Inulinase Activity	References
	<i>Aspergillus terreus</i>	Artichoke leaves- 3g, KH ₂ PO ₄ - 0.8%, Ca ²⁺ - 6mM	Artichoke leaves	SSF	21.058 U/gds	Abd-El Aty et al., 2014
	<i>Kluyveromyces marxianus</i> var.	CSL- 0.06072 g/gds Urea- 0.01916 g/gds Beef extract-0.00957	Pressmud	SSF	288 U/gds	Dilipkumar et al., 2014a
	<i>Bacillus safensis</i> AS-	Raw inulin-1%, YE - 1%,	Dahlia tubers	SmF	28.67 U/ml	Singh and Singh,
	<i>Penicillium rugulosum</i>	K ₂ HPO ₄ - 0.0047 g/gds, ZnSO ₄ .7H ₂ O- 0.02677 g/gds, Soya bean cake-	Copra waste	SSF	239U/gds	Dilipkumar et al., 2014b
	<i>K. marxianus</i> MTCC 188	Air flow rate- 0.82 L/min,	Pressmud	SSF	300.5 U/gds	Dilipkumar et al.,
	<i>Thermomyces lanuginosus</i>	Jerusalem artichoke extract- 1.8 % (m/v),	Jerusalem Artichoke	SmF	6.09 U/ml	Nguyen et al., 2013

<i>Aspergillus tubingensis</i>	Period of fermentation- 103 h,	Wheat Bran	SSF	257 U/gds	Trivedi et al., 2012
<i>Rhizoctonia</i> sp. I-PN4	Buckwheat- 6% YE- 5.82%	Buckwheat	SmF	1.87 U/ml	Bonciu et al., 2012
<i>Aspergillus ficuum</i> JNSP5-06	Extraction solvent- 0.1 M sodium acetate buffer (pH 4.5)	Wheat bran	SSF	189.02 U/gds	Chen et al., 2011
<i>A. ficuum</i> JNSP5-06	Inulin- 11.47%, NH ₄ H ₂ PO ₄ - 0.76%,	Wheat Bran	SSF	205.63 U/gds	Chen et al., 2011
<i>Candida guilliermondi</i> TISTR 5844	Inulin- 1%, NH ₄ Cl- 2.4%, MgSO ₄ ·7H ₂ O- 1.2%	Inulin	SmF	473.26 U/ml	Songpim et al., 2011
<i>K. marxianus</i>	Ethanol precipitation- 55% (w/w)	Sugarcane Bagasse	SSF	86.1 % Yield	Golunski et al.
<i>Penicillium rugulosum</i>	CSL - 0.05813 g/gds, FeSO ₄ ·7H ₂ O- 0.00011	Garlic Powder	SSF	268 U/gds	Dilipkumar et al.,
<i>K. marxianus</i> NRRL	Inlet temp- 30°C, Air flow rate- 2.2 m ³ /h,	Sugarcane Bagasse	SSF	436.7 J/gds	Mazutti et al., 2010b
<i>K. marxianus</i> MTCC 188	CSL - 0.0560 g/gds, ZnSO ₄ ·7H ₂ O- 0.0084	Copra waste	SSF	372 U/gds	Dilipkumar et al.,
<i>Pichia guilliermondi</i>	Inulin- 20.0 g/L YE - 5.0 g/L	Inulin	SmF	127.7 U/ml	Yu et al., 2009

<i>K. marxianus</i> NRRL Y-7571	Molasses- 250 g/L , CSL – 80 g/L, YE – 6 g/L,			Molasses	SmF	1317 U/ml	Treichel et al., 2009
<i>Geotricum candidum</i>	Incubation period- 48 h, Sucrose conc- 5.0 g/L,			Sucrose	SmF	71.85 U/ml	Mughal et al., 2009
<i>K. marxianus</i> NCYC 587, NRRLY- 7571	Molasse s CSL YE pH	NCYC C 90 g/L 45 g/L 4.0 g/L 4.5	NRR L Y 5-4.5 L 5.7	Molasses	SmF	735 U/ml (NCYC) 722 U/ml (NRRL Y)	Makino et al., 2009
<i>K. marxianus</i>	Sugarcane molasses-			Sugarcane	SmF	1,139	Sguarezi

Table 3 Heterologous expression of inulinase genes

	Source of	Host/vector	Properties of cloned gene	Reference
1	<i>Kluyveromyces cicerisporus</i>	<i>Pichia pastoris</i> X-33	Exoinulinase, MW- 79.5 kDa, Activity- 45.2 U/ml pH- 4.5 Temp- 55°C	Ma et al., 2015
2	<i>Sphingomonas</i> sp. JB 13	<i>Escherichia coli</i> , BL 21 (DE3), <i>pEASY-T1</i> , <i>pEASY-E2</i>	Exoinulinase, MW- 55.6 kDa, Enzyme – rInuAJB13 (408 bp), pH- 5.5, Temp- 55°C Inu Lev Suc Raff Stac K_m (mg/mL) 10 11 6.5 5.9 7.3 V_{max} (μmol/min/mg) 294 22 1666 213 164 K_{cat} (s ⁻¹) 274 21 1552 198 153 K_{cat}/K_m (mLmg/s) 27 2.0 239 34 21 Activator- SDS, CTAB, Tween- 80, Triton X-100, FeSO ₄ , MnSO ₄ ,	Zhou et al., 2015

			β -ME (Retained activity 76.5%), Inhibitors- HgCl_2	
3	<i>Aspergillus fumigatus</i> Cl1	<i>P. pastoris</i> <i>E. coli</i> DH5 α	Endoinulinase, ORF- 1561bp, Code- 507aa MW- 60 kDa, Activity-3860 U/ml pH- 6.0, Temp- 55°C, K_m - 2.18 mM, V_{max} - 1590 $\mu\text{mol/min/mg}$, Metal ion- Activation- MnSO_4 , CaCl_2 , FeSO_4 , Inhibition- AgNO_3 , SDS, EDTA, CuCl_2	Chen et al., 2015
4	<i>Aspergillus niger</i> CICIM F0620	<i>P. pastoris</i> , <i>E. coli</i> JM109	Endoinulinase Gene- <i>EnInu</i> , MW- 59 kDa,	He et al., 2014

5	<i>A. niger</i>	<i>S. cerevisiae</i> JZ1C-J, <i>E. coli</i> DH5 α , pMD19-	Endoinulinase Gene- <i>inuA</i> Activity- 3.1 U/L	Yuan et al., 2013
6	<i>Aspergillus ficuum</i> JNSP5-06	<i>E. coli</i> JM 109 and BL 21 (DE3) pET-28 a (+), PMD 19	Exoinulinase MW- 63 kDa Metal ion- Activation- Mn ²⁺ Inhibition Fe ³⁺ , Fe ²⁺ , Ni ²⁺ , Zn ²⁺ , Mg ²⁺	Chen et al., 2013
7	<i>A. ficuum</i> JNSP5- 06	<i>E. coli</i> JM 109 and BL 21 (DE3) pET-28 a (+),	Endoinulinase ORF- 1482bp MW- 60 kDa Metal ion- Activation- Zn ²⁺	Chen et al., 2012
8	<i>Lactobacillus</i> <i>casei</i> IAM 1045	<i>E. coli</i> XL 1-Blue	Exoinulinase MW- 85 kDa	Kuzuwa et al., 2012
9	<i>Arthrobacter</i> sp. S37	<i>Y. lipolytica</i> pINAI317, pMD 19-T	Endoinulinase MW- 78.9 kDa, Gene- Enl A K_m - 37.1 mg/mL, V_{max} - 3.9	Li et al., 2012
1	<i>Kluyveromyces</i> <i>marxianus</i>	<i>P. pastoris</i> <i>E.</i> <i>coli</i> DH5 α , pMD18-T,	Exoinulinase MW- 60 kDa Gene- <i>rKmInu</i>	Zhang et al., 2012
1	<i>K. cicerisporus</i>	<i>K. lactis</i>	<i>Kcinu</i> Activity- 391 U/ml	Yu et al., 2010

1	<i>K. marxianus</i> CBS 6556	<i>Y. lipolytica</i>	Gene- <i>INU1</i> Activity- 22.6 U/mg	Liu et al., 2010
1	<i>Pichia guilliermondii</i>	<i>P. pastoris</i> X-33	pPICZaA-Inu1 MW- 57.6 kDa	Zhang et al., 2009a
1	<i>P. guilliermondii</i>	<i>P. pastoris</i> X-33	pPICZalphaA ; MW- 60 kDa Activity- 58.7 U/ml	Zhang et al., 2009b

Table 4 Production and properties of thermostable, cold active and halotolerant inulinases

S N	Extremophilic microorganism	Type of enzyme	Extremophilic conditions				Referenc e
			Tem p	pH	Half-life inactivatio n period(τ 1/ 2)	Characteristi cs	
	<i>Sphingomonas</i>	Exo	10 to	5.5	135-163%	MW- 55.6	Zhou et
	<i>Arthrobacter</i> sp. MN8	Exo	0 to 20°C	-	More than 60 min at 50°C, rInuAMN8	Relative activity % at 0°C, 10°C and 20°C is	Zhou et al., 2014a
	<i>Sphingobacteriu</i> <i>m</i> sp. GN25	Exo	0 to 0°C	-	More than 60 min at 45°C	55.8% maximum retained	Zhou, et al., 2014b
	<i>Bacillus</i> <i>licheniformis</i> ATCC 14580	Levanase, inulinase	50°C	-	2,3- butandiol in 30 h	103.0 g/L 2,3 butanediol production	Li et al., 2014
	<i>Nocardiopsis</i> sp. DN-K15	Exo	60°C	8.0	81% active at 60°C for 1h over	25.1 U/ml	Lu et al., 2013

					wide pH 5.0-11.0 range		
	<i>Microbulbifer</i> sp.	Inulinase, Fructofuranosida	0 to 35°C	-	More than 50 min at 50°C	Relative activity % at 0°C-100°C	Kobayas hi et al., 2012
	<i>Bacillus smithii</i> T7	Endo	70°C	4.0 -	9h/ 2.5h at 70/80°C	Inulinase yiel d of 135.2	Gao et al. 2009

Table 5 Use of immobilized inulinase for efficient production of industrially important products

SN	Organism	Immobilization properties	Carrier used	Product formed	Reference
1	<i>Aspergillus niger</i>	Hydrolysis of sucrose,	Sepabeads EC-EA and EC-HA	Glucose	Fernandez-Arrojo et
2	<i>A. niger</i> 20 OSM	Continuous hydrolysis of	Eupergit C	Fructose and Fructooligosaccharide	Trytek et al., 2015
3	<i>A. niger</i> AUMC 9375	Continuous hydrolysis of inulin,	Sodium alginate	Fructooligosaccharide s and Fructose	Housseiny, 2014
4	<i>A. niger</i>	Continuous hydrolysis of inulin, pH - 5.0,	Concanavalin A- attached poly (ethylene glycoldimethacrylate	Fructose syrup	Altunbas et al., 2013
5	<i>A. niger</i> (Fructozyme)	Hydrolysis of inulin and sucrose	Montmorillonite	Fructooligosaccharide s	Kuhn et al., 2012
6	<i>A. niger</i> (Megazyme)	Continuous hydrolysis of	Chitosan	Oligofructose syrup	Nguyen et al., 2011

SN	Microorganism	Biotechnologically important end product ^s	Enzyme type	Fermentable sub
----	---------------	--	-------------	-----------------

Table 6 Potential uses of inulinases for generation of industrially and nutritionally important products

1.	<i>Aspergillus</i> spp., <i>Penicillium</i> spp., <i>Fusarium</i> sp., <i>Arthrini</i> sp.,	F, IOS, FOS	Exo, Endo	Chicory inulin, sucrose
2.	<i>Kluyveromyces marxianus</i>	Ethanol	Inu	JA
3.	<i>Lactobacillus paracasei</i> DSm 23505	LA	Inu	Chicory flour
4.	<i>Penicillium</i> sp. NFCC 2768	F, IOS	Endo, Exo	Dahlia inulin
5.	<i>Meyerozyma guilliermondii</i> ,	Ethanol	Exo	Inulin
6.	<i>Rhodospiridium toruloides</i> 2F5	SCO	Inu	Inulin
7.	<i>Aspergillus niger</i> CICIM F0620	IOS	Endo	Inulin
8.	<i>Arthrobacter</i> sp., <i>Saccharomyces</i> W0*	Ethanol	Endo	JA
9.	<i>A. niger</i> , <i>S. cerevisiae</i> *	Ethanol	Invertase	Inulin
10	<i>Pichia guilliermondii</i> Pcl22	SCO	Inu	Inulin
11	<i>K. marxianus</i>	Ethanol	Inu	JA
12	<i>Aspergillus niger</i> (Fructozyme)	FOS	Inu	Inulin, Sucrose
13	<i>K. marxianus</i> , <i>S. cerevisiae</i> *	Ethanol	Inu	JA
14	<i>K. marxianus</i> , <i>Yarrowia lipolytica</i> *	SCP	Exo	Inulin
15	<i>A. niger</i> (Megazyme)	Oligofructose syrup	Endo	JA
16	<i>Rhodotorula mucilaginosa</i> TJY15a	SCO	Inu	JA
17	Fructozyme L	Tequila	Endo, Exo	<i>Agave tequilana</i>
18	<i>Cryptococcus aureus</i> G7a	SCP	Inu	Yacon
19	<i>K. marxianus</i> , <i>Y. lipolytica</i> *	Citric acid	Exo	Inulin
20	<i>P. guilliermondii</i> , <i>Saccharomyces</i> sp.*	Ethanol	Inu	JA
21	<i>K. marxianus</i> , <i>Y. lipolytica</i> *	SCO	Exo	JA , Inulin
22	<i>Pichia pastoris</i> X-33*, <i>Saccharomyces</i> *	Ethanol	Exo	JA , Inulin
23	<i>Paenibacillus polymyxa</i> ZJ-9	2,3-Butanediol	Inu	JA
24	<i>R. mucilaginosa</i> TJY15a	SCO	Inu	Cassava
25	<i>Klebsiella pneumonia</i>	2,3-Butanediol	Inu	JA
26	<i>S. cerevisiae</i> KCCM50549	Ethanol	Inu	JA

27	<i>A. niger</i> SL-09, <i>Lactobacillus</i> sp.G-02	LA	Inu	JA
28	<i>A. niger</i> NK 126	F, IOS	Exo, Endo	Chicory inulin
29	<i>A. niger</i>	IOS	Endo	Inulin
30	<i>K. marxianus</i> YS-1	F	Exo	<i>Asparagus officinalis</i>

[§]F- Fructose; IOS- Inulooligosaccharides; FOS- Fructooligosaccharides; SCP- Single cell protein;

LA- Lactic acid; SCO- Single cell oil;

CA- Citric acid; Exo- Exoinulinase; Endo- Endoinulinase; Inu- Inulinase; JA- Jerusalem

artichoke; * Cloned or expressed or recombinant inulinase