



## The impact of the micronutrient iodine in health and diseases

Ma. Cecilia Opazo , Irenice Coronado-Arrázola , Omar P. Vallejos , Rodrigo Moreno-Reyes , Carlos Fardella , Lorena Mosso , Alexis M. Kalergis , Susan M. Bueno & Claudia A. Riedel

To cite this article: Ma. Cecilia Opazo , Irenice Coronado-Arrázola , Omar P. Vallejos , Rodrigo Moreno-Reyes , Carlos Fardella , Lorena Mosso , Alexis M. Kalergis , Susan M. Bueno & Claudia A. Riedel (2020): The impact of the micronutrient iodine in health and diseases, Critical Reviews in Food Science and Nutrition, DOI: [10.1080/10408398.2020.1843398](https://doi.org/10.1080/10408398.2020.1843398)

To link to this article: <https://doi.org/10.1080/10408398.2020.1843398>



Published online: 23 Nov 2020.



Submit your article to this journal [↗](#)



Article views: 13



View related articles [↗](#)










View Crossmark data [↗](#)

REVIEW



## The impact of the micronutrient iodine in health and diseases

Ma. Cecilia Opazo<sup>a,b</sup> , Irenice Coronado-Arrázola<sup>c</sup> , Omar P. Vallejos<sup>c</sup> , Rodrigo Moreno-Reyes<sup>d</sup>, Carlos Fardella<sup>e,f,g</sup> , Lorena Mosso<sup>e,f</sup>, Alexis M. Kalergis<sup>c,f</sup> , Susan M. Bueno<sup>c</sup> , and Claudia A. Riedel<sup>a,b</sup> 

<sup>a</sup>Laboratorio de Endocrino-Immunología, Facultad de Ciencias de la Vida, Departamento de Ciencias Biológicas, Universidad Andres Bello, Santiago, Chile; <sup>b</sup>Millennium Institute on Immunology and Immunotherapy, Endocrine-Immunology Laboratory, Facultad de Ciencias de la Vida, Departamento de Ciencias Biológicas, Universidad Andres Bello, Santiago, Chile; <sup>c</sup>Millennium Institute on Immunology and Immunotherapy, Departamento de Genética Molecular y Microbiología, Facultad de Ciencias Biológicas, Pontificia Universidad Católica de Chile, Santiago, Chile; <sup>d</sup>Erasme Hospital, Department of Nuclear Medicine, Université Libre de Bruxelles, Bruxelles, Belgium; <sup>e</sup>Millennium Institute on Immunology and Immunotherapy (IMII), Departamento de Endocrinología, Facultad de Medicina, Pontificia Universidad Católica de Chile, Santiago, Chile; <sup>f</sup>Departamento de Endocrinología, Facultad de Medicina, Pontificia Universidad Católica de Chile, Santiago, Chile; <sup>g</sup>Center for Translational Research in Endocrinology (CETREN-UC), School of Medicine, Pontificia Universidad Católica de Chile, Santiago, Chile

### ABSTRACT

Adequate iodine nutrition is crucial for all mammals by playing his starring role as a component of thyroid hormones, which are key regulators of cellular processes for life such as differentiation, growth, function, and metabolism. Deficiency or excess of iodine in the diet are worldwide highly frequent conditions that are responsible of health problems like hypothyroidism, hypothyroxinemia, goiter, thyroiditis, hyperthyroidism, and autoimmune thyroid diseases among others. The incorporation of iodine in salt or other nutrients resolved the consequences of severe iodine deficiency like goiter, cretinism. However, this strategy in several countries led to other ailments like Hashimoto autoimmune thyroiditis, hyperthyroidism, and hypothyroidism. The goal of this review is to analyze and discuss the different aspects of iodine nutrition for human health comprising its biological role through thyroid hormones, pathogen control, and the regulation of the intestinal microbiota.

### KEYWORDS

Bactericide; intestinal microbiota; iodine; nutrition; thyroid diseases; thyroid hormones

## 1. Introduction

Iodine is a trace element on earth that mammals will incorporate in the chemical form of iodide ( $I^-$ ) (Zbigniew 2017). In this review we will use the term “iodine” for referring to the chemical name of the element (Zbigniew 2017). Adequate iodine nutrition is fundamental for all mammals. In fact, five grams of iodine will be enough to fulfill the needs of a human being for 70 years (Cavalieri 1997). Iodine is incorporated from the diet by the intestinal epithelial cells to the blood (Nicola et al. 2009). Then, iodine is transported from blood into thyroid gland, where the 80% of total iodine in the body is stored and used for the synthesis of thyroid hormones 3,3',5-triiodothyronine ( $T_3$ ) and thyroxine ( $T_4$ ) (Ahad and Ganie 2010; Mansourian 2011). Thyroid hormones are essential for the metabolism of all cells and for the proper fetal development. Besides being iodine a structural and functional component of the thyroid hormones, it also plays an important role as bactericide due to its oxidative properties (McDonnell and Russell 1999). This role is performed in the gastrointestinal tract, where iodine is secreted by epithelial cells of the stomach to the gastric acid and from the salivary glands to the saliva (Josefsson and Ekblad 2009). This review discusses the aspects of the iodine nutrition status worldwide, highlighting its roles for human health and microbiota.

### 1.1. Iodine sources and human iodine nutrition status

#### 1.2. Sources of iodine for the human diet

For humans an adequate iodine nutrition status has been difficult to achieve due to the low availability of this element in the common food. The amount of iodine in natural foods is highly variable and mainly depends on the soil where plants and animals grow (Fuge and Johnson 1986; Teas et al. 2004). The oceans are the main source of iodine (45–60 µg/L, at the surface) (Korzh 1984). For the contrary, the soil has small quantities of iodine and it is considered to be the second most important reservoir for mammals. Soils from coastal areas are more abundant in iodine than soils from the mountains (0.24 µg of iodine/g of soil), given it evaporates from the sea and precipitate in the surrounded areas (Whitehead 1984). Seaweeds have a great capacity to capture and store iodine from the sea and they are the richest iodine foods on earth (Teas et al. 2004). Even though, lettuce and spinach have low iodine content compared to seaweeds, they are considered to be a valuable source of iodine for human nutrition (see Table 1) (Weng et al. 2003; Zhu et al. 2003; Dai et al. 2004; Teas et al. 2004; Pearce et al. 2004; Zealand, Food Standards Australia New 2015; Leung and Braverman 2014; Johnson 2003). The spinach is capable to concentrate iodine proportionally to the available

**Table 1.** Iodide sources.

Source	I <sup>-</sup> content
Edible seaweeds	22–8165 µg/g (Teas et al. 2004)
Bread	0.09–23.31 µg/g (Pearce et al. 2004)
Milk	88–168 µg/250 mL (Pearce et al. 2004)
Eggs	22 µg/g (Zealand, Food Standards Australia New 2015)
Fish	0.73 µg/g (Leung and Braverman 2014)
Leafy plants	89 µg/Kg (Johnson 2003)

iodine in the soil, thus making this ailment a promising candidate for iodine biofortification (Adams et al. 2006; Bath et al. 2017). Moreover, the consumption of goitrogens will reduce the uptake of iodine by the thyroid gland and consequently will decrease the synthesis of thyroid hormones (Eastman and Li 2017). Examples of goitrogens are thioglycosides and cyanoglycosides found in broccoli, maize, and sweet potatoes; and thiocyanates and perchlorate found in cigars, tap-water, and vegetables (Gaitan 1990). Chronic ingestion of plants rich in goitrogens is considered a possible cause of goiter particularly in those regions with higher goiter rates (Bourdoux et al. 1978). Interestingly, Braverman showed with in a long term study that intermittent exposure to goitrogens did not altered thyroid hormone synthesis (Braverman et al. 2005). To overcome the poor iodine ingestion several strategies have been developed worldwide among these are fortification programs aiming to incorporate iodine in common food like milk, bread, water, and salt (Jong 2007; Sarah Davis et al. 2012; Aburto et al. 2014; Olivieri et al. 2017). According to the World Health Organization (WHO) the most successful strategy is the universal salt iodization (USI) program (WHO 2008). In this program, salt is used as a vehicle for iodine administration given that it is consumed fairly by most of the population through the year. The technologies required for this process are of easy implementation, do not affect the iodine organoleptic properties and the iodization process have reasonable costs (WHO 2004). However, after several years of implementation of USI programs in countries that previously suffered of mild or severe iodine deficiency were showing an increment of overt autoimmune hypothyroidism, subclinical hypothyroidism and thyroiditis (Palaniappan et al. 2017; Katagiri et al. 2017; Sun, Shan, and Teng 2014; Marwaha et al. 2012). Health authorities, medical doctors and scientists are reacting, discussing and rethinking new and more appropriate strategies to deal with iodine deficiency and iodine excess. It seems crucial to monitor continually the consumption of iodine in the population by age and gender. This parameter is necessary for taking the right health decisions to avoid the appearance of thyroid diseases due to iodine deficiency or excess.


### 1.3. Iodine nutrition status

Adequate iodine intake is crucial for proper thyroid function, being low iodine intake responsible for the highly frequent iodine deficiency disorders (IDD) (Zimmermann, Jooste, and Pandav 2008). Today and worldwide, more

**Table 2.** Recommended median daily intake and tolerable upper intake level (UIL) of iodide by age or population group (Maalouf et al. 2015; Dzhatdueva et al. 2005; Valeix et al. 2009; Manousou et al. 2017; Olivieri et al. 2017; Medicine and Institute of Iodine 2001; "Iodine: Fact Sheet by the Office of Dietary Supplements (ODS)" 2011; Institute of Medicine (US) Panel on Micronutrients 2001).

	I <sup>-</sup> intake (µg/day)	UIL (µg/day)
Children 0–6 months	110	Not possible to establish
Children 7–12 months	130	
Children 1–8 years	90	200–300
Children 9–13 years	120	600
Children ≥14 years and adults	150	900
Pregnant	220	900–1100
Lactating	290	900–1100

than the 90% of the human population have access to sufficient iodine from natural sources or from fortified food. However, iodine deficiency is still present in some remote populations that lack access to natural sources of iodine or to iodine fortified foods. The WHO/UNICEF/ICCIDD recommended a daily intake of 150 µg/day of iodine for an adult to avoid IDD. However, this recommendation varies depending of age and physiological status (see Table 2) to supply themselves, fetus and the newborn for thyroid hormones synthesis (Murcia 2018; Opazo and Haensgen, et al. 2017). Among the risk groups for iodine consumption are pregnant and lactating women, they require 50% and 100% more iodine than an adult, respectively (see Table 2) (Maalouf et al. 2015; Dzhatdueva et al. 2005; Valeix et al. 2009; Manousou et al. 2017; Olivieri et al. 2017; Medicine, Institute of Iodine 2001; "Iodine: Fact Sheet by the Office of Dietary Supplements (ODS)" 2011; Institute of Medicine (US) Panel on Micronutrients 2001). The fact that iodine is a scarce element on earth it is important to determine whether an individual is under proper iodine nutrition, especially pregnant and lactating women. Today, the median urinary iodine concentration (UIC) is a parameter used worldwide to monitor iodine nutrition status in a population (Pearce et al. 2016). The UIC value corresponds to the median urinary excretion of iodine obtained after 24 hours of urine recollection or a spot urine sample taken in the morning (Pearce et al. 2016). The UIC values are really used as a parameter for the population of iodine nutrition status. For an individual the UIC accounts for the daily intake of iodine, but not for individual nutrition status due to its high intra- and inter- individual variation (Pearce et al. 2016; Rohner et al. 2014). Regarding to that, a consensus published by WHO indicated that UIC values of a population lower than 100 µg/L corresponds to insufficiency iodine intake and therefore this population it instead is at high risk of suffering IDD (see Table 2) (WHO 2008). Based on the UIC values there are 22 developing countries at risk of suffering severe iodine deficiency (Figure 1) (IGN 2017). Iodine nutrition status of a population could vary inside of a country or region or person to person (Laurberg et al. 2010). It has been reported that countries like Finland, Italy and Russia have regions with iodine deficiency (IGN 2017). Even though, iodine supplementation programs had helped to reduce significantly areas with severe iodine deficiency, there are regions that still not achieve sufficient levels to fulfill the requirements



**Figure 1.** Global status of iodine intake and the legislation of salt iodization. The colors of the world map indicate the levels of iodine intake per person for each country. This was measured by the urinary iodine concentration (UIC). These data were obtained from children, teenagers and adults during the period between 1993 and 2011. Both national and subnational data from recent period was prioritized. The different fillings indicate per country the type of legislation of salt iodization that it has been or not implemented according to the Iodine Global Network (2016).

of iodine for pregnant women (Figure 1) (FAO and WHO 2001). In fact, there are countries, like United States, France Spain, England, Norway, Australia, Turkish, Romania, Poland, that today have an adequate iodine nutrition status for adults and school-age children, however insufficient for pregnant women (Maalouf et al. 2015; Dzhatdoeva et al. 2005; Valeix et al. 2009; Manousou et al. 2017; Olivieri et al. 2017) (IGN 2017). Thus, it is important to count with a biomarker to determine properly iodine nutrition status individually. The determination of TSH,  $T_3$  and  $T_4$  are insensitive indicators for nutritional iodine status. Specially, these hormones they cannot be used to detect mild or moderate iodine deficiency (Pearce et al. 2016). The determination of thyroglobulin (Tg) in blood is a promising biomarker for nutritional iodine status particularly in populations with mild iodine deficiency, however there is a lack of evidence supporting its use in populations other than children (Pearce et al. 2016).

## 2. Iodine for human health

### 2.1. Iodine incorporation into the thyroid hormones

Iodine forms part of the structure of thyroid hormones, which are essential for cell metabolism, differentiation, and for fetus development (Mullur, Liu, and Brent 2014).  $I^-$  disposition is a key factor that affects thyroid physiology and thyroid hormone synthesis (Chung 2014). All mammals capture iodine from the diet in the form of an anion like iodate ( $IO_3^-$ ) or iodide ( $I^-$ ) which are combined with sodium or potassium to form salt (Patrick 2008). Dietary  $I^-$  first is taken up from food and water through the  $Na^+/I^-$  symporter (NIS) that localizes at the intestinal epithelial cells of intestine (Nicola et al. 2009).  $I^-$  is transported by NIS from the blood to tissues like the thyroid gland and lactating mammary gland for thyroid hormone synthesis (Portulano, Paroder-Belenitsky, and Carrasco 2014). It is important to recall that iodine plays its main physiological role by being



part of thyroid hormones. Thyroid hormone synthesis occurs at the thyroid gland and the first and limiting step for the synthesis of thyroid hormones is the  $I^-$  uptake mediated by NIS (Ravera et al. 2017). NIS transports  $I^-$  against its electrochemical gradient by using the energy released by the co-transport of  $Na^+$ , in favor of its electrochemical gradient (Portulano, Paroder-Belenitsky, and Carrasco 2014). Once in the cytosol,  $I^-$  diffuses to the apical side of the cell, where it is transported to the colloid by Pendrin and other transporters (Bizhanova and Kopp 2009; Zbigniew 2017). In the colloid, Tg, the most abundant protein of the thyroid gland, provides the polypeptide backbone for the organification of  $I^-$  at specific tyrosine residues. This reaction is mediated by thyroid peroxidase (TPO) and dual oxidase 2 (DUOX2) (Di Jeso and Arvan 2016). Tg will then be endocytosed by thyroid follicular cells and these endosomes will fuse with lysosomes, allowing the proteases to digest Tg in  $T_3$  and  $T_4$ . These endosomes will fuse to the plasma membrane for the exocytosis of  $T_3$  and  $T_4$  into the blood (Becker 2002; Ahad and Ganie 2010). It has been described that the enzyme iodotyrosine dehalogenase (DEHAL1) is also present in apical side of the thyroid cell and can deiodinate Tg and release of iodotyrosines that contributes as an intracellular source of iodide allowing iodide recycling (Carvalho and Dupuy 2017). All these steps of thyroid hormone synthesis are positively regulated by thyroid stimulating hormone (TSH). Only 30% of the thyroid hormones secreted by the thyroid gland corresponds to  $T_3$  and 70% correspond to  $T_4$  (Bianco et al. 2002) and once in the blood 99.7% of  $T_3$  and 99.95% of  $T_4$  are found bound to proteins like thyroxine-binding globulin (TBG), transthyretin (TTR) and human serum albumin (HAS) (Janssen and Janssen 2017). In fact, this high percentage of thyroid hormones bound to proteins constitute a big reservoir of thyroid hormones for the organs (Janssen and Janssen 2017). It has been reported that both  $T_3$  and  $T_4$  to enter into the cell require carriers like the organic anion transporting polypeptides OATP1A2, OATP4A1, OATP1C1, the L type amino acid transporter LAT1 and LAT2 and monocarboxylate transporter MCT8 and MCT10 (Visser et al. 2011).  $T_3$  is the principal physiological active thyroid hormone (Bianco et al. 2002). To understand the physiological aspects of thyroid hormones homeostasis is necessary to emphasize that the active  $T_3$  in most cells is produced in those cells mainly by the conversion of  $T_4$  into  $T_3$  and not by  $T_3$  secreted from the thyroid gland (Rousset et al. 2015). Therefore, the requirement of  $T_3$  for each organ or tissue is controlled, managed and supplied through the deiodination of  $T_4$  into  $T_3$  for each tissue (Mondal et al. 2016). This process is carried out by deiodinases 1 (D1) and/or deiodinases 2 (D2) (Bianco et al. 2002). Each tissue regulates the expression of these enzymes (Bianco and da Conceição 2018) and their enzymatic activities depend on selenium which will be bound to D1 and D2 (Köhrle 2005).  $T_3$  will perform its biological role by binding to nuclear receptors that are transcription factors that will increase or repress the expression of genes (Cheng, Leonard, and Davis 2010). Thyroid hormones also function

by non-genomic mechanisms please refer to more specific reviews (Davis, Goglia, and Leonard 2016).

### 3. Compensation and diseases caused by unappropriated iodine nutrition

#### 3.1. Mechanisms of compensation under iodine deficiency or excess

Insufficiency or excess of  $I^-$  affects the physiological process of thyroid hormones synthesis and several compensatory mechanisms will run to maintain the synthesis of thyroid hormones (Leung and Braverman 2014). Under  $I^-$  deficiency the thyroid gland compensates  $I^-$  insufficiency by increasing the production of  $T_3$  and decreasing  $T_4$  (Chung 2014; Karmarkar et al. 1969). Authors discuss that the mechanisms behind, are related with turning on the production to monoiodotyrosines (MIT) than diiodotyrosines (DIT). Another mechanism is the increasing expression and activity of deiodinase 2 (D2) at the extrathyroid tissues having as a consequence a higher conversion of  $T_4$  into  $T_3$  (Wang et al. 2009). Interestingly, the opposite occurs at the hypothalamus and hypophysis where the expression and activity of D2 decrease (Mullur, Liu, and Brent 2014). Then the conversion of  $T_4$  into  $T_3$  decreases in these tissues and as consequence the less content of  $T_3$  will stimulate the synthesis of TSH and thyrotropin releasing hormone (TRH). TSH will stimulate the enlargement of the thyroid gland aiming to increase  $I^-$  uptake and thyroid hormone synthesis (Mullur, Liu, and Brent 2014). The excess of iodide in the diet for a short period (3 to 4 days) induces a compensatory mechanism that aims to reduce the uptake of iodide by the thyroid gland and the synthesis of thyroid hormones (Leung and Braverman 2012). This mechanism is named the Wolff-Chaikoff effect. However, if the ingestion of iodide remains higher for more than five days the thyroid gland escapes of the Wolff-Chaikoff effect and begins to capture the excess of  $I^-$  and the patient increases its risk to develop thyroiditis (Luo et al. 2014).

#### 3.2. Endocrine disruptors that affect iodine incorporation into thyroid hormones

Endocrine disruptors agents can alter the gland function at different levels (Schug et al. 2013). This is the case for endocrine disrupting chemicals (EDC) that affect iodine uptake in the thyroid gland (Diamanti-Kandarakis et al. 2009; Diamanti-Kandarakis et al. 2009). Perchlorate, a contaminant found in water and food (Murray et al. 2008), thiocyanate, a component found in plants, chlorate and nitrates are EDC because they compete with iodide by being transported by NIS inducing thyroid dysfunction (De Groef et al. 2006; Dohán et al. 2007). The same iodine uptake reduction has been observed when FTRL-5 cells were incubated with the soy isoflavone genistein (Schmutzler et al. 2007) that interferes with thyroid hormone synthesis by inhibiting TPO (Marini et al. 2012) and it has been also described that is able to inhibit iodine reutilization by the inhibition of

sulfotransferase enzymes (SULT1A1 and SULT1A13) which contributes to the processing of thyroid hormones for the reutilization of iodine (Ebmeier and Anderson 2004). Second, there are small non-coding RNAs (nc-RNA) which are able to control gene expression in a sequence specific way (Catalanotto, Cogoni, and Zardo 2016). The most studied are micro-RNAs (miRNAs) (Catalanotto, Cogoni, and Zardo 2016). Recent studies have identified several miRNAs that are able to modify the expression/function of NIS. Lakshmanan et al., demonstrated using an *in vitro* model that miR-339-5p is able to decrease NIS radioiodine uptake rate (RAUI) (Lakshmanan et al. 2015). Moreover, gene sequencing analyses showed that the products of miR-146, identified in papillary thyroid carcinoma cells, are also able to bind to the UTR region of the NIS gene impairing protein translation and in consequence reduce iodine uptake (Riesco-Eizaguirre et al. 2015). Another important factors that influence thyroid function are trace elements as iron, zinc, copper, calcium and selenium. Selenium is an interesting essential trace element with anti-inflammatory and antioxidant properties (Drutel, Archambeaud, and Caron 2013), able to interact with iodine to contribute to thyroid hormone production (Köhrle 2015). The intake of this micronutrient is highly variable around the world (Rayman 2012). Even though there are several sources such as vegetables, seafood, meat and nuts among others that contain selenium (Finley 2006). Selenium is incorporated as selenocysteine (Sec) to selenoproteins and it is highly concentrated in the thyroid gland where selenium dependent deiodinases are able to produce  $T_3$  from  $T_4$  (Schomburg and Köhrle 2008). In fact, selenium is incorporated to the iodothyronine deiodinase 1 (DIO1) decreasing the conversion of  $T_4$  to  $T_3$  as was observed by Kawai et al., in a study in a Japanese population with severe selenium deficiency where a decreased in the selenium intake is correlated with an increase in free  $T_4$  serum levels meanwhile supplementation of selenium was able to increase free  $T_4$  levels in these patients (Kawai et al. 2018).

### 3.3. Iodine deficiency disorders

As mentioned above, iodine is a key structural component of thyroid hormones. As it is well known iodine deficiency will cause diverse pathologies such as goiter (Gizak, Gorstein, and Andersson 2017), hypothyroidism (Niwattisaiwong, Burman, and Li-Ng 2017), and hypothyroxinemia (Dosiou and Medici 2017). Other consequences of iodine deficiency during pregnancy to the offspring are intellectual impairment like low intellectual coefficient, autism (Błażewicz et al. 2016), schizophrenia (Harper and Brown 2012), and mental retardation (Walker et al. 2007; Pharoah, Buttfeld, and Hetzel 1971). Recently, iodine deficiency has been associated also to anxiety (Młyniec et al. 2015), depression (Młyniec et al. 2015), obesity (Verheesen and Schweitzer 2008) and fibromyalgia (Verheesen and Schweitzer 2008). In the following section of this review we will discuss relevant clinical data of several diseases and health impairment associated with iodine deficiency.

#### 3.3.1. Goiter

Goiter affects a 15.8% of general population and it is characterized by an enlargement of the thyroid gland that occurs in response to high levels of TSH in the blood (Dauksiene et al. 2017). The high levels of TSH are product of low insufficient level of thyroid hormones that can be the result of iodine deficiency or inactivating mutations at Tg (Siffo et al. 2018) and TPO (Ris-Stalpers and Bikker 2010) genes. Goiter is higher in women than in men (Dauksiene et al. 2017) and it increases with age (Gizak, Gorstein, and Andersson 2017).

#### 3.3.2. Hypothyroidism and hypothyroxinemia

Hypothyroidism and hypothyroxinemia are conditions classified among iodine deficiency disorders (IDD). Hypothyroidism, is clinically classified as overt hypothyroidism where patients have low thyroid hormones levels and high TSH levels and affects 1–2% of the population (Seo and Chung 2015). Second, subclinical hypothyroidism is characterized by the solely increase of TSH levels in the blood and it has a 3–15% prevalence worldwide (Peeters 2017). Considering both types, hypothyroidism affects between 4% to 10% of the population (Vanderpump 2011; Taylor et al. 2018). Iodine deficiency is the most important cause of hypothyroidism worldwide, however in those countries where there is sufficient iodine in the diet the main cause of hypothyroidism is autoimmune Hashimoto thyroiditis (Ragusa et al. 2019). Other factors that can provoke hypothyroidism are an impaired function of hypothalamus, pituitary and/or thyroid gland (Kapil 2007). Iodine deficiency it also exposed the appearance of hypothyroidism in those patients that are heterozygous for mutations in DUOX oxidase (Moreno and Visser 2007). The authors emphasize that patients with these mutations require a more elevated amount of iodine in the diet to avoid the development of hypothyroidism. Patients that suffer overt hypothyroidism frequently deal with overweight, coldness, constipation, dry skin among other symptoms. Some patients could also show several health consequences due to the low levels of  $T_3$  down regulates cell metabolism and impair the expression of genes that are important for cell function. It has been reported that hypothyroid patients could develop depression (Larisch et al. 2004; Fardella et al. 2000; Gloger et al. 2001), infertility (La Vignera et al. 2017; Dunn and Turner 2016) and miscarriages during pregnancy (Weiss and Clapauch 2014). It has been observed in hypothyroid patients a systemic vascular resistance leading to an increase of diastolic blood pressure resulting in diastolic hypertension that is often sodium sensitive (Klein and Danzi 2016). It has been reported that patients that suffer of subclinical hypothyroidism also develop heart problems like congestive heart failure (Rodondi et al. 2005) or cardiomyopathy (Zoltowska et al. 2018). The hypothyroxinemia condition is an asymptomatic condition in adults and clinically characterized by the solely decrease of serum  $T_4$  (below the 2.5<sup>th</sup> percentile), while the levels of  $T_3$  and TSH remain normal. It has been proposed that hypothyroxinemia is a compensatory mechanism to overcome iodine deficiency, thus keeping normal the level of

active thyroid hormone  $T_3$  in the blood (Alexander et al. 2017; Henrichs et al. 2013). Hypothyroxinemia reaches higher relevance at the first trimester of pregnancy given that the fetus depends of maternal  $T_4$ . According to the available data, the prevalence of hypothyroxinemia worldwide varies from region to region and it is believed that this condition could be more frequent than hypothyroidism. In those territories with iodine deficiency maternal hypothyroxinemia varies from 3.2% to 23.9%, respectively (López-Muñoz et al. 2019). Impressive is the information that declares that hypothyroxinemia in pregnant woman is also highly prevalent in countries with sufficient iodide nutrition, such as the United States and the Netherlands with 1.3% and 4.3%, respectively (Moleti, Trimarchi, and Vermiglio 2011). Iodine deficiency during pregnancy could induce the occurrence of maternal hypothyroidism or maternal hypothyroxinemia (Negro et al. 2011). Several reports have shown that both conditions have irreversible consequences to the offspring given that the mother supplies the fetus with her thyroid hormones at least until gestational weeks 17–20 (Thompson et al. 2018). This is because the fetus thyroid gland and the axis hypothalamus, hypophysis and thyroid gland has not properly developed and mature to produce thyroid hormones (Opazo, Haensgen, et al. 2017). Thus, the fetus requires maternal thyroid hormones, especially  $T_4$ , because this thyroid hormone is the principal hormone that crosses the placental barrier (Patel et al. 2011). Maternal thyroid hormones are needed for the development of important structures of the fetus like central nervous system (CNS) (Stiles and Jernigan 2010), cells from the immune system (Foster et al. 1998) and gastrointestinal tract (Sirakov and Plateroti 2011) among other organs (Colony 1983). Severe iodine deficiency during pregnancy has been associated with cognitive impairment in the offspring being mental retardation and cretinism the most dramatic consequences (Redman et al. 2016; Dunn 1993). Several studies shown that mild and moderate iodine deficiency during pregnancy can impairs cognition in the offspring, being attention deficit hyperactivity disorder (ADHD), and low intellectual quotient the most frequent observations (Vermiglio 2004; Moleti et al. 2016; Zhou et al. 2015). However, there are controversial results whether maternal iodine supplementation during pregnancy can rescue the offspring from cognitive impairment (O'Donnell et al. 2002; Murcia 2018; Taylor et al. 2017; Moleti et al. 2008). There is an important issue to consider before concluding that iodine supplementation during pregnancy will not rescue cognitive damage in the offspring. That is iodine supplementation must start in women before or at the beginning of pregnancy given that the offspring CNS development begins earlier as the third week of pregnancy (Silbereis et al. 2016). It has been shown that the strongest association among gestational factors and cognitive damage in the offspring is the level of maternal  $T_4$  during pregnancy converting gestational hypothyroxinemia in a harmful condition for the offspring (Levie et al. 2018; Chang and Shin 2014; Korzeniewski et al. 2013; Román 2007). Studies using animal models have also shown a deleterious consequences for CNS development

(Lavado-Autric et al. 2003; Cuevas et al. 2005) and cognitive performance (Opazo et al. 2008). At molecular level, it has been described that maternal hypothyroxinemia alters the expression and localization of key proteins that participate in the long-term potentiation (LTP) process, which is essential for a proper cognitive capacity (Opazo et al. 2008; Cisternas et al. 2016). In the last 11 years have appeared studies in humans that gestational hypothyroxinemia increases the probability of the offspring to develop autism (Román et al. 2013; Hamza, Hewedi, and Sallam 2013; Levie et al. 2018; Chang and Shin 2014; Korzeniewski et al. 2013; Román 2007). The consequences of gestational hypothyroxinemia or gestational hypothyroidism seem to be beyond the CNS, in fact there are reports using animal models that indicated that the offspring is more susceptible to suffer inflammation (Albornoz et al. 2013; Haensgen et al. 2018). Albornoz et al. (2013) showed that adult offspring gestated under hypothyroid condition where more susceptible to develop the EAE, increased  $CD4^+$  T cell infiltration and demyelination in the spinal cord of the offspring gestated under hypothyroidism (Albornoz et al. 2013). Moreover, gestational hypothyroxinemia in mice impairs the suppressive capacity of T regulatory lymphocytes after EAE (Haensgen et al. 2018). Another study, using a murine model, reported that maternal hypothyroxinemia increases the inflammatory response of astrocytes to an inflammatory stimulus (Opazo, Gonzalez, et al. 2017). Still gestational hypothyroxinemia is not considered clinically a dreadful condition for diagnose and treatment, this decision is based on the evidences that showed that  $T_4$  therapy did not revert the harmful effects in the offspring (Alexander et al. 2017).

### 3.4. Consequences of iodine excess intake

Previously in this review it was mention that in most cases the thyroid gland is able to compensate acute periods of iodide excess intake. However, in certain individuals iodine excess for chronic periods or after a period of iodine deficiency could increase the susceptibility to suffer thyroid disorders like hyperthyroidism, hypothyroidism, goiter, and autoimmune diseases (Farebrother, Zimmermann, and Andersson 2019).

#### 3.4.1. Autoimmune thyroiditis

Iodine excess intake has been associated with the development of autoimmune thyroiditis (Leung and Braverman 2014; Katagiri et al. 2017). There are some hypotheses regarding how excess of iodine in the diet can induce autoimmune thyroid diseases. It has been proposed that excess of iodine will increase the iodine organification in thyrocytes leading to oxidative cell damage, recognition of thyrocytes molecules as dangerous for the immune system, and the recruitment of lymphocytes to the thyroid gland (Luo et al. 2014). It also has been proposed that the iodine excess could alter the conformation of Tg and when it is processed by antigen presenting cells (APC), will produce antigens that



will be recognized as hazardous for the immune system (Luo et al. 2014).

Also, cells from the immune system, mainly T cells, infiltrate the thyroid gland and induce cell death by secreting antibodies against thyroid proteins, such as TPO or Tg (Antonelli et al. 2015). Furthermore, recruitment of T helper 1 (Th1) lymphocytes leads to the secretion of inflammatory cytokines, such as TNF- $\alpha$  and IFN- $\gamma$ , which activate the synthesis of chemokines by the thyrocytes, which stimulates Th1 lymphocytes activation and perpetuates the immune response (Antonelli et al. 2014). Thus, patients that have been diagnosed with autoimmune thyroiditis its thyroid gland will not able to produce and secrete thyroid hormones due to cell thyrocytes death caused by inflammation. In fact, most of these patients are diagnose first as hypothyroid patients and they will be treated with thyroxine for life (Brenta et al. 2013).

### 3.4.2. Grave's disease

Graves's disease is an autoimmune disease characterized by antibodies against TSH receptor (Antonelli et al. 2015). These antibodies have the capacity to stimulate the thyroid gland for the overproduction of thyroid hormones and development of hyperthyroidism (Antonelli et al. 2015). This disease may be promoted by an increased iodine intake. However, this is not the only factor involved in the development of Grave's disease. For instance, genetic disposition seems to be the principal risk factor, although it may develop earlier when is high the iodine intake. Furthermore, in Graves' disease the thyroid gland will uptake more iodine and the patient will develop hyperthyroidism (Laurberg et al. 2010).

## 4. The effects of iodine in microorganisms

Although few studies have been performed on the role of iodine and thyroid hormones and microorganisms, in this chapter we have summarized the information available so far. Considering the growing numbers of studies supporting the role of microbiota in health and disease, it is possible that the amount of ingested iodine in the diet and the levels of thyroid hormones could have an important impact in the intestinal microbiota. Iodine has not been described as an essential trace element for microorganisms, but it has been described some strains that able to accumulate radioactive iodine (Amachi et al. 2005; Li et al. 2011). Thus, in the following sections we will discuss the effects of iodine in pathogen control and the effects of iodine intake for the intestinal microbiota.

### 4.1. Iodine for pathogen control

Iodine is the principal specie responsible for antimicrobial efficacy, being one of the main anti-pathogenic agents used today to kill bacteria, viruses and fungi. Iodophors or "iodine releasing agents" had been developed due to iodine itself is very unstable in aqueous solution it establishes a

complex equilibrium between iodine species and iodine. The principal iodophor is the povidone solution. The povidone acts as an active iodine reservoir, releasing iodine to attack microorganisms (McDonnell and Russell 1999). The antimicrobial mechanism of iodine in bacteria seems to be related to the interference with the electron chain transport, preventing respiration and ATP production in aerobic bacteria (Maris 1995). There are evidences that support that iodine also disrupts the cell wall of microorganisms and then attacks nuclear and cytoplasmic structures inside them (McDonnell and Russell 1999). A seminal work performed by Seymour Klebanoff showed that myeloperoxidase, an enzyme found in granules of polymorphonuclear plus hydrogen peroxide and iodine induces iodination of bacteria that accounts for its antimicrobial activity (Klebanoff 1967). The authors proposed that an active intermediate compound is formed in the presence of myeloperoxidase, hydrogen peroxide and iodine, which can be precipitated trichloroacetic acid (TCA) and showed bactericidal effects, showed bactericidal effects has a short half-life because this activity is observed only when bacteria are incubated within this compounds and not after the reaction has occurred (Klebanoff 1967). The authors also shown that for higher bactericidal activity the three compounds must be present (Klebanoff 1967). Another study by Simmons and Karnovsky (1973) have shown that iodination also increases the ability of phagocytes from different animals to uptake *Mycobacterium tuberculosis* (Simmons and Karnovsky 1973). Moreover, the intermediate antimicrobial compounds that show the bactericidal ability could be also generated when myeloperoxidase and hydrogen peroxide are incubated with thyroid hormones T<sub>3</sub> and T<sub>4</sub> (Klebanoff 1967). Also, it was demonstrated that human leukocyte myeloperoxidase, in the presence of hydrogen peroxide, degrades thyroid hormones and in turn promote iodination of microorganisms (Klebanoff and Green 1973). These early studies support the notion that iodination used by biological systems is key tool for the immune system to control bacterial burden. Further, these studies suggest that proper iodine nutrition and the presence of adequate levels of thyroid hormones might be relevant for fighting bacterial infections.

Iodine can also kill viruses due to its ability to destroy inner structural components, such as nucleoproteins and genetic material (Sriwilaijaroen et al. 2009). A recent study has also shown that the presence of povidone-iodine inhibits the entry of human and avian influenza virus into MDCK cells (Sriwilaijaroen et al. 2009), as well as the release of viral particles from infected cells. In this study, the mechanisms of viral infection and replication inhibition proposed was inhibition of virus binding to the sialo-glycoconjugate receptor and inactivation of the sialidase activity of influenza, both human and avian (Sriwilaijaroen et al. 2009). The primordial factor that determines the antimicrobial action of iodine is its working concentration. In fact, there is an optimal working concentration that determines its effectivity against each pathogen, without causing cytotoxicity in mammalian cells (Shirai et al. 2000; Sriwilaijaroen et al. 2009). Further, there are several studies suggesting that the



antimicrobial effect of iodine is a promising tool for infectious diseases that currently lack adequate treatment. This is the case for keratoconjunctivitis, an infection produced by adenovirus that lacks effective antiviral therapy. A clinical study showed that the combination of 1.0% povidone-iodine with 0.1% dexamethasone promoted an important decrease in viral titers, as compared to the other treatments tested, and eliminated the virus after seven days (Kovalyuk et al. 2017). Moreover, the antimicrobial effects of iodine against foodborne pathogens like *Listeria monocytogenes*, *Staphylococcus aureus* and *Escherichia coli* had been studied using biofilms and compared to other disinfectants (Cabeca, Pizzolitto, and Pizzolitto 2012). The disinfectants were in contact with the pathogens for 10 minutes after that the viable cells are registered in the biofilm. Bacteria like *L. monocytogenes*, *S. aureus*, and *E. coli* growth in the biofilms, 6.3, 5.9, and 4.7 CFU/cm<sup>2</sup>, while with 0.20% w/v iodine treatment these numbers decreased to 2.0, 2.4, and 0.8 CFU/cm<sup>2</sup>, respectively. Iodine was the best antimicrobial against these three bacteria as compared to biguanide, quaternary ammonium and peracetic acid (Cabeca, Pizzolitto, and Pizzolitto 2012). Another study showed that povidone-iodine nanoparticles had 100% effectivity as antimicrobial against *Staphylococcus aureus*, *Escherichia coli* and *Pseudomonas aeruginosa* grown in agar plates (Gao et al. 2017). Moreover, povidone-iodine combined with glue, ink and dye have also shown a total inhibition of *Escherichia coli* growth (Gao et al. 2017). The use of iodine in combination with fluorides, has been investigated as dental therapy, given that fluoride by itself is not enough to eliminate dental caries (Featherstone 2006). The effect of single and combined effect of sodium fluoride with iodine was tested against *Streptococcus* mutants, the main cause of the dental caries, showing that the combination of these two agents exerted a beneficial antimicrobial effect against this bacterium (Featherstone 2006). This study also indicated that health-care professionals must use this combination at lower concentrations to avoid irritation of the mucous membranes (Caufield and Wannemuehler 1982).

#### 4.2. Iodine for the regulation of normal microbiota

It is very relevant to evaluate the effect of iodine over normal microbiota, considering that iodine is a micronutrient and it has an antimicrobial effect. However, there are few studies in the literature that have evaluated the effect of iodine over the oral and the intestinal microbiota. It is possible that dietary iodine, even though in small quantities, could affect the intestinal flora. In support of this notion, a study evaluated the quality of milk and the quantity of microorganism in the milk of cows that were fed with marine brown algae *Ascophyllum nodosum* as an iodine supplement. The authors found that even though the milk of these cows has a high level of iodine, this level was below the maximum permitted. Interestingly, the consumption of *Ascophyllum nodosum* reduced the proportion of *Pseudomonas* genus in the milk from 73.19% to 32.87% and increased the proportion of *Lactococcus* from 22.14% to 66.37%. A remarkably

fact was that the lifespan of the milk increased due to the reduction of the genus *Pseudomonas* (Chaves Lopez et al. 2016). Another study by Shen et al. (2019) evaluated the effect of iodine supplementation (18 µg/K/day) in normal and obese rodents (Shen et al. 2019). The authors found that after 8 weeks of treatment normal mice showed an increase abundance of beneficial intestinal bacteria, while obese mice showed a oppose phenotype. It has been also evaluated whether vulvar disinfection of mother during vaginal delivery affects the composition of the oral microbiota in newborns (Li et al. 2019). This study describes that both cesarian-delivered and conventional disinfected-delivery groups have a reduced abundance of the genus *Lactobacillus* and increased abundance of Proteobacteria and Bacteroidetes. Based on these studies, it is possible to suggest that the consumption or use of iodine could affect the human microbiota composition, which could be beneficial or detrimental depending on other health conditions of the host. We should take into account the importance of the quantity of ingested iodine, thinking that there are regions in the world where the consumption of iodine surpass 500 µg per day. These areas are considered to be in risk of thyroid diseases due to diets with iodine excess (WHO 2004) and unexplored effect in the intestinal microbiota can also be expected.

#### 4.3. Effects of thyroid diseases on intestinal microbiota

Other studies have evaluated the effect of thyroid diseases in the intestinal microbiota. A study found that Bifidobacterium and *Lactobacillus* were decreased in hyperthyroid patients and that *Enterococcus* was increased (L. Zhou et al. 2014). Furthermore, it has been described that the intestinal microbiota of patients with Hashimoto's thyroiditis has important differences, as compared to healthy subjects (Ishaq et al. 2017). This study reported that some genera, such as Prevotella and Dialister, showed decreased abundance in Hashimoto's patients. In contrast, *Escherichia*, *Shigella* and *Parasutterella* were increased. These results confirmed dysbiosis in Hashimoto's Thyroiditis patients, which could be closely related with iodine intake (Ishaq et al. 2017).

Another recent study established a correlation between intestinal microbiota of Hashimoto's patients and clinical parameters. This study shows that Hashimoto's patients have greater gut microbiota diversity than healthy controls (Zhao et al. 2018). This result may be due to bacterial overgrowth in the gastrointestinal tract of patients. Furthermore, the authors described a positive correlation between eighteen microbiota genera and TPO and Tg antibodies (Zhao et al. 2018).

It has been also suggested that intestinal microbiota may play a role in the absorption of I<sup>-</sup>. Vought et al. (1972) made a first approach by treating rats with kanamycin and then measuring I<sup>-</sup> absorption. Results showed that radioiodine uptake was significantly lower in treated rats than untreated rats, at 3 hours and after 42 and 72 days of treatment (Vought et al. 1972). Also, it has been described that

*Escherichia coli* binds T<sub>3</sub> and uptake iodothyronines. These bacteria have been related to decarboxylation, deiodination and deamination of iodothyronines. Additionally, Virili and Centanni (2017) suggested that intestinal microbiota may modulate the absorption of oral T<sub>4</sub>. Finally, lipopolysaccharide (LPS) from Gram-negative bacteria can stimulate TSH-induced iodine uptake by thyroid cells, by activation of NF- $\kappa$ B pathway and expression of Na<sup>+</sup>/I<sup>-</sup> symporter (NIS) (Nicola et al. 2010). Therefore, it is possible that commensal bacteria have a huge impact in the balance of thyroid hormones, and that I<sup>-</sup> intake affect the composition and diversity of intestinal microbiota. Further studies are required to evaluate these hypotheses.

## 5. Concluding remarks

Iodine is an essential nutrient for human life and the reality that there is a great disparity in its consumption around the world. This causes that approximately 36.5% of the people suffer of IDD and paradoxically 10 countries have high iodine intake (FAO and WHO 2001). Moreover, there are countries that the general population had sufficient iodine ingestion, however pregnant women are suffering mild iodine deficiency. To resolve these problems the governments will have to do a real effort for improving iodine intake in their countries. These tasks will involve health professionals, epidemiologist and politicians aiming to develop health policies that will eradicate IDD or high iodine intake especially in pregnant women and neonates. Governments must establish adequate and strategic plans that will precisely regulate and monitor iodine intake. Given that, the needs for iodine consumption are different depending of age, physiological status or region, the designed health plans have to fulfill these different necessities. Moreover, gestational hypothyroxinemia is still considered a condition and not a disease and for that it is not diagnosed or treated, even though there are strong evidences showing that impairs the cognitive capacities of the offspring. In animal models it has been shown that gestational hypothyroxinemia also affects the offspring immune response, specially increasing the intensity of autoimmune diseases like multiple sclerosis (Haensgen et al. 2018). In addition, there are several ailments like autoimmune diseases, autism, gastrointestinal alterations and dysbiosis that are linked to IDD. Thus, the consequences of IDD are wide and for this reason are important not only to be aware of them but also to take actions in their prevention. Iodine nutritional status needs to be evaluated constantly in human population, because iodine is essential for humans and it is a common health issue problem around the world. Now, there is not one a simple way to resolve this problem, given that iodine nutritional status varies between regions of the word or in the same country. Thus, it is important for each country to be aware of the status of iodine nutrition in order to develop the proper strategies to eliminate IDD or to avoid thyroid disfunction due to iodine excess intake. Future research should focus in the development of biomarkers that allow the appropriate determination of iodine status in individuals.


## Conflicts of interest

The authors declare no conflict of interest.

## Funding

The authors are supported by the Millennium Institute on Immunology and Immunotherapy, Iniciativa Científica Milenio, Ministry of Economy, Government of Chile (P09/016-F for MCO, CR, SB and AMK), the ECOS France-Chile 860 grant, Fondo Nacional de Desarrollo Científico y Tecnológico, Ministry of Education, Government of Chile (FONDECYT # 1191300, 1130996, 1161525, 1110604, 1070352, 1050979, 1040349, 1100926, 1110397, 1100971, 1110604). CR: Nucleus project UNAB DI-471-15/N and DI-03-19/N. ICA is supported by a PhD Fellowship from the Comisión Nacional de Investigación Científica y Tecnológica, Ministry of Education, Government of Chile (#63140215). MCO: Fondecyt 11180739, Nucleus project UNAB DI-03-19/N.CCC VI Comisión mixta de Cooperación bilateral entre las regiones de Chile y Valonia-Bruselas periodo 2019-2021.

## ORCID

Ma. Cecilia Opazo  <http://orcid.org/0000-0003-4263-9294>  
 Irenice Coronado-Arrázola  <http://orcid.org/0000-0001-5418-2768>  
 Omar P. Vallejos  <http://orcid.org/0000-0002-5203-7062>  
 Carlos Fardella  <http://orcid.org/0000-0002-5500-4434>  
 Alexis M. Kalgiris  <http://orcid.org/0000-0001-7622-5263>  
 Susan M. Bueno  <http://orcid.org/0000-0002-7551-8088>  
 Claudia A. Riedel  <http://orcid.org/0000-0001-6168-8601>

## References

- Aburto, .N., M. Abudou, V. Candeias, T. Wu. 2014. Effect and safety of salt iodization to prevent iodine deficiency disorders: a systematic review with meta-analyses. WHO eLibrary of Evidence for Nutrition Actions (eLENA). Geneva: World Health Organization.
- Adams, J. B., C. E. Holloway, F. George, and D. Quig. 2006. Analyses of toxic metals and essential minerals in the hair of Arizona children with autism and associated conditions, and their mothers. *Biological Trace Element Research* 110 (3):193–209. doi: [10.1385/BTER.110.3.193](https://doi.org/10.1385/BTER.110.3.193).
- Ahad, F., and S. A. Ganie. 2010. Iodine, iodine metabolism and iodine deficiency disorders revisited. *Indian Journal of Endocrinology and Metabolism* 14 (1):13–7. <http://www.ncbi.nlm.nih.gov/pubmed/21448409>.
- Albornoz, E. A., L. J. Carreño, C. M. Cortes, P. A. Gonzalez, P. A. Cisternas, K. M. Cautivo, T. P. Catalán, M. C. Opazo, E. A. Eugenin, J. W. Berman, et al. 2013. Gestational hypothyroidism increases the severity of experimental autoimmune encephalomyelitis in adult offspring. *Thyroid* 23 (12):1627–37. doi: [10.1089/thy.2012.0401](https://doi.org/10.1089/thy.2012.0401).
- Alexander, E. K., E. N. Pearce, G. A. Brent, R. S. Brown, H. Chen, C. Dosiou, W. A. Grobman, P. Laurberg, J. H. Lazarus, S. J. Mandel, et al. 2017. 2017 guidelines of the American Thyroid Association for the diagnosis and management of thyroid disease during pregnancy and the postpartum. *Thyroid* 27 (3):315–89. doi: [10.1089/thy.2016.0457](https://doi.org/10.1089/thy.2016.0457).
- Amachi, S., Y. Mishima, H. Shinoyama, Y. Muramatsu, and T. Fujii. 2005. Active transport and accumulation of iodide by newly isolated marine bacteria. *Applied and Environmental Microbiology* 71 (2): 741–5. doi: [10.1128/AEM.71.2.741-745.2005](https://doi.org/10.1128/AEM.71.2.741-745.2005).
- Antonelli, A., S. M. Ferrari, A. Corrado, A. Di Domenicantonio, and P. Fallahi. 2015. Autoimmune thyroid disorders. *Autoimmunity Reviews* 14 (2):174–80. doi: [10.1016/j.autrev.2014.10.016](https://doi.org/10.1016/j.autrev.2014.10.016).

- Antonelli, A., S. M. Ferrari, D. Giuggioli, E. Ferrannini, C. Ferri, and P. Fallahi. 2014. Chemokine (C-X-C motif) ligand (CXCL)10 in autoimmune diseases. *Autoimmunity Reviews* 13 (3):272–80. doi: [10.1016/j.autrev.2013.10.010](https://doi.org/10.1016/j.autrev.2013.10.010).
- Bath, S. C., V. J. Pop, V. L. Furmidge-Owen, M. A. Broeren, and M. P. Rayman. 2017. Thyroglobulin as a functional biomarker of iodine status in a cohort study of pregnant women in the United Kingdom. *Thyroid* 27 (3):426–33. doi: [10.1089/thy.2016.0322](https://doi.org/10.1089/thy.2016.0322).
- Becker, L. K., J. P. Bilezikian, and Ovid Technologies, Inc. 2001. Part III: The thyroid gland. In *Principles and practice of endocrinology and metabolism*. Philadelphia, PA: Lippincott Williams & Wilkins Publishers.
- Bianco, A. C., and R. R. da Conceição. 2018. The deiodinase trio and thyroid hormone signaling. *Methods in molecular biology* 1801: 67–83. New York, NY: Humana Press Inc. doi: [10.1007/978-1-4939-7902-8\\_8](https://doi.org/10.1007/978-1-4939-7902-8_8).
- Bianco, A. C., D. Salvatore, B. Gereben, M. J. Berry, and P. R. Larsen. 2002. Biochemistry, cellular and molecular biology, and physiological roles of the iodothyronine selenodeiodinases. *Endocrine Reviews* 23 (1):38–89. doi: [10.1210/edrv.23.1.0455](https://doi.org/10.1210/edrv.23.1.0455).
- Bizhanova, A., and P. Kopp. 2009. Minireview: The sodium-iodide symporter NIS and pendrin in iodide homeostasis of the thyroid. *Endocrinology* 150 (3):1084–90. doi: [10.1210/en.2008-1437](https://doi.org/10.1210/en.2008-1437).
- Błażewicz, A., A. Makarewicz, I. Korona-Glowniak, W. Dolliver, and R. Kocjan. 2016. Iodine in autism spectrum disorders. *Journal of Trace Elements in Medicine and Biology* 34:32–7. doi: [10.1016/j.jtemb.2015.12.002](https://doi.org/10.1016/j.jtemb.2015.12.002).
- Bourdoux, P., F. Delange, M. Gerard, M. Mafuta, A. Hanson, and A. M. Ermans. 1978. Evidence that cassava ingestion increases thiocyanate formation: A possible etiologic factor in endemic goiter\*. *The Journal of Clinical Endocrinology and Metabolism* 46 (4):613–21. doi: [10.1210/jcem-46-4-613](https://doi.org/10.1210/jcem-46-4-613).
- Braverman, L. E., X. M. He, S. Pino, M. Cross, B. Magnani, S. H. Lamm, M. B. Kruse, A. Engel, K. S. Crump, and J. P. Gibbs. 2005. The effect of perchlorate, thiocyanate, and nitrate on thyroid function in workers exposed to perchlorate long-term. *The Journal of Clinical Endocrinology & Metabolism* 90 (2):700–6. doi: [10.1210/jc.2004-1821](https://doi.org/10.1210/jc.2004-1821).
- Brenta, G., M. Vaisman, J. A. Sgarbi, L. M. Bergoglio, N. C. Andrada, P. P. Bravo, A. M. Orlandi, H. Graf, and Society Task Force on Hypothyroidism of the Latin American Thyroid. 2013. Clinical practice guidelines for the management of hypothyroidism. *Arquivos Brasileiros de Endocrinologia e Metabologia* 57 (4):265–91. doi: [10.1590/s0004-27302013000400003](https://doi.org/10.1590/s0004-27302013000400003).
- Cabeca, T. K., A. C. Pizzolitto, and E. L. Pizzolitto. 2012. Activity of disinfectants against foodborne pathogens in suspension and adhered to stainless steel surfaces. *Brazilian Journal of Microbiology* 43 (3):1112–9. doi: [10.1590/S1517-838220120003000038](https://doi.org/10.1590/S1517-838220120003000038).
- Carvalho, D. P., and C. Dupuy. 2017. Thyroid hormone biosynthesis and release. *Molecular and Cellular Endocrinology* 458:6–15. doi: [10.1016/j.mce.2017.01.038](https://doi.org/10.1016/j.mce.2017.01.038).
- Catalanotto, C., C. Cogoni, and G. Zardo. 2016. MicroRNA in control of gene expression: An overview of nuclear functions. *International Journal of Molecular Sciences* 17 (10):1712. doi: [10.3390/ijms17101712](https://doi.org/10.3390/ijms17101712).
- Caufield, P. W., and Y. M. Wannemuehler. 1982. In vitro susceptibility of *Streptococcus mutans* 6715 to iodine and sodium fluoride, singly and in combination, at various PH values. *Antimicrobial Agents and Chemotherapy* 22 (1):115–9. doi: [10.1128/AAC.22.1.115](https://doi.org/10.1128/AAC.22.1.115).
- Cavalieri, R. R. 1997. Iodine metabolism and thyroid physiology: Current concepts. *Thyroid* 7 (2):177–81. doi: [10.1089/thy.1997.7.177](https://doi.org/10.1089/thy.1997.7.177).
- Chang, K., and J. I. Shin. 2014. Association of gestational maternal hypothyroxinemia and increased autism risk: The role of brain-derived neurotrophic factor. *Annals of Neurology* 75 (6):971. doi: [10.1002/ana.24143](https://doi.org/10.1002/ana.24143).
- Chaves Lopez, C., A. Serio, C. Rossi, G. Mazzarrino, S. Marchetti, F. Castellani, L. Grotta, F. P. Fiorentino, A. Paparella, and G. Martino. 2016. Effect of diet supplementation with *Ascochyllum nodosum* on cow milk composition and microbiota. *Journal of Dairy Science* 99 (8):6285–97. doi: [10.3168/jds.2015-10837](https://doi.org/10.3168/jds.2015-10837).
- Cheng, S.-Y., J. L. Leonard, and P. J. Davis. 2010. Molecular aspects of thyroid hormone actions. *Endocrine Reviews* 31 (2):139–70. doi: [10.1210/er.2009-0007](https://doi.org/10.1210/er.2009-0007).
- Chung, H. R. 2014. Iodine and thyroid function. *Annals of Pediatric Endocrinology & Metabolism* 19 (1):8–12. doi: [10.6065/apem.2014.19.1.8](https://doi.org/10.6065/apem.2014.19.1.8).
- Cisternas, P., A. Louveau, S. M. Bueno, A. M. Kalergis, H. Boudin, and C. A. Riedel. 2016. Gestational hypothyroxinemia affects glutamatergic synaptic protein distribution and neuronal plasticity through neuron-astrocyte interplay. *Molecular Neurobiology* 53 (10):7158–69. doi: [10.1007/s12035-015-9609-0](https://doi.org/10.1007/s12035-015-9609-0).
- Colony, P. C. 1983. *Successive phases of human fetal intestinal development. In Nutritional adaptation of the gastrointestinal tract of the newborn*, eds. N. Kretchmer and A. Minkows, 3–28. New York: Nestlé, Vevey/RavenPress.
- Cuevas, E., E. Auso, M. Telefont, G. Morreale de Escobar, C. Sotelo, and P. Berbel. 2005. Transient maternal hypothyroxinemia at onset of corticogenesis alters tangential migration of medial ganglionic eminence-derived neurons. *The European Journal of Neuroscience* 22 (3):541–51. doi: [10.1111/j.1460-9568.2005.04243.x](https://doi.org/10.1111/j.1460-9568.2005.04243.x).
- Dai, J. L., Y. G. Zhu, M. Zhang, and Y. Z. Huang. 2004. Selecting iodine-enriched vegetables and the residual effect of iodate application to soil. *Biological Trace Element Research* 101 (3):265–76. doi: [10.1385/BTER.101:3:265](https://doi.org/10.1385/BTER.101:3:265).
- Dauksiene, D., J. Petkeviciene, J. Klumbiene, R. Verkauskiene, J. Vainikonyte-Kristapone, A. Seibokaite, J. Ceponis, V. Sidlauskas, L. Daugintyte-Petrusiene, A. Norkus, et al. 2017. Factors associated with the prevalence of thyroid nodules and goiter in middle-aged euthyroid subjects. *International Journal of Endocrinology*. 2017:1–8. doi: [10.1155/2017/8401518](https://doi.org/10.1155/2017/8401518).
- Davis, P. J., F. Goglia, and J. L. Leonard. 2016. Nongenomic actions of thyroid hormone. *Nature Reviews. Endocrinology* 12 (2):111–21. doi: [10.1038/nrendo.2015.205](https://doi.org/10.1038/nrendo.2015.205).
- De Groef, B., B. R. Decallonne, S. V. d Geyten, V. M. Darras, and R. Bouillon. 2006. Perchlorate versus other environmental sodium/iodide symporter inhibitors: Potential thyroid-related health effects. *European Journal of Endocrinology* 155 (1):17–25. doi: [10.1530/eje.1.02190](https://doi.org/10.1530/eje.1.02190).
- Di Jeso, B., and P. Arvan. 2016. Thyroglobulin from molecular and cellular biology to clinical endocrinology. *Endocrine Reviews* 37 (1): 2–36. doi: [10.1210/er.2015-1090](https://doi.org/10.1210/er.2015-1090).
- Diamanti-Kandarakis, E., J. P. Bourguignon, L. C. Giudice, R. Hauser, G. S. Prins, A. M. Soto, R. T. Zoeller, and A. C. Gore. 2009. Endocrine-disrupting chemicals: An endocrine society scientific statement. *Endocrine Reviews* 30 (4):293–342. doi: [10.1210/er.2009-0002](https://doi.org/10.1210/er.2009-0002).
- Dohán, O., C. Portulano, C. Basquin, A. Reyna-Neyra, L. M. Amzel, and N. Carrasco. 2007. The Na<sup>+</sup>/I symporter (NIS) mediates electroneutral active transport of the environmental pollutant perchlorate. *Proceedings of the National Academy of Sciences of the United States of America* 104 (51):20250–55. doi: [10.1073/pnas.0707207104](https://doi.org/10.1073/pnas.0707207104).
- Dosiou, C., and M. Medici. 2017. MANAGEMENT OF ENDOCRINE DISEASE: Isolated maternal hypothyroxinemia during pregnancy: Knowns and unknowns. *European Journal of Endocrinology* 176 (1): R21–R38. doi: [10.1530/EJE-16-0354](https://doi.org/10.1530/EJE-16-0354).
- Drutel, A., F. Archambeaud, and P. Caron. 2013. Selenium and the thyroid gland: More good news for clinicians. *Clinical Endocrinology* 78 (2):155–164. doi: [10.1111/cen.12066](https://doi.org/10.1111/cen.12066).
- Dunn, D., and C. Turner. 2016. Hypothyroidism in Women. *Nursing for Women's Health* 20 (1):93–98. doi: [10.1016/j.nwh.2015.12.002](https://doi.org/10.1016/j.nwh.2015.12.002).
- Dunn, J. T. 1993. Iodine supplementation and the prevention of cretinism. *Annals of the New York Academy of Sciences* 678:158–68. doi: [10.1111/j.1749-6632.1993.tb26119.x](https://doi.org/10.1111/j.1749-6632.1993.tb26119.x).
- Dzhatdoeva, F. A., L. E. Syrtsova, G. A. Gerasimov, T. E. Zubrilova, and Z. N. Salpagarova. 2005. [Prevention of iodine-deficient diseases]. *Probl Sotsialnoi Gig Zdravookhranennii Istor Med Jan-Feb(1):25–26*. <http://www.ncbi.nlm.nih.gov/pubmed/15828388>.
- Eastman, C. J., and M. Li. 2017. Mild to moderate iodine deficiency. In *Iodine deficiency disorders and their elimination*, ed. E. N. Pearce,



- 59–74. Cham: Springer International Publishing. doi: [10.1007/978-3-319-49505-7\\_5](https://doi.org/10.1007/978-3-319-49505-7_5).
- Ebmeier, C. C., and R. J. Anderson. 2004. Human thyroid phenol sulfotransferase enzymes 1A1 and 1A3: Activities in normal and diseased thyroid glands, and inhibition by thyroid hormones and phytoestrogens. *The Journal of Clinical Endocrinology and Metabolism* 89 (11):5597–5605. doi: [10.1210/jc.2003-031939](https://doi.org/10.1210/jc.2003-031939).
- FAO and WHO. 2001. *FAO/WHO expert consultation on human vitamin and mineral requirements*. Report of a Joint FAO/WHO Expert Consultation.
- Fardella, C., S. Gloger, R. Figueroa, R. Santis, C. Gajardo, C. Salgado, S. Barroilhet, and A. Foradori. 2000. High prevalence of thyroid abnormalities in a Chilean psychiatric outpatient population. *Journal of Endocrinological Investigation* 23 (2):102–6. doi: [10.1007/BF03343687](https://doi.org/10.1007/BF03343687).
- Farebrother, J., M. B. Zimmermann, and M. Andersson. 2019. Excess iodine intake: Sources, assessment, and effects on thyroid function. *Annals of the New York Academy of Sciences* 1446 (1):44–65. doi: [10.1111/nyas.14041](https://doi.org/10.1111/nyas.14041).
- Featherstone, J. D. 2006. Delivery challenges for fluoride, chlorhexidine and xylitol. *BMC Oral Health* 6 Suppl 1:S8. doi: [10.1186/1472-6831-6-S1-S8](https://doi.org/10.1186/1472-6831-6-S1-S8).
- Finley, J. W. 2006. Bioavailability of selenium from foods. *Nutrition Reviews* 64 (3):146–151. doi: [10.1301/nr.2006.mar.146-151](https://doi.org/10.1301/nr.2006.mar.146-151).
- Foster, M., E. Montecino-Rodriguez, R. Clark, and K. Dorshkind. 1998. Regulation of B and T cell development by anterior pituitary hormones. *Cellular and Molecular Life Sciences* 54 (10):1076–82. doi: [10.1007/s000180050236](https://doi.org/10.1007/s000180050236).
- Fuge, R., and C. C. Johnson. 1986. The geochemistry of iodine—A review. *Environmental Geochemistry and Health* 8 (2):31–54. doi: [10.1007/BF02311063](https://doi.org/10.1007/BF02311063).
- Gaitan, E. 1990. Goitrogens in food and water. *Annual Review of Nutrition* 10:21–39. doi: [10.1146/annurev.nu.10.070190.000321](https://doi.org/10.1146/annurev.nu.10.070190.000321).
- Gao, T., H. Fan, X. Wang, Y. Gao, W. Liu, W. Chen, A. Dong, and Y. J. Wang. 2017. Povidone-iodine-based polymeric nanoparticles for antibacterial applications. *ACS Applied Materials & Interfaces* 9 (31):25738–46. doi: [10.1021/acsami.7b05622](https://doi.org/10.1021/acsami.7b05622).
- Gizak, M., J. Gorstein, and M. Andersson. 2017. Epidemiology of iodine deficiency. In *Iodine deficiency disorders and their elimination*, ed. E. Pearce. Cham: Springer. [10.1007/978-3-319-49505-7\\_3](https://doi.org/10.1007/978-3-319-49505-7_3).
- Gloger, S., C. Fardella, C. Gajardo, R. Figueroa, C. Salgado, R. Santis, and S. Barroilhet. 2001. [Panic disorder and hyperthyroidism. Report of 2 cases]. *Revista Medica de Chile* 129 (2):187–90. <http://www.ncbi.nlm.nih.gov/pubmed/11351471>.
- Haensgen, H., E. Albornoz, M. C. Opazo, K. Bugueño, E. L. Jara Fernández, R. Binzberger, T. Rivero-Castillo, L. F. Venegas Salas, F. Simon, C. Cabello-Verrugio, et al. 2018. Gestational hypothyroxinemia affects its offspring with a reduced suppressive capacity impairing the outcome of the experimental autoimmune encephalomyelitis. *Frontiers in Immunology* 9:1257. doi: [10.3389/fimmu.2018.01257](https://doi.org/10.3389/fimmu.2018.01257).
- Hamza, R. T., D. H. Hewedi, and M. T. Sallam. 2013. Iodine deficiency in Egyptian autistic children and their mothers: Relation to disease severity. *Archives of Medical Research* 44 (7):555–61. doi: [10.1016/j.arcmed.2013.09.012](https://doi.org/10.1016/j.arcmed.2013.09.012).
- Harper, K. N., and A. S. Brown. 2012. Prenatal nutrition and the etiology of schizophrenia. In *The origins of schizophrenia*, ed. A. S. Brown and P. H. Patterson, 58–95. New York: Columbia University Press.
- Henrichs, J., A. Ghassabian, R. P. Peeters, and H. Tiemeier. 2013. Maternal hypothyroxinemia and effects on cognitive functioning in childhood: How and why? *Clinical Endocrinology* 79 (2):152–62. doi: [10.1111/cen.12227](https://doi.org/10.1111/cen.12227).
- IGN. 2017. *Iodine Global Network. Global Scorecard of Iodine Nutrition in 2017 in the General Population and in Pregnant Women (PW)*. Zurich, Switzerland: IGN.
- Institute of Medicine (US) Panel on Micronutrients. 2001. *Iodine . In Dietary reference intakes for vitamin A, vitamin K, arsenic, boron, chromium, copper, iodine, iron, manganese, molybdenum, nickel, silicon, vanadium, and zinc*. Washington, DC: National Academies Press (US). <https://www.ncbi.nlm.nih.gov/books/NBK222310/>.
- Iodine: Fact sheet by the Office of Dietary Supplements (ODS). 2011. <https://ods.od.nih.gov/factsheets/Iodine-HealthProfessional/#en2>.
- Ishaq, H. M., I. S. Mohammad, H. Guo, M. Shahzad, Y. J. Hou, C. Ma, Z. Naseem, X. Wu, P. Shi, and J. Xu. 2017. Molecular estimation of alteration in intestinal microbial composition in Hashimoto's thyroiditis patients. *Biomedicine & Pharmacotherapy = Biomedecine & Pharmacotherapie* 95:865–74. doi: [10.1016/j.biopha.2017.08.101](https://doi.org/10.1016/j.biopha.2017.08.101).
- Janssen, S. T., and O. E. Janssen. 2017. Directional thyroid hormone distribution via the blood stream to target sites. *Molecular and Cellular Endocrinology* 458:16–21. doi: [10.1016/j.mce.2017.02.037](https://doi.org/10.1016/j.mce.2017.02.037).
- Johnson, C. C. 2003. *The geochemistry of iodine and its application to environmental strategies for reducing the risks from iodine deficiency disorders*. British Geological Survey Commissioned Report, CR/03/057N. 54 pp.
- Jong, J. M. d. 2007. Final report review of use of iodized salt in processed foods. Global Iodine Network. 2007. [http://www.iccid.org/cm\\_data/Salt\\_in\\_processed\\_foods.pdf](http://www.iccid.org/cm_data/Salt_in_processed_foods.pdf).
- Josefsson, M., and E. Ekblad. 2009. Chapter 22—Sodium iodide symporter (NIS) in gastric mucosa: Gastric iodide secretion A2—Preedy, Edited By Victor R. In *Comprehensive handbook of iodine*, ed. G. N. Burrow and R. Watson, 215–20. San Diego: Academic Press. [10.1016/B978-0-12-374135-6.00022-4](https://doi.org/10.1016/B978-0-12-374135-6.00022-4).
- Kapil, U. 2007. Health consequences of iodine deficiency. *Sultan Qaboos University Medical Journal* 7 (3):267–72. <http://www.ncbi.nlm.nih.gov/pubmed/21748117>.
- Karmarkar, M. G., N. Kochupillai, M. G. Deo, and V. Ramalingaswami. 1969. Adaptation of thyroid gland to iodine deficiency. *Life Sciences* 8 (21):1135–41. doi: [10.1016/0024-3205\(69\)90042-3](https://doi.org/10.1016/0024-3205(69)90042-3).
- Katagiri, R., X. Yuan, S. Kobayashi, and S. Sasaki. 2017. Effect of excess iodine intake on thyroid diseases in different populations: A systematic review and meta-analyses including observational studies. *PLoS One* 12 (3):e0173722. doi: [10.1371/journal.pone.0173722](https://doi.org/10.1371/journal.pone.0173722).
- Kawai, M., Y. Shoji, S. Onuma, Y. Etani, and S. Ida. 2018. Thyroid hormone status in patients with severe selenium deficiency. *Clinical Pediatric Endocrinology* 27 (2):67–74. doi: [10.1297/cpe.27.67](https://doi.org/10.1297/cpe.27.67).
- Klebanoff, S. J. 1967. Iodination of bacteria: A bactericidal mechanism. *The Journal of Experimental Medicine* 126 (6):1063–78. doi: [10.1084/jem.126.6.1063](https://doi.org/10.1084/jem.126.6.1063).
- Klebanoff, S. J., and W. L. Green. 1973. Degradation of thyroid hormones by phagocytosing human leukocytes. *The Journal of Clinical Investigation* 52 (1):60–72. doi: [10.1172/JCI107174](https://doi.org/10.1172/JCI107174).
- Klein, I., and S. Danzi. 2016. Thyroid disease and the heart. *Current Problems in Cardiology* 41 (2):65–92. doi: [10.1016/j.cpcardiol.2015.04.002](https://doi.org/10.1016/j.cpcardiol.2015.04.002).
- Köhrle, J. 2005. Selenium and the control of thyroid hormone metabolism. *Thyroid* 15 (8):841–53. doi: [10.1089/thy.2005.15.841](https://doi.org/10.1089/thy.2005.15.841).
- Köhrle, J. 2015. Selenium and the thyroid. *Current Opinion in Endocrinology, Diabetes, and Obesity* 22 (5):392–401. doi: [10.1097/MED.0000000000000190](https://doi.org/10.1097/MED.0000000000000190). PMID: 26313901
- Korzeniewski, S. J., J. A. Pinto-Martin, A. H. Whitaker, J. F. Feldman, J. M. Lorenz, S. E. Levy, T. Z. Movsas, A. Pappas, and N. Paneth. 2013. Association between transient hypothyroxinaemia of prematurity and adult autism spectrum disorder in a low-birthweight cohort: An exploratory study. *Paediatric and Perinatal Epidemiology* 27 (2):182–7. doi: [10.1111/ppe.12034](https://doi.org/10.1111/ppe.12034).
- Korzh, V. D. 1984. Ocean as a source of atmospheric iodine. *Atmospheric Environment* 18 (12):2707–10. doi: [10.1016/0004-6981\(84\)90336-6](https://doi.org/10.1016/0004-6981(84)90336-6).
- Kovalyuk, N., I. Kaiserman, M. Mimouni, O. Cohen, S. Levartovsky, H. Sherbany, and M. Mandelboim. 2017. Treatment of adenoviral keratoconjunctivitis with a combination of povidone-iodine 1.0% and dexamethasone 0.1% drops: A clinical prospective controlled randomized study. *Acta Ophthalmologica* 95 (8):e686. doi: [10.1111/aos.13416](https://doi.org/10.1111/aos.13416).
- La Vignera, S., R. Vita, R. A. Condorelli, L. M. Mongioi, S. Presti, S. Benvenga, and A. E. Calogero. 2017. Impact of thyroid disease on testicular function. *Endocrine* 58 (3):397–407. doi: [10.1007/s12020-017-1303-8](https://doi.org/10.1007/s12020-017-1303-8).



- Lakshmanan, A., A. Wojcicka, M. Kotlarek, X. Zhang, K. Jazdzewski, and S. M. Jhiang. 2015. MicroRNA-339-5p modulates Na<sup>+</sup>/I<sup>-</sup> symporter-mediated radioiodide uptake. *Endocrine-Related Cancer* 22 (1):11–21. doi: [10.1530/ERC-14-0439](https://doi.org/10.1530/ERC-14-0439).
- Larisch, R., K. Kley, S. Nikolaus, W. Sitte, M. Franz, H. Hautzel, W. Tress, and H. W. Muller. 2004. Depression and anxiety in different thyroid function states. *Hormone and Metabolic Research* 36 (9): 650–53. doi: [10.1055/s-2004-825925](https://doi.org/10.1055/s-2004-825925).
- Laurberg, P., C. Cerqueira, L. Ovesen, L. B. Rasmussen, H. Perrild, S. Andersen, I. B. Pedersen, and A. Carle. 2010. Iodine intake as a determinant of thyroid disorders in populations. *Best Practice & Research. Clinical Endocrinology & Metabolism* 24 (1):13–27. doi: [10.1016/j.beem.2009.08.013](https://doi.org/10.1016/j.beem.2009.08.013).
- Lavado-Autric, R., E. Ausó, J. V. García-Velasco, M. d C. Arufe, F. Escobar del Rey, P. Berbel, and G. Morreale de Escobar. 2003. Early maternal hypothyroxinemia alters histogenesis and cerebral cortex cytoarchitecture of the progeny. *The Journal of Clinical Investigation* 111 (7):1073–82. doi: [10.1172/JCI16262](https://doi.org/10.1172/JCI16262).
- Leung, A. M., and L. E. Braverman. 2012. Iodine-induced thyroid dysfunction. *Current Opinion in Endocrinology, Diabetes and Obesity* 19 (5):414–9. doi: [10.1097/MED.0b013e3283565bb2](https://doi.org/10.1097/MED.0b013e3283565bb2).
- Leung, A. M., and L. E. Braverman. 2014. Consequences of excess iodine. *Nature Reviews. Endocrinology* 10 (3):136–42. doi: [10.1038/nrendo.2013.251](https://doi.org/10.1038/nrendo.2013.251).
- Levie, D., T. I. M. Korevaar, S. C. Bath, A. Dalmau-Bueno, M. Murcia, M. Espada, M. Dineva, J. M. Ibarluzea, J. Sunyer, H. Tiemeier, et al. 2018. Thyroid function in early pregnancy, child IQ, and autistic traits: A meta-analysis of individual participant data. *The Journal of Clinical Endocrinology & Metabolism* 103 (8):2967–79. doi: [10.1210/jc.2018-00224](https://doi.org/10.1210/jc.2018-00224).
- Li, H., S. Chen, L. Wu, H. Wang, K. Xiao, Y. Gao, Y. Li, H. Li, B. Xiao, and Y. Zhu. 2019. The effects of perineal disinfection on infant's oral microflora after transvaginal examination during delivery. *BMC Pregnancy and Childbirth* 19 (1):213. doi: [10.1186/s12884-019-2350-3](https://doi.org/10.1186/s12884-019-2350-3).
- Li, H. P., R. Brinkmeyer, W. L. Jones, S. Zhang, C. Xu, K. A. Schwehr, P. H. Santschi, D. I. Kaplan, and C. M. Yeager. 2011. Iodide accumulation by aerobic bacteria isolated from subsurface sediments of a 129I-contaminated aquifer at the Savannah River Site, South Carolina. *Applied and Environmental Microbiology* 77 (6):2153–60. doi: [10.1128/AEM.02164-10](https://doi.org/10.1128/AEM.02164-10).
- López-Muñoz, E., L. Mateos-Sánchez, G. E. Mejía-Terrazas, and S. E. Bedwell-Cordero. 2019. Hypothyroidism and isolated hypothyroxinemia in pregnancy, from physiology to the clinic. *Taiwanese Journal of Obstetrics & Gynecology* 58 (6):757–763. doi: [10.1016/j.tjog.2019.09.005](https://doi.org/10.1016/j.tjog.2019.09.005).
- Luo, Y., A. Kawashima, Y. Ishido, A. Yoshihara, K. Oda, N. Hiroi, T. Ito, N. Ishii, and K. Suzuki. 2014. Iodine excess as an environmental risk factor for autoimmune thyroid disease. *International Journal of Molecular Sciences* 15 (7):12895–912. doi: [10.3390/ijms150712895](https://doi.org/10.3390/ijms150712895).
- Maalouf, J., J. Barron, J. P. Gunn, K. Yuan, C. G. Perrine, and M. E. Cogswell. 2015. Iodized salt sales in the United States. *Nutrients* 7 (3):1691–95. doi: [10.3390/nu7031691](https://doi.org/10.3390/nu7031691).
- Manousou, S., L. Dahl, B. H. Thuesen, L. Hulthen, and H. N. Filipsson. 2017. Iodine deficiency and nutrition in Scandinavia. *Minerva Medica* 108 (2):147–58. doi: [10.23736/S0026-4806.16.04849-7](https://doi.org/10.23736/S0026-4806.16.04849-7).
- Mansourian, A. R. 2011. Metabolic pathways of tetraiodothyronine and triiodothyronine production by thyroid gland: A review of articles. *Pakistan Journal of Biological Sciences* 14 (1):1–12. <http://www.ncbi.nlm.nih.gov/pubmed/21913492>. doi: [10.3923/pjbs.2011.1.12](https://doi.org/10.3923/pjbs.2011.1.12).
- Marini, H., F. Polito, E. B. Adamo, A. Bitto, F. Squadrito, and S. Benvenga. 2012. Update on genistein and thyroid: An overall message of safety. *Frontiers in Endocrinology* 3:94. doi: [10.3389/fendo.2012.00094](https://doi.org/10.3389/fendo.2012.00094).
- Maris, P. 1995. Modes of action of disinfectants. *Revue Scientifique et Technique* 14 (1):47–55. doi: [10.20506/rst.14.1.829](https://doi.org/10.20506/rst.14.1.829).
- Marwaha, R. K., N. Tandon, M. K. Garg, A. Desai, R. Kanwar, A. Sastry, A. Narang, S. Arora, and K. Bhadra. 2012. Thyroid status two decades after salt iodization: Country-wide data in school children from India. *Clinical Endocrinology* 76 (6):905–10. doi: [10.1111/j.1365-2265.2011.04307.x](https://doi.org/10.1111/j.1365-2265.2011.04307.x).
- McDonnell, G., and A. D. Russell. 1999. Antiseptics and disinfectants: Activity, action, and resistance. *Clinical Microbiology Reviews* 12 (1): 147–79. <http://www.ncbi.nlm.nih.gov/pubmed/9880479>.
- Medicine, Institute of Iodine. 2001. *Dietary reference intakes for vitamin A, vitamin K, arsenic, boron, chromium, copper, iodine, iron, manganese, molybdenum, nickel, silicon, vanadium, and zinc*. Washington, DC: National Academy Press. [10.17226/10026](https://doi.org/10.17226/10026).
- Rousset, B., C. Dupuy, F. Miot, J. Dumont. 2015. Chapter 2 Thyroid hormone synthesis and secretion. In *Endotext*, ed. L. J. De Groot, G. Chrousos, K. Dungan, K. R. Feingold, A. Grossman, J. M. Hershman, C. Koch, et al. South Dartmouth, MA: MDText.com, Inc. <http://www.ncbi.nlm.nih.gov/pubmed/25905405>.
- Młyniec, K., M. Gawęł, U. Doboszewska, G. Starowicz, K. Pytko, C. L. Davies, and B. Budziszewska. 2015. Essential elements in depression and anxiety. Part II. *Pharmacological Reports* 67 (2):187–94. doi: [10.1016/j.pharep.2014.09.009](https://doi.org/10.1016/j.pharep.2014.09.009).
- Moleti, M., V. P. Lo Presti, M. C. Campolo, F. Mattina, M. Galletti, M. Mandolfino, M. A. Violi, G. Giorgianni, D. De Domenico, F. Trimarchi, et al. 2008. Iodine prophylaxis using iodized salt and risk of maternal thyroid failure in conditions of mild iodine deficiency. *The Journal of Clinical Endocrinology & Metabolism* 93 (7):2616–21. doi: [10.1210/jc.2008-0352](https://doi.org/10.1210/jc.2008-0352).
- Moleti, M., F. Trimarchi, G. Tortorella, A. Candia Longo, G. Giorgianni, G. Sturniolo, A. Alibrandi, and F. Vermiglio. 2016. Effects of maternal iodine nutrition and thyroid status on cognitive development in offspring: A pilot study. *Thyroid* 26 (2):296–305. doi: [10.1089/thy.2015.0336](https://doi.org/10.1089/thy.2015.0336).
- Moleti, M., F. Trimarchi, and F. Vermiglio. 2011. Doubts and concerns about isolated maternal hypothyroxinemia. *Journal of Thyroid Research* 2011:463029. doi: [10.4061/2011/463029](https://doi.org/10.4061/2011/463029).
- Mondal, S., K. Raja, U. Schweizer, and G. Mughesh. 2016. Chemistry and biology in the biosynthesis and action of thyroid hormones. *Angewandte Chemie* 55 (27):7606–30. doi: [10.1002/anie.201601116](https://doi.org/10.1002/anie.201601116).
- Moreno, J. C., and T. J. Visser. 2007. New phenotypes in thyroid dys-hormonogenesis: hypothyroidism due to DUOX2 mutations. *Endocrine Development* 10:99–117. doi: [10.1159/000106822](https://doi.org/10.1159/000106822). PMID17684392
- Mullur, R., Y. Y. Liu, and G. A. Brent. 2014. Thyroid hormone regulation of metabolism. *Physiological Reviews* 94 (2):355–82. doi: [10.1152/physrev.00030.2013](https://doi.org/10.1152/physrev.00030.2013).
- Murcia, M., M. Espada, J. Julvez, S. Llop, M.-J. Lopez-Espinosa, J. Vioque, M. Basterrechea, I. Riano, L. González, M. Alvarez-Pedrerol, et al. 2018. Iodine intake from supplements and diet during pregnancy and child cognitive and motor development: The INMA mother and child cohort study. *Journal of Epidemiology and Community Health* 72 (3):216–222. doi: [10.1136/jech-2017-209830](https://doi.org/10.1136/jech-2017-209830).
- Murray, C. W., S. K. Egan, H. Kim, N. Beru, and P. M. Bolger. 2008. US Food and Drug Administration's Total diet study: Dietary intake of perchlorate and iodine. *Journal of Exposure Science & Environmental Epidemiology* 18 (6):571–80. doi: [10.1038/sj.jes.7500648](https://doi.org/10.1038/sj.jes.7500648).
- Negro, R., O. P. Soldin, M. J. Obregon, and A. Stagnaro-Green. 2011. Hypothyroxinemia and pregnancy. *Endocrine Practice* 17 (3):422–29. doi: [10.4158/EP10309.RA](https://doi.org/10.4158/EP10309.RA).
- Nicola, J. P., C. Basquin, C. Portulano, A. Reyna-Neyra, M. Paroder, and N. Carrasco. 2009. The Na<sup>+</sup>/I<sup>-</sup> symporter mediates active iodide uptake in the intestine. *American Journal of Physiology. Cell Physiology* 296 (4):C654–62. doi: [10.1152/ajpcell.00509.2008](https://doi.org/10.1152/ajpcell.00509.2008).
- Nicola, J. P., M. Nazar, I. D. Mascanfroni, C. G. Pellizas, and A. M. Masini-Repiso. 2010. NF-KappaB P65 subunit mediates lipopolysaccharide-induced Na<sup>(+)</sup>/I<sup>(-)</sup> symporter gene expression by involving functional interaction with the paired domain transcription factor Pax8. *Molecular Endocrinology* 24 (9):1846–62. doi: [10.1210/me.2010-0102](https://doi.org/10.1210/me.2010-0102).
- Niwattisaiwong, S., K. D. Burman, and M. Li-Ng. 2017. Iodine deficiency: Clinical implications. *Cleveland Clinic Journal of Medicine* 84 (3):236–44. doi: [10.3949/ccjm.84a.15053](https://doi.org/10.3949/ccjm.84a.15053).

- O'Donnell, K. J., M. A. Rakeman, D. Zhi-Hong, C. Xue-Yi, Z. Y. Mei, N. DeLong, G. Brenner, M. Tai, W. Dong, and G. R. DeLong. 2002. Effects of iodine supplementation during pregnancy on child growth and development at school age. *Developmental Medicine and Child Neurology* 44 (2):76–81. doi: [10.1017/s0012162201001712](https://doi.org/10.1017/s0012162201001712).
- Olivieri, A., C. D. I. Cosmo, S. D. E. Angelis, R. D. A. Cas, P. Stacchini, A. Pastorelli, P. Vitti, and Prevention Regional Observatories for Goiter. 2017. The way forward in Italy for iodine. *Minerva Medica* 108 (2):159–68. doi: [10.23736/S0026-4806.17.04877-7](https://doi.org/10.23736/S0026-4806.17.04877-7).
- Opazo, M. C., A. Gianini, F. Pancetti, G. Azkcona, L. Alarcón, R. Lizana, V. Noches, P. A. Gonzalez, M. P. Marassi, M. Porto, et al. 2008. Maternal hypothyroxinemia impairs spatial learning and synaptic nature and function in the offspring. *Endocrinology* 149 (10): 5097–5106. doi: [10.1210/en.2008-0560](https://doi.org/10.1210/en.2008-0560).
- Opazo, M. C., P. A. Gonzalez, B. D. Flores, L. F. Venegas, E. A. Alborno, P. Cisternas, K. Bohmwald, P. A. Nieto, S. M. Bueno, A. M. Kalergis, et al. 2017. Gestational hypothyroxinemia imprints a switch in the capacity of astrocytes and microglial cells of the offspring to react in inflammation. *Molecular Neurobiology* 55 (5): 4373–4387. doi: [10.1007/s12035-017-0627-y](https://doi.org/10.1007/s12035-017-0627-y).
- Opazo, M. C., H. Haengen, K. Bohmwald, L. F. Venegas, H. Boudin, A. A. Elorza, F. Simon, C. Fardella, S. M. Bueno, A. M. Kalergis, et al. 2017. Imprinting of maternal thyroid hormones in the offspring. *International Reviews of Immunology* 36 (4):240–55. doi: [10.1080/08830185.2016.1277216](https://doi.org/10.1080/08830185.2016.1277216).
- Palaniappan, S., L. Shanmughavelu, H. K. Prasad, S. Subramaniam, N. Krishnamoorthy, and L. Lakkappa. 2017. Improving iodine nutritional status and increasing prevalence of autoimmune thyroiditis in children. *Indian Journal of Endocrinology and Metabolism* 21 (1): 85–89. doi: [10.4103/2230-8210.195996](https://doi.org/10.4103/2230-8210.195996).
- Patel, J., K. Landers, H. Li, R. H. Mortimer, and K. Richard. 2011. Delivery of maternal thyroid hormones to the fetus. *Trends in Endocrinology and Metabolism: TEM* 22 (5):164–70. doi: [10.1016/j.tem.2011.02.002](https://doi.org/10.1016/j.tem.2011.02.002).
- Patrick, L. 2008. Iodine: Deficiency and therapeutic considerations. *Alternative Medicine Review* 13 (2):116–27. <http://www.ncbi.nlm.nih.gov/pubmed/18590348>.
- Pearce, E. N., J. H. Lazarus, R. Moreno-Reyes, and M. B. Zimmermann. 2016. Consequences of iodine deficiency and excess in pregnant women: An overview of current knowns and unknowns. *The American Journal of Clinical Nutrition* 104 (suppl\_3):918S–923S. doi: [10.3945/ajcn.115.110429](https://doi.org/10.3945/ajcn.115.110429).
- Pearce, E. N., S. Pino, X. He, H. R. Bazrafshan, S. L. Lee, and L. E. Braverman. 2004. Sources of dietary iodine: Bread, cows' milk, and infant formula in the Boston area. *The Journal of Clinical Endocrinology and Metabolism* 89 (7):3421–24. doi: [10.1210/jc.2003-032002](https://doi.org/10.1210/jc.2003-032002).
- Peeters, R. P. 2017. Subclinical hypothyroidism. *The New England Journal of Medicine* 376 (26):2556–65. doi: [10.1056/NEJMcp1611144](https://doi.org/10.1056/NEJMcp1611144).
- Pharoah, P. O. D., I. H. Butfield, and B. S. Hetzel. 1971. Neurological damage to the fetus resulting from severe iodine deficiency during pregnancy. *The Lancet* 297 (7694):308–10. doi: [10.1016/S0140-6736\(71\)91040-3](https://doi.org/10.1016/S0140-6736(71)91040-3).
- Portulano, C., M. Paroder-Belenitsky, and N. Carrasco. 2014. The Na<sup>+</sup>/I<sup>-</sup> symporter (NIS): Mechanism and medical impact. *Endocrine Reviews* 35 (1):106–49. doi: [10.1210/er.2012-1036](https://doi.org/10.1210/er.2012-1036).
- Ragusa, F., P. Fallahi, G. Elia, D. Gonnella, S. R. Paparo, C. Giusti, L. P. Churilov, S. M. Ferrari, and A. Antonelli. 2019. Hashimoto's thyroiditis: Epidemiology, pathogenesis, clinic and therapy. *Best Practice and Research: Clinical Endocrinology and Metabolism* 33 (6): 101367. doi: [10.1016/j.beem.2019.101367](https://doi.org/10.1016/j.beem.2019.101367).
- Ravera, S., A. Reyna-Neyra, G. Ferrandino, L. M. Amzel, and N. Carrasco. 2017. The sodium/iodide symporter (NIS): Molecular physiology and preclinical and clinical applications. *Annual Review of Physiology* 79:261–89. doi: [10.1146/annurev-physiol-022516-034125](https://doi.org/10.1146/annurev-physiol-022516-034125).
- Rayman, M. P. 2012. Selenium and human health. *The Lancet* 379 (9822):1256–68. doi: [10.1016/S0140-6736\(11\)61452-9](https://doi.org/10.1016/S0140-6736(11)61452-9).
- Redman, K., T. Ruffman, P. Fitzgerald, and S. Skeaff. 2016. Iodine deficiency and the brain: Effects and mechanisms. *Critical Reviews in Food Science and Nutrition* 56 (16):2695–713. doi: [10.1080/10408398.2014.922042](https://doi.org/10.1080/10408398.2014.922042).
- Riesco-Eizaguirre, G., L. Wert-Lamas, J. Perales-Paton, A. Sastre-Perona, L. P. Fernandez, and P. Santisteban. 2015. The MiR-146b-3p/PAX8/NIS regulatory circuit modulates the differentiation phenotype and function of thyroid cells during carcinogenesis. *Cancer Research* 75 (19):4119–30. doi: [10.1158/0008-5472.CAN-14-3547](https://doi.org/10.1158/0008-5472.CAN-14-3547).
- Ris-Stalpers, C., and H. Bikker. 2010. Genetics and phenomics of hypothyroidism and goiter due to TPO mutations. *Molecular and Cellular Endocrinology* 322 (1-2):38–43. doi: [10.1016/j.mce.2010.02.008](https://doi.org/10.1016/j.mce.2010.02.008).
- Rodondi, N., A. B. Newman, E. Vittinghoff, N. de Rekeneire, S. Satterfield, T. B. Harris, and D. C. Bauer. 2005. Subclinical hypothyroidism and the risk of heart failure, other cardiovascular events, and death. *Archives of Internal Medicine* 165 (21):2460–66. doi: [10.1001/archinte.165.21.2460](https://doi.org/10.1001/archinte.165.21.2460).
- Rohner, F., M. Zimmermann, P. Jooste, C. Pandav, K. Caldwell, R. Raghavan, and D. J. Raiten. 2014. Biomarkers of nutrition for development-iodine review. *The Journal of Nutrition* 144 (8):1322S–42S. doi: [10.3945/jn.113.181974](https://doi.org/10.3945/jn.113.181974).
- Román, G. C. 2007. Autism: Transient in utero hypothyroxinemia related to maternal flavonoid ingestion during pregnancy and to other environmental antithyroid agents. *Journal of the Neurological Sciences* 262 (1-2):15–26. doi: [10.1016/j.jns.2007.06.023](https://doi.org/10.1016/j.jns.2007.06.023).
- Román, G. C., A. Ghassabian, J. J. Bongers-Schokking, V. W. V. Jaddoe, A. Hofman, Y. B. Rijke, F. C. Verhulst, and H. Tiemeier. 2013. Association of gestational maternal hypothyroxinemia and increased autism risk. *Annals of Neurology* 74 (5):733–42. doi: [10.1002/ana.23976](https://doi.org/10.1002/ana.23976).
- Sarah Davis, O., S. Margaret, M. B. Jennifer, and B. Betty. 2012. Use of iodized salt in processed foods in select countries around the world and the role of food processors. *Comprehensive Reviews in Food Science and Food Safety* 11 (2):233–84. doi: [10.1111/j.1541-4337.2011.00182.x](https://doi.org/10.1111/j.1541-4337.2011.00182.x).
- Schmutzler, C., I. Gotthardt, P. J. Hofmann, B. Radovic, G. Kovacs, L. Stemmler, I. Nobis, A. Bacinski, B. Mentrup, P. Ambrugger, et al. 2007. Endocrine disruptors and the thyroid gland—A combined in vitro and in vivo analysis of potential new biomarkers. *Environmental Health Perspectives* 115 (Suppl 1):77–83. doi: [10.1289/ehp.9369](https://doi.org/10.1289/ehp.9369).
- Schomburg, L., and J. Köhrle. 2008. On the importance of selenium and iodine metabolism for thyroid hormone biosynthesis and human health. *Molecular Nutrition & Food Research* 52 (11): 1235–46. doi: [10.1002/mnfr.200700465](https://doi.org/10.1002/mnfr.200700465). PMID: 18686295
- Schug, T. T., R. Abagyan, B. Blumberg, T. J. Collins, D. Crews, P. L. DeFur, S. M. Dickerson, T. M. Edwards, A. C. Gore, L. J. Guille, et al. 2013. Designing endocrine disruption out of the next generation of chemicals. *Green Chemistry* 15 (1):181–98. doi: [10.1039/C2GC35055F](https://doi.org/10.1039/C2GC35055F).
- Seo, G. H., and J. H. Chung. 2015. Incidence and prevalence of overt hypothyroidism and causative diseases in Korea as determined using claims data provided by the health insurance review and assessment service. *Endocrinology and Metabolism* 30 (3):288–96. doi: [10.3803/EnM.2015.30.3.288](https://doi.org/10.3803/EnM.2015.30.3.288).
- Shen, H., J. Han, Y. Li, C. Lu, J. Zhou, Y. Li, and X. Su. 2019. Different host-specific responses in thyroid function and gut microbiota modulation between diet-induced obese and normal mice given the same dose of iodine. *Applied Microbiology and Biotechnology* 103 (8):3537–47. doi: [10.1007/s00253-019-09687-1](https://doi.org/10.1007/s00253-019-09687-1).
- Shirai, J., T. Kanno, Y. Tsuchiya, S. Mitsubayashi, and R. Seki. 2000. Effects of chlorine, iodine, and quaternary ammonium compound disinfectants on several exotic disease viruses. *The Journal of Veterinary Medical Science* 62 (1):85–92. doi: [10.1292/jvms.62.85](https://doi.org/10.1292/jvms.62.85).
- Siffo, S., E. Adrover, C. E. Citterio, M. B. Miras, V. A. Balbi, A. Chiesa, J. Weill, G. Sobrero, V. G. González, P. Papendieck, et al. 2018. Molecular analysis of thyroglobulin mutations found in patients with goiter and hypothyroidism. *Molecular and Cellular Endocrinology* 473:1–16. doi: [10.1016/j.mce.2017.12.009](https://doi.org/10.1016/j.mce.2017.12.009).

- Silbereis, J. C., S. Pochareddy, Y. Zhu, M. Li, and N. Sestan. 2016. The cellular and molecular landscapes of the developing human central nervous system. *Neuron* 89 (2):248–268. doi: [10.1016/j.neuron.2015.12.008](https://doi.org/10.1016/j.neuron.2015.12.008).
- Simmons, S. R., and M. L. Karnovsky. 1973. Iodinating ability of various leukocytes and their bactericidal activity. *The Journal of Experimental Medicine* 138 (1):44–63. doi: [10.1084/jem.138.1.44](https://doi.org/10.1084/jem.138.1.44).
- Sirakov, M., and M. Plateroti. 2011. The thyroid hormones and their nuclear receptors in the gut: From developmental biology to cancer. *Biochim Biophys Acta* 1812(8):938–46. doi: [10.1016/j.bbadis.2010.12.020](https://doi.org/10.1016/j.bbadis.2010.12.020). PMID: 21194566.
- Sriwilaijaroen, N., P. Wilairat, H. Hiramatsu, T. Takahashi, T. Suzuki, M. Ito, Y. Ito, M. Tashiro, and Y. Suzuki. 2009. Mechanisms of the action of povidone-iodine against human and avian influenza A viruses: Its effects on hemagglutination and sialidase activities. *Virology Journal* 6:124. doi: [10.1186/1743-422X-6-124](https://doi.org/10.1186/1743-422X-6-124).
- Stiles, J., and T. L. Jernigan. 2010. The basics of brain development. *Neuropsychology Review* 20 (4):327–48. doi: [10.1007/s11065-010-9148-4](https://doi.org/10.1007/s11065-010-9148-4).
- Sun, X., Z. Shan, and W. Teng. 2014. Effects of increased iodine intake on thyroid disorders. *Endocrinology and Metabolism* 29 (3):240–47. doi: [10.3803/EnM.2014.29.3.240](https://doi.org/10.3803/EnM.2014.29.3.240).
- Taylor, P. N., D. Albrecht, A. Scholz, G. Gutierrez-Buey, J. H. Lazarus, C. M. Dayan, and O. E. Okosieme. 2018. Global epidemiology of hyperthyroidism and hypothyroidism. *Nature Reviews. Endocrinology* 14 (5):301–16. doi: [10.1038/nrendo.2018.18](https://doi.org/10.1038/nrendo.2018.18).
- Taylor, R. M., S. M. Fealy, A. Bisquera, R. Smith, C. E. Collins, T. J. Evans, and A. J. Hure. 2017. Effects of nutritional interventions during pregnancy on infant and child cognitive outcomes: A systematic review and meta-analysis. *Nutrients* 9 (11):1265. doi: [10.3390/nu9111265](https://doi.org/10.3390/nu9111265).
- Teas, J., S. Pino, A. Critchley, and L. E. Braverman. 2004. Variability of iodine content in common commercially available edible seaweeds. *Thyroid* 14 (10):836–41. doi: [10.1089/thy.2004.14.836](https://doi.org/10.1089/thy.2004.14.836).
- Thompson, W., G. Russell, G. Baragwanath, J. Matthews, B. Vaidya, and J. Thompson-Coon. 2018. Maternal thyroid hormone insufficiency during pregnancy and risk of neurodevelopmental disorders in offspring: A systematic review and meta-analysis. *Clinical Endocrinology* 88 (4):575–84. doi: [10.1111/cen.13550](https://doi.org/10.1111/cen.13550).
- Valeix, P., P. Faure, S. Peneau, C. Estaquio, S. Hercberg, and S. Bertrais. 2009. Lifestyle factors related to iodine intakes in French adults. *Public Health Nutrition* 12 (12):2428–37. doi: [10.1017/S1368980009005515](https://doi.org/10.1017/S1368980009005515).
- Vanderpump, M. P. 2011. The epidemiology of thyroid disease. *British Medical Bulletin* 99:39–51. doi: [10.1093/bmb/ldr030](https://doi.org/10.1093/bmb/ldr030).
- Verheesen, R. H., and C. M. Schweitzer. 2008. Iodine deficiency, more than cretinism and goiter. *Medical Hypotheses* 71 (5):645–48. doi: [10.1016/j.mehy.2008.06.020](https://doi.org/10.1016/j.mehy.2008.06.020).
- Vermiglio, F., V. P. Lo Presti, M. Moleti, M. Sidoti, G. Tortorella, G. Scaffidi, M. G. Castagna, F. Mattina, M. A. Violi, A. Crisà, et al. 2004. Attention deficit and hyperactivity disorders in the offspring of mothers exposed to mild-moderate iodine deficiency: A possible novel iodine deficiency disorder in developed countries. *The Journal of Clinical Endocrinology and Metabolism* 89 (12):6054–60. doi: [10.1210/jc.2004-0571](https://doi.org/10.1210/jc.2004-0571).
- Virili, C., and M. Centanni. 2017. “With a little help from my friends”—The role of microbiota in thyroid hormone metabolism and enterohepatic recycling. *Molecular and Cellular Endocrinology* 458:39–43. doi: [10.1016/j.mce.2017.01.053](https://doi.org/10.1016/j.mce.2017.01.053).
- Visser, W. E., E. C. H. Friesema, and T. J. Visser. 2011. Minireview: Thyroid hormone transporters: The knowns and the unknowns. *Molecular Endocrinology* 25 (1):1–14. doi: [10.1210/me.2010-0095](https://doi.org/10.1210/me.2010-0095).
- Vought, R. L., F. A. Brown, K. H. Sibinovic, and E. G. McDaniel. 1972. Effect of changing intestinal bacterial flora on thyroid function in the rat. *Hormone and Metabolic Research = Hormon- Und Stoffwechselforschung = Hormones et Metabolisme* 4 (1):43–47. doi: [10.1055/s-0028-1094095](https://doi.org/10.1055/s-0028-1094095).
- Walker, S. P., T. D. Wachs, J. M. Gardner, B. Lozoff, G. A. Wasserman, E. Pollitt, J. A. Carter, and Group International Child Development Steering. 2007. Child development: Risk factors for adverse outcomes in developing countries. *Lancet* 369 (9556):145–57. doi: [10.1016/S0140-6736\(07\)60076-2](https://doi.org/10.1016/S0140-6736(07)60076-2).
- Wang, K., Y. N. Sun, J. Y. Liu, L. Zhang, Y. Ye, L. X. Lin, Y. Q. Yan, and Z. P. Chen. 2009. The impact of iodine excess on thyroid hormone biosynthesis and metabolism in rats. *Biological Trace Element Research* 130 (1):72–85. doi: [10.1007/s12011-009-8315-z](https://doi.org/10.1007/s12011-009-8315-z).
- Weiss, R. V., and R. Clapauch. 2014. Female infertility of endocrine origin. *Arquivos Brasileiros de Endocrinologia & Metabologia* 58 (2):144–52. <http://www.ncbi.nlm.nih.gov/pubmed/24830591>.
- Weng, H. X., J. K. Weng, W. B. Yong, X. W. Sun, and H. Zhong. 2003. Capacity and degree of iodine absorbed and enriched by vegetable from soil. *Journal of Environmental Sciences* 15 (1):107–11. <http://www.ncbi.nlm.nih.gov/pubmed/12602613>.
- Whitehead, D. C. 1984. The distribution and transformation of iodine in the environment. *Environment International* 10 (4):321–39. doi: [10.1016/0160-4120\(84\)90139-9](https://doi.org/10.1016/0160-4120(84)90139-9).
- WHO. 2004. Iodine status worldwide: WHO Global Database on Iodine De Cency. Geneva.
- WHO. 2008. WHO, UNICEF and ICCIDD. Assessment of the iodine deficiency disorders and monitoring their elimination. [http://www.who.int/nutrition/publications/micronutrients/iodine\\_deficiency/9789241595827/en/](http://www.who.int/nutrition/publications/micronutrients/iodine_deficiency/9789241595827/en/).
- Zbigniew, S. 2017. Role of iodine in metabolism. *Recent Patents on Endocrine, Metabolic & Immune Drug Discovery* 10 (2):123–26. doi: [10.2174/1872214811666170119110618](https://doi.org/10.2174/1872214811666170119110618).
- Zealand, Food Standards Australia New. 2015. Iodine in food and iodine requirements. <http://www.foodstandards.gov.au/consumer/nutrition/iodinefood/Pages/default.aspx>.
- Zhao, F., J. Feng, J. Li, L. Zhao, Y. Liu, H. Chen, Y. Jin, B. Zhu, and Y. Wei. 2018. Alterations of the gut microbiota in Hashimoto’s thyroiditis patients. *Thyroid* 28 (2):175–86.
- Zhou, L., X. Li, A. Ahmed, D. Wu, L. Liu, J. Qiu, Y. Yan, M. Jin, and Y. Xin. 2014. Gut microbe analysis between hyperthyroid and healthy individuals. *Current Microbiology* 69 (5):675–80. doi: [10.1007/s00284-014-0640-6](https://doi.org/10.1007/s00284-014-0640-6).
- Zhou, S. J., S. A. Skeaff, P. Ryan, L. W. Doyle, P. J. Anderson, L. Kornman, A. J. McPhee, L. N. Yelland, and M. Makrides. 2015. The effect of iodine supplementation in pregnancy on early childhood neurodevelopment and clinical outcomes: Results of an aborted randomised placebo-controlled trial. *Trials* 16:563. doi: [10.1186/s13063-015-1080-8](https://doi.org/10.1186/s13063-015-1080-8).
- Zhu, Y. G., Y. Z. Huang, Y. Hu, and Y. X. Liu. 2003. Iodine uptake by spinach (*Spinacia oleracea* L.) plants grown in solution culture: Effects of iodine species and solution concentrations. *Environment International* 29 (1):33–37. doi: [10.1016/S0160-4120\(02\)00129-0](https://doi.org/10.1016/S0160-4120(02)00129-0).
- Zimmermann, M. B., P. L. Jooste, and C. S. Pandav. 2008. Iodine-deficiency disorders. *Lancet* 372 (9645):1251–62. doi: [10.1016/S0140-6736\(08\)61005-3](https://doi.org/10.1016/S0140-6736(08)61005-3).
- Zoltowska, D., Y. Agrawal, S. Patri, S. Aggarwal, C. S. Reddy, N. Sareen, J. Kalavakunta, and V. Gupta. 2018. Association between hypothyroidism and Takotsubo cardiomyopathy: Analysis of nationwide inpatient sample database. *Reviews on Recent Clinical Trials* 13 (3):222–5. doi: [10.2174/1574887113666180402144600](https://doi.org/10.2174/1574887113666180402144600).