



Bringing the digestibility of prebiotics into focus: update of carbohydrate digestion models

Alvaro Ferreira-Lazarte , F. Javier Moreno & Mar Villamiel

To cite this article: Alvaro Ferreira-Lazarte , F. Javier Moreno & Mar Villamiel (2020): Bringing the digestibility of prebiotics into focus: update of carbohydrate digestion models, Critical Reviews in Food Science and Nutrition, DOI: [10.1080/10408398.2020.1798344](https://doi.org/10.1080/10408398.2020.1798344)

To link to this article: <https://doi.org/10.1080/10408398.2020.1798344>



Published online: 03 Aug 2020.



Submit your article to this journal [↗](#)



View related articles [↗](#)



View Crossmark data [↗](#)

REVIEW



Bringing the digestibility of prebiotics into focus: update of carbohydrate digestion models

Alvaro Ferreira-Lazarte, F. Javier Moreno, and Mar Villamiel

Instituto de Investigación en Ciencias de la Alimentación (CIAL), CSIC-UAM, Madrid, Spain

ABSTRACT

Oro-gastrointestinal digestion of dietary carbohydrates involves up to six different carbohydrases in a multistage process. Enzymes from the small intestinal brush border membrane play a major role in the digestibility of these substrates. However, to date, the inclusion of these small intestinal enzymes has been dismissed in most *in vitro* studies carried out, despite their importance in the degradation of carbohydrates. Several *in vitro* and *in vivo* studies have demonstrated the capability of brush border enzymes to degrade certain “non-digestible” carbohydrates to a different extent depending on their structural composition (monomeric composition, glycosidic linkage, etc.). In this sense, considering the available evidence, mucosal disaccharidases embedded in the small intestinal brush border membrane vesicles must be considered in addition to α -amylases; therefore, new approaches for the evaluation of the digestibility of carbohydrates have been recently reported. These new methods based on the utilization of the small intestinal enzymes present in the brush border membrane aim to fulfill the final and key step of the digestion of carbohydrates in the small intestine. Here, rat small intestinal extract enzymes as well as brush border membrane vesicles from pig have emerged as very reliable and useful tools to evaluate carbohydrate digestion. Thus, this review aims to go briefly through the most relevant digestion methods for carbohydrates that are currently available and to highlight the new improved methods, which include mammalian intestinal enzymes, and their current use in the evaluation of the digestibility of prebiotics.

KEYWORDS

Brush border membrane vesicles; carbohydrate digestion; mammalian intestinal enzymes; prebiotics; small intestinal digestion

Introduction

In the last decades, interest toward the modulation of the human microbiota has undergone huge growth due to its strong relationship with the regulation of health and well-being. To date, several illnesses have been related with the incorrect function or alterations in the composition of the human microbiota (Guinane and Cotter 2013; Jandhyala et al. 2015; Villanueva-Millán, Pérez-Matute, and Oteo 2015; Lerner, Neidhöfer, and Matthias 2017). One of the most commonly used approaches to regulate homeostasis, maintaining the proper behavior of microbiota, is the use of “prebiotics” which are selectively fermented in the gut, giving rise to positive changes not only in the singular ecosystem that inhabits the colon but also at a systemic level (Gibson et al. 2017). Thereby, since prebiotics were first defined, relevant advances have been made in the development of new prebiotic compounds. In this regard, one focus has pointed toward obtaining molecules with new structures that provide new and improved properties (Gullón et al. 2013; Díez-Municio et al. 2014; Míguez et al. 2016).

However, despite the great progress that has been made toward prebiotics and their benefits on human health, few studies have been made concerning changes to them during

their passage through the upper gastrointestinal tract, even when structural differences could substantially affect their properties (Laparra et al. 2014; Li et al. 2015; Fernández et al. 2018; Larsen et al. 2019). The scarce information in this regard and the lack of specific methods to accurately determine their digestion, compel the use of general harmonized protocols such as the different AOAC or InfoGest methods, which could not reflect the whole range of enzymatic activities during the digestion of carbohydrates in the human small intestine.

The purpose of the present review is to compile the most important methods focused on carbohydrate digestion paying special attention to those based on the utilization of small intestinal extract from mammals in order to evaluate the potential partial digestion of prebiotics.

Resistance of prebiotic carbohydrates to the gastrointestinal digestion

One of the classical requirements for compounds to be considered as prebiotics is their resistance to upper gastrointestinal tract digestion. In this sense, ever since prebiotics were first defined in 1995, investigations have focused on their effect on the microbiota activity and/or composition.

However, despite the generally accepted concept that prebiotics pass through the upper gastrointestinal tract without modifications, few efforts have been made toward the study of the resistance of these compounds to digestion and conditions in the upper gastrointestinal stage. It is important whether or not chemical or structural changes occur when they are exposed to this environment, because even minor differences could substantially affect their properties and, therefore, their impact on colonic microbiota (Laparra et al. 2014; Li et al. 2015; Fernández et al. 2018; Larsen et al. 2019).

Human oro-gastrointestinal degradation of carbohydrates constitutes a multistage process starting in the oral cavity, where the food is converted into a homogenous mass during mastication, with very scarce hydrolysis by the α -amylase present in the saliva. The main carbohydrate degradation and absorption occurs in the small intestine, being the main site for carbohydrate digestion involving the α -amylase secreted by the pancreas. In addition, brush border membranes of the intestinal mucosa contain several key enzymes for carbohydrate digestion present as multienzyme complexes, i.e., sucrase-isomaltase, lactase-phlorizin hydrolase, maltase-glucoamylase and trehalase being responsible for the final stage of luminal digestion (Figure 1) (Holmes and Lobley 1989; Hooton et al. 2015; Picariello, Ferranti, and Addeo 2016). Therefore, dietary carbohydrate digestion involves up to six different carbohydrases produced by three different organs (Table 1). Moreover, the mammalian upper gastrointestinal tract that contains a variety of distinct microbial populations could also affect the resistance of these substrates. In this sense, the small intestine, which has more acidic conditions and higher levels of oxygen and antimicrobials than the colon, is dominated by fast-growing facultative anaerobes, such as Lactobacillaceae or Enterobacteraceae, that tolerate the combined effect of antimicrobials and bile salts (Donaldson, Lee, and Mazmanian 2016). Moreover, conditions such as the presence of oxygen, antimicrobial compounds (bile salts) or pH limit the bacterial density below 10⁴ colony-forming units (cfu/g) and only at the distal end of the small intestine, in the terminal ileum, bacterial densities reach levels similar to those found in the large intestine. With all these characteristics, it is not misplaced to consider that prebiotic compounds could suffer changes that could affect their properties when they reach the colon to be fermented.

Models for the assessment of digestibility and standardized protocols

To date, only scarce and fragmented information on their passage through the small intestine is available. As is well-known, *in vivo* feeding methods, using animals or humans, usually provide the most accurate results. In this sense, few *in vivo* studies have provided evidence of the partial degradation of prebiotic compounds after their passage through the small intestine. Holloway, Tasman-Jones, and Maher (1983) pointed out a recovery of 68% of pectin at the human terminal ileum in an ileostomy study, whereas Saito

et al. (2005) observed a recovery rate of 90% in the terminal ileum after colonic intubation of volunteers. Moreover, Hernández-Hernández et al. (2012) found 13–53% of small intestinal digestibility of purified galactooligosaccharides (GOS, with a degree of polymerization from 3 onwards) in an *in vivo* study with rats. However, these methods have several drawbacks, such as high costs, ethical constraints, limitations in sampling from the small intestine, and the fact that they are time consuming, which explains why much effort has been devoted to the development of *in vitro* procedures. *In vitro* digestion models may provide a very useful alternative to animal and human models by rapidly screening food ingredients. In this sense, the most frequently used approach to measure the digestion process is by continuous or static simulation of human physiological conditions as much as possible, taking into account the presence of digestive enzymes and their concentrations, pH, temperature, digestion time, and salt concentrations, among other factors (Table 2) (Kopf-Bolanz et al. 2012; Minekus et al. 2014). Very recently, Hernández-Hernández (2019) carried out an overview of different methodologies used for the assessment of the digestibility of carbohydrates.

Continuous *in vitro* digestion methods

Although the majority of models reported in the literature are static simulators, some computer-controlled multi-compartmental continuous system models can overcome some limitations present in static models allowing the simulation of dynamic aspects of digestion, such as the transport of digested components, variable enzyme concentrations, pH changes, peristaltic movements, continuous changes, and secretion flow rates (Ouwehand and Vaughan 2006). To date, these multi-compartmental models that mimic the different gastrointestinal stages in a continuous flow represent the most advanced attempt at simulating interdependent physiological functions within the stomach lumen, small intestine and human gut (Molly, Vande Woestyne, and Verstraete 1993; Minekus et al. 1995; Barroso et al. 2015; Dupont et al. 2019). However, these models require highly complex and substantial hardware and software and are still expensive to set up and maintain in a regular laboratory; therefore, static models have an economic and large throughput advantage.

Ferreira-Lazarte, Moreno, et al. (2019) studied the digestibility and fermentability of citrus pectin in a Dynamic Gastrointestinal Simulator (simgi®) and observed a 12% loss of pectin during its passage through the stomach and the small intestine after the evaluation of the remaining polymer by HPLC-ELSD analysis. Some authors attributed the loss of pectin during digestion to the effect of the bacteria that can be present in the digestive tract (Saito et al. 2005). As in the prototype of Ferreira-Lazarte, Gallego-Lobillo, et al. (2019), bacteria are only confined in the colon compartments, and the modifications observed in the structure of pectin could be attributed to the chemical effects of bile salts and pancreatic fluids.




Figure 1. Brush border membrane enzymes presence and formation of vesicles in the small intestine. Colored proteins inside the membrane bilayer (green and yellow), represent the brush border enzymes with peptide (gold and pale blue) and glycosylphosphatidylinositol (dark blue and purple) anchors. BBMVs can be released into the adjacent periapical space, however, it is likely that the mucus layer overlying the epithelia retards the egress of BBMVs from the periapical space into the lumen so that a significant amount of BBMVs would remain in close proximity to the intestinal mucosa.

Furthermore, Bellmann et al. (2018) presented a dynamic digestion model focused on carbohydrates, called TIMcarbo, which was based on the TNO GastroIntestinal Model (TIM) (Minekus et al. 1995). This model mimics carbohydrate uptake by the epithelium through dialysis of the products of digestion by a commercial rat small intestinal extract enzyme and the addition of a bacterial lactase given the low activity found in these extracts. Therefore, TIMcarbo would represent a highly relevant tool to evaluate the digestion of whole foods and meals where the physical changes of the digesta (e.g.,

viscosity or particle size reduction) are key determinants of the rate of carbohydrate digestion and absorption *in vivo* (Lifschitz, Grusak, and Butte 2002; Turnbull, Baxter, and Johnson 2005). However, this modified method is not so suitable for screening studies or for the evaluation of low available amounts of carbohydrates, which is typically a limiting factor when the digestibility of novel carbohydrates produced at laboratory scale needs to be assessed (Hernández-Hernández et al. 2019). In this sense, static *in vitro* methods can constitute a good alternative.

Table 1. Human carbohydrases involved in dietary carbohydrate digestion.

Digestive carbohydrases	Type of enzyme	Glycoside hydrolase family ^a	Production organ/main site of digestion	Glycosidic linkage specificity	Main substrates ^b	Main products ^b
Salivary α -amylase ^c	Secreted (α -glucosidase)	13	Salivary gland/mouth	Glc α (1 \rightarrow 4)Glc	Starch; linear maltooligosaccharides (n > 6)	Maltose; maltotriose; α -dextrins
Pancreatic α -amylase ^c	Secreted (α -glucosidase)	13	Pancreas/small intestine	Glc α (1 \rightarrow 4)Glc	Starch; linear maltooligosaccharides (n > 6)	Maltose; maltotriose; α -dextrins
Sucrase-isomaltase	Mucosal (α -glucosidase)	31	Small intestine (brush border membrane)/small intestine	Glc α (1 \leftrightarrow 2) β Fru Glc α (1 \rightarrow 4)Glc Glc α (1 \rightarrow 6)Glc	Sucrose; isomaltose; maltose; maltotriose; α -dextrins	Glucose; fructose
Maltase-glucoamylase	Mucosal (α -glucosidase)	31	Small intestine (brush border membrane)/small intestine	Glc α (1 \rightarrow 4)Glc Glc α (1 \rightarrow 6)Glc	Linear and branched maltooligosaccharides (n = 2–9)	Glucose
Lactase-phlorizin hydrolase	Mucosal (α -glycosidase)	1	Small intestine (brush border membrane)/small intestine	Glc α (1 \rightarrow 4)Gal Glc α (1 \rightarrow 4)Glc	Lactose, cellobiose, celotriose, cellulose	Glucose; galactose
Trehalase	Mucosal (α -glucosidase)	37	Small intestine (brush border membrane)/small intestine	Glc α (1 \leftrightarrow 1) α Glc	Trehalose	Glucose

Reproduced from Hernández-Hernández et al. (2019).

^aAccording to CAZy database (<http://www.cazy.org/>; Lombard et al. 2014).

^bBased on and updated from Alpers (2003).

^cHuman salivary and pancreatic amylases have 94% amino acid identity although they are encoded by different genes (Meisenberg and Simmons 2016).

Table 2. Overview on the *in vitro* intestinal digestion studies based on simulated fluids (without mucosal enzymes) for carbohydrates.

Carbohydrates evaluated	Digestion model	Digestion and fermentation stages/conditions	Degradation	References
Fructooligosaccharides	Static (InfoGest)	Oral, gastric and small intestinal digestion (pancreatin and porcine bile extract)	Slight degradation in GF ₃	Nobre et al. (2018)
Hydrolyzed Curdlan ((1→3)- β -D-glucan oligosaccharides)	Static (InfoGest)	Oral, gastric and small intestinal digestion (pancreatin and porcine bile extract)	No	Shi et al. (2018)
Fructooligosaccharides	Dynamic (SHIME)	Gastric, small intestinal digestion (pancreatin and bile salts) and colon (ascending transverse and descending)	No reported	Sivieri et al. (2014)
Citrus Pectin	Dynamic (simgi®)	Gastric, small intestinal digestion (pancreatin and bile salts) and colon (ascending, transverse and descending)	Slight Mw decrease after intestinal digestion	Ferreira-Lazarte, Moreno, et al. (2019)
Galactose rich oligosaccharides from Rhamnogalacturonan-I	Dynamic (TIM-1)	Gastric and Small Intestine (duodenum, jejunum, ileum - pancreatin, bile salts and dialysis)	~ 80% resistance	Khodaei et al. (2016)
Polysaccharide from <i>Hericium ernaceus</i> ((1→6)- α -D-galactopyranosyl backbone)	Static	Gastric, and small intestinal digestion (pancreatin and bile salts)	Mw decrease and free monosaccharides increase	Yang et al. (2018)
Polysaccharides from <i>Gracilaria rubra</i>	Static	Oral, gastric and small intestinal digestion (pancreatin and porcine bile extract)	No	Di et al. (2018)
Polysaccharides from <i>Rosa roxburgui</i>	Static	Oral, gastric and small intestinal digestion (pancreatin and porcine bile extract)	Slight Mw decrease after stomach and intestinal digestion	Wang, Li, et al. (2019)
Polysaccharides from Fuzhuan brick tea	Static	Oral, gastric and small intestinal digestion (pancreatin and porcine bile extract)	No	Chen, Xie, et al. (2018)
Polysaccharides from brown seaweed (<i>Ascophyllum nodosum</i>) and <i>Coralline pilulifera</i>	Static	Oral, gastric and small intestinal digestion (pancreatin and porcine bile extract)	No	Chen, Xu, et al. (2018) Wang, Chen, et al. (2019)
Polysaccharides from <i>Aloe vera</i>	Static	Small intestinal digestion (α -amylase)	No	Tornero-Martínez et al. (2019)
Polysaccharides from <i>Aspergillus cristatus</i>	Static	Oral, gastric and small intestinal digestion (pancreatin and porcine bile extract)	No	Rui et al. (2019)
Polysaccharides from <i>Gracilaria lemaneiformis</i>	Static	Oral, gastric and small intestinal digestion (pancreatin and porcine bile extract)	No	Han et al. (2020)

Static *in vitro* digestion methodologies

Static digestion methods represent the simplest techniques and can include two or three separate digestion steps (oral, gastric and intestinal). The oral phase is sometimes not taken into account given that the process in the mouth lasts from a few seconds to minutes, and since the salivary pH value is close to neutral, the degradation of significant compounds from food samples is not expected in this stage. To date, most *in vitro* techniques for the analysis of the glycaemic properties of foods have their roots in the earlier methods for the analysis of dietary fiber, total carbohydrates and resistant starch in foods (Woolnough et al. 2008). In this sense, current methods are based on those proposed by Southgate (1969), Englyst, Wiggins, and Cummings (1982), and/or Jenkins et al. (1982), where readily digestible starch was hydrolyzed with α -amylase and pullulanase, and resistant starch was subjected to more exhaustive hydrolysis with amyloglucosidase. Since then to the present day an extensive number of different methods for the digestibility of carbohydrates have been reported. In this regard, the most widely accepted standardized *in vitro* method for digestion was developed in 2014 as an attempt to unify the different

methodologies used until that moment and to produce data that are more comparable and reproducible between studies (Minekus et al. 2014; Egger et al. 2016; Brodkorb et al. 2019). The static protocol published by the InfoGest network represents an international consensus of scientists from 32 countries working in the field of digestion, and it has been widely used and supported in several works since its release (Egger et al. 2016, 2017; Bohn et al. 2018; Verhoeckx et al. 2019).

Concerning small intestinal digestion, this method presents two approaches regarding the enzymes used at this stage (Minekus et al. 2014). Firstly, the use of a pancreatic extract (pancreatin) containing all the relevant enzymes is suggested for reasons of simplicity. As an alternative, individual enzymes, such as trypsin, chymotrypsin, pancreatin lipase, colipase, pancreatin amylase can also be used. However, despite the clear usefulness of this method for the evaluation of the digestibility of proteins, lipids and starchy carbohydrates, the complexity of the gastrointestinal process in the small intestine means that the enzymes proposed in this method for intestinal digestion cannot reflect the broad array of enzymatic activities of the human gut toward carbohydrates since small intestinal mucosal carbohydrases are

Table 3. Overview on the *in vitro* intestinal digestion studies based on the mammalian mucosal small intestinal enzymes (rat and pig) for carbohydrates.

Carbohydrates evaluated	Digestion model	Digestion and fermentation stages/conditions	Degradation	References
Isomalto/Malto-polysaccharides	Static	Small Intestine: Brush Border carbohydrases from rat intestinal acetone powder	Yes	Leemhuis et al. (2014)
GOS from lactose GOS from lactulose Lactulose Fructooligosaccharides Lactosucrose Lactose and Lactulose	Static	Small Intestine: Brush Border carbohydrases from rat intestinal acetone powder	Yes – structural differences determined the resistance to degradation	Ferreira-Lazarte, Olano, et al. (2017)
Fructans and α -galactooligosaccharides	Static	Small Intestine: Brush Border carbohydrases from rat intestinal acetone powder	Yes – structural differences determined the resistance to degradation	Gallego-Lobillo, Ferreira-Lazarte, Hernández-Hernández, and Villamiel (2020)
Isomaltoligosaccharides	Static	Enzyme mixture (pancreatin, invertase and amyloglucosidase) and brush border carbohydrases from rat intestinal acetone powder	Yes	Gallego-Lobillo, Ferreira-Lazarte, Hernández-Hernández, Montilla et al. (2020) Hu et al. (2017)
Maltose/Sucrose isomers	Static	Small Intestine: Brush Border α -glucosidases from rat intestinal acetone powder	Yes	Lee et al. (2016)
Starch samples	Static	Oral, gastric and small intestinal (pancreatin, bile salts and brush border carbohydrases from rat intestinal acetone powder)	Yes	Garcia-Campayo et al. (2018)
GOS from lactose GOS from lactulose	Static (InfoGest modified)	Oral, gastric and small intestinal (pancreatin, bile salts and brush border carbohydrases from rat intestinal acetone powder)	Yes – milk matrix did not modify their digestibility	Ferreira-Lazarte, Montilla, et al. (2017)
Turanose and resistant starch	Static (InfoGest modified)	Oral, gastric and small intestinal (pancreatin, bile salts and brush border carbohydrases from rat intestinal acetone powder)	Yes	Park, Chandrasekaran, and Yoo (2019)
Starchy and fiber commercial food products	Dynamic (TIMcarbo)	Oral, gastric and small intestinal (pancreatin, bile salts) and brush border enzyme digestion (rat brush border enzyme extract and commercial lactase)	Yes	Bellmann et al. (2018)
GOS from lactose GOS from lactose GOS from lactose	Static Static	Small Intestine: brush border membrane vesicles from pig Small Intestine: brush border membrane vesicles from pig	Yes – $\beta(1\rightarrow3)$ linkages more hydrolyzed Slight GOS degradation	Ferreira-Lazarte, Gallego-Lobillo, et al. (2019) Tanabe, Nakamura, and Oku (2014) and Tanabe et al. (2015)
Sialyllactoses	Static (InfoGest modified)	Oral, Gastric, and Small Intestine (pancreatin, bile salts and brush border membrane vesicles from pig)	Yes	Moon et al. (2016)

not regularly taken into account (Table 1). There is therefore the lack of a prescribed final step during intestinal digestion in which disaccharidases from intestinal mucosa would complete carbohydrate digestion (Goodman 2010).

Studies on the digestion of different types of fiber such as resistant starch, FOS, pectic polysaccharides from kiwifruit, and sugar beet pectin using simulated gastric and intestinal fluids similar to the InfoGest method, have shown a partial degradation or changes in their structural features (Carnachan et al. 2012; Logan, Wright, and Goff 2015; Foucault et al. 2016; Sancho et al. 2017). Molecular weight diminution and decreases in their initial concentration were observed in these substrates, although the enzymes (pancreatin, α -amyloglucosidase or invertase) and concentration used

were different in each study, not showing any consensus between methods.

The Association of Official Analytical Chemists (AOAC) developed an integrated determination method for dietary fiber, including non-digestible oligosaccharides (NDOs) and resistant starch (AOAC 2009.01) (McCleary et al. 2010), which was later modified in 2015 (McCleary, Sloane, and Draga 2015). This method, as well as the others, is based on the use of isolated digestive enzymes. Porcine pancreatic α -amylase, mucosal α -glucosidase and a fungal amyloglucosidase from *Aspergillus niger* are used to produce the complete hydrolysis of digestible saccharides and, therefore, to distinguish between digestible and non-digestible carbohydrates. However, these enzymes cannot completely hydrolyze

digestible saccharides such as sucrose, lactose, panose, etc. since they do not represent the fully complex enzymatic environment of the small intestine. As a result, digestible saccharides are not fully degraded and are miscategorized as non-digestible carbohydrates leading to an inaccurate determination of these resistant carbohydrates (Tanabe, Nakamura, and Oku 2014).

New methods based on mammalian small intestine extracts for carbohydrate digestion

The scarce information about the use of mammalian intestinal enzymes has constituted a necessity for some authors in developing methods that may provide a better understanding of the processes occurring in the small intestine, so that it can be used in the evaluation of this functionality in foods.

In addition to the inclusion of intestinal brush border enzymes in the TIMcarbo, alternative methods based on the use of mammalian intestinal extracts, which could provide a more accurate approach of the intestinal process, have also been presented as a good option to overcome the disadvantages of the use of isolated enzymes (Table 3). In this sense, similarity between humans and rats with regard to the hydrolyzing activity of small intestinal disaccharidases was proved (Oku et al. 2011). Small intestinal mucosa obtained from healthy human donors and rats were used to test the digestibility of different oligosaccharides showing a similar enzymatic activity, providing a good alternative to the evaluation of functional food digestion in the small intestine.

Methods based on rat small intestinal extract

To date, there are limited reports regarding the use of a rat small intestinal extract for the evaluation of the digestibility of non-digestible carbohydrates. The most common use of these extracts refers to previous steps of digestion focusing on their fermentability (Ito et al. 2008; Kaulpiboon et al. 2015). However, other studies have also successfully applied these intestinal enzymes from rats for the assessment of the digestibility of resistant carbohydrates, such as fructooligosaccharides (FOS) (Oku, Tokunaga, and Hosoya 1984), GOS (Ohtsuka et al. 1990), as well as to a range of maltose and sucrose isomers/polymers (Lee et al. 2016; Shin et al. 2019).

Ferreira-Lazarte, Olano, et al. (2017) reported a new *in vitro* method to evaluate the digestibility of dietary oligosaccharides under physiological conditions using extracts derived from the small intestine of rats (RSIE) with the advantage of requiring small amounts of carbohydrate substrates to be assessed. The proposed method also included a reliable analytical method, such as GC-FID, where remaining carbohydrates as well as the products of digestion can be evaluated identifying the specific structural changes during digestion (type of glycosidic linkage hydrolyzed, monosaccharides released, etc.). This method represents a useful, simple and cost-effective tool to distinguish between digestible and non-digestible carbohydrates. The similarity found between rat intestinal disaccharidase activities (maltase,

sucrase, palatinase, trehalase and lactase) and their equivalent in humans (Oku et al. 2011), as well as the good correlation found between this method and data collected *in vivo* (Molis et al. 1996; Hernández-Hernández et al. 2012; Jantscher-Krenn, Marx, and Bode 2013), endorse the suitability of these mammalian extracts to assess the digestibility of dietary carbohydrates. The digestion rate of recognized and potential prebiotics (GOS, FOS, lactulose, novel lactulose-derived galactooligosaccharides (OsLu) and lactosucrose) was assessed and compared with this method. Prebiotic and potential prebiotic compounds were highly resistant to small intestine enzymes; however, structural differences exhibited an important effect on their susceptibility to degradation. Remarkably, different predominant linkages in GOS demonstrated a key role in resistance to degradation. For instance, $\beta(1\rightarrow6)$ linkages in GOS mixtures showed significantly higher resistance when compared to $\beta(1\rightarrow4)$ linkages, which were the most susceptible to degradation in all prebiotic samples tested. Monomeric composition was also a critical factor for digestibility, thus, oligosaccharides with a terminal fructose (those derived from lactulose) were less prone to hydrolysis than those with glucose (GOS). Lactulose and FOS resistance was higher as compared to GOS and OsLu, due to their monomer composition and resistant linkages. The molecular weight of OsLu, which has been related to slower fermentation by the colonic microbiota than lactulose, being able to reach the distal colon without major alterations (Cardelle-Cobas et al. 2011, 2012), seemed to also provide higher resistance to intestinal enzymes. In this sense, the trisaccharide fraction (mainly β -Gal-(1 \rightarrow 6)- β -Gal-(1 \rightarrow 4)-Fru) present in OsLu was almost not degraded as compared to lactulose after *in vitro* digestion, whereas the trisaccharide fraction in GOS, mainly β -Gal-(1 \rightarrow 4)- β -Gal-(1 \rightarrow 4)-Glc and β -Gal-(1 \rightarrow 6)- β -Gal-(1 \rightarrow 4)-Glc, was slightly more prone to degradation. The relationship between the structure of GOS and resistance to mammalian intestinal enzymes and their bioactive effect was already pointed out in previous reports (Rastall et al. 2005; Hernández-Hernández et al. 2012), however, no comparison of the digestibility of different types of GOS from lactose and lactulose using a mammalian intestinal extract was carried out before. Furthermore, a recent study carried out by Gallego-Lobillo, Ferreira-Lazarte, Hernández-Hernández, and Villamiel (2020), which evaluated the kinetics of degradation of digestible and non-digestible saccharides (lactose and lactulose, respectively) with RSIE under physiological conditions, showed the higher resistance of galactosyl-fructose linkages (with lower hydrolysis rate constants (K_m) and higher $t_{1/2}$), whereas lactose was highly degraded by these enzymes (higher K_m and lower $t_{1/2}$), supporting, therefore, the suitability of the use of RSIE for the simulation of carbohydrate digestion.

In addition, these small intestinal enzymes were also used to assess their impact on different types of functional fiber such as several types of fructans and α -galactooligosaccharides (Gallego-Lobillo, Ferreira-Lazarte, Hernández-Hernández, Montilla et al. 2020). In line with previous studies, inulin-type fructans and fructooligosaccharides were

found to be very resistant to digestion. Strikingly, the degree of hydrolysis of non-fructosylated α -galactooligosaccharides derived from peas was around 60%. Factors as Mw and type of linkage contributed to the resistance of fructans to digestion, whereas α -GOS were hydrolyzed considerably in a similar way to the degradation of β -GOS derived from lactose and lactulose observed in previous works.

Moreover, given that the mixtures of prebiotics are usually included in different foodstuffs, mainly milk and dairy products, there is the need to consider the effect of the food matrix on prebiotic digestibility. Therefore, Ferreira-Lazarte, Montilla, et al. (2017) reported the utilization of mammalian intestinal enzymes in a synergistic manner with the current standardized method (InfoGest) to evaluate the *in vitro* gastrointestinal digestion of prebiotics added to milk. The use of pancreatic fluids and bile salts to simulate intestinal digestion has provided evidence of the limitation of these models for carbohydrate digestion since hardly any modifications were observed in the carbohydrate fraction of the samples. Conversely, the utilization of the RSIE to digest samples after their gastric stage demonstrated its suitability to hydrolyze carbohydrates showing measurable decreases in this fraction. Firstly, higher amounts of lactose did not seem to affect prebiotic degradation after 2 hours of digestion with the RSIE. Degradation found in the oligosaccharide fraction of GOS (35%) and OsLu (15%) within the milk was similar to that obtained by Ferreira-Lazarte, Olano, et al. (2017), that is 34 and 18%, respectively. High decreases of lactose content were observed whereas lactulose remained as the least prone to disaccharide degradation. In line with the results of Ferreira-Lazarte, Olano, et al. (2017), oligosaccharides from the predominant $\beta(1\rightarrow4)$ -GOS mixture exhibited the highest degradation after digestion, whereas oligosaccharides from $\beta(1\rightarrow6)$ -OsLu stood as the most resistant structure with 85% of composition intact after 2 hours of digestion, highlighting the suitability of the inclusion of these substrates within a real food context.

Therefore, in a robust digestion method for carbohydrates the inclusion of a step using mammalian small intestinal extract is needed to fully cover the carbohydrase activities of the gastrointestinal system. In this sense, rat intestinal enzymes could provide a useful and reliable tool to determine the digestibility of dietary carbohydrates in static isolated models (Ferreira-Lazarte, Olano, et al. 2017; Shin et al. 2019) as well as a part of a complete gastrointestinal digestion model (Ferreira-Lazarte, Montilla, et al. 2017; Garcia-Campayo et al. 2018). However, despite the positive results obtained, the small size of these animals could create a challenge for their use as human models due to the very different anatomy and physiology as compared to humans.

Methods based on pig small intestinal enzymes

Pig has emerged as an important model due to their anatomical and physiological similarity to humans and the human genome (Humphray et al. 2007), as well as their larger litter size, and the fact that they are a food source may help avoid ethical concerns (Kuzmuk and Schook

2011). The use of pigs has proved to be a robust model for several studies such as tissue engineering, imaging, surgery, chemotherapy, radiation studies, cancer, atherosclerosis, myocardial infarction, and general cardiovascular models, which cannot be carried out accurately or have failed in small animals (Kuzmuk 2009; Jensen et al. 2010; Schook et al. 2015). The physiological similarity between humans and pigs in terms of digestive processes places the pig as a robust model for human digestive and colonic studies (Heinritz, Mosenthin, and Weiss 2013). Thus, the use of small intestinal materials from pigs could represent a more reliable method as well as a step forward toward the development of a robust method that is comparable to humans in order to gather information about dietary carbohydrate digestion. Nonetheless, the use of these extracts has still been sparsely used even though they have proved their successful utility on carbohydrate digestion and are similar to human intestinal activity (Lander et al. 2001; Humphray et al. 2007; Oku et al. 2011).

Alternative methods based on the use of mammalian intestinal enzymes derived from pigs (Tanabe, Nakamura, and Oku 2014) or weaning piglets (Strube et al. 2015) have been proposed. Given the limitations of the AOAC 2009.01 method, an improvement method by the addition of mammalian small intestinal enzymes from pigs was suggested (Tanabe, Nakamura, and Oku 2014). In this sense, the incomplete degradation of digestible saccharides that leads to overestimating NDOs was improved by the addition of the enzymes present at the small intestinal brush border from pigs, providing, therefore, a more realistic environment and an accurate quantification method of NDOs in processed food (Tanabe et al. 2015).

The application of intestinal disaccharidases, that are identified as glycoproteins linked to the apical membrane and are actively budded off as brush border membrane vesicles (BBMVs) (Figure 1) (McConnell et al. 2009; Hooton et al. 2015), from pig has recently been shown to be a reliable and useful approach to evaluate prebiotic carbohydrate digestibility, under physiological conditions of pH, temperature and time. Isolated BBMVs from the intestinal mucosa were used by Ferreira-Lazarte, Gallego-Lobillo, et al. (2019) to evaluate the digestibility of prebiotic oligosaccharides (GOS and OsLu). Regarding enzymatic activity, BBMVs showed higher values when compared to a commercial intestinal extract from rats mainly due to the exhaustive fractionation and purification carried out in the obtainment of the membrane vesicles. Digestibility assays, in line with previous reports (Ferreira-Lazarte, Montilla, et al. 2017; Ferreira-Lazarte, Olano, et al. 2017), confirmed the resistance of $\beta(1\rightarrow6)$ linkages as compared to $\beta(1\rightarrow4)$ linkages. Moreover, predominantly $\beta(1\rightarrow3)$ linkages in GOS that were also analyzed in this study, revealed the high susceptibility of these structures to intestinal disaccharidases with a degradation of 44%, whereas $\beta(1\rightarrow4)$ and $\beta(1\rightarrow6)$ showed a hydrolysis of 23 and 12%, respectively, after 2 hours of digestion.

Likewise, another recent study on the obtainment of GOS using the trans- β -galactosylation activity of the pig

β -galactosidase embedded in the BBMV has revealed the high preferable synthesis of GOS linked by $\beta(1\rightarrow3)$ as compared to those linked by $\beta(1\rightarrow4)$ or $\beta(1\rightarrow6)$ (Julio-Gonzalez et al. 2019). In good agreement with this, the β -Gal-(1 \rightarrow 3)-Gal- glycosidic bond was also identified, together with the β -Gal-(1 \rightarrow 5)-Fru linkage, as being preferentially digested when lactulose was incubated with pig BBMV (Julio-Gonzalez et al. 2020). Therefore, considering that enzymes can catalyze reversible reactions in either direction (Abdul Manas, Md Illias, and Mahadi 2018), data obtained by these authors support the hypothesis that most glycosidic linkages formed when intestinal β -galactosidase acts as transgalactosidase, are preferentially broken under hydrolytic conditions. In addition, specific monosaccharide composition provided a high resistance on several structures such as galactosyl-galactoses, galactosyl-fructoses and, especially, on the trisaccharide of OsLu ($\beta(1\rightarrow6)$) (only 9.8% of hydrolysis was reported after 2 hours), highlighting the key role of these structural features.

Conclusions

The lack of thorough studies on the intestinal degradation of non-digestible carbohydrates has led to the use of standardized official methods to determine their digestibility despite their limitation (McCleary et al. 2010; Egger et al. 2016; Drechsler and Bornhorst 2018). To solve this problem, important efforts have been made to develop an adequate *in vitro* model for carbohydrates, the intestinal mucosal enzymes being used only in few methods that have been recently developed (Ferreira-Lazarte, Olano, et al. 2017; Garcia-Campayo et al. 2018; Shin et al. 2019).

Evaluation of the digestibility of carbohydrates with new emerging methods based on the use of mammalian intestinal enzymes have revealed essential structure-function relationships in prebiotic carbohydrates, such as the strong resistance of $\beta(1\rightarrow6)$ oligosaccharides, especially of those derived from lactulose, which have been shown to be the most resistant GOS to *in vitro* digestion. Thus, when aiming for the potential development of tailored and new generation prebiotics, oligosaccharides derived from lactulose would represent the ideal candidates to be taken into account, given also the large amount of studies supporting their technological and biological properties (Villamiel et al. 2014; López-Sanz et al. 2015, 2018; Barroso et al. 2016; Fernández et al. 2018).

The *in vitro* digestion models based on the utilization of rat small intestinal extract proved to be a useful, reliable and efficient approach to evaluating the digestibility of dietary carbohydrates (digestible and non-digestible), overcoming the limitations of the current standardized methods for *in vitro* gastrointestinal digestion. Moreover, the utilization of a model based on small intestinal BBMV from pigs would be a more realistic approach to study the influence of glycosidic linkages, degree of polymerization and monomeric composition on resistance to intestinal digestion of selected prebiotics. Thus, although there is still work to be done in regard to developing more standardized conditions

concerning enzymatic activities, enzyme/substrate ratios or a regular supply of pig mucosal intestinal enzymes, the use of these enzyme material, combined with the standardized methodologies, clearly involves the next step toward a final and complete small intestinal digestion simulation for dietary carbohydrates.

Lastly, the prebiotic concept maintains that these substrates should be “non-digested” by intestinal enzymes, reaching the colon at least almost intact; however, findings obtained so far with current studies have revealed the partial degradation of some of these substrates. Therefore, the studies that have been discussed in the present review indicate that certain prebiotic oligosaccharides may not reach the colon fully intact, thus challenging the general belief about their complete non-digestibility, and, although more studies are required to focus, for instance, on the role of the food matrix or on the better understanding of the physiologically relevant activity of BBMV, a possible revision of the current concept of the digestibility of prebiotics could be envisaged.

Funding

This work has been funded by the Ministry of Economy and Competitiveness (MINECO) of Spain with Projects AGL2017-84614-C2-1-R.

References

- Abdul Manas, N. H., R. Md Illias, and N. M. Mahadi. 2018. Strategy in manipulating transglycosylation activity of glycosyl hydrolase for oligosaccharide production. *Critical Reviews in Biotechnology* 38 (2): 272–93. doi: 10.1080/07388551.2017.1339664.
- Alpers D. H. 2003. Carbohydrates/Digestion, absorption, and metabolism. In: *Encyclopedia of Food Sciences and Nutrition*, ed. B. Caballero, 2nd ed., 881–887. Cambridge, MA: Elsevier.
- Barroso, E., C. Cueva, C. Peláez, M. C. Martínez-Cuesta, and T. Requena. 2015. Development of human colonic microbiota in the computer-controlled dynamic SIMulator of the GastroIntestinal tract SIMGI. *LWT - Food Science and Technology* 61 (2):283–9. doi: 10.1016/j.lwt.2014.12.014.
- Barroso, E., A. Montilla, N. Corzo, C. Peláez, M. C. Martínez-Cuesta, and T. Requena. 2016. Effect of lactulose-derived oligosaccharides on intestinal microbiota during the shift between media with different energy contents. *Food Research International* 89 (Pt 1):302–8. doi: 10.1016/j.foodres.2016.08.025.
- Bellmann, S., M. Minekus, P. Sanders, S. Bosgra, and R. Havenaar. 2018. Human glycemic response curves after intake of carbohydrate foods are accurately predicted by combining *in vitro* gastrointestinal digestion with *in silico* kinetic modeling. *Clinical Nutrition Experimental* 17:8–22. doi: 10.1016/j.yclnex.2017.10.003.
- Bohn, T., F. Carriere, L. Day, A. Deglaire, L. Egger, D. Freitas, M. Golding, S. Le Feunteun, A. Macierzanka, O. Menard, et al. 2018. Correlation between *in vitro* and *in vivo* data on food digestion. What can we predict with static *in vitro* digestion models? *Critical Reviews in Food Science and Nutrition* 58 (13):2239–61. doi: 10.1080/10408398.2017.1315362.
- Brodkorb, A., L. Egger, M. Alminger, P. Alvito, R. Assunção, S. Ballance, T. Bohn, C. Bourlieu-Lacanal, R. Boutrou, F. Carrière, et al. 2019. INFOGEST static *in vitro* simulation of gastrointestinal food digestion. *Nature Protocols* 14 (4):991–1014. doi: 10.1038/s41596-018-0119-1.
- Cardelle-Cobas, A., N. Corzo, A. Olano, C. Peláez, T. Requena, and M. Ávila. 2011. Galactooligosaccharides derived from lactose and lactulose: Influence of structure on Lactobacillus, Streptococcus and

- Bifidobacterium growth. *International Journal of Food Microbiology* 149 (1):81–7. doi: [10.1016/j.ijfoodmicro.2011.05.026](https://doi.org/10.1016/j.ijfoodmicro.2011.05.026).
- Cardelle-Cobas, A., A. Olano, N. Corzo, M. Villamiel, M. Collins, S. Kolida, and R. A. Rastall. 2012. *In vitro* fermentation of lactulose-derived oligosaccharides by mixed fecal microbiota. *Journal of Agricultural and Food Chemistry* 60 (8):2024–32. doi: [10.1021/jf203622d](https://doi.org/10.1021/jf203622d).
- Carnachan, S. M., T. J. Bootten, S. Mishra, J. A. Monro, and I. M. Sims. 2012. Effects of simulated digestion *in vitro* on cell wall polysaccharides from kiwifruit (*Actinidia* spp.). *Food Chemistry* 133 (1): 132–9. doi: [10.1016/j.foodchem.2011.12.084](https://doi.org/10.1016/j.foodchem.2011.12.084).
- Chen, G., M. Xie, P. Wan, D. Chen, H. Ye, L. Chen, X. Zeng, and Z. Liu. 2018. Digestion under saliva, simulated gastric and small intestinal conditions and fermentation *in vitro* by human intestinal microbiota of polysaccharides from Fuzhuan brick tea. *Food Chemistry* 244:331–9. doi: [10.1016/j.foodchem.2017.10.074](https://doi.org/10.1016/j.foodchem.2017.10.074).
- Chen, L., W. Xu, D. Chen, G. Chen, J. Liu, X. Zeng, R. Shao, and H. Zhu. 2018. Digestibility of sulfated polysaccharide from the brown seaweed *Ascophyllum nodosum* and its effect on the human gut microbiota *in vitro*. *International Journal of Biological Macromolecules* 112:1055–61. doi: [10.1016/j.ijbiomac.2018.01.183](https://doi.org/10.1016/j.ijbiomac.2018.01.183).
- Di, T., G. Chen, Y. Sun, S. Ou, X. Zeng, and H. Ye. 2018. *In vitro* digestion by saliva, simulated gastric and small intestinal juices and fermentation by human fecal microbiota of sulfated polysaccharides from *Gracilaria rubra*. *Journal of Functional Foods* 40:18–27. doi: [10.1016/j.jff.2017.10.040](https://doi.org/10.1016/j.jff.2017.10.040).
- Díez-Municio, M., M. Herrero, A. Olano, and F. J. Moreno. 2014. Synthesis of novel bioactive lactose-derived oligosaccharides by microbial glycoside hydrolases. *Microbial Biotechnology* 7 (4): 315–31. doi: [10.1111/1751-7915.12124](https://doi.org/10.1111/1751-7915.12124).
- Donaldson, G. P., S. M. Lee, and S. K. Mazmanian. 2016. Gut biogeography of the bacterial microbiota. *Nature Reviews Microbiology* 14 (1):20–32. doi: [10.1038/nrmicro3552](https://doi.org/10.1038/nrmicro3552).
- Drechsler, K. C., and G. M. Bornhorst. 2018. Modeling the softening of carbohydrate-based foods during simulated gastric digestion. *Journal of Food Engineering* 222:38–48. doi: [10.1016/j.jfoodeng.2017.11.007](https://doi.org/10.1016/j.jfoodeng.2017.11.007).
- Dupont, D., M. Alric, S. Blanquet-Diot, G. Bornhorst, C. Cueva, A. Deglaire, S. Denis, M. Ferrua, R. Havenaar, J. Lelieveld, et al. 2019. Can dynamic *in vitro* digestion systems mimic the physiological reality? *Critical Reviews in Food Science and Nutrition* 59 (10): 1546–62. doi: [10.1080/10408398.2017.1421900](https://doi.org/10.1080/10408398.2017.1421900).
- Egger, L., O. Ménard, C. Delgado-Andrade, P. Alvito, R. Assunção, S. Balance, R. Barberá, A. Brodkorb, T. Cattenoz, A. Clemente, et al. 2016. The harmonized INFOGEST *in vitro* digestion method: From knowledge to action. *Food Research International* 88:217–25. doi: [10.1016/j.foodres.2015.12.006](https://doi.org/10.1016/j.foodres.2015.12.006).
- Egger, L., P. Schlegel, C. Baumann, H. Stoffers, D. Guggisberg, C. Brügger, D. Dürr, P. Stoll, G. Vergères, and R. Portmann. 2017. Physiological comparability of the harmonized INFOGEST *in vitro* digestion method to *in vivo* pig digestion. *Food Research International* 102:567–74. doi: [10.1016/j.foodres.2017.09.047](https://doi.org/10.1016/j.foodres.2017.09.047).
- Englyst, H., H. S. Wiggins, and J. H. Cummings. 1982. Determination of the non-starch polysaccharides in plant foods by gas-liquid chromatography of constituent sugars as alditol acetates. *The Analyst* 107 (1272):307–18. doi: [10.1039/an9820700307](https://doi.org/10.1039/an9820700307).
- Fernández, J., F. J. Moreno, A. Olano, A. Clemente, C. J. Villar, and F. Lombó. 2018. A galacto-oligosaccharides preparation derived from lactulose protects against colorectal cancer development in an animal model. *Frontiers in Microbiology* 9:1–14. doi: [10.3389/fmicb.2018.02004](https://doi.org/10.3389/fmicb.2018.02004).
- Ferreira-Lazarte, A., P. Gallego-Lobillo, F. J. Moreno, M. Villamiel, and O. Hernández-Hernández. 2019. *In vitro* digestibility of galactooligosaccharides: Effect of the structural features on their intestinal degradation. *Journal of Agricultural and Food Chemistry* 67 (16): 4662–70. doi: [10.1021/acs.jafc.9b00417](https://doi.org/10.1021/acs.jafc.9b00417).
- Ferreira-Lazarte, A., A. Montilla, A.-I. Mulet-Cabero, N. Rigby, A. Olano, A. Mackie, and M. Villamiel. 2017. Study on the digestion of milk with prebiotic carbohydrates in a simulated gastrointestinal model. *Journal of Functional Foods* 33:149–54. doi: [10.1016/j.jff.2017.03.031](https://doi.org/10.1016/j.jff.2017.03.031).
- Ferreira-Lazarte, A., F. J. Moreno, C. Cueva, I. Gil-Sánchez, and M. Villamiel. 2019. Behaviour of citrus pectin during its gastrointestinal digestion and fermentation in a dynamic simulator (simgi®). *Carbohydrate Polymers* 207:382–90. doi: [10.1016/j.carbpol.2018.11.088](https://doi.org/10.1016/j.carbpol.2018.11.088).
- Ferreira-Lazarte, A., A. Olano, M. Villamiel, and F. J. Moreno. 2017. Assessment of *in vitro* digestibility of dietary carbohydrates using rat small intestinal extract. *Journal of Agricultural and Food Chemistry* 65 (36):8046–53. doi: [10.1021/acs.jafc.7b01809](https://doi.org/10.1021/acs.jafc.7b01809).
- Foucault, M., J. Singh, R. B. Stewart, and H. Singh. 2016. Pilot scale production and *in vitro* gastro-small intestinal digestion of self-assembled recrystallised starch (SARS) structures. *Journal of Food Engineering* 191:95–7. doi: [10.1016/j.jfoodeng.2016.07.001](https://doi.org/10.1016/j.jfoodeng.2016.07.001).
- Gallego-Lobillo, P., A. Ferreira-Lazarte, O. Hernández-Hernández, A. Montilla, and M. Villamiel. 2020. Evaluation of the impact of a rat small intestinal extract on the digestion of four different functional fibers. *Food & function* 11 (5):4081–9. doi: [10.1039/d0fo00236d](https://doi.org/10.1039/d0fo00236d).
- Gallego-Lobillo, P., A. Ferreira-Lazarte, O. Hernández-Hernández, and M. Villamiel. 2020. Kinetic study on the digestibility of lactose and lactulose using small intestinal glycosidases. *Food Chemistry* 316: 126326. doi: [10.1016/j.foodchem.2020.126326](https://doi.org/10.1016/j.foodchem.2020.126326).
- García-Campayo, V., S. Han, R. Vercauteren, and A. Franck. 2018. Digestion of food ingredients and food using an *in vitro* model integrating intestinal mucosal enzymes. *Food and Nutrition Sciences* 9 (6):711–34. doi: [10.4236/fns.2018.96055](https://doi.org/10.4236/fns.2018.96055).
- Gibson, G. R., R. Hutkins, M. E. Sanders, S. L. Prescott, R. A. Reimer, S. J. Salminen, K. Scott, C. Stanton, K. S. Swanson, P. D. Cani, et al. 2017. Expert consensus document: The International Scientific Association for Probiotics and Prebiotics (ISAPP) consensus statement on the definition and scope of prebiotics. *Nature Reviews Gastroenterology & Hepatology* 14 (8):491–502. doi: [10.1038/nrgastro.2017.75](https://doi.org/10.1038/nrgastro.2017.75).
- Goodman, B. E. 2010. Insights into digestion and absorption of major nutrients in humans. *Advances in Physiology Education* 34 (2):44–53. doi: [10.1152/advan.00094.2009](https://doi.org/10.1152/advan.00094.2009).
- Guinane, C. M., and P. D. Cotter. 2013. Role of the gut microbiota in health and chronic gastrointestinal disease: Understanding a hidden metabolic organ. *Therapeutic Advances in Gastroenterology* 6 (4): 295–308. doi: [10.1177/1756283X13482996](https://doi.org/10.1177/1756283X13482996).
- Gullón, B., B. Gómez, M. Martínez-Sabajanes, R. Yáñez, J. C. Parajó, and J. L. Alonso. 2013. Pectic oligosaccharides: Manufacture and functional properties. *Trends in Food Science & Technology* 30 (2): 153–61. doi: [10.1016/j.tifs.2013.01.006](https://doi.org/10.1016/j.tifs.2013.01.006).
- Han, R., D. Pang, L. Wen, L. You, R. Huang, and V. Kulikouskaya. 2020. *In vitro* digestibility and prebiotic activities of a sulfated polysaccharide from *Gracilaria Lemaneiformis*. *Journal of Functional Foods* 64:103652. doi: [10.1016/j.jff.2019.103652](https://doi.org/10.1016/j.jff.2019.103652).
- Heinritz, S. N., R. Mosenthin, and E. Weiss. 2013. Use of pigs as a potential model for research into dietary modulation of the human gut microbiota. *Nutrition Research Reviews* 26 (2):191–209. doi: [10.1017/S0954422413000152](https://doi.org/10.1017/S0954422413000152).
- Hernández-Hernández, O. 2019. *In vitro* gastrointestinal models for prebiotic carbohydrates: A critical review. *Current Pharmaceutical Design* 25 (32):3478–6. doi: [10.2174/1381612825666191011094724](https://doi.org/10.2174/1381612825666191011094724).
- Hernández-Hernández, O., M. C. Marin-Manzano, L. A. Rubio, F. J. Moreno, M. L. Sanz, and A. Clemente. 2012. Monomer and linkage type of galacto-oligosaccharides affect their resistance to ileal digestion and prebiotic properties in rats. *The Journal of Nutrition* 142 (7):1232–9. doi: [10.3945/jn.111.155762](https://doi.org/10.3945/jn.111.155762).
- Hernández-Hernández, O., A. Olano, R. A. Rastall, and F. J. Moreno. 2019. *In vitro* digestibility of dietary carbohydrates: Toward a standardized methodology beyond amyolytic and microbial enzymes. *Frontiers in Nutrition* 6:61. doi: [10.3389/fnut.2019.00061](https://doi.org/10.3389/fnut.2019.00061).
- Holloway, W. D., C. Tasman-Jones, and K. Maher. 1983. Pectin digestion in humans. *The American journal of clinical nutrition* 37 (2): 253–5. doi: [10.1093/ajcn/37.2.253](https://doi.org/10.1093/ajcn/37.2.253).
- Holmes, R., and R. W. Lobley. 1989. Intestinal brush border revisited. *Gut* 30 (12):1667–78. doi: [10.1136/gut.30.12.1667](https://doi.org/10.1136/gut.30.12.1667).
- Hooton, D., R. Lentle, J. Monro, M. Wickham, and R. Simpson. 2015. The secretion and action of brush border enzymes in the

- mammalian small intestine. *Reviews of Physiology, Biochemistry and Pharmacology* 168:59–118. doi: [10.1007/112_2015_24](https://doi.org/10.1007/112_2015_24).
- Hu, Y., V. Winter, X. Y. Chen, and M. G. Gänzle. 2017. Effect of acceptor carbohydrates on oligosaccharide and polysaccharide synthesis by dextranase DsrM from *Weissella cibaria*. *Food Research International* 99 (Pt 1):603–11. doi: [10.1016/j.foodres.2017.06.026](https://doi.org/10.1016/j.foodres.2017.06.026).
- Humphray, S. J., C. E. Scott, R. Clark, B. Marron, C. Bender, N. Camm, J. Davis, A. Jenks, A. Noon, M. Patel, et al. 2007. A high utility integrated map of the pig genome. *Genome Biology* 8 (7): R139. doi: [10.1186/gb-2007-8-7-r139](https://doi.org/10.1186/gb-2007-8-7-r139).
- Ito, S., H. Taguchi, S. Hamada, S. Kawauchi, H. Ito, T. Senoura, J. Watanabe, M. Nishimukai, S. Ito, and H. Matsui. 2008. Enzymatic properties of cellobiose 2-epimerase from *Ruminococcus albus* and the synthesis of rare oligosaccharides by the enzyme. *Applied Microbiology and Biotechnology* 79 (3):433–41. doi: [10.1007/s00253-008-1449-7](https://doi.org/10.1007/s00253-008-1449-7).
- Jandhyala, S. M., R. Talukdar, C. Subramanyam, H. Vuyyuru, M. Sasikala, and D. Nageshwar Reddy. 2015. Role of the normal gut microbiota. *World Journal of Gastroenterology* 21 (29):8787–803. doi: [10.3748/wjg.v21.i29.8787](https://doi.org/10.3748/wjg.v21.i29.8787).
- Jantscher-Krenn, E., C. Marx, and L. Bode. 2013. Human milk oligosaccharides are differentially metabolised in neonatal rats. *The British Journal of Nutrition* 110 (4):640–50. doi: [10.1017/S0007114512005727](https://doi.org/10.1017/S0007114512005727).
- Jenkins, D. J., M. J. Thorne, K. Camelon, A. Jenkins, A. V. Rao, R. H. Taylor, L. U. Thompson, J. Kalmusky, R. Reichert, and T. Francis. 1982. Effect of processing on digestibility and the blood glucose response: A study of lentils. *The American Journal of Clinical Nutrition* 36 (6):1093–101. doi: [10.1093/ajcn/36.6.1093](https://doi.org/10.1093/ajcn/36.6.1093).
- Jensen, T. W., M. J. Mazur, J. E. Pettigew, V. G. Perez-Mendoza, J. Zachary, and L. B. Schook. 2010. A cloned pig model for examining atherosclerosis induced by high fat, high cholesterol diets. *Animal Biotechnology* 21 (3):179–87. doi: [10.1080/10495398.2010.490693](https://doi.org/10.1080/10495398.2010.490693).
- Julio-Gonzalez, L. C., O. Hernández-Hernández, F. J. Moreno, A. Olano, M. L. Jimeno, and N. Corzo. 2019. Trans- β -galactosidase activity of pig enzymes embedded in the small intestinal brush border membrane vesicles. *Scientific Reports* 9 (1):960. doi: [10.1038/s41598-018-37582-8](https://doi.org/10.1038/s41598-018-37582-8).
- Julio-Gonzalez, L. C., O. Hernández-Hernández, F. J. Moreno, M. L. Jimeno, E. García, A. Olano, and N. Corzo. 2020. Hydrolysis and transgalactosylation catalysed by β -galactosidase from brush border membrane vesicles isolated from pig small intestine: A study using lactulose and its mixtures with lactose or galactose as substrates. *Food Research International (Ottawa, Ont.)* 129:108811. doi: [10.1016/j.foodres.2019.108811](https://doi.org/10.1016/j.foodres.2019.108811).
- Kaulpiboon, J., P. Rudeekulthamrong, S. Watanasatitarpa, K. Ito, and P. Pongsawasdi. 2015. Synthesis of long-chain isomaltooligosaccharides from tapioca starch and an *in vitro* investigation of their prebiotic properties. *Journal of Molecular Catalysis B: Enzymatic* 120: 127–35. doi: [10.1016/j.molcatb.2015.07.004](https://doi.org/10.1016/j.molcatb.2015.07.004).
- Khodaei, N., B. Fernandez, I. Fliss, and S. Karboune. 2016. Digestibility and prebiotic properties of potato rhamnogalacturonan I polysaccharide and its galactose-rich oligosaccharides/oligomers. *Carbohydrate polymers* 136:1074–84. doi: [10.1016/j.carbpol.2015.09.106](https://doi.org/10.1016/j.carbpol.2015.09.106).
- Kopf-Bolanz, K. A., F. Schwander, M. Gijs, G. Verge, R. Portmann, and L. Egger. 2012. Validation of an *in vitro* digestive system for studying macronutrient decomposition in humans. *The Journal of Nutrition* 142 (2):245–50. doi: [10.3945/jn.111.148635](https://doi.org/10.3945/jn.111.148635).
- Kuzmuk, K. 2009. Animal models for elucidating human disease: Confronting cancer and other chronic diseases. *CAB Reviews: Perspectives in Agriculture, Veterinary Science, Nutrition and Natural Resources* 4:1–9.
- Kuzmuk, K. N., and L. B. Schook. 2011. Pigs as a model for biomedical sciences. In *The genetics of the pig*, ed. M. F. Rothschild and A. Ruvinsky, 2nd ed., 426–44. Wallingford, UK: CABI Publishing.
- Lander, E. S., L. M. Linton, B. Birren, C. Nusbaum, M. C. Zody, J. Baldwin, K. Devon, K. Dewar, M. Doyle, and W. FitzHugh. 2001. Initial sequencing and analysis of the human genome. *Nature* 409: 860–921.
- Laparra, J. M., M. Diez-Municio, M. Herrero, and F. J. Moreno. 2014. Structural differences of prebiotic oligosaccharides influence their capability to enhance iron absorption in deficient rats. *Food & Function* 5 (10):2430–7. doi: [10.1039/c4fo00504j](https://doi.org/10.1039/c4fo00504j).
- Larsen, N., C. Bussolo de Souza, L. Krych, T. Barbosa Cahú, M. Wiese, W. Kot, K. M. Hansen, A. Blennow, K. Venema, and L. Jespersen. 2019. Potential of pectins to beneficially modulate the gut microbiota depends on their structural properties. *Frontiers in Microbiology* 10:223–13. doi: [10.3389/fmicb.2019.00223](https://doi.org/10.3389/fmicb.2019.00223).
- Lee, B.-H., D. R. Rose, A. H.-M. Lin, R. Quezada-Calvillo, B. L. Nichols, and B. R. Hamaker. 2016. Contribution of the individual small intestinal α -glucosidases to digestion of unusual α -linked glycosidic disaccharides. *Journal of Agricultural and Food Chemistry* 64 (33):6487–94. doi: [10.1021/acs.jafc.6b01816](https://doi.org/10.1021/acs.jafc.6b01816).
- Leemhuis, H., J. M. Dobruchowska, M. Ebbelaar, F. Faber, P. L. Buwalda, M. J. E. C. Van Der Maarel, J. P. Kamerling, and L. Dijkhuizen. 2014. Isomalto/malto-polysaccharide, a novel soluble dietary fiber made via enzymatic conversion of starch. *Journal of Agricultural and Food Chemistry* 62 (49):12034–44. doi: [10.1021/jf503970a](https://doi.org/10.1021/jf503970a).
- Lerner, A., S. Neidhöfer, and T. Matthias. 2017. The gut microbiome feelings of the brain: A perspective for non-microbiologists. *Microorganisms* 5 (4):66–24. doi: [10.3390/microorganisms5040066](https://doi.org/10.3390/microorganisms5040066).
- Li, W., K. Wang, Y. Sun, H. Ye, B. Hu, and X. Zeng. 2015. Influences of structures of galactooligosaccharides and fructooligosaccharides on the fermentation *in vitro* by human intestinal microbiota. *Journal of Functional Foods* 13:158–68. doi: [10.1016/j.jff.2014.12.044](https://doi.org/10.1016/j.jff.2014.12.044).
- Lifschitz, C. H., M. A. Grusak, and N. F. Butte. 2002. Human nutrition and metabolism carbohydrate digestion in humans from a β -Glucan-enriched barley. *The Journal of Nutrition* 132 (9):2593–6. doi: [10.1093/jn/132.9.2593](https://doi.org/10.1093/jn/132.9.2593).
- Logan, K., A. J. Wright, and H. D. Goff. 2015. Correlating the structure and *in vitro* digestion viscosities of different pectin fibers to *in vivo* human satiety. *Food & Function* 6 (1):63–71. doi: [10.1039/c4fo00543k](https://doi.org/10.1039/c4fo00543k).
- Lombard, V., H. Golaconda Ramulu, E. Drula, P. M. Coutinho, and B. Henrissat. 2014. The carbohydrate-active enzymes database (CAZy) in 2013. *Nucleic Acids Research* 42:D490–5. doi: [10.1093/nar/gkt1178](https://doi.org/10.1093/nar/gkt1178).
- López-Sanz, S., A. Montilla, F. J. Moreno, and M. Villamiel. 2015. Stability of oligosaccharides derived from lactulose during the processing of milk and apple juice. *Food Chemistry* 183:64–71. doi: [10.1016/j.foodchem.2015.03.020](https://doi.org/10.1016/j.foodchem.2015.03.020).
- López-Sanz, S., R. Moreno, M. J. De La Mata, F. J. Moreno, and M. Villamiel. 2018. Stability of oligosaccharides derived from lactose and lactulose regarding rheological and thermal properties. *Journal of Food Quality* 2018:1–9. doi: [10.1155/2018/7597301](https://doi.org/10.1155/2018/7597301).
- McCleary, B., J. Rader, M. Champ, and K. Okuma. 2010. Determination of total dietary fiber (CODEX definition) by enzymatic-gravimetric method and liquid chromatography. *Journal of AOAC International* 93 (1):221–33.
- McCleary, B. V., N. Sloane, and A. Draga. 2015. Determination of total dietary fibre and available carbohydrates: A rapid integrated procedure that simulates *in vivo* digestion. *Starch - Stärke* 67 (9–10): 860–83. doi: [10.1002/star.201500017](https://doi.org/10.1002/star.201500017).
- McConnell, R. E., J. N. Higginbotham, D. A. Shifrin, D. L. Tabb, R. J. Coffey, and M. J. Tyska. 2009. The enterocyte microvillus is a vesicle-generating organelle. *The Journal of Cell Biology* 185 (7): 1285–98. doi: [10.1083/jcb.200902147](https://doi.org/10.1083/jcb.200902147).
- Meisenberg, G., and W. H. Simmons. 2016. Digestive enzymes. In: *Principles of Medical Biochemistry*, ed. G. Meisenberg and W. H. Simmons, 4th ed., 342–350. Amsterdam: Elsevier.
- Míguez, B., B. Gómez, P. Gullón, B. Gullón, and J. L. Alonso. 2016. Pectic oligosaccharides and other emerging prebiotics. In *Probiotics and prebiotics in human nutrition and health*, eds. V. Rao and L. Rao, 301–30. London, UK: InTechOpen.
- Minekus, M., M. Alving, P. Alvito, S. Ballance, T. Bohn, C. Bourlieu, F. Carrière, R. Boutrou, M. Corredig, D. Dupont, et al. 2014. A standardised static *in vitro* digestion method suitable for food – An

- international consensus. *Food & Function* 5 (6):1113–24. doi: [10.1039/c3fo60702j](https://doi.org/10.1039/c3fo60702j).
- Minekus, M., P. Marteau, R. Havenaar, and J. H. J. Huis in't Veld. 1995. A multicompartimental dynamic computer-controlled model simulating the stomach and small intestine. *Alternatives to Laboratory Animals* 23:197–209.
- Molis, C., B. Flourié, F. Ouarne, M. F. Gailing, S. Lartigue, A. Guibert, F. Bornet, and J. P. Galmiche. 1996. Digestion, excretion, and energy value of fructooligosaccharides in healthy humans. *The American Journal of Clinical Nutrition* 64 (3):324–8. doi: [10.1093/ajcn/64.3.324](https://doi.org/10.1093/ajcn/64.3.324).
- Molly, K., M. Vande Woestyne, and W. Verstraete. 1993. Development of a 5-step multi-chamber reactor as a simulation of the human intestinal microbial ecosystem. *Applied Microbiology and Biotechnology* 39 (2):254–8. doi: [10.1007/BF00228615](https://doi.org/10.1007/BF00228615).
- Moon, J. S., W. Joo, L. Ling, H. S. Choi, and N. S. Han. 2016. *In vitro* digestion and fermentation of sialyllactoses by infant gut microflora. *Journal of Functional Foods* 21:497–506. doi: [10.1016/j.jff.2015.12.002](https://doi.org/10.1016/j.jff.2015.12.002).
- Nobre, C., S. C. Sousa, S. P. Silva, A. C. Pinheiro, E. Coelho, A. A. Vicente, A. M. P. Gomes, M. A. Coimbra, J. A. Teixeira, and L. R. Rodrigues. 2018. *In vitro* digestibility and fermentability of fructooligosaccharides produced by *Aspergillus ibericus*. *Journal of Functional Foods* 46:278–87. doi: [10.1016/j.jff.2018.05.004](https://doi.org/10.1016/j.jff.2018.05.004).
- Ohtsuka, K., K. Tsuji, Y. Nakagawa, H. Ueda, O. Ozawa, T. Uchida, and T. Ichikawa. 1990. Availability of 4'-galactosylactose (O-beta-D-galactopyranosyl-(1)). *Journal of Nutritional Science and Vitaminology* 36 (3):265–76. doi: [10.3177/jnsv.36.265](https://doi.org/10.3177/jnsv.36.265).
- Oku, T., K. Tanabe, S. Ogawa, N. Sadamori, and S. Nakamura. 2011. Similarity of hydrolyzing activity of human and rat small intestinal disaccharidases. *Clinical and Experimental Gastroenterology* 4: 155–61. doi: [10.2147/CEG.S19961](https://doi.org/10.2147/CEG.S19961).
- Oku, T., T. Tokunaga, and N. Hosoya. 1984. Nondigestibility of a new sweetener, "Neosugar," in the rat. *The Journal of Nutrition* 114 (9): 1574–81. doi: [10.1093/jn/114.9.1574](https://doi.org/10.1093/jn/114.9.1574).
- Ouweland, A. C., and E. E. Vaughan. 2006. *Gastrointestinal microbiology*. New York, NY: Taylor & Francis.
- Park, M. O., M. Chandrasekaran, and S. H. Yoo. 2019. Production and characterization of low-calorie turanose and digestion-resistant starch by an amylase from *Neisseria subflava*. *Food Chemistry* 300:125225. doi: [10.1016/j.foodchem.2019.125225](https://doi.org/10.1016/j.foodchem.2019.125225).
- Picariello, G., P. Ferranti, and F. Addeo. 2016. Use of brush border membrane vesicles to simulate the human intestinal digestion. *Food Research International* 88:327–35. doi: [10.1016/j.foodres.2015.11.002](https://doi.org/10.1016/j.foodres.2015.11.002).
- Rastall, R. A., G. R. Gibson, H. S. Gill, F. Guarner, T. R. Klaenhammer, B. Pot, G. Reid, I. R. Rowland, and M. E. Sanders. 2005. Modulation of the microbial ecology of the human colon by probiotics, prebiotics and synbiotics to enhance human health: An overview of enabling science and potential applications. *FEMS Microbiology Ecology* 52 (2): 145–52. doi: [10.1016/j.femsec.2005.01.003](https://doi.org/10.1016/j.femsec.2005.01.003).
- Rui, Y., P. Wan, G. Chen, M. Xie, Y. Sun, X. Zeng, and Z. Liu. 2019. Simulated digestion and fermentation *in vitro* by human gut microbiota of intra- and extra-cellular polysaccharides from *Aspergillus cristatus*. *LWT - Food Science and Technology* 116:108508. doi: [10.1016/j.lwt.2019.108508](https://doi.org/10.1016/j.lwt.2019.108508).
- Saito, D., S. Nakaji, S. Fukuda, T. Shimoyama, J. Sakamoto, and K. Sugawara. 2005. Comparison of the amount of pectin in the human terminal ileum with the amount of orally administered pectin. *Nutrition (Burbank, Los Angeles County, Calif.)* 21 (9):914–9. doi: [10.1016/j.nut.2005.01.005](https://doi.org/10.1016/j.nut.2005.01.005).
- Sancho, R. A. S., J. D. R. P. Souza, F. A. de Lima, and G. M. Pastore. 2017. Evaluation of oligosaccharide profiles in selected cooked tubers and roots subjected to *in vitro* digestion. *LWT - Food Science and Technology* 76:270–7. doi: [10.1016/j.lwt.2016.07.046](https://doi.org/10.1016/j.lwt.2016.07.046).
- Schook, L. B., T. V. Collares, K. A. Darfour-Oduro, A. K. De, L. A. Rund, K. M. Schachtschneider, and F. K. Seixas. 2015. Unraveling the swine genome: Implications for human health. *Annual Review of Animal Biosciences* 3:219–44. doi: [10.1146/annurev-animal-022114-110815](https://doi.org/10.1146/annurev-animal-022114-110815).
- Shi, Y., J. Liu, Q. Yan, X. You, S. Yang, and Z. Jiang. 2018. *In vitro* digestibility and prebiotic potential of curdlan (1→3)-β-D-glucan oligosaccharides in *Lactobacillus* species. *Carbohydrate Polymers* 188: 17–26. doi: [10.1016/j.carbpol.2018.01.085](https://doi.org/10.1016/j.carbpol.2018.01.085).
- Shin, H., D. H. Seo, J. Seo, L. M. Lamothe, S. H. Yoo, and B. H. Lee. 2019. Optimization of *in vitro* carbohydrate digestion by mammalian mucosal α-glucosidases and its applications to hydrolyze the various sources of starches. *Food Hydrocolloids* 87:470–6. doi: [10.1016/j.foodhyd.2018.08.033](https://doi.org/10.1016/j.foodhyd.2018.08.033).
- Sivieri, K., M. L. V. Morales, S. M. I. Saad, M. A. T. Adorno, I. K. Sakamoto, and E. A. Rossi. 2014. Prebiotic effect of fructo-oligosaccharides in the simulator of the human intestinal microbial ecosystem (SHIME® model). *Journal of Medicinal Food* 17 (8):894–901. doi: [10.1089/jmf.2013.0092](https://doi.org/10.1089/jmf.2013.0092).
- Southgate, D. A. T. 1969. Determination of carbohydrates in foods I. – Available carbohydrate. *Journal of the Science of Food and Agriculture* 20:326–30.
- Strube, M. L., T. K. Jensen, A. S. Meyer, and M. Boye. 2015. *In situ* prebiotics: Enzymatic release of galacto-rhamnogalacturonan from potato pulp *in vivo* in the gastrointestinal tract of the weaning piglet. *AMB Express* 5 (1):8. doi: [10.1186/s13568-015-0152-1](https://doi.org/10.1186/s13568-015-0152-1).
- Tanabe, K., S. Nakamura, and T. Oku. 2014. Inaccuracy of AOAC method 2009.01 with amyloglucosidase for measuring non-digestible oligosaccharides and proposal for an improvement of the method. *Food Chemistry* 151:539–46. doi: [10.1016/j.foodchem.2013.11.121](https://doi.org/10.1016/j.foodchem.2013.11.121).
- Tanabe, K., S. Nakamura, K. Omagari, and T. Oku. 2015. Determination trial of nondigestible oligosaccharide in processed foods by improved AOAC Method 2009.01 using porcine small intestinal enzyme. *Journal of Agricultural and Food Chemistry* 63 (24):5747–52. doi: [10.1021/jf505844y](https://doi.org/10.1021/jf505844y).
- Tornero-Martínez, A., R. Cruz-Ortiz, M. E. Jaramillo-Flores, P. Osorio-Díaz, S. V. Ávila-Reyes, G. M. Alvarado-Jasso, and R. Mora-Escobedo. 2019. *In vitro* fermentation of polysaccharides from *Aloe vera* and the evaluation of antioxidant activity and production of short chain fatty acids. *Molecules* 24 (19):3605. doi: [10.3390/molecules24193605](https://doi.org/10.3390/molecules24193605).
- Turnbull, C. M., A. L. Baxter, and S. K. Johnson. 2005. Water-binding capacity and viscosity of Australian sweet lupin kernel fibre under *in vitro* conditions simulating the human upper gastrointestinal tract. *International Journal of Food Sciences and Nutrition* 56 (2): 87–94. doi: [10.1080/09637480500081080](https://doi.org/10.1080/09637480500081080).
- Verhoeckx, K., K. L. Bøgh, D. Dupont, L. Egger, G. Gadermaier, C. Larré, A. Mackie, O. Menard, K. Adel-Patient, G. Picariello, et al. 2019. The relevance of a digestibility evaluation in the allergenicity risk assessment of novel proteins. Opinion of a joint initiative of COST action ImpARAS and COST action INFOGEST. *Food and Chemical Toxicology* 129:405–23. doi: [10.1016/j.fct.2019.04.052](https://doi.org/10.1016/j.fct.2019.04.052).
- Villamiel, M., A. Montilla, A. Olano, and N. Corzo. 2014. Production and bioactivity of oligosaccharides derived from lactose. In *Food oligosaccharides: Production, analysis and bioactivity*, eds. F. J. Moreno and M. L. Sanz, 135–67. Hoboken, NJ: Wiley.
- Villanueva-Millán, M. J., P. Pérez-Matute, and J. A. Oteo. 2015. Gut microbiota: A key player in health and disease. A review focused on obesity. *Journal of Physiology and Biochemistry* 71 (3):509–25. doi: [10.1007/s13105-015-0390-3](https://doi.org/10.1007/s13105-015-0390-3).
- Woolnough, J. W., J. A. Monro, C. S. Brennan, and A. R. Bird. 2008. Simulating human carbohydrate digestion *in vitro*: A review of methods and the need for standardisation. *International Journal of Food Science & Technology* 43 (12):2245–56. doi: [10.1111/j.1365-2621.2008.01862.x](https://doi.org/10.1111/j.1365-2621.2008.01862.x).
- Wang, L., C. Li, Q. Huang, X. Fu, and R. H. Liu. 2019. *In vitro* digestibility and prebiotic potential of a novel polysaccharide from *Rosa roxburghii* Tratt fruit. *Journal of Functional Foods* 52:408–17. doi: [10.1016/j.jff.2018.11.021](https://doi.org/10.1016/j.jff.2018.11.021).
- Wang, Y., G. Chen, Y. Peng, Y. Rui, X. Zeng, and H. Ye. 2019. Simulated digestion and fermentation *in vitro* with human gut microbiota of polysaccharides from *Coralline pilulifera*. *LWT - Food Science and Technology* 100:167–74. doi: [10.1016/j.lwt.2018.10.028](https://doi.org/10.1016/j.lwt.2018.10.028).
- Yang, Y., C. Zhao, M. Diao, S. Zhong, M. Sun, B. Sun, H. Ye, and T. Zhang. 2018. The prebiotic activity of simulated gastric and intestinal digesta of polysaccharides from the *Hericium erinaceus*. *Molecules* 23 (12):3158–14. doi: [10.3390/molecules23123158](https://doi.org/10.3390/molecules23123158).