



## Aflatoxigenic fungi and mycotoxins in food: a review

Samuel Ayofemi Olalekan Adeyeye

To cite this article: Samuel Ayofemi Olalekan Adeyeye (2019): Aflatoxigenic fungi and mycotoxins in food: a review, Critical Reviews in Food Science and Nutrition, DOI: [10.1080/10408398.2018.1548429](https://doi.org/10.1080/10408398.2018.1548429)

To link to this article: <https://doi.org/10.1080/10408398.2018.1548429>



Published online: 28 Jan 2019.



Submit your article to this journal [↗](#)



View Crossmark data [↗](#)

REVIEW



## Aflatoxigenic fungi and mycotoxins in food: a review

Samuel Ayofemi Olalekan Adeyeye<sup>a,b</sup>

<sup>a</sup>Department for Management of Science and Technology Development, Ton Duc Thang University, Ho Chi Minh City, Vietnam; <sup>b</sup>Faculty of Environment and Labour Safety, Ton Duc Thang University, Ho Chi Minh City, Vietnam

### ABSTRACT

Food contamination is a common phenomenon in the production, distribution and consumption of processed and agricultural commodities all over the world. Food safety is now taking the frontal stage in food production, processing and distribution. This study assessed the presence of aflatoxigenic fungi and mycotoxins in foods, occurrence, control, socio-economic and health implications. This study also gave comprehensive information on the safety and mycological quality of foods as well the effect on the people and economy of various countries. The review revealed the various aflatoxigenic fungi in foods, their distribution in agricultural produce and their effects on reduction in yield and quality with attendant consumer health implications and resulting significant economic losses. In developing countries, majority of the people do not know the inherent dangers of consuming mouldy produce or food contaminated with fungi and moulds with possible contamination by mycotoxigenic fungi because of lack of awareness about the danger involved. In view of this, there is need for general and public education to sensitise the people on the economic and health hazards posed by mycotoxins. Control measures such as good agricultural practices, crop selection, proper washing and cooking practices of food commodities should be emphasized to the people. Regulatory control, fast and effective analyses and detection, good produce handling and storage should be encouraged as this will assist in mitigating the side effects of mycotoxins in foods particularly in the tropical and sub-tropical countries, and in Africa nations where there is enabling environment that promotes fungal growth, shortage of food and drought of modern storage and processing infrastructures.

### KEYWORDS

Aflatoxigenic fungi;  
mycotoxins; foods;  
occurrence; control

### Introduction

Fungi (*plural*, the *singular* noun is *fungus*) belong to the group of eukaryotic organisms which include yeasts, moulds and mushrooms (Moore, 1980, Hibbett et al., 2007). They are classified as a kingdom, Fungi, which differ from other eukaryotic life kingdoms of plants and animals (Hibbett et al., 2007).

Fungi are in a different kingdom from other plants, bacteria, and some protists due to the presence of chitin in their cell walls. This characteristic differentiates them from the likes of plants, bacteria, and some protists. Fungi are however, similar to animals because they are heterotrophs; possessing ability to absorb dissolved molecules digested through extracellular secretion of digestive enzymes into foods. Therefore, fungi do not have ability to produce their own food because they lack chlorophyll that is needed for photosynthetic processes (Hibbett et al., 2007).

Mycology is an emerging field of study which involves the study of fungi, fungal taxonomy, genetic and biochemical functions, and their use for food, medicinal and religious purposes and side effects of their secretions and metabolites (Struck, 2006).

Although some fungi are beneficial to man, however, several of them are known parasites to plants, animals (including man), and other fungi. Some fungi are pathogens to several crops causing extensive damage and agricultural losses. Common examples of fungal pathogens of importance are rice blast fungus *Magnaporthe oryzae* (Talbot, 2003), *Ophiostoma ulmi* and *Ophiostoma novo-ulmi* causing Dutch elm disease (Paoletti, et al., 2006) and *Cryphonectria parasitica* responsible for chestnut blight (Gryzenhout, et al., 2006). Crop pathogens of importance belong to these genera *Fusarium*, *Ustilago*, *Alternaria*, and *Cochliobolus* (Paszkowski, 2006). Some fungi are carnivores feeding on other organisms as predators, examples are *Paecilomyces lilacinus*, which feeds on nematodes, capturing them with constricting rings or adhesive nets (Yang et al., 2007). Several plant pathogenic fungi like *Phytophthora infestans*, could change from biotrophic i.e parasitic on living plants to necrotrophic i.e feeding on the dead tissues (Koeck, et al., 2011).

Some fungi are pathogenic to man and are disease causing organisms in humans, some of the diseases caused by them could be fatal if not treated. Such diseases include aspergillosis, candidiasis, coccidioidomycosis, cryptococcosis,

histoplasmosis, mycetomas, and paracoccidioidomycosis. Furthermore, immune-comprised people are particularly susceptible to disease by genera such as *Aspergillus*, *Candida*, *Cryptococcus* (Hube, 2004, Nielsen and Heitman, 2007, Brakhage, 2005), *Histoplasma* (Kauffman, 2007), and *Pneumocystis* (Cushion, *et al.*, 2007). Some fungi are also responsible for infections of the eyes, nails, hair, and skin, especially the so-called dermatophytic and keratinophilic fungi, and cause ringworm and athlete's foot infections (Cook and Zumla, 2008). Many allergic reactions and allergies are caused by fungal spores and fungi from different taxonomic groups (Simon-Nobbe, *et al.*, 2008).

## Uses of fungi

### Human use

Fungi have been used extensively for preparation and preservation of foods and other culinary applications. Mushrooms contain no cholesterol and are eaten as a good source of protein. Mushroom farming has become large enterprises in many countries of the world. Fungi are being used for large scale production of antibiotics, vitamins, and anti-cancer and cholesterol-lowering drugs and other natural products of economic importance (Brakhage, *et al.*, 2004, Fincham, 1989). Through genetic engineering fungal metabolism has been modified in recent times (Fincham, 1989), for example, some yeast species have genetically modified (Hawkins and Smolke, 2008) to make them grow faster during fermentation which has tremendous impacts on food and pharmaceutical production and beverage industries worldwide (Brakhage, *et al.*, 2004).

### Therapeutic uses

Many fungi produce metabolites that contain pharmacologically active drugs for treating man and animals. Among these important active ingredients are antibiotics, including the penicillins. However, artificial penicillins produced through chemical modifications of fungal synthesized penicillins have wider spectrum when compared with penicillin G (produced by *Penicillium chrysogenum*) with a narrower spectrum of biological activity. According to Pan, *et al.*, 2008, artificial penicillins are semisynthetic compounds which are gotten from fermentation cultures, and modified structurally to obtain specific desirable properties. Other antibiotics of importance produced by the same method are: ciclosporin, an immunosuppressant drug used in transplant surgery and fusidic acid, an antibiotic for controlling infection from methicillin-resistant *Staphylococcus aureus* bacteria (Fajardo and Martínez, 2008). Griseofulvin is also produced from *Penicillium griseofulvum* and for treating eczema, skin infections etc and other fungal infections (Manzoni and Rollini, 2002).

## Traditional and folk medicine

Fungi are used in many Asian countries for traditional and folk medicine; certain mushrooms such as *Agaricus subrufescens* (Yuen and Gohel, 2005, Paterson, 2006), *Ganoderma lucidum* (Paterson, 2008), and *Ophiocordyceps sinensis* (Kulp, 2000) are used for treating different ailments in traditional Chinese medicine.

## Cultured foods

In food processing, fungi have been used for centuries in baking. *Saccharomyces cerevisiae* or baker's yeast, a unicellular fungus, is used for production bread and other wheat-based products (Piskur, *et al.*, 2006). Also, yeast species of the genus *Saccharomyces* are used in fermentation processes to produce alcoholic beverages such beer, rum, stout etc. (Abe, *et al.*, 2006). In Asia, *Rhizopus* species are used for making tempeh from soybeans, a very important Asian food (Jørgensen, 2007).

## In food

Mushrooms are sources of good quality proteins. Edible mushrooms could be cultivated or wildy grown. Several mushrooms are consumed in different ways, Commonly cultivated and edible mushrooms are *Agaricus bisporus* used in preparing salads, soups, and many other dishes in the West. Some African and Asian fungi such as straw mushrooms (*Volvariella volvacea*), oyster mushrooms (*Pleurotus ostreatus*), shiitakes (*Lentinula edodes*), and enokitake (*Flammulina* spp) (Erdogan *et al.*, 2003) are very common and are now commercially grown and eaten in the West.

Many other wild mushroom species are also commonly eaten (Orr and Orr, 1979). Examples are milk mushrooms, morels, chanterelles, truffles, black trumpets, and *porcini* mushrooms (*Boletus edulis*) (Orr and Orr, 1979).

Some fungal species are used to inoculate milk curds to give characteristic flavour and texture to the cheese and they are very important in cheese production. An example of such fungi is *Penicillium roqueforti* which imparts the characteristic blue colour to cheese with such as Roquefort cheese, by inoculating milk with the fungi (Kinsella and Hwang, 1976; Orr and Orr, 1979). Some moulds used to produce cheese are non-toxic and safe for human consumption but some species could produce mycotoxins like aflatoxins, roquefortine C, patulin which may accumulate as a result of growth of mixed culture of fungi during cheese ripening or storage (Kinsella and Hwang, 1976, Hachmeister and Fung, 1993).

## Types of aflatoxigenic fungi

*Aspergillus* species are groups of moulds widely distributed all over the world. They can easily contaminate foodstuffs and animal feeds when conditions are favourable especially in the tropical and sub-tropical countries where relative humidity and temperature are very high. The genus

**Table 1.** Mycotoxins in staple grains and seeds.

Mycotoxin	Commodity	Fungal source(s)	Effects of ingestion	Reference
Deoxynivalenol/nivalenol	Wheat, maize, barley	<i>Fusarium graminearum</i> <i>Fusarium crookwellense</i>	Human toxicoses in India, China, Japan, and Korea. Toxic to animals, especially pigs	(FAO 2004)
Zearalenone	Maize, wheat	<i>Fusarium culmorum</i> <i>F. graminearum</i> <i>F. culmorum</i> <i>F. crookwellens</i>	Identified by the International Agency for Research on Cancer (IARC) as a possible human carcinogen. Affects reproductive system in female pigs	(FAO 2004)
Ochratoxin A	Barley, wheat, and many other commodities	<i>Aspergillus ochraceus</i> <i>Penicillium verrucosum</i>	Suspected by IARC as human carcinogen. Carcinogenic in laboratory animals and pigs	(FAO 2004)
Fumonisin B1	Maize	<i>Fusarium moniliforme</i> plus several less common species	Suspected by IARC as human carcinogen. Toxic to pigs and poultry. Cause of equine leukoencephalomalacia (ELEM), a fatal disease of horses	(FAO 2004)
Aflatoxin B1, B2 Aflatoxin B1, B2, G1, G2	Maize, peanuts, and many other commodities Maize, peanuts	<i>Aspergillus flavus</i> <i>Aspergillus parasiticus</i>	Aflatoxin B1, and naturally occurring mixtures of aflatoxins, identified as potent human carcinogens by IARC. Adverse effects in various animals, especially chickens	(FAO 2004)

Source: Adapted from Adeyeye (2016).

*Aspergillus* is an important mould in food processing and industrial food (Adejumo & Adejoro, 2014, Onions, *et al.*, 1981, Richard, 2007). *Aspergillus* species are moulds of importance in food spoilage. Spoilage due to contamination by *Aspergillus* species is responsible for changes like sensorial, nutritional and microbial such as pigmentation, discoloration, rotten, development of off odours and off flavours. Several *Aspergillus* species grew in foods very with low water activity and produced mycotoxins (Adejumo & Adejoro, 2014; Lee, *et al.*, 2004). *Aspergillus niger* are common mould all over the world and with wide distribution on a variety of substrates. *Aspergillus niger* is the most common mould involve in post-harvest decay and losses of foods and agricultural produce (Ashiq, 2015, Pitt, *et al.*, 2000).

*Penicillium* is also group of mould with wide distribution in foods and produce. *Penicillium* species have been found in fish and fish feed (Barbosa *et al.*, 2013). Some *Penicillium* species are responsible for producing toxic compounds like citrin and citreoviridin (Barbosa *et al.*, 2013, Richard, 2007).

*Fusarium* is another mould of importance which affects foods and feeds. It is a field mould which is widely distributed worldwide. Common mycotoxins produced by *Fusarium* are deoxynivalenol and zearalenone (Barbosa *et al.*, 2013, Richard, 2007). *Fusarium* is a group of fungi of economic importance because they produce highly potent mycotoxins.

### Mycotoxins produced by aflatoxigenic fungi

Mycotoxins are metabolites secreted by fungi growing on organic substrates of diverse nature which could produce fatal or side effects when ingested or consumed by humans or animals. Fatal effects could be vomiting, weight loss, tumors growth, and death. Several toxics secreted by fungi are well known, and majority of them occur in grain crops. Most mycotoxins affect organs in the body and could be fatal to blood, kidneys, skin, or central nervous system, and

while some are known carcinogens (Ashiq, 2015; Yin, *et al.*, 2008; Martins *et al.*, 2001).

Fungi are the major source of mycotoxins in foods. *Aspergillus*, *Penicillium*, and *Fusarium* are the fungi of greatest importance to humans with respect to food poison by mycotoxins. Aflatoxins are known as the most important fungal metabolites of a direct health hazard to humans. *Aspergillus flavus* is widely distributed in soil, air, and rotten plant residues. Aflatoxin production and contamination can take place in the field, in transit, or in storage. However, most infection takes places in the field while aflatoxin production takes place at any point under favourable conditions (Ashiq, 2015, Yin, *et al.*, 2008, Martins *et al.*, 2001).

### Major groups of mycotoxins in foods

Several toxigenic moulds have been identified, but only a few mycotoxin producing moulds, particularly those affecting cereals such as maize, wheat, barley, oats and rice and groundnuts are considered to be significant to human health (Bhat, *et al.*, 1989, Desjardins & Proctor, 2007, Jeswal & Kumar, 2015, Lewis *et al.*, 2005). Food contamination with mycotoxin could take place in the field as pre-harvest during growing period of crop plant or could occur after harvesting under favourable environmental conditions during processing, packaging, distribution, and storage (Pereira *et al.*, 2014, Bhatnagar *et al.*, 2006). Mycotoxins contamination occurs in crops such as cereals, legumes and oil seeds (Table 1) that are stored under high temperature and high relative humidity for a prolonged time (Bennett, 2003). Aflatoxins are the most common among the various mycotoxins found in most of agricultural produce for human consumption and animal feed.

Aflatoxins are the most common mycotoxin and potent human hepatocarcinogen. Aflatoxin is produced in maize and groundnuts in the field under drought and in stored. It is produced by *A. flavus* and *A. parasiticus* in tropical and

**Table 2.** Mycotoxins in Human blood/serum from different countries.

Country	Sampling period	Mycotoxin	Total samples	Positive samples	Detection technique	Range (µg/L)	Reference
Argentina	2004–2005	OTA	435	274	HPLC	0.15–0.43	(Pacin, et al. 2008)
Chile	2004–2005	OTA	88	62	HPLC	0.07–2.75	(Munoz et al. 2006)
Croatia (Kaniza)	2004–2005	OTA	4343	24.6%	HPLC	2–50	(Radic et al. 1997)
Czech Republic	2004–2005	OTA	115	115	HPLC/ELISA	0.037–1.13	(Dohnal et al. 2013)
Czech Republic	2004–2005	OTA	2206	2077	HPLC	0.1–13.7	(Ostry et al. 2005)
Czech Republic	2004–2005	OTA	100	100	HPLC	0.1–0.35	(Malir et al. 2013)
Egypt	2006	AFB1–Alb	34	98	ELISA	3.0–35.1 pg/mg	(Piekkola et al. 2012)
Egypt	1999, 2004	AFB1–Alb	24	24	ELISA	n.d.–32.8 pg/mg	(Turner et al. 2008)
Egypt	–	Total AFs (B1, B2, G1, M1, B2a, G2a,AFL)	60	60	Two–dimensional TLC	0.37–1.11 (Kwashiorkor) 0.15–0.38 (Marasmus)	(Hatem et al. 2005)
Germany	1990–1997	O TA	102	100/95	EIA/ HPLC	0.072–1.290	(Martlbauer et al. 2009)
Germany	2005–2006	O TA	61	61	HPLC	0.07–0.75	(Degen, Mayer, and Blaszkewicz 2007)
Ghana	–	AFB1–Alb	24	24	RIA	0.1–4.44 pmol/mg	(Tang et al. 2009)
Ghana	–	AFB1–Alb	64	64	RIA	0.3325–2.2703 pmol/mg	(Jiang et al. 2005)
Ghana (Kumasi)	2006	AFB1–Alb	507	507	HPLC	0.44–268.73 pg/mg	(Shuaib et al. 2010)
Guinea	2002	AFB1–Alb	250	245	ELISA	Nd–66 pg/mg	(Turner et al. 2005)
Italy	2007	O TA	130	129	HPLC	0.084 – 4.835	(Biasucci et al. 2011)
Italy	–	AFG1	71	1	LC/MS/MS	3.48	(Ritieni et al. 2010)
Italy (Tuscany)	1994–1996	O TA	138	97%	HPLC	0.12–2.84	(Palli et al. 1999)
Italy (Tuscany)	–	O TA	6	6	HPLC	0.94–3.28	(Iavicoli et al. 2002)
Ivory Coast	2001 and 2004	O TA	102	30	HPLC	0.01–5.81	(Sangare Tigori et al. 2006)
Japan	1992,1994–1996	O TA	184	156	ELISA	4–278	(Ueno et al. 1998)
Kenya	2007	AFB1–Alb	597	78%	LC/MS/MS	<0.02–211pg/mg	(Yard et al. 2013)
Lebanon	–	O TA	250	33%	HPLC	0.1– 0.87	(Assaf et al. 2004)
Malaysia	2008	AFB1–Alb	170	97%	HPLC	0.2–23.16 pg/mg	(Leong et al. 2012)
Pakistan	–	O TA	96	87	HPLC	0.032–3.409	(Aslam et al. 2005)
Pakistan	–	O TA	127	107	HPLC	0.03 to 3.41	(Aslam et al. 2012)
Poland	2005	O TA	25	25	HPLC	0.19–3.77	(Grajewski et al. 2007)
		O TA		25		0.4–60	(Jonsyn-Ellis 2007)
		O TB		6		0.03–5.6	
Sierra Leone	–	AFB1	131	23		0.2–74	
		AFB2		5		0.02–2.2	
		AFG1		29	HPLC	0.3–56	
		AFG2		22		0.01–1.2	
		AFM1		39		0.03–6.8	
		AFM2		51		0.04–48	
		AFL		30		0.01–3.2	
		O TA		325	HPLC	0.06–10.92	
		AFB1–Alb	325	119	ELISA	4.8–260.8 pg/mg	
		AFB1–Alb	119	179	HPLC	27.8 – 77.4 pg/mg	
Spain (Lleida)	2008–2009	O TA	357	109	HPLC	<0.1–5.8	(Coronel et al. 2011)
The Gambia	2000	O TA	109	27.5%	HPLC	0.1– 11.98	(Turner et al. 2007)
The Gambia	1992	O TA	40	68	HPLC	0.12 – 1.5	(Wild et al. 2000)
Tunisia	–	O TA	68	37	ELISA	0.59 (mean)	(Grosso et al. 2003)
Tunisia	2005	AFB1	37	410	HPLC	0.08–2.29	(Ghali et al. 2008)
Tunisia	–	AFB2	484	266		0.01–0.67	(Khalifa et al.2012)
Turkey	–	AFG1	325	409		0.03–1.24	(Giray et al. 2009)
Turkey	2007–2008	AFG2		401		0.01–1.03	(Aydin et al. 2011)
Turkey (Isparta)	–	O TA	133	133	HPLC	0.19–5.5	(Ozçelik, Koşar, and Soysal2001)
UK		AFB1–Alb	104	95%	HPLC/ELISA	28.1 ± 14.6 (mean) pg/mg	(Turner et al. 1998)
USA	2007–2008	AFB1–Alb	170	20.6%	HPLC	1.01–16.57 pg/mg	(Johnson et al. 2010)
USA	1999–2000	AFB1–Alb	27	2016	LC–MS/MS	≥ 0.02–0.20	(Schleicher et al. 2013)

Source: Adapted from Waseem et al. (2014).

(Key: OTA,OTB: ochratoxin A,B; AFB1,AFB2,AFG1, AFG2,AFM1,AFM2: aflatoxin B1,B2,G1,G2,M1,M2; AFL: aflatoxicol; AFB1 –Alb: aflatoxin B1 albumin adduct; HPLC: high performance liquid chromatography; LC: liquid chromatography, TLC: thin layer chromatography; ELISA: enzyme linked immune-sorbent assay; MS: mass spectrometry; RIA: radioimmunoassay; EIA: enzyme immunoassay)



## Aflatoxins




Figure 1. Structures of aflatoxins.




Figure 2. Ergotamine, a major mycotoxin produced by *Claviceps* species, which if ingested can cause gangrene, convulsions, and hallucinations.




Figure 3. Structure of patulin.

sub-tropical countries because of favourable weather conditions (Desjardins & Proctor, 2007, Jeswal & Kumar, 2015). Aflatoxins are known carcinogenic secondary metabolites of fungal origin (Adejumo and Adejoro, 2014) and also rarely produced by *A. nominus* and *A. pseudotamari* (Adejumo and Adejoro, 2014, Mazaheri, 2009). Among the 18 different types of aflatoxins identified, the four naturally occurring ones are Aflatoxin B<sub>1</sub> (AFB<sub>1</sub>), Aflatoxin B<sub>2</sub> (AFB<sub>2</sub>), Aflatoxin G<sub>1</sub> (AFG<sub>1</sub>) and Aflatoxin G<sub>2</sub> (AFG<sub>2</sub>) (Adejumo and Adejoro, 2014, Filazi and Sireli, 2013). Aflatoxin B<sub>1</sub> and B<sub>2</sub> are identified by strong blue fluorescence under ultraviolet light (Table 2), hence the 'B', while Aflatoxin G<sub>1</sub> and G<sub>2</sub> are identified by greenish yellow fluorescence (Adejumo and Adejoro, 2014, Kensler *et al.*, 2011). Other aflatoxins metabolites are Aflatoxin M<sub>1</sub> and M<sub>2</sub> and they are commonly

seen in foods of animal origin such as milk, eggs or blood of animals that are fed with AFB<sub>1</sub> and AFB<sub>2</sub>-contaminated feed (Adejumo and Adejoro, 2014, IARC, 2002, CAST, 2003).

Several species of *Fusarium* have been to be toxigenic and of importance in human and animal health. *Fusarium graminearum*, the causal organism of head blight and ear rot in cereals, is responsible for the production of deoxynivalenol, zearalenone and fusarin C. In Eastern Europe human toxicoses caused by ingesting T-2 toxin and related compounds has been linked to *F. sporotrichioides*. (Bayman & Baker, 2006, Desjardins & Proctor, 2007, Jeswal & Kumar, 2015).

It has been reported severally epidemics of head blight and maize ear rot of *Fusarium* which are common in Asia, Africa and South America where cereals are predominantly grown. The high rates of oesophageal cancer in southern Africa and parts of China linked with the consumption of maize contaminated with fusarins and fumonisins from *F. moniliforme* are of great concern at the moment (Bhat *et al.*, 1989, Desjardins & Proctor, 2007). Most of these mycotoxins produced by moulds that are of importance to human health are produced under poor storage conditions (Bhat *et al.*, 1989, Desjardins & Proctor, 2007, Jeswal & Kumar, 2015, Adeyeye, 2016).

Aflatoxins are chemical metabolites and mycotoxins produced by *Aspergillus* species of fungi, like *A. flavus* and *A. parasiticus* (Adejumo & Adejoro, 2014, Ashiq, 2015, Martins, *et al.*, 2001, Adeyeye, 2016). The common aflatoxins in foods and agricultural proce are (Figure 1) B<sub>1</sub>, B<sub>2</sub>, G<sub>1</sub>, and G<sub>2</sub> (Yin, *et al.*, 2008). Aflatoxin B<sub>1</sub>, the most toxic, is a potent carcinogen and has been directly correlated to adverse health effects, such as liver cancer, in many animal species (Ashiq, 2015, Martins *et al.*, 2001). Aflatoxins are




Figure 4. Structural formula of citrinin.




Figure 5. Structural formula of fumonisin B<sub>1</sub>.

mycotoxins largely related to agricultural produce produced in the tropics and subtropics under humid climate, cereals, legume and oil seeds (Ashiq, 2015, Martins *et al.*, 2001; Yin *et al.*, 2008, Adeyeye, 2016).

Three forms of ochratoxin have been identified. These are A, B, and C produced by *Penicillium* and *Aspergillus* species (Adeyeye, 2016). The difference between them is that Ochratoxin B (OTB) is a non-chlorinated form of Ochratoxin A (OTA) while Ochratoxin C (OTC) is an ethyl ester form Ochratoxin A (Ashiq, 2015, Bayman & Baker, 2006, Jeswal & Kumar, 2015, Adeyeye, 2016). Several species of *Aspergillus* have been implicated in secretion of ochratoxin. These include *Aspergillus ochraceus* responsible for contamination of beverages like beer and wine. *Aspergillus carbonarius* responsible for vine fruit contamination especially during juice making process. Ochratoxin A is a known carcinogen and nephrotoxin. Its activities have been seen as responsible for tumors in the human urinary tract but this needs further investigation (Ashiq, 2015, Bayman & Baker, 2006, Jeswal & Kumar, 2015, Adeyeye, 2016).

Citrinin has been detected *P. citrinum*, but currently several species of *Penicillium* and *Aspergillus* is implicated. Common ones among them is *Penicillium camemberti* used for cheese making and *Aspergillus oryzae* used for production of sake, miso, and soy sauce. Citrinin has been identified in yellowed rice disease in Japan and as a nephrotoxin in laboratory animals (Bennett & Klich, 2003, Adeyeye, 2016). Citrinin is widely linked with agricultural produce such as wheat, rice, corn, barley, oats, rye, and food coloured with Monascus pigment. Citrinin has been found to act synergistically with Ochratoxin A to depress RNA synthesis in murine kidneys but its side effects in human health is unknown. (Ashiq, 2015, Bennett & Klich, 2003, Jeswal & Kumar, 2015, Adeyeye, 2016).

Ergot has been detected as a toxin in the sclerotia of species of *Claviceps*. *Claviceps* infected cereals and consumption of cereal contaminated by ergot sclerotia especially bread produced from contaminated flour caused ergotism, a disease commonly referred to as St. Anthony's fire. Good

agricultural practices with modern storage techniques have eliminated ergotism as a human disease; but are still important in animal feed production. Ergot alkaloids are of importance in pharmaceutical industries (Ashiq, 2015, Bennett & Klich, 2003, Jeswal & Kumar, 2015).

Patulin has been isolated from *P. expansum*, *Aspergillus*, *Penicillium*, and *Paecilomyces* fungal species. *P. expansum* has been linked to moldy fruits and vegetables, especially rotting apples and Figs. (Moss, 2008, Trucksess & Scott, 2008, Adeyeye, 2016). However, fermentation process has been found to destroy it as it is not usually detected in apple beverages like cider. Patulin is non-carcinogenic but it has been implicated to damage the immune system of animals (Moss, 2008, Adeyeye, 2016).

Fusarium toxins have been detected developing cereal grains like maize, wheat etc and about 50 species of *Fusarium* have been implicated (Schaafsma & Hooker, 2007). Mycotoxins of importance from *Fusarium* are fumonisins, implicated in horses' nervous disorders and a proven carcinogen in rodents; trichothecenes, are known as causal of chronic and fatal toxicoses in animals and humans; and zearalenone, is not known for any fatal toxic effects in animals or humans. Other *Fusarium* toxins of importance are: beauvericin and enniatins, butenolide, equisetin, and fusarins (Desjardins & Proctor, 2007, Adeyeye, 2016).

### Health implications of eaten foods contaminated by mycotoxins

Mycotoxins are chemical metabolites of great economic importance and health implication to both animals and humans.

Aflatoxin ingestion has been implicated in hepatocellular carcinoma (liver cancer) (Adejumo and Adejoro, 2014, Williams *et al.*, 2004), reported to be the third-leading cause of cancer worldwide according to WHO (2008), with about 600,000 fresh cases each year. Adejumo and Adejoro, 2014, Hussein and Brussel, 2001 and Zain, 2011 have reported that aflatoxin contamination reduces feed intake, increase liver and kidney weights of farm animals, as well as induce immune-suppression and hepatitis as well as high mortality in farm animals.

Mycotoxins have been implicated in several countries for the outbreak of diseases. In India and Kenya mycotoxins have been implicated in the outbreak of aflatoxic hepatitis, in the outbreak of enteric ergotism in India; in the outbreak of vascular ergotism in Ethiopia; and in the outbreak of deoxynivalenol mycotoxicosis in India and China. The outbreaks have been linked to consumption of cereals that were grown and stored under drought and humid conditions during either the growing season or harvest or storage (Jelinek *et al.*, 1989).

Aflatoxins among all the mycotoxins were linked to human diseases such as liver cancer, Reye's syndrome, Indian childhood cirrhosis, chronic gastritis and kwashiorkor in many parts of the world, particularly in African and Asian countries where conditions for growth and storage




Figure 6. Structural formula of fumonisin B<sub>2</sub>.

encouraged fungi and mycotoxin production (Jelinek *et al.*, 1989, Palmgren & Hayes, 1987, Adeyeye, 2016).

Kenya recorded a serious and largest and most acute outbreaks of aflatoxicosis ever documented (CAST, 2003) in 2004, with 125 dead cases and over 317 others hospitalized for consuming aflatoxin-contaminated maize (Adeyeye, 2016). The nature of the outbreak was due to the consumption of maize contaminated with extremely toxic AFB1. The case-control studies carried out by Lewis *et al.*, (2005) showed that the affected people in the area consumed home-grown maize stored under humid conditions. During the outbreak, AFB1 individual daily exposure was 50mg/day (Probst *et al.*, 2007).

Several like Kashin Beck syndrome in the USSR, China and Viet Nam; Mseleni joint disease in southern Africa; endemic familial arthritis in India; alimentary toxic aleukia in the USSR; and oesophageal cancer in southern Africa were linked to *Fusarium* toxins. Also, Balkan endemic nephropathy and urinary tract tumours have been linked to Ochratoxins (Berry, 1988, Adeyeye, 2016).

### Economic implications of mycotoxins in foods

According to FAO, 1988, volume of global trade in agricultural produce like wheat, rice, barley, corn, sorghum, soybeans, groundnuts and oilseeds totalling several millions of tonnes on yearly basis. Agricultural produce are contaminated by mycotoxins due to unhygienic post-harvest handling of the crops and poor agricultural practices. Financial losses from mycotoxin-contamination of agricultural produce could be subjective, staggering and very difficult to obtain. Several other losses from mycotoxin contamination may include lower productivity; losses of foreign exchange earnings; costs involved in inspection, sampling and analysis of produce before and after shipments, all these could be large; losses to compensation paid in case of claims; farmers' subsidies to cover production losses; research, training and extension programme; costs of detoxification etc. could also be enormous when added together (Coulibaly, 1989, Adeyeye, 2016).

Economic impacts of mycotoxins on crops are very important, particularly cereals like wheat, maize, legumes like cowpea and groundnuts. According to Schmale III and Munkvold, (2018) over 25% of global harvest are involved on yearly basis, with total losses of 1 billion metric tons of foods and food products annually. The economic losses are as consequence of 1) crop yield losses as a result of toxigenic

fungi infestation; 2) value reduction due to mycotoxin contamination of crop; 3) animal productivity reduction due to mycotoxin-related animal health problems; and 4) costs of human health problems. Schmale III and Munkvold, (2018) also reported prevention, sampling, mitigation, litigation, and research costs.

Several researchers (Schmale III and Munkvold, 2018, Shephard, 2008, Richard, 2007, Munkvold, 2003) agreed that the economic impacts of mycotoxin contamination are felt by all stakeholders in food chain systems because of health care impacts and productivity losses. Schmale III and Munkvold, 2018 and Munkvold, 2003 estimated the costs of mycotoxins on produce in the USA and Canada annually to \$5 billion. However, this estimate does not include human health impacts or crop yield losses. They also estimated cost of aflatoxins in maize in the United States annually to \$225 million excluding costs of mitigation costs. The costs of aflatoxins in United States peanuts according to Schmale III and Munkvold, 2018 ranged over \$25.8 million losses per year between 1993 to 1996. Schmale III and Munkvold, 2018 also valued economic losses due to deoxynivalenol in the U.S.A to \$655 million annually, with the largest occurring in wheat. In developing countries, however, conflicting values are available, but according to Shephard, 2008 due to high aflatoxins contamination levels in the developing countries; economic losses may be far more than those obtaining in the United States. Shephard, 2008 revealed that economic losses in the three Asian countries of Indonesia, Philippines, and Thailand because of aflatoxins contamination exceeded \$900 million/year.

### Prevention and control of mycotoxins in foods

In several countries of the world, some preventive steps to reduce mycotoxin contamination in agricultural produce have been taken. The preventive measures are as follows:

#### Crop breeding

The contamination of produce could be reduced drastically to the barest minimum through effective crop breeding. By selection and adoption of mycotoxigenic fungi resistant crop varieties, mycotoxin contamination of the produce from the field could be drastically reduced. The adoption of this method has been used to reduce the problem of ergot contamination of cereals such as rye, wheat and pearl millet that are resistant to mycotoxigenic fungi. However, little success has been achieved in using the method to reduce aflatoxins contamination of corn and peanuts (Fox & Howlett, 2008, Jelinek *et al.*, 1989, Adeyeye, 2016).

#### Good Agricultural Practice

Aflatoxigenic fungi infestation on crops and subsequent mycotoxin formation in agricultural produce could be minimized during production and before harvest through adoption of good agricultural practices like crop rotation, farm irrigation, adequate weed control, adoption of mould-



resistant crop varieties, and use of bio-control like use of non-mycotoxigenic fungal strains.

### **Crop Rotation**

This is a cropping system that encourages farmers not plant crops that belong to the same family to follow each in a rotation. Crops that are affected by same toxigenic fungi should not be allowed to follow each other in a rotation. For example, wheat and maize that are highly susceptible to toxigenic *Fusarium* spp, attack should not be allowed to follow each other in a rotation system (Adejumo and Adejoro, 2014, CAC, 2003, Munkvold, 2003).

### **Sanitation**

In order to eliminate attack by toxigenic fungi on the produce, hygienic removal, destruction and disposal of wastes of previous harvested crops will assist in reducing toxigenic fungi contamination of the produce from the field (Adejumo and Adejoro, 2014). Cleaning of the stores and proper disinfection with appropriate fungicides before storing new produce has been found to reduce mycotoxin contamination of produce (Adejumo and Adejoro, 2014, Adeyeye, 2016, Hell *et al.*, 2000).

### **Insect Management**

Insect infestation and damages could be sources of entry for toxigenic fungi into the produce. According to Bankole and Adebajo, 2003, the extent of mycotoxins contamination of produce could be determined by the level of insect damage of the produce. Insect damage has been found to be a good entry point for *Fusarium* mycotoxins contamination of maize grains (Adejumo and Adejoro, 2014, Avataggio *et al.*, 2002). Insects could carry spores of mycotoxigenic fungi from crop surfaces to the interior of the produce through created wounds as a result of their feeding habits and thereby promoting mycotoxin contamination of produce (Adejumo and Adejoro, 2014, Munkvold, 2003, Ariyo *et al.*, 2013). Good pest control system could be an appropriate way to reduce mycotoxins contamination.

### **Proper Irrigation and Fertilization**

Mycotoxin development in produce has been linked to water stress during crop production. Adoption of irrigation and proper fertilization during crop production could reduce plants' stress which is sometimes responsible for mycotoxin development in produce. It has been reported that applying fertilizer and other soil conditioners to the soil to give adequate soil pH and enough plant nutrients are critical in mitigating against mycotoxin development in produce (Adejumo and Adejoro, 2014; CAC RC, 2003, Munkvold, 2003).

### **Early harvesting**

Fungal infection of crops in the field as well as post-harvest contamination of the harvested produce is greatly minimized by early harvesting. In Nigeria, unpredictable weather, lack of good storage facilities and space, labour constraint, farmers' financial need, attacks by rodents and thieves are among factors that force farmers to harvest their produce at inappropriate times (Amyot, 1983). Early harvesting and threshing of groundnuts have been found to reduce aflatoxin levels and increase gross returns on produce than in delayed harvesting according to Rachaputi *et al.*, (2002).

### **Proper drying**

Several researchers (Adejumo and Adejoro, 2014, Lanyasunya *et al.*, 2005, Ariyo *et al.*, 2013) revealed that proper and quick dehydration of produce to reduced moisture content is critical to reducing mycotoxin contamination of agricultural produce by creating unfavourable conditions for fungal growth and reproduction, insect attack and preserves produce longer. Rapid drying immediately after harvest has been found to mitigate against secretion of mycotoxins by fungi in produce. According to Ariyo *et al.*, 2013 and Adejumo and Adejoro, 2014, reported that reducing the water activity (aw) of produce to about 0.7 by reducing water content produce and maintaining produce below 0.7 aw is an effective way to control fungal spoilage and mycotoxin production in agricultural produce.

Reducing mycotoxins contaminations through postharvest steps include quick dehydration by mechanical means and maintaining crops in dried form. Sorting out contaminated produce by physical means, colour and proper washing with water have been found to reduce mycotoxins in produce.

Chemical methods of detoxification include ammoniation processes. Detoxification of aflatoxins in produce has been tried by many countries such as Senegal However, detoxification procedure should involve safety and efficacy tests for good result but this will surely increase handling costs (Fox & Howlett, 2008, Jelinek *et al.*, 1989).

According to Adejumo and Adejoro, 2014, several developing countries have undertaken some successful steps and measures to control mycotoxins like contaminated peanuts segregation in Malawi, peanut meal detoxification for export in Senegal, mycotoxins regulation in animal feed in Zimbabwe, adoption of resistant peanut varieties less susceptible to aflatoxin contamination in Burkina Faso, better produce-handling practices in Nigeria during the 1960s and in The Gambia during the 1990s.

### **Physical Separation**

Mycotoxin contamination in produce could be minimized through physical treatments by sorting, winnowing, washing, crushing and de-hulling to remove moulds and infested grains and thereby reducing mycotoxin concentration in the produce (Adejumo and Adejoro, 2014, Fandohan *et al.*, 2005).

## Biological Control

Mycotoxins are biologically controlled by using non-mycotoxigenic microorganisms to compete for space and nutrients, production of anti-mycotoxigenic metabolites by co-existing with mycotoxigenic microorganisms. Substratum pH changes coupled with a combination of other factors such as use of bio-control have been found to be effective in controlling fungal infestation and mycotoxin contamination in produce (Adejumo and Adejoro, 2014, Bianchini and Bullerman, 2010). Dorner and Cole (2002) and Adejumo and Adejoro, 2014 reported that post-harvest contamination of aflatoxins was minimized by a field application of non-toxigenic strains of *A. flavus* and *A. parasiticus*.

## Legislations

Legislations relating to mycotoxins have been introduced by several countries. However, attentions in those nations have been shifted to the major mycotoxins like aflatoxins, ergot alkaloids, deoxynivalenol and ochratoxins. Most of these legislative measures are yet to be harmonized but Codex Alimentarius Commission is making concerted efforts to establish international guidelines and rules to guide the amounts of mycotoxins, and aflatoxins in particular permissible in produce (Fox & Howlett, 2008, Jelinek *et al.*, 1989).

Implementation of mycotoxin control measures have been undertaken for agricultural produce involving in international export or in countries with large-scale buying and distribution systems for international trade. However, in many developing countries especially in Africa, where food production and consumption involved subsistence agriculture greater percentage of the population, such measures would be very difficult to implement (Adeyeye, 2016, Fox and Howlett, 2008, Jelinek *et al.*, 1989).

According to Adejumo and Adejoro, 2014, "the Codex Alimentarius Standard on aflatoxin limits are 0.05 µg/l for milk and milk products, 4 to 5 µg/kg for beans, 10 µg/kg in nuts such as peanuts and almonds, and 20 µg/kg for cereals. The acceptable limit for fumonisins is ≤1000 µg/kg (Adejumo and Adejoro, 2014, CAST, 2003). These limits are also adopted in Nigeria (Adejumo and Adejoro, 2014). Also, Adejumo and Adejoro, 2014 stated that the EU limit for OTA in raw cereal is 0.5 µg/kg, with a tolerable weekly intake (TWI) of 0.12 µg/kg, while maximum DON limits are 200 µg/kg for processed cereal based foods and baby infants formula and foods for young children, and 1750 µg/kg for unprocessed wheat, oats and maize (Adejumo and Adejoro, 2014, EC, 2006).

Apart from the above intervention strategies, Atanda *et al.*, (2013) recommended the following mycotoxins control strategies in Nigeria: a database collection of predominant fungi and mycotoxin in Nigeria, a mycotoxin occurrence map establishment to know the areas susceptible to mycotoxin contamination and a permanent culture collection media establishment (Adejumo and Adejoro, 2014). They believed that these strategies will reduce mycotoxin infection and reduce cancer prevalence in Nigeria.

## Conclusions

Fungi are very important organisms and their usefulness and economic importance cannot be overstated. However, many of them produced metabolites that are poisonous or dangerous to humans and animals. These metabolites are called mycotoxins. Mycotoxins are important because their implications in animal and human health are substantial and the economic importance widely studied. This review has shown the occurrence of mycotoxins in foods and feed in Nigeria and many parts of the world. Unfortunately, majority of the populace in these developing countries do not know the inherent dangers of consuming mouldy produce with possible contamination by mycotoxigenic fungi because of lack of awareness about the danger involved. In view of this, there is need for general and public education to sensitise the people on the economic and health hazards posed by mycotoxins. Control measures such as physical selection, proper washing and cooking practices of food commodities should be emphasized. Regulatory control, quick and effective analyses and detection, good produce handling and storage should be encouraged as this will assist in minimizing the side effects of mycotoxins contamination of foods especially in Africa where there is enabling environment for the growth of mycotoxigenic fungi, shortage of food and draught of modern storage and processing infrastructures.

## References

- Abe, K., K. Gomi, F. Hasegawa, and M. Machida. 2006. Impact of *aspergillus oryzae* genomics on industrial production of metabolites. *Mycopathologia* 162 (3):143–153. doi:10.1007/s11046-006-0049-2.
- Adejumo, T. O., and D. O. Adejoro. 2014. Incidence of aflatoxins, fumonisins, trichothecenes and ochratoxins in Nigerian foods and possible intervention strategies. *Food Science and Quality Management* 31 :127–146.
- Adeyeye, S. A. O. 2016. Fungal mycotoxins in foods: A review, *Cogent Food & Agriculture*, 2: 1213127.
- Amoyot, J. 1983. Social and economic aspects of dryer use for paddy and other agricultural produce in Thailand. Chulalongkorn University Social Research Institute and International Development Research Center.
- Ariyo, A.L., A.O. Peter, S.N. Abdulazeez, and O.H. Olatunde. 2013. Avian mycotoxicosis in developing countries. In *Mycotoxin and food safety in developing countries*, ed. H.A. Makun, 93–119. Rijeka, Croatia: InTech.
- Aslam, M., A. E. Beg, M. Blaszkewicz, G. H. Degen, and K. Golka. 2005. Ochratoxin a blood concentration in healthy subjects and bladder cancer cases from Pakistan. *Mycotoxin Research* 21 (3):164.
- Aslam, M., S. A. H. Rivzi, A. E. Beg, M. Blaszkewicz, K. Golka, and G. H. Degen. 2012. Analysis of ochratoxin a blood levels in bladder cancer cases and healthy persons from Pakistan. *Journal of Toxicology and Environmental Health. Part A* 75 (19–20):1176.
- Assaf, H., A. M. Betbeder, E. E. Creppy, M. Pallardy, and H. Azouri. 2004. Ochratoxin a levels in human plasma and foods in Lebanon. *Human & Experimental Toxicology* 23 (10):495.
- Ashiq, S. 2015. Natural occurrence of mycotoxins in food and feed: Pakistan perspective. *Comprehensive Reviews in Food Science and Food Safety* 14 :159–175. <http://dx.doi.org/10.1111/crf3.2015.14.issue-2>. doi:10.1111/1541-4337.12122.
- Atanda, O., H. A. Makun, I. M. Ogara, M. Edema, K. O. Idahor, M. E. Eshiett, and B. F. Oluwabamiwo. 2013. Fungal and mycotoxin contamination of Nigerian foods and feeds Countries. In *Mycotoxin and*

- food safety in developing countries, ed. H. A. Makun, 3–38. Rijeka, Croatia: InTech.
- Avantaggio, G., F. Quaranta, E. Desidero, and A. Visconti. 2002. Fumonisin contamination of maize hybrids visibly damaged by sesamia. *Journal of the Science of Food and Agriculture* 83 :13–18. doi:10.1002/jsfa.1272.
- Aydin A., H. Aksu, and U. Gunsen. 2011. Mycotoxin levels and incidence of mould in Turkish rice. *Environmental Monitoring and Assessment* 178: 271–280.
- Barbosa, T. S., C. M. Pereyra, C. A. Soleiro, E. O. Dias, A. A. Oliveira, K. M. Keller, P. P. O. Silva, L. R. Cavaglieri, and C. A. R. Rosa. 2013. Mycobiota and mycotoxins present in finished fish feeds from farms in the Rio De Janeiro state. *International Aquatic Research* 5 :1–9. doi:10.1186/2008-6970-5-3.
- Bayman, P., and J. L. Baker. 2006. Ochratoxins: A global perspective. *Mycopathologia* 162(3):215–223. doi:10.1007/s11046-006-0055-4.
- Berry, C. L. 1988. The pathology of mycotoxins. *The Journal of Pathology* 154 (4):301–311.
- Bhatnagar, D., J. W. Cary, K. Ehrlich, J. Yu, and T. E. Cleveland. 2006. Understanding the genetics of regulation of aflatoxin production and *Aspergillus flavus* development. *Mycopathologia* 162 (3):155–166. 2006. doi:10.1007/s11046-006-0050-9.
- Bhat, R. V., Y. Ramakrishna, S. R. Beedu, and K. L. Munshi. 1989. Outbreak of trichothecene mycotoxicosis associated with consumption of mould-damaged wheat products in Kashmir valley, India. *The Lancet* 333 (8628):35–37. doi:10.1016/S0140-6736(89)91684-X.
- Bianchini, A., and L.B. Bullerman. 2010. Biological Control of molds and mycotoxins in foods in mycotoxin prevention and control in agriculture. In *ACS symposium series 1031*, ed. M. Appell. Washington, DC: American Chemical Society.
- Biasucci, G., G. Calabrese, R. Di Giuseppe, G. Carrara, F. Colombo, B. Mandelli, M. Maj, T. Bertuzzi, A. Pietri, and F. Rossi. 2011. The presence of ochratoxin a in cord serum and in human milk and its correspondence with maternal dietary habits. *European Journal of Nutrition* 50 (3):211.
- Brakhage, A. A. 2005. Systemic fungal infections caused by *Aspergillus* species: Epidemiology, infection process and virulence determinants. *Current Drug Targets* 6 (8):875–886. doi:10.2174/138945005774912717.
- Brakhage, A. A., P. Spröte, Q. Al-Abdallah, A. Gehrke, H. Plattner, and A. Tüncher. 2004. Regulation of penicillin biosynthesis in filamentous fungi. *Advances in Biochemical Engineering/Biotechnology* 88 :45–90. doi:10.1007/b99257. ISBN 978-3-540-22032-9.
- Codex Alimentarius Commission (CAC). 2003. Code of practice for the prevention and reduction of mycotoxin contamination in cereals, including annexes on ochratoxin a, zearalenone, fumonisins and tricothecenes. Accessed 10 August, 2014. [http://www.fao.org/fileadmin/user\\_upload/agns/pdf/CXS\\_051e.pdf](http://www.fao.org/fileadmin/user_upload/agns/pdf/CXS_051e.pdf).
- Council for Agricultural Science and Technology (CAST). 2003. *Mycotoxins: Risks in plant, animal, and human systems*. Ames, IA: Task Force Report. 139.
- Cook, G. C., and A. I. Zumla. 2008. *Manson's tropical diseases: Expert consult*. Edinburgh, Scotland: Saunders Ltd. p. 347. ISBN 1-4160-4470-1
- Coronel, M. B., V. Sanchis, A. J. Ramos, and S. Marin. 2011. Ochratoxin a in adult population of lleida, Spain: Presence in blood plasma and consumption in different regions and seasons. *Food and Chemical Toxicology* 49(10):2697. doi:10.1016/j.fct.2011.07.045.
- Coulbaly, B. 1989. The problem of aflatoxin contamination of groundnut and groundnut products as seen by the African Groundnut Council. In *Aflatoxin contamination of groundnuts: Proceedings of the international workshop*, eds. D. McDonald and V. K. Mehan, 47–55. Patancheru: International Crops Research Institute for the Semi-Arid Tropics.
- Cushion, M. T., A. G. Smulian, B. E. Slaven, T. Sesterhenn, J. Arnold, C. Staben, A. Porollo, R. Adamczak, and J. Meller. 2007. Transcriptome of *Pneumocystis carinii* during fulminate infection: carbohydrate metabolism and the concept of a compatible parasite. *PLoS One* 2 (5):e423. Bibcode:2007PLoS...2.423C. doi:10.1371/journal.pone.0000423.PMC 1855432
- Degen, G. H., S. Mayer, and M. Blaszkewicz. 2007. Biomonitoring of ochratoxin a in grain workers. *Mycotoxin Research* 23(2):88).
- Desjardins, A. E., and R. H. Proctor. 2007. Molecular biology of fusarium mycotoxins. *International Journal of Food Microbiology* 119(1-2):47–50. doi:10.1016/j.ijfoodmicro.2007.07.024.
- Desjardins, D. E., B. A. Perry, D. J. Lodge, C. V. Stevani, and E. Nagasawa. 2010. Luminescent mycena: new and noteworthy species. *Mycologia* 102 (2):459–477. doi:10.3852/09-197.
- Dohnal, V., V. Dvořák, F. Malír, V. Ostrý, and T. Roubal. 2013. A comparison of ELISA and HPLC methods for determination of ochratoxin a in human blood serum in the Czech Republic. *Food and Chemical Toxicology: An International Journal Published for the British Industrial Biological Research Association* 62 :427
- Dorner, J. W., and R. J. Cole. 2002. Effect of application of nontoxigenic strain of *Aspergillus flavus* and *A. parasiticus* on subsequent aflatoxin contamination of peanuts in storage. *Journal of Stored Products Research* 38 (4):329–339. doi:10.1016/S0022-474X(01)00035-2.
- European Commission (EC). 2006. Commission regulation (EC) No 1881/2006 of 19 december 2006 setting maximum levels for certain contaminants in foodstuffs (text with EEA relevance). *Official Journal of the European Union* 364 :5–24.
- Erdogan, A., M. Gurses, and S. Sert. 2003. Isolation of moulds capable of producing mycotoxins from blue mouldy tulum cheeses produced in Turkey. *International Journal of Food Microbiology* 85 (1–2): 83–85. doi:10.1016/S0168-1605(02)00485-3.
- FAO. 1988. *FAO trade yearbook*, 1987, 380. Rome: Author. Food and Agriculture Organization (FAO). 2004. Worldwide regulations for mycotoxins in food and feed in 2003. (p. 81). Rome: Author.
- Fajardo, A., and J. L. Martinez. 2008. Antibiotics as signals that trigger specific bacterial responses. *Current Opinion in Microbiology* 11 (2): 161–167. doi:10.1016/j.mib.2008.02.006.
- Fandohan, P., D. Zoumenou, D. J. Hounhouigan, W. F. O. Marasas, M. J. Wingfield, and K. Hell. 2005. Fate of aflatoxins and fumonisins during the processing of maize into food products in Benin. *International Journal of Food Microbiology* 98 (3):249–259.
- Filazi, A., and U.T. Sireli. 2013. Occurrence of aflatoxins in food. In: *Aflatoxins – Recent advances and future prospects*, ed. Mehdi Razzaghi-Abyaneh, 143–170, Rijeka, Croatia: InTech.
- Fincham, J. R. 1989. Transformation in fungi. *Microbiological Reviews* 53 (1):148–170.
- Food and Drug Administration (FDA). 2000. Conference on mycotoxins in animal feeds, grains and food related to human and animal health, Rockville: Maryland.
- Fox, E. M., and B. J. Howlett. 2008. Secondary metabolism: Regulation and role in fungal biology. *Current Opinion in Microbiology* 11 (6): 481–487. doi:10.1016/j.mib.2008.10.007.
- Ghali, R., K. Hmaissia-Khlifa, C. Mezigh, H. Ghorbel, K. Maaroufi, S. Maschgoul, and A. Hedilli. 2008. HPLC determination of ochratoxin a in a low volume of human blood serum. *Analytical Letters* 41(5): 757. doi:10.1080/00032710801934528.
- Giray, B., P. Erkekoglu, S. Aydin, S. Sabuncuoglu, and G. Sahin. 2009. Determination of ochratoxin A and total aflatoxin levels in corn samples from Turkey by enzyme-linked immunosorbent assay. *Mycotoxin Research* 25 (2):113–116 doi:10.1007/s12550-009-0016-0.
- Grajewski, J., P. Jarzemski, M. Twaruzek, K. Kuzminska, and M. Trepala. 2007. The level of ochratoxin a in patients after nephrectomy. *Mycotoxin Research* 23 (1):22.
- Grosso, F., S. Said, I. Mabrouk, J. M. Fremy, M. Castegnaro, M. Jemmali, and S. Dragacci. 2003. New data on the occurrence of ochratoxin a in human sera from patients affected or not by renal diseases in Tunisia. *Food and Chemical Toxicology: An International Journal Published for the British Industrial Biological Research Association* 41 (8):1133.
- Gryzenhout, M., B. D. Wingfield, and M. J. Wingfield. 2006. New taxonomic concepts for the important forest pathogen *Cryphonectria parasitica* and related fungi. *FEMS Microbiology Letters* 258(2): 161–172. doi:10.1111/j.1574-6968.2006.00170.x.
- Hachmeister, K. A., and D. Y. Fung. 1993. Tempeh: A mold-modified indigenous fermented food made from soybeans and/or cereal



- grains. *Critical Reviews in Microbiology* 19 (3):137–188. doi:10.3109/10408419309113527.
- Hatem, N. L., H. Hassab, E. M. Abd Al-Rahman, S. A. El-Deeb, and R. L. El-Sayed Ahmed. 2005. Prevalence of aflatoxins in blood and urine of Egyptian infants with protein-energy malnutrition. *Food and Nutrition Bulletin* 26(1):49.
- Hawkins, K. M., and C. D. Smolke. 2008. Production of benzylisoquinoline alkaloids in *Saccharomyces cerevisiae*. *Nature Chemical Biology* 4 (9):564–573. doi:10.1038/nchembio.105.
- Hell, K., K. F. Cardwell, M. Setamou, and H. M. Poehling. 2000. The influence of storage practices on aflatoxin contamination in maize in four agroecological zones of Benin, West Africa. *Journal of Stored Products Research* 36 (4):365–382. doi:10.1016/S0022-474X(99)00056-9.
- Hibbett, D. S., M. Binder, J. F. Bischoff, M. Blackwell, P. F. Cannon, O. E. Eriksson, S. Huhndorf, T. James, P. M. Kirk, R. Lücking, et al. 2007. A higher-level phylogenetic classification of the fungi" (PDF). *Mycological Research* 111 (5):509–547. doi:10.1016/j.mycres.2007.03.004.
- Hube, B. 2004. From commensal to pathogen: Stage- and tissue-specific gene expression of *Candida albicans*. *Current Opinion in Microbiology* 7 (4):336–341. doi:10.1016/j.mib.2004.06.003.
- Hussein, H. S., and J. M. Brasel. 2001. Toxicity, metabolism, and impact of mycotoxins on humans and animals. *Toxicology* 167 (2):101–134.
- Iavicoli, I., C. Brera, G. Carelli, R. Caputi, A. Marinaccio, and M. Miraglia. 2002. External and internal dose in subjects occupationally exposed to ochratoxin A. *International Archives of Occupational and Environmental Health* 75 (6):381.
- International Agency for Research on Cancer (IARC). 2002. Some traditional herbal medicines, some mycotoxins, naphthalene and styrene. *IARC Monographs on the Evaluation of Carcinogenic Risks to Humans* 82 :1–556.
- Jelinek, C. F., A. E. Pohland, and G. E. Wood. 1989. Worldwide occurrence of mycotoxins in foods and feeds (an update). *Journal of the Association of Official Analytical Chemists* 72 :223–230.
- Jeswal, P., and D. Kumar. 2015. Mycobiota and natural incidence of aflatoxins, ochratoxin A, and citrinin in Indian spices confirmed by LC-MS/MS. *International Journal of Microbiology* 2015 :1. doi:10.1155/2015/242486.
- Jiang, Y., P. E. Jolly, W. O. Ellis, J. S. Wang, T. D. Phillips, and J. H. Williams. 2005. Aflatoxin B1 albumin adduct levels and cellular immune status in Ghanaians. *International Immunology* 17 (6):807.
- Johnson, N. M., G. Qian, L. Xu, D. Tietze, A. Marroquin-Cardona, A. Robinson, M. Rodriguez, L. Kaufman, K. Cunningham, J. Wittmer, et al. 2010. Aflatoxin and PAH exposure biomarkers in a U.S. population with a high incidence of hepatocellular carcinoma. *Science of the Total Environment* 408 (23):6027. doi:10.1016/j.scitotenv.2010.09.005.
- Jonsyn-Ellis, F. E. 2007. Aflatoxins and ochratoxins in serum samples of school children. *Journal of Nutritional and Environmental Medicine* 16 (1):52–58. doi:10.1080/13590840601016478
- Jørgensen, T. R. 2007. Identification and toxigenic potential of the industrially important fungi, *Aspergillus oryzae* and *Aspergillus sojae*. *Journal of Food Protection* 70 (12):2916–2934.
- Kauffman, C. A. 2007. Histoplasmosis: a clinical and laboratory update. *Clinical Microbiology Reviews* 20 (1):115–132. doi:10.1128/CMR.00027-06.PMC 1797635
- Kensler, T. W., B. D. Roebuck, G. N. Wogan, and J. D. Groopman. 2011. Aflatoxin: A 50-year odyssey of mechanistic and translational toxicology. *Toxicological Science* 120 (1):28–48.
- Kinsella, J. E., D. H. Hwang, and B. Dwivedi. 1976. Enzymes of *Penicillium roqueforti* involved in the biosynthesis of cheese flavor. *Critical Reviews in Food Science and Nutrition* 8 (2):191–228. doi:10.1080/10408397609527222.
- Koeck, M., A. R. Hardham, and P. N. Dodds. 2011. The role of effectors of biotrophic and hemibiotrophic fungi in infection. *Cellular Microbiology* 13 (12):1849–1857. doi:10.1111/j.1462-5822.2011.01665.x.
- Kulp, K. 2000. *Handbook of cereal science and technology*. Boca Raton, FL: CRC Press. ISBN 0-8247-8294-1.
- Lanyasunya, T. P., L. W. Wamae, H. H. Musa, O. Olowofeso, and I. K. Lokwaleput. 2005. The risk of mycotoxins contamination of dairy feed and milk on smallholder dairy farms in Kenya. *Pakistan Journal of Nutrition* 4 :162–169. doi:10.3923/pjn.2005.162.169.
- Lee, N. A., S. Wang, R. D. Allan, and I. R. Kennedy. 2004. A rapid aflatoxin B1 ELISA: Development and validation with reduced matrix effects for peanuts, corn, pistachio, and soybeans. *Journal of Agricultural and Food Chemistry* 52 (10):2746–2755. doi:10.1021/jf0354038.
- Leong, Y. H., A. Rosma, A. A. Latiff, and A. N. Izzah. 2012. Associations of serum aflatoxin B1–lysine adduct level with socio-demographic factors and aflatoxins intake from nuts and related nut products in Malaysia. *International Journal of Hygiene and Environmental Health* 215 (3):368. doi:10.1016/j.ijheh.2011.12.005.
- Lewis, L., M. Onsongo, H. Njapau, H. Schurz-Rogers, G. Lubet, S. Kieszak, J. Nyamongo, L. Backer, A. M. Dahiye, A. Misore, Kenya Aflatoxicosis Investigation Group, et al. 2005. Aflatoxin contamination of commercial maize products during an outbreak of acute aflatoxicosis in Eastern and Central Kenya. *Environmental Health Perspectives* 113 (12):1763–1767. doi:10.1289/ehp.7998.
- Manzoni, M., and M. Rollini. 2002. Biosynthesis and biotechnological production of statins by filamentous fungi and application of these cholesterol-lowering drugs. *Applied Microbiology and Biotechnology* 58 (5):555–564. doi:10.1007/s00253-002-0932-9.
- Malir, F., V. Ostry, M. Dofkova, T. Roubal, V. Dvorak, and V. Dohnal. 2013. Ochratoxin A levels in blood serum of czech women in the first trimester of pregnancy and its correspondence with dietary intake of the mycotoxin contaminant. *Biomarkers: Biochemical Indicators of Exposure, Response, and Susceptibility to Chemicals* 18 (8):673.
- Martlbauer, E., E. Usleber, R. Dietrich, and E. Schneider. 2009. Ochratoxin A in human blood serum—Retrospective long-term data. *Mycotoxin Research* 25 (4):175–86. doi:10.1007/s12550-009-0025-z
- Martins, M. L., H. M. Martins, and F. Bernardo. 2001. Aflatoxins in spices marketed in Portugal. *Food Additives and Contaminants* 18 (4):315–319. doi:10.1080/02652030120041.
- Mazaheri, M. 2009. Determination of aflatoxins in imported rice to Iran. *Food and Chemical Toxicology: An International Journal Published for the British Industrial Biological Research Association* 47 (8):2064–2066.
- Moore, R. T. 1980. Taxonomic proposals for the classification of marine yeasts and other yeast-like fungi including the smuts. *Botanica Marina* 23 :361–373.
- Moss, M. O. 2008. Fungi, quality and safety issues in fresh fruits and vegetables. *Journal of Applied Microbiology* 104(5):1239–1243. doi:10.1111/j.1365-2672.2007.03705.x.
- Nielsen, K., and J. Heitman. 2007. Sex and virulence of human pathogenic fungi. *Advances in Genetics* 57 :143–173. doi:10.1016/S0065-2660(06)57004-X. ISBN 978-0-12-017657-1.
- Munkvold, G. P. 2003. Mycotoxins in corn: occurrence, impacts, and management. In *Corn Chemistry and Technology*, eds. P. White and L. Johnson, 811–881. 2nd Ed. St. Paul, MN: American Association of Cereal Chemists.
- Munoz, K., M. Vega, G. Rios, S. Munoz, and R. Madariaga. 2006. Preliminary study of ochratoxin a in human plasma in agricultural zones of Chile and its relation to food consumption. *Food and Chemical Toxicology : An International Journal Published for the British Industrial Biological Research Association* 44 (11):1884.
- Onions, A. H. S., D. Allsopp, and H. O. W. Egging. 1981. *Smiths introduction to industrial mycology*, 398. UK: Cambridge University Press.
- Ostry, V., F. Malir, T. Roubal, J. Skarkova, J. Ruprich, M. Cerna, and E. E. Creppy. 2005. Monitoring of mycotoxin biomarkers in the Czech Republic. *Mycotoxin Research* 21(1):49.
- Orr, D. B., and R. T. Orr. 1979. *Mushrooms of Western North America*. Berkeley, California: University of California Press. p. 17. ISBN 0-520-03656-5.

- Ozçelik, N., A. Koşar, and D. Soysal. 2001. Ochratoxin a in human serum samples collected in Isparta-Turkey from healthy individuals and individuals suffering from different urinary disorders. *Toxicology Letters* 121 (1):9.
- Pacin, A. M., E. V. Ciancio Bovier, E. Motta, S. L. Resnik, D. Villa, and M. Olsen. 2008. Survey of Argentinean human plasma for ochratoxin A. *Food Additives & Contaminants. Part A, Chemistry, Analysis, Control, Exposure & Risk Assessment* 25 (5):635.
- Palmgren, M. S., and A. W. Hayes. 1987. Aflatoxins in food. In *Mycotoxins in food*, ed. P. Krogh, 65–95. Cambridge, MA: Academic Press.
- Palli, D., M. Miraglia, C. Saieva, G. Masala, E. Cava, M. Colatosti, A. M. Corsi, A. Russo, and C. Brera. 1999. Serum levels of ochratoxin a in healthy adults in tuscany: Correlation with individual characteristics and between repeat measurements. *Cancer Epidemiology, Biomarkers & Prevention: A Publication of the American Association for Cancer Research, Cosponsored by the American Society of Preventive Oncology* 8(3):265.
- Pan, A., S. Lorenzotti, and A. Zoncada. 2008. Registered and investigational drugs for the treatment of methicillin-resistant *Staphylococcus aureus* infection. *Recent Patents on anti-Infective Drug Discovery* 3 (1):10–33. doi:10.2174/157489108783413173.
- Paoletti, M., K. W. Buck, and C. M. Brasier. 2006. Selective acquisition of novel mating type and vegetative incompatibility genes via inter-species gene transfer in the globally invading eukaryote ophiostoma novo-ulmi. *Molecular Ecology* 15 (1):249–262. doi:10.1111/j.1365-294X.2005.02728.x.
- Paszkowski, U. 2006. Mutualism and parasitism: The yin and yang of plant symbioses. *Current Opinion in Plant Biology* 9 (4):364–370. doi:10.1016/j.pbi.2006.05.008.
- Paterson, R. R. M. 2008. Cordyceps: A traditional Chinese medicine and another fungal therapeutic biofactory? *Phytochemistry* 69 (7): 1469–1495. doi:10.1016/j.phytochem.2008.01.027.
- Paterson, R. R. M. 2006. Ganoderma – A therapeutic fungal biofactory. *Phytochemistry* 67 (18):1985–2001. doi:10.1016/j.phytochem.2006.07.004.
- Pereira, V. L., J. O. Fernandes, and S. C. Cunha. 2014. Mycotoxins in cereals and related foodstuffs: A review on occurrence and recent methods of analysis. *Trends in Food Science and Technology* 36(2): 96–136. doi:10.1016/j.tifs.2014.01.005.
- Piekkola, S., P. C. Turner, M. Abdel-Hamid, S. Ezzat, M. El-Daly, S. El-Kafrawy, E. Savchenko, T. Poussa, J. C. S. Woo, H. Mykkänen, and H. El-Nezami. 2012. Characterisation of aflatoxin and deoxynivalenol exposure among pregnant Egyptian women. *Food Additives & Contaminants. Part A, Chemistry, Analysis, Control, Exposure & Risk Assessment* 29 (6):962.
- Pitt, J. L., J. C. Basilio, M. L. Abarca, and C. Lopez. 2000. Mycotoxins and toxigenic fungi. *Medical Mycology* 38 (s1):41–46. doi:10.1080/mmy.38.1.41.46.
- Piskur, J., E. Rozpedowska, S. Polakova, A. Merico, and C. Compagno. 2006. How did *Saccharomyces* evolve to become a good brewer? *Trends in Genetics* 22 (4):183–186. doi:10.1016/j.tig.2006.02.002.
- Probst, C., H. Njapau, and P. J. Cotty. 2007. Outbreak of an acute aflatoxicosis in Kenya in 2004: Identification of the causal agent. *Applied and Environmental Microbiology* 73 (8):2762–2764.
- Rachaputi, N., G. C. Wright, and S. Krosch. 2002. Management practices to minimise pre-harvest aflatoxin contamination in Australian groundnuts. *Australian Journal of Experimental Agriculture* 42 (5): 595–605.
- Radic, B., R. Fuchs, M. Peraica, and A. Lucic. 1997. Ochratoxin A in human sera in the area with endemic nephropathy in Croatia. *Toxicology Letters* 91 :105. doi:10.1016/S0378-4274(97)03877-0.
- Richard, J. L. 2007. Some major mycotoxins and their mycotoxicoses—An overview. *International Journal of Food Microbiology* 119 (1–2): 3–10. doi:10.1016/j.ijfoodmicro.2007.07.019.
- Ritieni, A., A. Santini, M. Mussap, R. Ferracane, P. Bosco, D. Gazzolo, and F. Galvano. 2010. Simultaneous determination of mycotoxins in biological fluids by LC-MS/MS. *Frontiers in Bioscience (Elite Edition)* 2 :151.
- Sangare Tigori, B., S. Moukha, J. H. Kouadio, D. S. Dano, A. M. Betbeder, A. Achour, and E. E. Creppy. 2006. Ochratoxin a in human blood in Abidjan, Côte d'Ivoire. *Toxicon: Official Journal of the International Society on Toxinology* 47 (8):894.
- Schaafsma, A. W., and D. C. Hooker. 2007. Climatic models to predict occurrence of *Fusarium* toxins in wheat and maize. *International Journal of Food Microbiology* 119 (1–2):116–125.
- Schmale, D. G., III, and G. P. Munkvold. 2018. Mycotoxins in crops: A threat to human and domestic animal health. Topics in plant pathology, Mycotoxins: Economic impact. *The American Phytopathological Society*.
- Schleicher, R. L., L. F. McCoy, C. D. Powers, M. R. Sternberg, and C. M. Pfeiffer. 2013. Serum concentrations of an aflatoxin-albumin adduct in the national health and nutrition examination survey (NHANES) 1999–2000. *Clinica Chimica Acta; International Journal of Clinical Chemistry* 423 :46.
- Shephard, G. S. 2008. Determination of mycotoxins in human foods. *Chemical Society Reviews* 37 (11):2468–2477. doi:10.1039/b713084h
- Shephard, G. S. 2008. Impact of mycotoxins on human health in developing countries. *Food Additives & Contaminants. Part A, Chemistry, Analysis, Control, Exposure & Risk Assessment* 25 (2):146–151.
- Shuaib, F. M. B., P. E. Jolly, J. E. Ehiri, Y. Jiang, W. O. Ellis, J. K. Stiles, N. J. Yatich, E. Funkhouser, S. D. Person, C. Wilson, and J. H. Williams. 2010. Association between anemia and aflatoxin B1 biomarker levels among pregnant women in Kumasi, Ghana. *The American Journal of Tropical Medicine and Hygiene* 83 (5):1077.
- Simon-Nobbe, B., U. Denk, V. Pöll, R. Rid, and M. Breitenbach. 2008. The spectrum of fungal allergy. *International Archives of Allergy and Immunology* 145 (1):58–86. doi:10.1159/000107578.
- Struck, C. 2006. Infection strategies of plant parasitic fungi. In *The epidemiology of plant diseases*, eds. B. M. Cooke, D. G. Jones, and B. Kaye, 117. Berlin, Germany: Springer. ISBN 1-4020-4580-8.
- Talbot, N. J. 2003. On the trail of a cereal killer: Exploring the biology of *Magnaporthe grisea*. *Annual Review of Microbiology* 57 (1): 177–202. doi:10.1146/annurev.micro.57.030502.090957.
- Tang, L., L. Xu, E. Afriyie-Gyawu, W. Liu, P. Wang, Y. Tang, Z. Wang, H. J. Huebner, N.-A. Ankrah, D. Ofori-Adjei, et al. 2009. Aflatoxin-albumin adducts and correlation with decreased serum levels of vitamins a and E in an adult Ghanaian population. *Food Additives and Contaminants* 26 (1):108. doi:10.1080/02652030802308472.
- Trucksess, M. W., and P. M. Scott. 2008. Mycotoxins in botanicals and dried fruits: A review. *Food Additives & Contaminants: Part A* 25 :181–192. doi:10.1080/02652030701567459.
- Turner, P. C., Loffredo, C. Kafrawy, S. E. Ezzat, S. Eissa, L. S. A. Daly, M. E. Nada, O. Abdel-Hamid, and M. 2008. Pilot survey of aflatoxin-albumin adducts in sera from Egypt. *Food Additives & Contaminants. Part A, Chemistry, Analysis, Control, Exposure & Risk Assessment* 25 (5):583.
- Turner, P. C., A. C. Collinson, Y. B. Cheung, Y. Gong, A. J. Hall, A. M. Prentice, and C. P. Wild. 2007. Aflatoxin exposure in utero causes growth faltering in Gambian infants. *International Journal of Epidemiology* 36 (5):1119.
- Turner, P. C., A. Sylla, S. Y. Kuang, C. L. Marchant, M. S. Diallo, A. J. Hall, J. D. Groopman, and C. P. Wild. 2005. Absence of TP53 codon 249 mutations in young Guinean children with high aflatoxin exposure. *Cancer Epidemiology, Biomarkers & Prevention: A Publication of the American Association for Cancer Research, Cosponsored by the American Society of Preventive Oncology* 14(8):2053.
- Turner, P. C., K. H. Dingley, J. Coxhead, S. Russell, and C. R. Garner. 1998. Detectable levels of serum aflatoxin B1-albumin adducts in the United Kingdom population: implications for aflatoxin-B1 exposure in the United Kingdom. *Cancer Epidemiology, Biomarkers & Prevention: A Publication of the American Association for Cancer Research, Cosponsored by the American Society of Preventive Oncology* 7 (5):441.
- Ueno, Y., S. Maki, J. Lin, M. Furuya, Y. Sugiura, and O. Kawamura. 1998. A 4-year study of plasma ochratoxin a in a selected population in Tokyo by immunoassay and immunoaffinity column-linked HPLC. *Food and Chemical Toxicology: An International Journal Published for the British Industrial Biological Research Association* 36 (5):445.



- Waseem, A., S. A. Shah, A. Sajjad, A. R. Siddiqi, and M. Nafees. 2014. Human exposure to mycotoxins: A retrospective review of leading toxins and metabolites in human biological matrices. *Journal of The Chemical Society of Pakistan* 36 (6): 1196–1214.
- Wild, C. P., F. Yin, P. C. Turner, I. Chemin, B. Chapot, M. Mendy, H. Whittle, G. D. Kirk, and A. J. Hall. 2000. Environmental and genetic determinants of aflatoxin-albumin adducts in the Gambia. *International Journal of Cancer* 86 (1):1.
- Yang, Y., E. Yang, Z. An, and X. Liu. 2007. Evolution of nematode-trapping cells of predatory fungi of the orbiaceae based on evidence from rRNA-encoding DNA and multiprotein sequences. *Proceedings of the National Academy of Sciences of the United States of America* 104 (20):8379–8384. Bibcode:2007PNAS.104.8379Y. doi:10.1073/pnas.0702770104.
- Yard, E. E., J. H. Daniel, L. S. Lewis, M. E. Rybak, E. M. Paliakov, A. A. Kim, J. M. Montgomery, R. Bunnell, M. U. Abudo, W. Akhwale, et al. 2013. Human aflatoxin exposure in Kenya, 2007: A cross-sectional study. *Food Additives & Contaminants. Part A, Chemistry, Analysis, Control, Exposure & Risk Assessment* 30 (7):1322.
- Yin, Y. N., L. Y. Yan, J. H. Jiang, and Z. H. Ma. 2008. Biological control of aflatoxin contamination of crops. *Journal of Zhejiang University Science B* 9 (10):787–792. doi:10.1631/jzus.B0860003.
- Yuen, J. W., and M. D. Gohel. 2005. Anticancer effects of *Ganoderma lucidum*: A review of scientific evidence. *Nutrition and Cancer* 53 (1):11–17. doi:10.1207/s15327914nc5301\_2.
- Zain, E. M. 2011. Impact of mycotoxins on humans and animals. *Journal of Saudi Chemical Society* 15 (2):129–144. doi:10.1016/j.jscs.2010.06.006.