



Critical Reviews in Food Science and Nutrition

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/bfsn20>

Probiotic Lactic Acid Bacteria and Skin Health

Ji Hye Jeong^a, Chang Y. Lee^d & Dae Kyun Chung^{abcd}

^a Skin Biotechnology Center, Kyung Hee University, Yongin 449-701, South Korea

^b School of Biotechnology and Institute of Life Science and Resources, Kyung Hee University, Yongin 449-701, South Korea

^c RNA Inc., #308 College of Life Science, Kyung Hee University, Yongin 449-701, South Korea

^d Department of Food Science, Cornell University, Ithaca, NY 14850

Accepted author version posted online: 19 Aug 2015.



[Click for updates](#)

To cite this article: Ji Hye Jeong, Chang Y. Lee & Dae Kyun Chung (2015): Probiotic Lactic Acid Bacteria and Skin Health, Critical Reviews in Food Science and Nutrition, DOI: [10.1080/10408398.2013.834874](https://doi.org/10.1080/10408398.2013.834874)

To link to this article: <http://dx.doi.org/10.1080/10408398.2013.834874>

Disclaimer: This is a version of an unedited manuscript that has been accepted for publication. As a service to authors and researchers we are providing this version of the accepted manuscript (AM). Copyediting, typesetting, and review of the resulting proof will be undertaken on this manuscript before final publication of the Version of Record (VoR). During production and pre-press, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal relate to this version also.

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

Title page

Journal: Critical Reviews in Food Science and Nutrition

ISSN: 1040-8398

The paper title: Probiotic Lactic Acid Bacteria and Skin Health

Authors and address:

Ji Hye Jeong¹, Chang Y. Lee⁴ and Dae Kyun Chung^{1, 2, 3, 4}

¹ Skin Biotechnology Center, Kyung Hee University, Yongin 449-701, South Korea

² School of Biotechnology and Institute of Life Science and Resources, Kyung Hee University,
Yongin 449-701, South Korea

³RNA Inc., #308 College of Life Science, Kyung Hee University, Yongin 449-701, South Korea

⁴ Department of Food Science, Cornell University, Ithaca, NY 14850

Corresponding author:

Dae Kyun Chung,

E-mail address: dkchung@khu.ac.kr

Address: Skin Biotechnology Center, Kyung Hee University, Yongin 449-701, South Korea.

Probiotic Lactic Acid Bacteria and Skin Health**Abstract**

Human skin is the first defense barrier against the external environment, especially microbial pathogens and physical stimulation. Many studies on skin health with Lactic acid bacteria (LAB) have been published for many years, including prevention of skin disease and improvement of skin conditions. LAB, a major group of gram-positive bacteria, are known to be beneficial to human health by acting as probiotics. Recent studies have shown that LAB and their extracts have beneficial effects on maintenance and improvement of skin health. Oral administration of *Lactobacillus delbrueckii* inhibits the development of atopic disease. In addition, LAB and LAB extracts are known to have beneficial effects on intestinal diseases, with *Lactobacillus plantarum* having been shown to attenuate IL-10 deficient colitis. In addition to intestinal health, *L. plantarum* also has beneficial effects on skin. pLTA, which is lipoteichoic acid isolated from *L. plantarum*, has anti-photoaging effects on human skin cells by regulating the expression matrix metalloproteinase-1 (MMP-1) expression. While several studies have proposed a relationship between diseases of the skin and small intestines, there are currently no published reviews of the effects of LAB for skin health through regulation of intestinal conditions and the immune system. In this review, we discuss recent findings on the effects of LAB on skin health and its potential applications in beauty foods.

Keywords Lactic acid bacteria, Probiotic, Skin, Beauty food

Introduction

Probiotics refer to microorganisms that have beneficial effects on host health. Interestingly, the most common focus of probiotics is modulation of immune system potency [1]. Lactic acid bacteria (LAB) have a number of beneficial effects and have formed the basis of the term probiotics. More specifically, *Lactobacilli* and *Bifidobacteria* species, the most common strains of LAB, are known to have the ability to regulate Th1:Th2 cytokine balance. In addition, many studies have shown that probiotics have beneficial effects on physical barrier function, mucosal immune system, metabolism, and systemic immunity [2, 3]. Further, probiotics have been used to successfully treat various diseases including atopic diseases, inflammatory bowel disease (IBD), irritable bowel syndrome, cardiovascular diseases, and pathogen infections [4-8].

Skin is an effective defense barrier between organisms and the external environment, playing an important role in preventing invasion of pathogens and detecting physical stimulation. Skin acts as protector and immune-surveillance mechanism, and is an extensively exposed organ that functions as a unique stimulator for immune sentinel and effector cells. Keratinocytes, the major cell type of the epidermis, can sense harmful pathogens and distinguish them from harmless commensal organisms. In addition, following exposure to intensive UV irradiation, keratinocytes are activated and induce an immune response by secreting key pro-inflammatory cytokines [9]. As a physical barrier, skin barrier function is a central event in various skin diseases. Specifically, impaired skin barrier function can lead to pathogen infection and atopic dermatitis (AD) [10, 11]. Several studies have shown that skin health is associated with the nutritional condition of the host. Interestingly, the probiotic *L. paracasei* was shown to decrease skin sensitivity and increase

the recovery rate of skin barrier function [12]. Thus, the purpose of this review is to provide an overview of the recent findings on the effects of LAB on skin health and its potential applications in beauty foods.

I . Probiotics and skin

The skin is a complex network of interactions between microbes and epidermis cells. In healthy skin, probiotics can be considered as innocuous or even beneficial. However, some skin diseases are caused by an imbalance of microorganisms, including dry skin and mild AD [13]. A number of recent studies have looked at the effects of probiotics and maintenance of healthy skin conditions. In one study, oral bacteriotherapy with probiotics was suggested to be useful for treating certain skin diseases. Likewise, a combination of *L. rhamnosus* 19070-2 and *L. reuteri* DSM 122460 was shown to be an effective therapy for AD [14], while the administration of *B. subtilis* was shown to prevent atopic lesions induced by mite antigen in NC mice [15].

Probiotics can also help to maintain skin homeostasis by protecting against UV-induced skin damage. Oral supplementation with the *L. johnsonii* La1 has a beneficial effect on cutaneous immune status after acute UV irradiation [16, 17]. Further, oral administration of *Lactobacillus paracasei* improves skin barrier function and prevents local skin inflammation by modulating the immune system in a disease-dependent manner [18]. Interestingly, in addition to live probiotics, heat-killed probiotics can effectively modulate the immune system. Heat-killed R-037 and *Lactobacillus delbrueckii* subsp. *lacti* have an inhibitory effect on the development of AD [19]. Thus, it is clear that there are numerous strains of probiotics that have diverse effects on skin health. The relationships between probiotics and their effects on the skin are summarized in

Table 1.

The majority of probiotics in the human body are found in the stomach and small intestines. Indeed, probiotic LAB are closely associated with beneficial effects on gastroenteritis and diarrhea. It has been suggested that probiotics systemically modulate immune responses in the gut, and other studies have considered the possibility that probiotics can have similar effects in human skin [20].

II. Strategies to Improve Skin Health using Probiotics

1. Modulation of immune responses by probiotics

Lactobacilli, a major group of probiotics, can modulate the host immune response by binding to pattern recognition receptors (PRRs) expressed on a number of immune cells and other tissues including the intestinal epithelium. Probiotics, as microbe-associated molecular patterns (MAMPs), are recognized by PRRs, which in turn induce the production of cytokines and chemokines. PRR signaling receptors are divided into three families, namely, Toll-like receptors (TLRs), retinoic acid inducible gene-I (RIG-I)-like receptors, and nucleotide oligomerization domain-like (NOD) receptors (NLRs). Especially, TLR signaling is a major signaling pathway after MAMP binding and is activated according to the specific arrangement and sequence variations of extracellular leucine-rich repeat (LRR) domains. Dimerization of TLRs leads to the recruitment of adaptor molecules such as myeloid differentiation primary response gene 88 (MyD88). The TLR signaling pathway, through recruitment of MyD88, can subsequently activate the mitogen-activated protein kinase (MAPK) pathway and nuclear factor κ B (NF- κ B).

signaling cascade. Thus, TLR signaling is important for modulation of immune responses by probiotics [21].

LAB can induce different amounts and types of TLR signaling associated cytokines, namely, TNF- α , IFN- γ , IL-1 β , IL-4, IL-6, IL-10, and IL-12. Further, LAB-ligation of TLRs induces translocation of NF- κ B to the nucleus, resulting in production of proinflammatory cytokines. Similarly, *L. acidophilus* induces the release of high amounts of IFN- γ from natural killer (NK) cells and production of IL-12 in dendritic cells (DCs). However, not all species of LAB have a similar capacity for induction of IL-12 and IFN- γ production. For example, *B. bifisum* and *L. reuteri* inhibit *L. acidophilus*-induced secretion of IL-12 and IFN- γ secretion [22]. In addition, LABs are well known for their ability to modulate immune responses by altering the Th1/Th2 cytokine balance. Probiotic LAB modulates both protective Th1 type cytokines, such as IFN- γ , and immunosuppressive Th2 type cytokines, including IL-4. Both *L. rhammosus* GG and *Bifidobacterium bifibum* MF20/5 improve Th1/Th2 polarization by decreasing IL-4 secretion in both unstimulated and *Mycobacterium tuberculosis* antigen-stimulated peripheral blood mononuclear cells (PBMCs) [23]. Likewise, lipoteichoic acid (LTA) isolated from *L. plantarum* has been shown to significantly inhibit LPS-induced TNF- α production [24]. Similarly, fermentation by *Lactobacillus* enhances inhibition of LPS-induced NO, PGE₂, TNF- α and IL-6 production by abrogating the activities of NF- κ B and MAPK [25]. In addition, pre-treatment with *Lactobacillus acidophilus* has an anti-inflammation effect, reducing IL-8 and TNF- α secretion by inactivating the NF- κ B pathway [26]. *Lactobacillus paracasei* KW3110 induces IL-12 expression and inhibits IgE secretion [27], while *L. rhamnosus* GR-1 increases levels of TNF and pro-inflammatory cytokine and decreases secretion of IL-6 and CXCL8. Lastly, the NF- κ B

signaling pathway is also activated by the probiotic *L. rhamnosus* GG [28].

In summary, the immune modulatory effects of probiotics are coincident with their ability to stimulate PRR-associated cell signaling by acting as MAMPs. Interestingly, immune responses by probiotics are strain-specific, which has not been discussed in previous reviews.

2. Beneficial effects of probiotics for maintaining good skin condition

Recently, many researchers have studied the relationship between probiotics and skin health. The consensus of these studies is that probiotics have beneficial effects for improving skin health. Probiotics can improve skin conditions through two processes, either by acting as microflora on the skin or as probiotics in the intestines [29, 30]. In particular, resistant microflora of the skin, mainly Gram-positive bacteria, are regarded as “beneficial” to the “normal health” of the host. Indeed, such microflora can influence skin barrier function and the immune system [29]. Probiotics in the intestines can also have beneficial effects on skin health and can be used as a treatment by administration to the host. For example, *Bifidobacterium infantis* prevents skin disorders-associated immune abnormalities, while *Lactobacillus reuteri* protects epidermal keratinocytes from *staphylococcus*. Further, other LAB can act as probiotics by alleviating infantile eczema [30-32].

Blocking the release of pro-inflammatory cytokine is an important strategy for the treatment of skin inflammation [33]. Inhibition of pro-inflammatory cytokines is a well-known property of probiotics, especially LAB. Specifically, LAB including *L. plantarum* and *Lactobacillus acidophilus* strongly inhibit the potential activity of inflammatory mediators, inflammation associated cytokines, and related signaling pathways [22-26]. *L. paracasei* accelerates the

recovery of skin barrier function and inhibits P-induced skin inflammation by modulating reactive skin-associated inflammatory mechanisms. Specifically, *L. paracasei* was shown to abrogate the release of substance P-induced TNF- α expression [34]. In addition, probiotics have beneficial effects on the alleviation of AD lesions. Heat-killed *L. brevis* SBC8803 has been shown to have an inhibitory effect on development of AD by preventing increased IgE production [35]. Probiotic-mediated improvement of AD is also associated with Th1 and Th2 cytokines, and adults with AD treated with the probiotic *Lactobacillus salivarius* exhibit a statistically significant improvement of lesions that is associated with a significant reduction of Th1 cytokines (IL-12+IFN γ) and the Th1/Th2 ratio (IL-12+IFN γ /IL-4+IL-5) [36]. Another cytokine, IL-4, has been strongly implicated in skin diseases such as AD. The pathological features of AD are dependent on IL-4, in that they are a consequence of IL-4 overexpression in the skin. Importantly, several probiotics have been shown to decrease levels of IL-4 production [23, 37].

Keratinocytes are first responders to the external environment and serve as initiators in skin innate immune systems by modulating expression of cytokine/chemokines including IL-1 β , TNF- α , IL-8, and granulocyte/macrophage colony-stimulating factor (GM-CSF) in the skin [38-40]. Indeed, the immune initiation potency of keratinocytes is facilitated through TLRs such as TLR1, TLR2, TLR3, TLR4, TLR5, TLR6 and TLR 10. Notably, TLR2, a major recognition receptor of LPS, is considered to be a key receptor in epithelial cells. Further, one research group has reported that TLR2 is the only such receptor expressed on the surface of cultured normal human keratinocytes [41, 42]. Cytokine/chemokines production via TLRs results in protection of the host from bacteria pathogens and fungi by stimulating skin inflammation [43].

Indeed, LAB are known to function as immune modulators, with this property having been achieved through regulation of pro-inflammatory cytokine production through TLR signaling. Further, many studies have shown that LAB have a beneficial effect on skin inflammation, with some studies suggesting that T cells are important for mediating the effects of probiotics on skin. *L. casei* has been shown to alleviate allergic skin inflammation by inhibiting both effector and regulatory T cells, and treatment with *L. casei* reduces recruitment of CD8⁺ effector T cells into the skin, enhances the frequency of FoxP3⁺ Tregs, and increases production of IL-10 by CD4⁺CD25⁺ regulatory T cells in the skin [44].

While recent research has demonstrated a strong relationship between skin health and probiotics, the mechanisms by which probiotics facilitate improved skin health remains unclear. Here, we propose a mechanism to explain how skin health is improved by probiotics (Fig. 1). As described above, skin function is closely tied with inflammation. Modulation of skin immunity and inflammation responses are associated with common cell signaling, NF- κ B. Here between the skin immune and inflammatory responses in the presence of a common signal transduction factors to the NF- κ B signal [45]. Aspects of the skin immune, ROS (reactive oxygen species) is expressed by certain internal or external stress in skin keratinocytes. This ROS activate the cellular signaling through regulate inflammatory signal, NF- κ B [46]. Then activated NF- κ B enters the nucleus and induces expression of MMPs, factor that affect skin function through degradation of extracellular matrix proteins. In a perspective of inflammation reaction, NF- κ B influences the expression of inflammatory cytokines such as TNF- α , induce MMP expression. It means specific factor associated with skin is regulated through the NF- κ B signal [45, 47, 48]. In the previous studies, probiotic is well known the expression of inflammatory cytokines via the

NF- κ B signal [25]. Thus, we suggest that probiotics can affect skin and intestine health through regulation of inflammatory signals such as NF- κ B.

3. Skin health reflects the effect of probiotics on intestinal health

Until now, research groups have studied the beneficial effects of probiotics by focusing on their ability to maintain intestinal health and prevent diseases of the gut [49]. However, the current focus of probiotic studies has moved from looking only at intestinal health to studying relationships between the internal immune system and other immunological tissues in the host, such as intestine-skin and intestine-brain interactions. Importantly, the notion that intestinal microflora and inflammatory skin conditions are physiologically intertwined is not a new one. Stokes-Pillsbury previously suggested in 1930 that the skin is influenced by emotional states and microbial flora, reporting that gut microflora can influence the degree of acne by a systemic effect on inflammation as well as by controlling oxidative stress through direct and indirect mechanisms [50]. In addition, Fry defined the association between the small intestines and skin diseases to three subgroups, namely, i) a non-specific relationship, ii) a specific relationship, and iii) a generalized disease process [51].

As previously mentioned, probiotics enhance skin conditions by improving gut health (Fig. 1). The majority of the beneficial effects of probiotics on skin via intestines are facilitated through systemic immune responses [3, 20, 50]. In particular, probiotics regulate specific T-cells such as Th1 cells and modulate activity of Th1-type cytokines in the gut mucosa, which may influence immune responses in other tissues for which they play an important role [34-37]. Indeed, orally administered *Lactobacillus casei* has been shown to alleviate skin inflammation by reducing the

number of CD8⁺ effector cells such that only CD4⁺ T-cells are present [52]. The clinically relevant improvement of acne following administration of a mixture of *L. acidophilus* and *L. bulgaricus* provides further clinical evidence of the skin-gut connection of probiotics [53]. In addition, probiotics have been shown to attenuate release of biologically active peptides, including substance P release in both the intestinal tract and skin [32, 54]. Interestingly, the influence of probiotics through the inhibition of substance P production has been suggested as a mechanism of communication between the gut and skin [55]. Another study reported amelioration of skin homeostasis by oral administration of probiotics following stress-induced skin inflammation. Together, these observations strongly support the existence of a probiotic link in the gut-brain-skin axis [58]. This concept is supported by a number of studies, including identification of peptide-containing cells in skin [59]. In addition, the association of inflammatory bowel diseases and skin diseases provides further evidence to strengthen the relationship between the gut and skin [58, 59].

III. Probiotic extracts, including LTA, and skin health

Previous studies have suggested that live probiotics are the most effective for modulating the immune response of the host [60-62]. However, extracts of probiotics have been demonstrated to have similar or even more effective activity for stimulating and modulating immune response compared to live probiotics. Specifically, *Lactobacillus casei* extract is a potent inducer of NF- κ B through the TLR-2/MyD88 signaling pathway [63]. Likewise, heat-killed *Lactobacillus casei* generate an enhanced adaptive immune response when administered nasally with a pneumococcal antigen or live bacteria [64]. A clinical trial of a probiotic lysate of

Bifidobacterium longum sp. used a non-replicating bacteria extract applied to the skin, rather than live probiotics, and was shown to improve sensitive skin. Following this study, a test cream containing a 10% bacterial extract was shown to have a clinically relevant effect on skin dryness, and was shown to increase skin resistance against physical and chemical aggression. Together, these results suggest that specific bacteria extracts have beneficial effect on reactive skin [65]. In addition, many published studies have demonstrated that specific bacterial extracts, including those of *Lactobacillus johnsonii*, *Lactobacillus casei*, *Lactobacillus plantarum*, and *Lactobacillus acidophilus* have beneficial effects on anti-adhesion and anti-microbial potency on cutaneous and mucous surfaces, respectively [66, 67].

As mentioned above, regulation of TLR signaling pathways and skin immune responses are important to protect skin from immune associated skin disease, such as AD [41, 42, 68]. Specific elements of probiotics modulate immune responses triggered by TLR ligands. In particular, lipoteichoic acid (LTA), a major component of the Gram-positive bacteria cell wall, is important for modulating the activity of the innate immune response [69]. LTA is well known for its potent immune modulatory activity. For example, pLTA (isolated from *Lactobacillus plantarum*) has been shown to have an inhibitory effect on LPS-induced TNF- α production [24]. Research on the inhibition of TLR3-dependent inflammation by probiotic bacteria through a TLR2-dependent mechanism in skin injury suggests that LTA may be effective for treating skin inflammation [70]. Further, several research groups have consistently demonstrated the relationship between LTA and skin health by showing that purified LTA from *Lactobacillus rhamnosus* GG has an immune modulatory effect against UV radiation and inhibition of skin tumor development [71]. In addition, our research group has shown that pLTA has an anti-photoaging effect on UV-

irradiated human skin through regulation of MMP-1 expression (not published). Thus, we propose that LTA, which is a component of many probiotic extracts, is an important element of probiotics for maintenance of skin health.

IV. Conclusion

The international cosmetic market for beauty foods is expanding, and similarly the interest for probiotic LAB is growing very fast. LAB is a remarkable group of probiotics that have anti-inflammatory properties and are capable of improving skin health. Although the precise mechanisms by which LAB improve skin health is unknown yet, its beneficial effects on skin health have been demonstrated by numerous studies, and the literature continues to grow. LAB have been shown to modulate the expression of MMPs and their extracts, rather than live whole LAB, can have beneficial effects on immune modulation, including skin health. Probiotics directly influence human skin. In addition, modulation of immunity in the gut may be relevant for maintaining the health of human skin through a systemic mechanism. Recent years have witnessed a proliferation of published clinical studies involving probiotic foods and beverages. At the same time, the global market for nutricosmetics is growing rapidly due to changes in lifestyle, aging population and increased consumer awareness. Therefore, academic and industry interests on probiotic lactic acid bacteria on skin health are expected to grow rapidly.

Acknowledgements

This research was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (No. 2010-0012091)

References

- [1] West, N.P., Pyne, D.B., Peake, J.M., and Cripps, A.W. (2009). Probiotics, immunity and exercise: a review. *Exerc Immunol Rev.* **15**: 107-26.
- [2] Salminen, S.J., Gueimonde, M., and Isolauri, E. (2005). Probiotics that modify disease risk. *J Nutr.* **135**(5): 1294-1298.
- [3] Wallace, T.C., Guarner, F., Madsen, K., Cabana, M.D., Gibson, G., Hentges, E., and Sanders, M.E. (2011). Human gut microbiota and its relationship to health and disease. *Nutr Rev.* **69**(7): 392-403.
- [4] Whorwell, P.J. (2009). Do probiotics improve symptoms in patients with irritable bowel syndrome? *Therap Adv Gastroenterol.* **2**(4): 37-44.
- [5] Parvez, S., Malik, K.A., Ah Kang, S., and Kim, H.Y. (2006). Probiotics and their fermented food products are beneficial for health. *J Appl Microbiol.* **100**(6): 1171-1185.
- [6] Dominguez-Bello, M.G., and Blaser, M.J. (2008). Do you have a probiotic in your future? *Microbes Infect.* **10**(9): 1072-1076.
- [7] Ouwehand, A.C., Salminen, S., and Isolauri, E. (2002). Probiotics: an overview of beneficial effects. *Antonie Van Leeuwenhoek.* **82**(1-4): 279-289.
- [8]. Habil, N., Al-Murrani, W., Beal, J., and Foey, A.D. (2011). Probiotic bacterial strains differentially modulate macrophage cytokine production in a strain-dependent and cell subset-specific manner. *Benef Microbes.* **2**(4): 283-293.
- [9] Nestle, F.O., Di Meglio, P., Qin, J.Z., and Nickoloff, B.J. (2009). Skin immune sentinels in health and disease. *Nat Rev Immunol.* **9**(10): 679-691
- [10] Proksch, E., Brandner, J.M., and Jensen, J.M. (2008). The skin: an indispensable barrier. *Exp Dermatol.* **17**(12): 1063-1072.

- [11] Jensen, J.M., and Proksch, E. (2009). The skin's barrier. *G Ital Dermatol Venereol.* **144**(6): 689-700.
- [12] Piccardi, N., and Manissier, P. (2009). Nutrition and nutritional supplementation: Impact on skin health and beauty. *Dermatoendocrinol.* **1**(5): 271-274.
- [13] Simmering, R., Breves, R. (2009). Pre- and Probiotic cosmetics. *Hautarzt.* **60**(10): 809-814.
- [14] Rosenfeldt, V., Benfeldt, E., Nielsen, S.D., Michaelsen, K.F., Jeppesen, D.L., Valerius, N.H., and Paerregaard, A. (2003). Effect of probiotic *Lactobacillus* strains in children with atopic dermatitis. *J Allergy Clin Immunol.* **111**(2): 389-395.
- [15] Goto, K., Iwasawa, D., Kamimura, Y., Yasuda, M., Matsumura, M., and Shimada, T. (2011). Clinical and histopathological evaluation of *Dermatophagoides farinae*-induced dermatitis in NC/Nga mice orally administered *Bacillus subtilis*. *J Vet Med Sci.* **73**(5): 649-654.
- [16] Guéniche, A., Benyacoub, J., Buetler, T.M., Smola, H., and Blum, S. (2006). Supplementation with oral probiotic bacteria maintains cutaneous immune homeostasis after UV exposure. *Eur J Dermatol.* **16**(5): 511-517.
- [17] Guéniche, A., Philippe, D., Bastien, P., Blum, S., Buyukpamukcu, E., and Castiel-Higounenc, I. (2009). Probiotics for photoprotection. *Dermatoendocrinol.* **1**(5): 275-279.
- [18] Philippe, D., Blum, S., and Benyacoub, J. (2011). Oral *Lactobacillus paracasei* improves skin barrier function recovery and reduces local skin inflammation. *Eur J Dermatol.* **21**(2): 279-280.
- [19] Watanabe, T., Hamada, K., Tategaki, A., Kishida, H., Tanaka, H., Kitano, M., and Miyamoto, T. (2009). Oral administration of lactic acid bacteria isolated from traditional South Asian fermented milk 'dahi' inhibits the development of atopic dermatitis in NC/Nga mice. *J Nutr Sci Vitaminol (Tokyo).* **55**(3): 271-278.

- [20] Krutmann, J. (2012). Pre- and probiotics for human skin. *Clin Plast Surg.* **39**(1): 59-64.
- [21] Wells, J.M. (2011). Immunomodulatory mechanisms of lactobacilli. *Microb Cell Fact.* **10** Suppl 1: S17.
- [22] Rizzello, V., Bonaccorsi, I., Dongarrà, M.L., Fink, L.N., and Ferlazzo, G. (2011). Role of Natural Killer and Dendritic Cell Crosstalk in Immunomodulation by Commensal Bacteria Probiotics. *J Biomed Biotechnol.* **2011**: 473097
- [23] Ghadimi, D., de Vrese, M., Heller, K.J., and Schrezenmeir, J. (2010). Lactic acid bacteria enhance autophagic ability of mononuclear phagocytes by increasing Th1 autophagy-promoting cytokine (IFN-gamma) and nitric oxide (NO) levels and reducing Th2 autophagy-restraining cytokines (IL-4 and IL-13) in response to Mycobacterium tuberculosis antigen. *Int Immunopharmacol.* **10**(6): 694-706.
- [24] Kim, H.G., Kim, N.R., Gim, M.G., Lee, J.M., Lee, S.Y., Ko, M.Y., Kim, J.Y., Han, S.H., and Chung, D.K. (2008). Lipoteichoic acid isolated from *Lactobacillus plantarum* inhibits lipopolysaccharide-induced TNF-alpha production in THP-1 cells and endotoxin shock in mice. *J Immunol.* **180**(4): 2553-2561.
- [25] Oh, Y.C., Cho, W.K., Oh, J.H., Im, G.Y., Jeong, Y.H., Yang, M.C., and Ma, J.Y. (2012). Fermentation by Lactobacillus enhances anti-inflammatory effect of Oyaksungisan on LPS-stimulated RAW 264.7 mouse macrophage cells. *BMC Complement Altern Med.* **12**: 17.
- [26] Yang, Y.J., Chuang, C.C., Yang, H.B., Lu, C.C., and Sheu, B.S. (2012). Lactobacillus acidophilus ameliorates H. pylori-induced gastric inflammation by inactivating the Smad7 and NFkB pathways. *BMC Microbiol.* **12**: 38.
- [27] Ichikawa, S., Miyake, M., Fujii, R., and Konishi, Y. (2009). Orally administered

- Lactobacillus paracasei* KW3110 induces in vivo IL-12 production. *Biosci Biotechnol Biochem.* **73**(7): 1561-1565.
- [28] Karlsson, M., Scherbak, N., Reid, G., and Jass, J. (2012). *Lactobacillus rhamnosus* GR-1 enhances NF-kappaB activation in Escherichia coli-stimulated urinary bladder cells through TLR4. *BMC Microbiol.* **12**: 15.
- [29] Leyden, J.J., McGinley, K.J., Nordstrom, K.M., and Webster, G.F. (1987). Skin microflora. *J Invest Dermatol.* **88**: 65s-72s.
- [30] Prince, T., McBain, A.J., and O'Neill, C.A. (2012). *Lactobacillus reuteri* Protects Epidermal Keratinocytes from *Staphylococcus aureus* Induced Cell Death by Competitive Exclusion. *Appl Environ Microbiol.* **78**(15): 5119-5126.
- [31] Cinque, B., Di Marzio, L., Della Riccia, D.N., Bizzini, F., Giuliani, M., Fanini, D., De Simone, C., and Cifone, M.G. (2006). Effect of Bifidobacterium infantis on Interferon-gamma- induced keratinocyte apoptosis: a potential therapeutic approach to skin immune abnormalities. *Int J Immunopathol Pharmacol.* **19**(4): 775-786.
- [32] Zhu, D.L., Yang, W.X., and Yang, H.M. (2010). Meta analysis of lactic acid bacteria as probiotics for the primary prevention of infantile eczema. *Zhongguo Dang Dai Er Ke Za Zhi.* **12**(9): 734-739.
- [33] Oh, J., Yu, T., Choi, S.J., Yang, Y., Baek, H.S., An, S.A., Kwon, L.K., Kim, J., Rho, H.S., Shin, S.S., Choi, W.S., Hong, S., and Cho, J.Y. (2012). Syk/Src pathway-targeted inhibition of skin inflammatory responses by carnosic Acid. *Mediators Inflamm.* **2012**: 781375.
- [34] Gueniche, A., Benyacoub, J., Philippe, D., Bastien, P., Kusy, N., Breton, L., Blum, S., and Castiel-Higounenc, I. (2010). *Lactobacillus paracasei* CNCM I-2116 (ST11) inhibits

substance P-induced skin inflammation and accelerates skin barrier function recovery in vitro. *Eur J Dermatol.* **20**(6): 731-737.

- [35] Segawa, S., Hayashi, A., Nakakita, Y., Kaneda, H., Watari, J., and Yasui, H. (2008). Oral administration of heat-killed *Lactobacillus brevis* SBC8803 ameliorates the development of dermatitis and inhibits immunoglobulin E production in atopic dermatitis model NC/Nga mice. *Biol Pharm Bull.* **31**(5): 884-889.
- [36] Prescott, S.L., Dunstan, J.A., Hale, J., Breckler, L., Lehmann, H., Weston, S., and Richmond, P. (2005). Clinical effects of probiotics are associated with increased interferon-gamma responses in very young children with atopic dermatitis. *Clin Exp Allergy.* **35**(12): 1557-1564.
- [37] Elbe-Bürger, A., Egyed, A., Olt, S., Klubal, R., Mann, U., Rappersberger, K., Rot, A., and Stingl, G. (2002). Overexpression of IL-4 alters the homeostasis in the skin. *J Invest Dermatol.* **118**(5): 767-778.
- [38] Sugita, K., Kabashima, K., Atarashi, K., Shimauchi, T., Kobayashi, M., and Tokura, Y. (2007). Innate immunity mediated by epidermal keratinocytes promotes acquired immunity involving Langerhans cells and T cells in the skin. *Clin Exp Immunol.* **147**: 176-183.
- [39] Dorschner, R.A., Pestonjamasp, V.K., Tamakuwala, S., Ohtake, T., Rudisill, J., Nizet, V., Agerberth, B., Gudmundsson, G.H., and Gallo, R.L. (2001). Cutaneous injury induces the release of cathelicidin anti-microbial peptides active against group A Streptococcus. *J Invest Dermatol.* **117**: 91-97.
- [40] Ali, R.S., Falconer, A., Ikram, M., Bissett, C.E., Cerio, R., and Quinn, A.G. (2001). Expression of the peptide antibiotics human beta defensin-1 and human beta defensin-2 in normal human skin. *J Invest Dermatol.* **117**: 106-111.

- [41] Kollisch, G., Kalali, B.N., Voelcker, V., Wallich, R., Behrendt, H., Ring, J., Bauer, S., Jakob, T., Mempel, M., and Ollert, M. (2005). Various members of the toll-like receptor family contribute to the innate immune response of human epidermal keratinocytes. *Immunology* **114**: 531-541.
- [42] Kawai, K., Shimura, H., Minagawa, M., Ito, A., Tomiyama, K., and Ito, M. (2002). Expression of functional toll-like receptor 2 on human epidermal keratinocytes. *J Dermatol Sci.* **30**: 185–194.
- [43] Kobayashi, M., Yoshiki, R., Sakabe, J., Kabashima, K., Nakamura, M., and Tokura, Y. (2009). Expression of toll-like receptor 2, NOD2 and dectin-1 and stimulatory effects of their ligands and histamine in normal human keratinocytes. *Br J Dermatol.* **160**(2): 297-304.
- [44] Hacini-Rachinel, F., Gheit, H., Le Luduec, J.B., Dif, F., Nancey, S., and Kaiserlian, D. (2009). Oral probiotic control skin inflammation by acting on both effector and regulatory T cells. *PLoS One.* **4**(3): e4903.
- [45] Kim, H.H., Lee, Y., Eun, H.C., and Chung, J.H. (2008). Eicosapentaenoic acid inhibits TNF-alpha-induced matrix metalloproteinase-9 expression in human keratinocytes, HaCaT cells. *Biochem. Biophys. Res. Commun.* **36**: 343-349.
- [46] Kleniewska, P., Piechota-Polanczyk, A., Michalski, L., Michalska, M., Balcerczak, E., Zebrowska, M., and Goraca, A. (2013). Influence of block of NF-kappa B signaling pathway on oxidative stress in the liver homogenates. *Oxid Med Cell Longev.* **2013**: 308358
- [47] La, V.D., Lazzarin, F., Ricci, D., Fraternale, D., Genovese, S., Epifano, F., and Grenier, D. (2010). Active principles of *Grindelia robusta* exert antiinflammatory properties in a macrophage model. *Phytother Res.* **24**(11): 1687-1692.

- [48] Hwang, Y.P., Kim, H.G., Choi, J.H., Han, E.H., Kwon, K.I., Lee, Y.C., Choi, J.M., Chung, Y.C., Jeong, T.C., and Jeong, H.G. (2011). Saponins from the roots of *Platycodon grandiflorum* suppress ultraviolet A-induced matrix metalloproteinase-1 expression via MAPKs and NF- κ B/AP-1-dependent signaling in HaCaT cells. *Food Chem Toxicol.* **49**(12): 3374-3382
- [49] Mack, D.R. (2011). Probiotics in inflammatory bowel diseases and associated conditions. *Nutrients.* **3**(2): 245-264.
- [50] Stokes, J.H., and Pillsbury, D.H. (1930). The effect on the skin of emotional and nervous states: theoretical and practical consideration of a gastrointestinal mechanism. *Arch Dermatol Syphilol.* **22**: 962-993.
- [51] Fry, L. (1970). The gut and the skin. *Postgrad Med J.* **46**(541): 664-670.
- [52] Chapat, L., Chemin, K., Dubois, B., Bourdet-Sicard, R., and Kaiserlian, D. (2004). *Lactobacillus casei* reduces CD8⁺ T-cell-mediated skin inflammation. *Eur J Immunol.* **34**: 2520-2528.
- [53] Siver, R.H. (1961). *Lactobacillus* for the control of acne. *J Med Soc New Jersey.* **59**: 52-53.
- [54] Verdú, E.F., Bercik, P., Verma-Gandhu, M., Huang, X.X., Blennerhassett, P., Jackson, W., Mao, Y., Wang, L., Rochat, F., and Collins, S.M. (2006). Specific probiotic therapy attenuates antibiotic induced visceral hypersensitivity in mice. *Gut.* **55**: 182-190.
- [55] Teitelman, G., Joh, T.H., and Reis, D.J. (1981). Linkage of the brain-skin-gut axis: islet cells originate from dopaminergic precursors. *Peptides.* **2**(Suppl 2): 157-168.
- [56] Bowe, W.P., and Logan, A.C. (2011). Acne vulgaris, probiotics and the gut-brain-skin axis - back to the future? *Gut Pathog.* **3**(1): 1.

- [57] Arck, P., Handjiski, B., Hagen, E., Pincus, M., Bruenahl, C., Bienenstock, J., and Paus, R. (2010). Is there a 'gut-brain-skin axis'? *Exp Dermatol*. **19**(5): 401-405.
- [58] Gregory, B., and Ho, V.C. (1992). Cutaneous manifestations of gastrointestinal disorders. Part I. *J Am Acad Dermatol*. **26** (2 Pt 1): 153-166.
- [59] Gregory, B., and Ho, V.C. (1992). Cutaneous manifestations of gastrointestinal disorders. Part II. *J Am Acad Dermatol*. **26** (3 Pt 2): 371-383.
- [60] Girardin, M., and Seidman, E.G. (2011). Indications for the use of probiotics in gastrointestinal diseases. *Dig Dis*. **29**(6): 574-587
- [61] Youn, H.N., Lee, D.H., Lee, Y.N., Park, J.K., Yuk, S.S., Yang, S.Y., Lee, H.J., Woo, S.H., Kim, H.M., Lee, J.B., Park, S.Y., Choi, I.S., and Song, C.S. (2012). Intranasal administration of live *Lactobacillus* species facilitates protection against influenza virus infection in mice. *Antiviral Res*. **93**(1): 138-143.
- [62] Islam, M.A., Yun, C.H., Choi, Y.J., and Cho, C.S. (2010). Microencapsulation of live probiotic bacteria. *J Microbiol Biotechnol*. **20**(10): 1367-1377.
- [63] Rosenkranz, M.E., Schulte, D.J., Agle, L.M., Wong, M.H., Zhang, W., Ivashkiv, L., Doherty, T.M., Fishbein, M.C., Lehman, T.J., Michelsen, K.S., and Arditi, M. (2005). TLR2 and MyD88 contribute to *Lactobacillus casei* extract-induced focal coronary arteritis in a mouse model of Kawasaki disease. *Circulation*. **112**(19): 2966-2973.
- [64] Vintiñi, E.O., and Medina, M.S. (2011). Host immunity in the protective response to nasal immunization with a pneumococcal antigen associated to live and heat-killed *Lactobacillus casei*. *BMC Immunol*. **12**: 46.

- [65] Guéniche, A., Bastien, P., Ovigne, J.M., Kermici, M., Courchay, G., Chevalier, V., Breton, L., and Castiel-Higounenc, I. (2010). *Bifidobacterium longum* lysate, a new ingredient for reactive skin. *Exp Dermatol.* **19**(8): e1-8.
- [66] Ouwehand, A.C., Batsman, A., and Salminen, S. (2003). Probiotics for the skin: a new area of potential application? *Lett Appl Microbiol.* **36**: 327-331.
- [67] Rodrigues, K.L., Caputo, L.R., Carvalho, J.C., Evangelista, J., and Schneedorf, J.M. (2005). Antimicrobial and healing activity of kefir and kefir extract. *Int J Antimicrobial Agents* **25**: 404-408.
- [68] Niebuhr, M., Heratizadeh, A., Wichmann, K., Satzger, I., and Werfel, T. (2011). Intrinsic alterations of pro-inflammatory mediators in unstimulated and TLR-2 stimulated keratinocytes from atopic dermatitis patients. *Exp Dermatol.* **20**(6): 468-472.
- [69] Seo, H.S., Michalek, S.M., and Nahm, M.H. (2008). Lipoteichoic acid is important in innate immune responses to gram-positive bacteria. *Infect Immun.* **76**(1): 206-213
- [70] Lai, Y., Di Nardo, A., Nakatsuji, T., Leichtle, A., Yang, Y., Cogen, A.L., Wu, Z.R., Hooper, L.V., Schmidt, R.R., von Aulock, S., Radek, K.A., Huang, C.M., Ryan, A.F., and Gallo, R.L. (2009). Commensal bacteria regulate Toll-like receptor 3-dependent inflammation after skin injury. *Nat Med.* **15**(12): 1377-1382
- [71] Weill, F.S., Cela, E.M., Paz, M.L., Ferrari, A., Leoni, J., and González Maglio, D.H. (2012). Lipoteichoic acid from *Lactobacillus rhamnosus* GG as an oral photoprotective agent against UV-induced carcinogenesis. *Br J Nutr.* Epub ahead of print.

Tables and Figures Legend

Table 1. Various therapeutic effect on skin health by probiotics

Probiotics	Associated actions	Reference
<i>Bifidobacteria species</i>	Prevention of skin disorders – associated with immune abnormalities & improve sensitive skin	(32)
<i>Bacillus subtilis</i>	Prevention of atopic lesions	(15)
<i>Lactobacillus acidophilus</i>	Clinical improvement of acne	(50)
<i>Lactobacillus bulgaricus</i>	Clinical improvement of acne	(50)
<i>Lactobacillus casei</i>	Alleviation of skin inflammation	(45)
<i>Lactobacillus delbreuckii</i>	Inhibition of atopic dermatitis development	(19)
	Decrease of skin sensitivity	(12)
<i>Lactobacillus paracasei</i>	Acceleration of skin barrier function recovery	(35)
	Therapy for atopic dermatitis	(14)
<i>Lactobacillus rhamnosus</i>	Prevention against UV-induced skin damage	(16)
	Therapy for atopic dermatitis	(14)
<i>Lactobacillus reuteri</i>	Protection of epidermal keratinocytes	(31)
<i>Lactobacillus salivarius</i>	Therapy for atopic dermatitis	(37)




Figure 1. Probiotics can regulate the skin immunity through the inflammatory signals, NF-κB