



Hop bioactive compounds in prevention of nutrition-related noncommunicable diseases

Alejandro Bravo Iniguez & Mei-Jun Zhu

To cite this article: Alejandro Bravo Iniguez & Mei-Jun Zhu (2020): Hop bioactive compounds in prevention of nutrition-related noncommunicable diseases, Critical Reviews in Food Science and Nutrition, DOI: [10.1080/10408398.2020.1767537](https://doi.org/10.1080/10408398.2020.1767537)

To link to this article: <https://doi.org/10.1080/10408398.2020.1767537>



Published online: 28 May 2020.



Submit your article to this journal [↗](#)



View related articles [↗](#)



View Crossmark data [↗](#)

REVIEW



Hop bioactive compounds in prevention of nutrition-related noncommunicable diseases

Alejandro Bravo Iniguez and Mei-Jun Zhu

School of Food Science, Washington State University, Pullman, Washington, USA

ABSTRACT

Nutrition-related noncommunicable diseases (NR-NCDs) such as cardiovascular disease and type 2 diabetes both negatively impact the quality of life of many individuals and generate a substantial burden on society, demonstrating a need for intervention. Phytochemicals are investigated as a potential approach for combating NR-NCDs, and those found in hops have gained increased attention in recent decades. Hops, the strobile of the plant *Humulus lupulus*, are grown primarily for the brewing industry as they confer taste and increased shelf-life. The bitter acids represent the main compounds of interest for improving beer quality. Additionally, bitter acids as well as the prenylated chalcone xanthohumol, exhibit a wide range of health beneficial properties. This review summarizes those beneficial effects of bitter acids and xanthohumol on NR-NCDs, including inflammatory and immune diseases, obesity and metabolic disorders, as well as cancer prevention.

KEYWORDS

Hops; bitter acids; xanthohumol; obesity; cancer; intestinal health

Introduction

Globally, since the latter part of the twentieth century, the burden generated by nutrition-related noncommunicable disease (NR-NCD), which includes cancer, diabetes, and cardiovascular disease, has increased as a result of demographic and lifestyle changes (Popkin 2002; Popkin and Gordon-Larsen 2004). The rise of sedentary lifestyle, consumption of diets high in fats and sugar, and increased lifespans all contribute to the higher prevalence of NR-NCD (Musaiger and Al-Hazzaa 2012). In 2012, almost 50% of all noninstitutionalized adults in the United States suffered from at least one chronic condition. Among this group of adults, around half of them had two or more chronic conditions (Ward et al. 2014). Thus, it is imperative to reduce the strain resulting from NR-NCDs. Phytochemicals found in commonly cultivated plants display bioactive properties with the potential to help combat NR-NCDs (Zhang et al. 2015).

Hops (*Humulus lupulus*) have interacted with human civilization for centuries and possess unique chemical constituents which have received attention in the last decades. Belonging to the family Cannabaceae, *Humulus* most likely originated in China before spreading elsewhere (Murakami et al. 2006). The genus contains three species, *H. lupulus*, *H. japonicus*, and *H. yunnanensis*, all of which are dioecious, herbaceous vines (Small 1978). Documentation over the cultivation of hops goes as far back as the ninth century AD. Cultivation surged following the Middle Ages, once it became established as the principal flavoring agent of beer (Behre 1999). Outside of brewing, historic usage of hops includes serving as a remedy for different ailments. In various parts of the world, traditional medicine included the

intake of hops to treat inflammation, pneumonia, indigestion, insomnia, and dysentery (Zanoli and Zavatti 2008).

The glandular trichomes of the female inflorescences are their most important components when it comes to the production of secondary metabolites of interest. They are also referred to as the lupulin glands and reside on the bracteoles, producing large quantities of bitter acids and prenylated chalcones (De Keukeleire et al. 2003; Naya and Kotake 1971). Bitter acids and xanthohumol, the most abundant prenylated flavonoid in hops (Stevens and Page 2004), are the most widely studied components with regards to health beneficial properties. The present review summarizes the beneficial effects of bitter acids and xanthohumol exert on NR-NCDs.

Structure and absorption

Bitter acids

Structure and chemistry

The bitter acids represent the group of compounds of greatest interest to brewers. Beer relies on bitter acids for flavor, stability, and protection against spoilage microorganisms (De Keukeleire 2000). The bitter acids include two kinds of prenylated phloroglucinols, α - (humulone) and β -acids (lupulone) (Figure 1(A–C)). Humulones contain two prenyl groups, while lupulones differ by containing three prenyl groups. Together humulones and lupulones contribute up to 20% or more of the dry weight of hop cones (De Keukeleire 2000). Within hops, they occur as mixtures with the main forms being humulone/lupulone, cohumulone/colupulone, and adhumulone/adlupulone. Other forms present in lower




Figure 1. Structure of humulone/lupulone analogues and isohumulone. A: humulone analogues; B: lupulone analogues; C: chemical formula of analogue R-group; D: isohumulone.

concentrations include posthumulone/postlupulone, prehumulone/prelupulone and adprehumulone/adprelupulone (Zhang et al. 2004). When exposed to increased heat, such as those utilized during the worst boiling stage of beer production, α -acids isomerize to either *cis*- or *trans*- iso- α -acids (Figure 1(D)) (Jaskula et al. 2008). The iso- α -acids possess superior bitterness than their precursors, making their contribution to the flavor of beer greater (De Keukeleire 2000).

Absorption and metabolism

The human colon adenocarcinoma line, Caco-2, undergoes spontaneous differentiation, ultimately resembling enterocytes and functioning as a model for intestinal absorption (Balimane et al. 2000). In Caco-2 cell monolayers, humulone displays greater absorption ($P_{\text{app}} = 14 \pm 2 \times 10^{-6} \text{ cm/s}$) than lupulone ($P_{\text{app}} = 0.9 \pm 0.1 \times 10^{-6} \text{ cm/s}$). The absorption of lupulone resembles that of atenolol ($P_{\text{app}} = 0.45 \pm 0.18 \times 10^{-6} \text{ cm/s}$), a marker of lower permeability. Among the analogues of humulone and lupulone, different absorption values occur. Absorption of cohumulone ($P_{\text{app}} = 41 \pm 2 \times 10^{-6} \text{ cm/s}$) surpasses that of humulone. Colupulone ($P_{\text{app}} = 2.1 \pm 0.9 \times 10^{-6} \text{ cm/s}$) demonstrates a similar pattern with that of lupulone (Cattoor et al. 2010). The absorption of isohumulone in Caco-2 cells ($P_{\text{app}} = 4.62 \pm 1.29 \times 10^{-6} \text{ cm/s}$) was greater than that of lupulone (Cattoor et al. 2011).

Following intake, humulones, lupulones, and isohumulones all undergo biotransformation resulting in different metabolites. After being incubated with either human or rabbit liver microsomes for 2 hours, isohumulones displayed

the greatest resistance to metabolism (Cattoor et al. 2013). Human liver microsomes metabolized $73 \pm 3\%$, $95 \pm 1\%$, and $48 \pm 2\%$ of incubated humulone, lupulone, and isohumulone, respectively, producing humulinones, hulu-pones, and humulinic acid (Cattoor et al. 2013).

Xanthohumol

Structure and chemistry

Xanthohumol is a prenylated chalcone (Figure 2). Chalcones are open-chain flavonoids with two aromatic rings (Nowakowska 2007). Xanthohumol contributes to 0.1–1% of hop cones' dry weight, making it the major prenylflavonoid constituent (Stevens and Page 2004). Like the bitter acids found in hops, xanthohumol undergoes isomerization to iso-xanthohumol (Figure 2) during the brewing process, becoming less biologically active (Stevens et al. 1999). Combined with low solubility, isomerization results in low levels of xanthohumol present in beer (Karabín et al. 2013).

Absorption and metabolism

Xanthohumol is poorly absorbed ($P_{\text{app}} = 0.133 \pm 0.003 \times 10^{-6} \text{ cm/s}$), similar to sucrose ($P_{\text{app}} = 0.196 \pm 0.05 \times 10^{-6} \text{ cm/s}$), another marker of low permeability (Pang et al. 2007). Pang et al. (2007) also found that xanthohumol primarily accumulates in the cytosol upon entering enterocytes and binds to cytosolic proteins. Following oral administration, the majority of xanthohumol gets excreted in feces (Avula et al. 2004). The bioavailability of xanthohumol decreases with increased dosage, a trend




Figure 2. Structure of xanthohumol and isoxanthohumol. A: xanthohumol; B: isoxanthohumol.

exhibited by other flavonoids, with low, medium, and high doses resulting in 33%, 13%, and 11% bioavailability, respectively in rats (Legette et al. 2012). The maximum plasma concentrations of xanthohumol 1 hour following a single oral intake at a dose of 20, 60, or 180 mg were $45 \pm 7 \mu\text{g/L}$, $67 \pm 11 \mu\text{g/L}$, and $133 \pm 23 \mu\text{g/L}$, respectively (Legette et al. 2014).

Out of the prenylated compounds present in hops, 8-prenylnarigenin exerts the greatest estrogenic activity (Milligan et al. 2000). Since it can be produced from isoxanthohumol, there is concern that consumption of dietary xanthohumol and isoxanthohumol results in increased 8-prenylnarigenin exposure (Possemiers et al. 2005). In a human study, test subjects who took xanthohumol had either trace amounts or no circulating 8-prenylnarigenin (Legette et al. 2014). In addition, formation of 8-prenylnarigenin depends on gut microbiota. A study comparing germ-free mice with human microbiota-associated mice found that only the latter contained 8-prenylnarigenin in their excrements (Hanske et al. 2010). Incubation of hop prenylflavonoids with 12 different human intestinal bacteria isolated from feces showed isoxanthohumol converting to 8-prenylnarigenin in one third of the samples (Possemiers et al. 2005). Intestinal microbes involved with the metabolism of xanthohumol and other prenylated chalcones include *Eubacterium limosum* that possess the ability to activate isoxanthohumol to 8-prenylnarigenin through O-demethylation and *Eubacterium ramulus* that metabolizes both xanthohumol and 8-prenylnarigenin to desmethyloxanthohumol (Paraiso et al. 2019; Possemiers et al. 2005).

Obesity and metabolic syndrome

Metabolic syndrome encompasses a variety of risk factors, and depending on the organization, gets defined differently. Typically, metabolic syndrome's prognosis involves a combination of some or all the following risk factors: hypertension, hyperglycemia, obesity, low levels of high-density lipoprotein, and high triglyceride levels. It is estimated that about one-third of the population of the United States are obese, which is closely associated with metabolic syndrome, and have an increased relative risk of mortality (Saklayen 2018; Wu et al. 2010). This obesity is mainly caused by excessive energy intake, excessive white tissue build-up, and adipocyte hypertrophy, detrimental for human health metabolically (Bartelt and Heeren 2014). The potential of hop-

derived bioactive components to mitigate weight gain (Table 1) and other markers of metabolic syndrome (Figure 3) were summarized in this section.

Antiobesity effects

The supplementation of isomerized hop extract to a high fat diet (HFD) attenuated weight and adipose tissue gain in C57BL/6N mice in a dose-dependent manner (Ayabe et al. 2018; Yajima et al. 2005). Similarly, isomerized hop extract supplementation reduced weight gain in KK-A^y mice given a HFD (Yajima et al. 2005). In a randomized, double-blind placebo-controlled human study, matured bitter acid extract reduced both visceral and total body fat in overweight participants (Morimoto-Kobayashi et al. 2015). In addition, dietary xanthohumol supplementation reduced weight gain in HFD-fed male C57BL/6J mice (Miranda et al. 2016). The addition of a xanthohumol-rich extract to HFD fed male Wistar rats also resulted in decreased weight gain compared to the HFD control (Yui et al. 2014).


Brown adipose tissue

Brown adipose tissue (BAT) expresses uncoupling protein 1 (UCP1) which uncouples oxidative phosphorylation to dissipate energy as heat, and functions as a key site for preventing obesity and metabolic dysfunction. BAT can be separated into classic brown adipose tissue that develops during the fetal stage and inducible brown adipocytes (also called beige, brown-in-white adipocytes) which are introduced later in life (Bartelt and Heeren 2014). BAT activity is activated during cold exposure and found to be higher in lean individuals (van Marken Lichtenbelt et al. 2009). BAT recruitment can increase energy expenditure and assist with body fat reduction. It can be initiated by both cold exposure and certain chemicals such as capsinoid analogs (Yoneshiro et al. 2012) and β 3-adrenergic receptor agonists (Cypess et al. 2015; Yoneshiro et al. 2013). Dietary polyphenols such as resveratrol (Wang et al. 2015), epigallocatechin-3-gallate (Lee et al. 2017), berberine (Zhang et al. 2014), as well as polyphenol-rich raspberry (Xing et al. 2018) supplementation promote white adipose tissue browning and lipid oxidation.

Consistently, the matured bitter acids stimulate BAT through enhancing BAT sympathetic nerve activity and UCP1 (Morimoto-Kobayashi et al. 2015). A subsequent

Table 1. Anti-obesity effects of bitter acids and xanthohumol.

Compound	Model	Outcomes	References
Matured hop extract	Overweight human participants	Decrease of total fat area Decrease of visceral fat area	(Morimoto-Kobayashi et al. 2015)
Iso- α -acids	Male C57BL/6J mice	Decreased weight and adipose tissue gain resulting from consumption of a high fat diet	(Ayabe et al. 2018)
Isomerized hop extract	Female C57BL/6N mice	Decreased weight gain for both mice receiving standard diets and mice receiving high fat diets	(Yajima et al. 2005)
Isomerized hop extract	Male KK- A^y mice	Decreased weight gain resulting from consumption of a high fat diet.	(Yajima et al. 2005)
Xanthohumol	Male and female Zucker obese fa/fa rats	Decreased weight gain among male rats	(Legette et al. 2013)
Xanthohumol	Male KK- A^y mice	Non-significant decreased weight gain	(Nozawa 2005)

**Figure 3.** Potential mechanisms of hop phytochemicals in prevention of obesity and metabolic syndromes. Hop phytochemicals regulate energy expenditure through manipulation of skeletal muscle respiration, brown/beige adipose tissue thermogenesis, and bile acid receptor signaling via altering gut microbiome, combating weight gain. Furthermore, hop phytochemicals mitigate inflammation and decrease insulin resistance. Green arrows indicate promotive effects. Red arrows indicate inhibitory effects.

study found that matured bitter acid treatment induces cholecystokinin (CCK) secretion and CCK signaling is required for the upregulation of BAT sympathetic nerve activity, which increases BAT thermogenesis (Yamazaki et al. 2019). Xanthohumol induced AMP-activated protein kinase (AMPK) regulated being through promotion of proteins such as UCP1 (Samuels et al. 2018).

Skeletal muscle respiration

Respiration of skeletal muscle is a potential avenue for combating obesity as skeletal muscle impacts resting metabolic rate, and mice generated to overexpress UCP in their skeletal muscle resisted HFD induced weight gain (Li et al. 2000). Resveratrol is known to exert an effect on mitochondrial function and increases expression of UCP3 in skeletal muscle of mice (Lagouge et al. 2006). Treatment with 16.9 mg/kg BW of xanthohumol decreased weight gain in male Zucker fa/fa rats (Legette et al. 2013). Such effects are likely through mitochondrial uncoupling and stress response induction (Kirkwood et al. 2013).

Bile acid signaling

The farnesoid X receptor (FXR) is a nuclear hormone receptor that interacts with bile acids to help regulate energy expenditure (Ma and Patti 2014). Mice receiving HFD supplemented with a gut-restricted FXR agonist displayed ameliorated weight gain and enhanced browning of white

adipose tissue (Fang et al. 2015). Berberine safeguards against HFD induced obesity in wild-type mice but not their FXR knockout counterparts (Sun et al. 2017). Intake of a basal diet containing 1% xanthohumol lowered white adipose tissue weight in HFD KK- A^y mice (Nozawa 2005). The same study identified xanthohumol as a ligand of FXR (Nozawa 2005). In addition, bioactive compounds from hops can alter gut microbiota, which converts primary bile acids to secondary bile acids, which preferably binds to another bile acid receptor, TGR5, to activate intracellular thyroid hormone signaling and prevent obesity and metabolic diseases (Watanabe et al. 2006).

Markers of metabolic syndrome

Antioxidative effects

Adiposity correlates with increased systemic oxidative stress in both humans and animals (Furukawa et al. 2004). Damage from oxidative stress caused by reactive oxygen species (ROS) and reactive nitrogen species (RNS) plays an important role in aging and the progression of different disease states (Lee et al. 2004). Excessive ROS and RNS production generates oxidative stress (Valko et al. 2007), which results in lipid peroxidation, the formation of DNA adducts, and protein damage. Antioxidants prevent the oxidation through various means such as reducing molecular oxygen, removing harmful metal ions, and scavenging radicals that trigger the initiation of oxidation (Pisoschi and Pop 2015).

Table 2. Anti-inflammatory effects of bitter acids and xanthohumol.

Compound	Model	Mechanisms of action	References
Humulone	L929sA fibrosarcoma and COS1L2A fibroblast cells	Inhibited TNF-induced activity of NF- κ B, AP-1, and CREB.	(Van Cleemput et al. 2009)
	MC3T3-E1 osteoblast cells	Suppression of TNF- α -induced COX-2 activity	(Yamamoto et al. 2000)
	Mouse skin exposed to TPA	Downregulation of COX-2 expression Suppression of NF- κ B, and AP-1 DNA binding Inhibition of IKK and JNK	(Lee et al. 2007)
Lupulone	L929sA fibrosarcoma and COS1L2A fibroblast cells	Inhibited TNF- α -induced activity of NF- κ B, AP-1, and CREB.	(Van Cleemput et al. 2009)
	RAW 264.7 macrophage cells	Inhibition of LPS-induced NO and PGE ₂ activity Down regulation of iNOS and COX-2 Suppression of PI3K/Akt/IKK and ERK1/2	(Tang et al. 2011)
Isohumulone	L929sA fibrosarcoma and COS1L2A fibroblast cells	Inhibition of TNF- α -induced activity of NF- κ B, AP-1, and CREB.	(Van Cleemput et al. 2009)
	RAW 264.7 macrophage cells	Suppression of PGE2 Inhibition of NF- κ B activity	(Konda et al. 2009; Nozawa et al. 2005)
	Male Fischer 344 rats treated with AOM	Suppression of PGE2	(Nozawa et al. 2005)
Xanthohumol	RAW 264.7 macrophage cells	Suppression of NF- κ B, STAT-1 α , and IRF-1 activity	(Cho et al. 2008)
	Primary human hepatocytes and hepatic stellate cells	Suppression of NF- κ B	(Dorn et al. 2010)
	Male C57BL/6 mice treated with DSS	Inhibition of TNF- α and IL-1 β induction Inhibition of I κ B α phosphorylation Downregulation of COX-2	(Cho et al. 2018)

AP-1: activator protein-1; COX-2: cyclooxygenase-2; CREB: cAMP-response element-binding protein; ERK: extracellular signal-regulated kinase; I κ B: inhibitor κ B; IKK: I κ B kinase; IL: interleukin; iNOS: inducible nitric oxide synthase; IRF-1: IFN regulatory factor-1; JNK: c-Jun-N-terminal kinase; LPS: lipopolysaccharide; NF- κ B: Nuclear factor-kappa B; PGE2: prostaglandin E2; PI3K: phosphatidylinositol 3-kinase; STAT: signal transducer and activator of transcription; TNF: tumor necrosis factor; TPA: tumor promoter 12-O-tetradecanoylphorbol-13-acetate.

The antioxidant capacity of humulone and lupulone was analyzed utilizing 2,2-diphenyl-1-picrylhydrazyl (DPPH), and it was found that their radical scavenging activities (3.2×10^{-5} and 2.5×10^{-5} M, IC₅₀, respectively) are comparable to those of α -tocopherol and ascorbic acid (Tagashira et al. 1995). Humulone and lupulone inhibited lipid peroxidation with an IC₅₀ of 7.9×10^{-6} and 3.9×10^{-5} M, respectively (Tagashira et al. 1995). Using the oxygen radical absorbance capacity (ORAC) assay, xanthohumol was found to be 8.9 times more effective at scavenging hydroxyl radicals and 2.9 times more effective at scavenging peroxy radicals than Trolox, a vitamin E analogue and reference compound (Gerhauser et al. 2002). Comparing the antioxidant capacity of xanthohumol, humulone, and lupulone, humulone possessed the most effective scavenging activity against nitrite radical and DPPH, while xanthohumol displayed the strongest potential against hydroxyl radical, peroxy radical, superoxide, and ferric iron (Yamaguchi et al. 2009). In a different analysis of several hop constituents, humulone displayed the greatest hydrogen peroxide scavenging ability (Gorjanović et al. 2013). Xanthohumol pretreatment reduced lipopolysaccharides (LPS) induced reactive oxygen species formation and lessened the depletion of two antioxidant enzymes, superoxide dismutase and glutathione, in the bronchoalveolar lavage fluid and lungs of C57BL/6 mice, respectively (Lv et al. 2017).

Anti-inflammatory effects

Excessive white adipose tissue accumulation and adipocyte hypertrophy lead to low grade chronic inflammation.

Obesity results in increased quantities of macrophages present in adipose tissue contributing to inflammation (Weisberg et al. 2003; Xing et al. 2018; Xu et al. 2003). The macrophages in the white adipose tissue of obese mice also display reduced expression of anti-inflammatory cytokine, IL-10 (Lumeng et al. 2007). Phytochemical and phytochemical-rich dietary supplements suppress inflammatory signaling resulting from obesity. Quercetin suppressed macrophage infiltration into epididymis adipose tissue and inflammation in C57BL/6mice (Dong et al. 2014). Dietary red raspberry supplementation mitigated macrophage infiltration and inflammatory signaling in white adipose tissues of HFD fed mice (Xing et al. 2018). Similarly, bitter acids and xanthohumol are capable of mitigating inflammation by targeting different inflammatory signaling pathways (Table 2).

Nuclear factor- κ B (NF- κ B), a mediator of inflammatory genes plays a role in the pathogenesis of various inflammatory diseases (Liu et al. 2017). Humulones, lupulones, and isohumulones inhibit NF- κ B-dependent gene transcription, with humulone and lupulone being more effective than isohumulone (Van Cleemput et al. 2009). Humulone supplementation suppressed tumor necrosis factor- α (TNF- α) induced transcription of cyclooxygenase-2 (COX-2) in MC3T3-E1 cells (Yamamoto et al. 2000). Humulone down-regulated COX-2 in 12-O-tetradecanoylphorbol-13 acetate (TPA)-treated mouse skin, through suppression of NF- κ B and activator protein-1 (AP-1), attributed to inhibition of inhibitory kappaB (I κ B) kinase (IKK) and c-Jun-N-terminal kinase (JNK) (Lee et al. 2007). Reduced derivatives of lupulone displayed a capacity to downregulate phosphorylation

of extracellular signal-regulated kinase (ERK)1/2 and phosphatidylinositol 3-kinase (PI3K)/protein kinase B (Akt)/IKK in macrophages treated with LPS, suppressing NF- κ B signaling and downregulating the expression of inducible nitric oxide synthase (iNOS) and COX-2 (Tang et al. 2011).

Humulone's isomerized counterparts can also lessen the severity of inflammation through suppression of proinflammatory mediators. In RAW 264.7 macrophage cells, rho iso- α -acids suppressed expression of NF- κ B and its downstream targets and reduced glycogen synthase kinase 3 α (GSK3 α) activity and β -catenin phosphorylation (Konda et al. 2009). Isohumulone supplementation decreased prostaglandin E2 generation in LPS/IFN- γ treated RAW 264.7 cells and azoxymethane (AOM)-treated rats (Nozawa et al. 2005).

Inflammation contributes to the increased risk of cognitive impairment within metabolic syndrome (Yaffe et al. 2004). There is a growing interest in studying the capacity of iso- α -acids in decreasing inflammation in the tissue of the central nervous system. Iso- α -acids reduced the levels of IL-1 β , IL-16, IL-12p40, macrophage inflammatory protein-1 alpha (MIP-1 α), monocyte chemoattractant protein-1 (MCP-1), and TNF- α in microglial cells treated with LPS (Ano, Yoshikawa, et al. 2019). Dietary iso- α -acids conferred beneficial effects on cerebral inflammation of aged mice. Long-term dietary supplementation of 0.05% (w/w) isohumulones decreased age-related (Ano, Ohya, et al. 2019) and obesity induced (Ayabe et al. 2018) inflammation through decreasing proinflammatory cytokines such as of IL-1 β and TNF- α . Furthermore, short-term (3 or 7 days) daily oral administration of 1 mg/kg of isohumulones attenuated inflammatory cytokines such as IL-12p40 and MIP-1 α in Alzheimer's disease model mice (Ano, Yoshikawa, et al. 2019). Similarly, xanthohumol lessened LPS-induced inflammation in microglial cells through inhibition of NF- κ B activation, and increased the translocation of nuclear factor erythroid 2-related factor 2 (NRF2) into the nucleus (Lee et al. 2011). In senescence-accelerated prone mice (SAMP8), xanthohumol supplementation ameliorated age-related increase of TNF- α and IL-1 β in a dose-dependent manner (Rancán et al. 2017).

In RAW 264.7 cells treated with LPS, xanthohumol inhibited NF- κ B transcriptional activity in a dose-dependent manner (Cho et al. 2008). The suppression of NF- κ B by xanthohumol involved the degradation of I κ B α and suppressed p65 translocation in a cysteine-dependent manner (Harikumar et al. 2009). Xanthohumol supplementation decreased levels of MCP-1 and TNF- α in macrophages and in mice fed a nonalcoholic steatohepatitis diet (Dorn et al. 2010; Lupinacci et al. 2009). Similarly, xanthohumol supplementation suppressed NF- κ B's signaling pathway and proinflammatory cytokines, TNF- α , IL-1 β , and malondialdehyde production in Dextran sulfate sodium (DSS)-induced colitis mice, and further mitigated inflammation-induced damage (Cho et al. 2018).

Insulin resistance and glucose tolerance

A study involving both animal and human test subjects found that isohumulones activate both PPAR α and PPAR γ impacting insulin resistance (Yajima et al. 2004). A two-

week supplementation of isohumulones lowered non-fasting plasma glucose and lower insulin resistance in KK-A y mice and HFD fed C57BL/6N mice (Yajima et al. 2004). In a randomized double-blind placebo-controlled study with patients diagnosed with prediabetes, 32 mg of isohumulones per day improved hyperglycemia (Obara et al. 2009).

Furthermore, dietary xanthohumol supplementation lowered plasma glucose in male Zucker fa/fa rats (Legette et al. 2013). A 20-week dietary supplementation of xanthohumol or 8-prenylnaringenin (10 mg/L in 0.1% ethanol) reduced weight gain, glucose intolerance, and insulin resistance in male C57BL/6 mice fed a HFD which was associated with AMPK activation and downregulated 6-phosphofructo-2-kinase fructose-2,6-biphosphatase 3 (PFKFB3) in liver and skeletal muscle (Costa et al. 2017). *In vitro* study showed that xanthohumol can effectively inhibit α -glucosidase, an enzyme that hydrolyzes carbohydrates and impacts blood glucose levels (Liu et al. 2014).

Cardiovascular disease

Worldwide, the leading cause of mortality is ischemic heart disease (IHD). IHD falls under the category of atherosclerotic cardiovascular disease (ACD), which also includes ischemic stroke, another leading cause of global mortality (Barquera et al. 2015). In atherosclerosis, atheroma and plaque develop. The two ultimately contribute to artery occlusion after rupturing (Hansson and Libby 2006; Herrington et al. 2016). Known risk factors for atherosclerosis related to nutrition include adiposity, blood pressure, and blood cholesterol (Herrington et al. 2016).

The addition of isohumulone to rats fed a diet high in salt helped mitigate the rise in blood pressure induced by high sodium intake (Namikoshi et al. 2007). Feeding isohumulones to mice also increases high-density lipoprotein, and supplementation of 500 mg/kg BW of isomerized hop extract resulted in a decreased atherosclerosis index (Miura et al. 2005). When comparing the ability of different hop constituents to inhibit cholesteryl ester transfer protein (CETP), xanthohumol possessed the greatest activity, which was three times greater than that of humulone and lupulone (Hirata et al. 2012a). An *in vivo* study reported that xanthohumol decreased the accumulation of cholesterol in CETP-Tg mice fed high cholesterol diets (Hirata et al. 2012b).

Gut health

A wide breadth of conditions such as inflammatory bowel disease (IBD), gastritis, and colorectal cancer fall under the umbrella of gastrointestinal diseases, and in 2015, the economic burden generated by gastrointestinal diseases totaled \$135.9 billion (Peery et al. 2019). Maintenance of gut health is considered to be largely dependent on both the gastrointestinal microbiome and barrier (Bischoff 2011). Many chronic diseases involving dysregulated inflammatory responses appear to be influenced by host intestinal microbial communities (West et al. 2015). Evidence points toward proper formation of intestinal immune system requiring the




Figure 4. Potential mechanisms of hop phytochemicals on critical events during cancer development and progression. Hop phytochemicals suppress enzymes such as cytochrome P450 1A (CYP1A) that activate procarcinogens, and they enhance detoxifying enzyme quinone reductase; these compounds induce cell cycle arrest by targeting p53 and p21, and apoptosis through activation of the caspase-cascade. Furthermore, hop phytochemicals suppress cancer aggressiveness and metastasis by downregulating matrix metalloproteinases (MMP)-2 and 9, and decrease the development of new blood vessels by inhibiting vascular endothelial growth factor (VEGF). Green arrows indicate promotive effects. Red arrows indicate inhibitory effects.

presence of microorganisms. For example, *Bacteroides fragilis* colonization regulates the immune system by activating development of T-helper cells and correcting cytokine production (Mazmanian et al. 2005). Inflammation also impacts epithelial barrier function, and barrier disruption is a key etiological factor for IBD and a number of associated diseases (Vetrano et al. 2008). Research exploring the potential of hop derived compounds to modulate the gut microbiome and improve intestinal barrier integrity is brief and summarized in this section.

Effects against pathogenic bacteria and gut microbiota

Dysregulated microflora is associated with many immune-related diseases. Children who suffer from IBD (Maukonen et al. 2015), type 1 diabetes (Murri et al. 2013), and allergies (Candela et al. 2012) show compositional changes in their gut microbial communities compared to their disease-free peers. *Clostridium difficile* infection negatively impacts the levels of commensal intestinal bacteria including butyrate-producers (Antharam et al. 2013); hop bitter acids and xanthohumol exhibit inhibitory effects on pathogenic *C. difficile* isolated from patients with diarrhea, with xanthohumol being the most effective (Cermak et al. 2017). There is limited information regarding the impacts of hops and hop related bioactives on gut microbial composition. In chickens challenged with *Clostridium perfringens*, the addition of lupulone in drinking water decreased the abundance of *C. perfringens* (Tillman et al. 2011). In pigs, spent hops feeding reduced the abundance of *Streptococcus* spp. and *Clostridium* cluster XIVa in the fecal microbiota (Fiesel et al. 2014). However, the intestinal microbial composition of rats did not change following a dosage of 100 mg/kg/body weight of xanthohumol (Hanske et al. 2005). In contrast, a high level of supercritical CO₂ hop extract inhibited probiotic *Bifidobacterium* and butyrate-producing bacteria while increasing both *Enterobacteriaceae* and *Akkermansia*

(Blatchford et al. 2019), showing that the effects of hop bioactive compounds on gut microbiota are concentration dependent.

Effects on epithelial barrier function

The epithelium is in a constant state of renewal, requiring a careful balance of proliferation, differentiation, and apoptosis to function properly (Coskun et al. 2011). It not only assists with the transport of nutrients and waste but also serves as a protective barrier against foreign agents present in the luminal environment (Odenwald and Turner 2017). Like dysbiosis, impaired barrier function can be found in conditions such as IBD (König et al. 2016). Biopsies taken from colons of patients diagnosed with Crohn's diseases display impaired barrier function accompanied with decreased expression of tight junction proteins (Zeissig et al. 2007). Barrier function is largely dependent on junction proteins (Marchiando et al. 2011). Tight junction proteins reside on the lateral sides of epithelial cells close to the apical region, binding cells together and functioning as a selectively permeable barrier in the intracellular space (Farquhar and Palade 1963). Plant-derived chemicals and plant extracts such as quercetin (Suzuki and Hara 2009), kaempferol (Suzuki et al. 2011), naringenin (Noda et al. 2013) and polyphenol-rich purple potato extract (Sun et al. 2018) upregulate tight junction expression and enhance barrier function.

Consistently, 8 weeks of reduced iso-humulone supplementation in HFD mice increased intestinal alkaline phosphatase activity, a well-known epithelial differentiation marker, and mRNA expression of tight junction proteins, occludin and zonula occludens-1 (Everard et al. 2012). In Caco-2 cells, 8-prenylnaringenin, but not xanthohumol, pretreatment protected against TNF- α induced tight junction dysfunction (Luescher et al. 2017).

Table 3. Effects of bitter acids and xanthohumol on apoptosis and cell cycle.

Compound	Model	Mechanisms of action	References
Humulone	HL-60 cells	Enhanced expression of Fas and FasL Decreased expression of Bcl-2 and Bcl-X _L Mitochondrial dysfunction	(Chen and Lin 2004)
Lupulone	HL-60 cells	Enhanced expression of Fas and FasL Mitochondrial dysfunction	(Chen and Lin 2004)
	SW620 cells	Enhanced expression of Fas, FasL, TRAIL-R1 and R2 proteins. Mitochondrial dysfunction	(Lamy et al. 2007)
Xanthohumol	40-16 cells	Inhibition of antiapoptotic Bcl-2 Increased activation of caspases-3, -7, and -9	(Pan et al. 2005)
	SKOV3 and OVCAR3 cells	Increased expression of p21 Increased activation of caspase-3	(Drenzek et al. 2011)
	T98G cells	Increased activation of caspase-3 Inhibition of antiapoptotic Bcl-2 Increased intracellular levels of ROS	(Festa et al. 2011)
	Mouse bone marrow dendritic cells	Increased activation of caspases-3 and -8	(Xuan et al. 2010)

Bcl: B-cell lymphoma; Bcl-X_L: Bcl-extra large; Fas: Fas receptor; FasL: Fas ligand; ROS: reactive oxygen species; TRAIL: TNF-related apoptosis-inducing ligand; TRAIL-R1: TRAIL receptor 1/death receptor 4; TRAIL-R2: TRAIL receptor 2/death receptor 5.

Anticarcinogenic effects

Cancer imposes a significant burden on a global scale. It was estimated that in 2018, 18.1 million new cases of cancer and 9.6 million deaths occurred (Bray et al. 2018). The magnitude of cancer cases worldwide and epidemiological data pointing toward lifestyle behaviors such as diet impacting prevalence of certain kinds of cancer have resulted in increased interest in exploring the anticancer potential of various phytochemicals (Kotecha et al. 2016). Dietary anticarcinogenic intervention strives to offer protection of normal cells and disrupt cancer cell progression. The chemopreventative capacity of phytochemicals includes targeting the mechanisms behind mutagenesis, proliferation, angiogenesis, and mutagenesis of cancer cells (Surh 2003) (Figure 4). Carcinogenesis is a multistage process and plant derived chemicals such as polyphenols exert beneficial effects throughout the various stages (Ramos 2008). The capacity of hop derived compounds to exert anticarcinogenic effects was summarized below.

Biotransformation of carcinogens

Certain members of the cytochrome P450 (CYP) family activate various procarcinogens (Shimada et al. 1996). Inhibiting the activity of these enzymes is one strategy of chemoprevention. A study evaluating the potential of various hop prenylated flavonoids to inhibit P450 enzymes identified xanthohumol as a potent inhibitor of CYP1A1 and CYP1B1, both of which are responsible for the activation of carcinogenic polycyclic aromatic hydrocarbons (Henderson et al. 2000). A later study found that in rat hepatoma cells, both xanthohumol and isoxanthohumol inhibit CYP1A ($0.022 \pm 0.002 \mu\text{M}$ and $0.30 \pm 0.13 \mu\text{M}$ IC₅₀, respectively) (Gerhauser et al. 2002). Unlike xanthohumol's inhibitory effects on P450 enzymes that activate procarcinogens, colupone induces detoxification P450 enzymes. Mice fed a diet supplemented with colupulone showed increased expression of P4503A and P4502B (Mannering et al. 1992).

Along with affecting phase I enzymes, enhancing detoxifying phase II enzymes, such as quinone reductase (QR), serves as an approach for chemoprevention (Cuendet et al. 2006). Xanthohumol enhanced the activity of QR in Hepa 1c1c7 murine hepatoma cells (Dietz et al. 2005; Gerhauser et al. 2002; Miranda et al. 2000). Dietz et al. (2005) showed that the addition of dicumarol, a QR inhibitor, suppressed the ability of xanthohumol to prevent menadione-induced DNA damage. In HepG2 human liver cells, xanthohumol at 0.01 μM protected against the genotoxic effects of benzo(a)-pyrene and 2-amino-3-methylimidazo[4,5-f]quinoline (Plazar et al. 2007). Both humulone and isohumulone enhanced the activity of QR in Hepa1c1c7 cells, but the former also exhibited cytotoxicity (Bohr et al. 2008).

Induction of apoptosis and cell cycle arrest

In different cancer cell lines, hop constituents target an array of pathways to disrupt function and growth, culminating in cell death (Table 3). Utilizing human leukemia cell lines HL-60 and U937, a mixture containing both humulones and lupulones inhibited cell growth, with U937 being more resilient (Chen and Lin 2004). Treatment with bitter acids also altered mitochondrial function and upregulated Fas and FasL (Chen and Lin 2004). Similarly, in SW620 colon carcinoma cells, lupulones displayed the ability to induce apoptosis through activation of Fas as well as the TNF α -related apoptosis-inducing ligand (TRAIL) pathways (Lamy et al. 2007). A subsequent study found that lupulone interacts with death receptor 4 (DR4)/DR5 to activate both the extrinsic and intrinsic pathways, resulting in increased caspase cascade activity (Lamy et al. 2008).

Xanthohumol decreased the viability of various cancer cells such as prostate, breast, colon, and ovarian (Colgate et al. 2007; Miranda et al. 1999). Death-receptor and mitochondrial-mediated pathways of programmed cell death both activated as a result of xanthohumol treatment in colon cancer cells (Pan et al. 2005). Xanthohumol also targets the Notch pathway, resulting in cell cycle arrest and apoptosis

in ovarian cancer cells (Drenzek et al. 2011). In dendritic cells and the T98G human malignant glioblastoma cell line, xanthohumol induced caspase activity (Festa et al. 2011; Xuan et al. 2010). In the latter, xanthohumol generated ROS and induced apoptosis through the mitochondrial pathway (Festa et al. 2011).

Inhibition of angiogenesis and metastasis

The ability of cancer cells to develop vascular networks grants them access to nutrients needed for growth and a route by which to invade other parts of the body. Various factors play a role in the progression of angiogenesis. Vascular endothelial growth factor (VEGF) is a mitogen that functions as a main mediator of angiogenesis. Inhibition of VEGF results in suppressed tumor growth (Kim et al. 1993).

The antiangiogenic properties of humulone were discovered a decade ago (Shimamura et al. 2001). Work done *in vitro* found that humulone decreased endothelial cell expression of VEGF and vascular endothelial cell tube formation (Shimamura et al. 2001). Shimamura et al. (2001) also found that humulone inhibited angiogenesis in chick embryo chorioallantoic membranes in a dose-dependent manner. Similarly, lupulone inhibits angiogenesis, supplementation of 0.01% lupulone in drinking water reduced angiogenesis by 50% in mice compared to the control mice without lupulone (Siegel et al. 2008). iNOS has been implicated as a key regulator of VEGF since it generates nitric oxide, which participates in the production of VEGF. Lupulone and xanthohumol both suppressed iNOS in RAW 264.7 cells, with the latter being more effective (Zhao et al. 2003).

Xanthohumol suppressed TNF- α -induced NF- κ B activation in leukemia cells (Dell'Eva et al. 2007) and downregulated NF- κ B signaling in pancreatic cancer cells, resulting in decreased VEGF expression and production (Saito et al. 2018). In endothelial cells, antiangiogenic effects of xanthohumol involved AMPK activation mediated by calmodulin-dependent kinase kinase β (CAMMK β) (Gallo et al. 2016). In breast cancer cells, xanthohumol reduced factor VIII, an endothelial marker and inhibited NF- κ B signaling (Monteiro et al. 2008). Supplementation of 2 μ M or 20 μ M xanthohumol in drinking water before injection of Matrigel or Kaposi's sarcoma tumors, respectively, suppressed angiogenesis and tumor growth in mice compared to their respective controls (Albini et al. 2006). In addition, xanthohumol affected the adhesion of breast cancer cells by enhancing the E-cadherin/catenin complex (Vanhoecke et al. 2005).

Among the types of cancer, matrix metalloproteinases (MMP) are almost ubiquitously upregulated, and the various members of the MMP family contribute to the progression of cancer during different stages of development (Egeblad and Werb 2002). MMP-2 and -9 contribute to angiogenesis and invasion, demonstrated by mice with MMP-2 and -9 deficiencies exhibiting impaired tumor invasion and vascularization (Masson et al. 2005). In fibrosarcoma cells, xanthohumol suppressed the activity of MMP-2 and -9 (Kim et al. 2009). In triple negative breast cancer cells xanthohumol suppressed MMP-9 but not MMP-2 (Kim et al. 2013).

Conclusion and perspective

Chronic diseases are both prevalent among the population and generate a significant burden on society. Phytochemicals present in commonly grown agricultural commodities have gained increasing interests in the last several years as potential bioactive compounds capable of ameliorating the complications generated by NR-NCDs. Unlike other sources of health beneficial phytochemicals, hops are not consumed whole. The main source of dietary bitter acids and xanthohumol is beer, and levels tend to be low. An alternative to beer drinking is the development of new hop-derived therapeutics. Hop derived compounds show promise as therapeutics; both *in vitro* and *in vivo* studies support the notion that bitter acids and xanthohumol mitigate symptoms of NR-NCDs. Hop phytochemicals show potential in ameliorating hallmarks of metabolic syndrome by influencing energy expenditure and display anti-cancer properties capable of targeting carcinogenesis at different stages (Figures 3 and 4). While studies have begun elucidating the pathways responsible for the anti-inflammatory, antiobesity, and chemopreventative effects of hop compounds, few have deeply explored their effects on intestinal health and microbiota. Intestinal health impacts overall health, making it an important target for disease prevention, and it remains unclear if manipulation of the gut microbiome is involved in the improvement of biomarkers generated by hops. Moving forward, emphasis should focus on safe, effective dosage for humans and its impacts on facets of intestinal health.

Disclosure statement

Bravo Iniguez and Zhu have no conflicts of interests.

Funding

This work was financially supported by National Institutes of USDA-National Institute of Food and Agriculture (USDA-NIFA) (2018-67017-27517), and Washington State University Agricultural Research Center Emerging Research Issues Competitive Grant.

References

- Albini, A., R. Dell'Eva, R. Vené, N. Ferrari, D. R. Buhler, D. M. Noonan, and G. Fassina. 2006. Mechanisms of the antiangiogenic activity by the hop flavonoid xanthohumol: NF- κ B and Akt as targets. *FASEB Journal* 20 (3):527–9. doi: 10.1096/fj.05-5128fj.
- Ano, Y., R. Ohya, K. Kondo, and H. Nakayama. 2019. Iso- α -acids, hop-derived bitter components of beer, attenuate age-related inflammation and cognitive decline. *Frontiers in Aging Neuroscience* 11: 16.
- Ano, Y., M. Yoshikawa, Y. Takaichi, M. Michikawa, K. Uchida, H. Nakayama, and A. Takashima. 2019. Iso- α -acids, bitter components in beer, suppress inflammatory responses and attenuate neural hyperactivation in the hippocampus. *Frontiers in Pharmacology* 10: 81.
- Antharam, V. C., E. C. Li, A. Ishmael, A. Sharma, V. Mai, K. H. Rand, and G. P. Wang. 2013. Intestinal dysbiosis and depletion of butyrogenic bacteria in *Clostridium difficile* infection and nosocomial

- diarrhea. *Journal of Clinical Microbiology* 51 (9):2884–92. doi: [10.1128/JCM.00845-13](https://doi.org/10.1128/JCM.00845-13).
- Avula, B., M. Ganzera, J. E. Warnick, M. W. Feltenstein, K. J. Sufka, and I. A. Khan. 2004. High-performance liquid chromatographic determination of xanthohumol in rat plasma, urine, and fecal samples. *Journal of Chromatographic Science* 42 (7):378–82. doi: [10.1093/chromsci/42.7.378](https://doi.org/10.1093/chromsci/42.7.378).
- Ayabe, T., R. Ohya, K. Kondo, and Y. Ano. 2018. Iso- α -acids, bitter components of beer, prevent obesity-induced cognitive decline. *Scientific Reports* 8 (1):4760. doi: [10.1038/s41598-018-23213-9](https://doi.org/10.1038/s41598-018-23213-9).
- Balimane, P. V., S. Chong, and R. A. Morrison. 2000. Current methodologies used for evaluation of intestinal permeability and absorption. *Journal of Pharmacological and Toxicological Methods* 44 (1):301–12. doi: [10.1016/S1056-8719\(00\)00113-1](https://doi.org/10.1016/S1056-8719(00)00113-1).
- Barquera, S., A. Pedroza-Tobías, C. Medina, L. Hernández-Barrera, K. Bibbins-Domingo, R. Lozano, and A. E. Moran. 2015. Global overview of the epidemiology of atherosclerotic cardiovascular disease. *Archives of Medical Research* 46 (5):328–38. doi: [10.1016/j.arcmed.2015.06.006](https://doi.org/10.1016/j.arcmed.2015.06.006).
- Bartelt, A., and J. Heeren. 2014. Adipose tissue browning and metabolic health. *Nature Reviews Endocrinology* 10 (1):24–36. doi: [10.1038/nrendo.2013.204](https://doi.org/10.1038/nrendo.2013.204).
- Behre, K.-E. 1999. The history of beer additives in Europe — A review. *Vegetation History and Archaeobotany* 8 (1–2):35–48. doi: [10.1007/BF02042841](https://doi.org/10.1007/BF02042841).
- Bischoff, S. C. 2011. 'Gut health': A new objective in medicine? *BMC Medicine* 9:24. doi: [10.1186/1741-7015-9-24](https://doi.org/10.1186/1741-7015-9-24).
- Blatchford, P. A., S. G. Parkar, W. Hopkins, J. R. Ingram, and K. H. Sutton. 2019. Dose-dependent alterations to in vitro human microbiota composition and butyrate inhibition by a supercritical carbon dioxide hops extract. *Biomolecules* 9 (9):390. doi: [10.3390/biom9090390](https://doi.org/10.3390/biom9090390).
- Bohr, G., K. Klimo, J. Zapp, H. Becker, and C. Gerhäuser. 2008. Cancer chemopreventive potential of humulones and isohumulones (hops α - and iso- α -acids): Induction of NAD(P)H:Quinone Reductase as a novel mechanism. *Natural Products Communications* 3 (12):1971–6. doi: [10.1177/1934578X0800301206](https://doi.org/10.1177/1934578X0800301206).
- Bray, F., J. Ferlay, I. Soerjomataram, R. L. Siegel, L. A. Torre, and A. Jemal. 2018. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA: A cancer journal for clinicians* 68 (6):394–424. doi: [10.3322/caac.21492](https://doi.org/10.3322/caac.21492).
- Candela, M., S. Rampelli, S. Turroni, M. Severgnini, C. Consolandi, G. De Bellis, R. Masetti, G. Ricci, A. Pession, and P. Brigidi. 2012. Unbalance of intestinal microbiota in atopic children. *BMC Microbiology* 12:95. doi: [10.1186/1471-2180-12-95](https://doi.org/10.1186/1471-2180-12-95).
- Cattoor, K., M. Bracke, D. Deforce, D. De Keukeleire, and A. Heyerick. 2010. Transport of hop bitter acids across intestinal Caco-2 cell monolayers. *Journal of Agricultural and Food Chemistry* 58 (7):4132–40. doi: [10.1021/jf904079h](https://doi.org/10.1021/jf904079h).
- Cattoor, K., M. Dresel, L. De Bock, K. Boussey, J. Van Bocxlaer, J.-P. Remon, D. De Keukeleire, D. Deforce, T. Hofmann, and A. Heyerick. 2013. Metabolism of hop-derived bitter acids. *Journal of Agricultural and Food Chemistry* 61 (33):7916–24. doi: [10.1021/jf300018s](https://doi.org/10.1021/jf300018s).
- Cattoor, K., J.-P. Remon, K. Boussey, J. V. Bocxlaer, M. Bracke, D. De Keukeleire, D. Deforce, and A. Heyerick. 2011. Bioavailability of hop-derived iso- α -acids and reduced derivatives. *Food & Function* 2 (7):412–22. doi: [10.1039/c1fo10009b](https://doi.org/10.1039/c1fo10009b).
- Cermak, P., J. Olšovská, A. Mikyska, M. Dusek, Z. Kadleckova, J. Vaníček, O. Nyc, K. Sigler, V. Bostikova, and P. Bostik. 2017. Strong antimicrobial activity of xanthohumol and other derivatives from hops (*Humulus lupulus* L.) on gut anaerobic bacteria. *Acta Pathologica, Microbiologica et Immunologica Scandinavica* 125 (11):1033–8. doi: [10.1111/apm.12747](https://doi.org/10.1111/apm.12747).
- Chen, W.-J., and J.-K. Lin. 2004. Mechanisms of cancer chemoprevention by hop bitter acids (beer aroma) through induction of apoptosis mediated by Fas and caspase cascades. *Journal of Agricultural and Food Chemistry* 52 (1):55–64. doi: [10.1021/jf034737u](https://doi.org/10.1021/jf034737u).
- Cho, J.-M., S.-M. Yun, Y.-H. Choi, J. Heo, N.-J. Kim, S.-H. Kim, and E.-H. Kim. 2018. Xanthohumol prevents dextran sulfate sodium-induced colitis via inhibition of IKK β /NF- κ B signaling in mice. *Oncotarget* 9 (1):866–80. doi: [10.18632/oncotarget.23183](https://doi.org/10.18632/oncotarget.23183).
- Cho, Y.-C., H. J. Kim, Y.-J. Kim, K. Y. Lee, H. J. Choi, I.-S. Lee, and B. Y. Kang. 2008. Differential anti-inflammatory pathway by xanthohumol in IFN- γ and LPS-activated macrophages. *International Immunopharmacology* 8 (4):567–73. doi: [10.1016/j.intimp.2007.12.017](https://doi.org/10.1016/j.intimp.2007.12.017).
- Colgate, E. C., C. L. Miranda, J. F. Stevens, T. M. Bray, and E. Ho. 2007. Xanthohumol, a prenylflavonoid derived from hops induces apoptosis and inhibits NF- κ B activation in prostate epithelial cells. *Cancer Letters* 246 (1–2):201–9. doi: [10.1016/j.canlet.2006.02.015](https://doi.org/10.1016/j.canlet.2006.02.015).
- Coskun, M., J. T. Troelsen, and O. H. Nielsen. 2011. The role of CDX2 in intestinal homeostasis and inflammation. *Biochimica et Biophysica Acta* 1812 (3):283–9. doi: [10.1016/j.bbadis.2010.11.008](https://doi.org/10.1016/j.bbadis.2010.11.008).
- Costa, R., I. Rodrigues, L. Guardão, S. Rocha-Rodrigues, C. Silva, J. Magalhães, M. Ferreira-de-Almeida, R. Negrão, and R. Soares. 2017. Xanthohumol and 8-prenylnaringenin ameliorate diabetic-related metabolic dysfunctions in mice. *The Journal of Nutritional Biochemistry* 45:39–47. doi: [10.1016/j.jnutbio.2017.03.006](https://doi.org/10.1016/j.jnutbio.2017.03.006).
- Cuendet, M., C. P. Oteham, R. C. Moon, and J. M. Pezzuto. 2006. Quinone reductase induction as a biomarker for cancer chemoprevention. *Journal of Natural Products* 69 (3):460–3. doi: [10.1021/np050362q](https://doi.org/10.1021/np050362q).
- Cypess, A. M., L. S. Weiner, C. Roberts-Toler, E. F. Elia, S. H. Kessler, P. A. Kahn, J. English, K. Chatman, S. A. Trauger, A. Doria, et al. 2015. Activation of human brown adipose tissue by a β 3-Adrenergic receptor agonist. *Cell Metabolism* 21 (1):33–8. doi: [10.1016/j.cmet.2014.12.009](https://doi.org/10.1016/j.cmet.2014.12.009).
- De Keukeleire, D. 2000. Fundamentals of beer and hop chemistry. *Química Nova* 23 (1):108–12. doi: [10.1590/S0100-40422000000100019](https://doi.org/10.1590/S0100-40422000000100019).
- De Keukeleire, J., G. Ooms, A. Heyerick, I. Roldan-Ruiz, E. Van Bockstaele, and D. De Keukeleire. 2003. Formation and accumulation of alpha-acids, beta-acids, desmethylxanthohumol, and xanthohumol during flowering of hops (*Humulus lupulus* L.). *Journal of Agricultural and Food Chemistry* 51 (15):4436–41. doi: [10.1021/jf034263z](https://doi.org/10.1021/jf034263z).
- Dell'Eva, R., C. Ambrosini, N. Vannini, G. Piaggio, A. Albini, and N. Ferrari. 2007. AKT/NF- κ B inhibitor xanthohumol targets cell growth and angiogenesis in hematologic malignancies. *Cancer* 110 (9):2007–11. doi: [10.1002/cncr.23017](https://doi.org/10.1002/cncr.23017).
- Dietz, B. M., Y. H. Kang, G. Liu, A. L. Eggler, P. Yao, L. R. Chadwick, G. F. Pauli, N. R. Farnsworth, A. D. Mesecar, R. B. van Breemen, et al. 2005. Xanthohumol isolated from *Humulus lupulus* Inhibits menadione-induced DNA damage through induction of quinone reductase. *Chemical Research in Toxicology* 18 (8):1296–305. doi: [10.1021/tx050058x](https://doi.org/10.1021/tx050058x).
- Dong, J., X. Zhang, L. Zhang, H. X. Bian, N. Xu, B. Bao, and J. Liu. 2014. Quercetin reduces obesity-associated ATM infiltration and inflammation in mice: A mechanism including AMPK α 1/SIRT1. *Journal of Lipid Research* 55 (3):363–74. doi: [10.1194/jlr.M038786](https://doi.org/10.1194/jlr.M038786).
- Dorn, C., B. Kraus, M. Motyl, T. S. Weiss, M. Gehrig, J. Schölmerich, J. Heilmann, and C. Hellerbrand. 2010. Xanthohumol, a chalcone derived from hops, inhibits hepatic inflammation and fibrosis. *Molecular Nutrition & Food Research* 54 (S2):S205–S213. doi: [10.1002/mnfr.200900314](https://doi.org/10.1002/mnfr.200900314).
- Drenzek, J. G., N. L. Seiler, R. Jaskula-Sztul, M. M. Rausch, and S. L. Rose. 2011. Xanthohumol decreases Notch1 expression and cell growth by cell cycle arrest and induction of apoptosis in epithelial ovarian cancer cell lines. *Gynecologic Oncology* 122 (2):396–401. doi: [10.1016/j.ygyno.2011.04.027](https://doi.org/10.1016/j.ygyno.2011.04.027).
- Egeblad, M., and Z. Werb. 2002. New functions for the matrix metalloproteinases in cancer progression. *Nature Reviews Cancer* 2 (3):161–74. doi: [10.1038/nrc745](https://doi.org/10.1038/nrc745).
- Everard, A., L. Geurts, M. Van Roye, N. M. Delzenne, and P. D. Cani. 2012. Tetrahydro iso-alpha acids from hops improve glucose homeostasis and reduce body weight gain and metabolic

- endotoxemia in high-fat diet-fed mice. *PLoS One* 7 (3):e33858. doi: [10.1371/journal.pone.0033858](https://doi.org/10.1371/journal.pone.0033858).
- Fang, S., J. M. Suh, S. M. Reilly, E. Yu, O. Osborn, D. Lackey, E. Yoshihara, A. Perino, S. Jacinto, Y. Lukasheva, et al. 2015. Intestinal FXR agonism promotes adipose tissue browning and reduces obesity and insulin resistance. *Nature Medicine* 21 (2):159–65. doi: [10.1038/nm.3760](https://doi.org/10.1038/nm.3760).
- Farquhar, M. G., and G. E. Palade. 1963. Junctional complexes in various epithelia. *The Journal of Cell Biology* 17:375–412. doi: [10.1083/jcb.17.2.375](https://doi.org/10.1083/jcb.17.2.375).
- Festa, M., A. Capasso, C. W. D'Acunto, M. Masullo, A. G. Rossi, C. Pizza, and S. Piacente. 2011. Xanthohumol induces apoptosis in human malignant glioblastoma cells by increasing reactive oxygen species and activating MAPK pathways. *Journal of Natural Products* 74 (12):2505–13. doi: [10.1021/np200390x](https://doi.org/10.1021/np200390x).
- Fiesel, A., D. K. Gessner, E. Most, and K. Eder. 2014. Effects of dietary polyphenol-rich plant products from grape or hop on pro-inflammatory gene expression in the intestine, nutrient digestibility and faecal microbiota of weaned pigs. *BMC Veterinary Research* 10 (1):196. doi: [10.1186/s12917-014-0196-5](https://doi.org/10.1186/s12917-014-0196-5).
- Furukawa, S., T. Fujita, M. Shimabukuro, M. Iwaki, Y. Yamada, Y. Nakajima, O. Nakayama, M. Makishima, M. Matsuda, and I. Shimomura. 2004. Increased oxidative stress in obesity and its impact on metabolic syndrome. *The Journal of Clinical Investigation* 114 (12):1752–61. doi: [10.1172/JCI21625](https://doi.org/10.1172/JCI21625).
- Gallo, C., K. Dallaglio, B. Bassani, T. Rossi, A. Rossello, D. M. Noonan, G. D'Uva, A. Bruno, and A. Albini. 2016. Hop derived flavonoid xanthohumol inhibits endothelial cell functions via AMPK activation. *Oncotarget* 7 (37):59917–31. doi: [10.18632/oncotarget.10990](https://doi.org/10.18632/oncotarget.10990).
- Gerhauser, C., A. Alt, E. Heiss, A. Gamal-Eldeen, K. Klimo, J. Knauff, I. Neumann, H.-R. Scherf, N. Frank, H. Bartsch, et al. 2002. Cancer chemopreventive activity of xanthohumol, a natural product derived from hop. *Molecular Cancer Therapeutics* 1 (11):959–69.
- Gorjanović, S., F. T. Pastor, R. Vasić, M. Novaković, M. Simonović, S. Milić, and D. Sužnjević. 2013. Electrochemical versus spectrophotometric assessment of antioxidant activity of hop (*Humulus lupulus* L.) products and individual compounds. *Journal of Agricultural and Food Chemistry* 61 (38):9089–96. doi: [10.1021/jf401718z](https://doi.org/10.1021/jf401718z).
- Hanske, L., R. Hussong, N. Frank, C. Gerhäuser, M. Blaut, and A. Braune. 2005. Xanthohumol does not affect the composition of rat intestinal microbiota. *Molecular Nutrition & Food Research* 49 (9): 868–73. doi: [10.1002/mnfr.200500048](https://doi.org/10.1002/mnfr.200500048).
- Hanske, L., G. Loh, S. Szczesny, M. Blaut, and A. Braune. 2010. Recovery and metabolism of xanthohumol in germ-free and human microbiota-associated rats. *Molecular Nutrition & Food Research* 54 (10):1405–13. doi: [10.1002/mnfr.200900517](https://doi.org/10.1002/mnfr.200900517).
- Hansson, G. K., and P. Libby. 2006. The immune response in atherosclerosis: A double-edged sword. *Nature Reviews Immunology* 6 (7): 508–19. doi: [10.1038/nri1882](https://doi.org/10.1038/nri1882).
- Harikumar, K. B., A. B. Kunnumakkara, K. S. Ahn, P. Anand, S. Krishnan, S. Guha, and B. B. Aggarwal. 2009. Modification of the cysteine residues in IkappaBalpha kinase and NF-kappaB (p65) by xanthohumol leads to suppression of NF-kappaB-regulated gene products and potentiation of apoptosis in leukemia cells. *Blood* 113 (9):2003–13. doi: [10.1182/blood-2008-04-151944](https://doi.org/10.1182/blood-2008-04-151944).
- Henderson, M. C., C. L. Miranda, J. F. Stevens, M. L. Deinzer, and D. R. Buhler. 2000. In vitro inhibition of human P450 enzymes by prenylated flavonoids from hops, *Humulus lupulus*. *Xenobiotica* 30 (3):235–51. doi: [10.1080/004982500237631](https://doi.org/10.1080/004982500237631).
- Herrington, W., B. Lacey, P. Sherliker, J. Armitage, and S. Lewington. 2016. Epidemiology of atherosclerosis and the potential to reduce the global burden of atherothrombotic disease. *Circulation Research* 118 (4):535–46. doi: [10.1161/CIRCRESAHA.115.307611](https://doi.org/10.1161/CIRCRESAHA.115.307611).
- Hirata, H., K. Takazumi, S. Segawa, Y. Okada, N. Kobayashi, T. Shigyo, and H. Chiba. 2012a. Xanthohumol, a prenylated chalcone from *Humulus lupulus* L., inhibits cholesteryl ester transfer protein. *Food Chemistry* 134 (3):1432–7. doi: [10.1016/j.foodchem.2012.03.043](https://doi.org/10.1016/j.foodchem.2012.03.043).
- Hirata, H., Y. Yimin, Segawa, S. Ozaki, M. Kobayashi, N. Shigyo, T. and Chiba. H. 2012b. Xanthohumol prevents atherosclerosis by reducing arterial cholesterol content via CETP and apolipoprotein E in CETP-transgenic mice. *PLoS One* 7:e49415. doi: [10.1371/journal.pone.0049415](https://doi.org/10.1371/journal.pone.0049415).
- Jaskula, B., P. Kafarski, G. Aerts, and L. De Cooman. 2008. A kinetic study on the isomerization of hop alpha-acids. *Journal of Agricultural and Food Chemistry* 56 (15):6408–15. doi: [10.1021/jf8004965](https://doi.org/10.1021/jf8004965).
- Karabın, M., L. Jelínek, T. Kinčl, T. Hudcová, B. Kotlíková, and P. Dostálek. 2013. New approach to the production of xanthohumol-enriched beers. *Journal of the Institute of Brewing* 119:98–102.
- Kim, J.-A., Y. Kang, D. Thapa, J. S. Lee, M. A. Park, K. H. Lee, W. S. Lyoo, and Y. R. Lee. 2009. Anti-invasive and anti-angiogenic effects of xanthohumol and its synthetic derivatives. *Biomolecules and Therapeutics* 17 (4):422–9. doi: [10.4062/biomolther.2009.17.4.422](https://doi.org/10.4062/biomolther.2009.17.4.422).
- Kim, K. J., B. Li, J. Winer, M. Armanini, N. Gillett, H. S. Phillips, and N. Ferrara. 1993. Inhibition of vascular endothelial growth factor-induced angiogenesis suppresses tumour growth in vivo. *Nature* 362 (6423):841–4. doi: [10.1038/362841a0](https://doi.org/10.1038/362841a0).
- Kim, S. Y., I. S. Lee, and A. Moon. 2013. 2-Hydroxychalcone and xanthohumol inhibit invasion of triple negative breast cancer cells. *Chemico-Biological Interactions* 203 (3):565–72. doi: [10.1016/j.cbi.2013.03.012](https://doi.org/10.1016/j.cbi.2013.03.012).
- Kirkwood, J. S., L. L. Legette, C. L. Miranda, Y. Jiang, and J. F. Stevens. 2013. A metabolomics-driven elucidation of the anti-obesity mechanisms of xanthohumol. *The Journal of Biological Chemistry* 288 (26):19000–13. doi: [10.1074/jbc.M112.445452](https://doi.org/10.1074/jbc.M112.445452).
- Konda, V. R., A. Desai, G. Darland, J. S. Bland, and M. L. Tripp. 2009. Rho iso-alpha acids from hops inhibit the GSK-3/NF-kappaB pathway and reduce inflammatory markers associated with bone and cartilage degradation. *Journal of Inflammation (London, England)* 6: 26. doi: [10.1186/1476-9255-6-26](https://doi.org/10.1186/1476-9255-6-26).
- Kotecha, R., A. Takami, and J. L. Espinoza. 2016. Dietary phytochemicals and cancer chemoprevention: A review of the clinical evidence. *Oncotarget* 7 (32):52517–29. doi: [10.18632/oncotarget.9593](https://doi.org/10.18632/oncotarget.9593).
- König, J., J. Wells, P. D. Cani, C. L. García-Ródenas, T. MacDonald, A. Mercenier, J. Whyte, F. Troost, and R.-J. Brummer. 2016. Human intestinal barrier function in health and disease. *Clinical and Translational Gastroenterology* 7 (10):e196–e196. doi: [10.1038/ctg.2016.54](https://doi.org/10.1038/ctg.2016.54).
- Lagouge, M., C. Argmann, Z. Gerhart-Hines, H. Meziane, C. Lerin, F. Daussin, N. Messadeq, J. Milne, P. Lambert, P. Elliott, et al. 2006. Resveratrol improves mitochondrial function and protects against metabolic disease by activating SIRT1 and PGC-1alpha. *Cell* 127 (6): 1109–22. doi: [10.1016/j.cell.2006.11.013](https://doi.org/10.1016/j.cell.2006.11.013).
- Lamy, V., S. Roussi, M. Chaabi, F. Gossé, A. Lobstein, and F. Raul. 2008. Lupulone, a hop bitter acid, activates different death pathways involving apoptotic TRAIL-receptors, in human colon tumor cells and in their derived metastatic cells. *Apoptosis* 13 (10):1232–42. doi: [10.1007/s10495-008-0250-5](https://doi.org/10.1007/s10495-008-0250-5).
- Lamy, V., S. Roussi, M. Chaabi, F. Gossé, N. Schall, A. Lobstein, and F. Raul. 2007. Chemopreventive effects of lupulone, a hop β -acid, on human colon cancer-derived metastatic SW620 cells and in a rat model of colon carcinogenesis. *Carcinogenesis* 28 (7):1575–81. doi: [10.1093/carcin/bgm080](https://doi.org/10.1093/carcin/bgm080).
- Lee, I.-S., J. Lim, J. Gal, J. C. Kang, H. J. Kim, B. Y. Kang, and H. J. Choi. 2011. Anti-inflammatory activity of xanthohumol involves heme oxygenase-1 induction via NRF2-ARE signaling in microglial BV2 cells. *Neurochemistry International* 58 (2):153–60. doi: [10.1016/j.neuint.2010.11.008](https://doi.org/10.1016/j.neuint.2010.11.008).
- Lee, J., N. Koo, and D. B. Min. 2004. Reactive oxygen species, aging, and antioxidative nutraceuticals. *Comprehensive Reviews in Food Science and Food Safety* 3 (1):21–33. doi: [10.1111/j.1541-4337.2004.tb00058.x](https://doi.org/10.1111/j.1541-4337.2004.tb00058.x).
- Lee, J.-C., J. K. Kundu, D.-M. Hwang, H.-K. Na, and Y.-J. Surh. 2007. Humulone inhibits phorbol ester-induced COX-2 expression in mouse skin by blocking activation of NF-kappaB and AP-1: IkappaB kinase and c-Jun-N-terminal kinase as respective potential upstream targets. *Carcinogenesis* 28 (7):1491–8. doi: [10.1093/carcin/bgm054](https://doi.org/10.1093/carcin/bgm054).
- Lee, M. S., Y. Shin, S. Jung, and Y. Kim. 2017. Effects of epigallocatechin-3-gallate on thermogenesis and mitochondrial biogenesis in

- brown adipose tissues of diet-induced obese mice. *Food & Nutrition Research* 61 (1):1325307. doi: [10.1080/16546628.2017.1325307](https://doi.org/10.1080/16546628.2017.1325307).
- Legette, L., C. Karnpracha, R. L. Reed, J. Choi, G. Bobe, J. M. Christensen, R. Rodriguez-Proteau, J. Q. Purnell, and J. F. Stevens. 2014. Human pharmacokinetics of xanthohumol, an antihyperglycemic flavonoid from hops. *Molecular Nutrition & Food Research* 58 (2):248–55. doi: [10.1002/mnfr.201300333](https://doi.org/10.1002/mnfr.201300333).
- Legette, L., L. Ma, R. L. Reed, C. L. Miranda, J. M. Christensen, R. Rodriguez-Proteau, and J. F. Stevens. 2012. Pharmacokinetics of xanthohumol and metabolites in rats after oral and intravenous administration. *Molecular Nutrition & Food Research* 56 (3):466–74. doi: [10.1002/mnfr.201100554](https://doi.org/10.1002/mnfr.201100554).
- Legette, L. L., A. Y. Moreno Luna, R. L. Reed, C. L. Miranda, G. Bobe, R. R. Proteau, and J. F. Stevens. 2013. Xanthohumol lowers body weight and fasting plasma glucose in obese male Zucker fa/fa rats. *Phytochemistry* 91:236–41. doi: [10.1016/j.phytochem.2012.04.018](https://doi.org/10.1016/j.phytochem.2012.04.018).
- Li, B., L. A. Nolte, J. S. Ju, D. H. Han, T. Coleman, J. O. Holloszy, and C. F. Semenkovich. 2000. Skeletal muscle respiratory uncoupling prevents diet-induced obesity and insulin resistance in mice. *Nature Medicine* 6 (10):1115–20. doi: [10.1038/80450](https://doi.org/10.1038/80450).
- Liu, M., H. Yin, G. Liu, J. Dong, Z. Qian, and J. Miao. 2014. Xanthohumol, a prenylated chalcone from beer hops, acts as an α -glucosidase inhibitor in vitro. *Journal of Agricultural and Food Chemistry* 62 (24):5548–54. doi: [10.1021/jf500426z](https://doi.org/10.1021/jf500426z).
- Liu, T., L. Zhang, D. Joo, and S.-C. Sun. 2017. NF- κ B signaling in inflammation. *Signal Transduction and Targeted Therapy* 2:17023.
- Luescher, S., C. Urmann, and V. Butterweck. 2017. Effect of hops derived prenylated phenols on TNF- α induced barrier dysfunction in intestinal epithelial cells. *Journal of Natural Products* 80 (4): 925–31. doi: [10.1021/acs.jnatprod.6b00869](https://doi.org/10.1021/acs.jnatprod.6b00869).
- Lumeng, C. N., J. L. Bodzin, and A. R. Saltiel. 2007. Obesity induces a phenotypic switch in adipose tissue macrophage polarization. *The Journal of Clinical Investigation* 117 (1):175–84. doi: [10.1172/JCI29881](https://doi.org/10.1172/JCI29881).
- Lupinacci, E., J. Meijerink, J.-P. Vincken, B. Gabriele, H. Gruppen, and R. F. Witkamp. 2009. Xanthohumol from hop (*Humulus lupulus* L.) is an efficient inhibitor of monocyte chemoattractant protein-1 and tumor necrosis factor- α release in LPS-stimulated RAW 264.7 mouse macrophages and U937 human monocytes. *Journal of Agricultural and Food Chemistry* 57 (16):7274–81. doi: [10.1021/jf901244k](https://doi.org/10.1021/jf901244k).
- Lv, H., Q. Liu, Z. Wen, H. Feng, X. Deng, and X. Ci. 2017. Xanthohumol ameliorates lipopolysaccharide (LPS)-induced acute lung injury via induction of AMPK/GSK3 β -Nrf2 signal axis. *Redox Biology* 12:311–24. doi: [10.1016/j.redox.2017.03.001](https://doi.org/10.1016/j.redox.2017.03.001).
- Ma, H., and M. E. Patti. 2014. Bile acids, obesity, and the metabolic syndrome. *Best Practice & Research. Clinical Gastroenterology* 28 (4): 573–83. doi: [10.1016/j.bpg.2014.07.004](https://doi.org/10.1016/j.bpg.2014.07.004).
- Mannering, G. J., J. A. Shoeman, and L. B. Deloria. 1992. Identification of the antibiotic hops component, colupulone, as an inducer of hepatic cytochrome P-4503A in the mouse. *Drug Metabolism and Disposition: The Biological Fate of Chemicals* 20 (2):142–7.
- Marchiando, A. M., L. Shen, W. V. Graham, K. L. Edelblum, C. A. Duckworth, Y. Guan, M. H. Montrose, J. R. Turner, and A. J. Watson. 2011. The epithelial barrier is maintained by in vivo tight junction expansion during pathologic intestinal epithelial shedding. *Gastroenterology* 140 (4):1208–18. doi: [10.1053/j.gastro.2011.01.004](https://doi.org/10.1053/j.gastro.2011.01.004).
- Masson, V., L. R. de la Ballina, C. Munaut, B. Wielockx, M. Jost, C. Maillard, S. Blacher, K. Bajou, T. Itoh, S. Itoharu, et al. 2005. Contribution of host MMP-2 and MMP-9 to promote tumor vascularization and invasion of malignant keratinocytes. *FASEB Journal* 19 (2):234–6. doi: [10.1096/fj.04-2140fe](https://doi.org/10.1096/fj.04-2140fe).
- Maukonen, J., K.-L. Kolho, M. Paasela, J. Honkanen, P. Klemetti, O. Vaarala, and M. Saarela. 2015. Altered fecal microbiota in paediatric inflammatory bowel disease. *Journal of Crohn's and Colitis* 9 (12): 1088–95. doi: [10.1093/ecco-jcc/jjv147](https://doi.org/10.1093/ecco-jcc/jjv147).
- Mazmanian, S. K., C. H. Liu, A. O. Tzianabos, and D. L. Kasper. 2005. An immunomodulatory molecule of symbiotic bacteria directs maturation of the host immune system. *Cell* 122 (1):107–18. doi: [10.1016/j.cell.2005.05.007](https://doi.org/10.1016/j.cell.2005.05.007).
- Milligan, S. R., J. C. Kalita, V. Pocock, V. Van De Kauter, J. F. Stevens, M. L. Deinzer, H. Rong, and D. De Keukeleire. 2000. The endocrine activities of 8-prenylnaringenin and related hop (*Humulus lupulus* L.) flavonoids. *The Journal of Clinical Endocrinology and Metabolism* 85 (12):4912–15. doi: [10.1210/jcem.85.12.7168](https://doi.org/10.1210/jcem.85.12.7168).
- Miranda, C. L., G. L. Aponso, J. F. Stevens, M. L. Deinzer, and D. R. Buhler. 2000. Prenylated chalcones and flavanones as inducers of quinone reductase in mouse Hepa 1c1c7 cells. *Cancer Letters* 149 (1–2):21–9. doi: [10.1016/S0304-3835\(99\)00328-6](https://doi.org/10.1016/S0304-3835(99)00328-6).
- Miranda, C. L., V. D. Elias, J. J. Hay, J. Choi, R. L. Reed, and J. F. Stevens. 2016. Xanthohumol improves dysfunctional glucose and lipid metabolism in diet-induced obese C57BL/6J mice. *Archives of Biochemistry and Biophysics* 599:22–30. doi: [10.1016/j.abb.2016.03.008](https://doi.org/10.1016/j.abb.2016.03.008).
- Miranda, C. L., J. F. Stevens, A. Helmrich, M. C. Henderson, R. J. Rodriguez, Y. H. Yang, M. L. Deinzer, D. W. Barnes, and D. R. Buhler. 1999. Antiproliferative and cytotoxic effects of prenylated flavonoids from hops (*Humulus lupulus*) in human cancer cell lines. *Food and Chemical Toxicology* 37 (4):271–85. doi: [10.1016/S0278-6915\(99\)00019-8](https://doi.org/10.1016/S0278-6915(99)00019-8).
- Miura, Y., M. Hosono, C. Oyamada, H. Odai, S. Oikawa, and K. Kondo. 2005. Dietary isohumulones, the bitter components of beer, raise plasma HDL-cholesterol levels and reduce liver cholesterol and triacylglycerol contents similar to PPAR α activations in C57BL/6 mice. *British Journal of Nutrition* 93 (4):559–67. doi: [10.1079/BJN20041384](https://doi.org/10.1079/BJN20041384).
- Monteiro, R., C. Calhau, A. O. Silva, S. Pinheiro-Silva, S. Guerreiro, F. Gärtner, I. Azevedo, and R. Soares. 2008. Xanthohumol inhibits inflammatory factor production and angiogenesis in breast cancer xenografts. *Journal of Cellular Biochemistry* 104 (5):1699–707. doi: [10.1002/jcb.21738](https://doi.org/10.1002/jcb.21738).
- Morimoto-Kobayashi, Y., K. Ohara, H. Ashigai, T. Kanaya, K. Koizumi, F. Manabe, Y. Kaneko, Y. Taniguchi, M. Katayama, Y. Kowatari, et al. 2015. Matured hop extract reduces body fat in healthy overweight humans: A randomized, double-blind, placebo-controlled parallel group study. *Nutrition Journal* 15 (1):25. doi: [10.1186/s12937-016-0144-2](https://doi.org/10.1186/s12937-016-0144-2).
- Morimoto-Kobayashi, Y., K. Ohara, C. Takahashi, S. Kitao, G. Wang, Y. Taniguchi, M. Katayama, and K. Nagai. 2015. Matured hop bittering components induce thermogenesis in brown adipose tissue via sympathetic nerve activity. *PLoS One* 10 (6):e0131042. doi: [10.1371/journal.pone.0131042](https://doi.org/10.1371/journal.pone.0131042).
- Murakami, A., P. Darby, B. Javornik, M. S. Pais, E. Seigner, A. Lutz, and P. Svoboda. 2006. Molecular phylogeny of wild hops, *Humulus lupulus* L. *Heredity* 97 (1):66–74. doi: [10.1038/sj.hdy.6800839](https://doi.org/10.1038/sj.hdy.6800839).
- Murri, M., I. Leiva, J. M. Gomez-Zumaquero, F. J. Tinahones, F. Cardona, F. Soriguer, and M. I. Queipo-Ortuño. 2013. Gut microbiota in children with type 1 diabetes differs from that in healthy children: A case-control study. *BMC Medicine* 11:46. doi: [10.1186/1741-7015-11-46](https://doi.org/10.1186/1741-7015-11-46).
- Musaiger, A. O., and H. M. Al-Hazzaa. 2012. Prevalence and risk factors associated with nutrition-related noncommunicable diseases in the Eastern Mediterranean region. *International Journal of General Medicine* 5:199–217. doi: [10.2147/IJGM.S29663](https://doi.org/10.2147/IJGM.S29663).
- Namikoshi, T., N. Tomita, S. Fujimoto, Y. Haruna, M. Ohzeki, N. Komai, T. Sasaki, A. Yoshida, and N. Kashiara. 2007. Isohumulones derived from hops ameliorate renal injury via an anti-oxidative effect in dahl salt-sensitive rats. *Hypertension Research* 30 (2):175–84. doi: [10.1291/hypres.30.175](https://doi.org/10.1291/hypres.30.175).
- Naya, Y., and M. Kotake. 1971. The constituents of hops. VI studies of the volatile composition of *Humulus lupulus* L. during ripening. *Bulletin of the Chemical Society of Japan* 44 (11):3116–20. doi: [10.1246/bcsj.44.3116](https://doi.org/10.1246/bcsj.44.3116).
- Noda, S., S. Tanabe, and T. Suzuki. 2013. Naringenin enhances intestinal barrier function through the expression and cytoskeletal association of tight junction proteins in Caco-2 cells. *Molecular Nutrition & Food Research* 57 (11):2019–28. doi: [10.1002/mnfr.201300045](https://doi.org/10.1002/mnfr.201300045).
- Nowakowska, Z. 2007. A review of anti-infective and anti-inflammatory chalcones. *European Journal of Medicinal Chemistry*. 42 (2):125–37. doi: [10.1016/j.ejmech.2006.09.019](https://doi.org/10.1016/j.ejmech.2006.09.019).

- Nozawa, H. 2005. Xanthohumol, the chalcone from beer hops (*Humulus lupulus* L.), is the ligand for farnesoid X receptor and ameliorates lipid and glucose metabolism in KK-A(y) mice. *Biochemical and Biophysical Research Communications* 336 (3): 754–61. doi: [10.1016/j.bbrc.2005.08.159](https://doi.org/10.1016/j.bbrc.2005.08.159).
- Nozawa, H., W. Nakao, F. Zhao, and K. Kondo. 2005. Dietary supplement of isohumulones inhibits the formation of aberrant crypt foci with a concomitant decrease in prostaglandin E2 level in rat colon. *Molecular Nutrition & Food Research* 49 (8):772–8. doi: [10.1002/mnfr.200500027](https://doi.org/10.1002/mnfr.200500027).
- Obara, K., M. Mizutani, Y. Hitomi, H. Yajima, and K. Kondo. 2009. Isohumulones, the bitter component of beer, improve hyperglycemia and decrease body fat in Japanese subjects with prediabetes. *Clinical Nutrition (Edinburgh, Scotland)* 28 (3):278–84. doi: [10.1016/j.clnu.2009.03.012](https://doi.org/10.1016/j.clnu.2009.03.012).
- Odenwald, M. A., and J. R. Turner. 2017. The intestinal epithelial barrier: A therapeutic target? *Nature Reviews. Gastroenterology & Hepatology* 14 (1):9–21. doi: [10.1038/nrgastro.2016.169](https://doi.org/10.1038/nrgastro.2016.169).
- Pan, L., H. Becker, and C. Gerhäuser. 2005. Xanthohumol induces apoptosis in cultured 40–16 human colon cancer cells by activation of the death receptor- and mitochondrial pathway. *Molecular Nutrition & Food Research* 49 (9):837–43. doi: [10.1002/mnfr.200500065](https://doi.org/10.1002/mnfr.200500065).
- Pang, Y., D. Nikolic, D. Zhu, L. R. Chadwick, G. F. Pauli, N. R. Farnsworth, and R. Bv. Breemen. 2007. Binding of the hop (*Humulus lupulus* L.) chalcone xanthohumol to cytosolic proteins in Caco-2 intestinal epithelial cells. *Molecular Nutrition & Food Research* 51 (7):872–9. doi: [10.1002/mnfr.200600252](https://doi.org/10.1002/mnfr.200600252).
- Paraiso, I. L., L. S. Plagmann, L. Yang, R. Zielke, A. F. Gombart, C. S. Maier, A. E. Sikora, P. R. Blakemore, and J. F. Stevens. 2019. Reductive metabolism of xanthohumol and 8-Prenylnaringenin by the intestinal bacterium *Eubacterium ramulus*. *Molecular Nutrition & Food Research* 63 (2):e1800923. doi: [10.1002/mnfr.201800923](https://doi.org/10.1002/mnfr.201800923).
- Peery, A. F., S. D. Crockett, C. C. Murphy, J. L. Lund, E. S. Dellon, J. L. Williams, E. T. Jensen, N. J. Shaheen, A. S. Barritt, S. R. Lieber, et al. 2019. Burden and cost of gastrointestinal, liver, and pancreatic diseases in the United States: Update 2018. *Gastroenterology* 156 (1): 254–72. doi: [10.1053/j.gastro.2018.08.063](https://doi.org/10.1053/j.gastro.2018.08.063).
- Pisoschi, A. M., and A. Pop. 2015. The role of antioxidants in the chemistry of oxidative stress: A review. *European Journal of Medicinal Chemistry* 97:55–74. doi: [10.1016/j.ejmech.2015.04.040](https://doi.org/10.1016/j.ejmech.2015.04.040).
- Plazar, J., B. Zegura, T. T. Lah, and M. Filipic. 2007. Protective effects of xanthohumol against the genotoxicity of benzo(a)pyrene (BaP), 2-amino-3-methylimidazo[4,5-f]quinoline (IQ) and tert-butyl hydroperoxide (t-BOOH) in HepG2 human hepatoma cells. *Mutation Research* 632 (1–2):1–8. doi: [10.1016/j.mrgentox.2007.03.013](https://doi.org/10.1016/j.mrgentox.2007.03.013).
- Popkin, B. M. 2002. An overview on the nutrition transition and its health implications: The Bellagio meeting. *Public Health Nutrition* 5 (1A):93–103. doi: [10.1079/phn2001280](https://doi.org/10.1079/phn2001280).
- Popkin, B. M., and P. Gordon-Larsen. 2004. The nutrition transition: Worldwide obesity dynamics and their determinants. *International Journal of Obesity and Related Metabolic Disorders* 28 (Suppl 3): S2–S9. doi: [10.1038/sj.ijo.0802804](https://doi.org/10.1038/sj.ijo.0802804).
- Possemiers, S., A. Heyerick, V. Robbens, D. De Keukeleire, and W. Verstraete. 2005. Activation of proestrogens from hops (*Humulus lupulus* L.) by intestinal microbiota; conversion of isoxanthohumol into 8-prenylnaringenin. *Journal of Agricultural and Food Chemistry* 53 (16):6281–8. doi: [10.1021/jf0509714](https://doi.org/10.1021/jf0509714).
- Ramos, S. 2008. Cancer chemoprevention and chemotherapy: Dietary polyphenols and signalling pathways. *Molecular Nutrition & Food Research* 52 (5):507–26. doi: [10.1002/mnfr.200700326](https://doi.org/10.1002/mnfr.200700326).
- Rancán, L., S. D. Paredes, I. García, P. Muñoz, C. García, G. López de Hontanar, M. de la Fuente, E. Vara, and J. A. F. Tresguerres. 2017. Protective effect of xanthohumol against age-related brain damage. *Journal of Nutritional Biochemistry*. 49:133–40. doi: [10.1016/j.jnutbio.2017.07.011](https://doi.org/10.1016/j.jnutbio.2017.07.011).
- Saito, K., Y. Matsuo, H. Imafuji, T. Okubo, Y. Maeda, T. Sato, T. Shamoto, K. Tsuboi, M. Morimoto, H. Takahashi, et al. 2018. Xanthohumol inhibits angiogenesis by suppressing nuclear factor- κ B activation in pancreatic cancer. *Cancer Science* 109 (1):132–40. doi: [10.1111/cas.13441](https://doi.org/10.1111/cas.13441).
- Saklayen, M. G. 2018. The global epidemic of the metabolic syndrome. *Current Hypertension Reports* 20 (2):12. doi: [10.1007/s11906-018-0812-z](https://doi.org/10.1007/s11906-018-0812-z).
- Samuels, J. S., R. Shashidharamurthy, and S. Rayalam. 2018. Novel anti-obesity effects of beer hops compound xanthohumol: Role of AMPK signaling pathway. *Nutrition & Metabolism (London)* 15:42. doi: [10.1186/s12986-018-0277-8](https://doi.org/10.1186/s12986-018-0277-8).
- Shimada, T., C. L. Hayes, H. Yamazaki, S. Amin, S. S. Hecht, F. P. Guengerich, and T. R. Sutter. 1996. Activation of chemically diverse procarcinogens by human cytochrome P-450 1B1. *Cancer Research* 56 (13):2979–84.
- Shimamura, M., T. Hazato, H. Ashino, Y. Yamamoto, E. Iwasaki, H. Tobe, K. Yamamoto, and S. Yamamoto. 2001. Inhibition of angiogenesis by humulone, a bitter acid from beer hop. *Biochemical and Biophysical Research Communications* 289 (1):220–4. doi: [10.1006/bbrc.2001.5934](https://doi.org/10.1006/bbrc.2001.5934).
- Siegel, L., A. Miternique-Grosse, C. Griffon, C. Klein-Soyer, A. Lobstein, F. Raul, and D. Stephan. 2008. Antiangiogenic properties of lupulone, a bitter acid of hop cones. *Anticancer Research* 28 (1A): 289–94.
- Small, E. 1978. A numerical and nomenclatural analysis of morphogeographic taxa of *Humulus*. *Systematic Botany* 3 (1):37–76. doi: [10.2307/2418532](https://doi.org/10.2307/2418532).
- Stevens, J. F., and J. E. Page. 2004. Xanthohumol and related prenylflavonoids from hops and beer: To your good health! *Phytochemistry* 65 (10):1317–30. doi: [10.1016/j.phytochem.2004.04.025](https://doi.org/10.1016/j.phytochem.2004.04.025).
- Stevens, J. F., A. W. Taylor, J. E. Clawson, and M. L. Deinzer. 1999. Fate of xanthohumol and related prenylflavonoids from hops to beer. *Journal of Agricultural and Food Chemistry* 47 (6):2421–8. doi: [10.1021/jf990101k](https://doi.org/10.1021/jf990101k).
- Sun, R., N. Yang, B. Kong, B. Cao, D. Feng, X. Yu, C. Ge, J. Huang, J. Shen, P. Wang, et al. 2017. Orally administered berberine modulates hepatic lipid metabolism by altering microbial bile acid metabolism and the intestinal FXR signaling pathway. *Molecular Pharmacology* 91 (2):110–22. doi: [10.1124/mol.116.106617](https://doi.org/10.1124/mol.116.106617).
- Sun, X., M. Du, D. A. Navarre, and M.-J. Zhu. 2018. Purple potato extract promotes intestinal epithelial differentiation and barrier function by activating AMP-activated protein kinase. *Molecular Nutrition & Food Research* 62 (4):1700536. doi: [10.1002/mnfr.201700536](https://doi.org/10.1002/mnfr.201700536).
- Surh, Y.-J. 2003. Cancer chemoprevention with dietary phytochemicals. *Nature Reviews Cancer* 3 (10):768–80. doi: [10.1038/nrc1189](https://doi.org/10.1038/nrc1189).
- Suzuki, T., and H. Hara. 2009. Quercetin enhances intestinal barrier function through the assembly of zonula [corrected] occludens-2, occludin, and claudin-1 and the expression of claudin-4 in Caco-2 cells. *The Journal of Nutrition* 139 (5):965–74. doi: [10.3945/jn.108.100867](https://doi.org/10.3945/jn.108.100867).
- Suzuki, T., S. Tanabe, and H. Hara. 2011. Kaempferol enhances intestinal barrier function through the cytoskeletal association and expression of tight junction proteins in Caco-2 cells. *The Journal of Nutrition* 141 (1):87–94. doi: [10.3945/jn.110.125633](https://doi.org/10.3945/jn.110.125633).
- Tagashira, M., M. Watanabe, and N. Uemitsu. 1995. Antioxidative activity of hop bitter acids and their analogues. *Bioscience, Biotechnology, and Biochemistry* 59 (4):740–2. doi: [10.1271/bbb.59.740](https://doi.org/10.1271/bbb.59.740).
- Tang, W., L.-H. Chen, H. Daun, C.-T. Ho, and M.-H. Pan. 2011. Inhibitory effects of hexahydro- β -acids in LPS-stimulated murine macrophage. *Journal of Functional Foods*. 3 (3):215–22. doi: [10.1016/j.jff.2011.04.004](https://doi.org/10.1016/j.jff.2011.04.004).
- Tillman, G. E., G. J. Haas, M. G. Wise, B. Oakley, M. A. Smith, and G. R. Siragusa. 2011. Chicken intestine microbiota following the administration of lupulone, a hop-based antimicrobial. *FEMS Microbiology Ecology* 77 (2):395–403. doi: [10.1111/j.1574-6941.2011.01119.x](https://doi.org/10.1111/j.1574-6941.2011.01119.x).
- Valko, M., D. Leibfritz, J. Moncol, M. T. Cronin, M. Mazur, and J. Telser. 2007. Free radicals and antioxidants in normal physiological functions and human disease. *The International Journal of*

- Biochemistry & Cell Biology* 39 (1):44–84. doi: [10.1016/j.biocel.2006.07.001](https://doi.org/10.1016/j.biocel.2006.07.001).
- Van Cleemput, M., A. Heyerick, C. Libert, K. Swerts, J. Philippé, D. D. Keukeleire, G. Haegeman, and K. D. Bosscher. 2009. Hop bitter acids efficiently block inflammation independent of GR α , PPAR α , or PPAR γ . *Molecular Nutrition & Food Research* 53 (9):1143–55. doi: [10.1002/mnfr.200800493](https://doi.org/10.1002/mnfr.200800493).
- van Marken Lichtenbelt, W. D., J. W. Vanhommerig, N. M. Smulders, J. M. Drossaerts, G. J. Kemerink, N. D. Bouvy, P. Schrauwen, and G. J. Teule. 2009. Cold-activated brown adipose tissue in healthy men. *New England Journal of Medicine* 360 (15):1500–8. doi: [10.1056/NEJMoa0808718](https://doi.org/10.1056/NEJMoa0808718).
- Vanhoeck, B., L. Derycke, V. V. Marck, H. Depypere, D. D. Keukeleire, and M. Bracke. 2005. Antiinvasive effect of xanthohumol, a prenylated chalcone present in hops (*Humulus lupulus* L.) and beer. *International Journal of Cancer* 117 (6):889–95. doi: [10.1002/ijc.21249](https://doi.org/10.1002/ijc.21249).
- Vetrano, S., M. Rescigno, M. R. Cera, C. Correale, C. Rumio, A. Doni, M. Fantini, A. Sturm, E. Borroni, A. Repici, et al. 2008. Unique role of junctional adhesion molecule-1 in maintaining mucosal homeostasis in inflammatory bowel disease. *Gastroenterology* 135 (1):173–84. doi: [10.1053/j.gastro.2008.04.002](https://doi.org/10.1053/j.gastro.2008.04.002).
- Wang, S., X. Liang, Q. Yang, X. Fu, C. J. Rogers, M. Zhu, B. D. Rodgers, Q. Jiang, M. V. Dodson, and M. Du. 2015. Resveratrol induces brown-like adipocyte formation in white fat through activation of AMP-activated protein kinase (AMPK) α 1. *International Journal of Obesity* (2005) 39 (6):967–76. doi: [10.1038/ijo.2015.23](https://doi.org/10.1038/ijo.2015.23).
- Ward, B. W., J. S. Schiller, and R. A. Goodman. 2014. Multiple chronic conditions among US adults: A 2012 update. *Preventing Chronic Disease* 11:E62. doi: [10.5888/pcd11.130389](https://doi.org/10.5888/pcd11.130389).
- Watanabe, M., S. M. Houten, C. Mataka, M. A. Christoffolete, B. W. Kim, H. Sato, N. Messaddeq, J. W. Harney, O. Ezaki, T. Kodama, et al. 2006. Bile acids induce energy expenditure by promoting intracellular thyroid hormone activation. *Nature* 439 (7075):484–9. doi: [10.1038/nature04330](https://doi.org/10.1038/nature04330).
- Weisberg, S. P., D. McCann, M. Desai, M. Rosenbaum, R. L. Leibel, and A. W. Ferrante. 2003. Obesity is associated with macrophage accumulation in adipose tissue. *The Journal of Clinical Investigation* 112 (12):1796–808. doi: [10.1172/JCI200319246](https://doi.org/10.1172/JCI200319246).
- West, C. E., Renz, H. Jenmalm, M. C. Kozyrskyj, A. L. Allen, K. J. Vuillermin, P. Prescott, S. L., in-FLAME Microbiome Interest Group. 2015. The gut microbiota and inflammatory noncommunicable diseases: Associations and potentials for gut microbiota therapies. *The Journal of Allergy and Clinical Immunology* 135 (1):3–13; quiz 14. doi: [10.1016/j.jaci.2014.11.012](https://doi.org/10.1016/j.jaci.2014.11.012).
- Wu, S. H., W. S. Hui, Z. Liu, and S. C. Ho. 2010. Metabolic syndrome and all-cause mortality: A meta-analysis of prospective cohort studies. *European Journal of Epidemiology* 25 (6):375–84. doi: [10.1007/s10654-010-9459-z](https://doi.org/10.1007/s10654-010-9459-z).
- Xing, T., Y. Kang, X. Xu, B. Wang, M. Du, and M. J. Zhu. 2018. Raspberry supplementation improves insulin signaling and promotes brown-like adipocyte development in white adipose tissue of obese mice. *Molecular Nutrition & Food Research* 62:1701035.
- Xu, H., G. T. Barnes, Q. Yang, G. Tan, D. Yang, C. J. Chou, J. Sole, A. Nichols, J. S. Ross, L. A. Tartaglia, et al. 2003. Chronic inflammation in fat plays a crucial role in the development of obesity-related insulin resistance. *Journal of Clinical Investigation* 112 (12):1821–30. doi: [10.1172/JCI200319451](https://doi.org/10.1172/JCI200319451).
- Xuan, N. T., E. Shumilina, E. Gulbins, S. Gu, F. Götz, and F. Lang. 2010. Triggering of dendritic cell apoptosis by xanthohumol. *Molecular Nutrition & Food Research* 54 (S2):S214–S224. doi: [10.1002/mnfr.200900324](https://doi.org/10.1002/mnfr.200900324).
- Yaffe, K., A. Kanaya, K. Lindquist, E. M. Simonsick, T. Harris, R. I. Shorr, F. A. Tykavsky, and A. B. Newman. 2004. The metabolic syndrome, inflammation, and risk of cognitive decline. *JAMA* 292 (18):2237–42. doi: [10.1001/jama.292.18.2237](https://doi.org/10.1001/jama.292.18.2237).
- Yajima, H., E. Ikeshima, M. Shiraki, T. Kanaya, D. Fujiwara, H. Odai, N. Tsuboyama-Kasaoka, O. Ezaki, S. Oikawa, and K. Kondo. 2004. Isohumulones, bitter acids derived from hops, activate both peroxisome proliferator-activated receptor α and γ and reduce insulin resistance. *The Journal of Biological Chemistry* 279 (32):33456–62. doi: [10.1074/jbc.M403456200](https://doi.org/10.1074/jbc.M403456200).
- Yajima, H., T. Noguchi, E. Ikeshima, M. Shiraki, T. Kanaya, N. Tsuboyama-Kasaoka, O. Ezaki, S. Oikawa, and K. Kondo. 2005. Prevention of diet-induced obesity by dietary isomerized hop extract containing isohumulones, in rodents. *International Journal of Obesity* (2005) 29 (8):991–7. doi: [10.1038/sj.sjo.0802965](https://doi.org/10.1038/sj.sjo.0802965).
- Yamaguchi, N., K. Satoh-Yamaguchi, and M. Ono. 2009. In vitro evaluation of antibacterial, anticollagenase, and antioxidant activities of hop components (*Humulus lupulus*) addressing acne vulgaris. *Phytomedicine* 16 (4):369–76. doi: [10.1016/j.phymed.2008.12.021](https://doi.org/10.1016/j.phymed.2008.12.021).
- Yamamoto, K., J. Wang, S. Yamamoto, and H. Tobe. 2000. Suppression of cyclooxygenase-2 gene transcription by humulon of beer hop extract studied with reference to glucocorticoid. *FEBS Letters* 465 (2–3):103–6. doi: [10.1016/S0014-5793\(99\)01727-5](https://doi.org/10.1016/S0014-5793(99)01727-5).
- Yamazaki, T., Y. Morimoto-Kobayashi, K. Koizumi, C. Takahashi, S. Nakajima, S. Kitao, Y. Taniguchi, M. Katayama, and Y. Ogawa. 2019. Secretion of a gastrointestinal hormone, cholecystokinin, by hop-derived bitter components activates sympathetic nerves in brown adipose tissue. *The Journal of Nutritional Biochemistry* 64:80–7. doi: [10.1016/j.jnutbio.2018.10.009](https://doi.org/10.1016/j.jnutbio.2018.10.009).
- Yoneshiro, T., S. Aita, Y. Kawai, T. Iwanaga, and M. Saito. 2012. Nonpungent capsaicin analogs (capsinoids) increase energy expenditure through the activation of brown adipose tissue in humans. *The American Journal of Clinical Nutrition* 95 (4):845–50. doi: [10.3945/ajcn.111.018606](https://doi.org/10.3945/ajcn.111.018606).
- Yoneshiro, T., S. Aita, M. Matsushita, T. Kayahara, T. Kameya, Y. Kawai, T. Iwanaga, and M. Saito. 2013. Recruited brown adipose tissue as an antiobesity agent in humans. *The Journal of Clinical Investigation* 123 (8):3404–8. doi: [10.1172/JCI67803](https://doi.org/10.1172/JCI67803).
- Yui, K., A. Kiyofuji, and K. Osada. 2014. Effects of xanthohumol-rich extract from the hop on fatty acid metabolism in rats fed a high-fat diet. *Journal of Oleo Science* 63 (2):159–68. doi: [10.5650/jos.ess13136](https://doi.org/10.5650/jos.ess13136).
- Zanoli, P., and M. Zavatti. 2008. Pharmacognostic and pharmacological profile of *Humulus lupulus* L. *Journal of Ethnopharmacology* 116 (3):383–96. doi: [10.1016/j.jep.2008.01.011](https://doi.org/10.1016/j.jep.2008.01.011).
- Zeissig, S., N. Burgel, D. Gunzel, J. Richter, J. Mankertz, U. Wahnschaffe, A. J. Kroesen, M. Zeitz, M. Fromm, and J. D. Schulzke. 2007. Changes in expression and distribution of claudin 2, 5 and 8 lead to discontinuous tight junctions and barrier dysfunction in active Crohn's disease. *Gut* 56 (1):61–72. doi: [10.1136/gut.2006.094375](https://doi.org/10.1136/gut.2006.094375).
- Zhang, X., X. Liang, H. Xiao, and Q. Xu. 2004. Direct characterization of bitter acids in a crude hop extract by liquid chromatography-atmospheric pressure chemical ionization mass spectrometry. *Journal of the American Society for Mass Spectrometry* 15 (2):180–7. doi: [10.1016/j.jasms.2003.09.014](https://doi.org/10.1016/j.jasms.2003.09.014).
- Zhang, Y. J., R. Y. Gan, S. Li, Y. Zhou, A. N. Li, D. P. Xu, and H. B. Li. 2015. Antioxidant phytochemicals for the prevention and treatment of chronic diseases. *Molecules (Basel, Switzerland)* 20 (12):21138–56. doi: [10.3390/molecules201219753](https://doi.org/10.3390/molecules201219753).
- Zhang, Z., H. Zhang, B. Li, X. Meng, J. Wang, Y. Zhang, S. Yao, Q. Ma, L. Jin, J. Yang, et al. 2014. Berberine activates thermogenesis in white and brown adipose tissue. *Nature Communications* 5:5493. doi: [10.1038/ncomms6493](https://doi.org/10.1038/ncomms6493).
- Zhao, F., H. Nozawa, A. Daikonnaya, K. Kondo, and S. Kitanaka. 2003. Inhibitors of nitric oxide production from hops (*Humulus lupulus* L.). *Biological and Pharmaceutical Bulletin* 26 (1):61–5. doi: [10.1248/bpb.26.61](https://doi.org/10.1248/bpb.26.61).