



## Critical Reviews in Food Science and Nutrition

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/bfsn20>

### Analyzing B-Vitamins in Human Milk: Methodological Approaches

Daniela Hampel<sup>a</sup> & Lindsay H. Allen<sup>a</sup>

<sup>a</sup> USDA, ARS Western Human Nutrition Research Center, 430 W. Health Sciences Drive, Davis, CA 95616, USA

Accepted author version posted online: 15 Jan 2015.



[Click for updates](#)

To cite this article: Daniela Hampel & Lindsay H. Allen (2015): Analyzing B-Vitamins in Human Milk: Methodological Approaches, Critical Reviews in Food Science and Nutrition, DOI: [10.1080/10408398.2013.783550](https://doi.org/10.1080/10408398.2013.783550)

To link to this article: <http://dx.doi.org/10.1080/10408398.2013.783550>

Disclaimer: This is a version of an unedited manuscript that has been accepted for publication. As a service to authors and researchers we are providing this version of the accepted manuscript (AM). Copyediting, typesetting, and review of the resulting proof will be undertaken on this manuscript before final publication of the Version of Record (VoR). During production and pre-press, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal relate to this version also.

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

**Analyzing B-Vitamins in Human Milk: Methodological Approaches**

Daniela Hampel and Lindsay H. Allen

USDA, ARS Western Human Nutrition Research Center, 430 W. Health Sciences Drive, Davis, CA 95616, USA

Address correspondence to Daniela Hampel, USDA, ARS Western Human Nutrition Research Center, 430 W. Health Sciences Drive, Davis, CA 95616, USA. E-mail: daniela.hampel@ars.usda.gov

***ABSTRACT***

According to the World Health Organization (WHO) infants should be exclusively breastfed for the first six months of life. However, there is insufficient information about the concentration of nutrients in human milk. For some nutrients, including B-vitamins, maternal intake affects their concentration in human milk but the extent to which inadequate maternal diets affect milk B-vitamin content is poorly documented. Little is known about infant requirements for B-vitamins; recommendations are generally set as Adequate Intakes (AI) calculated based on the mean volume of milk (0.78L/day) consumed by infants exclusively fed with human milk from well-nourished mothers during the first six months, and the concentration of each vitamin in milk based on reported values. Methods used for analyzing B-vitamins, commonly microbiological, radioisotope dilution or more recently chromatographic, coupled with UV, fluorometric and MS detection, have rarely been validated for the complex human milk matrix. Thus the validity, accuracy and sensitivity of analytical methods is important for understanding infant requirements

for these nutrients, the maternal intakes needed to support adequate concentrations in breast milk. This review summarizes current knowledge on methods used for analyzing the B-vitamins thiamin, riboflavin, niacin, vitamin B-6 and pantothenic acid, vitamin B-12, folate, biotin, and choline in human milk, their chemical and physical properties, the different forms and changes in concentration during lactation, and the effects of deficiency on the infant.

***KEYWORDS***

Human milk, infant nutrient requirements, B-vitamin deficiency, microbiological assays, chromatographic analyses

## ***INTRODUCTION***

According to the World Health Organization (WHO) infants should be exclusively breastfed for the first 6 months of life rather than the 4-6 months previously recommended. This advice was based substantially on findings that exclusive breastfeeding has protective effects against gastrointestinal infections (WHO 2001). A breastfed infant's nutritional status, and especially B-vitamin status (Allen 2012) therefore relies on the amounts of the vitamins in breast milk, which are highly influenced by maternal dietary intake. Poor maternal status or inadequate intake of B-vitamins during lactation is usually reflected in lower breast milk concentrations and can subsequently cause infant deficiency.

Lack of reliable data reflecting the infant's response to different levels of nutrient intake inhibits the setting of Estimated Average Requirements (EARs) and therefore the determination of Recommended Dietary Allowances (RDAs). Thus, the infant's recommended intakes of B-vitamins (and other nutrients) are generally set as Adequate Intakes (AI), calculated as the mean intakes by full-term infants exclusively fed with human milk from well-nourished mothers during the first six months (I.O.M, 1998). This calculation is based on the mean concentration of the vitamins from 2 - 6 months of lactation using consensus values from several reported studies, and an average milk intake of 0.78L/day (I.O.M. 1998). Depending on the vitamin in question the studies used for setting the AIs include as few as 22 women (thiamin) and up to 111 women (vitamin B-12) and were conducted between 1951 and 1997 (I.O.M. 1998). Thus, the current AIs are based on data from outdated studies with a low number of subjects, indicating a potential need for re-evaluation. Additional problems with existing data include inconsistent timing and

methods of milk collection, different stages of lactation, and lack of information about the effects of any maternal supplements.

Methods used for analysis of B-vitamins are commonly microbiological, or in more recent years, chromatographic coupled with UV, fluorometric or MS detection (Sakurai, et al. 2005, Shi, et al. 2011, Hampel, et al. 2012). Other techniques include radioisotope dilution assays, animal studies and voltammetry (McCosh, et al. 1931, Thomas, et al. 1979, Mikheeva, et al. 2009). However, these approaches are known to be suboptimal regarding time, sample volumes and cost, and are not necessarily validated for analysis of B-vitamins in the complex human milk matrix (Tamura and Picciano 2006b). In some cases the results obtained with different methods were not in agreement (Shane, et al. 1980, Roughead and McCormick 1990). Therefore, evaluating the suitability of methods for the task on hand is crucial.

In this review, different methodological approaches for analyzing the B-vitamins thiamin, riboflavin, niacin, vitamin B-6 and pantothenic acid, vitamin B-12, folate, biotin, and choline in human milk are discussed including the potential advantages and disadvantages of each method. Relevant physical and chemical properties of each of the vitamins are described to assist in optimizing sample preparation or detection. Moreover, information on the different forms in milk and their change in concentration during lactation are described as well as their effects of deficiency on the infant.

## ***CURRENT KNOWLEDGE***

### ***Thiamin (Vitamin B1)***

Thiamin (3-(4-aminopyrimidin-2-ylmethyl)-5-(2-hydroxyethyl)-4-methylthiazole) was first isolated and structurally characterized in 1926. This water-soluble vitamin ( $S_w$ : 100g/100mL) exists naturally as free thiamin and phosphorylated as thiamin monophosphate (TMP) and thiamin pyrophosphate (TPP). Vitamin B1 is very susceptible to heat and to neutral and alkaline pH. Mild acidic conditions (pH 2 to 4) are preferred for storage in solution, but degradation still occurs when exposed to heat. The spectral properties of thiamin vary with pH; thiamin hydrochloride, a form of vitamin B1 used for food fortification and pharmaceuticals, shows a single maximum at 246nm at pH 2.9. Between pH 5 and 7, two absorption maxima are observed at 234nm and 264nm, representing the pyrimidine and thiazole rings. Above pH 8, thiamin and its phosphate esters can be oxidized quantitatively to thiochrome and thiochrome phosphate esters respectively. Unlike the substrate, the biologically inactive oxidation products are a strongly fluorescent ( $E_x$  : 375nm,  $E_m$  : 432-435nm), a quality which has been widely used for quantitative analysis of vitamin B1 in various matrices since the 1930s (Ellefson 1985, Ball 1994, Eitenmiller and Landen 1999). TPP is essential for normal carbohydrate, nucleic acid, and amino acid metabolism, e.g. decarboxylation of  $\alpha$ -keto acids, the transketolase reaction in the pentose phosphate pathway and synthesis of the primary neurotransmitter acetylcholine. Thiamin phosphates are also involved in nerve impulse transmission (Rindi 1982, Gibson 1990, Gerrits, et al. 1997, Eitenmiller and Landen 1999). Beriberi is the classical syndrome of thiamin deficiency, but cardiovascular, muscular, nervous and gastrointestinal systems are also affected (Gibson 1990). Thiamin concentrations in human milk depend on maternal status but maternal supplementation results in only a limited increase (Pratt and Hamil 1951, Nail, et al. 1980, Thomas, et al. 1980, Lönnerdal 1986). During pregnancy its deficiency can cause problems such

as alterations in glucose tolerance and intrauterine growth, increasing the risk of low birth-weight in the offspring (Baker, et al. 2000, Ortega, et al. 2004) and it is also thought to be a possible cause of sudden infant death syndrome (Bernshaw 1991). Thiamin nutritional status is usually measured by transketolase activity in erythrocytes which reflects the adequacy of body stores (Gibson 1990).

Free thiamin and thiamin monophosphate are the main forms of vitamin B1 in human milk with increasing concentrations as lactation proceeds (Emmett and Rogers 1997, Picciano 1998, Stuetz, et al. 2011). Maternal supplementation of thiamin is not reflected in milk concentrations but in increased urinary excretion, limiting the amount that can be transferred into milk (Slater and Rial 1942, Lönnerdal 1986). Methods for analysis include the classic thiochrome reaction, and microbiological and HPLC methods. However, the first quantitative measurement of vitamin B1 (and also riboflavin) was carried out by measuring the growth of young rats whose diet was supplemented with human milk as the only vitamin B source (McCosh, et al. 1931, Donelson 1934). By feeding various amounts of pooled and individual breast milk to young albino rats, the so-called unit-growth was determined as well as the influence of more adequate vitamin intake (Bourquin and Sherman 1931, Chase and Sherman 1931). By nature, this approach offers only a measurement of physiologically available thiamin. Besides being a very time consuming and expensive process, the results vary considerably. Furthermore, certain bacteria in the intestinal tract are able to synthesize thiamin and provide it to the animal, which adds to the complexity and inaccuracy of this biological assay (Ellefson 1985).

Thiamin quantitation by microbiological assays was first developed using *Lactobacillus fermenti*. However, the assay is compromised by the bacteria's susceptibility to both stimulatory and inhibitory matrix constituents such as sugars, reducing agents and calcium. Other organisms tested include *S. cerevisiae*, *O. malhamensis* and *L. viridescens* (ATCC 12706). The latter provides results comparable to quantification obtained by the thiochrome assay and is commonly accepted for food analysis. The extraction procedures for microbiological assays generally follow the thiochrome analysis procedures such as extraction under acidic conditions (0.33 M H<sub>2</sub>SO<sub>4</sub>) and heating prior to inoculation (Ford, et al. 1983). Microbiological assays require enzymatic hydrolysis of the phosphate esters due to differential growth response to TMP, TDP and TPP (Deibel, et al. 1957, Banhidi 1958, Ford, et al. 1983, Eitenmiller and Landen 1999, Fernández-Muiño, et al. 2008).

Since its development (Barger, et al. 1935), the thiochrome method has been widely used for thiamin analysis in biological matrices. For analyzing free and total thiamin in human milk samples were subjected to acidic hydrolysis (0.1N H<sub>2</sub>SO<sub>4</sub>) and heat followed by incubation overnight with (total thiamin) or without (free thiamin) clarase at pH 4.5. The vitamin was isolated by column chromatography prior to oxidation with potassium ferrocyanide yielding thiochrome. After extraction with isobutanol the samples were ready for analysis with a photofluorometer (Roderuck, et al. 1945). Although thiamin-phosphates are also converted under these conditions, the resulting thiochrome phosphates are not soluble in isobutanol and are therefore undetectable (Kinnorsley and Peters 1938) allowing differentiation between free thiamin and total thiamin in the sample as described. Use of the thiochrome method for human milk has been reported at intervals over the last 60 years (Pearson, et al. 1946, Solov'eva and



Khrustaleva 1954, Ortega, et al. 2004). On the other hand, Hoshi (1956) described a colorimetric analysis using diazotized p-aminoacetophenone to form a colored precipitate under acidic conditions. However heavy metal salts, potassium ferricyanide, cysteine, iodine and other compounds interfere with the analysis resulting in artificially lower thiamin values (Prebluda and McCollum 1939, Sealock and Goodland 1944).

In the last 30 years, HPLC methods have become a commonly used technique for thiamin analysis in human milk still imbedding the well-known thiochrome reaction followed by fluorescence detection (Wienders and Mink 1983, Böhm, et al. 1997, McGready, et al. 2001, Sakurai, et al. 2005); however, HPLC-UV analysis has also been reported for human milk (Shi, et al. 2011). Using the thiochrome approach, HPLC methodology offers pre- or post-column derivatization of thiamin and its phosphate esters. While earlier reports used a post-column modification of the vitamin (Wienders and Mink 1983, Böhm, et al. 1997), pre-column derivatization of thiamin analysis in human milk has been described for the last ten years (McGready, et al. 2001, Stuetz, et al. 2011, Stuetz, et al. 2012). Recently Stuetz *et al.* (2011, 2012) analyzed thiamin and its monophosphate in human milk based on pre-column derivatization, which can be conducted fully automated. The analysis is then carried out by reversed-phase liquid chromatography coupled with fluorescence detection. Most recently we have developed an UPLC-MS/MS method for human milk samples analyzing thiamin simultaneously with other B-vitamins, specifically vitamin B2, niacin and pyridoxal after removal of proteins and non-polar constituents (Hampel, et al. 2012). Since pre- or post-column derivatization has been rendered obsolete when using this method and multiple vitamins are analyzed within one analytical run, more information is available in less time and with lower

cost. To account for matrix effects influencing analyte recovery stable isotope internal standards were used for quantitation (Hampel et al. 2012).

A similar approach has been described for the simultaneous determination of thiamin, riboflavin, nicotinamide, nicotinic acid, pantothenic acid, pyridoxal, and pyridoxine using LC/MS-TOF in nutritional yeast powder as a model food matrix (Hälvin et al., 2013). A comparison between various enzyme and acid treatments was conducted to optimize the release of the B-vitamins from their co-enzymatic forms. While thiamin and riboflavin both could be liberated using enzyme preparations such as takadiastase,  $\alpha$ -amylase,  $\beta$ -glucosidase and acid phosphatase, acid treatment appeared to be ineffective but then again supported the release of pyridoxal, which was only marginally liberated by the enzymes (Hälvin et al., 2013). Neither nicotinamide nor pantothenic acid could be liberated under any tested conditions, illustrating the complexity and difficulties of simultaneous analysis of multiple analytes present in free and bound forms.

### ***Riboflavin (Vitamin B2)***

Riboflavin (7,8-dimethyl-10-ribityl-isoalloxazine) was originally isolated from milk whey simultaneously with thiamin in 1879 and given the name lactochrome or vitamin G (Powers 2003, Buehler 2011). The coenzymes flavin adenine dinucleotide (FAD) and flavin mononucleotide (FMN), the most important biologically active forms of riboflavin, were characterized between 1934 and 1938. While relatively stable under heat and acidic conditions, vitamin B2 is readily degraded above pH 7.0 and by UV- and visible light. Both coenzymes are easily converted to riboflavin below pH 5.0. Unlike FAD and FMN ( $S_{W(\text{Na salt})}$ : 30 - 67g/L<sub>25°C</sub>)

riboflavin's yellow-orange crystals are poorly soluble in water ( $S_w \approx 0.1 \text{ g/L}_{25^\circ\text{C}}$ ). In aqueous solution, riboflavin shows absorption-maxima at 223, 226, 373 and 445nm, but flavins also possess a strong native fluorescence ( $\lambda_{\text{ex}}$  : 400 to 500,  $\lambda_{\text{em}}$  : 520 to 530). On an equimolar basis, riboflavin and FMN fluoresce similarly, while FAD shows only 10-20% of the intensity. However, when flavins are bound to proteins a dramatic quenching of fluorescence is observed (Maggiorelli 1958, Shah 1985, Russell and Vanderslice 1992, Ball 1994, Eitenmiller and Landen 1999, Zielenkiewicz, et al. 2010). Humans require riboflavin for DNA repair, energy production, fatty acid and amino acid synthesis, folic acid activation, the formation of the free radical scavenger glutathione, and the conversion of tryptophan to niacin in the liver. Since this vitamin is poorly stored and not synthesized by vertebrates it must be constantly supplied by diet (e.g. meat, milk and dairy products). Therefore, maternal riboflavin deficiency rapidly results in low milk concentrations of the vitamin and biochemical signs of depletion occur within only a few days of dietary deprivation (Bates, et al. 1982b, Powers 2003). Biochemical methods, such as the analysis of the FAD-dependent erythrocyte glutathione reductase (EGR), are essential for confirming clinical and subclinical cases of ariboflavinosis. Riboflavin deficiency usually occurs along with other water-soluble vitamin deficiencies and causes growth retardation, anemia, skin lesions or degenerative changes in the nervous system (Gibson 1990, Van Herwaarden, et al. 2007, Buehler 2011).

FAD and riboflavin are the prevalent forms of vitamin B2 in human milk (Roughead and McCormick 1990, Sakurai, et al. 2005) with an average concentration of 0.35mg/L in well-nourished women (Allen 2005). Other flavins present include 10-hydroxy-ethylflavin (2-10%; formed during the anaerobic bacterial degradation of riboflavin) and traces of 10-formyl-

methylflavin, 7 $\alpha$ -hydroxyriboflavin, 8 $\alpha$ -hydroxy-riboflavin (both products of riboflavin catabolism) and FMN (Roughead and McCormick 1990, Sakurai, et al. 2005). Throughout lactation the vitamin B2 concentration in human milk remains constant (Emmett and Rogers 1997). Quantitative methods for analysis of vitamin B2 in human milk include animal assays, and microbiological and spectroscopic (UV, fluorescence) techniques (McCosh, et al. 1931, Donelson 1934, Thomas, et al. 1980, Ford, et al. 1983, Ronnholm 1986, Toyosaki, et al. 1986, Roughead and McCormick 1990). As already mentioned, the first quantitative measurement of ðvitamin Gö combined with thiamin was described by Donelson and Macy (1934) and McCosh *et al.* (1931) (McCosh, et al. 1931, Donelson 1934) based on the growth of young rats, which used to be the original standard for other methods (Shah 1985). However, the results of this assay tend to be overestimates due to additional amounts of the vitamin in the food. Such animal studies are time consuming, expensive and least accurate, but they best reflect riboflavin bioavailability (Cosner and Schuck 1948, Shah 1985).

Most microbiological assays for riboflavin have been performed with *Lactobacillus rhamnosis* (formerly *L. casei*) ATCC 7469 and were first described for milk from different animal species in the 1940s (Williams, et al. 1942, Pearson, et al. 1946). Sample preparations consist of acidic hydrolysis, protein precipitation and neutralization prior to incubation with the growth medium. Riboflavin-dependent cell growth is used for indirect quantification by turbidimetry. However, common matrix constituents such as starch, protein degradation products or free fatty acids affect the growth of the culture. Furthermore, the response to FMN or FAD differs from the growth promotion by riboflavin; therefore all methods based on microbiological analysis are for total riboflavin (Bates, et al. 1985, Eitenmiller and Landen 1999).

Fluorometric techniques take advantage of the fact that riboflavin undergoes different routes of degradation depending on the pH. In acidic and neutral conditions the loss of the ribityl side chain results in lumichrome (6,7-dimethyl alloxazine) formation, while under alkaline pH irradiation generates lumiflavin (6,7,9-trimethylisoalloxazine) by cleaving the deoxy-end carbon of the ribityl side chain. The latter is an important analytical tool for flavin detection due to the significantly stronger fluorescence of lumiflavin compared to the native riboflavin. Even though the specificity can be enhanced by additional preparation steps (e.g. chloroform extraction of lumiflavin), the actual reaction does not proceed quantitatively and depends strongly on the experimental conditions and instrumental set-up (Strohecker and Henning 1965, Bates, et al. 1982a, Woodcock, et al. 1982, Eitenmiller and Landen 1999).

In the last 20 years HPLC methods have been introduced for riboflavin analysis in human milk utilizing the spectral properties of vitamin B2 for detection. In 1945, Roderuck and Coryell described the analysis of free and bound riboflavin in human milk by acidic hydrolysis with and without subsequent enzymatic digestion. The samples were purified by column chromatography and analyzed by fluorescence reading with appropriate filters (Roderuck and Coryell 1945), a method which was modified from riboflavin analysis in food products (Conner and Straub 1941). Due to the degradation of FAD and FMN into riboflavin below pH 5.0, the so-called free riboflavin in this study not only refers to actual freely available riboflavin but also to converted FAD and FMN, while the bound fraction, analyzed after enzymatic degradation, also captures riboflavin previously bound by other constituents in the matrix (Watts, et al. 1948). Flavin analysis in human milk by HPLC and fluorescence detection described by Roughead and McCormick (1990) follows the same analytical principle. The

sample preparation, adapted from a goat and cow's milk work-up procedure (Owen and West 1971), includes protein precipitation with trichloro-acetic acid (TCA) followed by phenol extraction of the flavins. Adding water to the phenol phase and back-extracting the organic solvent with diethyl-ether results in an aqueous sample ready for analysis by RP-HPLC. The authors pointed out that the values obtained have to be corrected for the internal quenching in FAD caused by the formation of an intramolecular complex between the isoalloxazine ring and the adenine moieties, which might not have been considered in earlier investigations and be a possible explanation for the discrepancy in their findings (McCormick 1968, Roughead and McCormick 1990). In contrast, studies utilizing an enzymatic treatment for a quantitative conversion of FAD to riboflavin in human milk prior to fluorometric analysis show comparable results (Rettenmaier and Vuilleumier 1983, Ronnholm 1986, Roughead and McCormick 1990). HPLC coupled with fluorescence detection has been described for flavin analysis in human milk for the last two decades and has been used recently (Sakurai, et al. 2005, Van Herwaarden, et al. 2007, Israel-Ballard, et al. 2008). However, riboflavin and FAD can also be analyzed simultaneously with thiamin, niacin and pyridoxal using UPLC-MS/MS; thus both prevalent forms can be analyzed directly in their native forms allowing distinction between the free vitamin and its coenzyme (Hampel, et al. 2012).

In 2009 Mikheeva *et al.* developed a rapid riboflavin analysis in breast milk by voltammetry (VA). It is less labor-intensive and does not require large amounts of reagents and expensive equipment, compared to the previously discussed methods (Mikheeva, et al. 2009). This procedure is based on the oxidizability of riboflavin at a glassy-carbon (GC) indicator electrode. The samples are subjected to mild acidic hydrolysis (0.1N HCl), followed by an

ammonium-sulfate protein precipitation sediment removal prior to analysis. For quantitation purposes, the anodic differential voltammogram is recorded. However, the complexity of the pH dependence of the potential of vitamin B2 has to be considered, given that pH influences both the rate of the electrode process and its mechanism (Mikheeva, et al. 2009).

### *Niacin*

Niacin refers to nicotinic acid (pyridine-3-carboxylic acid) and nicotinamide (pyridine-3-carboxylic acid amide) and was recognized as a curative factor for pellagra in 1937 (Spies, et al. 1938). Symptoms of marginal deficiency include insomnia, loss of appetite, weakness, soreness of mouth and tongue and indigestion. Dietary intake of the vitamin can be established from the ratio of urinary excretion of N $\alpha$ -methylnicotinamide (NMN) and N $\alpha$ -methyl-2-pyridone-5-carboxamide (2-pyridone). However the niacin metabolites only reflect niacin status so potential depletion might not be recognized (Gibson 1990). Humans produce nicotinic acid *in vivo* in the liver utilizing 60mg of the amino acid tryptophan to produce 1mg of nicotinic acid, a process which can be inhibited by excess leucine in the diet (Heidelberger, et al. 1949, Gibson 1990, Ball 1997, Eitenmiller and Landen 1999). Rich sources of niacin include oil seeds, legumes, lean meat and cereals. The latter contain niacin bound as niacytin, of which only very little is biologically available after hydrolysis by gastric acid (Gibson 1990). Niacytin can be degraded by soaking cereals in calcium hydroxide to release the niacin, for example in the process called nixtamalization used in some Latin American countries. Both forms are interconvertible; the free acid is readily converted into nicotinamide, which serves as precursor for the synthesis of nicotinamide adenine dinucleotide (NAD) and nicotinamide adenine dinucleotide phosphate

(NADP). These coenzymes are important co-factors in oxidation-reduction reactions such as electron carriers for intracellular respiration, acting as a co-dehydrogenase involved in the oxidation of fuel molecules (NAD) or as a hydrogen donor in reductive processes for biosynthesis of fatty acids (NADP) (Eitenmiller and De Souza 1985, Eitenmiller and Landen 1999). Both free forms are stable in aqueous solution and not very susceptible to heat or pH. The amide is significantly more water-soluble ( $S_{\text{Wamide}} = 100\text{g}/100\text{mL}$ ,  $S_{\text{Wacid}} = 1.67\text{g}/100\text{mL}$ ) and hydrolyzes easily in acidic or basic conditions converting to the free acid (Eitenmiller and De Souza 1985). The coenzymes NAD and NADP show two UV maxima at 260nm (nicotinamide ring) and 340nm (adenine ring) as well as fluorescent qualities ( $E_m = 470\text{nm}$ , when excited at 260 or 340nm). Free niacin however lacks fluorescence and shows relatively low specificity for UV detection. Therefore, the spectral properties are suboptimal for quantitation purposes and require time consuming purification and isolation procedures to remove interferences. Nonetheless, due to its stability, acidic or alkaline hydrolysis as well as heat can be used to isolate niacin from coenzyme structures and destroy sample matrices (Eitenmiller and Landen 1999).

Nicotinamide, NAD and NADP have been reported as forms of niacin in human milk (Greer 2001, Sakurai, et al. 2005). However, there has been little interest regarding the content of this vitamin in human milk, and quantitation has usually occurred in the context of multi-vitamin studies (Coryell, et al. 1945, Ford, et al. 1983, Sakurai, et al. 2005, Shamsia 2009, Shi, et al. 2011). Its concentration increases with the duration of lactation from colostrum to mature milk (Emmett and Rogers 1997) but niacin deficiency remains challenging in dietary depleted areas where corn and other cereals are major dietary staples. Maternal supplementation with this



vitamin is reflected in the milk content and increases the amount available to the infant (Pratt and Hamil 1951, Lönnerdal 1986, Eitenmiller and Landen 1999). Methods routinely used for niacin analysis include chemical (colorimetric) and microbiological procedures, though other analytical techniques have become available in more recent years.

The basis of the colorimetric analysis is the König reaction (Pelletier and Campbell 1959), in which free niacin and cyanogen bromide form a pyridinium intermediate. By undergoing rearrangements the newly formed derivatives couple with aromatic amines to form colored polymethine products, which absorb at a VIS-maximum of 436nm. The analysis is susceptible to interferences by pyridine derivatives under certain conditions (e.g. trigonelline under alkaline pH) (Sarett, et al. 1940); therefore the choice of temperature, pH, and aromatic amine as a coupling-agent for maximum color development are critical factors for quantitation and need to be controlled (Pelletier and Campbell 1959, Strohecker and Henning 1965, Eitenmiller and De Souza 1985, Eitenmiller and Landen 1999). Although applied to niacin analysis in various matrices, this method does not appear to have been employed for human milk studies.

Most niacin analyses in human milk have been conducted using microbiological assays. Among others, *Lactobacillus plantarum*, *L. mesenteroides* and *Tetrahymena thermophila* have been used for niacin analysis in general. *L. plantarum* shows an equimolar response to nicotinic acid, nicotinamide and NAD offering a reliable analysis of total niacin in biological matrices, while *L. mesenteroides* only responds to nicotinic acid, and *Tetrahymena thermophila* equally but not additively responds to both free niacin forms (Eitenmiller and De Souza 1985, Eitenmiller and Landen 1999). With regard to human milk, *L. arabinosus* has been reported

many times to be the microorganism of choice (Coryell, et al. 1945, Macy 1949, Pratt and Hamil 1951, Ford, et al. 1983). Coryell *et al.* (1945) were the first group to analyze niacin in human milk using *L. arabinosus*, which had been introduced as a suitable microorganism by Snell and Wright (1941) and modified by Krehl *et al.* (1943) and subsequently used by other groups. The samples were autoclaved and hydrolyzed under acidic conditions (0.1N HCl) followed by protein precipitation at pH 4.6 and diluted before the basal medium was added. After inoculating with the bacterium the nicotinic acid concentration was determined by turbidimetry (Coryell, et al. 1945). Due to higher sensitivity and specificity, the microbiological method offers better detection limits for samples with low amounts of niacin than the colorimetric analysis. However, the assay is susceptible to growth-stimulating or growth-depressing interferences causing errors during the analysis (Eitenmiller and De Souza 1985, Eitenmiller and Landen 1999).

Only recently have HPLC methods been applied for niacin analysis in human milk. Sakurai *et al.* (2005) described the analysis of unbound nicotinamide in breast milk by HPLC using a UV detector at 260nm for quantitation. The proteins in the sample were precipitated with acetonitrile and removed by centrifugation prior to analysis, a procedure also employed by Shi *et al.* (2011). Protein precipitation was carried out using a precipitation solution consisting of zinc acetate, polyhydrated phosphotungstic acid and glacial acetic acid, a method adapted from the analysis of water-soluble vitamins in supplementary foods (Zafra-Gómez, et al. 2006). HPLC coupled to a diode array detector (DAD) using the method of the Association of Official Analytical Chemists has also been applied to human milk samples (AOAC, 1990, (Shamsia 2009). Using UPLC-MS/MS nicotinamide can be analyzed simultaneously with thiamin, riboflavin, FAD and pyridoxal after protein and fat-removal (Hampel, et al. 2012).

### ***Vitamin B6***

Vitamin B6 (2-methyl-3-hydroxy-5-hydroxy methyl pyridine derivatives) is the collective term for the biologically active equivalent and metabolically interconvertible pyridoxine (PN), pyridoxal (PL), and pyridoxamine (PM). In 1934, Gyorgy introduced the term vitamin B6 to differentiate between this factor and other hypothetical growth factors. Pyridoxine was the first B6 vitamers to be structurally identified in 1938 followed by pyridoxal and pyridoxamine in the 1940s. All vitamers exist in their phosphorylated forms i.e. pyridoxal-5 $\phi$ -phosphate (PLP), pyridoxine-5 $\phi$ -phosphate (PNP) and pyridoxamine-5 $\phi$ -phosphate (PMP). The latter two can be oxidized to the metabolically active form, PLP, which serves as a co-factor in numerous amino acid transformations such as transamination, decarboxylation or racemization by forming a Schiff base (Gyorgy 1934, Gyorgy and Eckardt 1939, Snell 1944, Polansky, et al. 1985, Eitenmiller and Landen 1999, Leklem 2001). PN is the most stable form of vitamin B6, followed by PL and PM. All vitamers are quite stable under acidic conditions but degrade greatly above neutral pH. All forms are light sensitive even under acidic conditions; near-UV and UV radiations cause especially significant degradation of the vitamers. The spectral properties are pH- and compound-dependent; all vitamers including their 5 $\phi$ -phosphates show a maximum absorption at 290nm when dissolved in 0.1N HCl, while at neutral pH the UV maxima of PL and PLP shift to 390nm. The remaining vitamers maximally absorb at 253 and 325nm under the given conditions. The non-phosphorylated vitamers as well as PNP and PMP possess a strong fluorescence (Ex : 310  $\phi$  365nm; Em : 365  $\phi$  430nm) from slightly acidic to strongly alkaline pH while PLP only shows weak fluorescent properties since the phosphate ester of the 5-position

of the pyridine ring prohibits the hemiacetal formation necessary for a strong fluorescence (Bridges, et al. 1966, Saidi and Warthesen 1983, Eitenmiller and Landen 1999, Ubbink 2000).

General symptoms of B6 deficiency include weakness, sleeplessness, appetite or growth depression and various dermatologic disorders. Clinical status tests include the analysis of plasma PLP and urinary measurements of PL and 4-pyridoxic acid (4-PA). Furthermore, PLP is also involved as a coenzyme in the kynurenine stage of niacin formation from tryptophan; therefore, B6 deficiency increases urinary excretion of kynurenine and 3-hydroxykynurenine, an effect also resulting from oral loading of tryptophan, known as the tryptophan load test. The latter combined with the plasma B6 test currently provides excellent biochemical confirmation of vitamin B6 status. The vitamers are found in all foods of both animal and plant origin such as meats, cereal grains, vegetables or nuts with extremely variable bioavailability (Polansky, et al. 1985, Eitenmiller and Landen 1999). Vitamin B6 is a crucial dietary factor during the early stages of central nervous system development in early infancy and the already limited pool of this vitamin in adult humans rapidly depletes with deficient intake. However, vitamin B6 intake of lactating women appears to influence the concentration of the vitamin in their milk and is a strong predictor of infant vitamin B6 status (West and Kirksey 1976, Morrison and Driskell 1985, Gibson 1990, Kang-Yoon, et al. 1992).

PL has been reported as the principal form of vitamin B6 in human milk, with possible contributions of PLP (7-64%), PMP, PN and PM (mostly traces) (Vanderslice, et al. 1983, Hamaker, et al. 1985, Morrison and Driskell 1985). Its concentration increases throughout lactation from colostrum to mature milk but decreases with lactation beyond 7 months (Karra, et al. 1986, Emmett and Rogers 1997). While maternal vitamin B6 status has been correlated with

milk concentrations, supplementation with this vitamin seems not to be reflected in the milk (Thomas, et al. 1980, Prentice, et al. 1983, Styslinger and Kirksey 1985, Lönnerdal 1986).

Microbiological assays and LC based methods have been established as the common techniques for vitamin B6 analysis. However, one of the first quantitative estimations of vitamin B6 carried out by Gyorgy (1936) utilized an animal assay to determine the volume of the so-called "Petersø eluate" (equivalent to 1 rat-day dose) which has a curative effect against dermatitis in rats (Gyorgy 1936).

The most common organism used for vitamin B6 analysis in biological samples by microbiological assay is *Saccharomyces uvarum* (ATCC 9080) and this has been used for human milk analysis (West and Kirksey 1976, Roepke and Kirksey 1979, Thomas, et al. 1980, Sneed, et al. 1981, Borschel, et al. 1986, Karra, et al. 1986, Eitenmiller and Landen 1999, Leklem 2001). Besides possible inhibition of growth due to high salt concentrations, different growth responses for PN, PL and PM add to the assay's complexity. Therefore, after hydrolyzing the samples the vitamers are separated chromatographically prior to addition to the yeast basal medium allowing the determination of each vitamer individually (Parrish, et al. 1955, Parrish, et al. 1956, Storvick, et al. 1964, Thiele and Brin 1966, Polansky 1981, Eitenmiller and Landen 1999, Fernández-Muñoz, et al. 2008). Other microorganisms such as *Kloeckera brevis* and *Lactobacillus casei* have also been described for vitamin B6 analysis in human milk (Barton-Wright 1962, Ford, et al. 1983, Wilson and Davis 1984). Since *Lactobacillus casei* only responds to pyridoxal, other forms of this vitamin have to be converted by hydrolysis with glycolic acid at 100°C for 15min (PM) or with manganese dioxide at room temperature (PN) (Wilson and Davis 1984). In any

case, the extensive sample preparation and intricacies of microbiological assays have led to the development of chromatographic methods for vitamin B6 analysis.

HPLC methods introduced for human milk analysis generally describe a chromatographic separation coupled with fluorometric detection. Vanderslice *et al.* (1983) treated their samples with sulfosalicylic acid as an extraction agent followed by further clean-up procedures. The B6 vitamers were separated by anion exchange chromatography prior to detection. Results obtained by HPLC analysis were in good agreement with the microbiological assay (Vanderslice, et al. 1983). Further methods described include (RP)-ion pair HPLC separation (Morrison and Driskell 1985) or a simple RP-HPLC separation, with subsequent post-column bisulfate derivatization to enhance PLP-detection. Samples were diluted in mobile phase buffer and proteins were precipitation with TCA. After centrifugation, the supernatant was collected and washed with diethyl ether to remove remaining TCA and fats prior to analysis (Hamaker, et al. 1985). The results obtained with the analytical method were in good agreement with the microbiological assay and also comparable to values obtained with the ion-exchange HPLC methods. RP-HPLC has continued to be described as a suitable method for vitamin B6 analysis in more recent years introducing a post-column photochemical conversion of PLP for improved detection (Gatti and Gioia 2005). However, UPLC-MS/MS allows the analysis of pyridoxal simultaneously with thiamin, riboflavin, FAD and niacin after protein and fat-removal directly without need for derivatization (Hampel, et al. 2012).

### ***Cobalamin (Vitamin B12)***

Vitamin B12 is the collective term for cobalt-containing corrinoids, which were first isolated in 1948 (Rickes, et al. 1948, Smith 1948) and structurally elucidated by Hodgkin and coworkers (Hodgkin, et al. 1956, Hodgkin, et al. 1962, White 1962, Brink-Shoemaker, et al. 1964). The molecule is composed of a central cobalt atom coordinated by a nearly planar corrin ring. Its axial coordination sites are occupied by dimethylbenzimidazole and various other groups such as cyano-, hydroxo-, nitrito- and sulfite-functions, a bonding that can be reversibly broken and reformed (Chin 1984, Eitenmiller and Landen 1999). The prevalent form of vitamin B12 in biological materials is coenzyme B12, in which the axial group is either occupied by a methyl group or an adenine nucleoside connected by a cobalt-carbon (Barker, et al. 1960, Lenhert and Hodgkin 1961, Lenhert 1968, I.o.M. 1998). These coenzymes are involved in the folate-dependent methylation of homocysteine to methionine and the conversion of methylmalonyl-coenzyme A to succinyl-coenzyme A (Gibson 1990, I.o.M. 1998). Animal source foods such as meats, sea food, eggs and dairy products represent the main source of B12, while plant derived foods usually provide none of this vitamin (Eitenmiller and Landen 1999, Allen 2009).

Being the most stable vitamer, cyanocobalamin (CNCbl) is used in vitamin preparations, supplements and medical or fortified foods. CNCbl is soluble in water (1g/80mL at 25°C), showing absorption maxima at 278, 361, and 551 nm. Further possible dissolving agents include alcohols, phenols and other polar solvents with a hydroxyl group. Its red crystals are stable when protected from light; they do not melt but decompose above 200°C. CNCbl is most stable between pH 4.0 and 4.5 but will be inactivated under strong alkaline or acidic conditions as well as by UV, strong visible light and oxidizing agents (Herbert 1996, Eitenmiller and Landen 1999, Lindemaus 2000, Beck 2001).

Since B12 is bound to proteins in food, the vitamin needs to be released by enzymatic and acidic hydrolysis in the stomach prior to absorption. Further steps include binding to salivary proteins (R-binders) and to the intrinsic factor secreted by gastric parietal cells prior to absorption in the ileum. B12 is either complexed with a delivery protein transcobalamin (TCII) or with the storage protein (haptocorrin, HC). B12 deficiency is generally caused by inadequate dietary intake, failure to secrete the gastric intrinsic factor necessary for B12 absorption (pernicious anemia, an autoimmune condition), or achlorhydria. Since vitamin B12 - as well as folate - is involved in DNA synthesis, B12 deficiency can result in megaloblastic anemia. Other symptoms include neurological disorders such as neuropathy or spinal cord dysfunction. General symptoms of pernicious anemia include glossitis, weakness, loss of appetite and weight, taste and smell, memory impairment, mild depression and hallucinations (Gibson 1990, Wardinsky, et al. 1995, Eitenmiller and Landen 1999, Allen 2009).

Even though biologically active (cobalamin) and inactive (cobinamide) forms of vitamin B12 are found in plasma, serum, bile and liver, only the biologically active cobalamins are selectively transported into milk (Kolhouse, et al. 1978, Adjalla, et al. 1993, Adjalla, et al. 1994). Compared to serum, human milk contains about 100-fold higher amounts of HC, mostly in the form of apoHC, which potentially interferes with B12 quantification (Sandberg, et al. 1981, Lildballe, et al. 2009). Methylcobalamin (MeCbl) was reported as being the dominant form of the B12 vitamers in human milk followed by 5 $\alpha$ -deoxyadenosylcobalamin (AdoCbl) and small amounts of hydroxocobalamin (OHCbl) and CNCbl, all bound to haptocorrin (Craft, et al. 1971, Sandberg, et al. 1981, Adjalla, et al. 1994). Throughout lactation vitamin B12 decreases in



concentration and low maternal intake is reflected in low milk concentrations (Lönnerdal 1986, Trugo and Sardinha 1994, Emmett and Rogers 1997, Deegan, et al. 2012).

Early approaches for analysis of B12 in human milk were carried out by microbiological assays using *Euglena gracilis* as the test organism (Jadhav, et al. 1962, Jathar, et al. 1970, Craft, et al. 1971, Sandberg, et al. 1981) based on a method developed for body fluids (Ross 1952) but adding enzymatic digestion with papain to release the B12 from binding to haptocorrin. Further steps include the conversion of the different forms into CNCbl by sodium cyanide prior to analysis. Alternatively, *Lactobacillus leichmanii* (NCIB 8118) has also been used to assay B12 microbiologically, and proteins were precipitated by heat in the presence of acetate buffer (Samson and McClelland 1980). However, microbiological assays using Lactobacilli can be influenced by desoxyribonucleosides such as thymidine and other compounds, which have been found to replace B12 as a growth factor resulting in a lower specificity and an overestimate of B12 concentrations (Shive, et al. 1951, Ross 1952).

Alternatively, the radioisotope dilution assay (RIDA) first described by Lau *et al.* (1965) for serum B12 has also been applied for human milk analysis (Areekul, et al. 1977, Thomas, et al. 1979, Thomas, et al. 1980, Van Zoeren-Grobbe, et al. 1987, McPhee, et al. 1988, Donangelo, et al. 1989, Specker, et al. 1990, Adjalla, et al. 1994, Trugo and Sardinha 1994, Casterline, et al. 1997, Patel and Lovelady 1998). The method is based on competitive binding of endogenous vitamin B12 and added radioactive B12 to limited binding sites on intrinsic factor. Therefore, the amount of bound radioactive B12 is inversely correlated to the amount of endogenous B12 (Lau, et al. 1965, Chin 1985). While it has been mentioned that modified versions of the radioisotope dilution procedure using commercially available radio immunoassay

kits are not validated for human milk, it has also been stated that this approach for vitamin B12 quantitation in human milk is acceptable (Areekul, et al. 1977, Sneed, et al. 1981). However, several different commercially available radio assay kits have been used for analysis, and none of them has been developed for the human milk matrix. Modifications to the method include a hot extraction of the sample with ethanol or digestions using papain prior to the radioassay (Specker, et al. 1990, Adjalla, et al. 1994, Keizer, et al. 1995). Analysis of human milk from a mother of a diagnosed B12-deficient infant using a competitive protein binding assay resulted in very low B12 concentrations compared to values obtained with the same method from well-nourished lactating women (McPhee, et al. 1988). Since no validation was carried out prior to application, the accuracy of the values remains uncertain.

Lildballe *et al.* (2009) proposed the removal of apo-haptocorrin (apoHC) by a cobinamide-sepharose affinity column prior to B12 analysis due to the fact that the large amounts of the free form of haptocorrin in human milk may interfere with the analysis. Analyzing untreated samples with high amounts of apoHC using different commercially available immunoassay systems coupled with chemiluminescence detection resulted in spuriously high (Centaur analyzer, Siemens) or low (Architect i2000 analyzer, Abbott; Cobas 6000 E immunoassay system, Roche Diagnostics) concentrations of B12, while concentrations of apoHC below 10nM appeared not to influence the analysis (Lildballe, et al. 2009). Competitive protein binding coupled with chemiluminescence detection with and without apoHC removal prior to analysis appears to be the method of choice for B12 analysis in human milk in recent years (Israel-Ballard, et al. 2008, Honzik, et al. 2010, Deegan, et al. 2012).

### *Folate*

Folate refers to the large group of heterocyclic compounds based on the pteric acid structure conjugated with one (folic acid) or more L-glutamates (polyglutamyl folate) linked to the  $\gamma$ -carboxyl of the amino acid, all exhibiting the biological activity of folic acid (pteroylglutamic acid). Folate deficiency was first recognized in 1931 and this anti-anemia factor was designated  $\beta$ -vitamin M $\ddot{o}$  (Day, et al. 1938). In 1946, folic acid was identified and synthesized using techniques previously used to study other pteridines, such as butterfly pigments. Even though folic acid is not found in nature, this slightly water-soluble form is used for food fortification and pharmaceuticals and must be reduced to dihydro- or the active co-enzyme form tetrahydro-folate (THF). As a co-enzyme, folate is involved in the transfer of single-carbon atom groups such as formyl- (CHO), methyl- (CH<sub>3</sub>) or formimino-functions (CH=NH) in the metabolism of amino acids and also in purine and pyrimidine synthesis in nucleic acid formation (Gibson 1990, Eitenmiller and Landen 1999, Brody and Shane 2001).

Folates are labile to changes in pH, oxidizing agents, heat and light exposure. They show characteristic UV absorption spectra generally with three absorbance maxima. The p-aminobenzoyl-glutamic acid (PABG) moiety influences the UV (UV<sub>max</sub> 270-280nm) as well as the fluorescence properties of folates while reduced pterins without PABG do not possess native fluorescence. While the reduced forms show excitation and emission maxima at  $\lambda_{Ex}$  300-320 nm and  $\lambda_{Em}$  360-425 nm, folate itself fluoresces at maxima of  $\lambda_{Ex}$  360-380 nm and  $\lambda_{Em}$  450-460 nm respectively. Fluorescence intensity is also significantly influenced by pH and buffer composition (Uyeda and Rabinowitz 1963, Ball 1994, Bates and Prentice 1994, Eitenmiller and Landen 1999, Fernández-Muiño, et al. 2008).

Signs of marginal deficiency include tiredness, irritability, and decreased appetite. Megaloblastic anemia indicates severe folate deficiency, which has been reported in pregnant and lactating women in developing countries where dietary intakes may not meet their high requirements. Moreover, low folate levels during the periconceptional period combined with genetic susceptibility can cause neural tube defects (NTD) in infants and may be deleterious to fetal growth and normal development of the fetal nervous system. The main forms of folate in foods are the polyglutamates found in liver, yeast, leafy vegetables or nuts (Donangelo, et al. 1989, Gibson 1990, Shaw, et al. 1995, Eitenmiller and Landen 1999).

In human milk, folates are quantitatively bound to whey binding proteins and predominantly present as pteroylpolyglutamates and as N-5 methyltetrahydrofolate (N-5-MeTHF) with a minor contribution of reduced folacin derivatives (Smith, et al. 1985, Brown, et al. 1986, Selhub 1989, O'Connor, et al. 1991, Tamura and Picciano 2006a). Recent experiments confirmed the presence of folic acid, *p*-aminobenzoylglutamate (*p*ABGA) and its acetamide derivative *a*-*p*ABGA in the picomolar-range in breast milk (Álvarez-Sánchez, et al. 2010). The folate content of human milk increases throughout lactation; however, a correlation between maternal intake and milk folate has not been found (Ramasastri 1965, Cooperman, et al. 1982, Smith, et al. 1983, Lönnerdal 1986, Greer 2001, Tamura and Picciano 2006a).

The commonly accepted method for folate quantitation in the past has been a microbiological assay using *Lactobacillus casei* ATTC 7469, since this bacterium responds to all forms of folate (O'Connor, et al. 1997). However, other microorganisms (*Streptococcus faecalis*, *Pediococcus cerevisiae*) have been suggested for folate measurement due to differential responses to different folate derivatives, and were tested initially on serum samples (Herbert

1961, Tamura, et al. 1972). Ramasastri (1965) suggested autoclaving the samples in the presence of ascorbic acid under neutral pH conditions to prevent oxidation of the analytes. Adjusting the pH to 4.5 resulted in clear filtrates before the pH was then adjusted to pH 6.8 prior to analysis by the microbiological assay. Using the different bacteria for analysis (*L. casei*, *S. faecalis*, *P. cerevisiae*) different growth responses were observed, presenting the first differential analysis of folate forms in human milk. Various studies have been conducted using *L. casei* and/or the other stated microorganisms, with and without folate conjugase treatment (to convert folate polyglutamates into the monoglutamate form) for total and free folate content respectively, or to differentiate between the different forms of folate in breast milk (Metz, et al. 1968, Jathar, et al. 1970, Tamura, et al. 1980, Butte and Calloway 1981, Sneed, et al. 1981, Cooperman, et al. 1982, Ek 1983, Ford, et al. 1983, Smith, et al. 1983, Eitenmiller, et al. 1984, Bank, et al. 1985, Smith, et al. 1985, Udipi, et al. 1985, Brown, et al. 1986, Udipi, et al. 1987, Swiatlo, et al. 1990, Keizer, et al. 1995). Lim et al. (1988) suggested a tri-enzyme treatment of the milk samples with  $\alpha$ -amylase and protease in addition to the folate conjugase prior to the microbiological assay resulting in increased values for the vitamin, an approach used in subsequent studies (Lim, et al. 1998, Mackey and Picciano 1999, Villalpando, et al. 2003, Hyun and Tamura 2005, Khambalia, et al. 2006, Han, et al. 2010).

Chromatographic methods have been used rarely for folate analysis. Sakurai et al. (2005) described a HPLC method using fluorometric detection, an approach described by Allfrey and coworkers in 1949 (Allfrey, et al. 1949, Sakurai, et al. 2005). However, this method appears to not be suitable since it was developed for analyzing the synthetic form of the vitamin, usually not present in biological samples. Furthermore, reduced folates were not protected by reducing

agents during the sample preparation and the described permanganate oxidation to cleave the C9-N10 bond is incomplete when a sample contains other folates such as 5-MeTHF and 10-formylTHF, the main folates found in biological samples. Therefore, results obtained most likely underestimate the actual folate content (Tamura and Picciano 2006b). Shi et al. (2011) also referred to a LC-fluorometric detection technique, described for the analysis of folic acid in vitamin-enriched (dairy) milk and infant nutrition products (Zafra-Gómez, et al. 2006). Therefore, the pteroylpolyglutamates and N-5 methyltetrahydrofolate present in human milk might not be reflected in the results. Recently, a new fully automated on-line SPE-HILIC-MS/MS has been introduced for determining folic acid and its catabolites *p*ABGA and the acetamide derivative *a-p*ABGA in human biofluids. Using only the whey fraction, the samples are subjected to solid-phase extraction (SPE) using ion exchange mode. Elution of the analytes was carried out with the mobile phase allowing the sample to be directly loaded onto the hydrophilic interaction liquid chromatography (HILIC)-system for analysis (Álvarez-Sánchez, et al. 2010). A high-throughput method for quantifying total folate by derivatization of the folates to *p*ABA (para-aminobenzoic acid) prior to analysis by LC-MS/MS using 96-well plates has also been reported (Owens et al. (2007). Even though the analysis is described for whole blood, this approach might be adaptable for breast milk analysis in the future.

Alternatively, competitive radio- and chemiluminescence assays have also been described for folate analysis (Van Zoeren-Grobbe, et al. 1987, Donangelo, et al. 1989, Israel-Ballard, et al. 2008) mostly using commercially available assay kits. However, a comparison between several radioassays and the microbiological technique revealed a potential overestimate of the free folacin content by radioassay when polyglutamate forms of 5-MeTHF are present in the

sample (Shane, et al. 1980). Additionally, kits available for folate analysis are commonly validated for serum, plasma or red blood cells and not tested for human milk.

### ***Pantothenic Acid***

Pantothenic acid (d(+)-N-(2,4-dihydroxy-3,3-dimethyl-1-oxobutyl)- $\beta$ -alanine) consists of pantoic acid coupled to  $\beta$ -alanine by an amide linkage. The optical activity derives from the hydroxylated carbon; however, only the d(+)-enantiomers are biologically active and present in nature. This compound was first isolated and identified as a growth factor from yeast in 1933 with wide distribution in biological tissues (Williams, et al. 1933). Later, this growth factor was related to the chick antidermatitis factor, establishing the status of pantothenic acid as a vitamin for animals (Jukes 1939, Woolley, et al. 1939). Unlike many vitamins, pantothenic acid is not very susceptible to oxidative conditions or light exposure. Since this vitamin does not contain a chromophore, only weak absorption occurs below 210 nm due to the carbonyl function. Thermal stability is best at pH 5-7. Pantothenic acid is highly hygroscopic; therefore, the calcium salt or the alcohol pantothenol are commonly used for food fortification and pharmaceutical products (Eitenmiller and Landen 1999, Plesofsky 2001).

Being a component of coenzyme A (CoA) and acyl carrier protein (ACP), pantothenic acid is vital to many enzymatic processes and the fatty acid synthase complex and is involved in the regulation of alcohols, amines, and amino acids, and the oxidation of pyruvate by the pyruvate hydroxylase complex to form acetyl-CoA. Pantothenic acid is essential to all forms of life and is readily available in good quantities in foods such as liver, kidney, yeast, egg yolk and broccoli. Deficiency of this vitamin may result in headaches, fatigue, insomnia, intestinal

disturbance and paresthesia (abnormal sensation) in hands and feet (Wyse, et al. 1985, Eitenmiller and Landen 1999, Plesofsky 2001).

In breast milk, pantothenic acid is mostly available in its free form (85-90%) while only a minor proportion is some bound form. Despite the importance of this vitamin in lipid metabolism, pantothenic acid is not entrapped by the large milk fat globules nor does it occur in significant amounts in the lipid fraction (Song, et al. 1984). Its concentrations in milk increase throughout lactation and appear to correlate with maternal intake (Lönnerdal 1986, Emmett and Rogers 1997). The majority of analyses carried out in human milk have been done by microbiological assays. Coryell et al. (1945) described a procedure using *Lactobacillus casei* no. 1769; the diluted samples were autoclaved, adjusted to pH 4.6 and boiled to precipitate the casein. After filtration, the basal medium was added and the sample autoclaved again prior to inoculating. The growth was measured by turbidimetry. Enzymatic treatment using mylase P or clarase did not seem to affect the assay. However *L. arabinosus* was also tested for suitability and showed a better growth response and reproducibility to pantothenic acid. Besides the mentioned microorganisms, *Lactobacillus plantarum* has also been employed for pantothenic acid analysis in human milk. Additional double enzyme treatment prior to sample preparation has been used to distinguish between free and total pantothenic acid. Enzymes applied include takadiastase/papain, intestinal phosphatase/pigeon liver extract or amylase/papain (Pearson, et al. 1946, Pratt and Hamil 1951, Ford, et al. 1983, Friend, et al. 1983, Goldsmith, et al. 1983, Sakurai, et al. 2005).

Alternatively, pantothenic acid has been quantified by radioimmunoassay (RIA) as described by Song et al. (1984). Samples were incubated at 37°C with bovine intestinal alkaline



phosphatase and pantetheinase. Additional testing of deproteinization methods revealed incomplete removal of the proteins by thermal treatment such as boiling or autoclaving compared to chemical treatment or dialysis prior to analysis by RIA as described for blood and other tissues (Wyse, et al. 1979). D-pantothenic acid is conjugated with bovine serum albumin via a bromoacetyl derivative of the vitamin for which the antibody was raised in rabbits. The sample and radiolabeled sodium D-pantothenate were then added to the antiserum, and incubated for 5min at room temperature before precipitation of the antibody-bound pantothenic acid by ammonium sulfate. The radioactivity of the precipitate was used for quantification (Wyse, et al. 1979).

Even though pantothenic acid does not possess a strong UV-absorption, HPLC coupled to UV-detection at 195 nm has been described for pantothenic acid analysis in supplemented foods such as vitamin-enriched milk and infant nutrition products and has been adapted for human milk. Samples were subjected to protein precipitation using aqueous acetic acid solution containing zinc and wolframium salts and filtered prior to analysis (Zafra-Gómez, et al. 2006, Shi, et al. 2011). Given the lack of chromophores in the molecule, this detection technique still allowed quantitation of 0.5 mg/kg (ppm).

### ***Biotin***

Biotin (*cis*-hexahydro-2-oxo-1H-thieno [3,4-d] imidazole-4-pentanoic acid) was found to be a requirement for mammals by Boas in 1927. So-called ðegg white injuryö was observed in rats fed raw egg white, while a curative effect was achieved by feeding them raw liver. In 1936, biotin was isolated and crystallized from egg yolk by Koegel and Tonnies, and Snell showed the

glycoprotein, avidin, functions as the biotin antagonist responsible for the toxicity of raw egg white in 1941. The structure was elucidated in 1942 with synthesis in 1943 (Boas 1927, Eitenmiller and Landen 1999, Mock 2001, Ball 2006). The bicyclic structure consists of an ureido ring fused to a tetrahydrothiophene ring with a valeric acid side chain. Of the eight known stereoisomers only d(+)-biotin exhibits biological activity and is found in nature. In solution, biotin is quite stable in a pH range of 4 to 9. This vitamin is insoluble in organic solvent and only sparingly water-soluble (20mg/100mL). The carbonyl group reveals only weak UV-absorption at  $\lambda_{\text{max}}$  204 nm. Both UV-absorbance and fluorescence can be used for detection but the vitamin needs to be derivatized in order to achieve reasonable sensitivity (Eitenmiller and Landen 1999).

Biotin functions as a covalently bound prosthetic group in four carboxylase enzymes that play key roles in gluconeogenesis, fatty acid biosynthesis, amino acid metabolism, and odd-chain fatty acid catabolism (Ball 2006). Symptoms of inadequate biotin levels include anorexia, nausea, vomiting, glossitis, pallor, mental depression, hair loss (alopecia), dry scaly dermatitis and an increase in serum cholesterol and bile pigments. Lack of biotinidase due to an inborn metabolic defect causes infant deficiency. This enzyme cleaves biotinyl-lysine bonds in biocytin resulting in inefficient release of protein bound biotin in the gastrointestinal tract if not available, less salvage of biotin at the cellular level and through increased renal loss. Organ meats, egg yolks and milk are good food sources of biotin (Eitenmiller and Landen 1999, Fernández-Muiño, et al. 2008).

In human milk, more than 95% of biotin is present in the skimmed milk fraction. Less than 3% is reversibly bound to macromolecules and less than 5% is covalently bound to macromolecules (Mock, et al. 1997). Forms found in early and transitional human milk include

biotin and its metabolites bisnorbiotin (~ 50%) and biotin sulfoxide (~ 10%); however, with maturation post partum the biotin concentration increases while bisnorbiotin and biotin sulfoxide only account for about 25% and 8% respectively. Biotin concentration remains fairly constant within one feeding but can vary substantially over 24h (Mock, et al. 1992a, Mock, et al. 1992b, Eitenmiller and Landen 1999, Mock 2001).

Microbiological assays for the determination of biotin have been described since the 1940s (Coryell, et al. 1945, Goldsmith, et al. 1982, Ford, et al. 1983, Friend, et al. 1983, Goldsmith, et al. 1983, Sakurai, et al. 2005). *Lactobacillus arabinosus* and *lactobacillus plantarum* are the common microorganisms used for the assay. Testing different conditions for hydrolysis such as hydrochloric acid or enzymatic treatment (papain, clarase, mylase P) did not produce a significant increase in biotin concentration compared to non-treated samples; however, enzymes used in the study contained large amounts of endogenous biotin compared to the milk sample causing inconclusive results (Coryell, et al. 1945). Moreover, growth-stimulating compounds such as oleic and aspartic acid can interfere with the determination resulting in an overestimate of the results (Hudson, et al. 1984).

Even though biotin itself does not contain an adequate chromophore for UV-detection, Hudson and coworkers developed a HPLC method using UV detection at 230 nm analyzing biotin in multivitamin products in the working range of 75 ó 300 g/mL. However, due to the limited number of mobile phase solvents with an appropriate UV cutoff and dissolved oxygen interference the analysis remains challenging. The latter can be resolved by degassing the solvents e.g. via sonication under vacuum. Besides limitations due to low UV-absorption, co-

elution with riboflavin and pyridoxine complicates quantification of biotin and this technique appears not to be applicable for the more complex human milk matrix.

Alternatively, Mock and DuBois developed a sequential solid phase assay using  $^{125}\text{I}$ -labeled avidin for biotin analysis in physiologic fluids, which has also been applied to human milk (Mock and DuBois 1986, Mock, et al. 1992a, Mock, et al. 1992b, Mock, et al. 1997). Earlier avidin-binding assays using  $^{14}\text{C}$ -carboxybiotin lack sensitivity for measuring physiologic concentrations of biotin due to the low specific activity of the radio labeled compound (Mock and DuBois 1986). A known amount of  $^{125}\text{I}$ -labeled avidin is incubated with varying amounts of biotin in buffer (standard curve) and with several dilutions of the unknown samples. An aliquot is then transferred to microtiter plates previously coated with biotin linked to albumin, which will occupy the remaining avidin binding sites. The wells are washed after incubation and counted individually. Increasing amounts of biotin in the standards or samples results in fewer counts bound to the plate (Mock and DuBois 1986).

### ***Choline***

Choline (*N*-trimethylethanolamine) was discovered by Strecker in 1862 and chemically synthesized in 1866. Its biosynthetic pathway was first described by duVigneaud in 1941 (Canty and Zeisel 1994, Zeisel and Holmes-McNary 2001). The colorless viscid liquid is soluble in water and in solvents such as methanol, ethanol or acetonitrile. Diluted aqueous solutions of choline are fairly stable compared to concentrated solutions, particularly when heated to 100°C. Being strongly alkaline, choline forms salts with organic and inorganic acids, which are stable in

aqueous solutions. UV and fluorometric properties have been used for detection after derivatization of choline or its metabolites (Venugopal 1985).

As a dietary component, choline is important for the structural integrity of cell membranes, methyl metabolism, cholinergic neurotransmission, transmembrane signaling and lipid and cholesterol transport and metabolism and is needed in relatively large quantities compared to the other vitamins. Choline is a precursor of the neurotransmitter acetylcholine, and betaine. Lack of choline causes cell death by apoptosis and can lead to fatty liver or liver, muscle, and DNA damage as well as in alterations in lymphocyte gene expression (Venugopal 1985, Koc, et al. 2002, Holm, et al. 2003, Zeisel and Caudill 2010). Since this vitamin is involved in the conversion of homocysteine to methionine, requirements for dietary choline must be considered in relation to methionine and folate metabolism (Zeisel 2006). Moreover, choline plays a critical role in fetal development, particularly in the brain as its availability appears to have an effect on neural tube closure and cognition. Therefore, low choline intake during pregnancy is associated with an increased risk of birth defects in the fetus. During pregnancy this nutrient is provided to the fetus across the placenta, and is secreted into human milk from the maternal circulation (Ilgol, et al. 2005, Caudill 2010, Fischer, et al. 2010).

Choline is commonly available in foods as free choline, choline esters or betaine. Good food sources include egg yolks, organ meats, spinach, nuts or wheat germ (Canty and Zeisel 1994, Zeisel and Caudill 2010). In human milk, forms of choline include free choline and its metabolites phosphocholine (PC) and glycerophosphocholine (GPC) as main contributors as well as lipophilic phosphatidylcholine (lecithin) and sphingomyelin (SM). Its concentration doubles 6

6 7 days after birth, mostly due to increasing amounts of phosphocholine and glycerophosphocholine (Holmes-McNary, et al. 1996, Holmes, et al. 2000).

One of the first approaches for analyzing choline in human milk was described by Zeisel and coworkers adapting a radioenzymatic assay introduced for choline and acetylcholine in brain. Samples were precipitated with chloroform/methanol, and the pellet washed and resuspended in chloroform/water prior to analysis. The radioenzymatic assay is based on the conversion of choline to phosphorylcholine- $^{32}\text{P}$  in the presence of choline kinase and ATP- $\gamma$ - $^{32}\text{P}$  (Reid, et al. 1971, Haubrich and Reid 1974, Haubrich, et al. 1975, Zeisel, et al. 1982, Zeisel, et al. 1986).

In the 1990s  $^1\text{H}$ -NMR was introduced for choline analysis by Holmes and coworkers. Water-soluble choline, PC and GPC were extracted with perchloric acid and neutralized with potassium hydroxide. To distinguish the metabolites from other components possessing similar chemical shift values such as carnitine, samples were run at pH 2 and 7. Field locking, quantitation and shift referencing was achieved by adding deuterated trimethylsilylpropionate. Fat-soluble metabolites were extracted using chloroform, dried and redissolved in deuterated chloroform/water containing TMS and 1,3,5-trichlorobenzene as chemical shift reference and quantitation standard respectively (Holmes, et al. 1996, Holmes, et al. 1998, Holmes, et al. 2000).

HPLC as well as GC-MS has been employed for choline analysis in human milk since the mid 1990s; more emphasis on these analytical applications has been given within the last decade (Holmes-McNary, et al. 1996, Koc, et al. 2002, Sakurai, et al. 2005, Fischer, et al. 2010). Holmes-McNary and coworkers analyzed chlorine and its metabolite present in human milk

using a GC-MS method described for tissues such as plasma, serum or liver (Pomfret, et al. 1989, Zeisel and DaCosta 1990). Samples were partitioned into an organic and aqueous phase; while the aqueous metabolites were purified via HPLC, the fat-soluble metabolites were isolated using thin-layer chromatography (TLC) prior to hydrolysis and derivatization into volatile propionyl ester. After removing the N-methylgroup, the sample is ready for GC-MS analysis. SM analysis was adapted from a spectrophotometric method described for plasma. Samples were digested in concentrated perchloric acid at 200°C and phosphorus was quantified after reaction with molybdate, spectrometrically at 825nm (Svanborg, et al. 1961). However, this laborious chromatographic approach requires different types of instrumentation for purifying each metabolite separately and their analysis and SM has to be analyzed separately, amounting to a very complex analysis. An alternative approach used HPLC analysis after hydrolysis of the sample and electrochemical detection after enzymatic reaction with choline oxidase on an immobilized enzyme column (Sakurai, et al. 2005).

Koc and co-workers introduced liquid chromatography/electrospray ionization-isotope dilution mass spectrometry (LC/ESI-IDMS) analyzing betaine, choline, acetylcholine, cytidine diphosphocholine, PC, GPC, lecithin and SM simultaneously in liver, plasma and various foods, a method also used in recent years for human milk. Samples are partitioned into organic and aqueous phases using methanol and chloroform and analyzed directly without the need for isolation and derivatization (Koc, et al. 2002, Fischer, et al. 2010). Other HPLC methods have been described for choline analysis, using different detection methods such as fluorometric detection or continuous-flow fast atom bombardment mass spectrometry (Ricny, et al. 1992, Ishimaru, et al. 1993). However, none of these methods have been used for human milk analysis.

**CONCLUSIONS:**

Multiple methods have been applied for analyzing B-vitamins in human milk samples. Commonly used methods include microbiological assays and chromatographic techniques. However, the form of the vitamin present in human milk dictates the choice of methodological approach for analysis (Table 1). Chromatographic methods are the most recent approaches for multiple B-vitamins such as thiamin, riboflavin, niacin, vitamin B6 and choline. These vitamins are usually not bound to proteins and can be analyzed individually or simultaneously, and even as different forms (i.e. riboflavin and its coenzymatic form FAD). Microbiological assays appear to be the analysis of choice for folate and panthothenic acid, although for different reasons. Pantothenic acid is mostly present in its unbound form, but the lack of chromophores in the molecule complicates the detection after chromatographic separation, whereas folate is bound to multiple enzymes and chromatographic approaches for quantitation have been shown to be inaccurate. Competitive protein binding assays using radiolabel or chemiluminescence for detection are the common approaches for analyzing biotin and vitamin B12. From this review it is evident that many methods have been used for the analysis of B vitamins in human milk. A substantial number do not give accurate information, and this review should be helpful in enabling users to better interpret the validity of published values in the literature.

**ACKNOWLEDGMENTS:**

USDA is an equal opportunity employer and provider.



## FUNDING

The authors would like to acknowledge funds by the Melinda and Bill Gates Foundation (OPP53107) provided by the University of North Carolina (Linda S. Adair and Margaret E. Bentley) and intramural USDA-ARS Project #5306-51000-003-00D.

**REFERENCES:**

- Adjalla, C., Benhayoun, S., Nicolas, J. P., Gueant, J. L. and Lambert, D. (1993). Existence of vitamin B12 analogs in biological samples: a reality. *The Journal of Nutritional Biochemistry*, **4**: 543-548.
- Adjalla, C., Lambert, D., Benhayoun, S., Berthelsen, J. G., Nicolas, J. P., Gueant, J. L. and Nexø, E. (1994). Forms of cobalamin and vitamin B12 analogs in maternal plasma, milk, and cord plasma. *The Journal of Nutritional Biochemistry*, **5**: 406-410.
- Allen, L. H. (2012). B vitamins in breast milk: relative importance of maternal status and intake, and effects on infant status and function. *Advances in Nutrition: An International Review Journal*, **3**: 362-369.
- Allen, L. H. (2009). How common is vitamin B-12 deficiency? *The American Journal of Clinical Nutrition*, **89**: 693S-696S.
- Allen, L. H. (2005). Multiple micronutrients in pregnancy and lactation: an overview. *American Journal of Clinical Nutrition*, **81**: 1206S-1212S.
- Allfrey, V., Teply, L. J., Geffen, C. and King, C. G. (1949). A fluorometric method for the determination of pteroylglutamic acid. *Journal of Biological Chemistry*, **178**: 465-481.
- Álvarez-Sánchez, B., Priego-Capote, F., Mata-Granados, J. M. and Luque de Castro, M. D. (2010). Automated determination of folate catabolites in human biofluids (urine, breast milk and serum) by on-line SPE-HILIC-MS/MS. *Journal of Chromatography A*, **1217**: 4688-4695.

- Areekul, S., Quarom, K. and Doungbarn, J. (1977). Determination of vitamin B12 and vitamin B12 binding proteins in human and cow's milk. *Modern Medicine of Asia*, **13**: 17-23.
- Baker, H., Hockstein, S., DeAngelis, B. and Holland, B. K. (2000). Thiamin status of gravidas treated for gestational diabetes mellitus compared to their neonates at parturition. *International journal for vitamin and nutrition research*, **70**: 317-320.
- Ball, G. F. M. (1997). Bioavailability and analysis of vitamins in foods. Chapman and Hall, New York, NY.
- Ball, G. F. M. (2006). Vitamins in foods: analysis, bioavailability, and stability. CRC Taylor & Francis, Boca Raton, FL.
- Ball, G. F. M. (1994). Water-soluble vitamin assays in human nutrition. Chapman & Hall, New York, NY.
- Banhidi, Z. G. (1958). Some aspects of the nutrition of *Lactobacillus fermenti* 36 in the tube assay of thiamine. *Acta Chemica Scandinavica*, **12**: 517-527.
- Bank, M. R., Kirksey, A., West, K. and Giacoia, G. (1985). Effect of storage time and temperature on folacin and vitamin C levels in term and preterm human milk. *The American Journal of Clinical Nutrition*, **41**: 235-242.
- Barger, G., Bergel, F. and Todd, R. A. (1935). Ueber das Thiochrom aus Vitamin B1 (Antineurin). *Chemische Berichte*, **67**: 2257-2262.
- Barker, H. A., Smyth, R. D., Weissbach, H., Munch-Petersen, A., Toohey, J. I., Ladd, J. N., Volcani, B. E. and Wilson, R. M. (1960). Assay, purification, and properties of the adenylocobamide coenzyme. *The Journal of Biological Chemistry*, **235**: 181-190.

- Barton-Wright, E. C. (1963). Practical methods for the microbiological assay of the vitamin B-complex and amino acids. United Trade Press, London,UK.
- Bates, C. J., Liu, D. S., Fuller, N. J. and Lucas, A. (1985). Susceptibility of riboflavin and vitamin A in breast milk to photodegradation and its implications for the use of banked breast milk in infant feeding. *Acta Paediatrica*, **74**: 40-44.
- Bates, C. J., Prentice, A. M., Watkinson, M., Morrell, P., Sutcliffe, B. A., Foord, F. A. and Whitehead, R. G. (1982a). Riboflavin requirements of lactating Gambian women: a controlled supplementation trial. *American Journal of Clinical Nutrition*, **35**: 701-709.
- Bates, C. J. and Prentice, A. (1994). Breast milk as a source of vitamins, essential minerals and trace elements. *Pharmacology and Therapeutics*, **62**: 193-220.
- Bates, C. J., Prentice, A. M., Paul, A. A., Prentice, A., Sutcliffe, B. A. and Whitehead, R. G. (1982b). Riboflavin status in infants born in rural Gambia, and the effect of a weaning food supplement. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, **76**: 253-258.
- Beck, W. S. (2001). Cobalamin (Vitamin B12). **In:** Handbook of Vitamins, pp. 463-512. Rucker, R. B., Suttie, J. W., McCormick, D. B., Machlin, L. J., Eds., Marcel Dekker Inc., New York, NY.
- Bernshaw, N. J. (1991). Does breastfeeding protect against sudden infant death syndrome? *Journal of Human Lactation*, **7**: 73-79.
- Boas, M. A. (1927). The effect of desiccation upon the nutritive properties of egg-white. *Biochemistry Journal*, **3**: 712-725.

- Böhm, V., Peiker, G., Starker, A., Weske, E., Schaarmann, G., Schubert, R., Bitsch, R. and Flachowsky, G. (1997). Vitamin B1, B2, A and E and beta-carotene content in transitional breast milk and comparative studies in maternal and umbilical cord blood. *Zeitschrift für Ernährungswissenschaft*, **36**: 214-219.
- Borschel, M. W., Kirksey, A. and Hannemann, R. E. (1986). Effects of vitamin B6 intake on nutriture and growth of young infants. *American Journal of Clinical Nutrition*, **43**: 7-15.
- Bourquin, A. and Sherman, H. C. (1931). Quantitative determination of vitamin G (B2). *Journal of the American Chemical Society*, **53**: 3501-3505.
- Bridges, J. W., Davies, D. S. and Williams, R. T. (1966). Fluorescence studies on some hydroxypyridines including compounds of the vitamin B6 group. *Biochemical Journal*, **98**: 451-468.
- Brink-Shoemaker, C., Cruickshank, D. W. J., Hodgkin, D. C., Kamper, M. J. and Pilling, D. (1964). The structure of vitamin B12. VI. The structure of crystals of vitamin B12 grown from and immersed in water. *Proceedings of the Royal Society of London A*, **278**: 1-26.
- Brody, T. and Shane, B. (2001). Folic acid. *Handbook of Vitamins*, pp. 427-462. Rucker, R. B., Suttie, J. W., McCormick, D. B., Machlin, L. J., Eds., Marcel Dekker Inc., New York, NY.
- Brown, C. M., Smith, A. M. and Picciano, M. F. (1986). Forms of human milk folacin and variation patterns. *Journal of Pediatric Gastroenterology and Nutrition*, **5**: 278-282.
- Buehler, B. A. (2011). Vitamin B2: Riboflavin. *Journal of Evidence-Based Complementary & Alternative Medicine*, **16**: 88-90.

- Butte, N. F. and Calloway, D. H. (1981). Evaluation of lactational performance of Navajo women. *The American Journal of Clinical Nutrition*, **34**: 2210-2215.
- Canty, D. J. and Zeisel, S. H. (1994). Lecithin and choline in human health and disease. *Nutrition Reviews*, **52**: 327-339.
- Casterline, J. E., Allen, L. H. and Ruel, M. T. (1997). Vitamin B-12 deficiency is very prevalent in lactating Guatemalan women and their infants at three months postpartum. *Journal of Nutrition*, **127**: 1966-1972.
- Caudill, M. A. (2010). Pre-and postnatal health: evidence of increased choline needs. *Journal of the American Dietetic Association*, **110**: 1198-1206.
- Chase, E. and Sherman, H. (1931). A quantitative study of the determination of the antineuritic vitamin B. *Journal of the American Chemical Society*, **53**: 3506-3510.
- Chin, H. B. (1985). Vitamin B12. **In**: Methods of Vitamin Assay, pp. 497-514. Augustin, J., Klein, B. P., Becker, D. and Venugopal, P., Eds., John Wiley & Sons, New York, NY.
- Conner, R. and Straub, G. (1941). Combined determination of riboflavin and thiamin in food products. *Industrial and Engineering Chemistry Analytical Edition*, **13**: 385-388.
- Cooperman, J. M., Dweck, H. S., Newman, L. J., Garbarino, C. and Lopez, R. (1982). The folate in human milk. *The American Journal of Clinical Nutrition*, **36**: 576-580.
- Coryell, M. N., Harris, M. E., Miller, S., Williams, H. H. and Macy, I. G. (1945). Human milk studies; nicotinic acid, pantothenic acid and biotin contents of colostrum and mature human milk. *American Journal of Diseases of Children*, **70**: 150-161.

- Cosner, L. B. and Schuck, C. (1948). Suitability of the Bourquin-Sherman diet for riboflavin assays and results obtained in the assay of evaporated milk. *The Journal of Nutrition*, **35**: 725-732.
- Craft, I. L., Matthews, D. M. and Linnell, J. C. (1971). Cobalamins in human pregnancy and lactation. *Journal of Clinical Pathology*, **24**: 449-455.
- Day, P. L., Langston, W. C. and Darby, W. J. (1938). Failure of Nicotinic Acid to Prevent Nutritional Cytopenia in the Monkey. *Proceedings of the Society for Experimental Biology and Medicine*, **38**: 860-863.
- Deegan, K. L., Jones, K. M., Zuleta, C., Ramirez-Zea, M., Lildballe, D. L., Nexo, E. and Allen, L. H. (2012). Breast milk vitamin B-12 concentrations in Guatemalan women are correlated with maternal but not infant vitamin B-12 status at 12 months postpartum. *Journal of Nutrition*, **142**: 112-116.
- Deibel, R. H., Evans, J. B. and Niven Jr, C. (1957). Microbiological assay for thiamin using *Lactobacillus viridescens*. *Journal of Bacteriology*, **74**: 818-821.
- Donangelo, C. M., Trugo, N. M., Koury, J. C., Barreto, S. M. I., Freitas, L. A., Feldheim, W. and Barth, C. (1989). Iron, zinc, folate and vitamin B12 nutritional status and milk composition of low-income Brazilian mothers. *European Journal of Clinical Nutrition*, **43**: 253-266.
- Donelson, E. G. and Macy, I.G. (1934). Human Milk Studies: XII. The vitamin B and vitamin G content before and during maternal consumption of yeast. *The Journal of Nutrition* **7**: 231-249.

- Eitenmiller, R. R. and De Souza, S. (1985). Niacin. **In:** Methods of Vitamin Assay, pp. 385-398. Augustin, J., Klein, B.P., Becker, D. and Venugopal, P., John Wiley & Sons, New York, NY.
- Eitenmiller, R. R., Bryan, W. D., Khalsa, I. K., Feeley, R. M. and Barnhart, H. M. (1984). Folate content of human milk during early lactational stages. *Nutrition Research*, **4**: 391-397.
- Eitenmiller, R. R. and Landen Jr., W.O. (1999). Vitamin Analysis for the Health and Food Sciences. CRC Press LLC, Athens, GA.
- Ek, J. (1983). Plasma, red cell, and breast milk folacin concentrations in lactating women. *American Journal of Clinical Nutrition* **38**: 929.
- Ellefson, W. C. (1985). Thiamin. **In:** Methods of Vitamin Assay, pp. 349-364. Augustin, J., Klein, B. P., Becker, D. and Venugopal, P., Eds., John Wiley & Sons, New York, NY.
- Emmett, P. M. and Rogers, I. S. (1997). Properties of human milk and their relationship with maternal nutrition. *Early Human Development*, **49**: S7-S28.
- Fernández-Muiño, M. Á., Sancho-Ortiz, M. T. and Valls-Garcia, F. (2008). Water-Soluble Vitamins. **In:** Methods of Analysis for Functional Foods and Nutraceuticals, pp. 401-432. Hurst, J., Ed., CRC Press, Boca Raton, FL.
- Fischer, L. M., da Costa, K. A., Galanko, J., Sha, W., Stephenson, B., Vick, J. and Zeisel, S. H. (2010). Choline intake and genetic polymorphisms influence choline metabolite concentrations in human breast milk and plasma. *The American Journal of Clinical Nutrition*, **92**: 336-346.



- Ford, J. E., Zechalko, A., Murphy, J. and Brooke, O. G. (1983). Comparison of the B vitamin composition of milk from mothers of preterm and term babies. *Archives of Disease in Childhood*, **58**: 367-372.
- Friend, B. A., Shahani, K. M., Long, C. A. and Vaughn, L. A. (1983). The effect of processing and storage on key enzymes, B vitamins, and lipids of mature human milk I. Evaluation of fresh samples and effects of freezing and frozen storage. *Pediatric Research*, **17**: 61-64.
- Gatti, R. and Gioia, M. G. (2005). Liquid chromatographic determination with fluorescence detection of B6 vitamers and riboflavin in milk and pharmaceuticals. *Analytica Chimica Acta*, **538**: 135-141.
- Gerrits, J., Eidhof, H., Brunnekreeft, J. W. I. and Hessels, J. (1997). Determination of thiamin and thiamin phosphates in whole blood by reversed-phase liquid chromatography with precolumn derivatization. *Methods in Enzymology*, **279**: 74-82.
- Gibson, R. S. (1990). Principles of Nutritional Assessment. Oxford University Press, New York, NY.
- Goldsmith, S. J., Eitenmiller, R. R., Toledo, R. T. and Barnhart, H. M. (1983). Effects of processing and storage on the water-soluble vitamin content of human milk. *Journal of Food Science*, **48**: 994-995.
- Goldsmith, S. J., Eitenmiller, R. R., Feeley, R. M., Barnhart, H. M. and Maddox, F. C. (1982). Biotin content of human milk during early lactational stages<sup>1</sup>. *Nutrition Research*, **2**: 579-583.

- Greer, F. R. (2001). Do breastfed infants need supplemental vitamins? *Pediatric Clinics of North America*, **48**: 415-423.
- Gyorgy, P. (1936). Quantitative estimation of lactoflavin and of vitamin B6 in cow's milk and in human milk. *Experimental Biology and Medicine*, **35**: 204-207.
- Gyorgy, P. (1934). Vitamin B2 and the pellagra-like dermanitis of rats. *Nature*, **133**: 498-499.
- Gyorgy, P. and Eckardt, R. E. (1939). Vitamin B6 and skin lesions in rats. *Nature*, **144**: 512-512.
- Hälvin, K., Paalme, T., Nisamedtinov, I. (2013). Comparison of different extraction methods for simultaneous determination of B complex vitamins in nutritional yeast using LC/MS-TOF and stable isotope dilution. *Analytical and Bioanalytical Chemistry*, **405**: 1213-1222.
- Hamaker, B., Kirksey, A., Ekanayake, A. and Borschel, M. (1985). Analysis of B-6 vitamers in human milk by reverse-phase liquid chromatography. *American Journal of Clinical Nutrition*, **42**: 650-655.
- Hampel, D., York, E. R. and Allen, L. H. (2012). Ultra-Performance Liquid Chromatography Tandem Mass-Spectrometry (UPLC-MS/MS) for the Rapid, Simultaneous Analysis of Thiamin, Riboflavin, Flavin Adenine Dinucleotide, Nicotinamide and Pyridoxal in Human Milk. *Journal of Chromatography B*, **903**: 7-13.
- Han, Y. H., Yon, M., Han, H. S., Kim, K. Y., Tamura, T. and Hyun, T. H. (2010). Folate contents in human milk and casein-based and soya-based formulas, and folate status in Korean infants. *British Journal of Nutrition*, **101**: 1769-1774.

- Haubrich, D. R. and Reid, W. D. (1974). Use of choline kinase in the radioisotopic estimation of brain choline and acetylcholine. **In:** Choline and acetylcholine: Handbook of chemical assay methods, pp. 33-45. Hanin, I. Ed., Raven Press, New York, NY.
- Haubrich, D. R., Wang, P. F. L., Clody, D. E. and Wedeking, P. W. (1975). Increase in rat brain acetylcholine induced by choline or deanol. *Life Sciences*, **17**: 975-980.
- Heidelberger, C., Abraham, E. P. and Lepkovsky, S. (1949). Tryptophan Metabolism: II. Concerning the mechanism of the mammalian conversion of tryptophan into nicotinic acid. *The Journal of Biological Chemistry*, **179**: 151-155.
- Herbert, V. (1961). The assay and nature of folic acid activity in human serum. *Journal of Clinical Investigation*, **40**: 81-91.
- Herbert, V. (1996). Vitamin B12. **In:** Present Knowledge in Nutrition, pp. 191-205. Ziegler, E. and Filer, L., Jr., Eds., International Life Sciences Institute Press, Washington, DC.
- Hodgkin, D. C., Kamper, J., Mackay, M., Pickworth, J., Trueblood, K. N. and White, J. G. (1956). Structure of vitamin B12. *Nature*, **178**: 64-66.
- Hodgkin, D. C., Lindsey, J., Mackay, M. and Trueblood, K. N. (1962). The structure of vitamin B12. IV. The X-ray analysis of air-dried crystals of B12. *Proceedings of the Royal Society of London A*, **266**: 475-493.
- Holm, P. I., Ueland, P. M., Kvalheim, G. and Lien, E. A. (2003). Determination of choline, betaine, and dimethylglycine in plasma by a high-throughput method based on normal-phase chromatography-tandem mass spectrometry. *Clinical Chemistry*, **49**: 286-294.

- Holmes-McNary, M. Q., Cheng, W. L., Mar, M. H., Fussell, S. and Zeisel, S. H. (1996). Choline and choline esters in human and rat milk and in infant formulas. *The American Journal of Clinical Nutrition*, **64**: 572-576.
- Holmes, H. C., Snodgrass, G. J. A. I. and Iles, R. A. (2000). Changes in the choline content of human breast milk in the first 3 weeks after birth. *European Journal of Pediatrics*, **159**: 198-204.
- Holmes, H., Snodgrass, G. J. A. I. and Iles, R. A. (1996). The choline content of human breast milk expressed during the first few weeks of lactation. *Biochemical Society Transactions*, **24**: 350S.
- Holmes, H., Graeme, J. and Iles, R. (1998). Choline metabolism in the neonatal period. *Biochemical Society Transactions*, **26**: S94.
- Honzik, T., Adamovicova, M., Smolka, V., Magner, M., Hrubá, E. and Zeman, J. (2010). Clinical presentation and metabolic consequences in 40 breastfed infants with nutritional vitamin B12 deficiency-What have we learned? *European Journal of Paediatric Neurology*, **14**: 488-495.
- Hoshi, S. (1956). The detection of vitamin B1 in mother's milk. *Sendai-shi Hokenjo Gyoho*, **3**: 133-134.
- Hudson, T. S., Subramanian, S. and Allen, R. J. (1984). Determination of pantothenic acid, biotin, and vitamin B12 in nutritional products. *Journal of the Association of Official Analytical Chemists*, **67**: 994-998.

- Hyun, T. H. and Tamura, T. (2005). Trienzyme extraction in combination with microbiologic assay in food folate analysis: an updated review. *Experimental Biology and Medicine*, **230**: 444-454.
- I.O.M. (1998). Dietary Reference Intakes for thiamin, riboflavin, niacin, vitamin B6, folate, vitamin B12, pantothenic acid, biotin, and choline. National Academy Press, Washington, DC.
- Ilcol, Y. O., Ozbek, R., Hamurtekin, E. and Ulus, I. H. (2005). Choline status in newborns, infants, children, breast-feeding women, breast-fed infants and human breast milk. *The Journal of Nutritional Biochemistry*, **16**: 489-499.
- Ishimaru, H., Ikarashi, Y. and Maruyama, Y. (1993). Use of high-performance liquid chromatography continuous-flow fast atom bombardment mass spectrometry for simultaneous determination of choline and acetylcholine in rodent brain regions. *Biological Mass Spectrometry*, **22**: 681-686.
- Israel-Ballard, K. A., Abrams, B. F., Coutsoudis, A., Sibeko, L. N., Cheryk, L. A. and Chantry, C. J. (2008). Vitamin content of breast milk from HIV-1 infected mothers before and after flash-heat treatment. *Journal of Acquired Immune Deficiency Syndromes*, **48**: 444-449.
- Jadhav, M., Webb, J. K. G., Vaishnava, S. and Baker, S. J. (1962). Vitamin-B12 deficiency in Indian infants. *The Lancet*, **280**: 903-907.
- Jathar, V. S., Kamath, S. A., Parikh, M. N., Rege, D. V. and Satoskar, R. S. (1970). Maternal milk and serum vitamin B12, folic acid, and protein levels in Indian subjects. *Archives of Disease in Childhood*, **45**: 236-241.

- Jukes, T. H. (1939). Pantothenic acid and the filtrate (chick anti-dermatitis) factor. *Journal of the American Chemical Society*, **61**: 975-976.
- Kang-Yoon, S. A., Kirksey, A., Giacoia, G. and West, K. (1992). Vitamin B-6 status of breast-fed neonates: influence of pyridoxine supplementation on mothers and neonates. *American Journal of Clinical Nutrition*, **56**: 548-558.
- Karra, M. V., Udipi, S. A., Kirksey, A. and Roepke, J. L. (1986). Changes in specific nutrients in breast milk during extended lactation. *American Journal of Clinical Nutrition*, **43**: 495-503.
- Keizer, S. E., Gibson, R. S. and O'Connor, D. L. (1995). Postpartum folic acid supplementation of adolescents: impact on maternal folate and zinc status and milk composition. *American Journal of Clinical Nutrition*, **62**: 377-384.
- Khambalia, A., Latulippe, M. E., Campos, C., Merlos, C., Villalpando, S., Picciano, M. F. and O'Connor, D. L. (2006). Milk folate secretion is not impaired during iron deficiency in humans. *The Journal of Nutrition*, **136**: 2617-2624.
- Kinnersley, H. W. and Peters, R. A. (1938). Note upon the preparation of crude cocarboxylase from vitamin B1 by yeast. *Biochemical Journal*, **32**: 697-698.
- Koc, H., Mar, M. H., Ranasinghe, A., Swenberg, J. A. and Zeisel, S. H. (2002). Quantitation of choline and its metabolites in tissues and foods by liquid chromatography/electrospray ionization-isotope dilution mass spectrometry. *Analytical Chemistry*, **74**: 4734-4740.
- Kolhouse, J. F., Kondo, H., Allen, N. C., Podell, E. and Allen, R. H. (1978). Cobalamin analogues are present in human plasma and can mask cobalamin deficiency because

- current radioisotope dilution assays are not specific for true cobalamin. *New England Journal of Medicine*, **299**: 785-792.
- Krehl, W. A., Strong, F. M. and Elvehjem, C. A. (1943). Determination of nicotinic acid. Modifications in the microbiological method. *Industrial and Engineering Chemistry*, **15**: 471-475.
- Lau, K. S., Gottlieb, C., Wasserman, L. R. and Herbert, V. (1965). Measurement of serum vitamin B12 level using radioisotope dilution and coated charcoal. *Blood*, **26**: 202-214.
- Leklem, J. E. (2001). Vitamin B6. . **In**: Handbook of Vitamins, pp. 339-396. Rucker, R. B., Suttie, J. W., McCormick, D. B., Machlin, L. J., Eds., Marcel Dekker Inc., New York, NY.
- Lenhert, P. G. (1968). The Structure of Vitamin B12. VII. The X-ray analysis of the vitamin B12 coenzyme. *Proceedings of the Royal Society of London A*, **303**: 45-84.
- Lenhert, P. G. and Hodgkin, D. C. (1961). Structure of the 5, 6-dimethylbenzimidazolylcobamide coenzyme. *Nature*, **192**: 937-938.
- Lildballe, D. L., Hardlei, T. F., Allen, L. H. and Nexø, E. (2009). High concentrations of haptocorrin interfere with routine measurement of cobalamins in human serum and milk. A problem and its solution. *Clinical Chemistry and Laboratory Medicine*, **47**: 182-187.
- Lim, H. S., Mackey, A. D., Tamura, T., Wong, S. C. and Picciano, M. F. (1998). Measurable human milk folate is increased by treatment with  $\alpha$ -amylase and protease in addition to folate conjugase. *Food Chemistry*, **63**: 401-407.

- Lindemaus, J. (2000). Cobalamins. **In:** Modern Chromatographic Analysis of Vitamins, pp. 511-554. De Leenheer, A. P., Lambert, W. E., Van Bocxlaer, J. F. Eds., Marcel Dekker Inc., New York, NY.
- Lönnerdal, B. (1986). Effects of maternal dietary intake on human milk composition. *The Journal of Nutrition*, **116**: 499-513.
- Mackey, A. D. and Picciano, M. F. (1999). Maternal folate status during extended lactation and the effect of supplemental folic acid. *The American Journal of Clinical Nutrition*, **69**: 285-292.
- Macy, I. G. (1949). Composition of human colostrum and milk. *Archives of Pediatrics and Adolescent Medicine*, **78**: 589-603.
- Maggiorelli, E. (1958). Sodium salts of riboflavin-5-phosphoric acid; physical, chemical & spectrophotometric properties; quantitative determination in various types of vials for hypodermic use. *Bollettino Chimico Farmaceutico*, **97**: 481-490.
- McCormick, D. B. (1968). Nature of the intramolecular complex of flavine adenine dinucleotide. **In:** Molecular Associations in Biology, pp. 377-392. Pullman, B., Ed., Academic Press, New York, NY.
- McCosh, S. S., Macy, I. G. and Hunscher, H. A. (1931). Human Milk Studies: VI. Vitamin potency as influenced by supplementing the maternal diet with yeast. *The Journal of Biological Chemistry*, **90**: 1-13.
- McGready, R., Simpson, J. A., Cho, T., Dubowitz, L., Changbumrung, S., Böhm, V., Munger, R. G., Sauberlich, H. E., White, N. J. and Nosten, F. (2001). Postpartum thiamine deficiency in a Karen displaced population. *American Journal of Clinical Nutrition*, **74**: 808-813.



- McPhee, A. J., Davidson, G. P., Leahy, M. and Beare, T. (1988). Vitamin B12 deficiency in a breast fed infant. *Archives of Disease in Childhood*, **63**: 921-923.
- Metz, J., Zalusky, R. and Herbert, V. (1968). Folic acid binding by serum and milk. *The American Journal of Clinical Nutrition*, **21**: 289-297.
- Mikheeva, E. V., Martynyuk, O. A., Slepchenko, G. B. and Anisimova, L. S. (2009). Study of the voltammetric behavior of vitamin B 2 and the development of a procedure for its determination in breast milk. *Journal of Analytical Chemistry*, **64**: 731-734.
- Mock, D. M. and DuBois, D. B. (1986). A sequential, solid-phase assay for biotin in physiologic fluids that correlates with expected biotin status. *Analytical Biochemistry*, **153**: 272-278.
- Mock, D. M., Mock, N. I. and Dankle, J. A. (1992a). Secretory patterns of biotin in human milk. *The Journal of Nutrition*, **122**: 546-552.
- Mock, D. M., Mock, N. I. and Langbehn, S. E. (1992b). Biotin in human milk: methods, location, and chemical form. *The Journal of Nutrition*, **122**: 535-545.
- Mock, D. M., Mock, N. I. and Stratton, S. L. (1997). Concentrations of biotin metabolites in human milk. *The Journal of Pediatrics*, **131**: 456-458.
- Mock, D. M. (2001). Biotin. **In:** Handbook of Vitamins, pp. 397-426. Rucker, R. B., Suttie, J. W., McCormick, D. B., Machlin, L. J., Eds., Marcel Dekker Inc., New York, NY.
- Morrison, L. A. and Driskell, J. A. (1985). Quantities of B6 vitamers in human milk by high-performance liquid chromatography. Influence of maternal vitamin B6 status. *Journal of Chromatography*, **337**: 249-258.

- Nail, P. A., Thomas, M. R. and Eakin, R. (1980). The effect of thiamin and riboflavin supplementation on the level of those vitamins in human breast milk and urine. *American Journal of Clinical Nutrition*, **33**: 198-204.
- O'Connor, D. L., Green, T. and Picciano, M. F. (1997). Maternal folate status and lactation. *Journal of Mammary Gland Biology and Neoplasia*, **2**: 279-289.
- O'Connor, D. L., Tamura, T. and Picciano, M. F. (1991). Pteroylpolyglutamates in human milk. *The American Journal of Clinical Nutrition*, **53**: 930-934.
- Ortega, R. M., Martínez, R. M., Andrés, P., Marín-Arias, L. and López-Sobaler, A. M. (2004). Thiamin status during the third trimester of pregnancy and its influence on thiamin concentrations in transition and mature breast milk. *British Journal of Nutrition*, **92**: 129-136.
- Owen, E. C. and West, D. W. (1971). Isolation and identification of 7, 8-dimethyl-10-(2'-hydroxyethyl) isoalloxazine from natural sources. *Methods in Enzymology*, **18**: 574-579.
- Owens, J. O., Holstege, D. M., Clifford, A. J. (2007). High-Throughput Method for the Quantitation of Total Folate in Whole Blood Using LC-MS/MS. *Journal of Agricultural and Food Chemistry*, **55**: 3292-3297.
- Parrish, W. P., Loy, H. W. and Kline, O. L. (1956). Further studies on the yeast method for vitamin B-6. *Journal of the Association of Official Agricultural Chemists*, **39**: 157-161.
- Parrish, W. P., Loy, H. W. and Kline, O. L. (1955). A study of the yeast method for vitamin B-6. *Journal of the Association of Official Agricultural Chemists*, **38**: 506-513.
- Patel, K. D. and Lovelady, C. A. (1998). Vitamin B12 status of East Indian vegetarian lactating women living in the United States. *Nutrition Research*, **18**: 1839-1846.

- Pearson, P. B., Darnell, A. L. and Weir, J. (1946). The thiamine, riboflavin, nicotinic acid and pantothenic acid content of colostrum and milk of the cow and ewe. *The Journal of Nutrition*, **31**: 51-57.
- Pelletier, O. and Campbell, J. A. (1959). A modified procedure for the determination of niacin in cereal products. *Journal of the Association of Official Analytical Chemists*, **42**: 625-630.
- Picciano, M. F. (1998). Human milk: nutritional aspects of a dynamic food. *Biology of the Neonate*, **74**: 84-93.
- Plesofsky, N. S. (2001). Pantothenic acid. **In:** Handbook of Vitamins, pp. 317-338. Rucker, R. B., Suttie, J. W., McCormick, D. B., Machlin, L. J., Eds., Marcel Dekker Inc., New York, NY.
- Polansky, M. M. (1981). Microbiological assay of vitamin B-6 in foods. **In:** Methods in vitamin B6 nutrition, pp. 21-44. Leklem, J.E., Reynolds, R.D., Plenum Press, New York, NY.
- Polansky, M. M., Reynolds, R. D. and Vanderslice, J. T. (1985). Vitamin B6. **In:** Methods of Vitamin Assay, pp. 417. Augustin, J., Klein, B., Becker, D. and Venugopal, P., John Wiley & Sons, New York, NY.
- Pomfret, E. A., daCosta, K. A., Schurman, L. L. and Zeisel, S. H. (1989). Measurement of choline and choline metabolite concentrations using high-pressure liquid chromatography and gas chromatography-mass spectrometry. *Analytical Biochemistry*, **180**: 85-90.
- Powers, H. J. (2003). Riboflavin (vitamin B-2) and health. *American Journal of Clinical Nutrition*, **77**: 1352-1360.
- Pratt, J. P. and Hamil, B. M. (1951). Metabolism of Women during the Reproductive Cycle. *The Journal of Nutrition*, **44**: 141-157.

- Prebluda, H. J. and McCollum, E. V. (1939). A chemical reagent for thiamine. *The Journal of Biological Chemistry*, **127**: 495-503.
- Prentice, A. M., Roberts, S. B., Prentice, A., Paul, A. A., Watkinson, M., Watkinson, A. A. and Whitehead, R. G. (1983). Dietary supplementation of lactating Gambian women. I. Effect on breast-milk volume and quality. *Human Nutrition Clinical Nutrition*, **37**: 53-64.
- Ramasastri, B. V. (1965). Folate activity in human milk. *British Journal of Nutrition*, **19**: 581-586.
- Reid, W. D., Haubrich, D. R. and Krishna, G. (1971). Enzymic radioassay for acetylcholine and choline in brain. *Analytical Biochemistry*, **42**: 390-397.
- Rettenmaier, R. and Vuilleumier, J. P. (1983). A simple method for the determination of riboflavin in human milk. *International Journal of Vitamin and Nutrition Research* **53**: 32-35.
- Rickes, E. L., Brink, N. G., Koniusky, F. R., Wood, T. R. and Folkers, K. (1948). Crystalline Vitamin B12. *Science*, **107**: 396-397.
- í ny, J., Tu ek, S. and Vin-, I. (1992). Sensitive method for HPLC determination of acetylcholine, choline and their analogues using fluorometric detection. *Journal of Neuroscience Methods*, **41**: 11-17.
- Rindi, G. (1982). Metabolism of thiamin and its phosphoric esters in different regions of the nervous system: a new approach. *Acta Vitaminologica et Enzymologica*, **4**: 59-68.
- Roderuck, C. E. and Coryell, M. N. (1945). Human milk studies; free and total riboflavin contents of colostrum and mature human milk. *American Journal of Diseases of Children*, **70**: 171-175.

- Roderuck, C. E., Williams, H. and Macy, I. G. (1945). Human milk studies; free and total thiamine contents of colostrum and mature human milk. *American Journal of Diseases of Children*, **70**: 162-170.
- Roepke, J. L. and Kirksey, A. (1979). Vitamin B6 nutriture during pregnancy and lactation. I. Vitamin B6 intake, levels of the vitamin in biological fluids, and condition of the infant at birth. *American Journal of Clinical Nutrition*, **32**: 2249-2256.
- Ronnholm, K. A. (1986). Need for riboflavin supplementation in small prematures fed with human milk. *American Journal of Clinical Nutrition*, **43**: 1-6.
- Ross, G. I. M. (1952). Vitamin B12 assay in body fluids using *Euglena gracilis*. *Journal of Clinical Pathology*, **5**: 250-256.
- Roughead, Z. K. and McCormick, D. B. (1990). Flavin composition of human milk. *American Journal of Clinical Nutrition*, **52**: 854-857.
- Russell, L. F. and Vanderslice, J. T. (1992). Non-degradative extraction and simultaneous quantitation of riboflavin, flavin mononucleotide, and flavin adenine dinucleotide in foods by HPLC. *Food Chemistry*, **43**: 151-162.
- Saidi, B. and Warthesen, J. J. (1983). Influence of pH and light on the kinetics of vitamin B6 degradation. *Journal of Agricultural and Food Chemistry*, **31**: 876-880.
- Sakurai, T., Furukawa, M., Asoh, M., Kanno, T., Kojima, T. and Yonekubo, A. (2005). Fat-soluble and water-soluble vitamin contents of breast milk from Japanese women. *Journal of Nutritional Science and Vitaminology*, **51**: 239-247.

- Samson, R. R. and McClelland, D. B. (1980). Vitamin B12 in human colostrum and milk. Quantitation of the vitamin and its binder and the uptake of bound vitamin B12 by intestinal bacteria. *Acta Paediatrica Scandinavica*, **69**: 93-99.
- Sandberg, D. P., Begley, J. A. and Hall, C. A. (1981). The content, binding, and forms of vitamin B12 in milk. *American Journal of Clinical Nutrition*, **34**: 1717-1724.
- Sarett, H. P., Perlzweig, W. A. and Levy, E. D. (1940). Synthesis and Excretion of Trigonelline. *The Journal of Biological Chemistry*, **135**: 483-485.
- Sealock, R. R. and Goodland, R. L. (1944). Interference in the Determination of Thiamine with the diazotized p-Aminophenone Reagent. *The Journal of Biological Chemistry*, **154**: 63-68.
- Selhub, J. (1989). Determination of tissue folate composition by affinity chromatography followed by high-pressure ion pair liquid chromatography. *Analytical Biochemistry*, **182**: 84-93.
- Shah, J. J. (1985). Riboflavin. **In**: Methods of Vitamin Assay, pp. 365-384. Augustin, J., Klein, B. P., Becker, D. and Venugopal, P., John Wiley & Sons, New York, NY.
- Shamsia, S. M. (2009). Nutritional and therapeutic properties of camel and human milks. *International Journal of Genetics and Molecular Biology*, **1**: 52-58.
- Shane, B., Tamura, T. and Stokstad, E. L. (1980). Folate assay: a comparison of radioassay and microbiological methods. *Clinica Chimica Acta*, **100**: 13-19.
- Shaw, G. M., Schaffer, D., Velie, E. M., Morland, K. and Harris, J. A. (1995). Periconceptional vitamin use, dietary folate, and the occurrence of neural tube defects. *Epidemiology*, **6**: 219-226.

- Shi, Y., Sun, G., Zhang, Z., Deng, X., Kang, X., Liu, Z., Ma, Y. and Sheng, Q. (2011). The chemical composition of human milk from Inner Mongolia of China. *Food Chemistry*, **127**: 1193-1198.
- Shive, W., Sibley, M. E. and Rogers, L. L. (1951). Replacement of vitamin B12 by desoxynucleotides in promoting growth of certain lactobacilli. *Journal of the American Chemical Society*, **73**: 867-868.
- Slater, E. C. and Rial, J. (1942). Thiamine (vitamin Bi) content of human milk. *Medicinal Journal of Australia*, **1**: 3-12.
- Smith, A. M., Picciano, M. F. and Deering, R. H. (1985). Folate intake and blood concentrations of term infants. *The American Journal of Clinical Nutrition*, **41**: 590-598.
- Smith, A. M., Picciano, M. F. and Deering, R. H. (1983). Folate supplementation during lactation: maternal folate status, human milk folate content, and their relationship to infant folate status. *Journal of Pediatric Gastroenterology and Nutrition*, **2**: 622-628.
- Smith, E. L. (1948). Purification of anti-pernicious anemia factors from liver. *Nature*, **161**: 638-639.
- Sneed, S. M., Zane, C. and Thomas, M. R. (1981). The effects of ascorbic acid, vitamin B6, vitamin B12, and folic acid supplementation on the breast milk and maternal nutritional status of low socioeconomic lactating women. *The American Journal of Clinical Nutrition*, **34**: 1338-1346.
- Snell, E. E. (1944). The vitamin activities of "pyridoxal" and "pyridoxamine". *The Journal of Biological Chemistry*, **154**: 313-314.

- Snell, E. E. and Wright, L. D. (1941). A microbiological method for the determination of nicotinic acid. *The Journal of Biological Chemistry*, **139**: 675-686.
- Solov'eva, E. I. and Khrustaleva, V. P. (1954). Certain data on thiamine content in human milk. *Voprosy Pitania*, **13**: 16-20.
- Song, W. O., Chan, G. M., Wyse, B. W. and Hansen, R. G. (1984). Effect of pantothenic acid status on the content of the vitamin in human milk. *The American Journal of Clinical Nutrition*, **40**: 317-324.
- Specker, B. L., Black, A., Allen, L. and Morrow, F. (1990). Vitamin B-12: low milk concentrations are related to low serum concentrations in vegetarian women and to methylmalonic aciduria in their infants. *The American Journal of Clinical Nutrition*, **52**: 1073-1076.
- Spies, T. O., Cooper, C. and Blankenhorn, M. A. (1938). The use of nicotinic acid in the treatment of pellagra. *Journal of the American Medical Association*, **110**: 622-627.
- Storvick, C. A., Benson, E. M., Edwards, M. A. and Woodring, M. J. (1964). Chemical and microbiological determination of vitamin B6. *Methods of Biochemical Analysis*, **12**: 183-276.
- Strohecker, R. and Henning, H. M. (1965). Vitamin Assay Tested Methods. Verlag Chemie, Darmstadt, GER.
- Stuetz, W., Carrara, V. I., McGready, R., Lee, S. J., Erhardt, J. G., Breuer, J., Biesalski, H. K. and Nosten, F. H. (2011). Micronutrient status in lactating mothers before and after introduction of fortified flour: cross-sectional surveys in Maela refugee camp. *European Journal of Nutrition*, **51**: 425-434.



- Stuetz, W., Carrara, V. I., McGready, R., Lee, S. J., Biesalski, H. K. and Nosten, F. H. (2012). Thiamine Diphosphate in Whole Blood, Thiamine and Thiamine Monophosphate in Breast-Milk in a Refugee Population. *PLoS ONE*, **7**:e36280.
- Styslinger, L. and Kirksey, A. (1985). Effects of different levels of vitamin B-6 supplementation on vitamin B-6 concentrations in human milk and vitamin B-6 intakes of breastfed infants. *American Journal of Clinical Nutrition*, **41**: 21-31.
- Svanborg, A., Svennerholm, L., Myrén, I., Soomägi, M. R., Claesson, B. and Rohman, S. (1961). Plasma total lipid, cholesterol, triglycerides, phospholipids and free fatty acids in a healthy Scandinavian population. *Acta Medica Scandinavica*, **169**: 43-49.
- Swiatlo, N., O'Connor, D. L., Andrews, J. and Picciano, M. F. (1990). Relative folate bioavailability from diets containing human, bovine and goat milk. *Journal of Nutrition*, **120**: 172-177.
- Tamura, T. and Picciano, M. F. (2006a). Folate and human reproduction. *The American Journal of Clinical Nutrition*, **83**: 993-1016.
- Tamura, T. and Picciano, M. F. (2006b). Folate Determination in Human Milk. *Journal of Nutritional Science and Vitaminology*, **52**: 161-161.
- Tamura, T., Shin, Y., Williams, M. and Stokstad, E. (1972). *Lactobacillus casei* response to pteroylpolyglutamates. *Analytical Biochemistry*, **49**: 517-521.
- Tamura, T., Yoshimura, Y. and Arakawa, T. (1980). Human milk folate and folate status in lactating mothers and their infants. *The American Journal of Clinical Nutrition*, **33**: 193-197.

- Thiele, V. F. and Brin, M. (1966). Chromatographic Separation and Microbiologic Assay of Vitamin B6 in Tissues from Normal and Vitamin B6-depleted Rats. *The Journal of Nutrition*, **90**: 347-353.
- Thomas, M. R., Kawamoto, J., Sneed, S. M. and Eakin, R. (1979). The effects of vitamin C, vitamin B6, and vitamin B12 supplementation on the breast milk and maternal status of well-nourished women. *American Journal of Clinical Nutrition*, **32**: 1679-1685.
- Thomas, M. R., Sneed, S. M., Wei, C., Nail, P. A., Wilson, M. and Sprinkle, E. (1980). The effects of vitamin C, vitamin B6, vitamin B12, folic acid, riboflavin, and thiamin on the breast milk and maternal status of well-nourished women at 6 months postpartum. *American Journal of Clinical Nutrition*, **33**: 2151-2156.
- Toyosaki, T., Yamamoto, A. and Mineshita, T. (1986). Simultaneous analysis of riboflavin and its decomposition products in various milks by high-performance liquid chromatography. *Journal of Micronutrient Analysis*, **2**: 117-123.
- Trugo, N. M. F. and Sardinha, F. (1994). Cobalamin and cobalamin-binding capacity in human milk. *Nutrition Research*, **14**: 23-33.
- Ubbink, J. B. (2000). Vitamin B 6. **In**: Modern Chromatographic Analysis of Vitamins, pp. 435-478. De Leenheer, A. P., Lambert, W. E., Van Bocxlaer, J. F. Eds., Marcel Dekker Inc., New York, NY.
- Udipi, S. A., Kirksey, A. and Roepke, J. L. (1987). Diurnal variations in folacin levels of human milk: use of a single sample to represent folacin concentration in milk during a 24h period. *The American Journal of Clinical Nutrition*, **45**: 770-779.

- Udipi, S. A., Kirksey, A., West, K. and Giacoia, G. (1985). Vitamin B6, vitamin C and folacin levels in milk from mothers of term and preterm infants during the neonatal period. *American Journal of Clinical Nutrition*, **42**: 522-530.
- Uyeda, K. and Rabinowitz, J. C. (1963). Fluorescence properties of tetrahydrofolate and related compounds. *Analytical Biochemistry*, **6**: 100-108.
- Van Herwaarden, A. E., Wagenaar, E., Merino, G., Jonker, J. W., Rosing, H., Beijnen, J. H. and Schinkel, A. H. (2007). Multidrug transporter ABCG2/breast cancer resistance protein secretes riboflavin (vitamin B2) into milk. *Molecular and Cellular Biology*, **27**: 1247-1253.
- Van Zoeren-Grobbe, D., Schrijver, J., Van den Berg, H. and Berger, H. M. (1987). Human milk vitamin content after pasteurisation, storage, or tube feeding. *Archives of Disease in Childhood*, **62**: 161-165.
- Vanderslice, J. T., Brownlee, S. G., Maire, C. E., Reynolds, R. D. and Polansky, M. (1983). Forms of vitamin B6 in human milk. *American Journal of Clinical Nutrition*, **37**: 867-871.
- Venugopal, P. B. (1985). Choline. **In:** Methods of Vitamin Assay, pp. 555-578. Augustin, J., Klein, B. P., Becker, D. and Venugopal, P., Eds., John Wiley & Sons, New York, NY.
- Villalpando, S., Latulippe, M. E., Rosas, G., Irurita, M. J., Picciano, M. F. and O'Connor, D. L. (2003). Milk folate but not milk iron concentrations may be inadequate for some infants in a rural farming community in San Mateo, Capulhuac, Mexico. *The American Journal of Clinical Nutrition*, **78**: 782-789.

- Wardinsky, T. D., Montes, R. G., Friederich, R. L., Broadhurst, R. B., Sinnhuber, V. and Bartholomew, D. (1995). Vitamin B12 deficiency associated with low breast-milk vitamin B12 concentration in an infant following maternal gastric bypass surgery. *Archives of Pediatrics and Adolescent Medicine*, **149**: 1281-1284.
- Watts, B. M., Peng, D. H. and Esselbaugh, N. C. (1948). The Enzymatic Extraction of Riboflavin from Pork for the Fluorometric Determination. *The Journal of Biological Chemistry*, **172**: 707-712.
- West, K. D. and Kirksey, A. (1976). Influence of vitamin B6 intake on the content of the vitamin in human milk. *American Journal of Clinical Nutrition*, **29**: 961-969.
- White, J. (1962). The Structure of Vitamin B 12 III. The crystal structure analysis of the air-dried B 12 crystals. *Proceedings of the Royal Society of London A*, **266**: 440-474.
- WHO (2001). The optimal duration of exclusive breastfeeding. Report of an expert consultation. WHO, Geneva.
- Wielders, J. P. M. and Mink, C. (1983). Quantitative analysis of total thiamine in human blood, milk and cerebrospinal fluid by reversed-phase ion-pair high-performance liquid chromatography. *Journal of Chromatography B*, **277**: 145-156.
- Williams, R. J., Cheldelin, V. H. and Mitchell, H. K. (1942). The B vitamin content of milk from animals of different species. *The University of Texas Pub*, **4237**: 97-104.
- Williams, R. J., Lyman, C. M., Goodyear, G. H., Truesdail, J. H. and Holaday, D. (1933). Pantothenic acid, a growth determinant of universal biological occurrence. *Journal of the American Chemical Society*, **55**: 2912-2927.

- Wilson, R. G. and Davis, R. E. (1984). Vitamin B6 intake and plasma pyridoxal phosphate concentrations in the first 2 weeks of life. *Acta Paediatrica*, **73**: 218-224.
- Woodcock, E. A., Warthesen, J. J. and Labuza, T. P. (1982). Riboflavin photochemical degradation in pasta measured by high performance liquid chromatography. *Journal of Food Science*, **47**: 545-549.
- Woolley, D. W., Waisman, H. A. and Elvehjem, C. A. (1939). Studies on the structure of the chick antidermatitis factor. *Journal of Biological Chemistry*, **129**: 673-679.
- Wyse, B. W., Wittwer, C. and Hansen, R. G. (1979). Radioimmunoassay for pantothenic acid in blood and other tissues. *Clinical Chemistry*, **25**: 108-110.
- Wyse, B. W., Song, W. O., Walsh, J. H. and Hansen, R. G. (1985). Pantothenic acid. **In:** Methods of Vitamin Assay, pp. 399-416. Augustin, J., Klein, B. P., Becker, D. and Venugopal, P., Eds., John Wiley & Sons, New York, NY.
- Zafra-Gómez, A., Garballo, A., Morales, J. C. and García-Ayuso, L. E. (2006). Simultaneous determination of eight water-soluble vitamins in supplemented foods by liquid chromatography. *Journal of Agricultural and Food Chemistry*, **54**: 4531-4536.
- Zeisel, S. H. (2006). Choline: critical role during fetal development and dietary requirements in adults. *Annual Review of Nutrition*, **26**: 229-250.
- Zeisel, S. H. and Caudill, M. A. (2010). Choline. *Advances in Nutrition: An International Review Journal*, **1**: 46-48.
- Zeisel, S. H., Char, D. and Sheard, N. F. (1986). Choline, phosphatidylcholine and sphingomyelin in human and bovine milk and infant formulas. *The Journal of Nutrition*, **116**: 50-58.

- Zeisel, S. H. and DaCosta, K. A. (1990). Choline: determination using gas chromatography/mass spectrometry. *Journal of Nutritional Biochemistry*, **1**: 55-59.
- Zeisel, S. H., Holmes-McNary, M. (2001). Choline. **In:** Handbook of Vitamins, pp. 513-528. Rucker, R. B., Suttie, J. W., McCormick, D. B., Machlin, L. J., Eds., Marcel Dekker Inc., New York, NY.
- Zeisel, S. H., Stanbury, J. B., Wurtman, R. J., Brigida, M. and Fierro-Benitez, R. (1982). Choline content of mothers' milk in Ecuador and Boston. *The New England Journal of Medicine*, **306**: 175-176.
- Zielenkiewicz, W., Terekhova, I. V., Kobiala, M. and Kumeev, R. S. (2010). Thermodynamic study on inclusion complex formation of riboflavin with hydroxypropyl- $\beta$ -cyclodextrin in water. *Journal of Thermal Analysis and Calorimetry*, **101**: 595-600.

**Table 1:** Summary of the most useful, valid methods that can be applied today, in the opinion of these authors

Vitamin	Methodology	References
Thiamin	UPLC-MS/MS,	Hampel et al. 2012,
	HPLC-FLD	Stuetz et al. 2011
Riboflavin	UPLC-MS/MS,	Hampel et al. 2012,
	HPLC-FLD	Roughead and McCormick 1990
Niacin	UPLC-MS/MS	Hampel et al. 2012
Vitamin B6	UPLC-MS/MS,	Hampel et al. 2012,
	RPLC-FLD	Hamaker et al. 1985
Cobalamin	Competitive Protein Binding - Chemiluminescence	Lildballe et al. 2009
Folate	Microbiological Assay	Lim et al. 1988
Pantothenic Acid	Microbiological Assay	Coryell et al. 1945,
	Radioimmuno Assay	Song et al. 1984
Biotin	Competitive Protein Binding - Radiodetection	Mock and DuBois 1986
Choline	LC/ESI-IDMS	Koc et al. 2002
	GC-MS / HPLC- Radiodetection	Promfret et al. 1989

**Supplementary Table 2A:** Summary of thiamin concentrations in human milk as reported in the literature



**Supplementary Table 2B:** Summary of riboflavin concentrations human milk as reported in the literature

**Supplementary Table 2C:** Summary of niacin concentrations human milk as reported in the literature

**Supplementary Table 2D:** Summary of vitamin B6 concentrations human milk as reported in the literature

**Supplementary Table 2E:** Summary of cobalamin concentrations human milk as reported in the literature

**Supplementary Table 2F:** Summary of folate concentrations human milk as reported in the literature

**Supplementary Table 2G:** Summary of pantothenic acid concentrations human milk as reported in the literature

**Supplementary Table 2H:** Summary of biotin concentrations in human milk as reported in the literature

**Supplementary Table 2I:** Summary of choline concentrations in human milk as reported in the literature