

This article was downloaded by: [University Library Utrecht]

On: 06 August 2013, At: 15:38

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Critical Reviews in Food Science and Nutrition

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/bfsn20>

Rice Bran: A Novel Functional Ingredient

Mian Kamran Sharif^a, Masood Sadiq Butt^a, Faqir Muhammad Anjum^a & Saima Hafiz Khan^a

^a National Institute of Food Science and Technology, University of Agriculture, Faisalabad, 38040, Pakistan

Accepted author version posted online: 19 Feb 2013.

To cite this article: Critical Reviews in Food Science and Nutrition (2013): Rice Bran: A Novel Functional Ingredient, Critical Reviews in Food Science and Nutrition, DOI: 10.1080/10408398.2011.608586

To link to this article: <http://dx.doi.org/10.1080/10408398.2011.608586>

Disclaimer: This is a version of an unedited manuscript that has been accepted for publication. As a service to authors and researchers we are providing this version of the accepted manuscript (AM). Copyediting, typesetting, and review of the resulting proof will be undertaken on this manuscript before final publication of the Version of Record (VoR). During production and pre-press, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal relate to this version also.

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

Rice Bran: A Novel Functional Ingredient

MIAN KAMRAN SHARIF, MASOOD SADIQ BUTT, FAQIR MUHAMMAD ANJUM and
SAIMA HAFIZ KHAN

National Institute of Food Science and Technology, University of Agriculture, Faisalabad 38040,
Pakistan

ABSTRACT

Rice (Oryza sativa) is the most important staple food for a large part of the world's human population, especially in East and South Asia, the Middle East, Latin America, and the West Indies. It provides more than one fifth of the calories consumed worldwide by the human. It is the 2nd leading cereal crop and staple food of half of the world's population. It is grown in at least 114 countries with global production of 645 million tons; share of Asian farmers is about 90% of the total produce. Rice bran, brown outer layer of rice kernel, is mainly comprised of pericarp, aleuron, subaleurone layer and germ. It contains appreciable quantities of nutrients like protein, fat and dietary fiber. Furthermore, it contains substantial amount of minerals like K, Ca, Mg and Fe. Presence of antioxidants like tocopherols, tocotrienols and γ -oryzanol also brighten prospects of rice bran utilization for humans as functional ingredient to mitigate the life threatening disorders. Moreover, in the developing countries, budding dilemma of food crisis, arising due to lower crop yields and escalating population; needs to utilize each part of available resources. In order to provide enough food to all people, there is the holistic approach of using the by-products generated during food processing and preparations. Rice is being

processed in well established industry but the major apprehension is the utilization of its by-products; rice bran (5-8%) and polishing (2-3%) that are going as waste. Rice processing or milling produces several streams of materials including milled rice, bran and husk. In developing countries, rice bran is considered as a by-product of the milling process and commonly used in animal feed or discarded as a waste. The potential of producing rice bran at the global level is 29.3 million tons annually while the share of Pakistan is worked out to be 0.5 million tons. In present article attempt has been made to highlight the significance of this valuable but neglected ingredients under various heading.

Keywords rice bran, hypercholesterolemia, hyperglycemia, dietary fiber, rice bran protein isolates, rice bran oil, baked products

Address all correspondence to:

Dr. Mian Kamran Sharif, Assistant Professor, National Institute of Food Science and Technology, University of Agriculture, Faisalabad 38040, Pakistan.

Tel: +92-333-860-8341

Fax: +92-41-920-1105

E-mail: mks@uaf.edu.pk; mks_ft@yahoo.com

PHYSIOLOGY AND GENERAL CHARACTERISTICS OF RICE BRAN

Rough rice (paddy) is composed of a white starchy rice kernel tightly covered by a coating of bran, enclosed in a tough siliceous hull (Lakkakula et al., 2004). When husk is removed, bran layer comes in direct contact with air, resulting in the development of off-flavor in brown rice due to its endogenous lipase. Moreover, the appearance of brown rice is not appealing due to its color (Saunders, 1990). Hence further processing of rice is required to remove the bran from brown rice to produce white rice (Hu et al., 1996). It is consumed after appropriate polishing to give a desired degree of whiteness (Juliano, 1985). Rice bran constitutes about 10% of the weight of rough rice. It is comprised of pericarp, aleurone, sub-aleurone, seed coat, nucellus along with the germ and a small portion of endosperm (Salunkhe et al., 1992; Hargrove, 1994; Hu et al., 1996). The percentage and composition of rice bran vary according to the rice variety, pretreatment before milling, type of milling system and the degree of milling (Saunders, 1990). Rice bran is light in color, sweet in taste, moderately oily and has a slightly toasted nutty flavor (Hu et al., 1996). Texture varies from a fine, powder-like consistency to a flake, depending on the stabilization process (Barber and Benedito-de, 1980).

TAXONOMIC INFORMATION

Botanical Name	<i>Oryza sativa</i>
Kingdom	Plantae
Division	Angiosperms
Class	Monocots

Order	Poales
Family	Poaceae
Genus	Oryza
Species	<i>O. sativa</i>

Rice bran contains 12-22% oil, 11-17% protein, 6-14% fiber, 10-15% moisture and 8-17% ash. It is rich in vitamins including vitamin E, thiamin, niacin and minerals like aluminum, calcium, chlorine, iron, magnesium, manganese, phosphorus, potassium, sodium and zinc (Saunders, 1990; Hu et al., 1996; Xu, 1998). Stabilized rice bran of Pakistani rice cultivar “Basmati-385” contains 6.68% moisture, 7.89% ash, 15.78% crude protein, 20.55% crude fat, 7.59% crude fiber and 41.51% nitrogen free extract (Sharif et al., 2005). Rice bran also contains a significant amount of nutraceutical compounds and approximately 4% unsaponifiables, mainly comprised of naturally occurring antioxidant such as tocopherols, tocotrienols and oryzanol (Ju and Vali, 2005). Rice bran proteins are of high nutritional value (Kennedy and Burlingame, 2003) and hypoallergenic (Tsuji et al., 2001). These proteins are rich in essential amino acids, especially lysine, hence can be used as an ingredients in food recipes (Wang et al., 1999). Stabilized rice bran is also a good source of both soluble and insoluble dietary fiber ranging from 20-51% (Saunders, 1990), which is almost twice as much as that of oat bran. Rice bran can be used as a stool bulking agent (Tomlin and Read, 1988) and for the enrichment of some foods (Burton, 2000).

ANTI-NUTRITIONAL FACTORS IN RICE BRAN

The effective utilization of rice bran is possible only by deactivating the lipase enzyme, responsible for the hydrolytic degradation of bran constituents and denaturation of anti-nutritional factors. Successful developments in the use of various techniques to stabilize rice bran have resulted in the emergence of rice bran as an important by-product of the rice milling industry. The anti-nutritional factors in rice bran include lipases, trypsin inhibitor, haemagglutinin-lectin and phytates. Lipases are enzymes that are primarily responsible for the hydrolysis of triglycerides into glycerol and fatty acids. Rice bran contains several types of lipases which results in significant increase of the free fatty acids (FFA) by hydrolyzing the oil. Rapid increase in the free fatty acid occurs within hours and reaches 7-8% within 24 hours, followed by about 5% increase per day (Ramezanzadeh et al., 1999; Rukmani, 2002). Lipase activity is greatly affected by moisture, temperature, pH, time and water activity (Dunford and King, 2001; Gangodavilage, 2002). The enzyme was active upto 40°C and the activity declined sharply to 65% at 60°C and then gradually decreased (Bhardwaj et al., 2001). In addition to native lipases, the microbial lipases also deteriorate the nutritional quality of the oil, making it unfit for human consumption. The hydrolytic rancidity severely affects the nutritive value and palatability of rice bran (Rajeshwara and Prakash, 1995). Trypsin inhibitors are also endogenous enzymes, which can form stable complex with proteolytic pancreatic enzymes i.e. trypsin and chemotrypsin. Due to complex formation, the activity of these enzymes decreases. Rice bran contains trypsin inhibitor (Kratzer and Payne, 1977; Deolankar and Singh, 1979). Approximately 85-95% trypsin inhibitor activity was found in rice embryo. One mole of rice bran trypsin inhibitor can inhibit two moles of trypsin.

Haemagglutinin-lectins are toxic globulin proteins present in the rice bran and agglutinate mammalian red blood cells (Ory et al., 1981). Similarly, lectin is a glycoprotein and is present in germ portion. It comprised of 27% carbohydrate, predominantly glucose (Takahashi et al., 1973) while another 10% carbohydrate is mainly in the form of xylose and arabinose (Indravathamma and Seshadri, 1980). The lectin also contains a large number of glycine and cystine residues (Tsuda, 1979). Phytates (1,2,3,4,5,6-hexaphosphate of myoinositol) occur in discrete regions of cereal grains and accounts for 85% of the total phosphorous content of grains. They reduce the bio-availability and digestibility of nutrients by forming complexes with minerals, protein, digestive enzymes and amino acids mainly lysine, methionine, arginine and histidine (Jangbloed et al., 1991; Bird, 1998). It is a rich source of minerals particularly phosphorous, zinc and ferrous (Farrell, 1994). Phytic acid showed strong chelating properties due to its structure (Ramzan, 2000). Phytates also affect the solubility, functionality and digestibility of proteins and carbohydrates.

PROCESSING OF RICE BRAN

The processing of rice bran was carried out to inactivate lipases as well as other nutritional inhibitors in such a way that their toxicity is ruined without damaging the protein quality of rice bran. Furthermore, it also destroys the field fungi, bacteria and insects infestation; ultimately the bran becomes safe from further deterioration which alternately enhanced its shelf life. The greatest restriction to the use of rice bran as a food ingredient is its instability during storage. Upon milling, the oil is exposed to lipases, causing rapid breakdown to free fatty acids @ 5–7% of the weight of oil per day. Hence due to the naturally occurring enzymatic activity and subsequent hydrolytic rancidity, it is necessary to stabilize the rice bran by suitable

techniques for controlling undesirable reactions. Bran, after proper stabilization, can serve as a good source of protein, essential fatty acids, calories, and nutrients such as tocopherols and ferulic acid derivatives. The commonly used stabilization techniques are thermal and chemical treatments (Randall et al., 1985; Kim et al., 1987). There are different types of heat stabilization procedures such as retained moisture heating (Lin and Carter, 1973), added moisture heating (Saunders, 1986), extrusion cooking (Sayre et al., 1982), microwave heating (Malekian et al., 2000) and Ohmic or electrical heating (Lakkakula et al., 2004).

Heat stabilization is accomplished commercially by wet or dry heating methods i.e. hot air, drum drying, dry extrusion and microwave (Prabhakar, 1987; Narisullah and Krishnamurthy, 1989). Although hot air drying is an effective method of stabilization, the non-uniform heating of material in the tray driers limits its application. Rice bran was stabilized by fluidized bed drying at 90-130°C (Fernando and Hewavitharana, 1993). Although fluidization provides uniform heating of bran; however, high air velocities are required for the process; making it uneconomical (Narisullah and Krishnamurthy, 1989). The stabilized rice bran was obtained by drum drying at 156°C (Delahaye et al., 2005). Parboiling also results in stabilization of rice bran by destroying lipase activity (Narisullah and Krishnamurthy, 1989). An edible acid (0.1-2.0% acetic acid) having anti-oxidative properties was added to parboiled rice bran to maintain the stability of the bran for at least 6 months at ambient conditions (Tao, 2001). The common drawbacks in heat treatment methods are: severe processing conditions capable of damaging valuable components, substantial moisture removal and inability to achieve irreversible inactivation of enzyme. To cope with these problems, moist heat treatment is suggested. Extrusion cooking has been found to produce stable rice bran by holding at 125-130 °C for few

seconds, then at 97-99 °C for 3 minutes prior to cooling (Randall et al., 1985). Heating in the presence of moisture is more effective for permanently denaturing lipases (Ramezanzadeh et al., 1999). Long-term storage studies with extrusion cooking indicate stability against FFA development upto 4 months (Carroll, 1990; Randall et al., 1985), in contrast to dry heat methods. Hence steaming is suitable method of bran pretreatment with respect to decrease in FFA development and the oil extractability in small-scale (Amarasinghe and Gangodavilage, 2004). However, less flexibility and higher initial and operating costs make the process uneconomical. Furthermore, moist heat results in agglomeration of bran, resulting in lumpy bran.

To achieve proper stabilization, every discrete bran particle must have uniform moisture content, depending upon time and temperature. In recent years, use of microwave energy as an inexpensive source of heat for thermal processing of foods has offered an alternative energy source for stabilization of rice bran. It is considered to be one of the most energy-efficient types and a rapid method for heating food items (Yoshida et al., 2003). Considering other heat treatments, microwave heating is efficient, economical, shorter in processing time, minor effects on the nutritional value and has a little or even no effect on the natural color of bran. Microwave heating is an effective method for stabilizing rice bran with the addition of moisture, which enables heating over 100°C to occur (Malekian et al., 2000). The water molecules in the rice bran are excited to spin by the electromagnetic waves resulting enhanced kinetic energy along with the friction. Since water molecules play an important role in this process, the initial moisture content is a critical factor in the microwave stabilization. Rice bran was stabilized by heating in a microwave for 4 minutes until the internal temperature reached 110-115°C to denature the enzymes (Zhu, 2000).

DIETARY FIBER OF RICE BRAN

Dietary fiber is the edible parts of plants or analogous carbohydrates that are resistant to digestion and absorption in the human small intestine with partial fermentation in the large intestine (CAC, 1998). These are plant food materials that are not hydrolyzed by enzymes secreted by the human digestive tract but may be digested by micro flora in the gut. These plant food materials include non-starch polysaccharides such as celluloses, some hemicelluloses, gums and pectins as well as resistant starches (DeVries, 2001). The components of dietary fiber include cellulose, hemicellulose, pectins, hydrocolloids and lignin. These can be classified into two major categories depending on their solubility in water. In humans, the structural or matrix fibers (lignins, cellulose, and some hemicelluloses) are insoluble, whereas the natural gel-forming fibers (pectins, gums, mucilages, and the remainder of the hemicelluloses) are soluble. Soluble fiber acts like a gel and insoluble fiber adds bulk or softens stool. Soluble fiber forms a gelatin like substance in the intestine and increases the water content in stool. It has been shown that soluble fiber decreases blood cholesterol and sugar after meals in diabetics (Yeager, 1998). Similarly, insoluble fiber is effective in increasing feeling of fullness, stool size, bulk and helps to reduce constipation and hemorrhoids. Good sources of soluble fibers include fruits, vegetables, legumes, psyllium seeds and oat bran whereas whole grains are good sources of insoluble fiber (Matz, 1991).

Fiber supplementation has been used to enhance the fiber content of array of foods. Traditionally, fiber supplementation has focused on the use of milling by-products of cereal grains like wheat, corn, sorghum and other grains (McKee and Latner, 2000). Nowadays, fiber

supplementation has focused in cookies, crackers, snack foods, beverages, spices, imitation cheeses, sauces, frozen foods, canned meats, meat analogues and many other cereal-based products (Hesser, 1994). The WHO recommendation for total dietary fiber intake is above 25 g/day (WHO, 2003). The total dietary fiber content in stabilized rice bran ranges from 25 to 40% depending on the product (Carroll, 1990). Rice bran's fiber comprised of a relatively low proportion of soluble fiber (7–13%) and the rest is insoluble fiber (Anderson et al., 1990). However, rice bran has high percentage of oil (12–23%) as compared to other bran sources, with 4.2% unsaponifiable matter (Sugano and Tsuji, 1997). Rice bran oil, possibly because of unsaponifiable fraction or its fatty acid content, lowers cholesterol levels in hamsters, rats, humans and non-human primates (Sharma and Rukmini, 1986; Seetharamaiah and Chandrasekhara, 1989; Nicolosi et al., 1991; Kahlon et al., 1992; Purushothama et al., 1995).

RICE BRAN OIL

Rice bran contains 15-22% oil by weight (Orthoefer, 1996; Patel and Walker, 2004). Crude rice bran oil contains 90-96% of saponifiable and about 4% unsaponifiable lipids. The saponifiable lipids include 68-71% triglycerides, 2-3% diglycerides, 5-6% monoglycerides, 2-3% free fatty acids, 2-3% waxes, 5-7% glycolipids and 3-4% phospholipids (McCaskill and Zhang, 1999) whereas the principal component of the unsaponifiable fraction is γ -oryzanol (Raghuram and Rukmini, 1995). Rice bran oil has excellent fatty acid profile. It has oleic acid (38.4 %), linoleic acid (34.4%) and linolenic acid (2.2%) as unsaturated fatty acids while palmitic acid (21.5%) and stearic acid (2.9%) as saturated fatty acids (Rukmini and Raghuram, 1991). The saturated, monounsaturated and polyunsaturated fatty acids are in the ratio of approximately 1: 2.2: 1.5 (Shin and Chung, 1998; Krishna, 2002). Three major fatty acids, palmitic, oleic and

linoleic make up 90% of the total fatty acids of the rice bran oil (Amarasinghe and Gangodavilage, 2004).

Rice bran oil (RBO) is traditionally consumed in Asian rice producing countries with growing interest in Western markets (Jariwalla, 2001; Kim and Godber, 2001; Nasirullah, 2001; Pszczola, 2001). It is in steady demand as “healthy oil” in Japan where approximately 80 thousand tons is consumed annually (Sugano and Tsuji, 1997). Traditionally, rice bran oil has been used for frying food, due to its oxidative stability and flavor; it is now considered as a good substitute for vegetable oils (Sayre and Saunders, 1985; Goenka, 1987). It is widely used in pharmaceutical, food and allied industries due to its unique properties, high medicinal value and therapeutical applications (Cicero and Gaddi, 2001; Amarasinghe and Gangodavilage, 2004). RBO is an unconventional vegetable oil believed to be healthy in some populations (Sugano et al., 1999) due to higher levels of antioxidants (Patel and Walker, 2004) and phytosterols (Stoggl et al., 2005). It is superior to other vegetable oils because it contains ω -3 and ω -6 fatty acids; particularly due to oryzanol and higher amounts of unsaponifiables (Krishna et al., 2005). Currently, efforts are being made to develop RBO with retained non-saponifiable components, while minimizing levels of problematic free fatty acids (Ginsberg et al., 1998).

RICE BRAN PROTEIN ISOLATES

Protein isolates can be prepared from a number of cereal by-products including rice, maize, barley, wheat and rye (Lasztity *et al.*, 1995). Functional properties of the protein play significant role in textural attributes of the final product (Neto et al., 2001). Optimum results can be obtained when both the protein and food's functional properties are well synchronized. Use of quality protein isolates furnishes food with rich amino acid profile. Plant proteins, being less

expensive than animal source, are used to supplement foods with desirable functional attributes (Wiseman and Price, 1987). Textural and functional properties of protein isolates may vary with the processing conditions used for extraction and product manufacturing (Gnanasambandam and Hetiarachchy, 1995; Wagner et al., 1996), nonetheless, thorough study of protein functionality is necessary for food applications.

Functional properties of rice bran protein concentrates are suitable for food application (Jiamyangyuen et al., 2005a). Rice bran protein isolates have promising emulsification properties to be used as natural emulsifier in food products (Lee et al., 2004). Emulsion properties and solubility of bran protein hydrolysates make them suitable for application in array of food products like coffee whiteners, toppings, beverages, confectionary, meat and bakery products (Hamada, 2000). Protein extracts from rice bran has been incorporated in liquid foods like milk and other drinks (Watchararuji et al., 2008). Protein content of baked products was increased by incorporating rice bran protein concentrates without compromising their final quality (Jiamyangyuen *et al.*, 2005b). Balanced amino acid profile and hypoallergenic nature of bran protein are suggestive for its application in infant foods (Wang et al., 1999). Rice protein concentrates in combination with polysaccharide pullulan are converted into edible films with good tensile strength (Shih, 1996). Addition of propylene glycol alginate (PGA) and oil provide resistance against water vapor creating water barrier properties in the films. Cross linking of amino acid and PGA improve films strength through covalent bonding under high pH conditions (Shih, 1994).

HYPOCHOLESTEROLEMIC EFFECTS OF RICE BRAN

Scientific studies support recommendations to increase dietary fiber as part of hyperlipidemia treatment. The hypocholesterolemic effects of rice bran have been demonstrated in experimental animals (Sharma and Rukmini 1987; Seetharamaiah and Chandrashekhara 1989; Kahlon et al., 1992) and humans (Hundemer et al., 1991; Sanders and Reddy, 1992; Hakala et al., 2002). The cholesterol lowering effects of rice bran (fullfat), soybean fiber, oat and barley bran were compared in mice adding 0.06% cholesterol in their diets. Both rice bran and soybean fiber diets had significantly lower total blood cholesterol compared with placebo. Rice bran was found to be the most effective supplement in reducing liver and plasma total cholesterol compared to the control diet. Moreover, mice consuming rice bran diet, demonstrated higher HDL to total cholesterol ratios (Hundemer et al., 1991). In another study, rats fed on diets containing rice and wheat bran showed significant reduction in liver cholesterol and triglycerides. The rice bran diet also increased LDL receptor activity in the liver more than that of wheat bran, hence, effectively lowering plasma cholesterol levels (Topping et al., 1990).

The cholesterol lowering effects of fullfat and defatted stabilized rice bran, parboiled rice bran and rice bran in combination with wheat bran were studied in hamsters fed on fiber diets with 0.5% added cholesterol. The liver cholesterol concentrations, in particular, were significantly lower in animals consuming fullfat stabilized rice bran than all other groups (Kahlon et al., 1990). Rice bran with extremely low β -glucan content is known to be as effective as high- β -glucan oat and barley brans in lowering serum cholesterol (Seetharamaiah and Chandrasekhara, 1989; Kahlon et al., 1990; 1992). The hypocholesterolemic effects of rice bran may be attributed to the unsaponifiable fraction of rice bran oil, primarily phytosterols, tocopherols (tocopherols and tocotrienols), γ -oryzanol, triterpene alcohol and other minor compounds

(Sharma and Rukmini, 1987; Yoshino et al., 1989; Nicolosi et al., 1991). In a similar study, benefits of bran addition from rice, oats, corn and wheat in diets fed to hamsters were evaluated at relatively high cholesterol level (0.3%). Liver cholesterol concentrations and liver weights were significantly lower for the rice bran diet than for either the corn or wheat bran diets. Animals fed on rice bran had significantly lower VLDL levels and the highest HDL to total cholesterol ratios when compared to all other bran due to greater lipid and sterol excretion (Kahlon et al., 1998).

Chicks were fed on 60% fullfat rice bran and corn/soy diets with 0.5% added cholesterol. Significant differences were found in total cholesterol, triglycerides, high-density and low-density lipoprotein cholesterol. Likewise, in a second study, chicks were fed on fullfat rice bran, defatted rice bran and corn/soy diets balanced for 18% protein, 14.47% total dietary fiber and 10.78% lipid with 0.5% added cholesterol. Total cholesterol and triglycerides were significantly lower in chicks fed on fullfat rice bran diets. Significant differences were found in HDL values for all diets with fullfat rice bran exhibiting the highest (155 mg/dL) and corn/soy exhibiting the lowest mean value (114 mg/dL). Fullfat rice bran appeared to increase HDL and lower LDL in chicks, but did not always affect (TC) total cholesterol (Newman et al., 1992). In an efficacy study, it was found that rice bran might lower cholesterol by increasing short chain fatty acid production in the cecum by hindering cholesterol absorption due to a change in intestinal fluid viscosity or by directly inhibiting cholesterol synthesis in the liver (Fukushima et al., 1999). Rice bran supplementation has been found effective in lowering total cholesterol and LDL levels in human subjects with moderate hypercholesterolemia (Hundemer et al., 1991). However, serum cholesterol was found to be decreased in patients with mild hypercholesterolemia who consumed

300g/d unpolished rice or 100g/d stabilized rice bran (Hakala et al., 2002). Fullfat rice bran was found to be more effective in lowering cholesterol than isolated rice bran fractions or their combinations (Kahlon et al., 1992).

Rice bran has been found to be equivalent to oat bran in lowering cholesterol. Mildly hypercholesterolemic subjects were fed on treatment diets (100g/day stabilized rice bran or oat bran) with 300mg/day added cholesterol. Total cholesterol levels were significantly reduced in both bran diets when compared to the control (Hegsted et al., 1993). Similarly, changes in plasma lipid levels were studied in men with slightly above the normal cholesterol levels providing test diets containing 35g/day of wheat bran, 60g/day of rice bran or 95g/day of oat bran. The varying amounts of the different brans provided a constant amount of total dietary fiber i.e. 11.8g/day. At baseline level, only oat bran was effective in reducing plasma cholesterol as compared to other treatments. However, the highest rise in HDL was associated with the rice bran diet, resulting in an improved HDL to total cholesterol ratio. Moreover, plasma triglycerides were also lower in case of rice bran compared to wheat bran diet (Kestin et al., 1990).

CHOLESTEROL-LOWERING MECHANISMS

The mechanism of action of rice bran and its oil on lipid metabolism is not yet evident. However, the most probable hypothesis of RBO hypolipidemic action is its specific content of phytosterols, polyphenols (γ -oryzanol) and tocopherols (tocopherols and tocotrienols). The cholesterol-lowering effects of RBO are possibly attributable to its relatively high unsaponifiables; physiologically bioactive in controlling cholesterol levels in subjects. These compounds have been found to work synergistically to exhibit hypocholesterolemic effects. The brief description of each is given below:

Phytosterols (campesterol, β -sitosterol and stigmasterol)

The phytosterols present in crude rice bran oil like campesterol, β -sitosterol and stigmasterol, have been proven effective in lowering plasma total and LDL-cholesterol without affecting HDL-cholesterol due to similarities in structures of plant sterols and cholesterol (Weststrate and Meijer, 1998). There are several mechanisms through which plant sterols affect cholesterol concentration in the body like formation of non-absorbable complex with cholesterol, altering the size and/or stability of the micelles, interferences with cholesterol esterification in the mucosal cell and interacting with protein receptors required in cholesterol absorption (Rong et al., 1997). It is generally assumed that plant sterols inhibit intestinal absorption of dietary and biliary cholesterol, because of the structural similarities with cholesterol. Some studies indicated that plant sterols contributed more hypocholesterolemic effects than unsaponifiables. In addition, some plant sterols may be more active than others (Wilson et al., 2000). Among the sterols, β -sitosterol has been recognized the predominant cholesterol-lowering component (Visser et al., 2000; Trautwein et al., 2002).

Polyphenols (oryzanol)

There are numerous mechanisms by which oryzanol lowers cholesterol levels such as: cholesterol-esterase inhibition by cycloartenol or by the inhibition of the accumulation of cholesterol-esters within macrophages or by the modulation of cholesterol acid esterase and acyl-CoA-cholesterol-acyltransferase (Rukmini and Raghuram, 1991); sterol moiety of γ -oryzanol is partly split off from the ferulic acid part in the small intestine by cholesterol esterase (Sugano and Tsuji, 1997); effect on biliary secretion resulting in increased faecal excretion of cholesterol and bile acids (Seetharamaiah et al., 1990); direct inhibition of lipid metabolism (Sakamoto et

al., 1987); increased fecal excretion of cholesterol and its metabolites (Wilson et al., 2007) and oryzanol exercises its effects on cholesterol metabolism at sites other than the intestine.

Tocols (tocopherol and tocotrienol)

In case of tocols, cholesterol lowering mechanisms include: antioxidant activity that inhibits cholesterol oxidation (Xu et al., 2001); inhibit HMG-CoA-R, a key enzyme in the endogenous synthesis of cholesterol, via increasing the controlled degradation of reductase protein and decreasing the efficiency of the translation of HMG-CoA-R messenger RNA (Parker et al., 1993; Khor et al., 1995); inhibit the activity of 3-hydroxy-3-methylglutaryl-coenzyme A (HMG-CoA) reductase, the liver enzyme that is critical to the rate at which cholesterol is synthesized (Khor and Ng, 2000); inhibit cholesterol synthesis by suppressing HMG-CoA reductase activity through a posttranscriptional mechanism in HepG2 cells (Pearce et al., 1992; Parker et al., 1993; Qureshi et al., 2000) and via decrease in serum total and LDL cholesterol by inhibiting the hepatic enzymic activity of β -hydroxy- β -methylglutaryl coenzyme A (Qureshi et al., 2002).

SUPPLEMENTATION IN BAKED PRODUCTS

Nowadays, people are becoming more conscious about their health and nutrition. Foods that are convenient with good taste, reasonably priced and carry a favorable nutritional image are in great demand like bakery products especially cakes and cookies. Wheat flour is the primary raw material which provides a matrix in which other ingredients are mixed to form dough or batter. In Pakistan, the predominant use of rice bran is an ingredient in livestock feed. Considering its importance, value-added processing technologies for rice bran have been sought by researchers. The products supplemented with rice bran can play an important role in the

existing food crises besides its associated health claims. From a marketing view point, the most abundant available rice bran derived product is the oil (Orthoefer, 1996). Rice bran oil has an impressive nutritional quality, which makes it suitable for nutraceutical products. It has industrial potential particularly in snack food preparation because of great frying stability and with a good fry life and nutty flavor (Sarkar, 1992). Production of margarine from rice bran oil has health benefits with reduced saturated fats and trans-fatty acids.

Rice bran fractions can be used to produce acceptable low fat, high fiber bakery products (Kennedy et al., 1996). *Torticas de Moron*, a traditional Cuban bakery product, was manufactured with 0, 20, 25 or 30% of the usual white flour replaced by parboiled rice bran. Protein, fat, crude fiber and ash were higher in the rice bran *Torticas* than in the control. It was further observed that 25% replacement of flour with rice bran resulted in a product with acceptable sensory properties, chemical composition and shelf life (Zumbado et al., 1997). Rice bran fiber has been reported to contain high amounts of functional proteins and fats along with antioxidants, vitamins and trace minerals in addition to being a concentrated source of fiber. The presence of these nutrients allows rice bran fiber to be used as both nutritional and functional ingredient. Chicken coated with stabilized rice bran fiber tend to absorb less fat during frying while the small amount of fat found naturally in rice bran fiber can act as a carrier for flavors (Hammond, 1994). Later, stabilized rice bran was incorporated @ 5, 10 and 20% in chutney powder. Addition of the rice bran had more affect on color and the least on texture with intermediate effects on aroma, taste and overall acceptability (Prakash and Jyothilakshmi, 1995).

Leavened Pan Bread

Bread is considered as one of the prime bakery products. Its large loaf volume and fine texture require formation of well developed, elastic dough structure. An essential element in this process is gluten present in the flour (Eneche, 1999). Rice bran is used by the food industry in the production of baked goods, snacks, crackers, breads, and cereals (Rukmani, 2002). The functional properties of fullfat and defatted rice bran were explored by blending rice bran in wheat flour @ 5, 10 and 15% to prepare leavened pan bread. Addition of any of the defatted and fullfat rice brans was associated with reduction in loaf volume and a decrease in overall acceptability of the bread. Breads containing upto 10% of either type of rice bran were still considered acceptable (Sekhon et al., 1997). In another study, stabilized rice bran was successfully incorporated upto 20% for the production of yeast bread. The hygroscopicity of the rice bran improved moisture retention in the baked products while foaming ability improved air incorporation and leavening (Carroll, 1990). In a similar study, bread volume decreased with blending of different types of rice bran; however, the decrease was more pronounced with the defatted bran. Stabilized fullfat rice bran upto 20% level and un-stabilized fullfat or stabilized defatted rice bran upto 10% was found suitable in various food products (Singh et al., 1995).

In commercial production of leavened pan bread, wheat flour was supplemented with rice bran from 15 to 30% and it was observed that it can be supplemented successfully upto 15% replacement level without affecting loaf weight, height or volume (Sharp and Kitchen, 1990). Likewise, leavened pan bread was made by supplementing 10 and 20% processed fullfat and defatted rice bran to study the functional behavior of bread compared with control. Texture profile analysis showed no significant differences as far as cohesive and springiness, but bread hardness, gumminess and chewiness increased with increasing levels of rice bran and were more

prominent in bread from defatted rice bran. Measurements of texture exhibited no detrimental effect of adding fullfat rice bran upto 10% and slight hardening of loaves with 20% level compared to control (Lima et al., 2002). Defatted rice bran (DRB) has great potential to be utilized for various bakery products without affecting the nutritional composition and sensory attributes. Overall quality and nutritional profile of pan bread was improved by DRB addition in various proportions. However, fiber and mineral enriched pan bread, with excellent quality, can be prepared by replacing the wheat flour with defatted rice bran upto 5% level without affecting internal and external characteristics of bread (Ajmal et al., 2006). The addition of rice bran to wheat flour increased proteins, lysine and dietary fiber in bread and cookies proportionately to the level of supplementation. In addition to color, flavor, protein extractability, solubility of bran and other properties like water & fat absorption, emulsifying & foaming capacity also showed improvement which further enlightens the potential use of bran in foods (Sharma and Chauhan, 2002). Defatted rice bran increases dough yield, contributes to an attractive tan crust and crumb, does not disturb fermentation or mixing tolerance of dough, causes baked products to remain fresher and more moist (Lynn, 1969). The flour strength and gluten quality decreased at different levels (5, 10, 15, 20 and 25%) of rice bran. The replacement of bran with decreasing amount of gluten has inverse correlation with flour strength. It was concluded that bran supplementation should not exceed to 15%, in flours containing low gluten content (Chumachenko et al., 1987).

Cookies

It is chemically leavened product, also known as “biscuit”. Generally the term biscuit is used in the European countries and cookies in the USA. Biscuits and biscuit like products have been made and eaten by man for centuries (Hoseney, 1986). Cookies are ideal for nutrient

availability, palatability, compactness and convenience. They differ from other baked products like bread and cakes because of having low moisture content, ensure comparatively free from microbial spoilage and confer a long shelf life of the product (Wade, 1988). Cookies are considered better for supplemented/composite flours due to their ready-to-eat form, wide consumption, relatively long shelf-life and good eating quality (Tsen et al., 1973). Cookies with high sensoric attributes have been produced from blends of millet/pigeon pea flour (Eneche, 1999), raw rice and wheat (Singh *et al.*, 1989), blackgram and wheat (Singh et al., 1993), chickpea and wheat (Singh et al., 1991), wheat, fonio and cowpea (McWatters et al., 2003) and soybean, chickpea or lupine with wheat (Hegazy and Faheid, 1990). Similarly, cookies with high sensory ratings have been produced from blends of wheat flour and rice bran.

Nutritional and functional properties of rice bran are well suited for baked products like cookies, muffins, breads, crackers, pastries and pancakes (Barber et al., 1981). The fullfat and defatted rice brans were blended in wheat flour @ 5, 10 and 15% to prepare cookies. There was improvement in spread of cookies with the addition of fullfat rice bran. In contrast, there was decrease in the spread of cookies after supplementation of defatted rice bran. Cookies supplemented with either type of rice bran were acceptable upto 10% supplementation level (Sekhon et al., 1997). In another study, cookies were successfully prepared from stabilized rice bran at levels of 20% (Carroll, 1990). In a similar study, stabilized fullfat rice bran upto 20% level and un-stabilized fullfat or stabilized defatted rice bran upto 10% was found suitable in various food products (Singh et al., 1995). Dry heat and extrusion stabilized rice bran was supplemented in wheat flour at 5 to 20% levels for the preparation of cookies (Sharma and Chauhan, 2002). Microwave stabilized defatted rice bran was supplemented in commercial

straight grade wheat flout @ 10, 20, 30, 40 and 50% supplementation level to prepare fiber and mineral enriched cookies. It was observed that defatted rice bran can be substituted from 10 to 20% in wheat flour to prepare rice bran supplemented cookies without adversely affecting quality attributes (Sharif et al., 2009).

CONCLUSION

Rice bran, as a co-product of the rice milling industry, is yet not to be efficiently utilized for human consumption. It is composed of pericarp, aleurone and subaleurone layers, parts of the germ and small portion of the starchy endosperm. It is rich in vitamins, minerals, amino acids, dietary fiber, essential fatty acids and plant sterols like γ -oryzanol, tocopherol and tocotrienol having promising health-related benefits. Due to its over all composition, nutritional profile, functional characteristics, and apparent hypoallergenicity, rice bran has many applications in a diet which is characterized by high in dietary fiber and low in saturated fat. It may be particularly beneficial to those individuals who show allergenicity to other cereal grains. Strong evidences are available that the consumption of rice bran may be beneficial in reducing the risk of cardiovascular diseases and colon cancer. There is dire need to consider rice bran as food rather feed ingredient to exploit full potential of this novel functional ingredient.

REFERENCES

Ajmal, M., Butt, M. S., Sharif, K., Nasir, M., and Nadeem, T. (2006). Preparation of fiber and mineral enriched pan bread by using defatted rice bran. *Int. J. Food Pro.* **9**(4): 623-636.

- Amarasinghe, B. M. W. P. K., and Gangodavilage, N. C. (2004). Rice bran oil extraction in Sri Lanka: Data for process equipment design. *Food Bio-prod. Process.* **82**(C1): 54-59.
- Anderson, J. W., Deakins, D. A., Floore, T. L. Smith B. M., and Whitis, S. E. (1990). Dietary fiber and coronary heart disease. *Crit. Rev. Food Sci. Nutr.* **29**: 95–147.
- Barber, S., and Benedito-de, B. C. (1980). Rice bran: chemistry and technology. In: Rice: Production and Utilization. Luh, D. S., Ed. AVI Publishing Co. Westport, C.T., USA.
- Barber, S., Benedito-de B. C., and Martenz, J. (1981). Rice bran proteins: Potential value of rice bran fractions as protein food ingredients. *Rev. Agroquim. Tecnol. Aliment.* **21**: 247-256.
- Bhardwaj, K., Raju, A., and Rajasekharan, R. (2001). Identification, purification, and characterization of a thermally stable lipase from rice bran: A new member of the (*Phospho*) lipase family. *Plant. Physiol.* **127**: 1728-1738.
- Bird, J. N. (1998). Phytase for Chinese poultry feeds. *Poult. Int.* **9**: 68-71.
- Burton, B. (2000). Dietary composition and obesity: do we need to look beyond dietary fat? *J. Nutri.* **130**: 272–274.
- CAC (Codex Alimentarius Commission). (1998). Codex Alimentarius: Food Labeling Complete Texts. FAO/WHO, Rome.
- Carroll, L. E. (1990). Functional properties and applications of stabilized rice bran in bakery products. *Food Technol.* **44**: 74-76.

- Chumachenko, N. A., Demehuk, A., Zagorodynaya, L., Zubko, L., and Shretsova, A. (1987). Technological indicators of quality of flour mixtures containing added bran. *Pishchevaye Promyshelmost.* **2**: 35-37.
- Cicero, A. F., and Gaddi, A. (2001). Rice bran oil and γ -oryzanol in the treatment of hyperlipoproteinaemias and other conditions. *Phytother. Res.* **15**: 277–289.
- Delahaye, E., Jimene, P., and Perez, E. (2005). Effect of enrichment with high content dietary fiber stabilized rice bran flour on chemical and functional properties of storage frozen pizzas. *J. Food Eng.* **68**: 1–7.
- Deolankar, R. P., and Singh, K. S. (1979). Trypsin inhibitor, mineral availability and performance of broiler chickens fed on diets based on rice bran. *Anim. Feed Sci. Technol.* **4**: 133-141.
- DeVries, J. W. (2001). AACC report: the definition of dietary fiber. *Cereal Foods World.* **46**: 112-126.
- Dunford, N. T., and King, J. W. (2001). Thermal gradient deacidification of crude rice bran oil utilizing supercritical carbon dioxide. *J. Am. Oil Chem. Soc.* **78**: 121-125.
- Eneche, E. H. (1999). Biscuit-making potential of millet/pigeon pea flour blends. *Plant Food. Hum. Nutri.* **54**: 21–27.
- Farrell, D. J. (1994). Utilization of rice bran in diets for domestic fowl and ducklings. *World Poult. Sci. J.* **5**: 116-131.

- Fernando, W. J. N., and Hewavitharana, L. G. (1993). Effect of fluidized bed drying on stabilization of rice bran. *Drying Technol.* **11**: 1115-1125.
- Fukushima, M., Fujii, S., Yoshimura, Y., Endo, T., and Nakano, M. (1999). Effect of rice bran oil on intrainestinal fermentation and cholesterol metabolism in cecectomized rats. *Nutri. Res.* **19**: 235-245.
- Gangodavilage, N. C. (2002). Optimal Operating Conditions and Equilibrium Characteristics of Rice Bran Oil Extraction. M.Sc. Thesis, Uni. Moratuwa, Sri Lanka.
- Ginsberg, H. N., Kris-Etherton, P., Dennis, B., Elmer, P. J., Ershow, A., Lefevre, M., Pearson, T., Roheim, P., Ramakrishnan, R., Reed, R., Stewart, K., Stewart, P., Phillips, K., and Anderson, N. (1998). Effects of reducing dietary saturated fatty acids on plasma lipids and lipoproteins in healthy subjects: the DELTA Study, protocol 1. *Arterioscler Thromb. Vasc. Biol.* **18**: 441–449.
- Gnanasambandam, R., and Hetiarachchy, N.S. (1995). Protein concentrates from unstabilized and stabilized rice bran: preparation and properties. *J. Food Sci.* **60**(5): 1066-1074.
- Goenka, O. P. (1987). Nutritional significance of RBO. In: Handbook on Rice Bran -Processing and Utilization of Products. Solvent Extractors Association of India, India.
- Hakala, P., Lampi, A. M., Ollilainen, V., Werner, U., Markova, M., Wahala, K., Karkola, S., and Piironen, V. (2002). Steryl phenolic acid esters in cereals and their milling fractions. *J. Agric. Food Chem.* **50**: 5300-5307.

- Hamada, J.S. (2000). Characterization and functional properties of rice bran proteins modified by commercial exoproteases and endoproteases. *J. Food Sci.* **65**(2): 305-310.
- Hammond, N. (1994). Functional and nutritional characteristics of rice bran extracts. *Cereal Foods World.* **39**: 752-754.
- Hargrove, K. L. (1994). Processing and utilization of rice bran in the United States. In: Rice Science and Technology. Marshall, W. E. and Wadsworth, J. I. (eds.). Marcel Dekker Inc., New York.
- Hegazy, N.A., and Faheid, S. M. N. (1990). Rheological and sensory characteristics of doughs and cookies based on wheat, soybean, chickpea and lupine flour. *Die Nahrung.* **34**: 835–841.
- Hegsted, M., Windhauser, M. M., Morris, S. K., and Lester, S. B. (1993). Stabilized rice bran and oat bran lower cholesterol in humans. *Nutr. Res.* **13**: 387-398.
- Hesser, J. M. (1994). Applications and usage of dietary fiber in the USA. *Int. Food Ing.* **2**: 50–52.
- Hoseney, R. C. (1986). Yeast Leavened Products. In: Principles of Cereal Science and Technology. pp. 203. Am. Assoc. Cereal Chem. Inc., St. Paul, Minnesota.
- Hu, W., Wells, J. H., Shin, T. S., and Godber, J. S. (1996). Comparison of isopropanol and hexane for extraction of vitamin E and oryzanols from stabilized rice bran. *J. Am. Oil Chem. Soci.* **73**: 1653–1656.

Hundemer, J. K., Nabar, S. P., Shriver, B. J., and Forman, L. P. (1991). Dietary fiber sources lower blood cholesterol in C57BL/6 mice. *J. Nutr.* **121**: 1360–1365.

Indravathamma, P., and Seshadri, H. S. (1980). Lectin from rice. *J. Biosci.* **2**: 29-36.

Jangbloed, A. W., Everts, H., and Kemme, P. A. (1991). Recent advances in animal nutrition. Butter-Worth, London.

Jariwalla, R. J. (2001). Rice bran products: Phytonutrients with potential applications in preventive and clinical medicine. *Drugs Exp. Clin. Res.* **27**: 17–26.

Jiamyangyuen, S., Harper, W.J., Srijesdaruk, V., and Kumthonglang, K. (2005a). Study of extraction and functional properties of rice bran protein concentrate. *Milchwissenschaft.* **60**(2): 192-195.

Jiamyangyuen, S., Srijesdaruk, V., and Harper, W.J. (2005b). Extraction of rice bran protein concentrate and its application in bread. *Songklanakarin J. Sci. Technol.* **27**(1): 55-64.

Ju, Y. H., and Vali, S. R. (2005). Rice bran oil as a potential resource for biodiesel. *J. Sci. Ind. Res.* **64**: 801-822.

Juliano, B. O. (1985). Rice: Chemistry and Technology, 2nd Ed. Am. Assoc. Cereal Chem. Inc., St. Paul. Minnesota, USA.

Kahlon, T. S., Chow, F. I., Sayre, S. R., and Betschart, A. A. (1992). Cholesterol lowering in hamsters fed rice bran at various levels, defatted rice bran and rice bran oil. *J. Nutr.* **122**: 513-519.

- Kahlon, T. S., Edwards, R. H., and Chow, F. I. (1998). Effect of extrusion on hypocholesterolemic properties of rice, oat, corn, and wheat bran diets in hamsters. *Cereal Chem.* **75**: 897-903.
- Kahlon, T. S., Saunders, R. M., Chow, F. I., Chiu, M. C., and Bestchart, A. A. (1990). Influence of rice bran, oat bran and wheat bran on cholesterol and triglycerides in hamsters. *Cereal Foods World.* **67**: 439-443.
- Kennedy, G., and Burlingame, B. (2003). Analysis of food composition data on rice from a plant genetic resources perspective. *Food Chem.* **80**: 589–596.
- Kennedy, L. L., Chung, H. Y., and Hegsted, M. (1996). New food products developed from rice bran fractions. In: IFT Book of Abstracts. IFT Annual Meeting: Book of Abst. pp. 56. Institute of Food Technologists, USA.
- Kestin, M., Moss, R., Clifton, P. M., and Nestle, P. J. (1990). Comparative effect of three cereal brans on plasma lipids, blood pressure and glucose metabolism in mildly hypercholesterolemic men. *Am. J. Clin. Nutr.* **52**:661-666.
- Khor, H. T., and Ng, T. T. (2000). Effects of administration of tocopherol and tocotrienols on serum lipids and liver HMG CoA reductase activity. *Int. J. Food Sci. Nutr.* **51**: S3–S11.
- Khor, H. T., Chieng, D. Y., and Ong, K. K. (1995). Tocotrienols inhibit HMG-CoA reductase activity in the guinea pig. *Nutr. Res.* **15**: 537–544.

- Kim, C. J., Byun, S. M., Cheigh, H. S., and Kwon, T. W. (1987). Optimization of extrusion of rice bran stabilization process. *J. Am. Oil Chem. Soc.* **64**: 514–516.
- Kim, J. S., and Godber, J. S. (2001). Oxidative stability and vitamin E levels increased in restructured beef roasts with added rice bran oil. *J. Food Qual.* **24**: 17–26.
- Kratzer, F. H., and Payne, C. G. (1977). Effect of autoclaving, hot water treatment, parboiling and addition of ethoxyquin on the value of rice bran as a dietary ingredient for chickens. *Brit. Poult. Sci.* **18**: 475–482.
- Krishna, A. G. G. (2002). Nutritional components of rice bran oil in relation to processing. *Lipid Technol.* **14**: 80–84.
- Krishna, A. G. G., Khatoon, S. and Babylatha, R. (2005). Frying performance of processed rice bran oils. *J. Food Lipids.* **12**: 1–11.
- Lakkakula, N. R., Lima, M., and Walker, T. (2004). Rice bran stabilization and rice bran oil extraction using ohmic heating. *Bioresour. Technol.* **92**: 157–161.
- Lasztity, R., Tomoskozi, S., Nagy, J., Bajkai, T., and Sarkadi, L. (1995). Investigation of some functional and nutritive properties of cereal germ proteins. *Per. Pol. Chem. Eng.* **39**(1): 63–76.
- Lee, S.K., Jang, I.S., Kim, M.K., Park, S.K., Lee, W.Y., Youn, K.S., and Bae, D.H. (2004). Changes in functional properties of rice bran and sesame meal proteins through chemical modifications. *Food Sci. Biotech.* **13**: 555–560.

- Lima, I., Guaraya, H., and Champagne, E. (2002). The functional effectiveness of reprocessed rice bran as an ingredient in bakery products. *Nahrung*. **46**: 112-117.
- Lin, S. H. C., and Carter, C. M. (1973). Effect of extrusion cooking on the formation of free fatty acids in rice bran. Food Protein R&D Center. College Station, Texas A&M Uni. Texas, U.S.A.
- Lynn, L. (1969). Edible rice bran foods. *Cereal Chem.* **46**: 54-172.
- Malekian, F., Rao, R. M., Prinyawiwatukul, W., Marshall, W. E., Windhauser, M., and Ahmedna, M. (2000). Lipase and lipoxygenase activity, functionality, and nutrient losses in rice bran during storage. Agricultural Center, Louisiana Agricultural Experiment Station, Baton Rouge. Bulletin No. **870**: 1-68.
- Matz, S. A. (1991). The Chemistry and Technology of Cereals as Food and Feed. pp. 731–741. Van Nostrand Reinhold Pub., New York.
- McCaskill, D. R., and Zhang, F. (1999). Use of rice bran oil in foods. *Food Technol.* **53**: 50–53.
- McKee, L. H., and Latner, T. A. (2000). Underutilized sources of dietary fiber: a review. *Plant Foods Hum. Nutr.* **55**: 285–304.
- McWatters, K. H., Ouedraogo, J. B., Resurreccion, A. V. A., Hung, Y. C., and Dixon, R. (2003). Physical and sensory characteristics of sugar cookies containing mixtures of wheat, fonio (*Digitaria exilis*) and cowpea (*Vigna unguiculata*) flours. *Int. J. Food Sci. Technol.* **38**: 403–410.

- Narisullah, and Krishnamurthy, M. N. (1989). Effect of stabilization on the quality characteristics of rice bran oil. *J. Am. Chem. Oil Soc.* **66**: 661-663.
- Nasirullah. (2001). Development of deep frying edible vegetable oils. *J. Food Lipids.* **8**: 295–304.
- Neto, V.Q., Narain, N., Silva, J.B., and Bora, P.S. (2001). Functional properties of raw and heat processed cashew nut (*Anacardium occidentale* L.) kernel protein isolates. *Nahrung.* **45**(4): 258-262.
- Newman, R. K., Betschart, A. A., Newman, C. W., and Hofer, P. J. (1992). Effect of full-fat or defatted rice bran on serum cholesterol. *Plant Foods Hum. Nutr.* **42**: 37-43.
- Nicolosi, R. J., Ausman, L. M., and Hegsted, D. M. (1991). Rice bran oil lowers serum total and low density lipoprotein cholesterol and apo B levels in nonhuman primates. *Atherosclerosis.* **88**: 133–142.
- Orthoefer, F. T. (1996). Rice bran oil: Healthy lipid source. *Food Technol.* **50**: 62- 64.
- Ory, R. L., Bog, H. T. C., and Mod, R. R. (1981). Properties of Haemagglutinin in rice and other cereal grains. In: Antineutrinos and Natural Toxicants in Foods. pp. 159-168. Food Nutr. Press, Inc., Westport.
- Parker, R. A., Pearce, B. C., Clark, R. W., Gordon D. A., and Wright, J. J. (1993). Tocotrienols regulate cholesterol production in mammalian cells by post-transcriptional suppression of 3-hydroxy-3-methylglutarylcoenzyme-A reductase. *J. Biol. Chem.* **268**: 11230–11238.

- Patel, P., and Walker, T. (2004). Effect of extraction method on the level of important antioxidants in rice bran oil. ASAE/CSAE Meeting Paper No. 046140. ASAE/CSAE Fairmont Chateau Laurier, the Westin, Government Centre Ottawa, Ontario, Canada.
- Pearce, B. C., Parker, R. A., Deason, M. E., Qureshi, A. A., and Wright, J. J. K. (1992). Hypocholesterolemic activity of synthetic and natural tocotrienols. *J. Med. Chem.* **35**: 3595–3606.
- Prabhakar, J. V. (1987). Stabilization of bran. In: Handbook on rice bran. Solvent Extractors Association of India, India.
- Prakash, J., and Jyothilakshmi, A. (1995). Utilizing fullfat rice bran in traditional foods. In: IFT Book of Abstracts. Abstract No. 81C-4. pp. 243. Institute of Food Technologists, USA.
- Pszczola, D. E. (2001). Rice: not just for throwing. *Food Technol.* **55**: 53–59.
- Purushothama, S., Raina, P. L., and Hariharan, K. (1995). Effect of long term feeding of rice bran oil upon lipids and lipoproteins in rats. *Mol. Cell. Biochem.* **146**: 63-69.
- Qureshi, A. A., Mo, H., Packer, L., and Peterson, D. M. (2000). Isolation and identification of novel tocotrienols from rice bran with hypocholesterolemic, antioxidant, and antitumor properties. *J. Agric. Food Chem.* **48**: 3130–3140.
- Qureshi, A. A., Sami, S. A., Salser, W. A., and Khan, F. A. (2002). Dose dependent suppression of serum cholesterol by tocotrienol rich fraction (TRF25) of rice bran in hypercholesterolemic humans. *Atherosclerosis.* **161**: 199–207.

Raghuram, T. C., and Rukmini, C. (1995). Nutritional significance of rice bran oil. *Indian J. Med. Res.* **102**: 241–244.

Rajeshwara, A. N., and Prakash, V. (1995). Purification and characterization of lipase from rice (*Oryza sativa* L.) bran. *Die Nahrung.* **39**: 406–418.

Ramezanzadeh, F. M., Rao, R. M., Windhauser, M., Prinyawiwatkul, W., and Marshall, W. E. (1999). Prevention of oxidative rancidity in rice bran during storage. *J. Agric. Food Chem.* **47**: 2997–3000.

Ramezanzadeh, F., Rao, R., Windhauser, M., Witoon, P., Tulley, R., and Marshall, W. (1999). Prevention of hydrolytic rancidity in rice bran during storage. *J. Agric. Food Chem.* **47**: 3050–3052.

Ramzan, M. (2000). Effect of Reduction of Crude Fiber and Phytic Acid in Treated Defatted Rice Bran on the Performance of Broilers. M. Sc. Thesis, C.V.S., Lahore, Univ. Agri. Faisalabad, Pakistan.

Randall, J. M., Sayre, R. N., Schultz, W. G., Fong, R. Y., Mossman, A. P., Tribelhorn, R. E., and Saunders, R. M. (1985). Rice bran stabilization by extrusion cooking for extraction of edible oil. *J. Food Sci.* **50**: 361–364.

Rong, N., Ausman, L. M., and Nicolosi, R. J. (1997) Oryzanol decreases cholesterol absorption and aortic fatty streaks in Hamsters. *Lipids.* **32**: 303-309.

Rukmani, C. (2002). What is rice bran (a web article on rice bran). Available from: www.moormans.com/equine/TechBulletins/WhatIsRiceBran.htm.

Rukmini, C., and Raghuram. (1991). Nutritional and biochemical aspects of the hypolipidemic action of rice bran oil: A review. *J. Am. Coll. Nutr.* **10**: 593-601.

Sakamoto, K., Tabata, T., Shirasaki, K., Inagaki, T., and Nakayama, S. (1987). Effects of gamma-oryzanol and cycloartenol ferulic acid ester on cholesterol diet induced hyperlipidemia in rats. *Jpn. J. Pharm.* **45**: 559-65.

Salunkhe, D. K., Chavan, J. K., Adsule, R. N., and Kadam, S. S. (1992). Rice. In: *World Oilseeds: Chemistry, Technology and Utilization*. pp. 424–448. Van Nostrand Reinhold, New York.

Sanders, T. A., and Reddy, S. (1992). The influence of rice bran on plasma lipids and lipoproteins in human volunteers. *Eur. J. Clin. Nutr.* **46**: 167-72.

Sarkar, S. (1992). Nutrition of rice bran oil in relation to its purification. *J. Am. Oil Chem. Soc.* **68**: 956-962.

Saunders, R. M. (1986). Rice bran composition and potential food uses. *Food Rev. Int.* **1**: 465-495.

Saunders, R. M. (1990). The properties of rice bran as a food stuff. *Cereal Foods World.* **35**: 632-639.

- Sayre, R. N., and Saunders, R. M. (1985). Extraction and refining of edible oil extrusion stabilized rice bran. *J. Oil Chem. Soc.* **62**: 1040-1043.
- Sayre, R. N., Saunders, R. M., Enochian, R. V., Schultz, W. G., and Beagle, E. C. (1982). Review of rice bran stabilization systems with emphasis on extrusion cooking. *Cereal Foods World.* **27**: 317.
- Seetharamaiah, G. S. and Chandrasekhara, N. (1989). Studies on hypocholesterolemic activity of rice bran oil. *Atherosclerosis.* **78**: 219–223.
- Seetharamaiah, G. S., Krishnakantha, T. P., and Chandrasekhara, N. (1990). Influence of oryzanol on platelet aggregation in rats. *J. Nutr. Sci. Vitaminol.* **36**: 291-297.
- Sekhon, K. S., Dhillon, S. S., Singh, N., and Singh, B. (1997). Functional suitability of commercially milled rice bran in India for use in different food products. *Plant Foods Hum. Nutr.* **50**: 127-140.
- Sharif, K., Butt, M. S., and Huma, N. (2005). Oil extraction from rice industrial waste and its effect on physico-chemical characteristics of cookies. *Nut. Food Sci.* **35**(6): 416-427.
- Sharif, M. K., Butt, M. S., Anjum, F. M., and Nawaz, H. (2009). Preparation of fiber and mineral enriched defatted rice bran supplemented cookies. 2009. *Pak. J. Nutrition.* **8**(5): 571-577.
- Sharma, H. R., and Chauhan, G. S. (2002). Effect of stabilized rice bran-fenugreek blends on the quality of breads and cookies. *J. Food Sci. Nutr.* **39**: 225-233.

Sharma, R. D., and Rukmini, C. (1986). Rice bran oil and hypocholesterolemia in rats. *Lipids*. **21**: 715-717.

Sharma, R. D., and Rukmini, C. (1987). Hypocholesterolemic activity of unsaponifiable fraction of rice bran oil. *Ind. J. Med. Res.* **85**: 278-281.

Sharp, C. Q., and Kitchen, K. J. (1990). Using rice bran in yeast bread in a home baker. *Cereal Foods World*. **35**: 1021-1028.

Shih, F.F. (1994). Interaction of soy isolate with polysaccharide and its effect on film properties. *J. Am. Oil Chem. Soc.* **71**: 1281-1285.

Shih, F.F. (1996). Edible films from rice protein concentrate and pullulan. *Cereal Chem.* **73**(3): 406-409.

Shin, D. H., and Chung, J. K. (1998). Changes during storage of rice germ oil and its fatty acid composition. *Korean J. Food. Sci. Technol.* **30**: 77-81.

Singh, B., Bajaj, M., Kaur, A., Sharma, S., and Sidhu, J. S. (1993). Studies on the development of high-protein biscuits from composite flour. *Plant Foods Hum. Nutr.* **43**: 181-189.

Singh, B., Sekhon, K. S., and Singh, N. (1995). Suitability of fullfat and defatted rice bran obtained from Indian rice for use in food products. *Plant Foods Hum. Nutr.* **47**: 191-200.

Singh, N., Harinder, K., Sekhon, K. S., and Kaur, B. (1991). Studies on the improvement of functional and baking properties of wheat-chickpea flour blends. *J. Food Proc. Pres.* **15**: 391-402.

- Singh, N., Kaur, A., Singh, R. P., and Sekhon, K. S. (1989). Rheological and cookie making studies on wheat-rice flour blends. *J. Food Sci. Tech.* **26**: 90–94.
- Stoggl, W., Christain, H., Surapote, W., Heimo, S., and Gonther, B. (2005). Simultaneous determination of carotenoids, tocopherols, and γ -oryzanol in crude rice bran oil by liquid chromatography coupled to diode array and mass spectrometric detection employing silica C₃₀ stationary phases. *J. Sep. Sci.* **28**: 1712–1718.
- Sugano, M., and Tsuji, E. (1997). Rice bran oil and cholesterol metabolism. *J. Nutr.* **127**: 521–524.
- Sugano, M., Koba, K., and Tsuji, E. (1999). Health benefits of rice bran oil. *Anticancer Res.* **19**: 3651–3657.
- Takahashi, T., Yamada, N., Iwamoto, L., Shimabayashi, Y., and Izutsu, K. (1973). Some properties and characterization of rice seed haemagglutinin. *Agri. Biol. Chem.* **37**: 29–36.
- Tao, J. (2001). Method of stabilization of rice bran by acid treatment and composition of the same. U.S. Patent No. 6245377.
- Tomlin, J., and Read, N. W. (1988). Comparison of the effects on clonic function caused by feeding rice bran and wheat bran. *Eur. J. Clin. Nutr.* **42**: 857.
- Topping, D. L., Illman, R. J., Roach, P. D., Trimble, R. P., Kambouris, A., and Nestel, P. J. (1990). Modulation of the hypolipidemic effect of fish oils by dietary fiber in rats: Studies with rice and wheat bran. *J. Nutr.* **120**: 325–330.

- Trautwein, E. A., Schulz, C., Rieckhoff, D., Rau, A. K., Erbersdobler, H. F., Groot, W. A., and Meijer, G. W. (2002). Effect of esterified 4-Desmethylsterols and 4-stanols or 4,4'-Dimethylsterols on cholesterol and bile acid metabolism in hamsters. *Br. J. Nutr.* **87**: 227-237.
- Tsen, C. C., Peters, E. M., Schaffer, T., and Hoover, W. J. (1973). High-protein cookies: Effect of soy fortification and surfactants. *Baker's Digest.* **47**: 34-39.
- Tsuda, M. (1979). Purification and characterization of a lectin from rice bran. *J. Biochem.* **86**: 1451-1461.
- Tsuji, H., Kimoto, M., and Natori, Y. (2001). Allergens in major crops. *Nutr. Res.* **21**: 925-934.
- Vissers, M. N., Peter, L. Z., Gert, W. M., and Martijn, B. K. (2000). Effect of plant sterols from rice bran oil and triterpene alcohols from sheanut oil on serum lipoprotein concentrations in humans. *Am. J. Clin. Nutr.* **72**: 1510-1515.
- Wade, P. (1988). Biscuit, Cookies and Crackers: The Principles of the Craft. Vol. I. Elsevier Applied Sci., London.
- Wagner, J.R., Sorgentini, D.A., and Anon, M.C. (1996). Thermal and electrophoretic behavior, hydrophobicity, and some functional properties of acid-treated soy isolates. *J. Agric. Food Chem.* **44**: 1881-1889.
- Wang, M., Hettiarachchy, N.S., Qi, M., Burks, W., and Siebenmorgen, T. (1999). Preparation and functional properties of rice bran protein isolate. *J. Agric. Food Chem.* **47**(2): 411-416.

- Watchararужи, K., Goto, M., Sasaki, M., and Shotipruk, A. (2008). Value-added subcritical water hydrolysate from rice bran and soybean meal. *Biores. Technol.* **99**: 6207-6213.
- Weststrate, J. A., and Meijer, G. W. (1998). Plant sterol-enriched margarines and reduction of plasma total and LDL-cholesterol concentrations in normocholesterolaemic and mildly hypercholesterolaemic subjects. *Eur. J. Clin. Nutr.* **52**: 334-343.
- WHO (World Health Organization). (2003). Diet, nutrition and the prevention of chronic diseases. Report of a joint WHO/FAO expert consultation. WHO Technical Report Series 916. World Health Organization, USA.
- Wilson, T. A., Ausman, L. M., Lawton, C. W., Hegsted, D. M., and Nicolosi, R. J. (2000). Comparative cholesterol lowering properties of vegetable oils: Beyond fatty acids. *J. Am. Coll. Nutr.* **19**: 601-607.
- Wilson, T. A., Nicolosi, R. J., Woolfrey, B., and Kritchevsky, D. (2007). Rice bran oil and oryzanol reduce plasma lipid and lipoprotein cholesterol concentrations and aortic cholesterol ester accumulation to a greater extent than ferulic acid in hypercholesterolemic hamsters. *J. Nutr. Biochem.* **18**: 105–112.
- Wiseman, M.O., and Price, R.L. (1987). Functional properties of protein concentrates from pressed jojoba meal. *Cereal Chem.* **64**(2): 94-97.
- Xu, Z. (1998). Purification and Antioxidant Properties of Rice Bran γ -Oryzanol Components. Ph.D. dissertation. Louisiana State Uni., U.S.A.

- Xu, Z., Hua, N., and Godber, J. S. (2001). Antioxidant activity of tocopherols, tocotrienols, and γ -oryzanol components from rice bran against cholesterol oxidation accelerated by 2,2V-azobis (2-methylpropionamidine) dihydrochloride. *J. Agric. Food Chem.* **49**: 2077-2081.
- Yeager, S. (1998). Fiber - the ultimate healer. In: *The Doctors Book of Food Remedies*. pp. 184-185. Rodale Press, Inc., Emmaus, Pennsylvania.
- Yoshida, Y., Niki, E., and Noguchi, N. (2003). Comparative study on the action of tocopherols and tocotrienols as antioxidant: chemical and physical effects. *Chem. Phys. Lipids*. **123**: 63-75.
- Yoshino, G., Kazumi, T., Amano, M., Takeiwa, M., Yamasaki, T., Takashima, S., Iwai, M., Hatanaka, H., and Baba, S. (1989). Effects of γ -oryzanol on hyperlipidemic subjects. *Current Therapy Res.* **45**: 543-552.
- Zhu, H. (2002). Utilization of rice bran by *Pythium irregulare* for lipid production. M.S. Thesis, Department of Biological and Agricultural Engineering, Agricultural and Mechanical College, Louisiana State Uni. U.S.A.
- Zumbado, H., Ledesma, L., Fuertes, S., and Ventura, J. (1997). Processing of a baked product with high levels of parboiled rice bran. *Alimentaria*. **280**: 21-23.