

Black Tea Polyphenols: A Mechanistic Treatise

M. S. Butt , A. Imran , M. K. Sharif , Rabia Shabir Ahmad , Hang Xiao , M. Imran & H. A. Rsool

To cite this article: M. S. Butt , A. Imran , M. K. Sharif , Rabia Shabir Ahmad , Hang Xiao , M. Imran & H. A. Rsool (2014) Black Tea Polyphenols: A Mechanistic Treatise, Critical Reviews in Food Science and Nutrition, 54:8, 1002-1011, DOI: [10.1080/10408398.2011.623198](https://doi.org/10.1080/10408398.2011.623198)

To link to this article: <https://doi.org/10.1080/10408398.2011.623198>



Accepted author version posted online: 08 Mar 2013.
Published online: 08 Mar 2013.



Submit your article to this journal [↗](#)



Article views: 973



View Crossmark data [↗](#)



Citing articles: 23 View citing articles [↗](#)

Black Tea Polyphenols: A Mechanistic Treatise

M. S. BUTT,¹ A. IMRAN,² M. K. SHARIF,¹ RABIA SHABIR AHMAD,²
HANG XIAO,³ M. IMRAN,¹ and H. A. RSOOL¹

¹National Institute of Food Science and Technology, University of Agriculture, Faisalabad, Pakistan

²Institute of Home and Food Sciences, GC University, Faisalabad, Pakistan

³Department of Food Science, University of Massachusetts, Amherst, MA, USA

Dietary interventions are among the emerging trends to curtail physiological malfunctioning like cancer, diabetes, cardiac complications, etc. The essence of phytonutrients has developed the concept of nutraceuticals at the junction of diet health linkages. In this context, theaflavin & thearubigins are the oxidized derivatives of black tea catechins during fermentation having nutraceutical potential owing to esterification of hydroxyl ring with digallate esters. Theaflavin may influence activation of transcription factors such as NF- κ B or AP-1 that ultimately hinder the formation of nitric oxide expression gene. Likewise, black tea contains a unique amino acid theanine acts as neurotransmitter owing to its ability to cross the blood–brain barrier. Moreover, it boasts immunity by enhancing the disease-fighting ability of gamma delta T cells. Theaflavin & thearubigins act as safeguard against oxidative stress thereby effective in the cardiac functioning. The mechanistic approach of these antioxidants is likely to be associated with inhibition of redox sensitive transcription factors & pro-oxidant enzymes such as xanthine oxidase or nitric oxide synthase. However, their involvement in antioxidative enzyme induction as in glutathione-S-transferases is also well documented. They act as curative agent against numerous pathological disorders by disrupting the electron chain thus inhibiting the progression of certain ailments. Black tea polyphenols established themselves as strong antioxidants due to their standard one-electron potential, and their vitality is dependent on the concentration of polyphenols and pH for their inclusive execution. Present review is an attempt to enrich the readers regarding the health promoting aspects of black tea polyphenols. Concomitantly, it needs core attention of researchers for the exploitations of black tea flavanols as an important dietary constituent for the vulnerable segment.

Keywords Black tea, polyphenols, flavanols, cancer, oxidative stress, theaflavin, thearubigins

INTRODUCTION

Dietary guidelines focus on the dynamic aspects of phytonutrients as they exert beneficial effects on human health particularly against cardiovascular complications, neurodegenerative diseases, diabetes, and numerous oncogenic events. This veracity has encouraged the supplementation of plant-derived nutraceuticals that, in turn, modulate the onset of chronic ailments. Epidemiological studies have correlated the capacity of numerous phytochemicals particularly polyphenols with improved health claims (Jayasekera et al., 2011). Owing to the health protective functions, black tea and its constituents are extensively investigated dietary source of polyphenols (Turkmen et al., 2006; Almajano et al., 2008).

Tea, a popular beverage after water, made from the leaves of tea plant (*Camellia sinensis*). The historians have linked its consumption almost 5000 years back (Yang et al., 2009). On the basis of fermentation, tea is mainly divided into four distinctive types white, green, oolong, and black tea that differ in terms of processing method and chemical profile (Cabrera et al., 2003). It has been reported that during 2008, the total production of tea was 3.7 billion kg (Letchoumy et al., 2008). Green tea account for approximately 20% of total tea produced and is consumed primarily in East and South East Asia. Contrarily, black tea that occupied approximately 78% of world tea production is consumed mainly in North America, Europe, and North Africa. Green tea is prepared by pan-frying or steaming fresh tea leaves at elevated temperature to inhibit polyphenol-oxidase mediated processes, whereas oolong tea is semifermented product (Imran et al., 2012). During the manufacturing of black tea, leaves are crushed and subjected to enzymatic oxidation process called fermentation (Hsu et al., in press). Subsequent oxidative

Address correspondence to Dr. Ali Imran, PhD, 100 Holdsworth Way, Amherst, MA 01003. E-mail: aliimran.ft@gmail.com

Table 1 Chemical composition of black tea

| Constituents | Concentration (%) |
|----------------|-------------------|
| Catechins | 3–10 |
| Theaflavins | 3–6 |
| Carbohydrates | 15 |
| Thearubigins | 12–18 |
| Protein | 1 |
| Flavonols | 6–8 |
| Mineral matter | 10 |
| Phenolic acids | 10–12 |
| Volatiles | 1 |
| Amino acids | 13–15 |
| Methylxanthine | 8–11 |

Source: Sharangi (2009).

condensation of the catechins during the fermentation leads to production of theaflavins (TFs) (benzotropolone dimers of catechins) as well as higher molecular weight polymers, i.e., thearubigins (TRs). Both constituents are responsible for specific taste and color of black tea. Polyphenolic compounds including substantial amounts of flavonoids dominate typical composition of black tea (Bursill et al., 2007). Flavonoids are classified into six groups based on structure and position of the heterocyclic oxygen carbon oxygen ring, i.e., flavones, flavanones, isoflavones, flavonols, flavanols, and anthocyanins. Flavanols consist of unoxidized catechins including TF and TRs, while flavonols are characterized by quercetin, kaempferol, and myricetin. Tea also contains phenolic acids (caffeic, quinic, and gallic acids), caffeine, theobromine, theophylline, and many flavors compounds enriched with linalool (Dimpfel et al., 2007). In addition, tea carries unique amino acid, theanine that has been extensively studied for biological activities. Tea composition is likely to be associated with origin, fermentation conditions, and processing (Table 1). Black tea health claims are allied to the TF, TRs, catechins, theanine, and flavonols (Mei et al., 2009). The brief description of tea phytochemicals is given below.

THEAFLAVIN

Theaflavins constitute a complicated structure of hydroxy-substituted benzotropolone ring that is synthesized by condensation of catechins in their hydroxylated B rings (di and tri forms), attaching with benzotropolone catechins (Ju et al., 2007). The reaction of abridgment initiates by B ring oxidation to the quinines, with addition of the gallocatechin quinone to the catechin quinone that ultimately brings down decarboxylation termed as Michael addition reaction (Kaur et al., 2007).

Theaflavin is a mixture of four compounds termed as theaflavin (TF1), theaflavin-3-gallate (TF2A), theaflavin-3'-gallate (TF2B), and theaflavin-3, 3'-digallate (TF3). The molecular formulas of TF1, TF2 (including TF2A and TF2B), and TF3 are $C_{29}H_{24}O_{12}$, $C_{39}H_{28}O_{16}$, and $C_{43}H_{32}O_{20}$, respectively. Their chemical structures and route of formation is shown in

Fig. 1. Several in vitro researches in TF chemistry have chalked out their new analogs. In this context theaflavate A is a famous resultant product of epicatechin gallate (EC) oxidation by potassium ferricyanide (Halliwell, 2007). Alongside to theaflavate A, other active forms of TFs including isotheaflavin-3'-o-gallate and neotheaflavin-3-o-gallate have also been reported containing a benzotropolone moiety common in their structure (Inami et al., 2007). TFs are natural antioxidant and metal chelators due to the presence of hydroxy groups; however, the gallic acid moiety is also a key contributor in this regard (Kumar et al., 2007; Lambert et al., 2007a). The majority of studies have reported elevated antioxidant efficiency of TFs by the esterification of hydroxyl ring with digallate esters (Lambert et al., 2007b). Furthermore, it has been revealed that the rate constant of TFs as a superoxide scavenging is higher in comparison with epigallocatechin gallate (EGCG) (Maeta et al., 2007). Likewise, preventive role of TFs in lipid preoxidation is mainly attributed to its ability to cease the chain reaction of lipid oxidation. Apart from free radical scavenging and metal chelating abilities, TF has potential to activate certain antioxidant enzymes like glutathione-S-transferases (GST), glutathione peroxides (GPX), superoxide dismutase (SOD), and catalase (CAT), which ultimately resulting lipid peroxidation reduction (Maki et al., 2009; Mazzanti et al., 2009).

THEARUBIGINS

Recent studies have documented TRs as the most copious group of phenolics found in black tea, contributing almost 60% of the solids in typical black tea (Nagao et al., 2009; Schneider and Segre, 2009). Among the black tea polyphenols, TRs having anti-initiating properties against cancer cells referred as polymeric black tea polyphenols (PBPs) (Patel et al., 2007). They are providing reddish tonality to black tea in contrast with green and white tea. The color of a black tea, however, is not only affected by the TRs, but some other intrinsic compounds like unoxidized polyphenols and TFs are also involved (Das et al., 2009; Schneider and Segre, 2009). Originally, TRs formation (Figure 1) is initiated after conversion of 75% the phenolic flavan-3-ol molecules from catechins. Nutraceutical properties of tea are accredited to the presence of TRs, which not only help mitigate various ailments but also enhance its credit and integrity toward the consumer acceptance. Numerous studies have investigated that the TRs in tea is intricate due to their varying concentrations that are often found in a chaotic manner as 20% in black tea leaf and 60% as solids in final black tea infusion (Gupta et al., 2009; Thielecke and Boschmann, 2009). TRs have gained special interest for their ability to activate phase II enzymes by inducing the transcriptional upregulation in liver and lungs after elevation of Nrf2-mediated antioxidant-responsive element binding. However, PBP extract has not been reported for the alteration of Nrf2 or Keap1 at the transcriptional level may increased by posttranslational modifications such as phosphorylation and decreased ubiquitination. Pretreatment of TRs is




Figure 1 Structure and mode of formation of flavanols.

responsible for the nuclear translocation of PKC and PI3-kinase-mediated phosphorylation of Nrf2, critical for the release of Nrf2 from Keap1 and its subsequent. Moreover, their anticancer concerns are probed due to their ability to decrease carcinogen DNA adducts by inactivation of carcinogen-induced phase I enzymes such as CYP450 1A1 and 1A2 (Patel and Maru, 2008).

THEANINE

Theanine (gamma-glutamylethylamide, or 5-N-ethyl-glutamine) (Fig. 2) is a glutamic acid analog commonly identified in tea (*Camellia sinensis*). Synonymously termed as L-theanine, (Table 2) due to its molecular orientation and clas-

sified as levorotary enantiomer. L-Theanine is a predominant amino acid, 1–2% of the dry weight of the leaves comprises about 50% of total protein (Palva and Palva, 2007). It was first identified by Sakato in tea leaves as the main free amino acid component that can influence the major umami constituents of tea (Hideyuki et al., 2002). Various health effects have been associated with L-theanine, such as neuroprotective effects, which ultimately enhanced nervous attention. On the basis of two parallel techniques, i.e., electroencephalographic (EEG) and behavioral modules indicate that L-theanine influences functionality of brain (De Mejia et al., 2009). Current consensus regarding theanine has explained its effect in mitigation of mental and physical stress due to its ability to cross the blood–brain barrier (Boris et al., 2009). Moreover, it enhances cognition and behavior in a systematic manner with caffeine.

Table 2 Promising Characteristic of black tea polyphenols



| Name of constituents | Silent features | Health claims | Sources | References |
|----------------------|--|--|--|----------------------|
| Catechin | White in color crystalline structure Mol. Wt. 302.2357 Boiling point 642.4 Density is 1.799 g/cm ³ | Act as strong antioxidant. Provide protection against cancer, obesity, and diabetic | Green tea, black tea, onion garlic, etc. | Boris et al., 2009 |
| Theaflavin | Orange-colored crystalline structure Mol. for C ₂₉ H ₂₄ O ₉ For. Wt. 516.5 Boiling point 1003.9° Density 1.87 g/cm ³ | Strong antioxidant provide protection against cancer and other life related disorders | Black tea | Popkin, 2010 |
| Thearubigin | Dark brown crystalline structure Heterogeneous in nature Molecular mass 700–40,000 Da. Water soluble and acidic | Provide multiplicity of health claims dominated by oxidative stress and cardiovascular compliances | Black tea | Popkin, 2010 |
| Theanine | Gammaglutamylethylamide analog amino acid levorotary enantiomer GRAS by FDA Mol. Wt. 174.20 mp. 202–215°C | Neurotransmitter enhance resistance against Alzheimer disease | Green and black tea | Rains et al., 2011 |
| Quercetin | Yellow crystalline powder mp. 310–317°C stable under normal conditions Mol. for C ₁₅ H ₁₀ O ₇ | Strong antioxidant status. Important in oral health reduce cholesterol | Vegetables like cabbages brocolis, onion, and green and black tea | Amić and Lučić, 2010 |
| Myricetin | White powder that was slightly soluble in water Mol. Wt. 318. 2351 mp. 300°C mp. 745.45°C | Metal chelating and cancer inhibiting properties | Black tea, green tea onion and mango peel | Amić and Lučić, 2010 |
| Kaempferol | Yellow solid having the crystalline structure. Boiling point 276–278°C. sparingly soluble in water readily soluble in hot organic solvents | Reduce cancer and act as antibacterial agent | Walnut is its richest source also reported in green and black tea | Amić and Lučić, 2010 |

In addition, L-theanine helps boost immunity against infection by enhancing the disease-fighting ability of gamma delta T cells (Sugiyama and Sadzuka, 2003). Recently, array of biochemical modulations proved theanine effective to enhance the activity of several antitumor agents, including doxorubicin (DOX) (Popkin, 2010). In addition to neurofunctionality and antitumor activities, its role as hepatic protective agent is also well documented. During oxidative stress it protects the liver damage due to increase in glutathione level via metabolism through glutaminase and c-GTP (Van der Pilji et al., 2010).

FLAVONOLS

Flavonols are often termed as aglycons due to their occurrence in plants in the form of glycosides. Mostly, glucose and galactose sugar moieties are attached with aglycons, re-

sponsible for their rapid metabolism (Baoru et al., 2009; Ou et al., 2002). Their composition and availability is influenced by numerous factors like plant variations, soil, season, light, degree of ripeness, food preparation, and processing (Ross et al., 2000). They are formed by the combination of derivatives synthesized from phenylalanine (via shikimic acid pathway) and acetic acid; nevertheless, formation of phenylalanine from phenylpyruvate is a crucial step. Furthermore, their absorption is dependent on attached sugar moiety on flavonol glycosides thereby effecting absorption rate (Noelia et al., 2010). They comprised kaempferol, myricetin, and quercetin (Fig. 3), whereas kaempferol is potentially present in broccoli, grapefruit, brussels sprout, and apple (Smith et al., 2000). It has yellowish tonality with boiling point ranging from 276–278°C

**Figure 2** Structural representation of theanine.

R = R' = H Kaempferol
R = H, R' = OH quercetin
R = R' = OH Myricetin

Figure 3 Structural representation of flavonols.

(Table 2). It is sparingly soluble in water but readily soluble in hot ethanol, diethyl ether, and other organic solvents (Rains et al., 2011). Its consumption through tea and other natural commodities has closely been associated with reduced cardiac complications and used in antidepressant drugs (Amić and Lučić, 2010).

Myricetin is predominantly found in many fruits like grapes, berry, some vegetables, herbs, and walnuts. In vitro research has proved myricetin as an active ingredient to reduce low-density lipoprotein (LDL) cholesterol by increasing the uptake and modifying the white blood cells that ultimately responsible for scavenging the free radicals that produced due to oxidation of LDL. In some scientific studies, high myricetin consumption resulted in lowering the onset of prostate cancer. Quercetin works as anti-inflammatory, antioxidant, and polar auxin transport inhibitor. Furthermore, in vivo study conducted on rats confirmed that its administration can increase energy expenditure (Amić et al., 2007a).

MECHANISTIC APPROACH OF BLACK TEA

The emphasis of phytochemical research has shifted researcher focus toward their disease preventive behavior by executing their functionality against lipid oxidation, nucleic acids, or proteins degeneration (Bai et al., 2007). Thereby, phytochemical pharmacokinetics work well in case of several chronic diseases like cancer, whereas more pronounced effects were observed in cardiovascular compliances, cataract, age-related macular degeneration, and aging (Cirico and Omaye, 2006). Although human cells are enriched with variety of endogenous antioxidants but still require some exogenous antioxidants to maintain an appropriate level of antioxidants to quench the reactive oxidative species (Maeta et al., 2007). Antioxidants protect the body by neutralizing the free radicals via donating electrons and thus ending the scavenger property (Amić et al., 2007b; Maeta et al., 2007). Black tea is considered as a dietary source of antioxidant nutrients like TFs, TRs, quercetin, myricetin, and kaempferol (Table 1). Black tea is a proven nutraceutical agent through inhibition of redox sensitive transcription factors and some "prooxidant" enzymes, like nitric oxide synthase, lipoxigenases, cyclooxygenases, and xanthine oxidase. Moreover, it can induce certain antioxidant enzymes mainly glutathione-S-transferases and superoxide dismutases (Amić and Lučić, 2010). Tea polyphenols with vicinal hydroxyl groups are also strong chelators of redoxactive transition metals (e.g., iron and copper) (Chlebda et al., 2010) may contribute to their overall antioxidant activity. It has been proposed that the stability of such complexes increases as a function of increasing pH as metals are not so efficient electrons acceptor as they are generally surrounded by negative charges. Studies have demonstrated antioxidant activity of black tea while using oxygen radical absorbance capacity (ORAC) assay and found to possess higher antioxidant activity against peroxy radicals as compared with vegetables like garlic, spinach, and kale (Mukai et al., 2005). Moreover, both

Table 3 Standard electron potential of some prominent antioxidants

| Antioxidants | Standard electron potential (mV) |
|------------------------------|----------------------------------|
| Ascorbate | 280 |
| Alpha tocopherol | 480 |
| Uric acid | 590 |
| Glutathione | 920 |
| (-)-Epigallocatechin gallate | 430 |
| (-)-Epigallocatechin | 430 |
| (-)-Epicatechin | 570 |
| (-)-Epicatechin gallate | 550 |
| Theaflavin | 510 |
| Theaflavin digallate | 540 |

Source: Luczaj and Skrzydlewska (2005).

catechins and TFs exhibited strong antioxidant properties as compared with that of glutathione and tocopherol (Kim et al., 2005). Individual fraction of theaflavin, (TF3) has shown higher antioxidant activity than EGCG, a precursor of TF3 (Kim et al., 2005; Phung et al., 2010) probably due to presence of more hydroxyl (OH) groups. Investigations regarding TF structure have shown that order of reactivity is dependent upon the position of hydroxyl groups within the TF molecules. Similarly, theaflavin TF3 showed higher antioxidant activity followed by TF2 and least is documented in TF1 (Auvichayapat et al., 2008). It has been proved that TF in their gallate form exhibits stronger antioxidant activity as compared with individual TF moiety (Nagao et al., 2009). It has also been reported that in vivo free radical scavenging activity is moderately influenced by the value of standard one-electron potential (Table 3), higher in black tea polyphenols (BTP) as compared with vitamin C and tocopherol (Henning et al., 2011). TF also possess ability to inhibit prooxidative enzymes. Study conducted on human leukemia cells HL-60 indicated that TF digallate efficiently reduced the xanthine oxidase activity, which catalyses oxidation of hypoxanthine and xanthine enzymes and convert into uric acid leading to superoxide radical and hydrogen superoxide formation. Furthermore, radical scavenging activity was proved through reduction of NADPH-cytochrome P-450 reductase activity after application of 2-amino-3-methylimidazo quinoline (Amić et al., 2008) as a reaction substrate (Cavia et al., 2010). Studies on human white blood cell (macrophages) exposed the ability of TFs as a nitric oxide synthase (NOS) inhibitor. Polyphenols of black tea also prevent the action of cancer promoting enzymes like cyclooxygenase COX-2 and 5-, 12-, and 15-lipoxygenase that can actively participate in enzymatic induced lipid peroxidation in human colon mucosa and tumor tissues (Lee et al., 2006; Martín et al., 2010). Strong interaction of polyphenols and transition metals in case of black tea is evidence of its antioxidative activity, forming complex with iron or copper ions and thus prevent free radicals initiation by inhibiting the lipid peroxidation mechanism (Young and Woodside, 2001; Mladěnka et al., 2010). However, iron binding ability of black tea polyphenols can impair the iron absorption in the gastrointestinal tract (Balentine et al., 1997; Mukhtar and Ahmad, 2000). This binding with iron can be

mitigated by enhancing the magnitude of ascorbic acid (Duffy et al., 2001; Widlansky et al., 2005). Another theory, regarding the antioxidative mechanism of black tea, revealed that its antioxidant response might be due to its ability to increase the activity and concentration of antioxidants in the living organism. This has been supported by research study, in which black tea dried leaves supplemented in rat's diets resulted in liver protection (Hodgson et al., 2002). Similarly, other experiments showed the improvement in serum superoxide dismutase (SOD) activity after oral administration of black tea to mice exposed to the strong carcinogen, i.e., 3-methylcolanthrene (Di et al., 2011; Hodgson et al., 2000). Many investigations revealed that black tea consumption causes an increase in plasma antioxidative activity from 20 to 40% depending upon condition and applied assay (Hakim et al., 2003; Rietveld and Wiseman, 2003). Evidence for antioxidative potential of black tea is obvious from attenuation of free radicals interaction with some cordial molecules like lipids, DNA, and proteins. It was proved that TF digallate has stronger inhibitory action against macrophage-mediated (white blood cell) LDL oxidation in comparison with EGCG and gallic acid due to its ability to protect membrane phospholipids oxidation (Maron et al., 2003; Sinisalo et al., 2010). Experiments on rats drinking black tea demonstrated inhibition of lipid peroxidation in their livers (Macedo et al., 2011). Antioxidative properties of black tea are also supported by the fact that resultant beverage as well as unoxidized catechins is able to inhibit changes in natural porosity of cell membranes resulting from oxidative stress. Moreover, black tea inhibition of cytochrome P-450 1A1 activity reduces the activation of benzopyrene (BaP), procarcinogen that participates in DNA damage (Roy et al., 2010). Similarly, black tea administration in rats for 10-day as gavages significantly decreased 8-OHdG generation (8-Oxo-2'-deoxyguanosine; a marker of DNA damage) induced by the colon carcinogen, 1, 2-dimethyl hydrazine (Huange et al., 2010; Miranda et al., 2008). However, remarkable decrease was found in the level of 8-OHdG in blood profile of smokers addicted to black tea. Experiments on RAW 264.7 mice macrophages revealed that TFs, especially TF3 effectively prevents activation of transcription factor (NF κ B) thus avoiding expression of nitric oxide synthase (iNOS) (Yang et al., 2009). Experiment conducted on prokaryotes macrophages (106 cells/well) disclosed that black tea TFs and catechins abrogate NO production up to 70%. Incubation of 21BES cells with TF digallate (TF3) showed inhibited c-jun protein phosphorylation considered responsible for the binding of transcription factor AP-1 that plays a vital role in transformation and proliferation of cells (Shen et al., 2010).

RECENT INNOVATION IN ANTIOXIDANT ACTIVITY OF BLACK TEA

In a recent study, antioxidant role of black tea polyphenols and bovine lactoferrin (bLF) was determined using the hamster buccal pouch (HBP) to evaluate the inhibitory effects of both substances on the onset of 7, 12-dimethylbenz anthracene

(DMBA)-induced cancer insurgence. The chemopreventive role of beta lactoferron and black tea polyphenols was accredited to their ability to modulate DMBA-induced DNA damage as well as the performance on different carcinoma proteins (von Staszewski et al., 2010). Moreover, a comparative trial was conducted to demonstrate the effects of consuming a high-flavonoid (HF) and low-flavonoid diet on oxidative damage in 32 healthy humans for testing the lipid peroxidation biomarkers (plasma F2-isoprostanes). Conclusion of study stressed upon the strong *ex vivo* activity of flavonoids as compared with *in vivo* activity mainly due to their low bioavailability (Jayasekera et al., 2011).

Likewise, role of black tea polyphenols in reducing oxidative stress was tested in a rat model of hepatocellular carcinoma (HCC). Administrations of polyphenol-B effectively suppressed hepatocarcinogenesis, as shown by reduced preneoplastic and neoplastic lesions, modulation of xenobiotic-metabolizing enzymes, and amelioration of oxidative stress (Adhami et al., 2004; Martin et al., 2009). Scantly, it can be concluded that polyphenons acts as an effective chain breaker and act as chemopreventive agent *in vivo* (Murugan et al., 2009). Moreover, Widlansky et al. (2005) evaluated the effect of black tea polyphenols on oxidation, inflammation, and antioxidant protection from 66 subjects after administration of two dosages as 450 ml of black tea (acute) and 900 ml of black tea per day (chronic). Results showed that it did not improve plasma antioxidant capacity, suggesting that the benefits of black tea consumption on endothelial function may not be attributable to tea dose (Ziyin et al., 2009).

PROOXIDANT ACTIVITIES

The field of prooxidant has employed the contradict attitude of black tea polyphenols with its antioxidative characteristics. Although antioxidative properties of black polyphenols have well documented, some opposing results also been reported proposing the prooxidative properties of tea. Recently, it has been reported that administration of TF on the human bronchial epithelial (21BES) cell lines, enhanced generation of hydrogen peroxide, and lead to apoptosis after 8 to 12 hours (Haslam, 2003; Lambert and Yang, 2003). Copper, which occurs in the mammalian cell nucleus, is believed to play a central role in the formation of oxygen species and produce DNA damage. Possible mechanism is copper ion (II) activated catechins autooxidation, subsequently generating copper ion (I) and semiquinone radical (Xianglin et al., 2000). This copper ion (I) combines with oxygen to produce hydrogen peroxide and forms bond with DNA to make DNA-Cu(I)OOH that might be responsible for the formation of hydroxyl radical resulting in DNA damage. Moreover, effect of copper ions (II) was also reported to catalyze the oxidation of other active black tea polyphenols, such as quercetin and myricetin, caused break up of DNA strand and formation of 8-OHdG complex (Agarwal et al., 1991; Feng et al., 2002; Leung et al., 2001). Moreover, retrospective data have illuminated that oxidized tea polyphenols, such as semiquinone

or benzoquinone radical undergo reduction that is classified as nonenzymatic through NADH, resulting in abundant redox substances formation that will yield reactive oxygen species. Coenzyme mediated (NADH) free radicals could also facilitate the catechins initiated DNA injuries. Prolonged in vivo administration of quercetin (Langley, 2000) leads to dysfunction of important antioxidant enzyme, such as glutathione, causes oxidative damages due to decrease glutathione reeducate activity and glutathione concentration. Thus, it could be said that antioxidant and prooxidant activity of black tea polyphenols is not consistent, but it dependent upon many factors like reducing potential, chelating ability, pH, solubility, bioavailability, and more important stability in tissues and cells (Meng et al., 2001).

CONCLUSIONS

Current scenario has established black tea as an effective nutraceutical product due to enriched polyphenols profile, an ultimate antioxidative entity. Although the cutting edge between functional and nutraceutical has not yet been illuminated, tea flavanols have enclosed features of an effective disease curative constituents. TF is one of the promising agents against cancer cells while TRs also contribute strongly toward the menace. Alongside, theanine and flavonols have affirmed significance and attained consumers' attention due to their minimal toxicity, high consumption safety, and health assuring perspectives. However, black tea polyphenols are also known to exhibit prooxidative properties in certain conditions. The manuscript artwork provides sufficient evidences to support black tea against various physiological threats. Furthermore, there is still a need of extensive exploration, corroboration, and refinement of respective nutraceuticals to address the remaining pitfalls and challenges in a novel way.

REFERENCES

- Adhami, V. M., Siddiqui, I. A., Ahmad, N., Gupta, S. and Mukhtar, H. (2004). Oral consumption of green tea polyphenols inhibits insulin-like growth factor-I-induced signaling in an autochthonous mouse model of prostate cancer. *Cancer Res.* **64**:8175–8122.
- Agarwal, R. and Mukhtar, H. (1993). Oxidative stress in skin chemical carcinogenesis. In: *Oxidative Stress in Dermatology*, pp. 207–241. Fuchs, J. and Packer, L., Eds., Marcel Dekker, New York.
- Almajano, M. P., Carboa, R., Jimenez, J. A. L. and Michael, H. (2008). Antioxidant and antimicrobial activities of tea infusion Gordon. *Food Chem.* **108**:55–63.
- Amić, D., Davidović-Amić, D., Bešlo, D., Rastija, V., Lučić, B. and Trinajstić, N. (2007a). SAR and QSAR of the antioxidant activity of flavonoids. *Curr. Med. Chem.* **14**:827–845.
- Amić, D., Lučić, B., de Lange, D. W., Verhoef, S., Gorter, G., Kraaijenhagen, R. J., van de Wiel, A. and Akkerman, J. W. (2007b). Polyphenolic grape extract inhibits platelet activation through PECAM-1: An explanation for the French paradox. *Alcohol Clin. Exp. Res.* **31**:1308–1314.
- Amić, D. and Lučić, B. (2010). Reliability of bond dissociation enthalpy calculated by the PM6 method and experimental TEAC values in antiradical QSAR of flavonoids. *Bioorg. Med. Chem.* **18**:28–35.
- Auvichayapat, P., Prapochanung, M., Tunkamnerdthai, O., Sripanidkulchai, B. O., Auvichayapat, N., Thinkhamrop, B., Kunhasura, S., Wongpratoom, S., Sinawat, S. and Hongprapas, P. (2008). Effectiveness of green tea on weight reduction in obese Thais: A randomized, controlled trial. *Physiol. Behav.* **93**:486–491.
- Bai, H. W., Shim, J. Y., Yu, J. and Zhu, B. T. (2007). Biochemical and molecular modeling studies of the O-methylation of various endogenous and exogenous catechol substrates catalyzed by recombinant human soluble and membrane-bound catechol-O-methyltransferases. *Chem. Res. Toxicol.* **20**:1409–1425.
- Balentine, D. A., Wiseman, S. A. and Bouwens, L. C. M. (1997). The chemistry of tea flavonoids. *Crit. Rev. Food Sci. Nutr.* **37**(8):693–704.
- Baor, Y., Teemu, H., Olli, R., Keith, P. and Heikki, K. (2009). Flavonol glycosides in wild and cultivated berries of three major subspecies of *Hippophaë rhamnoides* and changes during harvesting period. *Food Chem.* **115**:657–664.
- Boris, K., Nir, G., Amos, D. and Korczyn, H. (2009). Smoking and tea consumption delay onset of Parkinson's disease. *Parkinsonism and Related Disorders* **15**: 41–46 (E). 061.
- Bursill, C. A., Abbey, M. and Roach, P. D. (2007). A green tea extract lowers plasma cholesterol by inhibiting cholesterol synthesis and upregulating the LDL receptor in the cholesterol-fed rabbit. *Atherosclerosis.* **193**:83.
- Cabrera, C., Giménez, R. and López, M. C. (2003). Determination of tea components with antioxidant activity. *J. Agric. Food Chem.* **51**:4427–4435.
- Cao, G., Sofic, E. and Prior, R. (1996). Antioxidant capacity of tea and common vegetables. *J. Agri. Food Chem.* **44**:3426–3431.
- Cavia, S. M., Busto, M. D., Pilar-Izquierdo, M. C., Ortega, N., Perez-Mateos, M. and Muñoz, P. (2010). Antioxidant properties, radical scavenging activity and biomolecule protection capacity of flavonoid naringenin and its glycoside naringin: A comparative study. *J. Sci. Food Agric.* **90**:1238–1244.
- Chlebda, E., Magdalan, J., Merwid-Lad, A., Trocha, M. and Kopacz, M. (2010). Influence of watersoluble flavonoids, quercetin-5'-sulfonic acid sodium salt and morin-5'-sulfonic acid sodium salt, on antioxidant parameters in the sub acute cadmium intoxication mouse model. *Exp. Toxicol. Pathol.* **62**:105–108.
- Chow, H. H., Hakim, I. A., Vining, D. R., Crowell, J. A., Tome, M. E., Ranger-Moore, J., Cordova, C. A., Mikhael, D. M., Briehl, M. M. and Alberts, D. S. (2007). Modulation of human glutathione s-transferases by polyphenone intervention. *Cancer Epidemiol Biomarkers Prev.* **16**:1662–1666.
- Cirico, T. L. and Omaye, S. T. (2006). Additive or synergetic effects of phenolic compounds on human low density lipoprotein oxidation. *Food Chem. Toxicol.* **44**:510–516.
- Das, S. A., Mukherjee, M., Das, D. and Mitra, C. (2009). Protective action of aqueous black tea (*Camellia sinensis*) extract (BTE) against ovariectomy-induced oxidative stress of mononuclear cells and its associated progression of bone loss. *Phytother. Res.* **23**(9):1287–1294.
- De Mejia, E. G., Ramirez-Mares, M. V. and Puangpraphant, S. (2009). Bioactive components of tea: Cancer, inflammation and behavior. *Brain, Behavior, and Immunity.* **23**:721–731.
- Di, W., Ying, Z., Xiao, L., Shuang, W., Rong, X., Wei, B., Wei, Y., Hong, Y., Ping, Y. and Liegang, L. (2011). Pu-erh black tea supplementation decreases quinocetone-induced ROS generation and oxidative DNA damage in Balb/c mice. *Food Chem. Toxicol.* **49**:477–484.
- Dimpfel, W., Kler, A., Kriesl, E. and Lehnfeld, R. (2007). Theogallin and L-theanine as active ingredients in decaffeinated green tea extract: II. Characterization in the freely moving rat by means of quantitative field potential analysis. *J. Pharm. Pharmacol.* **59**:1397–1403.
- Duffy, S. J., Keaney Jr., J. F., Holbrook, M., Gokce, N., Swerdloff, P. L., Frei, B. and Vita, J. A. (2001). Short- and long-term black tea consumption reverses endothelial dysfunction in patients with coronary artery disease. *Circulation.* **104**:151–156.
- Dufresne, C. J. and Farnworth, E. R. (2001). A review of latest research findings on the health promotion properties of tea. *J. Nutr. Biochem.* **12**:404–412.
- Erba, D., Riso, P., Foti, P., Frigerio, F., Crisculi, F. and Testolin, G. (2003). Black tea extract supplementation decreases oxidative damage in Jurkat T cells. *Arch. Biochem. Biophys.* **416**:196–201.
- Feng, Q., Torii, Y., Uchida, K., Nakamura, Y., Hara, Y. and Osawa, T. (2002). Black tea polyphenols, theaflavins, prevent cellular DNA damage by

- inhibiting oxidative stress and suppressing cytochrome P450 1A1 in cell cultures. *J. Agric. Food Chem.* **50**:213–220.
- Flint, B. M. and Balz, F. (1999). Black tea increases the resistance of human plasma to lipid peroxidation in vitro, but not ex vivo. *Antonie van Leeuwenhoek J. Microbiol. Biotechnol.* **27**(3/4):381–387.
- Gardner, E. J., Ruxton, C. H. and Leeds, A. R. (2007). Black tea—helpful or harmful? A review of the evidence. *Eur. J. Clin. Nutr.* **61**:3–18.
- Gupta, J., Siddique, Y. H., Beg, T., Ara, G. and Afzal, M. (2009). Protective role of green tea extract against genotoxic damage induced by anabolic steroids in cultured human lymphocytes. *Biol. Med.* **1**(2):87–99.
- Hakim, I. A., Harris, R. B., Brown, S., Chow, H. H., Wiseman, S., Agarwal, S. and Talbot, W. (2003). Effect of increased tea consumption on oxidative DNA damage among smokers: A randomized controlled study. *J. Nutr.* **133**:3303S–3309.
- Halliwel, B. (2007). Dietary polyphenols: Good, bad, or indifferent for your health? *Cardiovasc. Res.* **73**:341–347.
- Haslam, E. (2003). Thoughts on thearubigins. *Phytochemistry*. **64**:61–73.
- Henning, S. M., Niu, Y., Lee, N. H., Thames, G. D., Minutti, R. R., Wang, H., Go, V. L. and Heber, D. (2004). Bioavailability and antioxidant activity of tea flavanols after consumption of green tea, black tea, or a green tea extract supplement. *Am. J. Clin. Nutr.* **80**(6):1558–1564.
- Henning, S. M., Wang, P. and Heber, D. (2011). Chemopreventive effects of tea in prostate cancer: Green tea versus black tea. *Mol. Nutr. Food Res.* **55**:1–16.
- Hideyuki, S., Shunsuke, I., Nobukazu, M. and Hidehiko, K. (2002). Enzymatic production of theanine, an “umami” component of tea, from glutamine and ethylamine with bacterial γ -glutamyl trans peptidase. *Enzyme Microb. Tech.* **31**:884–889.
- Hodgson, J. M., Puddey, I. B., Burke, V., Watts, G. F. and Beilin, L. J. (2002). Regular ingestion of black tea improves brachial artery vasodilator function. *Clin. Sci (Lond.)*. **102**:195–201.
- Hodgson, J. M., Puddey, I. B. and Croft, K. D. (2000). Acute effects of ingestion of black and green tea on lipoprotein oxidation. *Am. J. Clin. Nutr.* **71**:1103–1107.
- Hsu, A., Bruno, R. S., Löhr, C. V., Taylor, A. W., Dashwood, R. H., Bray, T. M. and Ho, E. (2011). Dietary soy and tea mitigate chronic inflammation and prostate cancer via NF κ B pathway in the Noble rat model. *J. Nutr. Biochem.* **22**:502–510.
- Huang, X. J., Zhang, H. H., Wang, X., Huang, L. L., Zhang, L. Y., Yan, C. X., Liu, Y. and Yuan, Z. H. (2010). ROS mediated cytotoxicity of porcine adrenocortical cells induced by QdNOs derivatives in vitro. *Chem. Biol. Interact.* **185**:227–234.
- Imran, A., Butt, M. S. and Sharif, M. K. (2012). Phytochemical Density of Some Promising Commercial Tea Brands. *International Journal of Food Properties* **15**:199–108.
- Inami, S., Takano, M., Yamamoto, M., Murakami, D., Tajika, K., Yodogawa, K., Yokoyama, S., Ohno, N., Ohba, T., Sano, J., Ibuki, C., Seino, Y. and Mizuno, K. (2007). Tea catechin consumption reduces circulating oxidized low-density lipoprotein. *Int. Heart J.* **48**:725–732.
- Jayasekera, S., Molan, A. L., Garg, M. and Moughan, P. J. (2011). Variation in antioxidant potential and total polyphenol content of fresh and fully-fermented Sri Lankan tea. *Food Chem.* **125**(2):536–541.
- Jovanovic, S. V., Hara, Y., Steenken, S. and Simic, M. G. (1997). Antioxidant potential of gallic catechins. A pulse radiolysis and laser photolysis study. *J. Am. Chem. Soc.* **119**:5337–5343.
- Ju, J., Lu, G., Lambert, J. D. and Yang, C. S. (2007). Inhibition of carcinogenesis by tea constituents. *Semin. Cancer Biol.* **17**:395–402.
- Katiyar, S. K. (2010). *Arch. Biochem. Biophys.*, In Press, Corrected Proof, Available online
- Kaur, S., Greaves, P., Cooke, D. N., Edwards, R., Steward, W. P., Gescher, A. J. and Marczylo, T. H. (2007). Breast cancer prevention by green tea catechins and black tea theaflavins in the C3(1) SV40 T,t antigen transgenic mouse model is accompanied by increased apoptosis and a decrease in oxidative DNA adducts. *J. Agric. Food Chem.* **55**:3378–3385.
- Kavirasan, K., Kalaivarsi, P. and Pugalandi, V. (2008). Antioxidant efficacy of flavonoid-rich fraction from *Spermocoe hispida* in hyperlipidemic rats. *J. Appl. Biomed.* **6**:165–176.
- Kim, B. H., Cho, S. M., Reddy, A. M., Kim, Y. S., Min, K. R., and Kim, Y. (2005). Down-regulatory effect of quercetin gallate on nuclear factor kappaB-dependent inducible nitric oxide synthase expression in lipopolysaccharide-stimulated macrophages RAW 264.7. *Biochem. Pharmacol.* **69**:1577–1583.
- Kumar, N., Shibata, D., Helm, J., Coppola, D. and Malafa, M. (2007). Green tea polyphenols in the prevention of colon cancer. *Front Biosci.* **12**:2309–2315.
- Lachman, J., Sulc, M. and Schilla, M. (2007). Comparison of the total antioxidant status of Bohemian wines during the wine-making process. *Food Chem.* **103**:802–807.
- Lambert, J. D., Sang, S. and Yang, C. S. (2007a). Biotransformation of green tea polyphenols and the biological activities of those metabolites. *Mol. Pharm.* **4**:819–825.
- Lambert, J. D., Sang, S. M. and Yang, C. S. (2007b). Possible controversy over dietary polyphenols: Benefits vs risks. *Chemical Research in Toxicology*. **20**:583–585.
- Lambert, J. D. and Yang, C. S. (2003). Cancer chemopreventive activity and bioavailability of tea and tea polyphenols. *Mutat Res.* **523–524**:201–208.
- Langley, S. C. (2000). Consumption of black tea elicits an increase in plasma antioxidant potential in humans. *Int. J. Food Sci. Nutr.* **51**:309–315.
- Lee-Hilz, Y. Y., Boerboom, A. M., Westphal, A. H., Berkel, W. J., Aarts, J. M. and Rietjens, I. M. (2006). Pro-oxidant activity of flavonoids induces EpRE-mediated gene expression. *Chem. Res. Toxicol.* **19**:1499–1505.
- Leenen, R., Roodenburg, A. J., Tijburg, L. B. and Wiseman, S. A. (2000). A single dose of tea with or without milk increases plasma antioxidant activity in humans. *Eur. J. Clin. Nutr.* **54**:87–92.
- Letchoumy, P. V., Mohan, K. V., Stegeman, J. J., Gelboin, H. V., Hara, Y. and Nagini, S. (2008). In vitro antioxidative potential of lactoferrin and black tea polyphenols and protective effects in vivo on carcinogen activation, DNA damage, proliferation, invasion, and angiogenesis during experimental oral carcinogenesis. *Oncol. Res.* **17**(5):193–203.
- Leung, L. K., Su, Y., Chen, R., Zhang, Z., Huang, Y. and Chen, Z. Y. (2001). Theaflavins in black tea and catechins in green tea are equally effective antioxidants. *J. Nutr.* **131**:2248–2251.
- Luczaj, W. and Skrzydlewska, E. (2005). Antioxidative properties of black tea. *Preventive Medicine*. **40**:910–918.
- Macedo, J. A., Battistin, V., Ribeiro, M. L. and Macedo, G. A. (2011). Increasing the antioxidant power of tea extracts by biotransformation of polyphenol. *Food Chemistry*. **126** 491–497.
- Maeta, K., Nomur, W., Takashume, Y., Izawa, S. and Inoue, Y. (2007). Green tea polyphenols function as prooxidants to activate oxidative-stress-responsive transcription factors in yeasts. *Appl. Environ. Microb.* **73**:572–580.
- Maeta, K., Nomura, W., Takatsume, Y., Izawa, S. and Inoue, Y. (2007). Green tea polyphenols function as prooxidants to activate oxidative-stress-responsive transcription factors in yeasts. *Appl. Environ. Microbiol.* **73**:572–580.
- Maki, K. C., Reeves, M. S., Farmer, M., Yasunaga, K., Matsuo, N., Katsuragi, Y., Komikado, M., Tokimitsu, I., Wilder, D., Jones, F., Blumberg, J. B. and Cartwright, Y. (2009). Green tea catechin consumption enhances exercise-induced abdominal fat loss in overweight and obese adults. *J. Nutr.* **139**:264–270.
- Maron, D. J., Lu, G. P., Gai, N. S., Wu, Z. G., Li, Y. H., Chen, H., Zhu, J. Q., Jin, X. J., Wouters, B. C. and Zhao, J. (2003). Cholesterol lowering effect of a theaflavin-enriched green tea extract: A randomised controlled trial. *Arch. Intern. Med.* **163**:1448–1453.
- Martín, M. Á., Serrano, A. B. G., Ramos, S., Pulido, M. I., Bravo, L. and Goya, L. (2010). Cocoa flavonoids up-regulate antioxidant enzyme activity via the ERK1/2 pathway to protect against oxidative stress-induced apoptosis in HepG2 cells. *J. Nutr. Biochem.* **21**:196–205.
- Martins, F., Suzan, A. J., Cerutti, S. M., Arçari, D. P., Ribeiro, L. M., Bastos, D. H. M. and Carvalho Pde O. (2009). Consumption of mate tea (*Ilex paraguariensis*) decreases the oxidation of unsaturated fatty acids in mouse liver. *Br. J. Nutr.* **101**:527–532.
- Mazzanti, G., Menniti-Ippolito, F., Moro, P. A., Cassetti, F., Raschetti, R., Santuccio, C. and Mastrangelo, S. (2009). Hepatotoxicity from green tea: A

- review of the literature and two unpublished cases. *Eur. J. Clin. Pharmacol.* **65**:331–341.
- Mei, X., Lin, D. H., Xu, Y., Wu, Y. Y. and Tu, Y. Y. (2009). Effects of phenanthrene on chemical composition and enzyme activity in fresh tea leaves. *Food Chem.* **115**(2):569, 573.
- Mei, X., Wu, Y.-Y., Mao, X. and Tu, Y.-Y. (2011). Antagonism of phenanthrene cytotoxicity for human embryo lung fibroblast cell line HFL-I by green tea polyphenol. *Environ. Pollut.* **159**:164–168.
- Meng, J., Ren, B., Xu, Y., Kamendulis, L. M., Dum, N. and Klaunig, J. E. (2001). Reduction of oxidative DNA damage (comet assay) in white blood-cells by black tea consumption in smokers and non-smokers. *Toxicol. Sci.* **60**: 411–412.
- Miranda, D. D. C., Arçari, D. P., Pedrazzoli J. Jr., Carvalho, P. O., Cerutti, S. M., Bastos, D. H. M. and Ribeiro, M. L. (2008). Protective effects of mate tea (*Ilex paraguariensis*) on H₂O₂-induced DNA damage and DNA repair in mice. *Mutagenesis*. **23**:261–265.
- Mladěnká, P., Zatloukalová, L., Filipický, T. and Hrdina, R. (2010). Cardiovascular effects of flavonoids are not caused only by direct antioxidant activity. *Free Radic. Biol. Med.* **15**:963–975.
- Montserrat, D., Ana, G.-P. and Celestino, S.-B. (2010). Method and experimental TEAC values in antiradical QSAR of flavonoids. *Bioorg. Med. Chem.* **18**:28–35.
- Mukai, K., Mitani, O. K. and Nagaoka, S. I. (2005). Structure-activity relationship of tocopherol regeneration reaction by catechin. *Free Radic. Biol. Med.* **38**:1243–1256.
- Mukhtar, H. and Ahmad, N. (2000). Tea polyphenols: Prevention of cancer and optimizing health. *Am. J. Clin. Nutr.* **71**:1698S–1702S (review).
- Murugan, R. S., Vinothini, G., Hara, Y. and Nagini, S. (2009). Black tea polyphenols target matrix metalloproteinases, RECK, proangiogenic molecules and histone deacetylase in a rat hepatocarcinogenesis model. *Anticancer Res.* **29**(6):2301–2305.
- Nagao, T., Meguro, S., Hase, T., Otsuka, K., Komikado, M., Tokimitsu, I., Yamamoto, T. and Yamamoto, K. A. (2009). Catechin-rich beverage improves obesity and blood glucose control in patients with type 2 diabetes. *Obesity (Silver Spring)*. **17**:310–317.
- Noelia, C.-M. A., Sergio, G.-A., Esteban, G.-R. and Isidro, H.-G. (2010). Flavonol profiles of *Vitis vinifera* white grape cultivars. *J. Food Comp. Anal.* **23**:699–705.
- Nurk, E., Refsum, H., Drevon, C. A., Tell, G. S., Nygaard, H. A., Engedal, K., Smith, A. D. (2009). Intake of flavonoid-rich wine, tea, and chocolate by elderly men and women is associated with better cognitive test performance. *J. Nutr.* **139**(1):120–127.
- Ou, B., Huang, D., Hampsch-Woodill, M., Flanagan, J. A. and Deemer, E. K. (2002). Analysis of antioxidant activities of common vegetables employing oxygen radical absorbance capacity (ORAC) and ferric reducing antioxidant power (FRAP) assays: A comparative study. *J. Agric. Food Chem.* **50**:3122–3128.
- Palva, S. and Palva, J. M. (2007). New vistas for alpha-frequency band oscillations. *Trends Neurosci.* **30**:150–158.
- Patel, R., Garg, R., Erande, S. and Maru, G. B. (2007). Chemopreventive herbal anti-oxidants: Current status and future perspectives. *J. Clin. Biochem. Nutr.* **40**:82–91.
- Patel, R. and Maru, G. (2008). Polymeric black tea polyphenols induce phase II enzymes via Nrf2 in mouse liver and lungs. *Free Radic. Biol. Med.* **44**:1897–1911.
- Phung, O. J., Baker, W. L., Matthews, L. J., Lanosa, M., Thorne, A. and Coleman, C. I. (2010). Effect of green tea catechins with or without caffeine on anthropometric measures: A systematic review and meta-analysis. *Am. J. Clin. Nutr.* **91**:73–81.
- Platt, K. L., Aderhold, S., Kulpe, K. and Fickler, M. (2008). Unexpected DNA damage caused by polycyclic aromatic hydrocarbons under standard laboratory conditions. *Mutat. Res.* **650**(2):96–103.
- Popkin, B. M. (2010). Recent dynamics suggest selected countries catching up to US obesity. *Am. J. Clin. Nutr.* **91**:1284S–1288S.
- Procházková, D., Boušová, I. and Wilhelmová, N. (2011). Antioxidant and prooxidant properties of flavonoids D. *Fitoterapia* **82**(4):513–523.
- Rains, T. M., Agarwal, S. and Maki, K. C. (2011). Antiobesity effects of green tea catechins: A mechanistic review. *J. Nutr. Biochem.* **22**:1–7.
- Rietveld, A. and Wiseman, S. (2003). Antioxidant effects of tea: Evidence from human clinical trials. *J. Nutr.* **133**:3285S–3292S.
- Ross, S., Ziska, D. S., Zhao, K. and ElSohly, M. A. (2000). Variance of common flavonoids by brand of grapefruit juice. *Fitoterapia*. **71**:154.
- Roy, M. K., Koide, M., Rao, T. P., Okubo, T., Ogasawara, W. and Juneja, L. R. (2010). ORAC and DPPH assay comparison to assess antioxidant capacity of tea infusions: Relationship between total polyphenol and individual catechins content. *Int. J. Food Sci. Nutr.* **61**:109–124.
- Schneider, C. and Segre, T. (2009). Green tea: Potential health benefits. *Am. Fam Physician*. **79**:591–594.
- Sharangi, A. B. (2009). Medicinal and therapeutic potentialities of tea (*Camellia sinensis* L.)—a review. *Food Res. Int.* **42**:529–535.
- Shen, J., Yang, C., Wu, C., Feng, P., Wang, Z., Li, Y., Li, Y. and Zhang, S. (2010). Identification of the major metabolites of quinocetone in swine urine using ultra-performance liquid chromatography/electrospray ionization quadrupole time-of-flight tandem mass spectrometry. *Rapid Commun. Mass Spectrom.* **24**:375–383.
- Shiraki, M., Hara, Y., Osawa, T., Kumon, H., Nakayama, T. and Kawakishi, S. (1994). Antioxidative and antimutagenic effects of theaflavins from black tea. *Mutat. Res.* **323**:29–34.
- Siddiqui, I. A., Raisuddin, S. and Shukla, Y. (2005). Protective effects of black tea extract on testosterone induced oxidative damage in prostate. *Cancer Let.* **227**:125–132.
- Sinisalo, M., Enkovaara, A.-L. and Kivisto, K. T. (2010). Possible hepatotoxic effect of rooibos tea: A case report. *Eur. J. Clin. Pharmacol.* **66**:427–428.
- Smith, G., Thomsen, S. J., Markham, K. R., Andary, C. and Cardon, D. (2000). The photostabilities of naturally occurring 5-hydroxyflavones, flavonols, their glycosides and their aluminium complexes. *J. Photochem. Photobiol. A*. **136**:87.
- Sugiyama, T. and Sadzuka, Y. (2003). Theanine and glutamate transporter inhibitors enhance the antitumor efficacy of chemotherapeutic agents. *Biochim. Biophys. Acta*. **1653**:47–59.
- Sun, C., Fu, J., Chen, J., Jiang, L. and Pan, Y. (2010). On-line HPLC method for screening of antioxidants against superoxide anion radical from complex mixtures. *J. Sep. Sci.* **33**:1018–1023.
- Suzuki, H. and Kumagaya, H. (2001). c-Glutamyltranspeptidase. *Protein, Nucleic Acid and Enzyme*. **46**:1842–1848.
- Takayama, T., Nakajima, T., Ohi, M., Miyanishi, K., Nobuoka, A., Hayashi, T., Henderson, C. J., Wolf, R. C. and Niitsu, Y. (2001). GST-pi as a cytoprotecting factor in colon carcinogenesis. *Proc. Am. Assoc. Cancer Res.* **42**: 2983.
- Thielecke, F. and Boschmann, M. (2009). The potential role of green tea catechins in the prevention of the metabolic syndrome—a review. *Phytochemistry*. **70**:11–24.
- Turkmen, N., Sari, F. and Velioglu, Y. S. (2006). Effects of extraction solvents on concentration and antioxidant activity of black and black mate tea polyphenols determined by ferrous tartrate and Folin-Ciocalteu methods. *Food Chem.* **99**:835–841.
- Van der Pijl, P., Chenb, L. and Muldera, T. P. J. (2010). Human disposition of L-theanine in tea or aqueous solution. *J. Funct. Foods*. **2**:239–244.
- Van het Hof, K. H., de Boer, H. S., Wiseman, S. A., Lien, N., Westrate, J. A. and Tijburg, L. B. (1997). Consumption of green or black tea does not increase resistance of low-density lipoprotein to oxidation in humans. *Am. J. Clin. Nutr.* **66**:1125–1132.
- von Staszewski, M., Pilosof, A. M. R. and Jagus, R. J. (2010). Antioxidant and antimicrobial performance of different Argentinean green tea Food varieties as affected by whey proteins. *Anticancer Res.* **29**(6):2301–2305.
- Wang, D., Xiao, R., Hu, X., Xu, K., Hou, Y., Zhong, Y., Meng, J., Fan, B. and Liu, L. (2010). Comparative safety evaluation of Chinese Pu-erh green

- tea extract and Pu-erh black tea extract in Wistar rats. *J. Agric. Food Chem.* **58**:1350–1358.
- Wang, H. and Helliwell, K. (2001). Determination of flavonols in green and black tea leaves and green tea infusions by high-performance liquid chromatography. *Food Res. Int.* **34**:223–227.
- Warden, B. A., Smith, L. S., Beecher, G. R., Balentine, D. A. and Clevidence, B. A. (2001). Catechins are bioavailable in men and women drinking black tea throughout the day. *J. Nutr.* **131**:1731–1737.
- Widlansky, M. E., Duffy, S. J., Hamburg, N. M., Gokce, N., Warden, B. A., Wiseman, S., Keaney, J. F., Jr, Frei, B. and Vita, J. A. (2005). Effects of black tea consumption on plasma catechins and markers of oxidative stress and inflammation in patients with coronary artery disease. *Free Radic. Biol. Med.* **38**:499–506.
- Xianglin, S., Ye, J., Leonard, S., et al. (2000). Antioxidant properties of (–)epicatechin-3-gallate and its inhibition of Cr(VI)-induced DNA damage and Cr(IV)- or TPA-stimulated NF- κ B activation. *Mol. Cell Biochem.* **206**:125–132.
- Yang, C. S., Wang, X., Lu, G. and Picinich, S. C. (2009). Cancer prevention by tea: Animal studies, molecular mechanisms and human relevance. *Nat. Rev. Cancer*. **9**(6):429–439.
- Young, I. S. and Woodside, J. V. (2001). Antioxidants in health and disease. *J. Clin. Pathol.* **54**:176–186.
- Ziyin, Y., Youying, Tu, Susanne, B., Fang, D., Yi, Xuc. and Naoharu, W. (2009). Isolation and identification of compounds from the ethanolic extract of flowers of the tea (*Camellia sinensis*) plant and their contribution to the antioxidant capacity. *LWT-Food Sci. Technol.* **42**:1439–1443.