



## The Health Potential of Fruits and Vegetables Phytochemicals: Notable Examples

Arantxa Rodriguez-Casado

To cite this article: Arantxa Rodriguez-Casado (2016) The Health Potential of Fruits and Vegetables Phytochemicals: Notable Examples, Critical Reviews in Food Science and Nutrition, 56:7, 1097-1107, DOI: [10.1080/10408398.2012.755149](https://doi.org/10.1080/10408398.2012.755149)

To link to this article: <https://doi.org/10.1080/10408398.2012.755149>



Accepted author version posted online: 16  
Sep 2014.  
Published online: 16 Sep 2014.



Submit your article to this journal [↗](#)



Article views: 1672



View Crossmark data [↗](#)



Citing articles: 34 View citing articles [↗](#)

# The Health Potential of Fruits and Vegetables Phytochemicals: Notable Examples

ARANTXA RODRIGUEZ-CASADO

IMDEA Food Institute, Madrid, Spain

*Fruit and vegetables are essential components of a healthy diet. The World Health Organization (WHO) recommends an intake of five to eight portions (400–600 g) daily of fruits and vegetables to reduce risk of cardiovascular disease, cancer, poor cognitive performance, and other diet-related diseases, as well as for the prevention of micronutrient deficiencies. Much of their potential for disease prevention is thought to be provided by phytochemicals, among which the preventive activity of antioxidants is most well documented. Since numerous meta-studies published indicate variable and often contradictory results about the impact of isolated phytochemicals on health, their consumption as supplements must be carried out with care, because doses may exceed the recommended nutritional intake. Nonetheless, there is a general consensus that whole fruit and vegetable intake is more important in providing health benefits than that of only one of their constituent, because of additive and synergistic effects. This review describes the most recent literature regarding the health benefits of some selected fruits and vegetables. Importantly, since some phytochemicals regulate the same genes and pathways targeted by drugs, diets rich in fruits and vegetables in combination with medical therapies are being considered as novel approaches to treatment. Therefore, phytochemicals in fruits and vegetable might be a promising tool for the prevention and/or amelioration of a wide range of diseases.*

**Keywords** Broccoli, dandelion, garlic, cocoa, olives, grapes

## INTRODUCTION

The onset and progression of certain diseases are clearly influenced by lifestyle factors. Nutrition is one of the most important determinants of health and increasing evidence suggests that diets rich in fruits and vegetables may prevent a wide range of diseases (Traka and Mithen, 2011; Aune et al., 2012a,b; Boeing et al., 2012). Although mechanisms underlying these effects have not been fully elucidated, the presence of secondary metabolites, commonly referred as phytochemicals, in fruits and vegetables might play major roles (Traka and Mithen, 2011; Aune et al., 2012a; Boeing et al., 2012). Many of such phytochemicals possess the ability to interfere with cellular functions, altering activation of transcription factors that regulate expression of genes, and change cellular metabolism in different ways (Tsao, 2010; Traka and Mithen, 2011). Accordingly, the past two decades witnessed increased scientific interest in fruits and vegetables and their putative health-promoting properties (Badimon et al., 2010; Ferrari et al., 2011; Speciale et al., 2011). Many lines of evidence

demonstrate that the majority of phytochemicals exhibit antioxidant (Lipinski, 2011; Speciale et al., 2011; Chiva-Blanch and Visioli, 2012; Giampieri et al., 2012) and anti-inflammatory (Zhao et al., 2011a,b; Singh et al., 2012) activities that confer them pharmacological properties likely important to prevent and maybe even treat (to a limited degree) different diseases (Baldrick et al., 2011). Among the thousands of phytochemicals found in our diets, phenolic compounds stand out as the most important group of natural antioxidants (De Kok et al., 2010; Chiva-Blanch and Visioli, 2012; Giampieri et al., 2012; Visioli, 2012).

Of the major areas of human health, cancer and cardiovascular disease are the best characterized ones. Free radicals, which induce oxidative stress, can cause DNA damage, which in turn can lead to base mutation, DNA cross-linking, and chromosomal breakage and rearrangement (De Kok et al., 2010). This damage may be limited by dietary antioxidants in fruits and vegetables through modulation of detoxification enzymes, scavenging of oxidative agents, stimulation of the immune system, hormone metabolism, and regulation of gene expression in cell proliferation and apoptosis (De Kok et al., 2010; Lipinski, 2011; Niki, 2011; Speciale et al., 2011). Indeed, epidemiologic studies

Address correspondence to Arantxa Rodriguez-Casado, IMDEA Food Institute, Carretera de Cantoblanco 8, 28049 Madrid, Spain. E-mail: arantxa.rodriguez@imdea.org

support an inverse relationship between a regular consumption of fruits and vegetables and the risk of cancers of the colon (Magalhaes et al., 2012), oesophagus (Berretta et al., 2012), oral cavity (Lu et al., 2011), stomach (Steevens et al., 2011), pancreas (Cappellani et al., 2012), prostatic (Sebastiano et al., 2012), breast (Aune et al., 2012a; Aune et al., 2012b), and ovary (Paxton et al., 2012). Cancer is a disease that features a strong genetic component and numerous genes that code for oncogenes or tumors suppressor genes have been identified. Extended treatment with physiologic concentrations of dietary phytochemicals results in altered gene expression, reduced growth, and apoptosis of cancer cells (McCullough and Giovannucci, 2004; Lu et al., 2011; Steevens et al., 2011; Aune et al., 2012a; Aune et al., 2012b; Berretta et al., 2012; Cappellani et al., 2012; Paxton et al., 2012; Sebastiano et al., 2012; Singh et al., 2012). On the other hand, many studies also indicate a strong link between fruit and vegetable intake and a reduced risk for cardiovascular disease (Badimon et al., 2010; Perez-Vizcaino and Duarte, 2010; Finks et al., 2012; Vasanthi et al., 2012). Likewise, the intake of this type of food has been inversely correlated with mortality from coronary disorders as well as incidence of risk factors such as diabetes, hypertension, and hyperlipidemia (Choudhary and Tran, 2011; El Haouari and Rosado, 2011). Oxidized low density lipoprotein (LDL) is considered an atherogenic factor in heart disease, promoting cholesterol ester accumulation and foam cell formation. Dietary antioxidants from fruits and vegetables get incorporated into LDL, and become oxidized themselves, thus preventing oxidation of polyunsaturated fatty acids, and modulate the synthesis and absorption of cholesterol (Choudhary and Tran, 2011; El Haouari and Rosado, 2011). Phytochemicals from fruit and vegetable also reduce platelet aggregation (Vita, 2005; El Haouari and Rosado, 2011), blood pressure (Galleano et al., 2010; Zhao et al., 2011a,b), insulin resistance index, and serum insulin concentration (Omar et al., 2010), and obesity (Calder et al., 2011), all of them important risk factor for cardiopathologies.

Moreover, fruits and vegetables also ameliorate systemic inflammation that may be a critical factor in cardiovascular disease (Calder et al., 2011; Zhao et al., 2011,b; Singh et al., 2012). Cytokines (IL-1, IL-6, TNF- $\alpha$ ), peptide hormones secreted by both inflammatory and adipose stromal cells that mediate the inflammatory response, are signals that stimulate tumor growth and are associated to obesity (Calder et al., 2011).

Finally, diets rich in fruits and vegetable in combination with medical therapies have been considered as novel treatment approaches since some phytochemicals regulate the same genes and pathways targeted by drugs (Shukla and George, 2011).

This review describes the most recent literature regarding the health benefits of some fruits and vegetable and their phytochemicals. Broccoli, for example, contains glucosinolates,

which have been associated with a decreased risk of cancer; dandelion possesses important anti-inflammatory properties; garlic contains allyl sulphides that inhibit cancer cell growth; cocoa intake is being linked with decreased risk of cardiovascular disease; olives might play cardio-protective roles due to their antioxidant and anti-inflammatory properties; and, finally, grapes might be a source of beneficial phytochemicals in Alzheimer's disease and age-related cognitive decline.

Research continues proving the effectiveness of a plant-based diet with regard to weight loss, body fat distribution, cholesterol and glucose levels, blood pressure, and protective against numerous chronic diseases including cancer, cardiovascular, and Alzheimer's diseases (Bach-Faig et al., 2011). The Mediterranean diet is characterized by a dietary pattern that has remained constant over time and space rich in plant foods (cereals, fruits, vegetables, legumes, tree nuts, seeds, and olives), with olive oil as the principal source of added fat, along with high to moderate intakes of fish and seafood, moderate consumption of eggs, poultry and dairy products, low consumption of red meat and a moderate intake of alcohol (Bach-Faig et al., 2011). However, the Mediterranean diet is something else food. It constitutes a set of skills, knowledge, practices, and traditions ranging that makes the Mediterranean diet is acknowledged as an Intangible Cultural Heritage of Humanity by UNESCO (2010) (Bach-Faig et al., 2011).

## HEALTH POTENTIAL OF SELECTED FRUITS AND VEGETABLES

Although multiple factors could contribute to incidence and prevalence of chronic diseases, the contribution of some fruit and vegetables is worthy being considered.

### Broccoli (*Brassica oleracea*)

Broccoli is a cruciferous vegetable that belongs the *Brassicaceae* (formerly called *Cruciferae*) family, together with cauliflower, Brussel's sprouts, kohlrabi, cabbage and mustard (Buck, 1956). The flower heads (commonly called broccoli, broccoli florets, or broccoli heads) are eaten before the flower buds open (Buck, 1956).

Principal components of broccoli are glucosinolates, which are present in all members of the *Brassicaceae* family (Buck, 1956). Glucosinolates may be hydrolyzed to yield a variety of biologically-active products, including isothiocyanates, thio-cyanates, nitriles, and oxazolidine-2-thiones (Buck, 1956; No authors listed, 2010).

These products influence a number of cellular processes through the regulation of transcription factor levels, signaling pathways, the cell cycle, and apoptosis (Buck, 1956; No authors listed, 2010). Broccoli contains high levels of the isothiocyanate sulforaphane and of other glucosinolate

derivatives thought to have beneficial health properties (Zhang et al., 1992; No authors listed, 2010).

Several other secondary plant metabolites have been identified in broccoli, and include a series of flavonoids, 2-caffeoylquinic acid derivatives, and cinnamic acid derivatives (Aires et al., 2011). These compounds are accompanied by ascorbic acid, xanthophylls (lutein, zeaxanthin), carotenoids, tocopherols, proteins, sugars, and chlorophyll (Aires et al., 2011). Broccoli is also rich in minerals such as selenium and calcium, fiber and vitamins as well as various amino acids (Aires et al., 2011). Broccoli has been highly valued due to its flavor, but also because of its beneficial effects on human health (Latté et al., 2011). Sulforaphane, an isothiocyanate biologically active metabolite of glucoraphanin, has been identified in broccoli as the compound responsible for many of their health benefits (Zhang et al., 1992). Glucoraphanin, also referred to as sulforaphane glucosinolate, is the most potent naturally occurring inducer of phase 2 detoxification enzymes (Zhang et al., 1992) and is a long-acting antioxidant (No authors listed, 2010).

Sulforaphane also exhibits broad-spectrum antimicrobial activity against numerous gram-positive and gram-negative bacteria (No authors listed, 2010). In addition, preclinical and clinical studies have concluded that sulforaphane possesses anti-inflammatory activity by inhibition of cytokine production (Juge et al., 2007). Sulforaphane's multiple molecular targets and clinical trials currently underway to assess its effects on various cancers (Kensler et al., 2012) and cardiovascular diseases (Park et al., 2012).

A wide range of studies in humans, animals, and in vitro has confirmed an inverse association between broccoli intake and risk of different cancers (Vasanthi et al., 2009; Kensler et al., 2012). Numerous in vitro studies on human colon (Gamet-Payraastre et al., 2000), leukemia (Fimognari et al., 2002), pancreatic (Pham et al., 2004), lung (Liang et al., 2008), and prostate (Singh et al., 2004) cancers have demonstrated the inhibitory effects of sulforaphane. Intake of broccoli sprouts has been associated with decreased incidence and tumor growth prostate and breast cancer in animal models (No authors listed, 2010). Epidemiological studies suggest that intake of more than one portion of cruciferous vegetables per week lowers both the incidence and the development of aggressive prostate cancer (Traka et al., 2008). Specifically, results suggest that by consuming—during 12 months—a broccoli-rich diet complex changes are produced in signaling pathways associated with inflammation and carcinogenesis (Traka et al., 2008). These changes may be mediated through the chemical interaction of isothiocyanates with signaling peptides in the plasma (Traka et al., 2008; Traka and Mithen, 2011; Dinkova-Kostova and Kostov, 2012). Apart from isothiocyanates and indoles (Riso et al., 2009), minerals (selenium) in broccoli play a critical key role in cancer prevention (Holst and Williamson, 2004).

Numerous clinical studies indicate that broccoli sprouts could have favorable effects on risk factors for cardiovascular disease.

A phase I trial involving 12 subjects consuming 100 g broccoli sprouts daily for one week, demonstrated decreased serum total and LDL cholesterol levels, and reductions in all oxidative stress markers in all subjects (No authors listed, 2010). Another study, a randomized clinical trial including 81 type II diabetic patients, has shown that broccoli sprout as supplementary treatment (10 g/day for four weeks) significantly decreased serum triglycerides, oxidized LDL, and the atherogenic index of plasma (Bahadoran et al., 2012a). In the same clinical trial, a significant decrease of insulin resistance index and serum insulin concentration was observed (Bahadoran et al., 2012b). Broccoli sprouts decrease renal and vascular oxidative stress and inflammation as well as blood pressure in spontaneously hypertensive stroke-prone rats (Senanayake et al., 2012).

Furthermore, research has suggested that the compounds in broccoli can protect the eyes against macular degeneration (Tanito et al., 2005), and alleviates inflammation, synovial hyperplasia, and bone destruction in mice suffering arthritis rheumatoid (Kong et al., 2010).

### *Dandelion (Taraxacum officinale)*

Dandelion (lion's tooth, blow ball, telltime) is a perennial plant that belongs to the *Compositae* family as well as *Artichokes* (Williams et al., 1996; Schütz et al., 2005; Gonzalez-Castejon et al., 2012). The young leaves of dandelion are consumed fresh as a salad, and the roots are roasted and are utilized as a coffee substitute (Williams et al., 1996; Schütz et al., 2005; Gonzalez-Castejon et al., 2012). In addition, flowers extracts are used as flavor components in various food products, including soft drinks, frozen dairy desserts, baked goods, puddings and cheese (Gonzalez-Castejon et al., 2012). Apart from being consumed as food, dandelion is traditionally used as herbal medicine for the treatment of digestive complaints, hepatobiliary disturbances, hyperlipidemia, dyspepsia, rheumatism, and renal disorders (Williams et al., 1996; Schütz et al., 2005; Gonzalez-Castejon et al., 2012).

The health-promoting potential of dandelion is attributed to its high phenolic content, mainly hydroxycinnamic acid derivatives and flavonoids like chlorogenic and chicoric acids, luteolin, and quercetin glycosides (Williams et al., 1996; Schütz et al., 2005; Gonzalez-Castejon et al., 2012). Furthermore, several sesquiterpene lactones have been identified as the bitter principles typical of *Compositae* in dandelion (Williams et al., 1996; Schütz et al., 2005; Gonzalez-Castejon et al., 2012). Besides these secondary metabolites, dandelion contains inulin, a characteristic carbohydrate of *Compositae*, minerals, and some amino-acids (Williams et al., 1996; Schütz et al., 2005; Gonzalez-Castejon et al., 2012).

Many studies have demonstrated the antioxidant activity of dandelion, both in vitro (Menghini et al., 2010; You et al., 2010) and in vivo (Choi et al., 2010; Menghini et al., 2010; You et al., 2010). In terms of constituents, dandelion flowers are a potentially exploitable source of antioxidants, because of

their elevated phenolic content attributed to caffeic and chlorogenic acids, and to the flavones luteolin and luteolin 7-*O*-glucoside (Hu and Kitts, 2005).

Mechanistically, dandelion downregulates the expression of certain pro-inflammatory mediators. Its antiinflammatory properties, in part attributed to sesquiterpene glucosides, have been proven showing a potent inhibitory activity of 86% against leukotriene B<sub>4</sub> (Kashiwada et al., 2001). The high content in polyphenols also mediates the downregulation of NO, prostaglandin E<sub>2</sub>, and proinflammatory cytokines production. Some researchers observed that luteolin and luteolin 7-*O*-glucoside (<20  $\mu$ M) from dandelion downregulate both inducible iNOS and COX-2 expression (Hu and Kitts, 2004; Koh et al., 2010). Very recent animal studies have reported protective antiinflammatory effects of dandelion on acute lung injury (Liu et al., 2010a,b) *via* blocking mitogen-activated protein kinase, extracellular signal-regulated kinases, and Akt-related signaling cascades and attenuation of neutrophil chemotaxis and respiratory burst (Lee et al., 2010).

A wide range of studies in animals and in vitro reported the positive effects of different dandelion extracts on various risk factors of cardiovascular disease (Choi et al., 2010; Gonzalez-Castejon et al., 2012). The hypoglycemic properties of dandelion have been studied in vitro and in vivo. It appears that certain dandelion extracts (40  $\mu$ g/mL) stimulate the release of insulin by pancreatic  $\beta$ -cell and hypoglycemic activities (Hussain et al., 2004). In vivo, a significant decreasing in serum glucose concentrations was observed in diabetic rats after administration of dandelion leaf extracts (Cho et al., 2002). Dandelion root extracts exert inhibitory effects on human platelet aggregation, at least in vitro (Neef et al., 1996). Such extracts exert a maximal inhibition of 85% at a concentration corresponding to 40 mg dried root/mL of human platelet-rich plasma. Low-molecular polysaccharides produce a 91% inhibition, while a fraction enriched in triterpenes and steroids showed an 80% inhibition of platelet aggregation, at equivalent concentrations (Neef et al., 1996).

Dandelion has also been considered as a potential anti-obesity agent with limited side effects. Its potent inhibitory activity on pancreatic lipase was evaluated in vitro and in vivo proving a pancreatic lipase inhibitory activity—at a concentration of 250  $\mu$ g/mL—about 90.2% of that of Orlistat, a pancreatic lipase inhibitor used as anti-obesity medication endowed with several untoward side effects (Zhang et al., 2008).

On the other hand, dandelion has been used in traditional medicine for the treatment of various types of cancer. In in vitro studies have been observed that extracts at concentrations of 0.2 mg/mL induce apoptosis in human hepatoma cells and exhibit cytotoxic activities in human intestinal carcinoma cells (Koo et al., 2004). Recent results indicate that dandelion extracts decrease the growth of breast cancer cells by 40%, after 96 hours of incubation, and block invasion of prostate cancer cells (Sigstedt et al., 2008). A recent study indicates that components of dandelion root act either alone or synergistically

to induce selectively cancer cell death in human leukemia with no toxicity to noncancerous cells (Ovadje et al., 2010).

Dandelion roots decrease lipid peroxidation, in turn protecting against alcoholic liver damage (You et al., 2010). Recent animal studies have revealed the hepatoprotective effect of two polysaccharides isolated from dandelion (Park et al., 2010; You et al., 2010). These polysaccharides (305 mg/kg body weight, for 7 days) attenuate hepatic damage in Sprague-Dawley rats through the regulation of NF- $\kappa$ B and its mediators such as iNOS, COX-2, TNF- $\alpha$ , and IL-1 (Domitrovic et al., 2010). Such polysaccharides promote the immune-modulator activities of dandelion (Yoon, 2008). The health benefits of dandelion roots have also been explored in relation to their proportion of inulin, which works as demulcent and prebiotic agent that strongly enhances immune function. An infusion of dandelion root stimulates—in vitro—the growth of 14 strains of bifidobacterias (Trojanová et al., 2004). Furthermore, an animal study showed that dandelion extract for six weeks upregulates the expression of estrogen and progesterone receptors, and follicle-stimulating hormone receptor in the adipose tissue and reproductive organs of mice, suggesting its potential application to treat reproductive hormone-related disturbances (Zhi et al., 2007).

### *Garlic (Allium sativum)*

*Allium sativum*, commonly known as garlic, belongs to the *Liliaceae* family (Block, 2010). Its close relatives include onions, shallots, leek, and chive (Block, 2010). Garlic has been used for both culinary and medicinal purposes for hundreds of years in many cultures (Simonetti, 1990; Newall et al., 1996). It contains a higher concentration of sulfur compounds such as diallyl sulphate, alliin, ajoene, allicin than any other *Allium* species, in addition to several enzymes, amino acids, and minerals such as selenium (Simonetti, 1990; Newall et al., 1996; Block, 2010). The sulfur compounds are responsible for both garlic's pungent odor and many of its medicinal effects (Simonetti, 1990; Newall et al., 1996; Block, 2010).

In in vitro studies, garlic has been found to have anti-bacterial, anti-viral, and anti-fungal properties (Sohn et al., 2009). Clinical trials have reported that fresh garlic has significant anti-microbial activities (Groppo et al., 2007; Sohn et al., 2009). In addition, scientific studies have shown that garlic has anti-inflammatory actions (Keiss et al., 2003; Sohn et al., 2009; Block, 2010), anti-cancer properties (Kaschula et al., 2010; Antony and Singh, 2011), and beneficial effects on several risk factors for cardiovascular disease (Block, 2010; Ginter and Simko, 2010; Sobenin et al., 2010).

Notably, human studies have evaluated garlic's lipid lowering effects (Durak et al., 2004). Case series and controlled trials in healthy adults given garlic supplements along with cholesterol-rich diets indicate that garlic reduces serum cholesterol levels and increases fibrinolytic activity (Durak et al., 2004). Furthermore, garlic helps reducing blood pressure

(Ried et al., 2010) and total lipid concentrations (Durak et al., 2004; Kojuri et al., 2007). Also, garlic improves the elasticity of blood vessels and increases blood circulation, relieving cramps and circulatory disorders (Chan et al., 2007; Kojuri et al., 2007). Several studies found that garlic supplementation reduces accumulation of cholesterol in the vascular walls of animals (Chan et al., 2007), inhibits vascular calcification in human patients with high blood cholesterol, and reduces platelet aggregation (Steiner and Lin, 1998; Durak et al., 2004; Rahman, 2007). Different studies also found that moderate amounts of garlic could help regulating blood glucose levels and potentially lessen diabetes complications, as well as fight infections (Kook et al., 2009).

Several epidemiological studies show an association between increased intakes of garlic and reduced risk of certain types of cancer (Fleischauer and Arab, 2001). An analysis of data from seven population studies showed that the higher the amount of garlic consumed, the lower the risk of stomach, esophagus, pancreatic, and colorectal cancer (Tanaka et al., 2004; Chan et al., 2005; Gonzalez et al., 2006). Evidence also indicates that increased (>10 g/day vs. <2.2 g/day) allium intake is associated with an approximately 50% reduction in the prostate cancer risk (Hsing et al., 2002). Mechanistic studies showed that garlic reduces the development of mammary cancer in animals and suppresses the growth of human breast cancer cells (Tsubura et al., 2011).

### **Cocoa (*Theobroma cacao*)**

Cocoa beans and their derived products such as chocolate are rich in antioxidants, mostly catechins, epicatechin, and especially the polymers procyanidins (Hooper et al., 2012). In addition to these phytochemicals, cocoa also contain flavonoids and xanthines, with theobromine as the main representative of the latter, which have vasodilatory properties (Hooper et al., 2012). Due to its high concentration of this series of phytochemicals with a variety of biological functions, cocoa as a plant and chocolate as food may have beneficial health effects against oxidative stress and chronic inflammation, in addition to ameliorate cardiovascular disorders, cancer, and other pathological conditions (Hooper et al., 2012; Ramos-Romero et al., 2012).

Many clinical and epidemiological studies indicate that appropriate intakes of cacao reduce the incidence of cardiovascular disease and of several of its risk factors (Corti et al., 2009). Chocolate consumption has a positive influence on hypertensive, inflammatory, atherogenic, and thrombotic health conditions as well as impact on insulin sensitivity and vascular endothelial function (Buijsse et al., 2010; Mostofsky et al., 2010; Oba et al., 2010; Djousse et al., 2011). As an example, the Stockholm Heart Epidemiology Program (Janszky et al., 2009) is a study including a cohort of 1169 patients who had suffered myocardial infarction, who are being followed for eight years. Its results show that chocolate

consumption is associated with lower cardiac mortality in a dose-dependent manner. Recent meta-analysis reported that higher levels of chocolate consumption are associated with a reduction of about a third in the risk of cardio-metabolic disorders (Desch et al., 2010; Buitrago-Lopez et al., 2011). This beneficial association was significant for any cardiovascular disease (37% reduction), diabetes (31% reduction), and stroke (29% reduction) (Buitrago-Lopez et al., 2011).

There are few epidemiological studies and scant data that correlate the incidence of cancer with consumption of cocoa. While several studies of intervention—despite their short duration—have reported favorable changes in biomarkers assessing antioxidant status, there is very little evidence related to cancer markers (Maskarinec, 2009). However, these metabolic studies indicate that regular intake of cocoa and its derivatives increase plasma level of antioxidants, which might be a desirable effect because it would protect against reactive oxygen species (ROS) (Weisburger, 2001).

### **Olives (*Olea europaea*)**

The olive tree, *Olea europaea*, is native to the Mediterranean countries and parts of Asia Minor. In folk medicine, the fruit and its oil have a wide range of therapeutic and culinary applications (López-Miranda et al., 2010; Belarbi et al., 2011; Visioli, 2012). Traditionally, olives have been used as emollients, laxatives, nutritive, sedatives, and tonics (Belarbi et al., 2011). Specific conditions traditionally treated include colic, alopecia, paralysis, rheumatic pain, sciatica, and hypertension (López-Miranda et al., 2010; Belarbi et al., 2011). Raw olives are very bitter and inedible and, therefore, must be treated before consumption (Waterman and Lockwood, 2007). They can be harvested when still unripe (green olives) or when fully ripe (black olives) (Waterman and Lockwood, 2007). Ripe olives are pressed to obtain olive oil (Waterman and Lockwood, 2007).

In vitro and animal studies show that olives have potent antioxidant activities and could partially account for the observed health benefits of the Mediterranean diet (Raederstorff, 2009; Kawaguchi et al., 2011).

Although the composition of olives is complex, the major groups of compounds thought to contribute to the observed health benefits include oleic acid, phenols, and squalene (Owen et al., 2000), all of which have been found to inhibit oxidative stress (Visioli and Bernardini, 2011). Major phenols include hydroxytyrosol, tyrosol, oleuropein (Perona et al., 2006). Hydroxytyrosol and oleuropein are more potent at scavenging free radicals than the endogenous antioxidant vitamin E and the exogenous antioxidants dimethyl sulfoxide and butylated hydroxytoluene (Owen et al., 2000; Visioli et al., 2002). Human intervention studies indicate that these two phenols inhibit LDL oxidation in plasma and regulate several biomarkers of oxidative damage in the micro-molar range (Baroni et al., 1999; Owen et al., 2000; Visioli et al., 2002).

The olive has been widely studied for its effects on cardiovascular disease, specifically for its ability to reduce blood pressure and LDL-cholesterol (Baroni et al., 1999; Waterman and Lockwood, 2007). In vitro studies have demonstrated hydroxytyrosol and oleuropein are capable of inhibiting production of isoprostanes, a marker of LDL oxidation (Salami et al., 1995). Several studies have also demonstrated the antihypertensive properties of olives and olive oil (Alonso et al., 2006; Belarbi et al., 2011), and since hydroxytyrosol has biological activity beyond its antioxidant capacity, it affects a wide range of enzymes, including cyclooxygenase and NAD(P)H oxidase, and reduce platelet aggregation (Fabiani et al., 2002; Visioli et al., 2002).

Strong epidemiological evidence suggests that regular intake of olive and derivative products lowers the incidence of certain cancers, including breast, skin, and colon (Owen et al., 2000; Harwood and Yaqoob, 2002). One mechanism associated with the anticancer effects of hydroxytyrosol and oleuropein is through the prevention of DNA damage that in turn prevents mutagenesis and carcinogenesis (Visioli et al., 2002). In addition, oleuropein has been shown to exert hepatoprotective activities (Domitrovic et al., 2012). In vitro studies have demonstrated the antimicrobial activity of hydroxytyrosol, tyrosol, and oleuropein against several strains of bacteria implicated in intestinal and respiratory infections (Romero et al., 2007). Furthermore, the administration of olive in lipid emulsions may exert beneficial effects on the immune system (Puertollano et al., 2010).

### *Grapes (Vitis vinifera L.)*

The health effects of grapes on cardiovascular disease risk, mainly on endothelial function, LDL oxidation, progression of atherosclerosis, and reduction in oxidative stress, have been thoroughly investigated (Folts, 2002; Dohadwala and Vita, 2009; Feringa et al., 2011; Weseler et al., 2011). There is evidence suggesting that moderate consumption of grapes can decrease mortality (Vislocky and Fernandez, 2010). Numerous studies in vitro as well as in animals and humans demonstrate beneficial effects of grape polyphenols on cardiovascular risk factors including hypertension (Schini-Kerth et al., 2011) and diabetes (Zunino, 2009; Chuang and McIntosh, 2011), among others.

Many studies have also suggested chemopreventive benefits (Athar et al., 2007; Neves et al., 2012). Data are particularly compelling in the area of skin cancer prevention (Athar et al., 2007; Neves et al., 2012). Further favorable effects of grapes on oral health (Wu, 2009), immune function (Percival, 2009), anti-inflammatory (Chuang and McIntosh, 2011; Zhou et al., 2011; Lee et al., 2012), and anti-allergic (Lee et al., 2012) systems have also been reported. Other studies show that grape phenolic compounds possess antibacterial (Baydar et al., 2004) and antiviral (Liu et al., 2010a,b) activities. Novel and exciting results from independent research demonstrate that

grapes have beneficial effects on other devastating illnesses, such as Alzheimer's disease, and age-related cognitive decline (Nakata et al., 2012; Wang et al., 2012).

These effects are often attributed to a valuable source of phytochemicals found in grapes with strong antioxidant activity that helps to prevent oxidative damage as well as other actions such as increasing nitric oxide production (Folts, 2002; Iriti and Faoro, 2009; Xia et al., 2010; Kawaguchi et al., 2011). Grapes contain high concentrations of polyphenols, especially flavonoids (Folts, 2002; Iriti and Faoro, 2009). The main grape polyphenols are anthocyanins in red grapes and flavan-3-ols in the white grapes; red grapes contain more total polyphenols than white grapes (Folts, 2002; Iriti and Faoro, 2009; Lee et al., 2012). Grape seeds contain significant amounts of proanthocyanidins or condensed tannins with therapeutic potential in the prevention and treatment of vascular complications in diabetic patients (Folts, 2002; Iriti and Faoro, 2009; Lee et al., 2012). Studies in human and experimental models demonstrate antioxidant properties closely associated with the maintenance of endothelial function, protection against LDL oxidation, and neuroprotective effects (Feringa et al., 2011). The flavonoids ellagic acid and the phytoalexin resveratrol have been suggested to reduce the risk of coronary heart disease (Folts, 2002; Iriti and Faoro, 2009; Xia et al., 2010; Ramprasath and Jones, 2010). Even though human evidence is lacking, resveratrol has been proposed to have heart-protective potential (Folts, 2002; Ramprasath and Jones, 2010), and cancer chemopreventive activities in different stages of carcinogenesis (Athar et al., 2007; Iriti and Faoro, 2009; Xia et al., 2010; Nakata et al., 2012; Neves et al., 2012). It must be underlined that these proposed activities await in vivo proof and that the concentrations of resveratrol in grapes are extremely low (Rotches-Ribalta et al., 2012).

### CONCLUSIONS

The influence of diet on the overall health status has been recognized since ancient times, when pathologic conditions were treated with plants and natural foods believed to have medicinal properties. Currently, epidemiological studies have confirmed the association between diets rich in fruits and vegetables and lower risk for chronic diseases such as cancer and cardiovascular disease. A reduced risk of obesity and better control of diabetes are some additional benefits from increased consumption of plant foods. In addition, evidence also suggests that diets rich in fruit and vegetables decrease the risk of premature mortality. Fruits and vegetables, apart from being rich sources of vitamins, minerals, and fiber, also contain a wide variety of phytochemicals with potent bioactivity, among which polyphenols are mostly well known. Recently, the consumer perception is that phytochemicals with antioxidant properties present in these foodstuffs might prevent some of the processes involved in the development of cancer (i.e. by protecting DNA from oxidative damage) and in the

development of cardiovascular disease (i.e. by inhibiting oxidative damage to LDL-cholesterol). However, epidemiologic studies on the impact of phytochemicals in human health have been inconclusive. Even when good evidence exists, results need to be interpreted and divulged carefully in relation to human health benefits, as phytochemicals may have limited bioavailability and may also be extensively metabolized. In addition, some phytochemicals can be toxic and mutagenic in some cell culture systems. Also, it should be borne in mind that fruits and vegetables contain a complex mixture of metabolites that cannot be simulated by single purified phytochemicals. In this context—and until more is known about the activity, bioavailability, and the distribution of phytochemicals in the organism—it appears correct to recommend fruits and vegetables in their entirety (minimum of 400 g/day) rather than isolated components as dietary supplements.

### ACKNOWLEDGMENTS

The author thanks Professor Visioli for his fruitful comments and revision of the manuscript.

### FUNDING

This work has been supported by the Spanish Ministry of Economy and Competitiveness (R&C2007-01920).

### STATEMENT OF INTEREST

The author declares no competing interests.

### REFERENCES

- Aires, A., Fernandes, C., Carvalho, R., Bennett, R. N., Saavedra, M. J. and Rosa, E. A. (2011). Seasonal effects on bioactive compounds and antioxidant capacity of six economically important *Brassica* vegetables. *Molecules*. **16**(8):6816–6832.
- Alonso, A., Ruiz-Gutierrez, V. and Martinez-Gonzalez, M. A. (2006). Mono-unsaturated fatty acids, olive oil and blood pressure: Epidemiological, clinical and experimental evidence. *Public Health Nutr.* **9**(2):251–257.
- Antony, M. L. and Singh, S. V. (2011). Molecular mechanisms and targets of cancer chemoprevention by garlic-derived bioactive compound diallyl trisulfide. *Indian J. Exp. Biol.* **49**(11):805–816.
- Athar, M., Back, J. H., Tang, X., Kim, K. H., Kopelovich, L., Bickers, D. R. and Kim, A. L. (2007). Resveratrol: A review of pre-clinical studies for human cancer prevention. *Toxicol. Appl. Pharmacol.* **224**(3):274–283.
- Aune, D., Chan, D. S., Vieira, A. R., Navarro Rosenblatt, D. A., Vieira, R., Greenwood, D. C. and Norat, T. (2012b). Dietary compared with blood concentrations of carotenoids and breast cancer risk: A systematic review and meta-analysis of prospective studies. *Am. J. Clin. Nutr.* **96**(2):356–373.
- Aune, D., Chan, D. S., Vieira, A. R., Rosenblatt, D. A., Vieira, R., Greenwood, D. C. and Norat, T. (2012a). Fruits, vegetables and breast cancer risk: A systematic review and meta-analysis of prospective studies. *Breast Cancer Res. Treat.* **134**(2):479–493.
- Bach-Faig, A., Berry, E. M., Lairon, D., Reguant, J., Trichopoulou, A., Dernini, S., Medina, F. X., Battino, M., Belahsen, R., Miranda, G. and Serra-Majem, L. (2011). Mediterranean Diet Foundation Expert Group. 1. Mediterranean diet pyramid today. Science and cultural updates. *Public Health Nutr.* **14**(12A):2274–2284.
- Badimon, L., Vilahur, G. and Padro, T. (2010). Nutraceuticals and atherosclerosis: Human trials. *Cardiovasc Ther.* **28**(4):202–215.
- Bahadoran, Z., Mirmiran, Hosseinpahan, P., Rajab F., Asghari G., and Azizi, F. (2012a). Broccoli sprouts powder could improve serum triglyceride and oxidized LDL/LDL-cholesterol ratio in type 2 diabetic patients: A randomized double-blind placebo-controlled clinical trial. *Diabetes Res. Clin. Pract.* **96**(3):348–354.
- Bahadoran, Z., Tohidi, M., Nazeri, P., Mehran, M., Azizi, F. and Mirmiran, P. (2012b). Effect of broccoli sprout on insulin resistance in type 2 diabetic patients: A randomized double-blind clinical trial. *Int. J. Food Sci. Nutr.* **63**(7):767–771.
- Baldrick, F. R., Woodside, J. V., Elborn, J. S., Young, I. S. and McKinley, M. C. (2011). Biomarkers of fruit and vegetable intake in human intervention studies: A systematic review. *Crit. Rev. Food Sci. Nutr.* **51**(9):795–815.
- Baroni, S. S., Amelio, M., Sangiorgi, Z., Gaddi, A. and Battino, M. (1999). Solid monounsaturated diet lowers LDL unsaturation trait and oxidisability in hypercholesterolemic (type IIb) patients. *Free Radic. Res.* **30**(4):275–285.
- Baydar, N. G., Ozkan, G. and Sagdic, O. (2004). Total phenolic contents and antibacterial activities of grape (*Vitis vinifera* L.) extracts. *Food Control*. **15**(5):335–339.
- Belarbi, M., Bendimerad, S., Sour, S., Soualem, Z., Baghdad, C., Hmimed, S., Chemat, F. and Visioli, F. (2011). Oleaster oil positively modulates plasma lipids in humans. *J. Agric. Food Chem.* **59**(16):8667–8669.
- Berretta, M., Lleshi, A., Fisichella, R., Berretta, S., Basile, F., Li Volti, G., Bolognese, A., Biondi, A., De Paoli, P., Tirelli, U. and Cappellani, A. (2012). The role of nutrition in the development of esophageal cancer: What do we know? *Front Biosci. (Elite Ed)* **4**:351–357.
- Block, E. (2010). *Garlic and Other Alliums: The Lore and the Science*. 1st ed. Royal Society of Chemistry, Cambridge.
- Boeing, H., Bechthold, A., Bub, A., Ellinger, S., Haller, D., Kroke, A., Leschik-Bonnet, E., Müller, M. J., Oberitter, H., Schulze, M., Stehle, P. and Watzl, B. (2012). Critical review: Vegetables and fruit in the prevention of chronic diseases. *Eur. J. Nutr.* **51**(6):637–663.
- Bruckert, E. and Rosenbaum, D. (2011). Lowering LDL-cholesterol through diet: Potential role in the statin era. *Curr. Opin. Lipidol.* **22**(1):43–48.
- Buck, P. A. (1956). Origin and Taxonomy of Broccoli. *Economic Botany*. **10**(3):250–253.
- Buijsse, B., Weikert, C., Drogan, D., Bergmann, M. and Boeing, H. (2010). Chocolate consumption in relation to blood pressure and risk of cardiovascular disease in German adults. *Eur. Heart J.* **31**(13):1616–1623.
- Buitrago-Lopez, A., Sanderson, J., Johnson, L., Warnakula, S., Wood, A., Di Angelantonio, E. and Franco, O. H. (2011). Chocolate consumption and cardiometabolic disorders: Systematic review and meta-analysis. *BMJ*. **343**:d4488.
- Calder, P. C., Ahluwalia, N., Brouns, F., Buetler, T., Clement, K., Cunningham, K., Esposito, K., Jönsson, L. S., Kolb, H., Lansink, M., Marcos, A., Margioris, A., Matusheski, N., Nordmann, H., O'Brien, J., Pugliese, G., Rizkalla, S., Schalkwijk, C., Tuomilehto, J., Wärnberg, J., Watzl, B. and Winkhofer-Roob, B. M. (2011). Dietary factors and low-grade inflammation in relation to overweight and obesity. *Br. J. Nutr.* **106**(Suppl 3):S5–S78.
- Cappellani, A., Cavallaro, A., Di Vita, M., Zanghi, A., Piccolo, G., Lo Menzo, E., Cavallaro, V., Malaguamera, M., Giaquinta, A., Veroux, M., Cimino, L. and Berretta, M. (2012). Diet and pancreatic cancer: Many questions with few certainties. *Eur. Rev. Med. Pharmacol. Sci.* **16**(2):192–206.
- Chan, J. M., Wang, F. and Holly, E. A. (2005). Vegetable and fruit intake and pancreatic cancer in a population-based case-control study in the San Francisco bay area. *Cancer Epidemiol. Biomarkers Prev.* **14**(9):2093–2097.
- Chan, K. C., Yin, M. C. and Chao, W. J. (2007). Effect of diallyl trisulfide-rich garlic oil on blood coagulation and plasma activity of anticoagulation factors in rats. *Food Chem. Toxicol.* **45**(3):502–507.
- Chiva-Blanch, G. and Visioli, F. (2012). Polyphenols and health: Moving beyond antioxidants. *J. Berry Res.* **2**(2):63–71.



- Cho, S. Y., Park, J. Y., Park, E. M., Choi, M. S., Lee, M. K., Jeon, S. M., Jang, M. K. and Kim, M. J. (2002). Alternation of hepatic antioxidant enzyme activities and lipid profile in streptozotocin-induced diabetic rats by supplementation of dandelion water extract. *Clin. Chim. Acta.* **317**(1–2):109–117.
- Choi, U. K., Lee, O. H., Yim, J. H., Cho, C. W., Rhee, Y. K., Lim, S. I. and Kim, Y. C. (2010). Hypolipidemic and antioxidant effects of dandelion (*Taraxacum officinale*) root and leaf on cholesterol-fed rabbits. *Int. J. Mol. Sci.* **11**(1):67–78.
- Choudhary, S. P. and Tran, L. S. (2011). Phytosterols: Perspectives in human nutrition and clinical therapy. *Curr. Med. Chem.* **18**(29):4557–4567.
- Chuang, C. C. and McIntosh, M. K. (2011). Potential mechanisms by which polyphenol-rich grapes prevent obesity-mediated inflammation and metabolic diseases. *Annu. Rev. Nutr.* **31**:155–176.
- Corti, R., Flammer, A. J., Hollenberg, N. K. and Luscher, T. F. (2009). Cocoa and cardiovascular health. *Circulation.* **119**(10):1433–1442.
- De Kok, T. M., De Waard, P., Wilms, L. C. and Van Breda, S. G. (2010). Antioxidative and antigenotoxic properties of vegetables and dietary phytochemicals: The value of genomics biomarkers in molecular epidemiology. *Mol. Nutr. Food Res.* **54**(2):208–217.
- Desch, S., Schmidt, J., Kobler, D., Sonnabend, M., Eitel, I., Sareban, M., Rahimi, K., Schuler, G. and Thiele, H. (2010). Effect of cocoa products on blood pressure: Systematic review and meta-analysis. *Am. J. Hypertens.* **23**(1):97–103.
- Dinkova-Kostova, A. T. and Kostov, R. V. (2012). Glucosinolates and isothiocyanates in health and disease. *Trends Mol. Med.* **18**(6):337–347.
- Djousse, L., Hopkins, P. N., Arnett, D. K., Pankow, J. S., Borecki, I., North, K. E. and Curtis Ellison, R. (2011). Chocolate consumption is inversely associated with calcified atherosclerotic plaque in the coronary arteries: The NHLBI Family Heart Study. *Clin. Nutr.* **30**(2):182–187.
- Dohadwala, M. M. and Vita, J. A. (2009). Grapes and cardiovascular disease. *J. Nutr.* **139**(9):1788S–1793S.
- Domitrovic, R., Jakovac, H., Marchesi, V. V., Sain, I., Romic, Z. and Rahelic, D. (2012). Preventive and therapeutic effects of oleuropein against carbon tetrachloride-induced liver damage in mice. *Pharmacol. Res.* **65**(4):451–464.
- Domitrovic, R., Jakovac, H., Romic, Z., Rahelic, D. and Tadic, Z. (2010). Anti-fibrotic activity of *Taraxacum officinale* root in carbon tetrachloride-induced liver damage in mice. *J. Ethnopharmacol.* **130**(3):569–577.
- Durak, I., Kavutcu, M., Aytac, B., Avci, A., Devrim, E. and Ozturk, H. (2004). Effects of garlic extract consumption on blood lipid and oxidant/antioxidant parameters in humans with high blood cholesterol. *J. Nutr. Biochem.* **15**(6):374–377.
- El Hachimi, M. and Rosado, J. A. (2011). Modulation of platelet function and signaling by flavonoids. *Mini Rev. Med. Chem.* **11**(2):131–142.
- Fabiani, R., De Bartolomeo, A., Rosignoli, P., Servili, M., Montedoro, G. F. and Morozzi, G. (2002). Cancer chemoprevention by hydroxytyrosol isolated from virgin olive oil through G1 cell cycle arrest and apoptosis. *Eur. J. Cancer Prev.* **11**(4):351–358.
- Feringa, H. H., Laskey, D. A., Dickson, J. E. and Coleman, C. I. (2011). The effect of grape seed extract on cardiovascular risk markers: A meta-analysis of randomized controlled trials. *J. Am. Diet. Assoc.* **111**(8):1173–1181.
- Ferrari, N., Tosetti, F., De Flora, S., Donatelli, F., Sogno, I. M., Noonan D. and Albini, A. (2011). Diet-derived phytochemicals: From cancer chemoprevention to cardio-oncological prevention. *Curr. Drug Targets.* **12**(13):1909–1924.
- Fimognari, C., Nüsse, M., Cesari, R., Lori, R., Cantelli-Forti, G. and Hrelia, P. (2002). Growth inhibition, cell-cycle arrest and apoptosis in human T-cell leukemia by the isothiocyanate sulforaphane. *Carcinogenesis.* **23**(4):581–586.
- Finks, S. W., Airee, A., Chow, S. L., Macaulay, T. E., Moranville, M. P., Rogers, K. C. and Trujillo, T. C. (2012). Key articles of dietary interventions that influence cardiovascular mortality. *Pharmacotherapy.* **32**(4):e54–e87.
- Fleischauer, A. T. and Arab, L. (2001). Garlic and cancer: A critical review of the epidemiologic literature. *J. Nutr.* **131**(3s):1032S–1040S.
- Folts, J. D. (2002). Potential health benefits from the flavonoids in grape products on vascular disease. *Adv. Exp. Med. Biol.* **505**:95–111.
- Galleano, M., Pechanova, O. and Fraga, C. G. (2010). Hypertension, nitric oxide, oxidants, and dietary plant polyphenols. *Curr. Pharm. Biotechnol.* **11**(8):837–848.
- Gamet-Payastre, L., Li, P., Lumeau, S., Cassar, G., Dupont, M. A., Chevolleau, S., Gasc, N., Tulliez, J. and Terce, F. (2000). Sulforaphane, a naturally occurring isothiocyanate, induces cell cycle arrest and apoptosis in HT29 human colon cancer cells. *Cancer Res.* **60**(5):1426–1433.
- Giampieri, F., Tulipani, S., Alvarez-Suarez, J. M., Quiles, J. L., Mezzetti, B. and Battino, M. (2012). The strawberry: Composition, nutritional quality, and impact on human health. *Nutrition.* **28**(1):9–19.
- Ginter, E. and Simko, V. (2010). Garlic (*Allium sativum* L.) and cardiovascular diseases. *Bratisl Lek Listy.* **111**(8):452–456.
- Gonzalez, C. A., Pera, G., Agudo, A., Bueno-de-Mesquita, H. B., Ceroti, M., Boeing, H., Schulz, M., Del Giudice, G., Plebani, M., Carneiro, F., Berrino, F., Sacerdote, C., Tumino, R., Panico, S., Berglund, G., Simán, H., Hallmans, G., Stenling, R., Martinez, C., Dorronsoro, M., Barricarte, A., Navarro, C., Quirós, J. R., Allen, N., Key, T. J., Bingham, S., Day, N. E., Linseisen, J., Nagel, G., Overvad, K., Jensen, M. K., Olsen, A., Tjønneland, A., Büchner, F. L., Peeters, P. H., Numans, M. E., Clavel-Chapelon, F., Boutron-Ruault, M. C., Roukos, D., Trichopoulos, A., Psaltopoulou, T., Lund, E., Casagrande, C., Slimani, N., Jenab, M. and Riboli, E. (2006). Fruit and vegetable intake and the risk of stomach and esophagus adenocarcinoma. European Prospective Investigation into Cancer and Nutrition (EPIC-EURGAST). *Int. J. Cancer.* **118**(10):2559–2566.
- Gonzalez-Castejon, M., Visioli, F. and Rodriguez-Casado, A. (2012). Diverse biological activities of dandelion. *Nutr. Rev.* **70**(9):534–547.
- Groppi, F., Ramacciato, J., Motta, R., Ferraresi, P. and Sartoratto, A. (2007). Antimicrobial activity of garlic against oral streptococci. *Int. J. Dent. Hyg.* **5**(2):109–115.
- Harwood, J. L. and Yaqoob, P. (2002). Nutritional and health aspects of olive oil. *Eur. J. Lipid Sci. Technol.* **104**(9–10):685–697.
- Holst, B. and Williamson, G. (2004). A critical review of the bioavailability of glucosinolates and related compounds. *Nat. Prod. Rep.* **21**(3):425–447.
- Hooper, L., Kay, C., Abdelhamid, A., Kroon, P. A., Cohn, J. S., Rimm, E. B. and Cassidy, A. (2012). Effects of chocolate, cocoa, and flavan-3-ols on cardiovascular health: A systematic review and meta-analysis of randomized trials. *Am. J. Clin. Nutr.* **95**(3):740–751.
- Hsing, A. W., Chokkalingam, A. P., Gao, Y. T., Madigan, M. P., Deng, J., Gridley, G. and Fraumeni, J. F., Jr. (2002). *Allium* vegetables and risk of prostate cancer: A population-based study. *J. Natl. Cancer Inst.* **94**(21):1648–1651.
- Hu, C. and Kitts, D. D. (2004). Luteolin and luteolin 7-O-glucoside from dandelion flower suppress iNOS and COX-2 in RAW264.7 cells. *Mol. Cell Biochem.* **265**(1–2):107–113.
- Hu, C. and Kitts, D. D. (2005). Dandelion (*Taraxacum officinale*) flower extract suppresses both reactive oxygen species and nitric oxide and prevents lipid oxidation *in vitro*. *Phytomedicine.* **12**(8):588–597.
- Hussain, Z., Waheed, A., Qureshi, R. A., Burdi, D. K., Verspohl, E. J., Khan, N. and Hasan, M. (2004). The effect of medicinal plants of Islamabad and Murree region of Pakistan on insulin secretion from INS-1 cells. *Phytother. Res.* **18**(1):73–77.
- Iriti, M. and Faoro, F. (2009). Bioactivity of grape chemicals for human health. *Nat. Prod. Commun.* **4**(5):611–634.
- Janszky, I., Mukamal, K. J., Ljung, R., Ahnve, S., Ahlbom, A. and Hallqvist, J. (2009). Chocolate consumption and mortality following a first acute myocardial infarction: The Stockholm Heart Epidemiology Program. *J. Intern. Med.* **266**(3):248–257.
- Juge, N., Mithen, R. F. and Traka, M. (2007). Molecular basis for chemoprevention by sulforaphane: A comprehensive review. *Cell. Mol. Life Sci.* **64**(9):1105–1127.
- Kaschula, C. H., Hunter, R., Hassan, H. T., Stellenboom, N., Cotton, J., Zhai, X. Q. and Parker, M. I. (2010). Anti-proliferation activity of synthetic

- ajoene analogues on cancer cell-lines. *Anticancer Agents Med. Chem.* **11** (3):260–266.
- Kashiwada, Y., Takanaka, K., Tsukada, H., Miwa, Y., Taga, T., Tanaka, S. and Ikeshiro, Y. (2001). Sesquiterpene glucosides from antileukotriene B4 release fraction of *Taraxacum officinale*. *J. Asian Nat. Prod. Res.* **3**(3): 191–197.
- Kawaguchi, K., Matsumoto, T. and Kumazawa, Y. (2011). Effects of antioxidant polyphenols on TNF- $\alpha$ -related diseases. *Curr. Top. Med. Chem.* **11** (14):1767–1779.
- Keiss, H. P., Dirsch, V. M., Hartung, T., Haffner, T., Trueman, L., Auger, J., Kahane, R. and Vollmar, A. M. (2003). Garlic (*Allium sativum* L.) modulates cytokine expression in lipopolysaccharide-activated human blood thereby inhibiting NF- $\kappa$ B activity. *J. Nutr.* **133**(7):2171–2175.
- Kensler, T. W., Egner, P. A., Agyeman, A. S., Visvanathan, K., Groopman, J. D., Chen, J. G., Chen, T. Y., Fahey, J. W. and Talalay, P. (2013). Keap1-Nrf2 Signaling: A target for cancer prevention by sulforaphane. *Top. Curr. Chem.* **329**:163–177.
- Koh, Y. J., Cha, D. S., Ko, J. S., Park, H. J. and Choi, H. D. (2010). Anti-inflammatory effect of *Taraxacum officinale* leaves on lipopolysaccharide-induced inflammatory responses in RAW264.7 Cells. *J. Med. Food.* **13** (4):870–878.
- Kojuri, J., Vosoughi, A. R. and Akrami, M. (2007). Effects of *Anethum graveolens* and garlic on lipid profile in hyperlipidemic patients. *Lipids Health Dis.* **1**(6):5.
- Koo, H. N., Hong, S. H., Song, B. K., Kim, C. H., Yoo, Y. H. and Kim, H. M. (2004). *Taraxacum officinale* induces cytotoxicity through TNF- $\alpha$  and IL-1 secretion in HepG2 cells. *Life Sci.* **74**(9):1149–1157.
- Kook, S., Kim, G. H. and Choi, K. (2009). The anti-diabetic effect of onion and garlic in experimental diabetic rats: Meta-analysis. *J. Med. Food.* **12** (3):552–560.
- Kong, J. S., Yoo, S. A., Kim, H. S., Kim, H. A., Yea, K., Ryu, S. H., Chung, Y. J., Cho, C. S. and Kim, W. U. (2010). Inhibition of synovial hyperplasia, rheumatoid T cell activation, and experimental arthritis in mice by sulforaphane, a naturally occurring isothiocyanate. *Arthritis Rheum.* **62**(1): 159–170.
- Latté, K. P., Appel, K. E. and Lampen, A. (2011). Health benefits and possible risks of broccoli—an overview. *Food Chem. Toxicol.* **49**(12):3287–3309.
- Lee, T., Kwon, H. S., Bang, B. R., Lee, Y. S., Park, M. Y., Moon, K. A., Kim, T. B., Lee, K. Y., Moon, H. B. and Cho, Y. S. (2012). Grape seed proanthocyanidin extract attenuates allergic inflammation in murine models of asthma. *J. Clin. Immunol. Dec.* **32**(6):1292–1304.
- Lee, J. P., Li, Y. C., Chen, H. Y., Lin, R. H., Huang, S. S., Chen, H. L., Kuan, P. C., Liao, M. F., Chen, C. J. and Kuan, Y. H. (2010). Protective effects of luteolin against lipopolysaccharide-induced acute lung injury involve inhibition of MEK/ERK and PI3K/Akt pathways in neutrophils. *Acta Pharmacol. Sin.* **31**(7):831–838.
- Liang, H., Lai, B. and Yuan, Q. (2008). Sulforaphane induces cell-cycle arrest and apoptosis in cultured human lung adenocarcinoma LTP-A2 cells and retards growth of STEP-A2 xenografts *in vivo*. *J. Nat. Prod.* **71**(11): 1911–1914.
- Lipinski, B. (2011). Hydroxyl radical and its scavengers in health and disease. *Oxid. Med. Cell Longev.* **2011**:809696.
- Liu, L., Xiong, H., Ping, J., Ju, Y. and Zhan, X. (2010a). *Taraxacum officinale* protects against lipopolysaccharide-induced acute lung injury in mice. *J. Ethnopharmacol.* **130**(2):392–397.
- Liu, T., Zhao, J., Li, H. and Ma, L. (2010b). Evaluation on anti-hepatitis viral activity of *Vitis vinifera* L. *Molecules.* **15**(10):7415–7422.
- López-Miranda, J., Pérez-Jiménez, F., Ros, E., De Caterina, R., Badimón, L., Covas, M. I., Escribá, E., Ordóñez, J. M., Soriguer, F., Abiá, R., Alarcón de la Lastra, C., Battino, M., Corella, D., Chamorro-Quirós J., Delgado-Lista, J., Giugliano, D., Esposito, K., Estruch, R., Fernández-Real, J. M., Gáforio, J. J., La Vecchia, C., Lairon, D., López-Segura, F., Mata, P., Menéndez, J. A., Muriana, F. J., Osada, J., Panagiotakos, D. B., Paniagua, J. A., Pérez-Martínez, P., Perona, J., Peinado, M. A., Pineda-Priego, M., Poulsen, H. E., Quiles, J. L., Ramírez-Tortosa, M. C., Ruano, J., Serra-Majem, L., Solá, R., Solanas, M., Solfrizzi, V., de la Torre-Fornell, R., Trichopoulou, A., Uceda, M., Villalba-Montoro, J. M., Villar-Ortiz, J. R., Visioli, F. and Yiannakouris, N. (2010). Olive oil and health: Summary of the II international conference on olive oil and health consensus report, Jaén and Córdoba (Spain) 2008. *Nutr. Metab. Cardiovasc Dis.* **20**(4):284–294.
- Lu, R., Dan, H., Wu, R., Meng, W., Liu, N., Jin, X., Zhou, M., Zeng, X., Zhou, G. and Chen, Q. (2011). Lycopene: Features and potential significance in the oral cancer and precancerous lesions. *J. Oral. Pathol. Med.* **40**(5): 361–368.
- Magalhaes, B., Peleteiro, B. and Lunet, N. (2012). Dietary patterns and colorectal cancer: Systematic review and meta-analysis. *Eur. J. Cancer Prev.* **21** (1):15–23.
- Maskarinec, G. (2009). Cancer protective properties of cocoa: A review of the epidemiologic evidence. *Nutr. Cancer.* **61**(5):573–579.
- McCullough, M. L. and Giovannucci, E. L. (2004). Diet and cancer prevention. *Oncogene.* **23**(38):6349–6364.
- Menghini, L., Genovese, S., Epifano, F., Tirillini, B., Ferrante, C. and Leporini, L. (2010). Anti-proliferative, protective and antioxidant effects of artichoke, dandelion, turmeric and rosemary extracts and their formulation. *Int. J. Immunopathol. Pharmacol.* **23**(2):601–610.
- Mostofsky, E., Levitan, E. B., Wolk, A. and Mittleman, M. A. (2010). Chocolate intake and incidence of heart failure: A population-based, prospective study of middle-aged and elderly women. *Circ. Heart Fail.* **3**(5):612–616.
- Nakata, R., Takahashi, S. and Inoue, H. (2012). Recent advances in the study on resveratrol. *Biol. Pharm. Bull.* **35**(3):273–279.
- Neef, H., Cilli, F., Declercq, P. J. and Laekeman, G. (1996). Platelet anti-aggregating activity of *Taraxacum officinale* Weber. *Phytother. Res.* **10**: S138–S140.
- Neves, A. R., Lucio, M., Lima, J. L. and Reis, S. (2012). Resveratrol in medicinal chemistry: A critical review of its pharmacokinetics, drug-delivery, and membrane interactions. *Curr. Med. Chem.* **19**(11):1663–1681.
- Newall, C. A., Anderson, L. A. and Phillipson, J. D. (1996). *Herbal medicines: A guide for health-care professionals*. Pharmaceutical Press, London.
- Niki, E. (2011). Antioxidant capacity: Which capacity and how to assess it? *J. Berry Res.* **1**(4):169–176.
- Oba, S., Nagata, C., Nakamura, K., Fujii, K., Kawachi, T., Takatsuka, N. and Shimizu, H. (2010). Consumption of coffee, green tea, oolong tea, black tea, chocolate snacks and the caffeine content in relation to risk of diabetes in Japanese men and women. *Br. J. Nutr.* **103**(3):453–459.
- Omar, E. A., Kam, A., Alqahtani, A., Li, K. M., Razmovski-Naumovski, V., Nammi, S., Chan, K., Roufogalis, B. D. and Li, G. Q. (2010). Herbal medicines and nutraceuticals for diabetic vascular complications: Mechanisms of action and bioactive phytochemicals. *Curr. Pharm. Des.* **16**(34):3776–3807.
- Ovadjee, P., Chatterjee, S., Griffin, C., Tran, C., Hamm, C. and Pandey, S. (2010). Selective induction of apoptosis through activation of caspase-8 in human leukemia cells (Jurkat) by dandelion root extract. *J. Ethnopharmacol.* **133**(1):86–91.
- Owen, R. W., Giacosa, A., Hull, W. E., Haubner, R., Wurtele, G., Spiegelhalder, B. and Bartsch, H. (2000). Olive oil consumption and health: The possible role of antioxidants. *Lancet Oncol.* **1**(2):107–112.
- Park, L. K., Friso, S. and Choi, S. W. (2012). Nutritional influences on epigenetics and age-related disease. *Proc. Nutr. Soc.* **71**(1):75–83.
- Park, C. M., Youn, H. J., Chang, H. K. and Song, Y. S. (2010). TOP1 and 2, polysaccharides from *Taraxacum officinale*, attenuate CCl<sub>4</sub>-induced hepatic damage through the modulation of NF- $\kappa$ B and its regulatory mediators. *Food Chem. Toxicol.* **48**(5):1255–1261.
- Paxton, R. J., Garcia-Prieto, C., Berglund, M., Hernandez, M., Hajek, R. A., Handy, B., Brown, J. and Jones, L. A. (2012). A randomized parallel-group dietary study for stages II–IV ovarian cancer survivors. *Gynecol. Oncol.* **124** (3):410–416.
- Percival, S. S. (2009). Grape consumption supports immunity in animals and humans. *J. Nutr.* **139**(9):1801S–1805S.
- Perez-Vizcaino, F. and Duarte, J. (2010). Flavonols and cardiovascular disease. *Mol. Aspects Med.* **31**(6):478–494.

- Perona, J. S., Cabello-Moruno, R. and Ruiz-Gutierrez, V. (2006). The role of virgin olive oil components in the modulation of endothelial function. *J. Nutr. Biochem.* **17**(7):429–445.
- Pham, N. A., Jacobberger, J. W., Schimmer, A. D., Cao, P., Gronda, M. and Hedley, D. W. (2004). The dietary isothiocyanate sulforaphane targets pathways of apoptosis, cell cycle arrest, and oxidative stress in human pancreatic cancer cells and inhibits tumor growth in severe combined immunodeficient mice. *Mol. Cancer Ther.* **3**(10):1239–1247.
- Puertollano, M. A., Puertollano, E., Alvarez de Cienfuegos, G. and De Pablo Martínez, M. A. (2010). Olive oil, immune system and infection. *Nutr. Hosp.* **25**(1):1–8.
- Raederstorff, D. (2009). Antioxidant activity of olive polyphenols in humans: A review. *Int. J. Vitam. Nutr. Res.* **79**(3):152–165.
- Rahman, K. (2007). Effects of garlic on platelet biochemistry and physiology. *Mol. Nutr. Food Res.* **51**(11):1335–1344.
- Ramos-Romero, S., Pérez-Cano, F. J., Ramiro-Puig, E., Franch, A. and Castell, M. (2012). Cocoa intake attenuates oxidative stress associated with rat adjuvant arthritis. *Pharmacol. Res.* **66**(3):207–212.
- Ramprasath, V. R. and Jones, P. J. (2010). Anti-atherogenic effects of resveratrol. *Eur. J. Clin. Nutr.* **64**(7):660–668.
- Ried, K., Frank, O. R. and Stocks, N. P. (2010). Aged garlic extract lowers blood pressure in patients with treated but uncontrolled hypertension: A randomized controlled trial. *Maturitas.* **67**(2):144–150.
- Riso, P., Martini, D., Visioli, F., Martinetti, A. and Porrini, M. (2009). Effect of broccoli intake on markers related to oxidative stress and cancer risk in healthy smokers and nonsmokers. *Nutr. Cancer.* **61**(2):232–237.
- Romero, C., Medina, E., Vargas, J., Brenes, M. and De Castro, A. (2007). In vitro activity of olive oil polyphenols against *Helicobacter pylori*. *J. Agric. Food Chem.* **55**(3):680–686.
- Rotches-Ribalta, M., Andres-Lacueva, C., Estruch, R., Escribano, E. and Urpí-Sarda, M. (2012). Pharmacokinetics of resveratrol metabolic profile in healthy humans after moderate consumption of red wine and grape extract tablets. *Pharmacol. Res.* **66**(5):375–382.
- Salami, M., Galli, C., De Angelis, L. and Visioli, F. (1995). Formation of F2-isoprostanes in oxidized low density lipoprotein: Inhibitory effect of hydroxytyrosol. *Pharmacol. Res.* **31**(3–4):275–279.
- Schini-Kerth, V. B., Etienne-Selloum, N., Chataigneau, T. and Auger, C. (2011). Vascular protection by natural product-derived polyphenols: In vitro and in vivo evidence. *Planta Med.* **77**(11):1161–1167.
- Schütz, K., Kammerer, D. R., Carle, R. and Schieber, A. (2005). Characterization of phenolic acids and flavonoids in dandelion (*Taraxacum officinale* WEB. ex WIGG.) root and herb by high-performance liquid chromatography/electrospray ionization mass spectrometry. *Rapid Commun. Mass. Spectrom.* **19**(2):179–186.
- Sebastiano, C., Vincenzo, F., Tommaso, C., Giuseppe, S., Marco, R., Ivana, C., Giorgio, R., Massimo, M. and Giuseppe, M. (2012). Dietary patterns and prostatic diseases. *Front. Biosci. (Elite Ed)* **4**:195–204.
- Senanayake, G. V., Banigesh, A., Wu, L., Lee, P. and Juurlink, B. H. (2012). The dietary phase 2 protein inducer sulforaphane can normalize the kidney epigenome and improve blood pressure in hypertensive rats. *Am. J. Hypertens.* **25**(2):229–235.
- Shukla, Y. and George, J. (2011). Combinatorial strategies employing nutraceuticals for cancer development. *Ann. N. Y. Acad. Sci.* **1229**(1):162–175.
- Sigstedt, S. C., Hooten, C. J., Callewaert, M. C., Jenkins, A. R., Romero, A. E., Pullin, M. J., Kornienko, A., Lowrey, T. K., Slambrouck, S. V. and Steelant, W. F. (2008). Evaluation of aqueous extracts of *Taraxacum officinale* on growth and invasion of breast and prostate cancer cells. *Int. J. Oncol.* **32**(5):1085–1090.
- Simonetti, G. (1990). In: Schuler, S. ed. *Simon & Schuster's Guide to Herbs and Spices* (Nature Guide Series). Inc, New York. pp. 261–262.
- Singh, M., Singh, P. and Shukla, Y. (2012). New strategies in cancer chemoprevention by phytochemicals. *Front. Biosci. (Elite Ed)* **4**:426–452.
- Singh, A. V., Xiao, D., Lew, K. L., Dhir, R. and Singh, S. V. (2004). Sulforaphane induces caspase-mediated apoptosis in cultured PC-3 human prostate cancer cells and retards growth of PC-3 xenografts *in vivo*. *Carcinogenesis.* **25**(1):83–90.
- Sobenin, I. A., Pryanishnikov, V. V., Kunnova, L. M., Rabinovich, Y. A., Martirosyan, D. M. and Orekhov, A. N. (2010). The effects of time-released garlic powder tablets on multifunctional cardiovascular risk in patients with coronary artery disease. *Lipids Health Dis.* **9**:119.
- Sohn, D. W., Han, C. H., Jung, Y. S., Kim, S. I., Kim, S. W. and Cho, Y. H. (2009). Anti-inflammatory and antimicrobial effects of garlic and synergistic effect between garlic and ciprofloxacin in a chronic bacterial prostatitis rat model. *Int. J. Antimicrob. Agents.* **34**(3):215–219.
- Speciale, A., Chirafisi, J., Saija, A. and Cimino, F. (2011). Nutritional antioxidants and adaptive cell responses: An update. *Curr. Mol. Med.* **11**(9):770–789.
- Steevens, J., Schouten, L. J., Goldbohm, R. A. and Van den Brandt, P. A. (2011). Vegetables and fruits consumption and risk of esophageal and gastric cancer subtypes in the Netherlands Cohort Study. *Int. J. Cancer.* **129**(11):2681–2693.
- Steiner, M. and Lin, R. S. (1998). Changes in platelet function and susceptibility of lipoproteins to oxidation associated with administration of aged garlic extract. *J. Cardiovasc. Pharmacol.* **31**(6):904–908.
- Tanaka, S., Haruma, K., Kunihiro, M., Nagata, S., Kitadai, Y., Manabe, N., Sumii, M., Yoshihara, M., Kajiyama, G. and Chayama, K. (2004). Effects of aged garlic extract (AGE) on colorectal adenomas: A double-blinded study. *Hiroshima J. Med. Sci.* **53**(3–4):39–45.
- Tanito, M., Masutani, H., Kim, Y. C., Nishikawa, M., Ohira, A. and Yodoi, J. (2005). Sulforaphane induces thioredoxin through the antioxidant responsive element and attenuates retinal light damage in mice. *Invest. Ophthalmol. Vis. Sci.* **46**(3):979–987.
- Traka, M., Gasper, A. V., Melchini, A., Bacon, J. R., Needs, P. W., Frost, V., Chantry, A., Jones, A. M. E., Ortori, C. A., Barrett, D. A., Ball, R. Y., Mills, R. D. and Mithen, R. F. (2008). Broccoli consumption interacts with GSTM1 to perturb oncogenic signaling pathways in the prostate. *PLoS ONE* **3**(7):e2568.
- Traka, M. H. and Mithen, R. F. (2011). Plant science and human nutrition: Challenges in assessing health-promoting properties of phytochemicals. *Plant Cell.* **23**(7):2483–2497.
- Trojanová, I., Rada, V., Kokoska, L. and Vlková, E. (2004). The bifidogenic effect of *Taraxacum officinale* root. *Fitoterapia.* **75**(7–8):760–763.
- Tsao, R. (2010). Chemistry and biochemistry of dietary polyphenols. *Nutrients.* **2**(12):1231–1246.
- Tsubura, A., Lai, Y. C., Kuwata, M., Uehara, N. and Yoshizawa, K. (2011). Anticancer effects of garlic and garlic-derived compounds for breast cancer control. *Anticancer Agents. Med. Chem.* **11**(3):249–253.
- Vasanthi, H. R., Mukherjee, S. and Das, D. K. (2009). Potential health benefits of broccoli — a chemico-biological overview. *Mini. Rev. Med. Chem.* **9**(6):749–759.
- Vasanthi, H. R., ShriShriMal, N. and Das, D. K. (2012). Phytochemicals from plants to combat cardiovascular disease. *Curr. Med. Chem.* **19**(14):2242–2251.
- Visioli, F. (2012). Olive oil phenolics: Where do we stand? Where should we go? *J. Sci. Food Agric.* **92**(10):2017–2019.
- Visioli, F. and Bernardini, E. (2011). Extra virgin olive oil's polyphenols: Biological activities. *Curr. Pharm. Des.* **17**(8):786–804.
- Visioli, F., Poli, A. and Gall, C. (2002). Antioxidant and other biological activities of phenols from olives and olive oil. *Med. Res. Rev.* **22**(1):65–75.
- Vislocky, L. M. and Fernandez, M. L. (2010). Biomedical effects of grape products. *Nutr. Rev.* **68**(11):656–670.
- Vita, J. A. (2005). Polyphenols and cardiovascular disease: Effects on endothelial and platelet function. *Am. J. Clin. Nutr.* **81**(Suppl 1):292S–297S.
- Wang, J., Ferruzzi, M. G., Ho, L., Blount, J., Janle, E. M., Gong, B., Pan, Y., Gowda, G. A. N., Raftery, D., Arrieta-Cruz, I., Sharma, V., Cooper, B., Lobo, J., Simon, J. E., Zhang, C., Cheng, A., Qian, X., Ono, K., Teplow, D. B., Pavlides, C., Dixon, R. A. and Pasinetti, G. M. (2012). Brain-targeted proanthocyanidin metabolites for Alzheimer's disease treatment. *J. Neurosci.* **32**(15):5144–5150.
- Waterman, E. and Lockwood, B. (2007). Active components and clinical applications of olive oil. *Altern. Med. Rev.* **12**(4):331–342.
- Weisburger, J. H. (2001). Chemopreventive effects of cocoa polyphenols on chronic diseases. *Exp. Biol. Med. (Maywood)* **226**(10):891–897.

- Weseler, A. R., Ruijters, E. J., Driessens, M. J., Reesink, K. D., Haenen, G. R. and Bast, A. (2011). Pleiotropic benefit of monomeric and oligomeric flavanols on vascular health: A randomized controlled clinical pilot study. *PLoS One*. **6**(12):e28460.
- Williams, C. A., Goldstone, F. and Greenham, J. (1996). Flavonoids, cinnamic acids and coumarins from the different tissues and medicinal preparations of *Taraxacum officinale*. *Phytochemistry*. **42**(1):121–127.
- Wu, C. D. (2009). Grape products and oral health. *J. Nutr.* **139**(9): 1818S–1823S.
- Xia, E. Q., Deng, G. F., Guo, Y. J. and Li, H. B. (2010). Biological activities of polyphenols from grapes. *Int. J. Mol. Sci.* **11**(2):622–646.
- Yoon, T. J. (2008). Effect of water extracts from root of *Taraxacum officinale* on innate and adaptive immune responses in mice. *Korean J. Food Nutr.* **21**(3):275–282.
- You, Y., Yoo, S., Yoon, H. G., Park, J., Lee, Y. H., Kim, S., Oh, K. T., Lee, J., Cho, H. Y. and Jun, W. (2010). *In vitro* and *in vivo* hepatoprotective effects of the aqueous extract from *Taraxacum officinale* (dandelion) root against alcohol-induced oxidative stress. *Food Chem. Toxicol.* **48**(6):1632–1637.
- Zhang, J., Kang, M. J., Kim, M. J., Kim, M. E., Song, J. H., Lee, Y. M. and Kim, J. I. (2008). Pancreatic lipase inhibitory activity of *Taraxacum officinale* *in vitro* and *in vivo*. *Nutr. Res. Pract.* **2**(4):200–203.
- Zhang, Y., Talalay, P., Cho, C. G. and Posner, G. H. (1992). A major inducer of anti-carcinogenic protective enzymes from broccoli: Isolation and elucidation of structure. *Proc. Natl. Acad. Sci. (USA)* **89**(6):2399–2403.
- Zhao, L., Lee, J. Y. and Hwang, D. H. (2011a). Inhibition of pattern recognition receptor-mediated inflammation by bioactive phytochemicals. *Nutr. Rev.* **69**(6):310–320.
- Zhao, D., Qi, Y., Zheng, Z., Wang, Y., Zhang, X. Y., Li, H. J., Liu, H. H., Zhang, X. T., Du, J. and Liu, J. (2011b). Dietary factors associated with hypertension. *Nat. Rev. Cardiol.* **8**(8):456–465.
- Zhi, X., Honda, K., Ozaki, K., Misugi, T., Sumi, T. and Ishiko, O. (2007). Dandelion T-1 extract up-regulates reproductive hormone receptor expression in mice. *Int. J. Mol. Med.* **20**(3):287–292.
- Zhou, D. Y., Du, Q., Li, R. R., Huang, M., Zhang, Q. and Wei, G. Z. (2011). Grape seed proanthocyanidin extract attenuates airway inflammation and hyperresponsiveness in a murine model of asthma by down regulating inducible nitric oxide synthase. *Planta Med.* **77** (14):1575–1581.
- Zunino, S. (2009). Type 2 diabetes and glycemic response to grapes or grape products. *J. Nutr.* **139**(9):1794S–1800S.
- [No authors listed]. (2010). Sulforaphane glucosinolate. *Monograph. Altern. Med. Rev. Dec*; **15**(4):352–360. Review. [PMID: 21194251].