



Alternative yeasts for winemaking: *Saccharomyces non-cerevisiae* and its hybrids

Roberto Pérez-Torrado, Eladio Barrio & Amparo Querol

To cite this article: Roberto Pérez-Torrado, Eladio Barrio & Amparo Querol (2017): Alternative yeasts for winemaking: *Saccharomyces non-cerevisiae* and its hybrids, Critical Reviews in Food Science and Nutrition, DOI: [10.1080/10408398.2017.1285751](https://doi.org/10.1080/10408398.2017.1285751)

To link to this article: <http://dx.doi.org/10.1080/10408398.2017.1285751>



Accepted author version posted online: 31 Mar 2017.



Submit your article to this journal [↗](#)



View related articles [↗](#)



View Crossmark data [↗](#)

S. non-cerevisiae in winemaking

Alternative yeasts for winemaking: *Saccharomyces non-cerevisiae* and its hybrids

Roberto Pérez-Torrado^{a,b}, Eladio Barrio^{a,b} and Amparo Querol^{a,*}

^aInstituto de Agroquímica y Tecnología de los Alimentos, IATA-CSIC. E-46980 Paterna, Spain

^bDepartament de Genètica, Universitat de València, Valencia, Spain

*Corresponding author: Departamento de Biotecnología, Instituto de Agroquímica y Tecnología de los Alimentos (CSIC). E-46980 Paterna, (Valencia), Spain, E-mail: aquerol@iata.csic.es (Amparo Querol)

Abstract

Wine fermentation has not significantly changed since ancient times and the most traditional aspects are seen by the market as elements that uplift wine nuances and quality. In recent years, new trends have emerged from the sector in line with consumer preferences, and due to the effects of global climate change on grape ripening. In the first cases the consumers are looking for wines with less ethanol and fruitier aromas and in the second cases the wineries want to reduce the wine alcohol levels and/or astringency. New yeast starters of alternative *Saccharomyces* species and their hybrids can help to solve some problems that wineries face. In this article we review several physiological and genetic aspects of *S. uvarum* and *S. kudriavzevii* and the hybrids, which are especially relevant during the winemaking process, such as their good fermentative capabilities at low temperatures, resulting in wines with lower alcohol and higher glycerol amounts.

Keywords

Winemaking; yeasts; *S. non-cerevisiae*; aroma; glycerol; cold fermentation

INTRODUCTION

Winemaking is one of the oldest food processing procedures that humans have performed since ancient civilizations. In fact, the earliest evidence of winemaking is traced to Iran at the Hajji Firuz Tepe site (5400-5000 BC) (This et al., 2006). The process of wine production is a conversion of grape juice made by microorganisms, mainly yeasts, where a huge number of compounds are metabolically consumed or produced. The result is one of the most widely consumed beverages. In the modern wine production process, selected yeasts are used to inoculate grape must in order to control the fermentation, reduce risk of contamination, increase the reproducibility and generate specific characteristics in the wine by selecting yeasts strains with specific abilities of fermentation. Only specific wine strains are able to outcompete and impose over other non-wine and spoilage microorganism and ensure a proper wine fermentation. Yeast starters companies have enormous interest in expand their strain catalogue with the objective to present wine strains with general wine strain type characteristics abut also with specific abilities that can be useful for wineries.

Wine yeasts are highly specialised organisms that evolved under restrictive environmental conditions created by human technology. During cellular adaptation to those manipulated environments, yeast strains were subjected to selective pressures generated by alcoholic fermentation (Querol et al., 2003), resulting in adaptative differences among them (Barrio et al., 2006). Most of commercial yeasts are *S. cerevisiae*, being the most frequently used in wine fermentations, as well as the most studied species. However other species of the *Saccharomyces* genus and even other species of the genus *Saccharomyces* (*S. uvarum*) and their

interspecific hybrids (*S. cerevisiae* x *S. uvarum* and *S. cerevisiae* x *S. kudriavzevii*) have shown their potential application to solve the new challenges of the winemaking industry, thus attracting high interest in last years by researchers in the field (González et al., 2007; Gangl et al., 2009; Tronchoni et al., 2009; Gamero et al., 2013). In addition, strain selection has been extended in recent years to non-*Saccharomyces* yeasts such as those belonging to genera *Candida*, *Kloeckera*, *Kluyveromyces*, *Debaryomyces*, *Hanseniaspora*, *Hansenula*, *Pichia*, *Metschnikowia*, *Schizosaccharomyces*, *Saccharomycodes*, *Starmerella*, *Torulaspora* or *Rhodotorula*. Although non-*Saccharomyces* species lack competitiveness under oenological conditions, mainly because they do not ferment so vigorously and display lower stress resistance than *Saccharomyces*, employing mixed starter cultures or sequential fermentations are using in the last years to enhanced glycerol, improve the aroma complexity, reduced acetic acid and ethanol content (Schuller and Casal, 2005; Comitini et al., 2011; Milanovic et al., 2012; Contreras et al., 2015; Morales et al., 2015; Lencioni et al., 2016; Ciani et al., 2016 a and b; Varela, 2016; Wang et al., 2016). Besides the scientific interest for the use of these species, only strains of the species *Kluyveromyces thermotolerans*, *Metschnikowia pulcherrima*, *Pichia kluyveri*, *Torulaspora delbrueckii* are commercialized.

One of the most important challenges in winemaking is the modification of the composition and properties of the grape must due to climate change (Borneman et al., 2013). Answers to these new demands require improvements in the enological practices, among which the development of new yeast starters adapted to the fermentation conditions imposed by climate change are of chief importance. Climate change impact on the winemaking industry has stress out the importance of ethanol yield in the fermentation (White et al., 2006). Temperature

increases force early maturation of grapes in Mediterranean regions and sugar contents in musts are higher and, as a consequence, alcohol contents in the resulting wine increase (Jones et al., 2005). An alternative is to harvest grapes earlier, when sugar content is optimum to obtain wines with the appropriated alcohol content. However, in this case, the ripening of balanced fruits becomes more difficult because the maturation stage of grape tannins and phenols is not reached and the outcome is very astringent wine. These wines with high ethanol contents or with unpleasant young unripe tannins are not well accepted by consumers. To avoid this lack of competitiveness, wineries demand yeasts exhibiting lower ethanol yields as well as higher glycerol and mannoprotein productions, because these compounds help to balance wine astringency (White et al., 2006).

New yeast starters, of non-*cerevisiae* strains (*S. uvarum*) or hybrids (*S. cerevisiae* x *S. uvarum* and *S. cerevisiae* x *S. kudriavzevii*) can contribute to solve some problems of the wineries related to climate change. Contrary to what happens with non-*Saccharomyces* species that have previously described, *S. uvarum* and hybrids among species of the genus *Saccharomyces* exhibit good fermentative capabilities at low temperatures, producing wines with lower alcohol and higher glycerol amounts, while fulfilling the requirements of the commercial yeasts, such as a good fermentative performance and aromatic profiles, of great interest for the wine industry (González et al., 2007; Tosi et al., 2009; Gamero et al., 2012). Moreover, some strains exhibit pectinolytic activity (Naumov et al., 2001) and, hence, produce more aromatic wines (Henschke and Rose, 1991).

Another important driving force in winemaking industry is the consumer's demands for new products with novel aromatic profiles. Nowadays, fermentations at low temperatures are a trend in winemaking because it improves the aromatic profile of the produced wines (Boulton et al., 1996). However, low fermentation temperatures have disadvantages because requires longer periods and the risks of halted or sluggish fermentations are bigger (Bisson, 1999). These problems can be avoided by providing yeasts better adapted to ferment at low temperatures.

BENEFICIAL ASPECTS OF *S. UVARUM*, *S. KUDRIAVZEVII* AND *SACCHAROMYCES HYBRIDS* FOR WINEMAKING

Low fermentation temperatures conducted by *S. cerevisiae* have disadvantages because longer processes are required, and the risks of halted or sluggish fermentations increase (Boulton et al., 1996; Bisson, 1999). These problems can be avoided with yeasts better adapted to ferment at low temperatures. Although supposedly “cryotolerant” *S. cerevisiae* yeasts are already available for the wine industry, e.g. QA23 from Lallemant Inc. or Fermol Cryophile and Fermol Reims Champagne from AEB, most of them do not offer desirable fermentation performances at low temperatures (10-15°C). Physiological and enological works have indicated the potential benefits of the use of *S. uvarum*, *S. kudriavzevii* and their hybrids with *S. cerevisiae* fermenting at low temperature (Figure 1) and have also shown its well-established cryotolerant character (Tronchoni et al., 2012; Gamero et al., 2013; López-Malo et al., 2013). As *S. uvarum*, *S. kudriavzevii* and hybrids strains are able to finish fermentations in musts with 250 g/L sugars, producing lower alcohol levels and increased glycerol content without increasing acetic acid, especially at low temperatures (González et al., 2006; González et al., 2007; Lopandic et al.,

2007; Tronchoni et al., 2009; Masneuf-Pomarede et al., 2010; Paget et al., 2014) which makes this species suitable to counteract problems derived from climate change. Furthermore, its sugar consumption rate, similar to that of *S. cerevisiae*, could make this organism a good candidate to compete for a place at low fermentation temperatures in the wine yeast industry (Figure 1).

S. kudriavzevii has been never found in winemaking probably due to low resistance to elevated concentrations of ethanol (Belloch et al., 2008), although its contribution is very relevant as a part of the *Saccharomyces* hybrids. However, all the published data has been obtained at the laboratory level, so the possibility of using *S. kudriavzevii* in wine fermentations must be tested at the industrial scale. There are some physiological properties of *S. kudriavzevii* that could explain the potential benefits in the use of *S. kudriavzevii* in winemaking as the ability to perform complete wine fermentations at laboratory scale with similar or even enhanced kinetics than *S. cerevisiae* (Gonzalez et al., 2006; Combina et al., 2012). On the other hand, other works have shown that *S. kudriavzevii* cannot outperform *S. cerevisiae* strains even at low temperature in synthetic media growth (Arroyo-López et al., 2011). Thus, the study of interaction among both species can be important to explain the absence of *S. kudriavzevii* in wine fermentations conditions. In spite of these results, is not clear that industrial wine fermentations inoculated with *S. kudriavzevii* selected strains will be completed by them or they will be competed out at some point by *S. cerevisiae* contaminant cells, which are present in different parts of the wineries. Thus, further research at industrial scale must be conducted to finally implement the use of the potentially beneficial species *S. kudriavzevii* in winemaking.

So far, it is known that *S. cerevisiae* and *S. kudriavzevii* natural hybrids are able to dominate industrial wine fermentations performed at low temperatures in regions of cold climates (Gonzalez et al., 2006; Gang et al., 2009; Erny et al., 2012). On the other hand, mixed cultures between *S. kudriavzevii* and *S. cerevisiae* auxotrophic mutants performed in synthetic must showed a clear *S. cerevisiae* imposition at high temperatures (31 °C) (Arroyo-López et al., 2011). But it is interesting to note that at low temperatures (17 °C), *S. kudriavzevii* is able to compete with *S. cerevisiae* in the first part of the fermentation when they are inoculated 50:50. This suggests that in controlled industrial wine fermentations at low temperatures, which contains very low levels of *S. cerevisiae* cells coming from grape grains and winery equipment, the inoculation of *S. kudriavzevii* could contribute significantly to the final wine characteristics, by increasing glycerol production and enhancing its aroma profile (Figure 1) (Gonzalez et al., 2007; Tronchoni et al., 2009; Arroyo-López et al., 2011).

In the next sections we will analyze the molecular mechanisms related with the adaptation of *S. uvarum* and *S. kudriavzevii* to growth at low temperatures and the differences in the contribution to the aroma profile with respect to *S. cerevisiae* (Figure 1).

Adaptation to low temperatures

Several studies have focused in understand the cryophilic character of *S. uvarum* and *S. kudriavzevii* at the molecular level, including transcriptomic and metabolomic studies (Combina et al., 2012; López-Malo et al., 2013). Some aspects of these species have been highlighted in relation to cold resistance and winemaking as glycerol accumulation (Arroyo-López et al., 2010;

Oliveira et al., 2014), membrane composition (Tronchoni et al., 2012), or translation efficiency (Tronchoni et al., 2014) and are addressed with more detail below.

Glycerol accumulation

The cryophilic species *S. uvarum* and *S. kudriavzevii* produce higher amounts of glycerol compared to *S. cerevisiae* in must fermentations (Arroyo-López et al., 2010; Oliveira et al., 2014). Glycerol contributes to wine quality by generating sweetness and fullness (Ribereau-Gayon et al., 1977; Eustace and Thornton, 1987). Also increase viscosity which can soften the astringency caused by unripened tannins present in early harvested grapes (Ishikawa and Noble, 1995, Llaudy et al., 2006). In addition, glycerol plays an important role in cold stress adaptation (Tulha et al., 2010), although other features have been related to yeasts adaptation to low temperatures, including synthesis of ribosomal proteins, changes in membrane lipid composition and a higher trehalose content (Aguilera et al., 2007). In *S. cerevisiae*, glycerol is synthesized after a cold shock via a regulatory mechanism involving the HOG (High Osmolarity Glycerol) pathway (Hayashi and Maeda, 2006; Panadero et al., 2006). Intracellular glycerol content was linked to cell survival in fermentations at low temperatures (Tulha et al., 2010) and also is involved in freeze/thawing stress resistance (Izawa et al., 2004).

Cryoprotectant glycerol is synthesized by a branch of glycolysis involving two steps (Ansell et al., 1997; Norbeck et al., 1997; Pahlman et al., 2001), which in the case of *Saccharomyces* yeasts are catalyzed by two isoenzymes each step: GPD for glycerol-3-phosphate dehydrogenases (Gpd1p and Gpd2p) and GPP for glycerol-3-phosphatases (Gpp1p/Rhr1p and Gpp2p/Hor2p). Flux modelling calculations of metabolic control values of glycerol synthesis

indicate that the glycerol-3-phosphate dehydrogenase reaction has a flux control coefficient of approximately 0.85, being the major responsible of the flux control of this pathway (Remize et al., 2001). Moreover, *GPD1* gene overexpression increases the glycerol levels produced while the overexpression of the other three enzymes does not (Nevoight and Stahl, 1996; Pahlman et al., 2001; Remize et al., 2001) whereas a reduction of *GPD1* expression leads to a reduced flux towards glycerol (Nevoight and Stahl, 1996; Hubmann et al., 2011). Furthermore, *GPD1* is activated in response to cold stress (Panadero et al., 2006). All this data supported the important role of *GPD1* in the cold stress response in *Saccharomyces*.

Recently, our research focused in deciphering the mechanisms by which the cryophilic species *S. kudriavzevii* produce more glycerol and its putative role in the better adaptation of this species to cold environments (Figure 1). Interestingly, it has been shown that *S. kudriavzevii* has a higher *GPD1* expression and that Gpd1p presents enhanced enzymatic parameters and an increased activity (Oliveira et al., 2014). All these differences suggest that *S. kudriavzevii* has changed its metabolism to promote the branch of the glycolytic pathway involved in the glycerol production to adapt to low temperature environments (Figure 1) and maintain the NAD^+/NADH ratio in alcoholic fermentations. In agreement with these results, new studies suggested that temperature-induced redox imbalances could be compensated by increasing glycerol accumulation but also by the production of cytosolic acetaldehyde through the deletion of *GUT2* or *ADH3* respectively (Paget et al., 2014).

Membrane lipid composition

As previously indicated, the lipid composition of *S. kudriavzevii* exhibits some features that may explain its adaptation to low temperatures (Tronchoni et al., 2012). Yeast membranes are a lipid bilayer consisting of two amphipathic molecules, phospholipids (PL) and sphingolipids, with an apolar phase of fatty acids (FA). The fatty acids of *S. cerevisiae* mainly correspond both to unsaturated fatty acids (UFA), oleic acid and palmitoleic acid, and saturated fatty acids (SFA), palmitic acid and stearic acid. The medium chain fatty acids (MCFA) are found in lower proportions but their concentrations increase when yeasts are grown under anaerobic conditions, such as the wine fermentation. Sterols are also important, being ergosterol the main sterol in fungi (Henschke and Rose, 1991). Changes in any of these lipids can significantly disturb the membrane function and alter the role of membrane-associated proteins. Thus, many organisms have developed mechanisms to keep the appropriate membrane fluidity regardless of the temperature (Cossins and Prosser, 1978). It has been described that low temperatures increase in the degree of fatty acid (FA) unsaturation in *S. cerevisiae* (Sakamoto and Murata, 2002), producing a higher membrane fluidity. Another way to increase membrane fluidity is to reduce the FA chain length (Torija et al., 2003). Beltran et al. (2006) reported some of these effects in *S. cerevisiae* strains during an industrial wine fermentation. Redón et al. (2011) also reported common changes in the lipid composition of different industrial species and strains of *Saccharomyces* after growth at low temperature showing that the MCFA and triacylglyceride content increased, whereas the phosphatidic acid content and the phosphatidylcholine/phosphatidylethanolamine ratio decreased, as occurs in *S. uvarum* compared to *S. cerevisiae* (Figure 1). A similar response was observed in *S. kudriavzevii*, although sterol esters and squalene were also increased (Tronchoni et al., 2012). These results indicate that the

membrane of *S. kudriavzevii* has a composition that allows its adaptation to growth at low temperatures due to two mechanisms already observed in the other species (Figure 1). Regardless the growth temperature, the *S. kudriavzevii* strains had higher medium-chain fatty acids and squalene percentages and also shorter chain lengths (Tronchoni et al., 2012). This differential lipid content has a significant impact in the enhanced fitness of *S. kudriavzevii* at low temperatures. However, more research must be done in order to clarify the role of the different membrane components in the physiology of these yeasts.

Translation efficiency

At low temperatures, ribosome maturation may be compromised due to a reduction in RNA structure plasticity by RNA hyperstabilization (Zavanelli et al., 1994; Fortner et al., 1994; Li and Brow, 1996; Staley and Guthrie, 1999; Hilliker et al., 2007; Perriman and Ares, 2007; Kurata et al., 2013). A transcriptome comparison of *S. kudriavzevii* with *S. cerevisiae* shed light on the response of this cryotolerant yeast to low temperature (Tronchoni et al., 2014). A similar low temperature stress response for both species is the presence of up-regulated genes related to translational machinery, although *S. kudriavzevii* shows a higher response compared to *S. cerevisiae*. Tronchoni et al (2014) postulated that this higher response could be the result of changes in the stability of a functional RNA conformation in relation to a competing structure (Zavanelli et al., 1994). However, the different paromomycin susceptibility (Tronchoni et al., 2012) suggests that other mechanisms can also be involved related to *S. kudriavzevii* translation machinery at low temperatures. Also, paromomycin resistance can be the result of enhanced translation efficiency (Perriman and Ares, 2007). This study suggests that *S. kudriavzevii* has

increased translation efficiency due to higher ribosome availability after adaptation to cold shock (Figure 1) (Tronchoni et al., 2012). García-Rios et al (2016) analyzing the proteome profiling of *S. cerevisiae* and cryotolerant species *S. uvarum* and *S. kudriavzevii* during low-temperature wine fermentation we by iTRAQ-based showed that the main differences among the proteomic profiling of the three *Saccharomyces* strains grown at 12 °C and 28 °C lay in translation, glycolysis and amino acid metabolism. These data corroborate previous transcriptomic results (Tronchoni et al., 2014) , which suggest again that *S. kudriavzevii* is better adapted to grow at low temperature as a result of enhanced more efficient translation.

Metabolic balance

Although no similar transcriptomic data is available for *S. uvarum*, the metabolome comparison of *S. cerevisiae*, *S. uvarum* and *S. kudriavzevii* growing at 12 °C revealed that the main differences between the two cryotolerant species and *S. cerevisiae* were in carbohydrate metabolism, mainly fructose. However, these two species have developed different strategies for cold resistance. *S. uvarum* presented elevated shikimate pathway activity, while *S. kudriavzevii* displayed increased NAD⁺ synthesis (López-Malo et al., 2013). García-Rios et al (2016) analyzing the proteome profiling of these species and comparing with *S. cerevisiae* observed that amino acid biosynthetic pathways can also be mechanisms that better explain biomass yield in cryotolerant strains and also can justify the differences in the aroma synthesis. However, at low temperature, *S. cerevisiae* showed higher concentrations of glycolytic and alcoholic fermentation enzymes.

Contribution to the wine aroma

Wine fermentation aromas result from a complex mixture of chemical compounds produced by yeast secondary metabolism (Lilly et al., 2000; Swiegers and Pretorius, 2005). Higher alcohols (fusel, marzipan and floral aromas) as well as acetate and ethyl esters (fruity and floral aromas) are the main scents in young wines. Higher alcohols are generated from sugar metabolism intermediates, through anabolic reactions, or from branched-chain amino acids, through the Ehrlich pathway, a multistep catabolic reaction (Boulton et al., 1996; Dickinson et al., 1997; Eden et al., 2001; Dickinson et al., 2003). Ethyl ester compounds are produced by condensation of alcohols and coenzyme-A-activated acids (Swiegers and Pretorius, 2005), while acetate esters result from the combination of acetyl-CoA with an alcohol, by the action of alcohol acetyl transferases (Lilly et al., 2000). The amount and nature of these aroma compounds are influenced by many factors, such as must nitrogen content, fermentation temperature and the yeast strain (Lilly et al., 2000; Swiegers and Pretorius, 2005). Several studies have characterized the impact of *S. uvarum* and *S. kudriavzevii* on the fermentation processes and the by-product formation and aroma profile (Gonzalez et al., 2007; Henschke et al., 2000; Gamero et al., 2011a; Gamero et al., 2011b; Gamero et al., 2012; Gamero et al., 2013). These studies showed that *S. uvarum* and *S. kudriavzevii* strains present different aroma compound accumulation patterns during wine fermentation compared to those of *S. cerevisiae* strains (Figure 1). Hence, the utilization of *S. uvarum* and *S. kudriavzevii* strains in winemaking can contribute to a new aromatic composition in the final wines, by increasing the higher alcohol and ester amounts. Specifically, *S. uvarum* generates larger amounts of 2-phenylethanol, 2-phenylethyl acetate and ethyl lactate (Henschke et al., 2000; Masneuf-Pomarede et al., 2010; Gamero et al., 2011a; Gamero et al., 2012) and *S. kudriavzevii* produce larger amounts of higher alcohols and 2-phenylethanol at 12°C (Henschke

et al., 2000; Gamero et al., 2011a; Gamero et al., 2013; Stribny et al., 2015). These differences in aroma production have been correlated with differences in gene sequences and in gene regulations. For example, *S. kudriavzevii* strains exhibited an up-regulation in acyltransferase *EHT1* and a down-regulation in acyltransferase *EEB1* (the genes involved in ethyl esters formation) during fermentation at 28°C (Gamero et al., 2014). The low ethyl ester production in *S. kudriavzevii* also suggests that acyltransferase *EEB1* is more important in the production of these aromatic compounds than *EHT1*, as previously described (Saerens et al., 2008). Also it is interesting to note that the higher level of geraniol uptake, which is then bioconverted into linalool and alpha-terpineol, is one of the most significant abilities of *S. kudriavzevii* among others (Gamero et al., 2011b). By comparative genomic analysis, among the available genome sequences of *S. cerevisiae*, *S. uvarum* and *S. kudriavzevii*, of genes involved in aroma synthesis, Stribny et al (2015) observed important differences at nucleotide level in the genes *ARO10* (encoding a phenylpyruvate decarboxylase), *ATF1* and *ATF2* (coding for alcohol acetyltransferases 1 and 2, respectively). The heterologous expression of these genes from *S. kudriavzevii* and *S. uvarum* in 3 deletant *S. cerevisiae* T73 strains ($\Delta ATF1$, $\Delta ATF2$, and $\Delta ARO10$ strains) confirmed the higher production of several aromatic compounds by *S. kudriavzevii* and *S. uvarum* such as isoamyl alcohol, isobutanol and their esters (Stribny et al., 2016a and 2016b). As occurred with Gpd1p (Oliverira et al., 2014), it has also been observed interesting differences in enzymatic activities between these proteins in the cases of *S. kudriavzevii* and *S. uvarum*. The enzymatic activity of SkAro10p with phenylpyruvate as a substrate was half that ScAro10p whereas the activities of SkAro10p for the other tested substrates were more than three times higher. Differences in the enzymatic properties of ScAtf1p, SkAtf1p and SuAtf1p were also

detected (Stribny et al., 2016a). Hence, the utilization of *S. uvarum*, *S. kudriavzevii* or their hybrids strains in winemaking can contribute to a new aromatic composition in the final wines.

THE ROLE OF *S. CEREVISIAE* HYBRIDS IN WINEMAKING

In the case of *Saccharomyces* genus, one of the most interesting mechanisms observed in the adaptation of these yeasts to industrial process is the formation of interspecific hybrids. Allopolyploidy and introgression by interspecific hybridization are the main mechanisms of lateral gene transfer in eukaryotes (de Barros Lopes et al., 2002; Barrio et al., 2006). Interspecific hybridization generates new gene combinations of potential adaptive value conferring, under fluctuating or intermediate environmental conditions, selective advantages to the hybrids with respect to their parental species (Masneuf et al., 1998). Hybrids between *S. cerevisiae* and other *Saccharomyces* species such as the cryotolerant *S. uvarum* (Naumov et al., 2000; Greig et al., 2002; Le Jeune et al., 2007) and *S. kudriavzevii* (Bradbury et al., 2006; González et al 2006; Lopandic et al., 2007; Erny et al., 2012; Peris et al., 2012a) have been isolated from wine, cider and brewing fermentations, and other sources.

On the origin of *S. kudriavzevii* wine hybrids

In recent years, hybrids between *S. cerevisiae* and *S. kudriavzevii* have been isolated not only from wine, but also from cider and brewing fermentations, as well as other sources (Bradbury et al., 2006; González et al 2006; Lopandic et al., 2007; Erny et al., 2012; Peris et al., 2012a) suggesting interesting human interactions in their origin and propagation (Peris et al., 2012b). Researchers have looked for *S. kudriavzevii* wine hybrids worldwide but they were only isolated in specific locations in Europe (Figure 2). By combining the phylogenetic analysis of gene

sequences with all the available information on genetic and genomic characterization of *S. cerevisiae* x *S. kudriavzevii* hybrids a total of seven potential hybridization events were predicted as the *S. kudriavzevii* wine hybrids origins (Peris et al., 2012b). Most hybrids seem to have been generated by rare-mating events involving a diploid *S. cerevisiae* strain and a haploid strain of *S. kudriavzevii* generating different chimerical genomes with ploidy values close to 3n (Table 1) (Peris et al., 2012a; Peris et al., 2012b). Two of these hybridization events generated the most frequent wine hybrid types (Figure 2), one was found in Wädenswill, Switzerland, and is phylogenetically related to Trappist brewing hybrids. The other type is widely distributed from the Rhine valley (Alsace and Germany) to the Danube valley (Austria, Croatia and Hungary) wine regions.

After ~6,000 years of adaptation to wine fermentation conditions in warm climates of the Mediterranean and Fertile Crescent regions, wine *S. cerevisiae* yeasts were taken by the Romans, together with the vines and winemaking tools, to the limits of their empire, the Rhine and Danube Rivers. In these regions of Oceanic and Continental climates (Figure 2), these *S. cerevisiae* wine strains could have problems to perform wine fermentations at low temperatures at which other *Saccharomyces* species are better adapted. In such climate conditions, however, hybrids have clear advantages over the parental species.

These *S. cerevisiae* x *S. kudriavzevii* hybrids likely originated several times but they probably spread during the Middle Age when Christian monks spread the viticulture and enology practices all over Europe (Figure 2) (Burton and Kerr, 2011). For example, in Central Europe Cistercian monks extended the Burgundian family of grape varieties, mainly Chardonnay and

Pinots, as well as German varieties, and with them also the main lineage of hybrid yeasts responsible of wine fermentation (Erny et al., 2012; Peris et al., 2012a; Peris et al., 2012b).

Molecular characterization of enological *Saccharomyces* hybrids

At the moment, very few information about the physiological and molecular part of *S. uvarum* hybrids are available. However, some more studies have been done on the *S. kudriavzevii* hybrids. Oenological characterization of *S. cerevisiae* and *S. uvarum* and *S. cerevisiae* x *S. kudriavzevii* hybrid strains has demonstrated that the hybrids are well adapted to ferment at low and intermediate temperatures, producing moderate or higher levels of glycerol and less acetic acid and more aromas (higher alcohols and esters) with regard to reference strains of *S. cerevisiae* and *S. kudriavzevii* (Masneuf et al., 1998; Gamero et al., 2013; González et al., 2007).

Natural hybrids between *S. cerevisiae* and *S. kudriavzevii* conducting wine fermentations have been characterized by genetic approaches (Belloch et al., 2008; Peris et al., 2012a; Peris et al., 2012b, Borneman et al., 2013). It is interesting to note that some of the advantages of this hybrids (fermenting at low temperature and produce more glycerol) can be correlated with the genome composition (Figure 2 and Table 1). Thus, it has been observed that most natural hybrids present as less, one *S. kudriavzevii* allele for the genes related with the cold stress, ergosterol and glycerol metabolism (Table 1). But also have been correlated some of these differences with gene expression (Combina et al., 2012; Gamero et al., 2015). The commercial hybrid W27 induced eight genes related to these functional groups compared with *S. cerevisiae* reference species. When expression of these W27 hybrid genes was compared with *S. kudriavzevii*, significant differences were observed in genes *PGII* and *TIP1* involved in glycerol biosynthesis;

in genes mainly belonged to the *PAU*, *DAN/TIR* families, linked to cold shock adaptation, while one gene (*ARE1*) is involved in sterol metabolic processes and in aroma synthesis (Figure 2). Besides, the quantitative contribution to the overall gene expression of the alleles coming from one parental strain or the other was clearly determined by the fermentation temperature for some genes as has been demonstrated for *ARO1* and *ATF2* genes where *S. kudriavzevii* allele was more expressed than that of *S. cerevisiae* particularly at 12°C (Gamero et al., 2015).

At present, the only contribution of *S. kudriavzevii* to winemaking is through its interspecific hybrids with other *Saccharomyces species*, most of them with *S. cerevisiae*, that are present, and very frequently predominant, in wine fermentations of European regions of Continental and Oceanic climates (Figure 2). Several wine hybrids were physiologically and genetically characterized (Figure 2), indicating that these hybrids take advantage of their chimerical genomes (Table 1), coming from both parental species, to adapt to changes in their environments (González et al., 2007; Gangl et al., 2009; Belloch et al., 2008; Arroyo-López et al., 2009), especially to grow at lower temperatures, a trait acquired from the *S. kudriavzevii* parent (González et al., 2006, González et al., 2007; Gangl et al., 2009; Salvadó et al., 2011; Tronchoni et al., 2012). The enological characterization of some of those *S. cerevisiae* x *S. kudriavzevii* hybrids have shown that *S. cerevisiae* provide ethanol tolerance and high fermentative capacity (Arroyo-López et al., 2010; Le Jeune et al., 2007; Bradbury et al., 2006) whereas *S. kudriavzevii* genome contributes to their adaptation to low temperatures (Belloch et al., 2008; Arroyo-López et al., 2009). In fact, glycerol production in the hybrids seems intermediate between both parentals in must fermentations, which suggest that the adaptation to low temperatures by increased glycerol production has been partially inherited from *S.*

kudriavzevii (González et al., 2007; Gangl et al., 2009). Regarding ethanol, hybrids usually produce similar or higher levels compared to *S. cerevisiae* (Gangl et al., 2009). On the other hand, both species showed a differentiated pattern of aroma compounds accumulation in winemaking which can generate a diverse aroma profile in *S. cerevisiae* x *S. kudriavzevii* hybrids. Aroma production by some hybrids (W27 and HA1841) showed that aroma production profile of these hybrids was similar to that of *S. kudriavzevii* at low fermentation temperature, whereas at moderate or high fermentation temperatures, they showed higher similarities with *S. cerevisiae* (González et al., 2007; Gangl et al., 2009). Other studies have shown differences between aroma profiles of hybrids and parental species except for higher alcohols production that were comparable to those of *S. cerevisiae* at 28°C and *S. kudriavzevii* at 12°C (Bellon et al., 2011; Cus and Jenko, 2013; Gamero et al., 2013).

GENERATION OF ARTIFICIAL HYBRIDS

Genomic blind approaches, as artificial hybridization, are the most adequate methodologies to generate of new industrial strains (Giudici et al., 2005). The most promising examples of wine yeast strains improvement involved generation of interspecies hybrids within the species of the *Saccharomyces* genus (Bellon et al., 2011) or further inbreed program of them (Bizaj et al., 2012). The final destination of the generated hybrid should be considered to select the hybridization method since the use of genetically modified organisms (GMOs) in food is limited by current legislations in different countries, as well as by public concern (Schilter and Constable, 2002; Pretorius and Hoj, 2005; Cebollero et al., 2007). According to Directive 2001/18/EC of the European Parliament and the Council of the European Union, hybrids

generated by mating of spores and rare-mating—based on the natural rare event of mating type switching in industrial yeasts—must not be considered as GMOs. Recent studies, have focused on optimize and develop programs of generation of artificial hybrids (Bellon et al., 2011, Pérez-Través et al., 2012; Pérez-Través et al., 2014; da Silva et al., 2015; Solieri et al., 2015; Lopandic et al., 2016). It has been described that hybrids obtained by rare-mating are easily obtained and contain a complete set of chromosomes of both parents (Pérez-Través et al., 2012; Pérez-Través et al., 2014). These artificial hybrids have a more complete subset of genetic material inherited from each parental strain just before they are generated. Consequently, they possess an extremely high genetic plasticity which could render a potentially better adaption to the environment. Due to the fact that a loss of genetic material occurs during hybrids generation and genetic stabilization, hybrids possessing a high amount of DNA became a better resource to obtain the best suitable hybrid strain for industrial purposes. However, to guarantee the genetic invariability of recently generated hybrids during future industrial utilization, the genetic stabilization is a last crucial aspect to be considered in every hybridization program (Pérez-Través et al., 2012; Pérez-Través et al., 2014). After generating the hybrid, 30-50 generations were enough to obtain genetically stable interspecific *S. cerevisiae*-*S. kudriavzevii* and *S. cerevisiae* intraspecific hybrids, respectively (Pérez-Través et al., 2012). Indeed, Pérez-Través et al. (2012) presented evidences of the existence of extensive genetic rearrangements among genetically similar genomes during hybrid genetic stabilization.

CONCLUSIONS

Yeasts starters are essential for the winemaking industry in order to obtain high quality standards. *S. cerevisiae* strains has been classically used but wineries require new strains with different physiological characteristics to face new consumer demands and other problems as the ones derived from climate change. Several studies have shown that wine *Saccharomyces* non-*cerevisiae* as *S. kudriavzevii* and *S. uvarum* are suitable for winemaking due to their cryophilic character, based on increased glycerol production, adapted membrane composition and enhanced translation efficiency, and the ability to produce valuable aromatic compounds. These species have great potential to solve some of these problems either as a part of natural or artificial *S. cerevisiae* x *S. kudriavzevii* hybrids, in co-cultures or directly as a new starter.

FUNDING

This work has supported by grants AGL2015-67504-C3-1-R from the Spanish Government and FEDER and Generalitat Valenciana PROMETEO (PROMETEOII/2014/042) to AQ.

REFERENCES

1. Aguilera, J., Randez-Gil, F., and Prieto, J. A. (2007). Cold response in *Saccharomyces cerevisiae*: new functions for old mechanisms. *FEMS Microbiol. Rev.* **31**:327-341.
2. Ale, C. E., Bru, E., Strasser de Saad, A. M., and Pasteris, S.E. (2014). Effect of physicochemical factors on glycerol production by simultaneous cultures of wine microorganisms using the response surface method. *J. Appl. Microbiol.* **117**:1336-47.
3. Ansell, R., Granath, K., Hohmann, S., Thevelein, J. M., and Adler, L. (1997). The two isoenzymes for yeast NAD⁺-dependent glycerol 3-phosphate dehydrogenase encoded by *GPD1* and *GPD2* have distinct roles in osmoadaptation and redox regulation. *EMBO J.* **16**:2179-2187.
4. Arroyo-López, F. N., Querol, A., and Barrio, E. (2009). Effects of temperature, pH and sugar concentration on the growth parameters of *Saccharomyces cerevisiae*, *S. kudriavzevii* and their interspecific hybrid. *Int. J. Food Microbiol.* **131**:120-7.
5. Arroyo-López, F. N., Pérez-Torrado, R., Querol, A., and Barrio, E. (2010). Modulation of the glycerol and ethanol syntheses in the yeast *Saccharomyces kudriavzevii* differs from that exhibited by *Saccharomyces cerevisiae* and their hybrid. *Food Microbiol.* **27**:628-37.
6. Arroyo-López, F.N., Pérez-Través, L., Querol, A., and Barrio, E. (2011). Exclusion of *Saccharomyces kudriavzevii* from a wine model system mediated by *Saccharomyces cerevisiae*. *Yeast.* **28**:423-35.

7. Barrio, E. González, S.S., Arias, A., Belloch, C., and Querol, A. (2006). *Molecular mechanisms involved in the adaptive evolution of industrial yeasts*. In: Querol, A., Fleet, G.H. eds, *Yeasts in Food and Beverages*, Ed 1st Vol 2. Springer-Verlag, 153-174.
8. Belda, I., Navascués, E., Marquina, D., Santos, A., Calderón F. (2016). Outlining the influence of non- conventional yeasts in wine ageing over lees. *Yeast* 33: 329–338
9. Belloch, C. Orlic, S., Barrio, E., and Querol, A. (2008). Fermentative stress adaptation of hybrids within the *Saccharomyces sensu stricto* complex. *Int. J. Food Microbiol.* **122**:188–195.
10. Bellon, J. R., Eglinton, J. M., Siebert, T., Pollnitz, A. P., Rose, L., De Barros Lopes, M., and Chambers, P. J. (2011). Newly generated interspecific wine yeast hybrids introduce flavour and aroma diversity to wines. *Appl. Microbiol. Biotechnol.* **91**:603–612.
11. Beltran, G., Novo, M., Leberre, V., Sokol, S., Labourdette, D., Guillamón, J. M., Mas, A., François, J., and Rozès, N. (2006). Integration of transcriptomic and metabolic analyses for understanding the global responses of low-temperature winemaking fermentations. *FEMS Yeast Res.* **6**:1167–118.
12. Bisson, L. F. (1999). Stuck and Sluggish Fermentations. *Am. J. Enol. Vitic.* **50**:107–119.
13. Bizaj, E. Cordente, A. G., Bellon, J. R., Raspor, P., Curtin, C. D., and Pretorius, I. S. (2012). A breeding strategy to harness flavor diversity of *Saccharomyces* interspecific hybrids and minimize hydrogen sulfide production. *FEMS Yeast Res.* **12**:456-65.
14. Borneman, A. R., Schmidt, S. A., and Pretorius, I. S. (2013), At the cutting-edge of grape and wine biotechnology. *Trends Genet.* **29**:263-71.

15. Boulton, R. B., Singleton, V. L., and Bisson, L. F. (1996). Principle and Practices of Winemaking. Chapman & Hall.
16. Bradbury, J.E., Richards, K. D., Niederer, H. A., Lee, S. A., Rod Dunbar, P., and Gardner, R. C. (2006). A homozygous diploid subset of commercial wine yeast strains *Antonie Van Leeuwenhoek*. **89**:27-37.
17. Burton, J., and Kerr, J. (2011) *The Cistercians in the Middle Ages*. The Boydell Press.
18. Canonico, L., Comitini, F., Oro, L., and Ciani, M. (2016). Sequential fermentation with selected immobilized non-*Saccharomyces* yeast for reduction of ethanol content in wine. *Front. Microbiol.* **7**:278.
19. Cebollero, E. D., González-Ramos, L., Tabera, R., and González (2007). González Transgenic wine yeast technology comes of age: is it time for transgenic wine? *Biotechnol. Lett.* **29**:191–200.
20. Ciani, M., Capece, A., Comitini, F., Canonico, L., Siesto, G., Romano, P. (2016). Yeast Interactions in Inoculated Wine Fermentation. *Front Microbiol.* **22**:7:555.
21. Ciani, M., Morales, P., Comitini, F., Tronchoni, J., Canonico, L., Curiel, JA., Oro, L., Rodrigues, AJ., Gonzalez, R. (2016). Non-conventional Yeast Species for Lowering Ethanol Content of Wines. *Front Microbiol.* **4**:7:642.
22. Combina, M., Pérez-Torrado, R., Tronchoni, J., Belloch, C., and Querol, A. (2012). Genome-wide gene expression of a natural hybrid between *Saccharomyces cerevisiae* and *S. kudriavzevii* under enological conditions. *Int. J. Food Microbiol.* **157**:340-5.
23. Comitini, F., Gobbi, M., Domizio, P., Romani, C., Lencioni, L., Mannazzu, I. and Ciani,

- M. (2011). Selected non-*Saccharomyces* wine yeasts in controlled multistarter fermentations with *Saccharomyces cerevisiae*. *Food Microbiol.* 2011 **28**(5):873-82.
24. Contreras, A. C., Hidalgo, P. A., Henschke, P. J., Chambers, C., Curtin, C., and Varela, C. (2015). Evaluation of non-*Saccharomyces* yeasts for the reduction of alcohol content in wine. *Appl. Environ. Microbiol.* **80**:1670–1678.
25. Cossins, A. R., and Prosser, C. L. (1978). Evolutionary adaptation of membranes to temperature. *Proc. Natl. Acad. Sci. U S A.* **75**:2040–2043.
26. Cus, F., and Jenko, M. (2013). The Influence of Yeast Strains on the Composition and Sensory Quality of Gewürztraminer Wine. *Food Technol. Biotechnol.* **51**:547–553.
27. da Silva, T., Albertin, W., Dillmann, C., Bely, M., la Guerche, S., Giraud, C., Huet, S., Sicard, D., Masneuf-Pomarede, I., de Vienne, D., Marullo, P. (2015). Hybridization within *Saccharomyces* Genus Results in Homoeostasis and Phenotypic Novelty in Winemaking Conditions PLoS One. 6;10(5): 1-24
28. de Barros Lopes, M., Bellon, J. R., Shirly, N. J., and Ganter, P. F. (2002). Evidence for multiple interspecific hybridization in *Saccharomyces sensu stricto* species. *FEMS Yeast Res.* **1**:323-331.
29. Dickinson, J. R., Lanterman, M. M., Danner, D. J., Pearson, B. M., Sanz, P., Harrison, S. J., and Hewlins, M. J. (1997). ¹³C nuclear magnetic resonance investigation of the metabolism of leucine to isoamyl alcohol in *Saccharomyces cerevisiae*. *J. Biol. Chem.* **272**:26871-8.

30. Dickinson, J. R., Eshantha, L., Salgado, J., and Hewlins, M. J. E. (2003). The catabolism of amino acids to long chain and complex alcohols in *Saccharomyces cerevisiae*. J. Biol. Chem. **278**:8028-34.
31. Eden, A., Van Nedervelde, L., Drukker, M., Benvenisty, N., and Debourg, A. (2001). Involvement of branched-chain amino acid aminotransferases in the production of fusel alcohols during fermentation in yeast. Appl. Microbiol. Biotechnol. **55**:296-300.
32. Erny, C., Raoult, P., Alais, A., Butterlin, G., Delobel, P., Matei-Radoi, F., Casaregola, S., and Legras, J. L. (2012). Ecological success of a group of *Saccharomyces cerevisiae*/*Saccharomyces kudriavzevii* hybrids in the Northern European wine making environment. Appl. Environ. Microbiol. **78**:3256-65.
33. Eustace, R., and Thornton R. J. (1987). Selective hybridization of wine yeasts for higher yields of glycerol. Can. J. Microbiol. **33**:112-117.
34. Fortner, D. M., Troy, R. G., and Brow, D. A. (1994). A stem/loop in U6 RNA defines a conformational switch required for pre-mRNA splicing. Genes Dev. **8**:221-233.
35. Gamero, A., Hernández-Orte, P., Querol, A., and Ferreira, V. (2011a). Effect of aromatic precursor addition to wine fermentations carried out with different *Saccharomyces species* and their hybrids. Int. J. Food Microbiol. **147**:33-44.
36. Gamero, A., Manzanares, P., Querol, A., and Belloch, C. (2011b). Monoterpene alcohols release and bioconversion by *Saccharomyces* species and hybrids. Int. J. Food Microbiol. **145**:92-7.
37. Gamero, A., Tronchoni, J., Querol, A., and Belloch, C. (2013a). Production of aroma compounds by cryotolerant *Saccharomyces* species and hybrids at low and moderate

- fermentation temperatures. J. Appl. Microbiol. **114**:1405-14.
38. Gamero, A., Wesselink, W., and de Jong, C. (2013b). Comparison of the sensitivity of different aroma extraction techniques in combination with gas chromatography-mass spectrometry to detect minor aroma compounds in wine. J. Chromatogr. A. **1272**:1–7.
 39. Gamero, A., Belloch C., Ibáñez C., and Querol A. (2014). Molecular analysis of the genes involved in aroma synthesis in the species *S. cerevisiae*, *S. kudriavzevii* and *S. bayanus* var. *uvarum* in winemaking conditions. PLoS One. **9**:e97626.
 40. Gamero, A., Belloch, C., and Querol, A. (2015). Genomic and transcriptomic analysis of aroma synthesis in two hybrids between *Saccharomyces cerevisiae* and *S. kudriavzevii* in winemaking conditions. Microb. Cell Fact. **14**:1-10.
 41. Gangl, H., Batusic, M., Tscheik, G., Tiefenbrunner, W., Hack, C., and Lopandic, K. (2009). Exceptional fermentation characteristics of natural hybrids from *Saccharomyces cerevisiae* and *S. kudriavzevii*. N. Biotechnol. **25**:244-51.
 42. García-Ríos, E., Querol A, Guillamón, JM. (2016). iTRAQ-based proteome profiling of *Saccharomyces cerevisiae* and cryotolerant species *Saccharomyces uvarum* and *Saccharomyces kudriavzevii* during low-temperature wine fermentation. J. Proteomics. **146**, 70–79
 43. Giudici, P., Solieri, A., Pulvirenti, M., and Cassanelli S. (2005). Strategies and perspectives for genetic improvement of wine yeasts. Appl. Microbiol. Biotechnol. **66**:622–628.
 44. González, S. S., Barrio, E., Gafner, J., and Querol, A. (2006). Natural hybrids from *Saccharomyces cerevisiae*, *Saccharomyces bayanus* and *Saccharomyces kudriavzevii* in

wine fermentations. FEMS Yeast Res. **6**:1221-34.

45. González, S. S., Barrio, E., and Querol A (2007). Enological characterization of natural hybrids from *Saccharomyces cerevisiae* and *S. kudriavzevii*. Int. J. Food Microbiol. **116**:11-8.
46. Greig, D., Louis, E. J., Borts, R. H., and Travisano, M. (2002). Hybrid speciation in experimental populations of yeast. Science. **298**:1773-1775.
47. Hayashi, M., and Maeda, T. (2006). Activation of the HOG pathway upon cold stress in *Saccharomyces cerevisiae*. J. Biochem. **139**:797-803.
48. Henschke, P.A., McWilliam, S., Fogarty, M., Francis, L., Kwiatkowsli, M., Høj, P., and Henschke, P. (2000). The effect of *Saccharomyces bayanus* var. *uvarum*-mediated fermentation on the chemical composition and aroma profile of Chardonnay wine. Aust. J. Grape Wine Res. **6**:190–196.
49. Henschke, P.A., and Rose A. H. (1991). *Plasma membrane* In: Rose, A.H. Harrison, J.S. (Eds.) *The yeasts, Yeast Organelles, IV*, Academic Press Limited, 297–345.
50. Hilliker, A. K., Mefford, M. A., and Staley, J. P. (2007). U2 toggles iteratively between the stem IIa and stem IIc conformations to promote pre-mRNA splicing. Genes Dev. **21**:821–834.
51. Hubmann, G., Guillouet, S., and Nevoigt, E. (2011). Gpd1 and Gpd2 fine-tuning for sustainable reduction of glycerol formation in *Saccharomyces cerevisiae*. Appl. Environ. Microbiol. **77**: 5857-67.

52. Ishikawa, T., and Noble, A. C. (1995). Temporal perception of astringency and sweetness in red wine. *Food Qual. Pref.* **6**:27–33.
53. Izawa, S., Sato, M., Yokoigawa, K., and Inoue, Y. (2004). Intracellular glycerol influences resistance to freeze stress in *Saccharomyces cerevisiae*: analysis of a quadruple mutant in glycerol dehydrogenase genes and glycerol-enriched cells. *Appl. Microbiol. Biotechnol.* **66**:108-114.
54. Jones, G. V., White, M. A., Cooper, O. R., and Storchmann, K. (2005). Climate change and global wine quality. *Climate Change.* **73**:319-343.
55. Kurata, S., Shen, B., Liu, J. O., Takeuchi, N., Kaji, A., and Kaji, H. (2013). Possible steps of complete disassembly of post-termination complex by yeast eEF3 deduced from inhibition by translocation inhibitors. *Nucleic Acids Res.* **41**:264–276.
56. Lencioni, L., Romani, C., Gobbi, M., Comitini, F., Ciani, M., Domizio, P. (2016). Controlled mixed fermentation at winery scale using *Zygorhizopus florentinus* and *Saccharomyces cerevisiae*. *Int J Food Microbiol.* 2016 **3;234**:36-44.
57. Le Jeune, C., Lollier, M., Demuyter, C., Erny, C., Legras, J. L., Aigle, M., and Masneuf-Pomarède, I. (2007). Characterization of natural hybrids of *Saccharomyces cerevisiae* and *Saccharomyces bayanus* var. *uvarum*. *FEMS Yeast Res.* **7**:540–549
58. Li, Z., and Brow, D.A. (1996). A spontaneous duplication in U6 spliceosomal RNA uncouples the early and late functions of the ACAGA element in vivo. *RNA.* **2**:879–894.
59. Lilly, M., Lambrechts, M. G., and Pretorius I. S. (2000). Effect of increased yeast alcohol acetyltransferase activity on flavor profiles of wine and distillates. *Appl. Environ. Microbiol.* **66**:744-53.

60. Llaudy M. del C., Canals, R., González-Manzano, S., Canals, J. M., Santos-Buelga, C., and Zamora, F. (2006). Influence of micro-oxygenation treatment before oak aging on phenolic compounds composition, astringency, and color of red wine. *Agric. Food Chem.* **54**:4246-52.
61. Lopandic, K., Gangl, H., Wallner, E., Tscheik, G., Leitner, G., Querol, A., Borth, N., Breitenbach, M., Prillinger, H., and Tiefenbrunner, W. (2007). Genetically different wine yeasts isolated from Austrian vine-growing regions influence wine aroma differently and contain putative hybrids between *Saccharomyces cerevisiae* and *Saccharomyces kudriavzevii*. *FEMS Yeast Res.* **7**:953-65.
62. Lopandic, K., Pfliegler, WP., Tiefenbrunner, W., Gangl, H., Sipiczki, M., Sterflinger, K. (2016). Genotypic and phenotypic evolution of yeast interspecies hybrids during high-sugar fermentation. *Appl Microbiol Biotechnol.* **100**(14):6331-43.
63. López-Malo, M., Querol, A., and Guillamon, J. M. (2013). Metabolomic comparison of *Saccharomyces cerevisiae* and the cryotolerant species *S. bayanus* var. *uvarum* and *S. kudriavzevii* during wine fermentation at low temperature. *PLoS One.* **8**:e60135.
64. Masneuf, I., Hansen, J., Groth, C., Piskur, J., and Dubourdieu, D. (1998). New hybrids between *Saccharomyces sensu stricto* yeast species found among wine and cider production strains. *Appl. Environ. Microbiol.* **64**:3887-3892.
65. Masneuf-Pomarede, I., Bely, M., Marullo, P., Lonvaud-Funel, A., and Dubourdieu, D. (2010). Reassessment of phenotypic traits for *Saccharomyces bayanus* var *uvarum* wine yeast strains. *Int. J. Food Microbiol.* **139**:79-86.

66. Milanovic, V., Maurizio, C., Oro, L., and Comitini F. (2012). *Starmerella bombicola* influences the metabolism of *Saccharomyces cerevisiae* at pyruvate decarboxylase and alcohol dehydrogenase level during mixed wine fermentation. *Microb. Cell Fact.* **11**:1-18
67. Morales, P., Rojas, V., Quirós, M., and Gonzalez, R. (2015). The impact of oxygen on the final alcohol content of wine fermented by a mixed starter culture. *Appl. Microbiol. Biotechnol.* **99**:3993-4003.
68. Naumov, G., James, S. A., Naumova, E. S., Louis, E. J., and Roberts, I. N. (2000). Association of *Saccharomyces bayanus* var. *uvarum* with some French wines: genetic analysis of yeast populations. *Res. Microbiol.* **151**:683–691.
69. Naumov, G.I., Naumova, E. S., Aigle, M., Masneuf, I., and Belarbi, A. (2001). Genetic reidentification of the pectinolytic yeast strain SCPP as *Saccharomyces bayanus* var. *uvarum*. *Appl. Microbiol. Biotechnol.* **55**:108-11.
70. Norbeck, J., and Blomberg, A. (1997). Metabolic and regulatory changes associated with growth of *Saccharomyces cerevisiae* in 1.4 M NaCl. Evidence for osmotic induction of glycerol dissimilation via the dihydroxyacetone pathway. *J. Biol. Chem.* **272**:5544-5554.
71. Oliveira, B. M., Barrio, E., Querol, A., and Pérez-Torrado, R. (2014). Enhanced enzymatic activity of glycerol-3-phosphate dehydrogenase from the cryophilic *Saccharomyces kudriavzevii*. *PLoS One.* **9**:e87290.
72. Paget, C. M., Schwartz, J. M., and Delneri, D. (2014). Environmental systems biology of cold-tolerant phenotype in *Saccharomyces* species adapted to grow at different temperatures. *Mol Ecol.* **23**:5241-57.

73. Pahlman, A. K. Granath, K., Ansell, R., Hohmann, S., and Adler, L. (2001). The yeast glycerol 3-phosphatases Gpp1p and Gpp2p are required for glycerol biosynthesis and differentially involved in the cellular responses to osmotic, anaerobic, and oxidative stress. *J. Biol. Chem.* **276**:3555-3563.
74. Panadero, J., Pallotti, C., Rodriguez-Vargas, S., Rande-Gil, F., and Prieto, J. A. (2006). A downshift in temperature activates the high osmolarity glycerol HOG pathway, which determines freeze tolerance in *Saccharomyces cerevisiae*. *J. Biol. Chem.* **281**:4638-4645.
75. Pérez-Través, L. Lopes, C. A., Barrio, E., and Querol, A. (2012). Evaluation of different genetic procedures for the generation of artificial hybrids in *Saccharomyces* genus for winemaking. *Int. J. Food Microbiol.* **156**:102-11.
76. Pérez-Través, L. Lopes, C. A., Barrio, E., and Querol A. (2014). Stabilization process in *Saccharomyces* intra and interspecific hybrids in fermentative conditions. *Int. Microbiol.* **17**:213-24.
77. Peris, D., Lopes, C. A., Belloch, C., Querol A., and Barrio E. (2012a). Comparative genomics among *Saccharomyces cerevisiae* x *Saccharomyces kudriavzevii* natural hybrid strains isolated from wine and beer reveals different origins. *BMC Genomics.* **13**:407.
78. Peris, D. Lopes, C. A., Arias, A., and Barrio, E. (2012b). Reconstruction of the evolutionary history of *Saccharomyces cerevisiae* x *S. kudriavzevii* hybrids based on multilocus sequence analysis. *PLoS One.* **7**, e45527.
79. Perriman, R. J., and Ares, M. (2007). Rearrangement of competing U2 RNA helices within the spliceosome promotes multiple steps in splicing. *Genes Dev.* **21**:811–820.

80. Pretorius, I. S., and Hoj, P. B. (2005). Grape and wine biotechnology: challenges, opportunities and potential benefits. *Aust. J. Grape Wine Res. Res.* **11**:83–108.
81. Pretorius, I. S. (2000). Tailoring wine yeast for the new millennium: novel approaches to the ancient art of winemaking. *Yeast* **16**:675-729.
82. Querol, A., Fernández-Espinar, M. T., del Olmo, M., and Barrio, E. (2003). Adaptive evolution of wine yeast. *Int. J. Food Microbiol.* **86**:3-10.
83. Redón, M., Guillamón, J. M., Mas, A., and Rozès, N. (2011). Effect of growth temperature on yeast lipid composition and alcoholic fermentation at low temperature. *Eur. Food Res. Technol.* **232**:517-527.
84. Remize, F., Barnavon, L., and Dequin, S. (2001). Glycerol export and glycerol-3-phosphate dehydrogenase, but not glycerol phosphatase, are rate limiting for glycerol production in *Saccharomyces cerevisiae*. *Metab. Eng.* **3**:301-312.
85. Ribereau-Gayon, J., Peynaud, E., Ribereau-Gayon, P., and Sudraud P. (1977). In. *Traité D'oenologie. Sciences et Techniques Du Vin*, pp. 353-361. Paris: Dunod.
86. Saerens, S. M. G., Delvaux, F., Verstrepen, K. J., Van Dijck, P., Thevelein, J. M., and Delvaux, F. R. (2008). Parameters affecting ethyl ester production by *Saccharomyces cerevisiae* during fermentation. *Appl. Environ. Microbiol.* **74**:454-61.
87. Sakamoto, T., and Murata, N. (2002). Regulation of the desaturation of fatty acids and its role in tolerance to cold and salt stress. *Curr. Opin. Microbiol.* **5**:208-10.
88. Salvadó, Z., Arroyo-López, F. N., Guillamón, J. M., Salazar, G., Querol, A., and Barrio, E. (2011). Temperature adaptation markedly determines evolution within the genus *Saccharomyces*. *Appl. Environ Microbiol.* **77**:2292-302.

89. Schilter, B., and Constable A. (2002). Regulatory control of genetically modified (GM) foods: likely developments Toxicol. Lett. **127**:341–349.
90. Schuller, D., and Casal, M. (2005). The use of genetically modified *Saccharomyces cerevisiae* strains in the wine industry. Appl. Microbiol. Biotechnol. **68**:292-304.
91. Solieri, L., Verspohl, A., Bonciani, T., Caggia, C., and Giudici, P. (2015). Fast method for identifying inter- and intra-species *Saccharomyces* hybrids in extensive genetic improvement programs based on yeast breeding. J. Appl. Microbiol. **119**:149-61.
92. Staley, J. P., and Guthrie, C. (1999). An RNA switch at the 5' splice site requires ATP and the DEAD box protein Prp28p. Mol. Cell. **3**:55–64.
93. Stribny, J., Gamero, A., Pérez-Torrado, R., and Querol, A. (2015). *Saccharomyces kudriavzevii* and *Saccharomyces uvarum* differ from *Saccharomyces cerevisiae* during the production of aroma-active higher alcohols and acetate esters using their amino acidic precursors. Int. J. Food Microbiol. **205**:41-6.
94. Stribny, J., Querol, A., and Pérez-Torrado, R. 2016a. Differences in Enzymatic Properties of the *Saccharomyces kudriavzevii* and *Saccharomyces uvarum* Alcohol Acetyltransferases and Their Impact on Aroma-Active Compounds Production. Front. Microbiol. **7**:897.
95. Stribny, J., Romagnoli, G., Pérez-Torrado, R., Daran, J. M., and Querol, A. 2016b. Characterisation of the broad substrate specificity 2-keto acid decarboxylase Aro10p of *Saccharomyces kudriavzevii* and its implication in aroma development. Microb. Cell Fact. **15**:51.

96. Swiegers, J. H., and Pretorius, I. S. (2005). Yeast modulation of wine flavor. *Adv. Appl. Microbiol.* **57**:131-75.
97. This, P., Lacombe, T., and Thomas, M. R. (2006). Historical origins and genetic diversity of wine grapes. *Trends Genet.* **22**:511–519.
98. Torija, M.J., Beltran, G., Novo, M., Poblet, M., Guillamón, J. M., Mas, A., and Rozès, N. (2003). Effects of fermentation temperature and *Saccharomyces* species on the cell fatty acid composition and presence of volatile compounds in wine. *Int. J. Food Microbiol.* **85**:127-36.
99. Tosi, E., Azzolini, M., Guzzo, F., and Zapparoli, G. (2009). Evidence of different fermentation behaviours of two indigenous strains of *Saccharomyces cerevisiae* and *Saccharomyces uvarum* isolated from Amarone wine. *J. Appl. Microbiol.* **107**:210-8.
100. Tronchoni, J., Arroyo-Lopez, F. N., Gamero, A., Querol, A., and Barrio, E. (2009). Differences in the glucose and fructose consumption profiles in diverse *Saccharomyces* wine species and their hybrids during grape juice fermentation. *Int. J. Food Microbiol.* **134**:237-243.
101. Tronchoni, J., Rozès, N., Querol, A., and Guillamón, J. M. (2012). Lipid composition of wine strains of *Saccharomyces kudriavzevii* and *Saccharomyces cerevisiae* grown at low temperature. *Int. J. Food Microbiol.* **155**:191–198.
102. Tronchoni, J., Medina, V., Guillamón, J. M., Querol, A., and Pérez-Torrado, R. (2014). Transcriptomics of cryophilic *Saccharomyces kudriavzevii* reveals the key role of gene translation efficiency in cold stress adaptations. *BMC Genomics.* **15**:432.

103. Tulha, J., Lima, A., Lucas, C., and Ferreira, C. (2010). *Saccharomyces cerevisiae* glycerol/H⁺ symporter Stl1p is essential for cold/near-freeze and freeze stress adaptation. A simple recipe with high biotechnological potential is given. *Microb. Cell Fact.* **9**:82.
104. Varela, C. (2016). The impact of non-*Saccharomyces* yeasts in the production of alcoholic beverages. *Appl Microbiol Biotechnol.* 2016 **100(23)**:9861-9874.
105. Wang, C., Mas, A., Esteve-Zarzoso, B. (2016). The Interaction between *Saccharomyces cerevisiae* and Non-*Saccharomyces* Yeast during Alcoholic Fermentation Is Species and Strain Specific. *Front Microbiol.*
106. White, M. A., Diffenbaugh, N. S., Jones, G. V., Pal, J. S., and Giorgi F. (2006). Extreme heat reduces and shifts United States premium wine production in the 21st century. *Proc. Natl. Acad. Sci. U S A.* **103**:11217-22.
107. Zavanelli, M., Britton, J., Igel, A., and Ares, M. Jr. (1994). Mutations in an essential U2 small nuclear RNA structure cause cold-sensitive U2 small nuclear ribonucleoprotein function by favoring competing alternative U2 RNA structures. *Mol. Cell Biol.* **14**:1689.

Table 1. Genome composition for the natural hybrids between *S. cerevisiae* and *S. kudriavzevii* for genes involved in their physiological differences (Peris et al., 2012a).

ORF	Gene	AMH	PB7	Vin7	HA [*]	HA 1841	SOY3	W27	W46	SPG 16-91	441
Cold stress											
YNL112W	<i>DBP2</i>	CC	CK	CK	CK	CK	CK	CCK	CCK	CCK	CCK
YJL153C	<i>INO1</i>	CC	CK	CK	CK	CK	CK	CK	CK	CK	CK
YJL223C	<i>PAU1</i>	CC	CK	CK	CK	CK	CK	CK	CK	CK	CK
YEL049W	<i>PAU2</i>	CC	CK	CK	CK	CC	CK	CCK	CCK	CCK	CCK
YCR104W	<i>PAU3</i>	CC	CK	CC	CK	CK	CK	CK	CK	CK	CK
YLR461W	<i>PAU4</i>	CC	CK	CK	CK	CK	CK	CK	CK	CC	CK
YNR076W	<i>PAU6</i>	CC	CK	CK	CK	CK	CK	CC	CCC	CCC	CCC

YAR020C	<i>PAU7</i>	C	CK	CK	CK	CK	CK	CK	CK	CK	CC
YBR067C	<i>TIP1</i>	CC	CK	CK	CK	CK	CK	CK	CK	CK	CK
YER011W	<i>TIR1</i>	CC	CK	CK	CK	CC	CK	CKK	CKK	CKK	CKK
Ergosterol metabolism											
YGR175C	<i>ERG1</i>	CC	CK	CC	CCK	CK	CK	CK	CK	CK	CK
YHR007C	<i>ERG11</i>	C	CK	CK	CK	CK	CK	CK	CK	CK	CCK
YGR060W	<i>ERG25</i>	CK	CK	CK	CCK	CK	CK	CK	CK	CK	CK
YGL001C	<i>ERG26</i>	CK	CK	CK	CCK	CK	CK	CK	CK	CK	CK
YLR100W	<i>ERG27</i>	CC	CK	CK	CK	CK	CK	CK	CK	CK	CK
YLR056W	<i>ERG3</i>	CC	CK	CK	CK	CK	CK	CK	CK	CK	CK
YGL012W	<i>ERG4</i>	CK	CK	CK	CCK	CK	CK	CK	CK	CK	CK

YHR072W	<i>ERG7</i>	C	CK	CK	CK	CK	CK	CK	CK	CK	CKK
YNR043W	<i>MVD1</i>	CC	CK	CK	CK	CK	CK	CKK	CKK	CKK	CKK
YHR042W	<i>NCP1</i>	C	CK	CK	CK	CK	CK	CK	CK	CK	CKK
Glycerol metabolism											
YBR196C	<i>PGI1</i>	CC	CK	CK	CK	CK	CK	CK	CK	CK	CK
YDR050C	<i>TPH1</i>	CC	CK	CK	CK	CK	CC	CC	CC	CC	CC
Osmotic stress											
YGR175C	<i>ERG1</i>	CC	CK	CC	CCK	CK	CK	CK	CK	CK	CK
YLR056W	<i>ERG3</i>	CC	CK	CK	CK	CK	CK	CK	CK	CK	CK
YDL022W	<i>GPD1</i>	CC	CK	CK	CK	CK	CK	CC	CC	CC	CC
YLR362W	<i>STE11</i>	CC	CK	CK	CK	CK	CK	CK	CK	CK	CK

Oxidative stress											
YHR183W	<i>GND1</i>	C	CK	CK	CK	CK	CK	CK	CK	CK	CCK
YOL151W	<i>GRE2</i>	CK	CK	CK	CK	CK	CK	CC	CC	CC	CC
YJL101C	<i>GSH1</i>	CC	CK	CK	CK	CK	CK	CK	CK	CK	CK
YGR209C	<i>TRX2</i>	CC	CK	CC	CCK	CK	CK	CK	CK	CK	CK
YGR019W	<i>UGA1</i>	CK	CK	CK	CCK	CK	CK	CK	CK	CK	CK
Membrane fluidity											
YLR153C	<i>ACS2</i>	CC	CK	CK	CK	CK	CK	CK	CK	CK	CK
YER026C	<i>CHO1</i>	CC	CK	CK	CK	CC	CK	CKK	CKK	CKK	CKK
YGL055W	<i>OLE1</i>	CK	CK	CK	CCK	CK	CK	CK	CK	CK	CK
YNL169C	<i>PSD1</i>	CC	CK	CK	CK	CK	CK	CCK	CCK	CCK	CCK

*Include strains HA1835, HA1837, HA1842. Colors indicate absence of *S. kudriavzevii* (K) alleles with one (grey), two

(magenta) or three (orange) *S. cerevisiae* (C) gene copies; or two *S. kudriavzevii* alleles with one copy of *S. cerevisiae* (green); or one *S. kudriavzevii* allele with two copy of *S. cerevisiae* (blue).




Figure1. Summary of new challenges for the winemaking industry and the potential benefits of using *S. kudriavzevii* and *S. uvarum* as a starter. The experience acquired from the *S. kudriavzevii* and *S. uvarum* wine performances suggest relevant benefits that can potentially solve most of these new problems. such as glycerol accumulation, changes in membrane composition (higher levels of medium chain fatty acids (MCFA) and squalene (SQ), whereas phosphatidic acid (PA) content and the phosphatidylcholine/phosphatidylethanolamine ratio (PC/PE) lowered) or enhanced translation efficiency (ribosome recycling). They may

considerably contribute to the aroma profile for *S. cerevisiae* by increasing the levels of higher alcohols and esters in the final wine. Genes involved overexpressed in these species are depicted.




Figure 2. Distribution of the *S. cerevisiae* x *S. kudriavzevii* hybrids. The geographic origin and sources from which the different hybrids were isolated are indicated on a climatic map of Europe. These hybrids are linked according to their group assignments, based on previous phylogenetic analyses (Erny et al., 2012; Peris et al., 2012a; Peris et al., 2012b). Within squares numbers indicate the number of strains described until now (up) and the range of *S. kudriavzevii* genome present in % (down).