



Industrial potential of carotenoid pigments from microalgae: Current trends and future prospects

Ranga Rao Ambati, Deepika Gogisetty, Ravishankar Gokare
Aswathanarayana, Sarada Ravi, Panduranga Narasimharao Bikkina, Lei Bo &
Su Yuepeng

To cite this article: Ranga Rao Ambati, Deepika Gogisetty, Ravishankar Gokare
Aswathanarayana, Sarada Ravi, Panduranga Narasimharao Bikkina, Lei Bo & Su Yuepeng (2018):
Industrial potential of carotenoid pigments from microalgae: Current trends and future prospects,
Critical Reviews in Food Science and Nutrition, DOI: [10.1080/10408398.2018.1432561](https://doi.org/10.1080/10408398.2018.1432561)

To link to this article: <https://doi.org/10.1080/10408398.2018.1432561>



Accepted author version posted online: 25
Jan 2018.
Published online: 16 Feb 2018.



Submit your article to this journal [↗](#)



Article views: 375



View Crossmark data [↗](#)



Industrial potential of carotenoid pigments from microalgae: Current trends and future prospects

Ranga Rao Ambati^{a,b,c}, Deepika Gogisetty^d, Ravishankar Gokare Aswathanarayana^e, Sarada Ravi^f, Panduranga Narasimharao Bikkina^g, Lei Bo^a, and Su Yuepeng^b

^aFood Science and Technology Programme, Beijing Normal University-Hong Kong Baptist University United International College, Tangjiawan, Zhuhai, Guangdong, China; ^bEstuarine Fisheries Research Institute, Doumen, Zhuhai, Guangdong, China; ^cDepartment of Biotechnology, Vignan's Foundation for Science, Technology and Research (Deemed to be University), Vadlamudi, Guntur, Andhra Pradesh, India; ^dDepartment of Chemistry, Sri Chaitanya Junior College, Tenali, Guntur, Andhra Pradesh, India; ^eDr. C. D. Sagar Center for Life Sciences, Dayananda Sagar Institutions, Bangalore, Karnataka, India; ^fPlant Cell Biotechnology Department, Central Food Technological Research Institute, (Constituent Laboratory of Council of Scientific & Industrial Research), Mysore, Karnataka, India; ^gRegional Centre, Indira Gandhi National Open University, Mumbai, Maharashtra, India

ABSTRACT

Microalgae are rich source of various bioactive molecules such as carotenoids, lipids, fatty acids, hydrocarbons, proteins, carbohydrates, amino acids, etc. and in recent Years carotenoids from algae gained commercial recognition in the global market for food and cosmeceutical applications. However, the production of carotenoids from algae is not yet fully cost effective to compete with synthetic ones. In this context the present review examines the technologies/methods in relation to mass production of algae, cell harvesting for extraction of carotenoids, optimizing extraction methods etc. Research studies from different microalgal species such as *Spirulina platensis*, *Haematococcus pluvialis*, *Dunaliella salina*, *Chlorella* sps., *Nannochloropsis* sps., *Scenedesmus* sps., *Chlorococcum* sps., *Botryococcus braunii* and *Diatoms* in relation to carotenoid content, chemical structure, extraction and processing of carotenoids are discussed. Further these carotenoid pigments, are useful in various health applications and their use in food, feed, nutraceutical, pharmaceutical and cosmeceutical industries was briefly touched upon. The commercial value of algal carotenoids has also been discussed in this review. Possible recommendations for future research studies are proposed.

KEYWORDS

microalgae; carotenoids; processing; biological activities; applications

CHEMICAL COMPOUNDS STUDIED IN THIS ARTICLE

Neoxanthin (PubChem CID: 5281247); Violaxanthin (PubChem CID: 448438); Astaxanthin (PubChem CID: 5281224); Lutein (PubChem CID: 5281243); Zeaxanthin (PubChem CID: 5280899); β -cryptoxanthin (PubChem CID: 5281235); α -carotene (PubChem CID: 6419725); β -carotene (PubChem CID: 5280489); Canthaxanthin (PubChem CID: 5281227); Fucoxanthin (PubChem CID: 5281239)

1. Introduction

Carotenoids are known for their industrial applications (Dufosse et al., 2005; Rodriguez-Amaya, 2016). While natural carotenoids are produced by plants (Giuliano et al., 2008; Ramel et al., 2012; Carbonell-Capella et al., 2014), synthetic ones are mainly by products from coal distillation. However, there is growing demand to phase out the use of synthetic carotenoids in many countries due to health considerations. Use of natural bioactive compounds in functional foods is beneficial as they will be able to protect cells from oxidative damage (Ranga Rao et al., 2013a, 2014a; Carbonell-Capella et al., 2014). They are also valuable in protecting deterioration of food products during the storage and processing. It is also a known fact that carotenoid pigments can influence signaling and regulation of many biological pathways (Stahl et al., 2002). More than 600 naturally occurring carotenoid pigments are identified and characterized and among them astaxanthin, β -carotene and lutein are considered to be important carotenoids for their potential applications in foods (Zhang et al., 2014). Microalgal species are also known to produce various bioactive compounds

and fine chemicals (Pignolet et al., 2013; Phang et al., 2015; Ranga Rao et al., 2016). Certain algal species are exploited commercially for proteins, vitamins, pigments, fatty acids, lipids and phenolic compounds (Spolaore et al., 2006; Larkum et al., 2012). Table 1 shows recent review of literature on microalgal species for various uses. Recently, carotenoid pigments from algae have received more attention in health food applications (Guedes et al., 2011a; 2010b; Buono et al., 2014; Zhang et al., 2014). Among the algal species, *Haematococcus pluvialis*, *Dunaliella salina*, *Chlorella* sps., *Scenedesmus* sps., *Spirulina platensis*, *Botryococcus braunii*, and *Diatoms* are well known for the production of β -carotene, lutein, canthaxanthin, astaxanthin, and fucoxanthin (Lamers et al., 2008; Ranga Rao et al., 2010a; Zhang et al., 2014). However, the pigment composition of algae vary from species to species and also according to environmental conditions. It is observed that most of the algal species accumulate more amounts of pigments under stress conditions like nutrient deficiency (Ranga Rao et al., 2010b; Sarada et al., 2012). The present review provides information on the

Table 1. Recent literature on microalgae and their use in various applications.

Micro algal species	Focus on	References
<i>S. platensis</i> , <i>H. pluvialis</i> , <i>D. salina</i> , <i>Chlorella</i> sps, <i>Nannochloropsis</i> sps, <i>Scenedesmus</i> sps, <i>Chlorococcum</i> sps, <i>B. braunii</i> , <i>Diatoms</i>	Carotenoid pigments, chemical structure, extraction and processing, commercial applications, carotenoid producing companies, global market, current trends and future prospects	Current article
Not specified algal species	Carotenoids, biosynthesis pathway, genetic engineering, transcription factors, metabolic engineering, nutritional fatty acids, lipids and carotenoids.	Coesel et al., 2008; Vigani et al., 2015; Odjadjare et al., 2015; Varela et al., 2015; Wang et al., 2015; Henriquez et al., 2016; Gong and Bassi, 2016; D Alessandro and Antoniosi Filho, 2016; Hamed, 2016; Minhas et al., 2016; Schweiggert and Carle, 2016; Bajhaiya et al., 2017; Ranga Rao et al., 2014a; Shah et al., 2016; Kishimoto et al., 2016; Bilbao et al., 2016; Panis and Carreon 2016; Zhang et al., 2014
<i>H. pluvialis</i>	Astaxanthin production, astaxanthin sources, stability, nutritional properties, commercial applications.	
<i>H. pluvialis</i> , <i>D. salina</i> , <i>Chlorella</i> sps, <i>Scenedesmus</i> sps	Carotenoid health benefits.	
<i>S. platensis</i> , <i>H. pluvialis</i> , <i>Chlorella</i> sps, <i>Scenedesmus</i> sps, <i>Isochrysis</i> sps; <i>Nannochloropsis</i> sps, <i>Porphyridium</i> sps, <i>Tetraselmis</i> sps, <i>S. platensis</i>	Bioactive molecules for livestock feed, poultry, aquaculture, and bioremediation	Yaakob et al., 2014a; Yaakob et al., 2014b; Roy and Pal, 2015
	Nutritional, Medical applications	Eriksen, 2008; Kuddus et al., 2013; Hoseini et al., 2013
<i>S. platensis</i> , <i>H. pluvialis</i> , <i>D. salina</i> , <i>Chlorella</i> sps, <i>Scenedesmus</i> sps, <i>Isochrysis</i> sps	Functional properties of carotenoids; Functional ingredients; Novel food products	Gouveia et al., 2008; Plaza et al., 2009; Christaki et al., 2012; Vilchez et al., 2011; Buono et al., 2014; Guedes et al., 2011a; 2011b; Del Campo et al., 2007
<i>H. pluvialis</i> , <i>D. salina</i> , <i>Chlorella</i> sps, <i>Scenedesmus</i> sps, <i>Muriellopsis</i> sps, <i>Chlorella</i> sps, <i>Scenedesmus</i> sps, <i>Chlorococcum</i> sps, <i>Muriellopsis</i> sps	Lutein production	Fernandez et al., 2010.
<i>S. platensis</i> , <i>H. pluvialis</i> , <i>D. salina</i> , <i>Chlorella</i> sps, <i>Nannochloropsis</i> sps, <i>Isochrysis</i> sps	Commercial applications	Spolaore et al., 2006.
<i>S. platensis</i> , <i>H. pluvialis</i> , <i>D. salina</i> , <i>Diatoms</i>	Micro-organisms for pigments Pigments, fucoxanthin, biosynthesis pathway, biological properties, applications,	Dufosse et al., 2005. Lee et al., 2006; Kroger and Poulsen, 2008; Bozarth et al., 2009; Prestegard et al., 2009; Xia et al., 2013; Kuczyńska et al., 2015; Fu et al., 2015; Lelyana, 2016; Guo et al., 2016; Medlin, 2016.

carotenoid pigments including pigment production, chemical structure, extraction process, and commercial applications from select algal species from Cyanophyceae, Chlorophyceae, Rhodophyceae and diatoms. The select microalgal species include *S. platensis*, *H. pluvialis*, *D. salina*, *Chlorella* sps, *Nannochloropsis* sps, *Scenedesmus* sps, *Chlorococcum* sps, *Botryococcus braunii*, and *Diatoms*. Further, their market values and the future prospects of carotenoids are discussed.

2. Structure of carotenoids

Carotenoids are classified into two types based on their chemical structure. Oxygenated carotenoids are referred as xanthophyll's while other carotenoids are hydrocarbon carotenoids or simply referred as carotenoids. Carotenoids are fat soluble pigments and are tetraterpenoids (C₄₀). C₄₀ carbon atoms are considered as the backbone of the carotenoid molecule. The content and composition of carotenoid pigments produced by microalgae species vary and are influenced by the culture conditions (Sarada et al., 2012; Ranga Rao et al., 2014a; Wells et al., 2016). Figure 1 shows common chemical structure of carotenoids in microalgal species.

3. Carotenoid pigments from microalgae

Microalgae are photosynthetic micro-organisms of around 30,000 species. Algal groups are classified as Cyanophyta(e.g. blue-green algae), Phaeophyta(e.g. brown algae), Rhodophyta

(e.g. red algae) and Chlorophyta(e.g. green algae) based on their morphological features. However, based on the carotenoids, ten major classes of algae are identified (Table 2)

Though synthetic pigments are banned in many countries due to their adverse effects on human health, still they are used in various applications in many other countries (Mohammed and Mohd, 2011). The world demand of carotenoids is increasing due to consumer preference of natural carotenoids to synthetic ones. Based on the class to which the alga belongs and based on the species either carotenoids, or chlorophylls or phycobiliproteins may be dominant pigment in a particular alga. All these algae based pigments (Table 1) are used in various applications (Dufosse et al., 2005; Spolaore et al., 2006; Gantar and Svircev, 2008; Ceron et al., 2008; Ranga Rao et al., 2009; Ranga Rao et al., 2010b; Guedes et al., 2011a; 2011b; Pignolet et al., 2013; Yaakob et al., 2014a; Roy and Pal, 2015). Different stress conditions such as temperature, salinity, and irradiance may induce carotenoid production in algal species (Ranga Rao et al., 2010b; Sarada et al., 2012; Fu et al., 2014; Minhas et al., 2016). Accumulation of carotenoids in algae under nutrient deficiency culture conditions was reported by various authors (Orosa et al., 2001; Ranga Rao et al., 2010b; Sarada et al., 2012; Fu et al., 2013; Tran et al., 2014). Carotenoids from various microalgal species are discussed in the following section. Table 3, 4 and 5 summarizes culture conditions for carotenoid production, major carotenoid pigments, and their content for the selected microalgal species.




Figure 1. Common chemical structures of carotenoid pigments in microalgae. (a). Neoxanthin (3S,5R,6R,3'S,5'R,6'S)-6,7-didehydro-5',6'-epoxy-5,6,5',6'-tetrahydro- β,β -carotene-3,5,3'-triol), (b). Violaxanthin (3S,5R,6R,3'S,5'R,6'S)-5,6, 5',6'-diepoxy-5,6,5', 6'-tetrahydro- β,β -carotene-3,3'-diol), (c). Astaxanthin ((3S,3'S)-3,3'-dihydroxy- β,β -carotene-4,4'-dione), (d). Lutein (3R,3'R,6'R)- β,β -carotene-3,3'-diol), (e). Zeaxanthin ((3R,3'R)- β,β -carotene-3,3'-diol), (f). β -cryptoxanthin ((3R)- β,β -caroten-3-ol), (g). α -carotene ((6'R)- β,β -carotene), (h). β -carotene (β,β -carotene), (i). Canthaxanthin (β,β -carotene-4,4'-dione), and (j). Fucoxanthin ((3S,5R,6S,3'S,5'R,6'R)-5,6-epoxy-3,3',5'-trihydroxy-6',7'-didehydro-5,6,7,8,5',6'-hexahydro- β,β -caroten-8-one 3'-acetate).

4. Select microalgal species for carotenoid pigments

4.1. Cyanobacteria (Cyanophyceae forms)

4.1.1. *Spirulina platensis*

S. platensis is studied extensively as it contains high amount of protein ranging from 55 to 65% (Coca et al., 2014). *Spirulina* biomass is used commercially since it contains higher amounts of proteins, fatty acids including essential fatty acids and pigments (Hoseini et al., 2013). Carotenoid content of this alga was reported to vary from 0.1 to 0.4 mg/g (Ranga Rao et al., 2010a; Kumar et al., 2013). Carotenoids such as β -carotene, echinenone, β -cryptoxanthin, β -carotene-5,6-epoxide, 3-hydroxyechinenone, lutein, zeaxanthin, diatoxanthin, canthaxanthin, myxoxanthophyll, and oscillaxanthin were isolated and identified from *Spirulina* sps (Hanaa et al., 2003; Ranga Rao et al., 2010a; 2013b). Major carotenoids identified in *S. platensis* are β -carotene, zeaxanthin and β -cryptoxanthin (Ranga Rao et al., 2010a). In addition to carotenoids, *S. platensis* contains phycobiliprotein, the content of which varies based on the culture conditions (Simeunovic et al., 2012; Simeunovic et al.,

2013). Upon consumption, these carotenoids help in controlling photo-oxidative damage of cells, immune enhancement, scavenging free radicals, hormone regulation, and enhancement of cell growth, cell maturation and multiplication (Erikssen, 2008).

4.2. Chlorophyceae forms

4.2.1. *Haematococcus pluvialis*

H. pluvialis, green alga is well-known to produce astaxanthin and astaxanthin esters (Ranga Rao et al., 2013a). It is grown in autotrophic and heterotrophic culture conditions (Kang et al., 2005; Sarada et al., 2012). The carotenogenesis stage in this alga involves formation of thick walled aplanospores and accumulation of astaxanthin to a level of 2–4% of the dry weight (Ranga Rao et al., 2010a, 2013a, 2014a). Previously, we reported carotenoids in *H. pluvialis* as violaxanthin, astaxanthin, astaxanthin esters, lutein, zeaxanthin, α -carotene, and β -carotene (Ranga Rao et al., 2009, 2013a). Astaxanthin is permitted as color additive in fish feed by United States of Food and Drug

Table 2. Classification of algae and their carotenoid pigments.

Algae name	Scientific name	Carotenoid pigments
Blue-green algae	Cyanophyceae/Myxophyceae	Chlorophyll-a, β -carotene, c-phycoerythrin, c-phycoyanin,
Green algae	Chlorophyceae	Chlorophyll-a, chlorophyll-b, α -carotene, β -carotene, γ -carotene, lutein, siphonoxanthin and siphonein
Yellow-green algae	Xanthophyceae	Chlorophyll-a, β -carotene, diatoxanthin, diadinoxanthin, heteroxanthin
Golden-brown algae	Chrysophyceae	Chlorophyll-a, chlorophyll-c1, chlorophyll-c2 and fucoxanthin
Diatoms	Bacillariophyceae	Chlorophyll-a, chlorophyll-c, β -carotene, fucoxanthin, diatoxanthin and diadinoxanthin,
Cryptomonads	Cryptophyceae	Chlorophyll-a, chlorophyll-c, α -carotene, diatoxanthin, phycoerythrin and phycocyanin
Dinoflagellates	Dinophyceae	Chlorophyll-a, chlorophyll-c2, β -carotene, peridinin and neoperidinin
Euglenoids	Euglenineae Euglenophyceae	Chlorophyll-a, chlorophyll-b, α -carotene, β -carotene, astaxanthin, antheraxanthin, diadinoxanthin and neoxanthin
Brown algae	Phaeophyceae	Chlorophyll-a, chlorophyll-c1, chlorophyll-c2, and β -carotene
Red algae	Rhodophyceae	Chlorophyll-a, r-phycoyanin, allo-phycoyanin, c-phycoerythrin, α -carotene, β -carotene

Administration (USFDA) (Pashkow et al., 2008). European countries have also approved astaxanthin as ingredient in human dietary supplement (Roche, 1987).

4.2.2. *Dunaliella salina*

D. salina is a motile green microalga which habitats lakes, oceans and brackish water bodies. It is a unicellular algal genus belonging to the family *Dunaliellaceae* (Feng et al., 2014; Lao et al., 2014). It is rich in carotenoids content that varies based on the culture conditions (Murthy et al., 2005; Tan and Suter, 2011; Fu et al., 2013; 2014; Tran et al., 2014). *Dunaliella* contains high amount of carotenoids (Fu et al., 2013). Thus, *Dunaliella* pro-vitamin A carotenes exhibit bright red color. Recently, carotenoids from *D. salina* are approved by USFDA as food color, and is recognized as safe natural colour (Dufosse et al., 2005; Yang et al., 2013). The provitamin-A carotenoids of *Dunaliella* are used as food additive (Hosseini and Shariati, 2009).

4.2.3. *Chlorella* sps

Chlorella is a green microalga producing good quantities of proteins, vitamins, minerals, carotenoids, and polysaccharides.

(Sharma et al., 2012; Liu and Hu, 2013). *Chlorella* sps are also reported to produce xanthophylls such as violaxanthin, lutein and zeaxanthin, with lutein as the major carotenoid (Del Campo et al., 2004; Gouveia et al., 2006; Cha et al., 2008). *Chlorella vulgaris* and *Chlorella protothecoides* are mass cultivated both heterotrophically and autotrophically for biomass and carotenoid production (Shi et al., 2002; Wei et al. 2008; Seyfabadi et al., 2010; Campenni et al., 2013). In heterotrophic culture conditions, algal cells utilize organic carbon source available as energy whereas under autotrophic conditions cells collect the energy through photosynthesis (Larsdotter, 2006). *C. vulgaris* cells double in eight hours at a temperature of 20–35°C under autotrophic culture conditions (Gouveia et al., 1995). Carotenoids accumulation in *C. vulgaris* is affected by various factors such as light, salinity, nitrogen and phosphorus limitation (Gouveia et al., 1995; Zhang et al., 2017). Carotenoid pigments and their contents were reported to vary under different growth conditions (Safi et al., 2014; Abreu et al., 2012). *C. vulgaris* was grown in flat-plate photo-bioreactor, tubular photo-bioreactor and column photo-bioreactor for biomass, lipids, fatty acids, and pigments (Safi et al., 2014). Degen et al.,

Table 3. Possible carotenoid pigments from microalgae species.

Microalgae	Carotenoid pigments	References
<i>S. platensis</i> <i>H. pluvialis</i>	Lutein, zeaxanthin, β -carotene, Astaxanthin, astaxanthin esters, lutein, zeaxanthin, α -carotene, β -carotene, violaxanthin,	Hanaa, 2003; Ranga Rao et al., 2010a; 2013a Sandesh Kamath et al., 2008; Ranga Rao et al., 2010a; 2013b
<i>D. salina</i>	Phytoene, β -carotene, lutein, zeaxanthin, <i>cis</i> and <i>trans</i> isomers,	Borowitzka and Borowitzka, 1989; Murthy et al., 2005; Dufosse et al., 2005; Borowitzka, 2010; Tan and Suter, 2011; Yang et al., 2013.
<i>Chlorella</i> sps; <i>Chlorella vulgaris</i> ; <i>Chlorella protothecoides</i>	β -carotene, lutein, antheraxanthin, zeaxanthin, violaxanthin, canthaxanthin, astaxanthin, chlorophyll-a, chlorophyll-b, pheophytin-a, pheophytin-b	Gouveia et al., 1995; Shi et al., 2002; Pelah et al., 2004; Del Campo et al., 2004; Cha et al., 2008; Wei et al., 2008; Koo et al., 2012; Campenni et al., 2013; Liu et al., 2014; Zhang et al., 2014; Safi et al., 2014
<i>Nannochloropsis</i> sps	Zeaxanthin, astaxanthin, canthaxanthin	Lubian et al., 2000; Rebolloso-Fuentes et al., 2001; Macias-Sanchez et al., 2005; Forjan et al., 2007; Solovchenko et al., 2014.
<i>Scenedesmus</i> sps	Lutein, canthaxanthin isomer, canthaxanthin, astaxanthin, astaxanthin isomers, β -carotene, echinenone, adonirubin	Burczyk et al., 1981; Qin et al., 2008; Ceron et al., 2008; Sanchez et al., 2008; Pirastru et al., 2012; Ho et al., 2014.
<i>Chlorococcum</i> sps	Astaxanthin, astaxanthin esters, lutein,	Zhang and Lee, 1997; Zhang et al., 1997; Masojidek et al., 2000; Yuan et al., 2002; Sivathanu and Palaniswamy, 2012
<i>B. braunii</i>	Violaxanthin, astaxanthin, lutein, zeaxanthin, α -carotene, β -carotene	Grung et al., 1989; 1994; Okada et al., 1996; 1997; 1998; Muntean et al., 2008; Ranga Rao et al., 2010a; 2010b.
Diatoms- <i>Phaeodactylum tricornutum</i> , <i>Hasle ostrearia</i> ; <i>Haslea Karadagensis</i>	Fucoxanthin, β -carotene, diadinoxanthin, diatoxanthin, violaxanthin, antheraxanthin, zeaxanthin, marennine, chlorophyll-a, chlorophyll-c,	Kraay et al., 1992; Prestegard et al., 2009; Bozarth et al., 2009; Peng et al., 2011; Gastineau et al., 2012; 2014; Kuczynska et al., 2015; Fu et al., 2015; Guo et al., 2016; Lelyana, 2016;

Table 4. Carotenoid content in the selected microalgae species.

Algal species	#Pigment (%)	#Major pigment or total carotenoids	References
<i>Haematococcus pluvialis</i>	3–7%	AX	Hata et al., 2001; Steinbrenner et al., 2001; Kang et al., 2005; Nobre et al., 2006; Ceron et al., 2007; Ranga Rao et al., 2011; 2014a; Regnier et al., 2015;
<i>Chlorella vulgaris</i>	12.5% TC	AX	Mendes et al., 1995; Cha et al., 2008
<i>Chlorella vulgaris</i>	55.5% TC	AX	Mendes et al., 2003; Singh and Gu, 2010; Chacon-Lee et al., 2010; Cha et al., 2010
<i>Chlorella zofingiensis</i>	0.7%	AX	Bar et al., 1995
<i>Coelastrella striolata</i> Var. <i>multistriata</i>	0.15%	AX	Abe et al., 2007
<i>Dunaliella salina</i>	3–13%	BC	El-Baz et al., 2002;
<i>Chlorella zofingiensis</i>	0.9%	BC	Bar et al., 1995
<i>Coelastrella striolata</i> Var. <i>multistriata</i>	0.7%	BC	Abe et al., 2007
<i>Spirulina platensis</i>	70–80% TC	BC	Miranda et al., 1998; El-Baky et al., 2003; Jaime et al., 2005; Ranga Rao et al., 2010
<i>Chlorella pyrenoidosa</i>	0.2–0.4%	LT	Wu et al., 2007
<i>Botryococcus braunii</i>	0.16%	LT	Tonegawa et al., 1998
<i>Botryococcus braunii</i>	75% TC	LT	Ranga Rao et al., 2006; 2007a; 2010a; 2010b; 2013b
<i>Chlorella vulgaris</i>	45% TC	LT CX	Mendes et al., 2003; Singh and Gu, 2010; Chacon-Lee et al., 2010; Cha et al., 2010
<i>Phaeodactylum tricornutum</i>	1.65%	FX	Ragni et al., 2007; Kim et al., 2012; Dambek et al., 2012
<i>Isochrysis aff. galbana</i>	1.8%	FX	Kim et al., 2012
<i>Cylindrotheca closterium</i>	0.5%	FX	Rijstenbil et al., 2003
<i>Odontella aurita</i>	2.2%	FX	Xia et al., 2013
<i>Coelastrella striolata</i> Var. <i>multistriata</i>	4.7%	CX	Abe et al., 2007
<i>Chlorella zofingiensis</i>	25% TC	CX	Bar et al., 1995
<i>Chlorella vulgaris</i>	36% TC	CX	Li et al., 2002; Singh et al., 2010; Chacon-Lee et al., 2010; Kong et al., 2012
<i>Botryococcus braunii</i>	0.17%	ECN	Tonegawa et al., 1998
<i>Nannochloropsis</i> sps	0.1%	TC	Lubian et al., 2000; Macias-Sanchez et al., 2005; Forjan et al., 2007; Nobre et al., 2013; Solovchenko et al., 2014
<i>Scenedesmus</i> sps	0.69%	TC	Qin et al., 2008; Ceron et al., 2008; Pirastru et al., 2012; Chan et al., 2013; Guedes et al., 2013; Ho et al., 2014;
<i>Chlorococcum</i> sps	0.25%	TC	Zhang and Lee, 1997; Zhang et al., 1997; Masojidek et al., 2000; Yuan et al., 2002; Sivathanu and Palaniswamy, 2012;

carotenoid contents in algae species varied upon their culture conditions; AX, astaxanthin; BC, β -carotene; LT, lutein; CX, canthaxanthin; ECN, echineone; TC, total carotenoids.

(2001) reported biomass productivity of 0.11 g/L/h when *C. vulgaris* was cultured in a flat panel photobioreactor under continuous light. High biomass productivity was obtained (up to 0.25 g/L/d) when *C. vulgaris* was grown in stirred bioreactor (Ogawa and Aiba, 1981; Martinez et al., 1991; Liang et al., 2009). *Chlorella* was widely cultivated for health food and nutrition supplements (Merchant and Andre, 2001; Merchant et al., 2002; Kitada et al., 2009). *Chlorella* contains very rigid cell wall and hence cells are processed in order to break the cell wall before using it as a food supplement in various applications (Tokusoglu and Unal, 2003). The bioavailability of carotenoids from *Chlorella* in humans is yet to be confirmed. However, as a good source of pro-vitamin A, the carotenoids from *Chlorella* could be converted in to vitamin A in humans.

4.2.4. *Nannochloropsis* sps

Nannochloropsis is a unicellular photosynthetic microalga with diameter of 2–3 μ m and is found in marine, fresh and brackish waters (Brown, 1987). These species are non-motile, and cannot be distinguished by either light or electron microscopy. The characterization of this alga is done by *rbcl* gene and 18S rDNA sequence analysis (Starkenbourg et al., 2014). The alga is known to produce various compounds such as fatty acids, lipids, including pigments such as zeaxanthin, astaxanthin and canthaxanthin (Reboloso-Fuentes et al., 2001; Solovchenko et al., 2014). *Nannochloropsis* is considered as an important

natural source in various applications due to its high essential fatty acids content (Hu and Gao, 2006; Gog et al., 2012; Nobre et al., 2013). *Nannochloropsis* is used as feed for fish larvae and rotifers (Bae and Hur, 2011).

4.2.5. *Scenedesmus* sps

Scenedesmus is a green microalga and is investigated for its possible applications in fish feed, as supplement in human nutrition, and in production of pharmaceuticals (Yaakob et al., 2014a; Roy and Pal, 2015). It is one of the dominant species in freshwater bodies, lakes, and rivers (Mandotra et al., 2014). *Scenedesmus* sps are cultivated in continuous and semi-continuous culture mode using raceway and photobioreactors for obtaining high quantities of bioactive molecules (Wu et al., 2013; Abomohra et al., 2014). Physiological changes were observed in the cells under various culture conditions (Pancha et al., 2014). Under stress conditions, *Scenedesmus* accumulates β -carotene, astaxanthin isomers along with lutein and canthaxanthin (Burczyk et al., 1981; Ceron et al., 2008; Qin et al., 2008; Sanchez et al., 2008; Pirastru et al., 2012; Ho et al., 2014).

4.2.6. *Chlorococcum* sps

Chlorococcum is a green microalga, belonging to *Chlorophyta*, and is found in freshwater bodies and lakes. Under stress conditions *Chlorococcum* culture colour changes from green to orange as the cells accumulate secondary carotenoids (Zhang

Table 5. Culture conditions for carotenoid production in the selected microalgae using various photobioreactors.

Algal species	Culture conditions	Mode of cultivation and reactor	^a Carotenoid productivity (mg/m ² /d or mg/L/d or mg/g/L/d)	Reference
<i>Haematococcus pluvialis</i>	T: 28°C; LI: 345 μ mol photons/m ² /s Day light cycle T: 15–25°C; LI: 2000 μ mol photons/m ² /s T: 20–25°C; aeration: 1.5%v/v CO ₂ ; day light T: 25°C; pH: 7.5 \pm 0.5; LI: 281 \pm 89 μ mol photons/m ² /s	Batch (1L) Continuous chemostat, Tubular PBR (50 L) Enclosed PBR, outdoor (25000 L) Glass column, batch mode; outdoor Semi-continuous outdoor, closed tubular PBR (55 L)	AX: 98 mg/g AX: 8.0 mg/L/d AX: 13 g/m ² /d AX: 17.1 mg/L/d TC: 102.5 \pm 33.1 mg/m ² /d;	Dominguez-Bocanegra et al., 2004; García-Malea et al., 2009 Olaizola, 2000 Wang et al., 2013 García-Gonzalez et al., 2005
<i>Dunaliella salina</i>	T: 30°C; pH: 7.5; LI: 200–1200 μ mol photons/m ² /s T: 30°C; pH: 7.5; LI: 200–1200 μ mol photons/m ² /s	Continuous turbidostat, flat-panel PBR (2.5 L) Continuous turbidostat flat-panel PBR (1.9L) with in situ extraction	BC: 13.5 mg/L/d BC: 0.7–8.3 mg/L/d	Del Campo et al., 2001 Del Campo et al., 2001
<i>Chlorella protothecoides</i> <i>Chlorella zofingiensis</i>	T: 28°C; pH: 6.5; LI: absence of light; heterotrophic T: 28°C; pH: 7; LI: 200 μ mol photons/m ² /s; AF: 50–100 L/h (1% v/v CO ₂) T: 30°C; pH: 6.5; LI: darkness; SR: 130 rpm; heterotrophic	Batch (16 L) Batch (0.2 L) Batch (250 mL)	LT: 10 mg/L/d LT: 3.4 mg/L/d AX: 1.03 mg/L	Wei et al., 2008 Kleinegris et al., 2011 Ip and Chen, 2005
<i>Scenedesmus almeriensis</i>	T: 30°C; pH: 7.5; LI: 1700 μ E/m ² /s; AF: 0.5 (v/v)/min/s; or T: 35°C; LI: 1900 μ E/m ² /s	Continuous (2L) or Continuous outdoor, tubular PBR	LT: 4.9 mg/L/d; LT: 5.31 mg/m ² /d	Sanchez et al., 2008
<i>Chlorococcum citriniforme</i>	T: 28°C; pH: 7; LI: 200 μ mol photons/m ² /s; AF: 50–100 L/h (1% v/v CO ₂)	Batch (0.2 L)	LT: 1.05 mg/L/h	Kleinegris et al., 2011
<i>Muriellopsis</i> sp.	T: 28°C; pH: 6.5; LI: 460 μ mol photons/m ² /s; T: 28°C; pH: 7.0; LI: continuous 200 μ mol photons/m ² /s; AF: 50–100 L/h (1% v/v CO ₂)	Batch (0.2 L) Continuous outdoor, tubular PBR (55 L); Semi continuous outdoor, open tank (100 L)	LT: 5.5 mg/g/L/d LT: 0.8–1.4 mg/L/d LT: 5.5 mg/g/L/d or LT: 100 mg/g/L/d	Kleinegris et al., 2011 García-Gonzalez et al., 2005

^aCarotenoid productivity in algal species varied upon their culture conditions; T, temperature; LI, light irradiance; AF, air flow; PBR, photobioreactors; TC, total carotenoids; AX, astaxanthin; LT, lutein; BC, β -carotene;

and Lee, 1997; Masojidek et al., 2000). *Chlorococcum* cells contain free astaxanthin, astaxanthin esters, free adonixanthin, adonixanthin esters, canthaxanthin, lutein, β -carotene and *cis*-isomers of keto-carotenoids (Zhang et al., 1997; Yuan et al., 2002). *Chlorococcum* produces high amounts of carotenoid esters like *Haematococcus pluvialis* (Ranga Rao et al., 2013a). Astaxanthin esters, free canthaxanthin and adonixanthin are the major pigments in *Chlorococcum* (Zhang and Lee, 1997; Sivathanu and Palaniswamy, 2012). *Chlorococcum* cells may be able to synthesize carotenoids from β -carotene via canthaxanthin and adonixanthin pathway (Yuan et al., 2002).

4.2.7. *Botryococcus braunii*

B. braunii is a Chlorophycean microalga and is found usually in fresh water reservoirs, ponds and brackish lakes (Metzger and Largeau et al., 2005). It is grouped into race 'A', race 'B', and race 'L' based on hydrocarbons they produce (Banerjee et al., 2002; Ranga Rao et al., 2016). Apart from hydrocarbons, *B. braunii* produces carotenoids, lipids, and exopolysaccharides (Ranga Rao et al., 2007a, 2007b, 2012a, 2013a; Bayona and Garcés, 2014). Accumulation of carotenoids in race 'A', 'B' and 'L' of *B. braunii* was reported (Grung et al., 1989, 1994; Okada et al., 1996, 1997, 1998; Muntean et al., 2008; Ranga Rao et al., 2010). *B. braunii* culture changes to brown, red, orange, and pale yellow from green due to the carotenoids accumulation in the cells (Ranga Rao et al., 2016). Braunixanthin-1, braunixanthin-2, α -botryoxanthin, botryoxanthin-A, and botryoxanthin-B were isolated from 'B' race (Okada et al., 1996, 1991, 1998). Total carotenoids content in race 'A' was 0.28% as reported by Ranga Rao et al. (2010b). Carotenoid pigments such as lutein, violaxanthin, zeaxanthin, β -carotene and astaxanthin were reported in 'A' race (Muntean et al., 2008), and Ranga Rao et al. (2006) reported lutein as major pigment in 'A' race. Recently, our research group reported carotenoid production and other molecules of interest from *B. braunii* for commercial applications (Ranga Rao et al., 2017).

4.2.8. *Diatoms*

Diatoms are prominent microalgal forms containing chlorophylls, carotenoids, fatty acids, lipids, and polysaccharides. The production of lipids, polysaccharides and pigments are regulated by abiotic stress factors and genetic modification of metabolic pathways (Perl et al., 1990; Lee et al., 2006; Prestegard et al., 2009; Peng, 2014). Diatoms have novel metabolic pathways in the evolutionary history of eukaryotes and show ecological success (Kuczyńska et al., 2015). Diatoms are linked with the silicon metabolism involved in the biogeochemical cycling of silicon (Si) in the sea. The siliceous structures of their cell wall create unique morphologies which are used as taxonomic keys (Martin-Jezequel et al., 2000). Diatom cell physiology displays a significant difference in stationary, exponential and declining growth phases (Vidoudez and Pohnert, 2012). In both stationary and exponential growth phases, the influence of blue-green light and white light on accumulation of carotenoids in diatom species was reported (Sanchez-saavedra and voltolina, 2002; Brunet et al., 2014). Diatom cells accumulate high content of carotenoids under various stress conditions such as low and high irradiance, salinity, and nitrogen limitation (Xia et al., 2013). Diatoms contain chlorophylls and carotenoids

which play a major role in photo-protection (Jeffrey and Veski, 1997). The major component such as chlorophyll-a and chlorophyll-c together with fucoxanthin form fucoxanthin-chlorophyll protein complexes in diatoms species (Gelzinis et al., 2015). Fucoxanthin acts as light harvesting carotenoid pigment in diatoms which is usually found more in brown algal species. Carotenoids such as fucoxanthin, diadinoxanthin, diatoxanthin, zeaxanthin, β -carotene, astaxanthin, and violaxanthin were found in diatoms (Kuczyńska et al., 2015). The carotenoid content in diatoms is different from that found in other algal species. The carotenoid composition is similar in various diatom species; however their levels among the species are varied. Carotenoid pigments from diatoms are used in health food applications (Bozar et al., 2009; Fu et al., 2015; Kuczyńska et al., 2015). Changes in carotenoid content in diatom species under various culture conditions is presented in Table 6.

5. Extraction and processing of carotenoid pigments

Extraction and processing of carotenoid pigments from microalga is very challenging task as it involves harvesting of algal cells, processing, extraction and purification. Harvesting, analysis, extraction, purification and downstream processing of carotenoids from microalgae biomass are very important for commercialization process. Some of the microalgal cell walls are thick resisting easy disruption. Cell wall disruption can be achieved using any one or more of the several methods-mortar and pestle, two-phase extraction, milling, ultra-sonication, microwave, thawing, freezing, supercritical fluid extraction and also with edible oils. The selection of suitable extraction method depends on the specific algal species (McMillan et al., 2013). Carotenoids were extracted from *D. salina*, *H. pluvialis*, *S. platensis*, *B. braunii* and *Chlorococcum* sps cells with organic solvents using mortar and pestle (Zhang and Lee, 1997; Murthy et al., 2005; Sarada et al., 2006; Ranga Rao et al., 2013b). This method suits for bench scale process and not for scale-up process of industrial applications. Carotenoids extracted from the algal biomass using two phase extraction method of alkaline treatment were reported in different algal species (Sarada et al., 2006; Ceron et al., 2008; Ishida and Chapman, 2009). Extraction with ethanol and hexane allows easy removal of residual solvent and yielding extracts with high lutein content (Fernandez-Sevilla et al., 2010). Astaxanthin was extracted using dodecane and methanol solvent extraction method, and in this process free astaxanthin was recovered up to 85% (Kang and Sim, 2007). Another extraction method was developed for astaxanthin from *H. pluvialis* using vegetable oils. 90% of astaxanthin was recovered in olive oil extraction method (Kang and Sim, 2008). Carotenoid extraction with solvents was standardized to meet commercial specifications. However selective precipitation using supercritical CO₂ is a promising alternative. Carotenoids extraction from algal biomass is time consuming with many steps and requires large quantities of solvents which are expensive and harmful. Supercritical carbon-dioxide extraction method provides an efficient method for carotenoid extraction. This is a very rapid, efficient, non-flammable, non-toxic and an inexpensive extraction method. Carotenoids were extracted from various microalgal biomasses using supercritical carbon-dioxide (CO₂), with satisfactory results (Mendes et al.,



Table 6. Carotenoid pigments changes in diatoms species upon culture conditions.

Diatoms species	Culture conditions	Chl-a	Chl-c	BC	FX	DDX	DX	VX	References
<i>Cyclotella meneghiniana</i>	140 $\mu\text{mol photons/m}^2/\text{s}$; 40 $\mu\text{mol photons/m}^2/\text{s}$; 16 h; 8 h (L:D)	-	++	++	++	+	+	-	Alexandre et al., 2014
<i>Cyclotella meneghiniana</i>	140 $\mu\text{mol photons/m}^2/\text{s}$; Iron rich or replete medium	+	-	-	-	+++	+++	-	Beer et al., 2011
<i>Cyclotella meneghiniana</i>	40 $\mu\text{mol photons/m}^2/\text{s}$; Iron rich or replete medium	+	-	-	-	++	++	-	Brotas and Plante-Cuny, 2003
<i>Pseudo-nitzschia multistriata</i>	450 PFD with red and blue light ratio 0.25	+++	+	++	+++	+++	+++	-	Brunet et al., 2014
<i>Pseudo-nitzschia multistriata</i>	250 PFD with red and blue light ratio 0.25	+++	+	++	+++	+++	+++	-	Costa et al., 2013
<i>Phaeodactylum tricornutum</i>	300 $\mu\text{mol photons/m}^2/\text{s}$; 50 $\mu\text{mol photons/m}^2/\text{s}$; 14 h; 10 h (L:D)	-	+++	+	+++	+	+	-	Brunet et al., 2014
<i>Phaeodactylum tricornutum</i>	24 or 40 $\mu\text{mol photons/m}^2/\text{s}$ with blue and white light,	+	++	++	++	++	-	+++	Costa et al., 2013
<i>Phaeodactylum tricornutum</i>	40 or 41 $\mu\text{mol photons/m}^2/\text{s}$ with red and white light,	+	+++	+	+++	+++	-	+++	Costa et al., 2013
<i>Phaeodactylum tricornutum</i>	1250 $\mu\text{mol photons/m}^2/\text{s}$; 40 $\mu\text{mol photons/m}^2/\text{s}$; 12 h; 12 h (L:D)	+++	++	++	++	+++	+	-	Domingues et al., 2012
<i>Phaeodactylum tricornutum</i>	24 or 40 $\mu\text{mol photons/m}^2/\text{s}$ with blue and red light	+	-	-	-	+++	-	-	Jungandreas et al., 2014
<i>Phaeodactylum tricornutum</i>	24 or 40 $\mu\text{mol photons/m}^2/\text{s}$ with blue and red light	++	-	-	-	+	-	-	Jungandreas et al., 2014
<i>Cyclotella meneghiniana</i>	700 or 40 $\mu\text{mol photons/m}^2/\text{s}$; 16 h; 8 h (L:D)	-	-	-	-	+++	+++	+++	Lohr et al., 1999
<i>Odontella aurita</i>	100 $\mu\text{mol photons/m}^2/\text{s}$ with low or high nitrogen culture	-	-	-	+++	-	-	-	Xia et al., 2013

*Carotenoid pigments changes in diatoms species upon their culture conditions; Chl-a, chlorophyll-a; Chl-c, chlorophyll-c; BC, β -carotene; FX, fucoxanthin; VX, violaxanthin; DDX, diadinoxanthin; DX, diatoxanthin; "+", increase; "++", increase; "+++", increase; "L", light; "D", dark; PFD, photon flux density.

1995; Careri et al., 2001; Macias-Sanchez et al., 2005; 2007; 2009). 80% of astaxanthin was recovered from *Haematococcus* cells when treated with hydrochloric acid at 80°C for 2 min. Similarly 70% of astaxanthin was recovered when cells were treated with 40% acetone at 80°C for 2 min (Kobayashi et al., 1997; Sarada et al., 2006). 95% of fucoxanthin in *Isochrysis galbana* was recovered using ethanol solvent extraction method (Kim et al., 2012).

Downstream processing is the most important step for algal carotenoid production and various strategies are available (Gong and Bassi, 2016). For cell harvesting, physical (Grima et al., 2003; Chen et al., 2011) or chemical methods (Uduman et al., 2010; Gorin et al., 2015; Hu et al., 2013; Pragya et al., 2013; McMillan et al., 2013; Utomo et al., 2013) can be used. Several extraction methods such as grinding (Hu et al., 2013), cryogenic grinding (Grima et al., 2003; Zheng et al., 2011), bead milling (Chan et al., 2013; Taucher et al., 2016), high pressure homogenizer (Kim et al., 2015), autoclave (Chan et al., 2013), microwave (McMillan et al., 2013; Pasquet et al., 2011; Zheng et al., 2011), enzymatic hydrolysis (Deenu et al., 2013; Kadam et al., 2013), pulsed electric field (Lai et al., 2014), alkaline treatment (Halim et al., 2012a; 2012b), ionic liquids (Park

et al., 2015), conventional solvent extraction (Gil-Chavez et al., 2013), and supercritical carbon dioxide (Guedes et al., 2013; Yen et al., 2015; Liao et al., 2010) are used in carotenoid extraction. For cost effective downstream processing of carotenoids for food, feed and health application, an efficient method is to be adopted/developed based on the literature. Figure 2 presents downstream processing of carotenoids from algae for various applications.

6. Industrial application of carotenoid pigments

Carotenoids are natural pigments that occur widely in nature and shown to be beneficial in human health (Rao and Rao, 2007; Wells et al., 2016; Cooperstone and Schwartz, 2016). Various health benefits of carotenoids cover different ailments such as cancer (Sharoni et al., 2012), degenerative diseases (Ma et al., 2012; Aronow and Chew, 2014), cardiovascular diseases (Gaziano et al., 1995; Omenn et al., 1996), diabetes (Luvizotto et al., 2013), ulcer (Zhou et al., 2016). They also known to have ultraviolet light protection property (Stahl et al., 2000; Cesarini et al., 2003; Stahl and Sies, 2012). Carotenoids are bioactive compounds and play a part in important biological activities




Figure 2. Processing of microalgal carotenoid pigments for commercial applications.

like bioavailability process (Castenmiller and West, 1998; Cooperstone et al., 2015), pro-vitamin A activity (Ross and Harrison, 2014), regulation of gene transcription (Palozza et al., 2003, 2012), induction of phase-II enzymes (Sharoni et al., 2004; Lian and Wang, 2008), antioxidant activity (Krinsky, 1989, 1991), alteration of immune function (Burton-Freeman et al., 2012), and also enhancement of gap junction communication (Stahl et al., 1997). Carotenoids are used in feed, food, and cosmeceutical applications (Vilchez et al., 2011; Tominaga et al., 2012; Ranga Rao et al., 2013a; Zhang et al., 2014; Yaakob et al., 2014a). Currently, the world demand for natural carotenoid pigments is increasing. Biotechnological process of β -carotene, astaxanthin, lutein, fucoxanthin and other pigments has high demand in the global market (www.bccresearch.com). Natural carotenoid pigments are preferred by the health market because they are mixture of cis/trans isomer forms, esters, and xanthophylls with high biological properties (Murthy et al., 2005; Hsu et al., 2008; Ranga Rao et al., 2013b). Microalgae rich in carotenoids are used as food colorants, food additives, and vitamin supplements (Zhang et al., 2014; Ranga Rao et al., 2014a). The most widely used algal species for this are *S. platensis*, *H. pluvialis*, *D. salina*, *B. braunii*, *Chlorella* sps, *Chlorococcum* sps, and *Scenedesmus* sps. Various cultivation systems (photo-bioreactors and raceway ponds) are used for mass cultivation of algae to obtain higher biomass yield and carotenoids for commercial applications (Olaizola, 2000; Del Campo et al., 2007; Ugwu et al., 2008; Chen et al., 2011; Ranga Rao et al., 2012a, 2012b; Havlik et al., 2013; Klok et al., 2014; Poonkum et al., 2015). Algae accumulate pigments in their biomass and also offer biological activities such as anti-oxidant activity, anti-cancer activity and anti-bacterial activity (Plaza et al., 2008, 2009; Buono et al., 2014). Among the algae species, β -carotene in *S. platensis*; β -carotene in and *D. salina*; astaxanthin and its esters in *H. pluvialis* and *Chlorococcum* sps; lutein in *Chlorella* sps and *B. braunii*; and fucoxanthin in diatoms have been used for commercial applications. Carotenoid pigments from algal species have been evaluated for biological properties in *in vitro* and *in vivo* models (Dufosse et al., 2005; Murthy et al., 2005; Vilchez et al., 2011; Zhang et al., 2014; Ranga Rao et al., 2014a).

Spirulina biomass is very interesting in food applications because of its nutritional properties (Eriksen, 2008; Hoseini et al., 2013). It is used as food supplements for humans, and as feed for animals as it contains nutritional constituents (Holman and Malau Adlui, 2013). It is also used in animal feed for pigmentation and for improvement of integumentary color of cultured fishes such as tilapia and sweet melt (Yaakob et al., 2014a). *S. platensis* has been used as a feed component in quality broilers and layer diets to enhance yolk color and flesh (Tan and Suter, 2011; Yaakob et al., 2014a). In *S. platensis*, β -carotene has been found to be the major carotenoid pigment (Del Campo et al., 2007). β -carotene is an important carotenoid which acts as pro-vitamin-A. It is used as color additive in foods and as an antioxidant. β -carotene from natural sources showed potential antioxidant properties compared to synthetic carotene (Murthy et al., 2005). *S. platensis* and *D. salina* are two algae which are the rich sources for β -carotene (Del Campo et al., 2007). Products derived from *D. salina* are either whole cell biomass or β -carotene extracts and are used for

human consumption and also for animal feed (Spolaore et al., 2006; Gouveia et al., 2006). β -carotene from *D. salina* is used as natural colorant in foods and cosmetics (Hosseini and Shariati, 2009). β -carotene is the most efficient quencher and involved in the scavenging of reactive oxygen species, peroxy radicals, singlet oxygen against lipid peroxidation (Stahl et al., 2005). It is also an efficient stimulator of gap junctional communication associated with cell differentiation, growth, apoptosis and also inhibition of lipid peroxidation (Palozza and Krinsky, 1992).

H. pluvialis, *Chlorella* sps and *Chlorococcum* sps are rich source for astaxanthin and its esters (Zhang et al., 1997; Gouveia et al., 2006; Ranga Rao et al., 2013b; Liu et al., 2014). Astaxanthin is a very valuable pigment, used as pigmentation source in aquaculture, nutraceutical, feed and food industries (Tominaga et al., 2012; Buono et al., 2014). Astaxanthin showed potential biological activities in *in vitro* and *in vivo* models (Sandeesh Kamath et al., 2008; Ranga Rao et al., 2014a). Recently our research group has investigated biological activities of astaxanthin and astaxanthin esters against skin cancer in rat model, and the results showed better anticancer activity in both astaxanthin and astaxanthin esters treated animals (Ranga Rao et al., 2013a).

Chlorella sps are used as food supplements with various applications (Liu et al., 2014). It is used as nutritional supplements to the treatment of hypertension, fibromyalgia, and ulcerative colitis (Merchant and Andre, 2001). *Chlorella* sps showed potential antioxidant and anticancer properties in cell culture models (Gouveia et al., 2006; Cha et al., 2008). *C. vulgaris* is an important microalga used in biofuel, human nutrition, animal feed, food colorant, emulsifier agent, pharmaceuticals, and bio-fertilizers (Safi et al., 2014). *C. vulgaris* products are available in the form of capsules, tablets, extracts and powder in the market (Yamaguchi et al., 1996). *C. vulgaris* is rich in fatty acids, polysaccharides, pigments, vitamins, proteins, and is important in its use for nutraceutical purpose (Grobbelaar, 2003). *C. vulgaris* gives a better preservative activity compared to butylated hydroxytoluene and butylated hydroxyanisole (Rodriguez-Garcial and Guil-Guerrero, 2008). *C. vulgaris* accumulates pigments used as animal feed and showed an interesting pigmentation property for fish flesh and egg yolk in poultry industry (Gouveia et al., 1996; Gouveia et al., 2002). *C. vulgaris* showed protective activity against heavy metals and also harmful compounds by reducing oxidative stress in the tested animals (Vijayavel et al., 2007; Shim et al., 2008; Yun et al., 2011). The liquid extract of *C. vulgaris* used as bio-fertilizer in plants. It improved the seed germination, plant growth and productivity (Gonzalez and Bashan, 2000; Faheed and Abdel Fattah, 2008).

Nannochloropsis sps is used for nutraceutical applications as it contains vitamins, minerals, pigments, and polyunsaturated fatty acids (Markovits et al., 1992; Bishop and Zubeck, 2012; Nobre et al., 2013; Kent et al., 2015). *Scenedesmus* sps biomass is used as nutritional supplement for various applications (Ho et al., 2014; Kumar et al., 2015; Kent et al., 2015). Further, *Scenedesmus* sps is considered as potential source of feed stock for biofuel applications (Mandotra et al., 2014). *Chlorococcum* sps. is rich in carotenoids, polyphenols, fatty acids and showed potential biological activities in *in vitro* and *in vivo* models (Bhagavathy et al., 2011a, 2011b, 2012). Lutein accumulation in *B. braunii* and *Muriellopsis* sps. is exploited for commercial application (Del Campo et al., 2000, 2004, 2007; Ranga Rao

et al., 2010a; 2010b). Lutein and zeaxanthin are responsible for the coloration of the macula lutea in the retina, and are extremely important in ophthalmology for protection against age-related macular degeneration (Krinsky et al., 2003). *B. braunii* extracts showed potential anti-bacterial properties (Ranga Rao et al., 2010c). Antioxidant enzymes levels in rats were increased after feeding *B. braunii* as source of lutein (Ranga Rao et al., 2006, 2010a, 2013b).

Diatoms have great application in pharmaceutical, nutraceutical, and cosmeceutical industry (Perl et al., 1990; Lee et al., 2006; Prestegard et al., 2009) as they contain good quantities of chlorophylls, carotenoids, blue pigments, amino acids and fatty acids. There is much interest created for use of diatoms in health foods, pharmaceuticals, biofuels, and bioremediation applications (Bozarth et al., 2009). Diatom species also accumulates significant amount of fucoxanthin. Fucoxanthin is a carotenoid pigment with very high commercial value in the global market, and is used in biological activities such as anti-obesity, anti-oxidant, anti-angiogenic, anti-inflammatory, anti-cancer, photoprotective and neuro-protective activity (Pangestuti and Kim et al., 2011). Fucoxanthin may inhibit reactive oxygen formation, DNA damage, and apoptosis induced by hydrogen peroxide, and its antioxidant activity was comparable to α -tocopherol (Peng et al., 2011). Fucoxanthin showed anti-proliferative activity in human cell lines such as HL-60, PC-3, HT-29, DLD-1 and Caco-2 (Hosokawa et al., 1999; Asai et al., 2004). Fucoxanthin has the ability to reduce the nitric oxide levels, prostaglandin-E2, interleukin-6, tumor necrosis factor- α , histamine and interleukin-1 β in *in vivo* models (Heo et al., 2008; Kim et al., 2010). Fucoxanthin is a promising bioactive compound which may help in the prevention of obesity through upregulation of uncoupling protein-1 (UCPI) which stimulates energy expenditure by thermogenesis (Maeda et al., 2005). Diatoms are rich in naviculan and domoic acid which showed anti-viral, neuroexcitatory, cytotoxic and blood platelet inhibitory activity (Perl et al., 1990; Lee et al., 2006; Prestegard et al., 2009). Marine diatom species such as *Haslea ostrearia* and *Haslea karadagensis* contain marennine (blue pigment) and showed anti-bacterial activity, anti-viral activity, and anti-oxidant activity in *in vitro* and *in vivo* experimental models (Gastineau et al., 2014). These unusual pigments

accumulate in select diatoms species and these pigments have great potential applications in cosmetics and food supplements. Pigments from diatoms showed a major role in photo-protection (Wichuk et al., 2014; Fu et al., 2015; Kuczyńska et al., 2015). Benefits of carotenoid pigments from algae are presented in Table 7.

7. Carotenoids market value and its manufactures

The importance and use of synthetic carotenoid pigments started declining due to their potential toxic effects. Nowadays, a transition has been stimulated towards green solutions and natural products and global market for natural products is estimated to reach \$1.5 billion by 2020. One of the safe alternatives to synthetic pigments is carotenoids from microbial source (Dufosse et al., 2005; Yang et al., 2013). Naturally derived carotenoid pigments from microbial sources such as fungi, bacteria and algae have been exploited for commercial applications (Borowitzka, 2010). Carotenoid pigments such as astaxanthin, β -carotene, fucoxanthin and lutein from algae have received much more attention as they are shown to have potential use in nutraceuticals, pharmaceuticals, food, animal feed, dietary supplements, and cosmetics sector (Ranga Rao et al., 2014a; Bilbao et al., 2016; Lelyana, 2016). For nutritional importance, astaxanthin is considered as one of the best powerful antioxidant in nature that plays a major role in scavenging free radicals in human body (Roche, 1987; Guerin et al., 2003; Hussein et al., 2006; Park et al., 2010; Kent et al., 2015). The market value of carotenoids was expected to reach \$1.53 billion by 2021. The demand for astaxanthin, β -carotene and lutein is now emerging in the global market. Astaxanthin is consumed by the salmon feed industry and the annual market value of the pigment is at \$200 million with cost of \$2500 per kg. Astaxanthin market for animal feed was \$300 million, and for nutraceutical (as an antioxidant agent) was \$30 million in 2009. It is expected to increase to \$800 million and \$300 million by 2020 for animal feed and for nutraceuticals respectively. Market value of β -carotene and lutein is expected to reach \$334 and \$309 million by 2018 respectively. The global market for fucoxanthin production was about 500 tonnes in 2015, and it is expected to reach at 5.3% CAGR by 2021 (www.marketresearchstore.com).

Table 7. Benefits of carotenoid pigments from microalgae.

Carotenoid pigments	Benefits	References
Astaxanthin	Anti-oxidants, protect from UV rays, bioavailability, food colorant, animal feed, enhance immune functions, improve eye/skin health, anti-aging property, pharmaceutical, nutraceutical, cosmeceutical	Lorenz and Cysewski, 2000; Guerin et al., 2003; Higuera-Ciapara et al., 2003; Hussein et al., 2006; Lu et al., 2010; Park et al., 2010; Ranga Rao et al., 2010a; 2011; Choi et al., 2011; Kidd, 2011; Kim et al., 2011; Chan et al., 2012; Maoka et al., 2012; Chew et al., 2013; Dong et al., 2013; Huangfu et al., 2013; Ishiki et al., 2013; Yamashita, 2013; Yang et al., 2013; Ranga Rao et al., 2013a; 2014a;
β -carotene,	Vitamin A precursor, anti-oxidant, food color, malnutrition, prevent macular degeneration, skin burn from UV rays, depression, asthma, infertility, psoriasis, high blood pressure, pharmaceutical, cosmetics, animal feed, fish feed, pet foods	Fujii et al., 1993; Ben Amotz and Levy, 1996; Curtain, 2000; Kar, 2002; Murthy et al., 2005; Chidambara Murthy et al., 2005; Dufosse et al., 2005; Shaish et al., 2006; Mogedas et al., 2009; Ranga Rao et al., 2010a; Weber and Grune, 2012; Jayappriyan et al., 2013; Zhang et al., 2014; Donhowe and Kong, 2014;
Lutein, Zeaxanthin	Anti-oxidant, food colorant, fish feed, eye disease, health, age related macular degeneration, cataracts, pharmaceutical, nutraceutical, cosmetics, animal feed.	Landrum and Bone, 2001; Ranga Rao et al., 2006; Ceron et al., 2008; Fernandez-Sevilla et al., 2010; Ranga Rao et al., 2010a; Christaki et al., 2012; Maoka et al., 2012;
Fucoxanthin	Anti-oxidant, anti-viral, anti-diabetes, anti-obesity, anti-cancer, anti-inflammatory, anti-allergic, anti-osteoporotic	Maeda et al., 2009; Heo et al., 2010; Peng et al., 2011; Lee et al., 2013; Rokkaku et al., 2013;

As per the global carotenoids market analysis, Europe has a strong and potential market due to the increasing demand for animal feed, health supplements and Cosmetics. Involvement of leading cosmetic industries such as Unilever, L'Oreal, Henkel, and Beiersdorf are expected to improve the growth of carotenoid market value in the Europe market. Asia pacific region is projected to enhance the use of carotenoids in food supplements and animal feed in future. Several countries such as Japan, China and India are striving to improve the technology for carotenoid production economically to improve the market for carotenoids in Asia pacific region. A number of key vendors are playing a major role in producing carotenoid pigments across the globe such as Lycored, Divis Laboratories, Naturex SA, BASF Corporation, FMC Corporation, DSM Nutritional Products, Kemira Industries Inc., Allied Biotech Corp., Brenntag, Royal DSM N.V.Ch. Hansen A/S, Allied Biotech, Cognis, Carotech, D.D. Williamson & Co. Inc., Dohler Group, and ExcelVite SDN. BHD. Astaxanthin, β -carotene, fucoxanthin and whole cell biomass from algae are manufactured by various companies for commercial applications in food, feed, nutraceutical, pharmaceutical, and cosmeceutical sectors (Table 8).

8. Current trends and future prospects

Multi-steps are being considered towards screening, selection, production, cost and applications of carotenoids from microalgae. The important questions to be addressed are as follows: How to design micro-bioreactor technologies for emerging biotechnological applications to reduce bioprocess cost and time line? How to make a micro-photobioreactor design with high powered light emitting diodes (LEDs) with adjustable light intensity for massive carotenoid production and products for human health benefits? How to improve advanced sustainable

engineering tools for proteomics, genomics, metabolomics and also applied for strain development for high value chemicals? How to make sustainable mass algal culture technologies with a control system for photosynthetic organisms for efficient biomass and carotenoid productivity? How to select mixed algal culture for high value compounds for health benefits? How to control potential predators in the open mass algal culture system (raceway/circular/photobioreactors)? How to develop early detection methods of controlling predators in mass algal culture? How to enhance the carotenoid production of the selected microalgae? How to develop sustainable technologies for carotenoid production of commercial process? How to enhance high quality carotenoid pigments in the global market for human consumption? How to exploit carotenoids for drug discovery and other novel applications? New integrated emerging technologies are required for microalgae to produce bioenergy molecules for health food applications. Novel technologies are required for cell harvesting, cell disruption, and downstream processing of microalgal carotenoid pigments. To save energy cost on the freeze or oven drier, wet extraction methods will have to be developed for carotenoids from algal biomass. The understanding of carotenoid structure, function relationships will enable researchers to tailor new microalgae carotenoids for biotechnological applications. Due to the high cost of the currently available technologies for microalgal carotenoid pigments production on industrial scale, there is a requirement to develop sustainable cost effective process of the production of carotenoids to compete and replace the synthetic ones. Developments in microalgae research are expected through interaction between biotechnologists, biochemists, chemists, chemical engineers and as well as medical professionals. Micro algal carotenoids represent microorganisms capable of variety of important applications, thereby presenting a fascinating field for future research. The carotenoids from algae can be scalable in cost effective manner, so that it can be continued to expand pigment production for the developing a new nutraceutical industry.

Table 8. List of companies producing carotenoid pigments and algae biomass.

Company name	Carotenoids or Biomass	Countries
E.I.D Parry Ltd.,	Astaxanthin	India
Algae Technologies	Astaxanthin	Israel
Mera Pharmaceuticals Inc.,	Astaxanthin	USA
Cyanotech Corporation	Astaxanthin	USA
Valensa International	Astaxanthin	USA
Fuji Chemicals Industry Co., Ltd.,	Astaxanthin	Japan, Sweden
Jingzhou Natural Astaxanthin Inc.,	Astaxanthin	China
Aquacarotene Ltd.,	β -carotene	USA
Cognis Nutrition and Health	β -carotene	Australia
Nikken Sohonsa Corporation	β -carotene	Japan
Tianjin Norland Biotech Co., Ltd.,	β -carotene	China
E.I.D Parry Ltd.,	β -carotene	India
Seambiotic	β -carotene	Israel
Muradel Pty Ltd.,	β -carotene	Australia
Cyanotech Corporation	β -carotene,	USA
AlgaNova International	Fucoxanthin	China
Leili Natural Products Co., Ltd.,	Fucoxanthin	China
Cyanotech Corporation	<i>Spirulina</i>	USA
E.I.D Parry Ltd.,	<i>Spirulina</i>	India
Hydrolina Biotech Pvt., Ltd.,	<i>Spirulina</i>	India
Nutrex Hawaii Inc.,	<i>Spirulina</i>	USA
Sun Chlorella Corporation	<i>Chlorella</i>	Japan
Yaeyama Shokusan Co., Ltd.,	<i>Chlorella</i>	Japan
Maypro Industries Inc.,	<i>Chlorella</i>	USA
Taiwan Chlorella Manufacturing Co., Ltd.	<i>Chlorella</i>	Taiwan
Far East Microalgae Ind Co., Ltd.	<i>Chlorella</i>	Taiwan
Roquette Klotze GmbH and Co. KG	<i>Chlorella</i>	Germany

9. Conclusion

Use of carotenoids from microalgae for health benefits is a promising field with large economic potential. Carotenoids from natural resources have been in demand in the global market. Microalgae species produce various carotenoid pigments. Major carotenoid pigments such as astaxanthin and astaxanthin esters in *H. pluvialis*, *Chlorococcum* sps and *Chlorella* sps, β -carotene in *D. salina*, *S. platensis*, and *Scenedesmus* sps, lutein in *B. braunii*, canthaxanthin in *Nannochloropsis* sps and fucoxanthin from diatoms were reported in microalgae. These carotenoids have high demand in the global market for health food applications. Astaxanthin and β -carotene have been well recognized in prevention and treatment of various diseases. There is lack of research studies on the astaxanthin esters, *cis* and *trans* forms of carotenoids, lutein and fucoxanthin in *in vitro* and *in vivo* models. Further, biochemistry, physiology, molecular genetics of carotenoid pigments needs to be investigated which will have major impact on the development and optimization of microalgae technologies. In view of this the genes encoding enzymes that are directly involved in specific carotenoid synthesis may be assessed. Further, mass algal cultivation systems

are required to be developed for production of high quality carotenoid pigments at economic cost. Currently, carotenoid pigments from algae are a very competitive, attractive, and the most popular growing field for bio-based economy. There is a need for better engineering tools to make the process better and cost effective manner.

Declaration of interest

The authors declare that they have no conflicts of interest.

Acknowledgments

The first author thank Beijing Normal University-Hong Kong Baptist University United International College, Zhuhai Key Laboratory of Agricultural Product Quality and Food Safety (R1053 and R201632), and Vignan's Foundation for Science, Technology and Research University for providing financial support for execute this work. Authors acknowledge to reviewers and editors for providing very valuable comments and suggestions on the manuscript.

Author's contribution

ARR wrote the manuscript, GD involved in literature collection and chemical structures; GAR, RS, BPRN, BL, and SYP have been involved in improving the manuscript, and revising it critically. Finally all authors are in approval of the article submission.

Funding

Zhuhai Key Laboratory of Agricultural Product Quality and Food Safety ID: R1053,R201632.

References

- Abe, K., H. Hattori, and M. Hirano. 2007. Accumulation and antioxidant activity of secondary carotenoids in the aerial microalga *Coelastrum striolatum* var. *multistriatum*. *Food Chemistry* 100:656–661. doi:10.1016/j.foodchem.2005.10.026.
- Abomohra, A. E. F., M. El-Sheekh, and D. Hanelt. 2014. Pilot cultivation of the chlorophyte microalga *Scenedesmus obliquus* as a promising feedstock for biofuel. *Biomass and Bioenergy* 64:237–244. doi:10.1016/j.biombioe.2014.03.049.
- Abreu, A. P., B. Fernandes, A. A. Vicente, J. Teixeira, and G. Dragone. 2012. Mixotrophic cultivation of *Chlorella vulgaris* using industrial dairy waste as organic carbon source. *Bioresource Technology* 118:61–66. doi:10.1016/j.biortech.2012.05.055.
- Alexandre, M., K. Gundermann, A. Pascal, R. Van Grondelle, C. Büchel, and B. Robert. 2014. Probing the carotenoid content of intact *Cyclotella* cells by resonance raman spectroscopy. *Photosynthesis Research* 119:273–281. doi:10.1007/s11120-013-9942-y.
- Aronow, M. E., and E. Y. Chew. 2014. Age-related eye disease study 2: perspectives, recommendations, and unanswered questions. *Current Opinion in Ophthalmology* 25:186–190. doi:10.1097/ICU.0000000000000046.
- Asai, A., T. Sugawara, H. Ono, and A. Nagao. 2004. Biotransformation of fucoxanthinol into amarouci-xanthin in mice and HepG2 cells: Formation and cytotoxicity of fucoxanthin metabolites. *Drug Metabolism and Disposition* 32:205–211. doi:10.1124/dmd.32.2.205.
- Bae, J. H., and S. B. Hur. 2011. Selection of suitable species of *Chlorella*, *Nannochloris*, and *Nannochloropsis* in high and low temperature seasons for mass culture of the rotifer *Bachionus plicatilis*. *Fisheries and Aquatic Sciences* 14:323–332. doi:10.5657/FAS.2011.0323.
- Bajhaiya, A. K., M. J. Ziehe, and J. K. Pittman. 2017. Transcriptional engineering of microalgae: Prospects for high value chemicals. *Trends in Biotechnology* 35:95–99. doi:10.1016/j.tibtech.2016.06.001.
- Banerjee, A., R. Sharma, Y. Chisti, and U. C. Banerjee. 2002. *Botryococcus braunii*: a renewable source of hydrocarbons and other chemicals. *Critical Reviews in Biotechnology* 22:245–279. doi:10.1080/07388550290789513.
- Bar, E., M. Rise, M. Vishkautsan, and S. Arad. 1995. Pigment and structural changes in *Chlorella zofingiensis* upon light and nitrogen stress. *Journal of Plant Physiology* 146:527–534. doi:10.1016/S0176-1617(11)82019-5.
- Bayona, K. C. D., and L. A. Graces. 2014. Effect of different media on exopolysaccharide and biomass production by the green microalga *Botryococcus braunii*. *Journal of Applied Phycology* 26:2087–2095. doi:10.1007/s10811-014-0242-5.
- Beer, A., M. Juhas, and C. Buchel. 2011. Influence of different light intensities and different iron nutrition on the photosynthetic apparatus in the diatom *Cyclotella meneghiniana* (Bacillariophyceae). *Journal of Phycology* 47:1266–1273. doi:10.1111/j.1529-8817.2011.01060.x.
- Ben-Amotz, A., and Y. Levy. 1996. Bioavailability of a natural isomer mixture compared with synthetic all-trans β -carotene in human serum. *American Journal of Clinical Nutrition* 63:729–734. doi:10.1093/ajcn/63.5.729.
- Bhagavathy, S., and P. Sumathi. 2012. Stabilization of membrane bound ATPases and lipid peroxidation by carotenoids from *Chlorococcum humicola* benzo(a)pyrene induced toxicity. *Asian Pacific Journal of Tropical Biomedicine* 2:380–384. doi:10.1016/S2221-1691(12)60060-4.
- Bhagavathy, S., P. Sumathi and B. I. Jancy Sherene. 2011a. Green algae *Chlorococcum humicola*-a new source of bioactive compounds with anti-microbial activity. *Asian Pacific Journal of Tropical Biomedicine* 1: S1–S7. doi:10.1016/S2221-1691(11)60111-1.
- Bhagavathy, S., P. Sumathi, and M. Madhushree. 2011b. Antimutagenic assay of carotenoids from green algae *Chlorococcum humicola* using *Salmonella typhimurium* TA98, TA100 and TA102. *Asian Pacific Journal of Tropical Biomedicine* 1:308–316. doi:10.1016/S2222-1808(11)60073-X.
- Bilbao, P. G. S., C. Damiani, G. A. Salvador, and P. Leonardi. 2016. *Haematococcus pluvialis* as a source of fatty acids and phytosterols: Potential nutritional and biological implications. *Journal of Applied Phycology* 28:3283–3294. doi:10.1007/s10811-016-0899-z.
- Bishop, W. M., and H. M. Zubeck. 2012. Evaluation of microalgae for use as nutraceuticals and nutritional supplements. *Journal of Nutrition and Food Sciences* 2:147. doi:10.4172/2155-9600.1000147.
- Borowitzka, L. J., and M. A. Borowitzka. 1989. β -Carotene (pro-vitamin A) production with algae. In *Biotechnology of Vitamins, Pigments and Growth Factors*, Ed. E. J. Vandamme, pp. 15–26. London: Elsevier Applied Science.
- Borowitzka, M. A. 2010. Carotenoid production using microorganisms. In *Single Cell Oils: Microbial and Algal oils*, Eds. Z. Cohen and C. Ratledge, pp. 225–240. Urbana: AOCS Press.
- Bozarth, A., U. G. Maier, and S. Zauner. 2009. Diatoms in biotechnology: Modern tools and applications. *Applied Microbiology and Biotechnology* 82:195–201. doi:10.1007/s00253-008-1804-8.
- Brotas, V., and M. R. Plante-Cuny. 2003. The use of HPLC pigment analysis to study microphytobenthos communities. *Acta Oecologica* 24:109–115. doi:10.1016/S1146-609X(03)00013-4.
- Brown, J. S. 1987. Functional organization of chlorophyll a and carotenoids in alga, *Nannochloropsis salina*. *Plant Physiology* 83:434–437. doi:10.1104/pp.83.2.434.
- Brunet, C., R. Chandrasekaran, L. Barra, V. Giovagnetti, F. Corato, and A. V. Ruban. 2014. Spectral radiation dependent photo-protective mechanism in the diatom *Pseudo-nitzschia multistriata*. *Plos One* 9:1–10. doi:10.1371/journal.pone.0087015.
- Buono, S., A. L. Langellotti, F. Rinna, A. Martello, and V. Fagliano. 2014. Functional ingredients from microalgae. *Food and Function* 5:1669–1685. doi:10.1039/C4FO00125G.
- Burczyk, J., H. Szkawran, I. Zontek, and F. C. Czygan. 1981. Carotenoids in the outer cell-wall layer of *Scenedesmus* (Chlorophyceae). *Planta* 151:247–250. doi:10.1007/BF00395176.
- Burton-Freeman, B., J. Talbot, E. Park, S. Krishnankutty, and I. Edirisinghe. 2012. Protective activity of processed tomato products on

- postprandial oxidation and inflammation: a clinical trial in healthy weight men and women. *Molecular Nutrition and Food Research* 56:622–631. doi:10.1002/mnfr.201100649.
- Campenni, L., B. P. Nobre, C. A. Santos, A. C. Oliveira, M. R. Aires-Barros, A. M. F. Palavra, and L. Gouveia. 2013. Carotenoid and lipid production by the autotrophic microalga *Chlorella protothecoides* under nutritional, salinity, and luminosity stress conditions. *Applied Microbiology and Biotechnology* 97:1383–1393. doi:10.1007/s00253-012-4570-6.
- Carbonell-Capella, J. M., M. Buniowska, F. J. Barba, M. J. Esteve, and A. Frigola. 2014. Analytical methods for determining bio-availability and bio-accessibility of bioactive compounds from fruits and vegetables: A review. *Comprehensive Reviews in Food Science and Food Safety* 13:155–171. doi:10.1111/1541-4337.12049.
- Careri, M., L. Furlattini, A. Mangia, M. Musci, E. Anklam, A. Theobald, and C. Von Holst. 2001. Supercritical fluid extraction for liquid chromatographic determination of carotenoids in *Spirulina pacifica* algae: A chemometric approach. *Journal of Chromatography A* 12:61–71. doi:10.1016/S0021-9673(01)00545-3.
- Castenmiller, J. J. M., and C. E. West. 1998. Bioavailability and bioconversion of carotenoids. *Annual Review of Nutrition* 18:19–38. doi:10.1146/annurev.nutr.18.1.19.
- Ceron, M. C., I. Campos, J. F. Sánchez, F. G. Acien, E. Molina, and J. M. Fernández-Sevill. 2008. Recovery of lutein from microalgae biomass: Development of a process for *Scenedesmus almeriensis* biomass. *Journal of Agricultural and Food Chemistry* 56:11761–11766. doi:10.1021/jf8025875.
- Ceron, M. C., M. C. Garcia-Malea, J. Rivas, F. G. Acien, J. M. Fernandez, E. del Rio, M. G. Guerrero, and E. Molina. 2007. Antioxidant activity of *Haematococcus pluvialis* cells grown in continuous culture as a function of their carotenoid and fatty acid content. *Applied Microbiology and Biotechnology* 74:1112–1119. doi:10.1007/s00253-006-0743-5.
- Cesarini, J. P., L. Michel, J. M. Maurette, H. Adhoue, and M. Bejot. 2003. Immediate effects of UV radiation on the skin: Modification by an antioxidant complex containing carotenoids. *Photodermatology, Photoimmunology and Photomedicine* 19:182–189. doi:10.1034/j.1600-0781.2003.00044.x.
- Cha, K. H., S. Y. Koo, and D. U. Lee. 2008. Anti-proliferative effects of carotenoids extracted from *Chlorella ellipsoidea* and *Chlorella vulgaris* on human colon cancer cells. *Journal of Agricultural and Food Chemistry* 56:10521–10526. doi:10.1021/jf802111x.
- Cha, K. H., H. J. Lee, S. Y. Koo, D. G. Song, D. U. Lee, and C. H. Pan. 2010. Optimization of pressurized liquid extraction of carotenoids and chlorophylls from *Chlorella vulgaris*. *Journal of Agricultural and Food Chemistry* 58:793–797. doi:10.1021/jf902628j.
- Chacon-Lee, T. L., and E. Gonzalez-Marino G. 2010. Microalgae for healthy foods-possibilities and challenges. *Comprehensive Reviews in Food Science and Food Safety* 9:655–675. doi:10.1111/j.1541-4337.2010.00132.x.
- Chan, K. C., P. J. Pen, and M. C. Yin. 2012. Anti-coagulatory and anti-inflammatory effects of astaxanthin in diabetic rats. *Journal of Food Science* 77:H76–H80. doi:10.1111/j.1750-3841.2011.02558.x.
- Chan, M. M. C., S. S. H. Ho, D. D. J. Lee, C. C. Y. Chen, C. C. Huang, and J. S. Chang. 2013. Characterization, extraction and purification of lutein produced by an Indigenous microalga *Scenedesmus obliquus* CNW-N. *Biochemical Engineering Journal* 78:24–31. doi:10.1016/j.bej.2012.11.017.
- Chen, C. Y., K. L. Yeh, R. Aisyah, D. J. Lee, and J. S. Chang. 2011. Cultivation, photobioreactor design and harvesting of microalgae for biodiesel production: A critical review. *Bioresource Technology* 102:71–81. doi:10.1016/j.biortech.2010.06.159.
- Chew, W., B. D. Mathison, L. L. Kimble, P. F. Mixer, and B. P. Chew. 2013. Astaxanthin decreases inflammatory biomarkers associated with cardiovascular disease in human umbilical vein endothelial cells. *American Journal of Advanced Food Science and Technology* 1:1–17. doi:10.7726/ajafst.2013.1001.
- Chidambara Murthy, K. N., J. Rajesha, A. Vanitha, M. M. Swamy, and G. A. Ravishankar. 2005. Protective effect of *Dunaliella salina*-A marine microalga, against carbon tetrachloride-induced hepatotoxicity in rats. *Hepatology Research* 33:313–319. doi:10.1016/j.hepres.2005.08.008.
- Choi, H. D., H. E. Kang, S. H. Yang, M. G. Lee, and W. G. Shin. 2011. Pharmacokinetics and first-pass metabolism of astaxanthin in rats. *British Journal of Nutrition* 105:220–227. doi:10.1017/S0007114510003454.
- Christaki, E., E. Bonos, I. Giannenas, and P. Florou-Paneri. 2012. Functional properties of carotenoids originating from algae. *Journal of the Science of Food and Agriculture* 93:5–11. doi:10.1002/jsfa.5902.
- Chu, W. L. 2012. Biotechnological applications of microalgae. *Te/SME* 6: S24–S37.
- Coca, M., V. M. Barrocal, S. Lucas, G. González-Benito, and M. T. Garcia-Cubero. 2014. Protein production in *Spirulina platensis* biomass using beet vinasse supplemented culture media. *Food and Bioproducts Processing* 94:306–312. doi:10.1016/j.fbp.2014.03.012.
- Coesel, S., M. Obornik, J. Varela, A. Falcioratore, and C. Bowler. 2008. Evolutionary origins and functions of the carotenoid biosynthetic pathway in marine diatoms. *Plos One* 3:e2896. doi:10.1371/journal.pone.0002896.
- Cooperstone, J. L., and S. J. Schwartz. 2016. Recent insights into health benefits of carotenoids. In *Handbook on Natural Pigments in Food and Beverages: Industrial Applications for Improving Food Color*, Eds. R. Carle and R. M. Schweiggert, pp. 473–497, Duxford, UK: Woodhead Publishing, Elsevier.
- Cooperstone, J. L., R. A. Ralston, K. M. Riedl, T. C. Haufe, R. M. Schweiggert, S. A. King, C. D. Timmers, D. M. Francis, G. B. Lesinski, S. K. Clinton, and S. J. Schwartz. 2015. Enhanced bioavailability of lycopene in humans when consumed as *cis*-isomers from tangerine tomatoes compared to red tomato juices, a randomized, cross-over clinical trial. *Molecular Nutrition and Food Research* 59:658–669. doi:10.1002/mnfr.201400658.
- Costa, B. S., A. Jungandreas, T. Jakob, W. Weisheit, M. Mittag, and C. Wilhelm. 2013. Blue light is essential for high light acclimation and methylation and chromatin patterning photoprotection in the diatom *Phaeodactylum tricornutum*. *Journal of Experimental Botany* 64:483–493. doi:10.1093/jxb/ers340.
- Curtain, C. 2000. Plant biotechnology the growth of Australia's algal β -carotene industry. *Australasian Biotechnology* 10:19–23.
- D'Alessandro, E. B., and N. R. Antoniosi Filho. 2016. Concepts and studies on lipid and pigments of microalgae: A review. *Renewable and Sustainable Energy Reviews* 58:832–841. doi:10.1016/j.rser.2015.12.162.
- Dambek, M., U. Eilers, J. Breitenbach, S. Steiger, C. Büchel, and G. Sandmann. 2012. Biosynthesis of fucoxanthin and diadinoxanthin and function of initial pathway genes in *Phaeodactylum tricornutum*. *Journal of Experimental Botany* 63:5607–5612. doi:10.1093/jxb/ers211.
- Deenu, A., S. Naruenartwongsakul, and S. M. Kim. 2013. Optimization and economic evaluation of ultrasound extraction of lutein from *Chlorella vulgaris*. *Biotechnology and Bioprocess Engineering* 18:1151–1162. doi:10.1007/s12257-013-0213-8.
- Degen, J., A. Uebele, A. Retze, U. Schmid-Staiger, and W. Trosch. 2001. A novel airlift photobioreactor with baffles for improved light utilization through the flashing light effect. *Journal of Biotechnology* 92:89–94. doi:10.1016/S0168-1656(01)00350-9.
- Del Campo, J. A., M. García-González, and M. G. Guerrero. 2007. Outdoor cultivation of microalgae for carotenoid production: Current state and perspectives. *Applied Microbiology and Biotechnology* 74:1163–1174. doi:10.1007/s00253-007-0844-9.
- Del Campo, J. A., J. Moreno, H. Rodriguez, M. A. Vargas, J. Rivas, and M. G. Guerrero. 2000. Carotenoid content of chlorophycean microalgae: Factors determining lutein accumulation in *Muriellopsis* sp. (*Chlorophyta*). *Journal of Biotechnology* 76:51–59. doi:10.1016/S0168-1656(99)00178-9.
- Del Campo, J. A., J. Moreno, H. Rodriguez, M. A. Vargas, J. Rivas, and M. G. Guerrero. 2001. Lutein production by *Muriellopsis* sp. in an outdoor tubular photobioreactor. *Journal of Biotechnology* 85:289–295. doi:10.1016/S0168-1656(00)00380-1.
- Del Campo, J. A., H. Rodriguez, J. Moreno, M. Á. Vargas, J. Rivas, and M. G. Guerrero. 2004. Accumulation of astaxanthin and lutein in *Chlorella zofingiensis* (*Chlorophyta*). *Applied Microbiology and Biotechnology* 64:848–854. doi:10.1007/s00253-003-1510-5.
- Domingues, N., A. R. Matos, J. M. da Silva, and P. Cartaxana. 2012. Response of the diatom *Phaeodactylum tricornutum* to photo oxidative stress resulting from high light exposure. *Plos One* 7:1–6. doi:10.1371/journal.pone.0038162.
- Dominguez-Bocanegra, A. R., L. I. Guerrero, F. M. Jeronimo, and A. T. Campocoso. 2004. Influence of environmental and nutritional factors in the production of astaxanthin from *Haematococcus pluvialis*. *Bioresource Technology* 92:209–214 doi:10.1016/j.biortech.2003.04.001.

- Dong, L. Y., J. Jin, G. Lu, and X. L. Kang. 2013. Astaxanthin attenuates the apoptosis of retinal ganglion cells in db/db mice by inhibition of oxidative stress. *Marine Drugs* 11:960–974. doi:10.3390/md11030960.
- Donhowe, E. G., and F. Kong. 2014. β -carotene: Digestion, microencapsulation, and *in vitro* bioavailability. *Food and Bioprocess Technology* 7:338–354. doi:10.1007/s11947-013-1244-z.
- Dufosse, L., P. Galaup, A. Yaron, S. M. Arad, P. Blanc, K. N. Chidambaram Murthy, and G. A. Ravishankar. 2005. Microorganisms and microalgae as sources of pigments for food use: a scientific oddity or an industrial reality? *Trends in Food Science and Technology* 16:389–406. doi:10.1016/j.tifs.2005.02.006.
- El-Baky, H. H. A., F. K. El-Baz, and G. S. El-Baroty. 2003. *Spirulina* species as a source of carotenoids and α -tocopherol and its anti-carcinoma factors. *Biotechnol* 2:222–240. doi:10.3923/biotech.2003.222.240.
- El-Baz, F. K., A. M. Abdoul-Enein, G. S. El-Baroty, A. M. Youssef, and H. H. A. El-Baky. 2002. Accumulation of antioxidant vitamins in *Dunaliella salina*. *Journal of Biological Sciences* 2:220–223. doi:10.3923/jbs.2002.220.223.
- Eriksen, N. T. 2008. Production of phycocyanin a pigment with applications in biology, biotechnology, foods and medicine. *Applied Microbiology and Biotechnology* 80:1–14. doi:10.1007/s00253-008-1542-y.
- Faheed, F., and Z. Abd el Fattah. 2008. Effect of *Chlorella vulgaris* as bio-fertilizer on growth parameters and metabolic aspects of lettuce plant. *Journal of Agriculture and Social Sciences* 4:165–169.
- Feng, S., X. Li, Z. Xu, and J. Qi. 2014. *Dunaliella salina* as a novel host for the production of recombinant proteins. *Applied Microbiology and Biotechnology* 98:4293–4300. doi:10.1007/s00253-014-5636-4.
- Fernandez-Sevilla, J. M., F. G. Acien-Fernandez, and E. Molina-Grima. 2010. Biotechnological production of lutein and its applications. *Applied Microbiology and Biotechnology* 86:27–40. doi:10.1007/s00253-009-2420-y.
- Forjan, L. E., N. I. Garbayo, B. C. Casal, and L. C. Vilchez. 2007. Enhancement of carotenoid production in *Nannochloropsis* by phosphate and sulphur limitation. In: *Communicating Current Research and Educational Topics and Trends in Applied Microbiology*, Ed. A. Mendez-Vilas, pp. 356–364, Espana: Microbiology Book Series, Formatex, Badajoz, Spain. ISBN-13: 978-84-611-9421-6(Series), ISBN-13: 978-84-611-9422-3(vol.1).
- Fu, W., O. Guomundsson, G. Paglia, G. Herjolfsson, O. S. Andresson, B. Q. Palsson, and S. Brynjolfsson. 2013. Enhancement of carotenoid biosynthesis in the green microalga *Dunaliella salina* with light-emitting diodes and adaptive laboratory evolution. *Applied Microbiology and Biotechnology* 97:2395–2403. doi:10.1007/s00253-012-4502-5.
- Fu, W., G. Paglia, M. Magnusdottir, E. A. Steinarsdottir, S. Gudmundsson, B. Q. Palsson, O. S. Andresson, and S. Brynjolfsson. 2014. Effect of abiotic stressors on lutein production in the green microalga *Dunaliella salina*. *Microbial Cell Factories* 13:3. doi:10.1186/1475-2859-13-3.
- Fu, W., K. Wichuk, and S. Brynjolfsson. 2015. Developing diatoms for value-added products: Challenges and opportunities. *New Biotechnology* 32:547–551. doi:10.1016/j.nbt.2015.03.016.
- Fujii, Y., A. Ben-Amotz, S. Sakamoto, and H. Nagasawa. 1993. Effects of β -carotene rich algae *Dunaliella bardawil* on the dynamic changes of normal and neoplastic mammary cells and general metabolism in mice. *Anticancer Research* 13:389–393.
- Gantar, M., and Z. Svircev. 2008. Microalgae and cyanobacteria: Food for thought. *Journal of Phycology* 44:260–268. doi:10.1111/j.1529-8817.2008.00469.x.
- Garcia-Gonzalez, M., J. Moreno, J. C. Manzano, F. J. Florencio, and M. G. Guerrero. 2005. Production of *Dunaliella salina* biomass rich in 9-cis- β -carotene and lutein in a closed tubular photobioreactor. *Journal of Biotechnology* 115:81–90. doi:10.1016/j.jbiotec.2004.07.010.
- Garcia-Malea, M. C., F. G. Acien, E. del Río, J. M. Fernandez, M. C. Ceron, M. G. Guerrero, and E. Molina-Grima. 2009. Production of astaxanthin by *Haematococcus pluvialis*: taking the one-step system outdoors. *Biotechnology and Bioengineering* 102:651–657. doi:10.1002/bit.22076.
- Gastineau, R., N. A. Davidovich, J. F. Bardeau, A. Caruso, V. Leignel, Y. Hardivillier, Y. Rince, B. Jacqueline, O. I. Davidovich, P. Gaudin, E. J. Cox, and J. L. Mouget. 2012. *Haslea karadagensis* (Bacillariophyta): A second blue diatom, recorded from the black sea and producing a novel blue pigment. *European Journal of Phycology* 47:469–479. doi:10.1080/09670262.2012.741713.
- Gastineau, R., F. Turcotte, J. B. Pouvreau, M. Moranc ais, J. Fleurence, E. Windarto, S. Arsad, F. S. Prasetya, P. Jaouen, M. Babin, et al. 2014. Marine, promising blue pigments from a widespread *Haslea* diatom species complex. *Marine Drugs* 12:3161–3189. doi:10.3390/md12063161.
- Gaziano, J. M., J. E. Manson, L. G. Branch, G. A. Colditz, W. C. Willett, and J. E. Buring. 1995. A prospective study of consumption of carotenoids in fruits and vegetables and decreased cardiovascular mortality in the elderly. *Annals of Epidemiology* 5:255–260. doi:10.1016/1047-2797(94)00090-G.
- Gelzinis, A., V. Butkus, E. Songaila, R. Augulis, A. Gall, C. B ichel, B. Robert, D. Abramavicius, D. Zigmantas, and L. Valkunas. 2015. Mapping energy transfer channels in fucoxanthin-chlorophyll protein complex. *BBA Bioenergetics* 1847:241–247. doi:10.1016/j.bbabio.2014.11.004.
- Gil-Chavez, G. J., J. A. Villa, J. Fernando Ayala-Zavala, J. Basilio Heredia, D. Sepulveda, E. M. Yahia, and G. A. Gonzalez-Aguilar. 2013. Technologies for extraction and production of bioactive compounds to be used as nutraceuticals and food ingredients: An overview. *Comprehensive Reviews in Food Science and Food Safety* 12:5–23. doi:10.1111/1541-4337.12005.
- Giuliano, G., R. Tavazza, G. Diretto, P. Beyer, and M. A. Taylor. 2008. Metabolic engineering of carotenoid biosynthesis in plants. *Trends in Biotechnology* 26:139–145. doi:10.1016/j.tibtech.2007.12.003.
- Gog, A., L. Senila, M. Roman, E. Luca, C. Roman, and F. D. Irimie. 2012. Oil extraction and fatty acid characterization of *Nannochloropsis oculata* microalgae for biodiesel applications. *Studia UBB Chemia* 57:111–118.
- Gong, M., and A. Bassi. 2016. Carotenoids from microalgae: A review of recent developments. *Biotechnology Advances* 34:1396–1412. doi:10.1016/j.biotechadv.2016.10.005.
- Gonzalez, L. E., and Y. Bashan. 2000. Increased growth of the microalga *Chlorella vulgaris* when co-immobilized and co-cultured in alginate beads with the plant- growth-promoting bacterium *Azospirillum brasilense*. *Applied and Environmental Microbiology* 66:1527–1531. doi:10.1128/AEM.66.4.1527-1531.2000.
- Gorin, K. V., Y. E. Sergeeva, V. V. Butylin, A. V. Komova, V. M. Pojidaev, G. U. Badranova, A. A. Shapovalova, I. A. Konova, and P. M. Gotovtsev. 2015. Methods coagulation/flocculation and flocculation with ballast agent for effective harvesting of microalgae. *Bioresource Technology* 193:178–184. doi:10.1016/j.biortech.2015.06.097.
- Gouveia, L., A. P. Batista, I. Sousa, A. Ray-mundo, and N. M. Bandarra. 2008. Micro-algae in novel food products. In *Food Chemistry Research Developments*, Ed. K. Papa-doupoulos, pp. 75–112. New York: Nova Science Publishers.
- Gouveia, L., G. Choubert, N. Pereira, J. Santinha, J. Empis, and E. Gomes. 2002. Pigmentation of gilthead seabream, *Sparus aurata* (L. 1875), using *Chlorella vulgaris* (Chlorophyta, Volvocales) microalga. *Journal of Aquaculture Research and Development* 33:987–993. doi:10.1046/j.1365-2109.2002.00751.x.
- Gouveia, L., A. Raymundo, A. P. Batista, I. Sousa, and J. Empis. 2006. *Chlorella vulgaris* and *Haematococcus pluvialis* biomass as colouring and antioxidant in food emulsions. *European Food Research and Technology* 222:362–367. doi:10.1007/s00217-005-0105-z.
- Gouveia, L., V. Veloso, A. Reis, H. Fernandes, J. Novais, and J. Empis. 1995. Evolution of pigment composition in *Chlorella vulgaris*. *Biore-source Technology* 57:157–163. doi:10.1016/0960-8524(96)00058-2.
- Gouveia, L., V. Veloso, A. Reis, H. Fernandes, J. Novais, and J. Empis. 1996. *Chlorella vulgaris* used to colour egg yolk. *Journal of the Science of Food and Agriculture* 70:167–172. doi:10.1002/(SICI)1097-0010(199602)70:2%3c167::AID-JSFA72%3e3.0.CO;2-2.
- Grima, E. M., E. Belarbi, F. G. A. Fernandez, A. R. Medina, and Y. Chisti. 2003. Recovery of microalgal biomass and metabolites: Process options and economics. *Biotechnology Advances* 20:491–515. doi:10.1016/S0734-9750(02)00050-2.
- Grobelaar, J. U. 2003. Quality control and assurance: crucial for the sustainability of the applied phycology industry. *Journal of Applied Phycology* 15:209–215. doi:10.1023/A:1023820711706.
- Grung, M., P. Metzger, and S. Liaen Jensen. 1989. Primary and secondary carotenoids in two races of green *Botryococcus braunii*. *Biochemical Systematics and Ecology* 17:263–269. doi:10.1016/0305-1978(89)90001-X.

- Grung, M., P. Metzger, and S. Liaaen Jensen. 1994. Algal carotenoids 53; secondary carotenoids of algae 4; secondary carotenoids in the green alga *Botryococcus braunii*, race L, new strain. *Biochemical Systematics and Ecology* 22:25–29. doi:10.1016/0305-1978(94)90111-2.
- Guedes, A. C., H. M. Amaro, and F. X. Malcata. 2011b. Microalgae as sources of carotenoids. *Marine Drugs* 9:625–644. doi:10.3390/md9040625.
- Guedes, A. C., M. S. Gíao, A. A. Matias, A. V. M. Nunes, M. E. Pintado, C. M. M. Duarte, and F. X. Malcata. 2013. Supercritical fluid extraction of carotenoids and chlorophylls a, b and c, from a wild strain of *Scenedesmus obliquus* for use in food processing. *Journal of Food Engineering* 116:478–482. doi:10.1016/j.jfoodeng.2012.12.015.
- Guedes, A., H. M. Amaro, and F. X. Malcata. 2011a. Microalgae as sources of high added value compounds—A brief review of recent work. *Biotechnology Progress* 27:597–613. doi:10.1002/btpr.575.
- Guerin, M., M. E. Huntley, and M. Olaizola. 2003. *Haematococcus* astaxanthin: Applications for human health and nutrition. *Trends in Biotechnology* 21:210–216. doi:10.1016/S0167-7799(03)00078-7.
- Guo, B., B. Liu, B. Yang, P. Sun, X. Lu, J. Liu, and F. Chen. 2016. Screening of diatom strain and characterization of *Cyclotella cryptica* as a potential fucoxanthin producer. *Marine Drugs* 14:125. doi:10.3390/md14070125.
- Halim, R., M. K. Danquah, and P. A. Webley. 2012a. Extraction of oil from microalgae for biodiesel production: A review. *Biotechnology Advances* 30:709–732. doi:10.1016/j.biotechadv.2012.01.001.
- Halim, R., R. Harun, M. K. Danquah and P. A. Webley. 2012b. Microalgal cell disruption for biofuel development. *Applied Energy* 91:116–121. doi:10.1016/j.apenergy.2011.08.048.
- Hamed, I. 2016. The evolution and versatility of microalgal biotechnology: A review. *Comprehensive Reviews in Food Science and Food Safety* 15:1104–1123. doi:10.1111/1541-4337.12227.
- Hanaa, H. A. E. B., K. D. B. Farouk, and G. S. E. Baroty. 2003. *Spirulina* species as a source of carotenoid and tocopherol and its anti-carcinoma factors. *Biotechnology* 2:222–240. doi:10.3923/biotech.2003.222.240.
- Hata, N., J. C. Ogbonna, Y. Hasegawa, H. Taroda, and H. Tanaka. 2001. Production of astaxanthin by *Haematococcus pluvialis* in a sequential heterotrophic-photoautotrophic culture. *Journal of Applied Phycology* 13:395–402. doi:10.1023/A:1011921329568.
- Havlik, I., P. Lindner, T. Scheper, and K. F. Reardon. 2013. On-line monitoring of large cultivations of microalgae and cyanobacteria. *Trends in Biotechnology* 31:406–414. doi:10.1016/j.tibtech.2013.04.005.
- Henriquez, V., C. Escobar, J. Galarza, and J. Gimpel. 2016. Carotenoids in microalgae. In *Carotenoids in Nature*, Ed. S. Claudia, pp. 219–237, Switzerland: Springer International Publishing.
- Heo, S. J., S. C. Ko, S. M. Kang, H. S. Kang, J. P. Kim, S. H. Kim, K. W. Lee, M. G. Cho, and Y. J. Jeon. 2008. Cytoprotective effect of fucoxanthin isolated from brown algae *Sargassum siliquastrum* against H₂O₂-induced cell damage. *European Food Research and Technology* 228:145–151. doi:10.1007/s00217-008-0918-7.
- Heo, S. J., W. J. Yoon, K. N. Kim, G. N. Ahn, S. M. Kang, D. H. Kang, C. Oh, W. K. Jung, and Y. J. Jeon. 2010. Evaluation of anti-inflammatory effect of fucoxanthin isolated from brown algae in lipopolysaccharide-stimulated RAW 264.7 macrophages. *Food and Chemical Toxicology* 48:2045–2051. doi:10.1016/j.fct.2010.05.003.
- Higuera-Ciupara, I., L. Felix-Valenzuela, and F. M. Goycoolea. 2006. Astaxanthin: A review of its chemistry and applications. *Critical Reviews in Food Science and Nutrition* 46:185–196. doi:10.1080/10408690590957188.
- Ho, S. H., M. C. Chan, C. C. Liu, C. Y. Chen, W. L. Lee, D. J. Lee, and J. S. Chang. 2014. Enhancing lutein productivity of an indigenous microalga *Scenedesmus obliquus* FSP-3 using light-related strategies. *Bioresource Technology* 152:275–282. doi:10.1016/j.biortech.2013.11.031.
- Holman, B. W. B., and A. E. O. Malau-Aduli. 2013. *Spirulina* as a livestock supplement and animal feed. *Journal of Animal Physiology and Animal Nutrition* 97:615–623. doi:10.1111/j.1439-0396.2012.01328.x.
- Hoseini, S. M., K. Khosravi-Darani, and M. R. Mozafari. 2013. Nutritional and medical applications of *Spirulina* microalgae. *Mini Reviews in Medicinal Chemistry* 13:1231–1237. doi:10.2174/1389557511313080009.
- Hosokawa, M., S. Wanezaki, K. Miyauchi, H. Kurihara, H. Kohno, J. Kawabata, S. Odashima, and K. Takahashi. 1999. Apoptosis-inducing effect of fucoxanthin on human leukemia cell line HL-60. *Food Science and Technology Research* 5:243–246. doi:10.3136/fstr.5.243.
- Hosseini, T. A., and M. Shariati. 2009. *Dunaliella* biotechnology: Methods and applications. *Journal of Applied Microbiology* 107:14–35. doi:10.1111/j.1365-2672.2009.04153.x.
- Hsu, Y. W., C. F. Tsai, W. H. Chang, Y. C. Ho, W. K. Chen, and F. J. Lu. 2008. Protective effects of *Dunaliella salina*-a carotenoid rich alga, against carbon tetrachloride induced hepatotoxicity in mice. *Food and Chemical Toxicology* 46:3311–3317. doi:10.1016/j.fct.2008.07.027.
- Hu, H., and K. Gao. 2006. Response of growth and fatty acid compositions of *Nannochloropsis* sp. to environmental factors under elevated CO₂ concentration. *Biotechnology Letters* 28:987–992. doi:10.1007/s10529-006-9026-6.
- Hu, Y. R., F. Wang, S. K. Wang, C. Z. Liu, and C. Guo. 2013. Efficient harvesting of marine microalgae *Nannochloropsis maritima* using magnetic nanoparticles. *Bioresource Technology* 138:387–390. doi:10.1016/j.biortech.2013.04.016.
- Huangfu, J., J. Liu, Z. Sun, M. Wang, Y. Jiang, Z. Y. Chen, and F. Chen. 2013. Anti-ageing effects of astaxanthin-rich alga *Haematococcus pluvialis* on fruit flies under oxidative stress. *Journal of Agricultural and Food Chemistry* 61:7800–7804. doi:10.1021/jf402224w.
- Hussein, G., U. Sankawa, H. Goto, K. Matsumoto, and H. Watanabe. 2006. Astaxanthin, a carotenoid with potential in human health and nutrition. *Journal of Natural Products* 69:443–449. doi:10.1021/np050354+.
- Ip, P. F., and F. Chen. 2005. Production of astaxanthin by the green microalga *Chlorella zofingensis* in the dark. *Process Biochemistry (Barking, UK)* 40:733–738. doi:10.1016/j.procbio.2004.01.039.
- Ishida, B. K., and M. H. Chapman. 2009. Carotenoid extraction from plants using a novel, environmentally friendly solvent. *Journal of Agricultural and Food Chemistry* 57:1051–1059.
- Ishiki, M., Y. Nishida, H. Ishibashi, T. Wada, S. Fujisaka, A. Takikawa, M. Urakaze, T. Sasaoka, I. Usui, and K. Tobe. 2013. Impact of divergent effects of astaxanthin on insulin signaling in l6 cells. *Endocrinol* 154:2600–2612. doi:10.1021/jf8026292. doi:10.1210/en.2012-2198.
- Jaime, L., J. A. Mendiola, M. Herrero, C. Soler-Rivas, S. Santoyo, F. J. Señorans, A. Cifuentes, and E. Ibanez. 2005. Separation and characterization of antioxidants from *Spirulina platensis* microalga combining pressurized liquid extraction, TLC, and HPLC-DAD. *Journal of Separation Science* 28:2111–2119. doi:10.1002/jssc.200500185.
- Jayappriyan, K. R., R. Rajkumar, V. Venkatakrishnan, S. Nagaraj, and R. Rengasamy. 2013. *In vitro* anticancer activity of natural-carotene from *Dunaliella salina* EU5891199 in PC-3 cells. *Biomedicine and Preventive Nutrition* 3:99–105. doi:10.1016/j.bionut.2012.08.003.
- Jeffrey, S., and M. Vesik. 1997. Introduction to marine phytoplankton and their pigment signatures. In *Phytoplankton Pigments in Oceanography: Guidelines to Modern Methods*, Eds. S. Jeffrey, R. Mantoura, and S. Wright, pp. 37–84, Paris, France: UNESCO Publishing.
- Jungandreas, A., B. S. Costa, T. Jakob, M. von Bergen, S. Baumann, and C. Wilhelm. 2014. The acclimation of *Phaeodactylum tricornutum* to blue and red light does not influence the photosynthetic light reaction but strongly disturbs the carbon allocation pattern. *Plos One* 9:1–14. doi:10.1371/journal.pone.0099727.
- Kadam, S. U., B. K. Tiwari, and C. P. O'Donnell. 2013. Application of novel extraction technologies for bioactives from marine algae. *Journal of Agricultural and Food Chemistry* 61:4667–4675. doi:10.1021/jf400819p.
- Kang, C. D., and S. J. Sim. 2008. Direct extraction of astaxanthin from *Haematococcus* culture using vegetable oils. *Biotechnology Letters* 30:441–444. doi:10.1007/s10529-007-9578-0.
- Kang, C. D., and S. J. Sim. 2007. Selective extraction of free astaxanthin from *Haematococcus* culture using a tandem organic solvent system. *Biotechnology Progress* 23:866–871. doi:10.1021/bp0700354.
- Kang, C. D., J. S. Lee, T. H. Park, and S. J. Sim. 2005. Comparison of heterotrophic and photoautotrophic induction on astaxanthin production by *Haematococcus pluvialis*. *Applied Microbiology and Biotechnology* 68:237–241. doi:10.1007/s00253-005-1889-2.
- Kar, H. K. 2002. Efficacy of β -carotene topical application in melisma: An open clinical trial. *Indian Journal of Dermatology, Venereology and Leprosology* 68:320–322.

- Kent, M., H. M. Welladsen, A. Mangott, and Y. Li. 2015. Nutritional evaluation of Australian microalgae as potential human health supplements. *Plos One*. 10:e0118985. doi:10.1371/journal.pone.0118985.
- Kidd, P. 2011. Astaxanthin, cell membrane nutrient with diverse clinical benefits and anti-aging potential. *Alternative Medicine Review* 16:355–364.
- Kim, D. Y., D. Vijayan, R. Praveenkumar, J. I. Han, K. Lee, J. Y. Park, W. S. Chang, J. S. Lee, and Y. K. Oh. 2015. Cell-wall disruption and lipid/astaxanthin extraction from microalgae: *Chlorella* and *Haematococcus*. *Biore-source Technology* 199:300–310. doi:10.1016/j.biortech.2015.08.107.
- Kim, J. H., M. J. Chang, H. D. Choi, Y. K. Youn, J. T. Kim, J. M. Oh, and W. G. Shin. 2011. Protective effects of *Haematococcus* astaxanthin on oxidative stress in healthy smokers. *Journal of Medicinal Food* 14:1469–1475. doi:10.1089/jmf.2011.1626.
- Kim, K. N., S. J. Heo, W. J. Yoon, S. M. Kang, G. Ahn, T. H. Yi, and Y. J. Jeon. 2010. Fucoxanthin inhibits the inflammatory response by suppressing the activation of NF- κ B and MAPKs in lipopolysaccharide-induced RAW 264.7 macrophages. *European Journal of Pharmacology* 649:369–375. doi:10.1016/j.ejphar.2010.09.032.
- Kim, S. M., Y. J. Jung, O. N. Kwon, K. H. Cha, B. H. Um, D. Chung, and C. H. A. Pan. 2012. Potential commercial source of fucoxanthin extracted from the microalga *Phaeodactylum tricornutum*. *Applied Biochemistry and Biotechnology* 166:1843–1855. doi:10.1007/s12010-012-9602-2.
- Kim, S. M., S. W. Kang, O. N. Kwon, D. Chung, and C. H. Pan. 2012. Fucoxanthin as a major carotenoid in *Isochrysis* aff. *galbana*: Characterization of extraction for commercial application. *Journal of the Korean Society for Applied Biological* 55:477–483. doi:10.1007/s13765-012-2108-3.
- Kishimoto, Y., H. Yoshida, and K. Kondo. 2016. Potential anti-atherosclerotic properties of astaxanthin. *Marine Drugs* 14:35. doi:10.3390/md14020035.
- Kitada, K., S. Machmudah, M. Sasaki, M. Goto, Y. Nakashima, S. Kumamoto, and T. Hasegawa. 2009. Supercritical CO₂ extraction of pigment components with pharmaceutical importance from *Chlorella vulgaris*. *Journal of Chemical Technology and Biotechnology* 84:657–661. doi:10.1002/jctb.2096.
- Kleinegriss, D. M. M., M. Janssen, W. A. Brandenburg, and R. H. Wijffels. 2011. Continuous production of carotenoids from *Dunaliella salina*. *Enzyme and Microbial Technology* 48:253–259. doi:10.1016/j.enzmictec.2010.11.005.
- Klok, A. J., P. P. Lamers, D. E. Martens, R. B. Draaisma, and R. H. Wijffels. 2014. Edible oils from microalgae: Insights in TAG accumulation. *Trends in Biotechnology* 32:521–528. doi:10.1016/j.tibtech.2014.07.004.
- Kobayashi, M., Y. Kurimura, Y. Sakamoto, and Y. Tsuji. 1997. Selective extraction of astaxanthin and chlorophyll from the green alga *Haematococcus pluvialis*. *Biotechnology Techniques* 11:657–660. doi:10.1023/A:1018455209445.
- Kong, W., N. Liu, J. Zhang, Q. Yang, S. Hua, H. Song, and C. Xia. 2014. Optimization of ultrasound assisted extraction parameters of chlorophyll from *Chlorella vulgaris* residue after lipid separation using response surface methodology. *Journal of Food Science and Technology* 51:2006–2013. doi:10.1007/s13197-012-0706-z.
- Koo, S., K. Cha, D. G. Song, D. Chung, and C. H. Pan. 2012. Optimization of pressurized liquid extraction of zeaxanthin from *Chlorella ellipsoidea*. *Journal of Applied Phycology* 24:725–730. doi:10.1007/s10811-011-9691-2.
- Kraay, G. W., M. Zapata, and M. J. W. Veldhuis. 1992. Separation of chlorophylls c1, c2, and c3 of marine phytoplankton by reversed-phase-C18-high-performance liquid chromatography. *Journal of Phycology* 28:708–712. doi:10.1111/j.0022-3646.1992.00708.x.
- Krinsky, N. I. 1989. Antioxidant functions of carotenoids. *Free Radical Biology and Medicine* 7:617–635. doi:10.1016/0891-5849(89)90143-3.
- Krinsky, N. I. 1991. Effects of carotenoids in cellular and animal systems. *American Journal of Clinical Nutrition* 53:238S–246S. doi:10.1093/ajcn/53.1.238S.
- Krinsky, N. I., J. T. Landrum, and R. A. Bone. 2003. Biologic mechanisms of the protective role of lutein and zeaxanthin in the eye. *Annual Review of Nutrition* 23:171–201. doi:10.1146/annurev.nutr.23.011702.073307.
- Kroger, N., and N. Poulsen. 2008. Diatoms-from cell wall biogenesis to nanotechnology. *Annual Review of Genetics* 42:83–107. doi:10.1146/annurev.genet.41.110306.130109.
- Kuczyńska, P., M. Jemiola-Rzeminska, and K. Strzalka. 2015. Photosynthetic pigments in diatoms. *Marine Drugs* 13:5847–5881. doi:10.3390/md13095847.
- Kuddus, M., P. Singh, G. Thomas, and A. Al-Hazimi. 2013. Recent developments in production and biotechnological applications of C-phyco-cyanin. *BioMed Research International* 2013:1–9. doi:10.1155/2013/742859.
- Kumar, D., N. Kumar, S. Pabbi, S. Walia, and D. W. Dhar. 2013. Protocol optimization for enhanced production of pigments in *Spirulina*. *Indian Journal of Plant Physiology* 18:308–312. doi:10.1007/s40502-013-0045-8.
- Kumar, S. A., M. W. Magnusson, L. C. Ward, N. A. Paul, and L. Brown. 2015. A green algae mixture of *Scenedesmus* and *Schroederiella attenuates* obesity-linked metabolic syndrome in rats. *Nutrients* 7:2771–2787. doi:10.3390/nut7042771.
- Lai, Y. S., P. Parameswaran, A. Li, M. Baez and B. E. Rittmann. 2014. Effects of pulsed electric field treatment on enhancing lipid recovery from the microalga, *Scenedesmus*. *Bioresource Technology* 173:457–461. doi:10.1016/j.biortech.2014.09.124.
- Lamers, P. P., M. Janssen, R. C. De Vos, R. J. Bino, and R. H. Wijffels. 2008. Exploring and exploiting carotenoid accumulation in *Dunaliella salina* for cell-factory applications. *Trends in Biotechnology* 26:631–638. doi:10.1016/j.tibtech.2008.07.002.
- Landrum, J. T., and R. A. Bone. 2001. Lutein, zeaxanthin, and the macular pigment. *Archives of Biochemistry and Biophysics* 385:28–40. doi:10.1006/abbi.2000.2171.
- Lao, Y. M., L. Xiao, L. X. Luo, and J. G. Jiang. 2014. Hypoosmotic expression of *Dunaliella bardawil* ζ -carotene desaturase is attributed to a hypoosmolarity-responsive element different from other key carotenogenic genes. *Plant Physiology* 165:359–372. doi:10.1104/pp.114.235390.
- Larkum, A. W., I. L. Ross, O. Kruse, and B. Hankamer. 2012. Selection, breeding and engineering of microalgae for bioenergy and biofuel production. *Trends in Biotechnology* 30:198–205. doi:10.1016/j.tibtech.2011.11.003.
- Larsdotter, K. 2006. Wastewater treatment with microalgae—a literature review. *Vatten* 62:31–38.
- Lee, J. B., K. Hayashi, M. Hirata, E. Kuroda, E. Suzuki, Y. Kubo, and T. Hayashi. 2006. Antiviral sulfated polysaccharide from *Navicula directa*, a diatom collected from deep-sea water in Toyama Bay. *Biological & Pharmaceutical Bulletin* 29:2135–2139. doi:10.1248/bpb.29.2135.
- Lee, J. C., M. F. Hou, H. W. Huang, F. R. Chang, C. C. Yeh, J. Y. Tang, and H. W. Chang. 2013. Marine algal natural products with anti-oxidative, anti-inflammatory, and anti-cancer properties. *Cancer Cell International* 13:55. doi:10.1186/1475-2867-13-55.
- Lelyana, R. 2016. Role of marine natural ingredient fucoxanthin on body's immune response of obesity. *Journal of Nanomedicine and Nanotechnology* 7:2. doi:10.4172/2157-7439.1000397.
- Li, H. B., Y. Jiang, and F. Chen. 2002. Isolation and purification of lutein from the microalga *Chlorella vulgaris* by extraction after saponification. *Journal of Agricultural and Food Chemistry* 50:1070–1072. doi:10.1021/jf010220b.
- Lian, F., and X. D. Wang. 2008. Enzymatic metabolites of lycopene induce Nrf2-mediated expression of phase-II detoxifying/antioxidant enzymes in human bronchial epithelial cells. *International Journal of Cancer* 123:1262–1268. doi:10.1002/ijc.23696.
- Liang, Y., N. Sarkany, and Y. Cui. 2009. Biomass and lipid productivities of *Chlorella vulgaris* under autotrophic, heterotrophic and mixotrophic growth conditions. *Biotechnology Letters* 31:1043–1049. doi:10.1007/s10529-009-9975-7.
- Liau, B. C., C. T. Shen, F. P. Liang, S. E. Hong, S. L. Hsu, T. T. Jong, and C. M. J. Chang. 2010. Supercritical fluids extraction and anti-solvent purification of carotenoids from microalgae and associated bioactivity. *Journal of Supercritical Fluids* 55:169–175. doi:10.1016/j.supflu.2010.07.002.
- Liu, J., and Q. Hu. 2013. *Chlorella*: Industrial production of cell mass and chemicals. In *Hand Book of Microalgal Culture: Applied Phycology and Biotechnology*, Eds. A. Richmond, and Q. Hu, pp. 329–338. 2 ed. Oxford, UK: John Wiley & Sons, Ltd.
- Liu, J., Z. Sun, H. Gerken, Z. Liu, Y. Jiang, and F. Chen. 2014. *Chlorella zofingiensis* as an alternative microalgal producer of astaxanthin: Biology and industrial potential. *Marine Drugs* 12:3487–3515. doi:10.3390/md12063487.

- Lohr, M., and C. Wilhelm. 1999. Algae displaying the diadinoxanthin cycle also possess the violaxanthin cycle. *Proceedings of the National Academy of Sciences (USA)* 96:8784–8789. doi:10.1073/pnas.96.15.8784.
- Lorenz, R. T., and G. R. Cysewski. 2000. Commercial potential for *Haematococcus* microalgae as a natural source of astaxanthin. *Trends in Biotechnology* 18:160–167. doi:10.1016/S0167-7799(00)01433-5.
- Lu, Y. P., S. Y. Liu, H. Sun, X. M. Wu, J. J. Li, and L. Zhu. 2010. Neuroprotective effect of astaxanthin on H₂O₂-induced neurotoxicity *in vitro* and on focal cerebral ischemia *in vivo*. *Brain Research* 1360:40–48. doi:10.1016/j.brainres.2010.09.016.
- Lubian, L. M., O. Montero, I. Moreno-Garrido, I. Emma Huertas, C. Sobrino, M. G. Valle, and G. Pares. 2000. *Nannochloropsis* (Eustigmatophyceae) as source of commercially valuable pigments. *Journal of Applied Phycology* 12:249–255. doi:10.1023/A:1008170915932.
- Luvizotto, R. D. A. M., A. F. Nascimento, E. Imaizumi, D. T. Pierine, S. J. Conde, C. R. Correa, K. J. Yeum, and A. L. A. Ferreira. 2013. Lycopene supplementation modulates plasma concentrations and epididymal adipose tissue mRNA of leptin, resistin and IL-6 in diet-induced obese rats. *British Journal of Nutrition* 110:1803–1809. doi:10.1017/S0007114513001256.
- Ma, L., H. L. Dou, Y. Q. Wu, Y. M. Huang, Y. B. Huang, X. R. Xu, Z. Y. Zou, and X. M. Lin. 2012. Lutein and zeaxanthin intake and the risk of age-related macular degeneration: A systematic review and meta-analysis. *British Journal of Nutrition*. 107:350–359. doi:10.1017/S0007114511004260.
- Macias-Sanchez, M. D., C. Mantell, M. Rodriguez, E. Martinez de la Ossa, L. M. Lubian, and O. Montero. 2009. Comparison of supercritical fluid and ultrasound-assisted extraction of carotenoids and chlorophyll-a from *Dunaliella salina*. *Talanta* 77:948–952. doi:10.1016/j.talanta.2008.07.032.
- Macias-Sanchez, M. D., C. Mantell, M. Rodriguez, E. Martinez de la Ossa, L. M. Lubian, and O. Montero. 2007. Supercritical fluid extraction of carotenoids and chlorophyll a from *Synechococcus* sp. *Journal of Supercritical Fluids* 39:323–329. doi:10.1016/j.supflu.2006.03.008.
- Macias-Sanchez, M. D., C. Mantell, M. Rodriguez, E. Martinez de la Ossa, L. M. Lubian, and O. Montero. 2005. Supercritical fluid extraction of carotenoids and chlorophyll-a from *Nannochloropsis gaditana*. *Journal of Food Engineering* 66:245–251. doi:10.1016/j.jfoodeng.2004.03.021.
- Maeda, H., M. Hosokawa, T. Sashima, K. Funayama, and K. Miyashita. 2005. Fucoxanthin from edible seaweed, *Undaria pinnatifida*, shows anti-obesity effect through UCP1 expression in white adipose tissues. *Biochemical and Biophysical Research Communications* 332:392–397. doi:10.1016/j.bbrc.2005.05.002.
- Maeda, H., M. Hosokawa, T. Sashima, K. Murakami-Funayama, and K. Miyashita. 2009. Anti-obesity and anti-diabetic effects of fucoxanthin on diet-induced obesity conditions in a murine model. *Molecular Medicine Reports* 2:897–902. doi:10.3892/mmr.00000189.
- Mandotra, S. K., P. Kumar, M. R. Suseela, and P. W. Ramteke. 2014. Fresh water green microalga *Scenedesmus abundans*: A potential feedstock for high quality biodiesel production. *Bioresource Technology* 156:42–47. doi:10.1016/j.biortech.2013.12.127.
- Maoka, T., H. Tokuda, N. Suzuki, H. Kato, and H. Etoh. 2012. Anti-oxidative, anti-tumor promoting, and anti-carcinogenesis activities of nitro-astaxanthin and nitrolutein, the reaction products of astaxanthin and lutein with peroxynitrite. *Marine Drugs* 10:1391–1399. doi:10.3390/md10061391.
- Markovits, A., R. Conejeros, L. López, and M. Lutz. 1992. Evaluation of marine microalga *Nannochloropsis* sp as a potential dietary supplement, chemical, nutritional and short term toxicological evaluation in rats. *Nutrition Research* 12:1273–1284. doi:10.1016/S0271-5317(05)80784-5.
- Martinez, F., C. Ascaso, and M. I. Orus. 1991. Morphometric and stereologic analysis of *Chlorella vulgaris* under heterotrophic growth conditions. *Annals of Botany* 67:239–245. doi:10.1093/oxfordjournals.aob.a088128.
- Martin-Jezequel, V., M. Hildebrand, and M. A. Brzezinski. 2000. Silicon metabolism in diatoms: Implications for growth. *Journal of Phycology* 36:821–840. doi:10.1046/j.1529-8817.2000.00019.x.
- Masojidek, J., G. Torzillo, J. Kopecky, M. Koblizek, L. Nidiaci, J. Komenda, and A. Sacchi. 2000. Changes in chlorophyll fluorescence quenching and pigment composition in the green alga *Chlorococcum* sp. grown under nitrogen deficiency and salinity stress. *Journal of Applied Phycology* 12:417–426. doi:10.1023/A:1008165900780.
- McMillan, J. R., I. A. Watson, M. Ali, and W. Jaafar. 2013. Evaluation and comparison of algal cell disruption methods: Microwave, water bath, blender, ultrasonic and laser treatment. *Applied Energy* 103:128–134. doi:10.1016/j.apenergy.2012.09.020.
- Medlin, L. K. 2016. Evolution of the diatoms: Major steps in their evolution and a review of the supporting molecular and morphological evidence. *Phycologia* 55:79–103. doi:10.2216/15-105.1.
- Mendes, R. L., H. L. Fernandes, J. P. Coelho, E. C. Reis, J. M. S. Cabral, J. M. Novais, and A. F. Palabra. 1995. Supercritical CO₂ extraction of carotenoids and other lipids from *Chlorella vulgaris*. *Food Chemistry* 53:99–103. doi:10.1016/0308-8146(95)95794-7.
- Mendes, R. L., B. P. Nobre, M. T. Cardoso, A. P. Pereira, and A. F. Palavra. 2003. Supercritical carbon dioxide extraction of compounds with pharmaceutical importance from microalgae. *Inorganica Chimica Acta* 356:328–334. doi:10.1016/S0020-1693(03)00363-3.
- Merchant, R. E., and C. A. Andre. 2001. A review of recent clinical trials of the nutritional supplement *Chlorella pyrenoidosa* in the treatment of fibromyalgia, hypertension, and ulcerative colitis. *Alternative Therapies in Health and Medicine* 7:79–91.
- Merchant, R. E., C. A. Andre, and D. A. Sica. 2002. Nutritional supplementation with *Chlorella pyrenoidosa* for mild to moderate hypertension. *Journal of Medicinal Food* 5:141–152. doi:10.1089/10966200260398170.
- Metzger, P., and C. Largeau. 2005. *Botryococcus braunii*: A rich source for hydrocarbons and related ether lipids. *Applied Microbiology and Biotechnology* 66:486–496. doi:10.1007/s00253-004-1779-z.
- Minhas, A. K., P. Hodgson, C. J. Barrow, and A. Adholeya. 2016. A review on the assessment of stress conditions for simultaneous production of microalgal lipids and carotenoids. *Frontiers in Microbiology* 7:546. doi:10.3389/fmicb.2016.00546.
- Miranda, M. S., R. G. Cintra, S. B. Barros, J. Mancini-Filho. 1998. Antioxidant activity of the microalga *Spirulina maxima*. *Brazilian Journal of Medical and Biological Research* 31:1075–1079. doi:10.1590/S0100-879X1998000800007.
- Mogedas, B., C. Casal, E. Forjan, and C. Vilchez. 2009. β -Carotene production enhancement by UV-A radiation in *Dunaliella bardawil* cultivated in laboratory reactors. *Journal of Bioscience and Bioengineering* 108:47–51. doi:10.1016/j.jbiosc.2009.02.022.
- Mohammed, M. K., and M. K. Mohd. 2011. Production of carotenoids in *Spirulina platensis* in response to indole acetic acid (IAA). *International Journal of Engineering Science and Technology* 3:4973–4979.
- Muntean, E., N. Muntean, N. Dragos, and V. Bercea. 2008. Carotenoids as biomarkers in *Botryococcus braunii* algae. *Agricultural Science Research Journal* 40:49–54.
- Murthy, K. N., A. Vanitha, J. Rajesha, M. M. Swamy, P. R. Sowmya, and G. A. Ravishanker. 2005. *In vivo* antioxidant activity of carotenoids from *Dunaliella salina* a green microalga. *Life Sciences* 76:1381–1390. doi:10.1016/j.lfs.2004.10.015.
- Nobre, B. P., F. Villalobos, B. E. Barragan, A. C. Oliveira, A. P. Batista, P. A. S. S. Marques, and L. Gouveia. 2013. A biorefinery from *Nannochloropsis* sp. microalga extraction of oils and pigments. Production of biohydrogen from the leftover biomass. *Bioresource Technology* 135:128–136. doi:10.1016/j.biortech.2012.11.084.
- Nobre, B., F. Marcelo, R. Passos, L. Beirao, A. Palavra, L. Gouveia, and R. Mendes. 2006. Supercritical carbon dioxide extraction of astaxanthin and other carotenoids from the microalga *Haematococcus pluvialis*. *European Food Research and Technology* 223:787–790. doi:10.1007/s00217-006-0270-8.
- Odjadjare, E. C., T. Mutanda, and A. O. Olaniran. 2015. Potential biotechnological application of microalgae: A critical review. *Critical Reviews in Biotechnology* 37:37–52. doi:10.3109/07388551.2015.1108956.
- Ogawa, T., and S. Aiba. 1981. Bioenergetic analysis of mixotrophic growth in *Chlorella vulgaris* and *Scenedesmus acutus*. *Biotechnology and Bioengineering* 23:1121–1132. doi:10.1002/bit.260230519.
- Okada, S., M. Mastuda, M. Murakami, and K. Yamaguchi. 1996. Botryoxanthin-A a new member of the new class of carotenoids from green microalga *Botryococcus braunii*, Berkeley. *Tetrahedron Letters* 37:1065–1068. doi:10.1016/0040-4039(95)02349-6.
- Okada, S., I. Tonegawa, M. Mastuda, M. Murakami, and K. Yamaguchi. 1997. Braunixanthins 1 and 2, new carotenoids from the green

- microalga *Botryococcus braunii*. *Tetrahedron* 53:11307–11316. doi:10.1016/S0040-4020(97)00705-9.
- Okada, S., I. Tonegawa, M. Mastuda, M. Murakami, and K. Yamaguchi. 1998. Botryoxanthin-B and alpha-botryoxanthin-A from green microalga *Botryococcus braunii*. *Phytochemistry* 47:1111–1115. doi:10.1016/S0031-9422(98)80082-4.
- Olaizola, M. 2000. Commercial production of astaxanthin from *Haematococcus pluvialis* using 25,000-liter outdoor photobioreactors. *Journal of Applied Phycology* 12:499–506. doi:10.1023/A:1008159127672.
- Omenn, G. S., G. E. Goodman, M. D. Thornquist, J. Balmes, M. R. Cullen, A. Glass, J. P. Keogh, F. L. Meyskens, B. Valanis, J. H. Williams, S. Barnhart, and S. Hammar. 1996. Effects of a combination of β -carotene and vitamin A on lung cancer and cardiovascular disease. *The New England Journal of Medicine* 334:1150–1155. doi:10.1056/NEJM199605023341802.
- Orosa, M., J. F. Valero, C. Herrero, and J. Abalde. 2001. Comparison of the accumulation of astaxanthin in *Haematococcus pluvialis* and other green microalgae under N-starvation and high light conditions. *Biotechnology Letters* 23:1079–1085. doi:10.1023/A:1010510508384.
- Palozza, P., and N. I. Krinsky. 1992. β -Carotene and α -tocopherol are synergistic antioxidants. *Archives of Biochemistry and Biophysics* 297:184–187. doi:10.1016/0003-9861(92)90658-J.
- Palozza, P., S. Serini, F. Di Nicuolo, E. Piccioni, and G. Calviello. 2003. Pro-oxidant effects of β -carotene in cultured cells. *Molecular Aspects of Medicine* 24:353–362. doi:10.1016/S0098-2997(03)00031-1.
- Palozza, P., R. Simone, A. Catalano, M. Russo, and V. Bohm. 2012. Lycopene modulation of molecular targets affected by smoking exposure. *Current Cancer Drug Targets* 12:640–657. doi:10.2174/156800912801784866.
- Pancha, I., K. Chokshi, B. George, T. Ghosh, C. Paliwal, R. Maurya, and S. Mishra. 2014. Nitrogen stress triggered biochemical and morphological changes in the microalgae *Scenedesmus* sp. CCNM 1077. *Bioresource Technology* 156:146–154. doi:10.1016/j.biortech.2014.01.025.
- Pangestuti, R., and S. K. Kim. 2011. Biological activities and health benefit effects of natural pigments derived from marine algae. *Journal of Functional Foods* 3:255–266. doi:10.1016/j.jff.2011.07.001.
- Panis, G., and J. R. Carreon. 2016. Commercial astaxanthin production derived by green alga *Haematococcus pluvialis*: A microalgae process model and a techno-economic assessment all through production line. *Algal Research* 18:175–190. doi:10.1016/j.algal.2016.06.007.
- Park, J. S., J. H. Chyun, Y. K. Kim, L. L. Line, and B. P. Chew. 2010. Astaxanthin decreased oxidative stress and inflammation and enhanced immune response in humans. *Nutrition and Metabolism* 7:1–10.
- Park, J. Y., M. S. Park, Y. C. Lee, and J. W. Yang. 2015. Advances in direct transesterification of algal oils from wet biomass. *Bioresource Technology* 184:267–275. doi:10.1016/j.biortech.2014.10.089.
- Pashkow, F. J., D. G. Watumull, and C. L. Campbell. 2008. Astaxanthin: A novel potential treatment for oxidative stress and inflammation in cardiovascular disease. *American Journal of Cardiology* 101:58D–68D. doi:10.1016/j.amjcard.2008.02.010.
- Pasquet, V., J. R. Cherouvrier, F. Farhat, V. Thiery, J. M. Piot, J. B. Berard, R. Kaas, B. Serive, T. Patrice, J. P. Cadoret, and L. Picot. 2011. Study on the microalgal pigments extraction process: Performance of microwave assisted extraction. *Process Biochemistry (Barking, UK)* 46:59–67. doi:10.1016/j.procbio.2010.07.009.
- Pelach, D., A. Sintov, and E. Cohen. 2004. The effect of salt stress on the production of canthaxanthin and astaxanthin by *Chlorella zofingiensis* grown under limited light intensity. *World Journal of Microbiology and Biotechnology* 20:483–486. doi:10.1023/B:WIBI.0000040398.93103.21.
- Peng, J., J. P. Yuan, C. F. Wu, and J. H. Wang. 2011. Fucoxanthin, a marine carotenoid present in brown seaweeds and diatoms: Metabolism and bioactivities relevant to human health. *Marine Drugs* 9:1806–1828. doi:10.3390/md9101806.
- Peng, K. T. 2014. Delta-5 fatty acid desaturase upregulates the synthesis of polyunsaturated fatty acids in the marine diatom *Phaeodactylum tricornutum*. *Journal of Agricultural and Food Chemistry* 62:8773–8776. doi:10.1021/jf5031086.
- Perl, T. M., L. Bedard, T. Kosatsky, J. C. Hockin, E. C. Todd, and R. S. Remis. 1990. An out break of toxic encephalopathy caused by eating mussels contaminated with domoic acid. *New England Journal of Medicine* 322:1775–1780. doi:10.1056/NEJM199006213222504.
- Phang, S. M., E. M. Mustafa, R. R. Ambati, N. M. Nik Sulaiman, P. E. Lim, X. Dommange, C. Schwob, and K. E. Liew. 2015. Check list of microalgae collected from different habitats in peninsular Malaysia for selection of algal biofuel feed stocks. *Malaysian Journal of Science* 34:148–174.
- Pignolet, O., S. Jubeau, C. Vaca-Garcia, and P. Michaud. 2013. Highly valuable microalgae: Biochemical and topological aspects. *Journal of Industrial Microbiology and Biotechnology* 40:781–96. doi:10.1007/s10295-013-1281-7.
- Pirastu, L., M. Darwish, F. L. Chu, F. Perreault, L. Sirois, L. Sleno, and R. Popovic. 2012. Carotenoid production and change of photosynthetic functions in *Scenedesmus* sp. exposed to nitrogen limitation and acetate treatment. *Journal of Applied Phycology* 24:117–124. doi:10.1007/s10811-011-9657-4.
- Plaza, M., A. Cifuentes, and E. Ibanez. 2008. In the search of new functional food ingredients from algae. *Trends in Food Science and Technology* 19:31–39. doi:10.1016/j.tifs.2007.07.012.
- Plaza, M., M. Herrero, A. Cifuentes, and E. Ibanez. 2009. Innovative natural functional ingredients from microalgae. *Journal of Agricultural and Food Chemistry* 57:7159–7170. doi:10.1021/jf901070g.
- Poonkum, W., S. Powtongsook, and P. Pavasant. 2015. Astaxanthin induction in microalga *Haematococcus pluvialis* with flat panel airlift photobioreactors under indoor and outdoor conditions. *Preparative Biochemistry and Biotechnology* 45:1–17. doi:10.1080/10826068.2013.867871.
- Pragya, N., K. K. Pandey, and P. K. Sahoo. 2013. A review on harvesting, oil extraction and biofuels production technologies from microalgae. *Renewable and Sustainable Energy Reviews* 24:159–171. doi:10.1016/j.rser.2013.03.034.
- Prestegard, S. K., L. Oftedal, G. Nygaard, K. H. Skjaerven, G. Knutsen, S. O. Doskeland, R. T. Coyne, and L. Herfindal. 2009. Marine benthic diatoms contain compounds able to induce leukemia cell death and modulate blood platelet activity. *Marine Drugs* 7:605–623. doi:10.3390/md7040605.
- Qin, S., G. X. Liu, and Z. Y. Hu. 2008. The accumulation and metabolism of astaxanthin in *Scenedesmus obliquus* (Chlorophyceae). *Process Biochemistry (Barking, UK)* 43:795–802. doi:10.1016/j.procbio.2008.03.010.
- Ragni, M., and M. R. Dalcá. 2007. Circadian variability in the photobiology of *Phaeodactylum tricornutum*: Pigment content. *Journal of Plankton Research* 29:141–156. doi:10.1093/plankt/fbm002.
- Ramel, F., S. Birtic, S. Cuine, C. Triantaphylides, J. L. Ravanat, and M. Havaux. 2012. Chemical quenching of singlet oxygen by carotenoids in plants. *Plant Physiology* 159:1267–1278. doi:10.1104/pp.111.182394.
- Ranga Rao, A. 2011. *Production of astaxanthin from cultured green alga Haematococcus pluvialis and its biological activities*. Ph.D Thesis, University of Mysore, Mysore, India.
- Ranga Rao, A., V. Baskaran, R. Sarada, and G. A. Ravishankar. 2013b. *In vivo* bioavailability and antioxidant activity of carotenoids from microalgal biomass- a repeated dose study. *Food Research International* 54:711–777. doi:10.1016/j.foodres.2013.07.067.
- Ranga Rao, A., C. Dayananda, R. Sarada, T. R. Shamala, and G. A. Ravishankar. 2007a. Effect of salinity on growth of green alga *Botryococcus braunii* and its constituents. *Bioresource Technology* 98:560–564. doi:10.1016/j.biortech.2006.02.007.
- Ranga Rao, A., G. Deepika, G. A. Ravishankar, R. Sarada, B. P. Narasimha Rao, Y. Su, and B. Lei. 2017. *Botryococcus* as an alternative source of carotenoids and its possible applications- an overview. *Critical Reviews in Biotechnology* 22:1–18 (In press).
- Ranga Rao, A., S. M. Phang, R. Sarada, and G. A. Ravishankar. 2014a. Astaxanthin: Sources, extraction, stability, biological activities and its commercial applications-A review. *Marine Drugs* 12:128–152. doi:10.3390/md12010128.
- Ranga Rao, A., R. L. Raghunath Reddy, V. Baskaran, R. Sarada, and G. A. Ravishankar. 2010a. Characterization of microalgal carotenoids by mass spectrometry and their bioavailability and antioxidant properties elucidated in rat model. *Journal of Agricultural and Food Chemistry* 58:8553–8559. doi:10.1021/jf101187k.
- Ranga Rao, A., G. A. Ravishankar, and R. Sarada. 2012a. Cultivation of green alga *Botryococcus braunii* in raceway, circular ponds under outdoor conditions and its growth, hydrocarbon production. *Bioresource Technology* 123:528–533. doi:10.1016/j.biortech.2012.07.009.

- Ranga Rao, A., A. H. Reddy, and S. M. Aradhya. 2010c. Antibacterial properties of *Spirulina platensis*, *Haematococcus pluvialis*, *Botryococcus braunii* micro algal extracts. *Current Trends in Biotechnology and Pharmacy* 4:809–819.
- Ranga Rao, A., R. Sarada, V. Baskaran, and G. A. Ravishankar. 2006. Antioxidant activity of *Botryococcus braunii* extract elucidated in vitro models. *Journal of Agricultural and Food Chemistry* 54:4593–4599. doi:10.1021/jf060799j.
- Ranga Rao, A., R. Sarada, and G. A. Ravishankar. 2010b. Enhancement of carotenoids in green alga-*Botryococcus braunii* in various autotrophic media under stress conditions. *International Journal of Biomedical Science* 4:87–92.
- Ranga Rao, A., R. Sarada, and G. A. Ravishankar. 2007b. Influence of CO₂ on growth and hydrocarbon production in *Botryococcus braunii*. *Journal of Microbiology and Biotechnology* 17:414–419.
- Ranga Rao, A., R. Sarada, V. Baskaran, and G. A. Ravishankar. 2009. Identification of carotenoids from green alga *Haematococcus pluvialis* by HPLC and LC-MS (APCI) and their antioxidant properties. *Journal of Microbiology and Biotechnology* 19:1333–1341.
- Ranga Rao, A., R. Sarada, G. A. Ravishankar, and S. M. Phang. 2016. Industrial production of microalgal cell-mass and bioactive constituents from green microalga-*Botryococcus braunii*. In *Recent Advances in Microalgal Biotechnology*. Eds. J. Liu, Z. Sun, and G. Henri, pp.103–126, 731 Gull Ave, Foster City. CA 94404, USA: OMICS Group Incorporation.
- Ranga Rao, A., H. N. Sindhuja, S. M. Dharmesh, K. U. Sankar, R. Sarada, and G. A. Ravishankar. 2013a. Effective inhibition of skin cancer, tyrosinase, and antioxidant properties by astaxanthin and astaxanthin esters from the green alga *Haematococcus pluvialis*. *Journal of Agricultural and Food Chemistry* 61:3842–3851. doi:10.1021/jf304609j.
- Ranga Rao, A., G. Yingchun, Z. Hu, M. Sommerfeld, and Q. Hu. 2012b. Comparative study of predators in mass cultures of *Chlorella zofingiensis*, *Scenedesmus dimorphus* and *Nannochloropsis oceanica* in ponds and photobioreactors. Colorado, USA: Algal Biomass Organization.
- Rao, A. V., and L. G. Rao. 2007. Carotenoids and human health. *Pharmacological Research* 55:207–216. doi:10.1016/j.phrs.2007.01.012.
- Rebollosa-Fuentes, M. M., A. Navarro-Perez, F. Garcia-Camacho, J. J. Ramos-Miras, and J. L. Guil-Guerrero. 2001. Biomass nutrient profiles of the microalga *Nannochloropsis*. *Journal of Agricultural and Food Chemistry* 49:2966–2972. doi:10.1021/jf0010376.
- Regnier, P., J. Bastias, V. Rodriguez-Ruiz, N. Caballero-Casero, C. Caballo, D. Sicilia, A. Fuentes, M. Maire, M. Crepin, and D. Letourneur, et al. 2015. Astaxanthin from *Haematococcus pluvialis* prevents oxidative stress on human endothelial cells without toxicity. *Marine Drugs* 13:2857–2874. doi:10.3390/md13052857.
- Rijstenbil, J. W. 2003. Effects of UVB radiation and salt stress on growth, pigments and oxidative defence of the marine diatom *Cylindrotheca closterium*. *Marine Ecology Progress Series* 254:37–48. doi:10.3354/meps254037.
- Roche, F. 1987. Astaxanthin: Human food safety summary. In: *Astaxanthin As a Pigment in Salmon Feed, Color Additive Petition 7CO₂ 1 1*, pp. 43. *United States Food and Drug Administration (USFDA)*. Basel, Switzerland: Hoffman-La Roche Ltd.
- Rodriguez-Amaya, D. B. 2016. Natural food pigments and colorants. *Current Opinion in Food Science* 7:20–26. doi:10.1016/j.cofs.2015.08.004.
- Rodriguez-Garcia, I., and J. L. Guil-Guerrero. 2008. Evaluation of the antioxidant activity of three micro algal species for use as dietary supplements and in the preservation of foods. *Food Chemistry* 108:1023–1026. doi:10.1016/j.foodchem.2007.11.059.
- Rokkaku, T., R. Kimura, C. Ishikawa, T. Yasumoto, M. Senba, F. Kanaya, and N. Mori. 2013. Anticancer effects of marine carotenoids, fucoxanthin and its deacetylated product, fucoxanthinol, on osteosarcoma. *International Journal of Oncology* 43:1176–1186. doi:10.3892/ijo.2013.2019.
- Ross, A. C., and E. H. Harrison. 2014. Vitamin A: Nutrition aspects of retinoids and carotenoids. In *Handbook of Vitamins*, eds. J. Zempleni, W. Suttie, J. F. Gregory, and P. J. Stover, pp.1–50. Boca Raton, FL: CRC Press.
- Roy, S. S., and R. Pal. 2015. Microalgae in aquaculture: A review with special references to nutritional value and fish dietetics. *Proceedings of the Zoological Society* 68:1–8. doi:10.1007/s12595-013-0089-9.
- Safi, C., B. Zebib, O. Merah, P. Y. Pontalier, and C. Vaca-Garcia. 2014. Morphology, composition, production, processing and applications of *Chlorella vulgaris*: a review. *Renewable and Sustainable Energy Reviews* 35:265–278. doi:10.1016/j.rser.2014.04.007.
- Sanchez, J. F., J. M. Fernandez, F. G. Acien, A. Rueda, J. Pérez-Parra, and E. Molina. 2008. Influence of culture conditions on the productivity and lutein content of the new strain *Scenedesmus almeriensis*. *Process Biochemistry (Barking, UK)* 43:398–405. doi:10.1016/j.procbio.2008.01.004.
- Sanchez-Saavedra, M. P., and D. Voltolina. 2002. Effect of photon fluence rates of white and blue-green light on growth efficiency and pigment content of three diatom species in batch cultures. *Ciencias Marinas* 28:273–279. doi:10.7773/cm.v28i3.225.
- Sandesh Kamath, B., R. Vidhyavathi, R. Sarada, and G. A. Ravishankar. 2008. Enhancement of carotenoids by mutation and stress induced carotenogenic genes in *Haematococcus pluvialis* mutants. *Bioresource Technology* 99:8667–8673. doi:10.1016/j.biortech.2008.04.013.
- Sarada, R., A. Ranga Rao, B. K. Sandesh, C. Dayananda, N. Anila, V. S. Chauhan, and G. A. Ravishankar. 2012. Influence of different culture conditions on yield of biomass and value added products in microalgae. *Dynamic Biochemistry, Process Biotechnology and Molecular Biology* 6:77–85.
- Sarada, R., R. Vidhyavathi, D. Usha, and G. A. Ravishankar. 2006. An efficient method for extraction of astaxanthin from green alga *Haematococcus pluvialis*. *Journal of Agricultural and Food Chemistry* 54:7585–7588. doi:10.1021/jf060737t.
- Schweiggert, R. M., and R. Carle. 2016. Carotenoid production by bacteria, microalgae, and fungi. In *Carotenoids: Nutrition, Analysis and Technology*, Eds. A. Kaczor, and M. Baranska, pp.217–240. Chichester, UK: John Wiley & Sons, Ltd.
- Seyfabadi, J., Z. Ramezanzpour, and Z. A. Khoeyi. 2010. Protein, fatty acid and pigment content of *Chlorella vulgaris* under different light regimes. *Journal of Applied Phycology* 23:721–726. doi:10.1007/s10811-010-9569-8.
- Shah, M. M. R., Y. Liang, J. J. Cheng, and M. Daroch. 2016. Astaxanthin-producing green microalga *Haematococcus pluvialis* from single cell to high value commercial products. *Frontiers in Plant Science* 7:531. doi:10.3389/fpls.2016.00531.
- Shaish, A., A. Harari, L. Hananshvili, H. Cohen, R. Bitzur, T. Luvish, E. Ulman, M. Golan, A. Ben-Amotz, D. Gavish, Z. Rotstein, and D. Harats. 2006. 9-cis-carotene-rich powder of the alga *Dunaliella bardawil* increases plasma HDL-cholesterol in fibrates-treated patients. *Atherosclerosis* 189:215–221. doi:10.1016/j.atherosclerosis.2005.12.004.
- Sharma, R., G. Pal Singh, and V. K. Sharma. 2012. Effect of culture conditions on growth and biochemical profile of *Chlorella vulgaris*. *Journal of Plant Pathology and Microbiology* 3:131. doi:10.4172/2157-7471.1000131.
- Sharoni, Y., M. Danilenko, N. Dubi, A. Ben-Dor, and J. Levy. 2004. Carotenoids and transcription. *Archives of Biochemistry and Biophysics* 430:89–96. doi:10.1016/j.abb.2004.03.009.
- Sharoni, Y., K. Linnewiel-Hermoni, G. Zango, M. Khanin, H. Salman, A. Veprik, M. Danilenko, and J. Levy. 2012. The role of lycopene and its derivatives in the regulation of transcription systems: Implications for cancer prevention. *American Journal of Clinical Nutrition* 96S:1173S–1178S. doi:10.3945/ajcn.112.034645.
- Shi, X. M., Y. Jiang, and F. Chen. 2002. High-yield production of lutein by the green microalga *Chlorella protothecoides* in heterotrophic fed batch culture. *Biotechnology Progress* 18:723–727. doi:10.1021/bp0101987.
- Shim, J. Y., H. S. Shin, J. G. Han, H. S. Park, B. L. Lim, K. W. Chung, and A. S. Om. 2008. Protective effects of *Chlorella vulgaris* on liver toxicity in cadmium-administered rats. *Journal of Medicinal Food* 11:479–485. doi:10.1089/jmf.2007.0075.
- Simeunovic, J., K. Beslin, Z. Svircev, D. Kovac, and O. Babic. 2013. Impact of nitrogen and drought on phycobiliprotein content in terrestrial cyanobacterial strains. *Journal of Applied Phycology* 25:597–607. doi:10.1007/s10811-012-9894-1.
- Simeunovic, J., S. Markovic, D. Kovac, A. Misan, A. Mandic, and Z. Svircev. 2012. Filamentous cyanobacteria from vojvodina region as source of phycobiliprotein pigments as potential natural colorants. *Food Feed Research* 39:23–31.

- Singh, J., and S. Gu. 2010. Commercialization potential of microalgae for biofuels production. *Renewable and Sustainable Energy Reviews* 14:2596–2610. doi:10.1016/j.rser.2010.06.014.
- Sivathanu, B., and S. Palaniswamy. 2012. Purification and characterization of carotenoids from green algae *Chlorococcum humicola* by HPLC-NMR and LC-MS-APCI. *Biomedicine and Preventive Nutrition* 2:276–282. doi:10.1016/j.bionut.2012.04.006.
- Solovchenko, A., A. Lukyanov, O. Solovchenko, S. Didi-Cohen, S. Bous-siba, and I. Khozin-Goldberg. 2014. Interactive effects of salinity, high light, and nitrogen starvation on fatty acid and carotenoid profiles in *Nannochloropsis oceanica* CCALA 804. *European Journal of Lipid Science and Technology* 166:635–644. doi:10.1002/ejlt.201300456.
- Spolaore, P., C. Joannis-Cassan, E. Duran, and A. Isambert. 2006. Commercial applications of microalgae. *Journal of Bioscience and Bioengineering* 10:87–96. doi:10.1263/jbb.101.87.
- Stahl, W., and H. Sies. 2012. β -carotene and other carotenoids in protection from sunlight. *American Journal of Clinical Nutrition* 96:1179–1184. doi:10.3945/ajcn.112.034819.
- Stahl, W., N. Ale-Agha, and M. C. Polidori. 2002. Non-antioxidant properties of carotenoids. *Biological Chemistry* 383:553–558. doi:10.1515/BC.2002.056.
- Stahl, W., and H. Sies. 2005. Bioactivity and protective effects of natural carotenoids. *Biochimica et Biophysica Acta - Molecular Basis of Disease* 1740:101–107. doi:10.1016/j.bbdis.2004.12.006.
- Stahl, W., U. Heinrich, H. Jungmann, H. Sies, and H. Tronnier. 2000. Carotenoids and carotenoids plus vitamin E protect against ultraviolet light induced erythema in humans. *American Journal of Clinical Nutrition* 71:795–798.
- Stahl, W., S. Nicolai, K. Briviba, M. Hanusch, G. Broszeit, M. Peters, H. D. Martin, and H. Sies. 1997. Biological activities of natural and synthetic carotenoids: Induction of gap junctional communication and singlet oxygen quenching. *Carcinogenesis* 18:89–92. doi:10.1093/carcin/18.1.89.
- Starkenburg, S. R., K. J. Kwon, R. K. Jha, C. McKay, M. Jacobs, O. Chertkov, and R. A. Cattolico. 2014. A pangenomic analysis of the *Nannochloropsis* organellar genomes reveals novel genetic variations in key metabolic genes. *BMC Genomics* 15:212. doi:10.1186/1471-2164-15-212.
- Steinbrenner, J., and H. Linden. 2001. Regulation of two carotenoid biosynthesis genes coding for phytoene synthase and carotenoid hydrolase during stress-induced astaxanthin formation in the green alga *Haematococcus pluvialis*. *Plant Physiology* 125:810–817. doi:10.1104/pp.125.2.810.
- Tan, G., and M. P. Suter. 2011. Vitamin A, nutrition, and health values of algae: *Spirulina*, *Chlorella* and *Dunaliella*. *Journal of Pharmacy and Nutrition Sciences* 1:111–118.
- Taucher, J., S. Baer, P. Schwerna, D. Hofmann, M. Hummer, R. Buchholz, and A. Becker. 2016. Cell disruption and pressurized liquid extraction of carotenoids from microalgae. *Journal of Thermodynamics and Catalysis* 7:1–7. doi:10.4172/2157-7544.1000158.
- Tokusoglu, O., and M. K. Unal. 2003. Biomass nutrient profiles of three microalgae: *Spirulina platensis*, *Chlorella vulgaris*, and *Isochrysis galbana*. *Journal of Food Science* 68:1144–1148. doi:10.1111/j.1365-2621.2003.tb09615.x.
- Tominaga, K., N. Hongo, M. Karato, and E. Yamashita. 2012. Cosmetic benefits of astaxanthin on humans subjects. *Acta Biochimica Polonica* 59:43–47.
- Tonegawa, I., S. Okada, M. Murakami, and K. Yamagushi. 1998. Pigment composition of the green microalga *Botryococcus braunii* Kawagushi-1. *Fisheries Science* 64:305–308. doi:10.2331/fishsci.64.305.
- Tran, D., N. Doan, C. Louime, M. Giordano, and S. Portilla. 2014. Growth, antioxidant capacity and total carotene of *Dunaliella salina* DCCBC15 in a low cost enriched natural seawater medium. *World Journal of Microbiology and Biotechnology* 30:317–322. doi:10.1007/s11274-013-1413-2.
- Uduman, N., Y. Qi, M. K. Danquah, G. M. Forde, and A. Hoadley. 2010. Dewatering of microalgal cultures: A major bottleneck to algae-based fuels. *Journal of Renewable and Sustainable Energy* 2:1–15. doi:10.1063/1.3294480.
- Ugwu, C. U., H. Aoyagi, and H. Uchiyama. 2008. Photobioreactors for mass cultivation of algae. *Bioresource Technology* 99:4021–4028. doi:10.1016/j.biortech.2007.01.046.
- Utomo, R. P., Y. R. Chang, D. J. Lee, and J. S. Chang. 2013. Lutein recovery from *Chlorella* sp. ESP-6 with coagulants. *Bioresource Technology* 139:176–180. doi:10.1016/j.biortech.2013.04.025.
- Varela, J. C., H. Pereira, M. Vila, and R. Leon. 2015. Production of carotenoids by microalgae: Achievements and challenges. *Photosynthesis Research* 125:423–436. doi:10.1007/s11220-015-0149-2.
- Vidoudez, C., and G. Pohnert. 2012. Comparative metabolomics of the diatom *Skeletonema marinoi* in different growth phases. *Metabolomics* 8:654–669. doi:10.1007/s11306-011-0356-6.
- Vigani, M., C. Parisi, E. Rodríguez-Cerezo, M. J. Barbosa, L. Sijsma, M. Ploeg, and C. Enzing. 2015. Food and feed products from microalgae: Market opportunities and challenges for the EU. *Trends in Food Science and Technology* 42:81–92. doi:10.1016/j.tifs.2014.12.004.
- Vijayavel, K., C. Anbuselvam, and M. P. Balasubramanian. 2007. Antioxidant effect of the marine algae *Chlorella vulgaris* against naphthalene-induced oxidative stress in the albinorats. *Molecular and Cellular Biochemistry* 303:39–44. doi:10.1007/s11010-007-9453-2.
- Vilchez, C., E. Forjan, M. Cuaresma, F. Bedmar, I. Garbayo, and J. M. Vega. 2011. Marine carotenoids: biological functions and commercial applications. *Marine Drugs* 9:319–333. doi:10.3390/md9030319.
- Wang, H. M. D., C. C. Chen, P. Huynh, and J. S. Chang. 2015. Exploring the potential of using algae in cosmetics. *Bioresource Technology* 184:355–362. doi:10.1016/j.biortech.2014.12.001.
- Wang, J., D. Han, M. R. Sommerfeld, C. Lu, and Q. Hu. 2013. Effect of initial biomass density on growth and astaxanthin production of *Haematococcus pluvialis* in an outdoor photobioreactor. *Journal of Applied Phycology* 25:253–260. doi:10.1007/s10811-012-9859-4.
- Weber, D., and T. Grune. 2012. The contribution of β -carotene to vitamin A supply of humans. *Molecular Nutrition and Food Research* 56:251–258. doi:10.1002/mnfr.201100230.
- Wei, D., F. Chen, G. Chen, X. W. Zhang, L. J. Liu, and H. Zhang. 2008. Enhanced production of lutein in heterotrophic *Chlorella protothecoides* by oxidative stress. *Science China Life Sciences* 51:1088–1093. doi:10.1007/s11427-008-0145-2.
- Wells, M. L., P. Potin, J. S. Craigie, J. A. Raven, S. S. Merchant, K. E. Hellwell, A. G. Smith, M. E. Camire, and S. H. Brawley. 2016. Algae as nutritional and functional food sources: revisiting our understanding. *Journal of Applied Phycology* 29:949–982. doi:10.1007/s10811-016-0974-5.
- Wichuk, K., S. Brynjolfsson, and W. Fu. 2014. Biotechnological production of value-added carotenoids from microalgae. *Bioengineered* 5:204–208. doi:10.4161/bioe.28720.
- Wong, Y. K. 2016. Feasibility of using *Chlorella vulgaris* for the production of algal lipids, for advancement towards a potential application in the manufacture of commodity chemicals and the treatment of wastewater. *Theses and Dissertations* pp. 254.
- Wu, Y. H., X. Li, Y. Yu, H. Y. Hu, T. Y. Zhang, and F. M. Li. 2013. An integrated micro algal growth model and its application to optimize the biomass production of *Scenedesmus* sp. LX1 in open pond under the nutrient level of domestic secondary effluent. *Bioresource Technol* 144:445–451. doi:10.1016/j.biortech.2013.06.065.
- Wu, Z., S. Wu, and X. Shi. 2007. Supercritical fluid extraction and determination of lutein in heterotrophically cultivated *Chlorella pyrenoidosa*. *Journal of Food Process Engineering* 30:174–185. doi:10.1111/j.1745-4530.2007.00102.x.
- Xia, S., K. Wang, L. Wan, A. Li, Q. Hu, and C. Zhang. 2013. Production, characterization, and antioxidant activity of fucoxanthin from the marine diatom *Odontella aurita*. *Marine Drugs* 11:2667–2681. doi:10.3390/md11072667.
- Yaakob, Z., E. Ali, A. Zainal, M. Mohamad, and M. S. Takriff. 2014a. An overview: biomolecules from microalgae for animal feed and aquaculture. *Journal of Biological Research-Thessaloniki* 21:1–6. doi:10.1186/2241-5793-21-6.
- Yaakob, Z., K. F. Kamarudin, R. Rajkumar, M. S. Takriff, and S. N. Badar. 2014b. The current methods for the biomass production of the microalgae from wastewaters: An overview. *World Applied Sciences Journal* 31:1744–1758.
- Yamaguchi, K. 1996. Recent advances in micro algal bioscience in Japan, with special reference to utilization of biomass and metabolites: a review. *Journal of Applied Phycology* 8:487–502. doi:10.1007/BF02186327.

- Yamashita, E. 2013. Astaxanthin as a medical food. *Functional Foods in Health and Disease* 3:254–258.
- Yang, D. J., J. T. Lin, Y. C. Chen, F. J. Lu, T. J. Chang, T. J. Wang, H. W. Lin, and Y. Y. Chang. 2013. Suppressive effect of carotenoid extract of *Dunaliella salina* alga on production of LPS-stimulated pro-inflammatory mediators in RAW264.7 cells via NF- κ B and JNK inactivation. *Journal of Functional Foods* 5:607–615. doi:10.1016/j.jff.2013.01.001.
- Yang, Y., B. Kim, and J. Y. Lee. 2013. Astaxanthin structure, metabolism, and health benefits. *Journal of Human Nutrition and Food Science* 1:1003.
- Yen, H. W., S. C. Yang, C. H. Chen, and J. S. Chang. 2015. Supercritical fluid extraction of valuable compounds from microalgal biomass. *Bioresource Technology* 184:291–296. doi:10.1016/j.biortech.2014.10.030.
- Yuan, J. P., F. Chen, X. Liu, and X. Z. Li. 2002. Carotenoid composition in the green microalga *Chlorococcum*. *Food Chemistry* 76:319–325. doi:10.1016/S0308-8146(01)00279-5.
- Yun, H., I. Kim, S. H. Kwon, J. S. Kang, and A. S. Om. 2011. Protective effect of *Chlorella vulgaris* against lead-induced oxidative stress in rat brains. *Journal of Health Sciences* 57:245–254. doi:10.1248/jhs.57.245.
- Zhang, D. H. and Y. K. Lee. 1997. Enhanced accumulation of secondary carotenoids in a mutant of the green alga, *Chlorococcum* sp. *Journal of Applied Phycology* 9:459–463. doi:10.1023/A:1007902103419.
- Zhang, D. H., Y. K. Ng, and S. M. Phang. 1997. Composition and accumulation of secondary carotenoids in *Chlorococcum* sp. *Journal of Applied Phycology* 9:147–155. doi:10.1023/A:1007926528388.
- Zhang, J., Z. Sun, P. Sun, T. Chen, and F. Chen. 2014. Microalgal carotenoids: Beneficial effects and potential in human health. *Food and Function* 5:413–425. doi:10.1039/c3fo60607d.
- Zhang, P., Z. Li, L. Lu, Y. Xiao, J. Liu, J. Guo, and F. Fang. 2017. Effects of stepwise nitrogen depletion on carotenoid content, fluorescence parameters and the cellular stoichiometry of *Chlorella vulgaris*. *Spectrochimica Acta, Part A: Molecular and Biomolecular Spectroscopy* 181:30–38. doi:10.1016/j.saa.2017.03.022.
- Zhang, Z., J. J. Huang, D. Sun, Y. Lee, and F. Chen. 2017. Two-step cultivation for production of astaxanthin in *Chlorella zofingiensis* using a patented energy-free rotating floating photobioreactor (RFP). *Bioresource Technology* 224:515–522. doi:10.1016/j.biortech.2016.10.081.
- Zheng, H., J. Yin, Z. Gao, H. Huang, X. Ji, and C. Dou. 2011. Disruption of *Chlorella vulgaris* cells for the release of biodiesel producing lipids: A comparison of grinding, ultrasonication, bead milling, enzymatic lysis, and microwaves. *Applied Biochemistry and Biotechnology* 164:1215–1224. doi:10.1007/s12010-011-9207-1.
- Zhou, Y., T. Wang, Q. Meng, and S. Zhai. 2016. Association of carotenoids with risk of gastric cancer: A meta-analysis. *Clinical Nutrition* 35:109–116. doi:10.1016/j.clnu.2015.02.003.