

Bounds of Validity for Bifurcations of Equilibria in a Class of Networked Dynamical Systems

Pranav Gupta¹, Ravi Banavar², and Anastasia Bizyaeva³

Abstract—Local bifurcation analysis plays a central role in understanding qualitative transitions in networked nonlinear dynamical systems, including dynamic neural network and opinion dynamics models. In this article we establish explicit bounds of validity for the classification of bifurcation diagrams in two classes of continuous-time networked dynamical systems, analogous in structure to the Hopfield and the Firing Rate dynamic neural network models. Our approach leverages recent advances in computing the bounds for the validity of Lyapunov–Schmidt reduction, a reduction method widely employed in nonlinear systems analysis. Using these bounds we rigorously characterize neighborhoods around bifurcation points where predictions from reduced-order models remain reliable. We further demonstrate how these bounds can be applied to an illustrative family of nonlinear opinion dynamics on k -regular graphs, which emerges as a special case of the general framework. These results provide new analytical tools for quantifying the robustness of bifurcation phenomena in dynamics over networked systems and highlight the interplay between network structure and nonlinear dynamical behavior.

I. INTRODUCTION

Local bifurcations are points of critical transitions in behavior of nonlinear dynamical systems that occur when a parameter crosses a threshold, causing the stability of an equilibrium point to change and new features such as new equilibria to emerge. Bifurcations are a common route to multistability in complex systems, and bifurcation analysis is used to characterize the emergence, stability, and basins of attraction of coexisting equilibria as parameters vary.

In this paper we study the robustness of bifurcation diagrams in two broad classes of nonlinear networked dynamical systems, analogous in their structure to Hopfield and firing rate dynamic neural network models. Systems of this form are commonly encountered as models of dynamic neural networks [1]–[4], opinion dynamics and decision-making [5]–[10], Lur’e feedback systems [11]–[13], gene regulatory networks [14]–[16], and other complex systems across application domains. Multistable equilibria are common desirable features in these models, representing meaningful states such as stored associative memories in a neural network or collective decision states in a group of communicating agents. Lyapunov–Schmidt reduction is a nonlinear analysis

technique commonly used to analyze bifurcation diagrams in nonlinear networks to classify such multistabilities [10], [17]. In this work we will bound the regions of validity for the conclusions of such bifurcation analysis.

In our prior work [18], we developed estimates for the domain of validity of reduced-order models derived through the Lyapunov–Schmidt reduction framework. These results built directly upon the theoretical foundation established in [19], where explicit bounds of validity were obtained for the application of Implicit Function theorem using tools from functional analysis. In the present work, we build on these results by specializing the general bounds to rigorously bound the validity region of reduced-order models in the context of bifurcations of equilibria in networked dynamical systems. Specifically, we apply this methodology to Hopfield and Firing Rate network models, relating the regions of validity of Lyapunov–Schmidt reduction-based analysis to network structure and choice of smooth nonlinear activation functions in these models.

Our contributions are the following. First, in Section III we present a simplified general bound on the regions of validity for Lyapunov–Schmidt reduction in an arbitrary nonlinear dynamical system, building on our previous work in [18]. Next, in Section IV we specialize this bound for the Hopfield and Firing Rate network models, presenting simplified computable expressions that leverage the structure of the model equations. Finally, in Section V we apply the derived bounds to consensus bifurcations in nonlinear opinion dynamics networks over regular graphs. We show that the resulting bounds involve interpretable quantities related to the structure of the underlying graph.

The remainder of the paper is organized as follows. In section II we introduce notation, relevant lemmas, and key background results on explicit bounds of validity for the Lyapunov–Schmidt reduction. In section III we derive a simplified statement of the general Lyapunov–Schmidt reduction bounds. In section IV, we specialize these bounds for two classes of continuous-time dynamic network models. In section V we apply the derived network bounds to consensus bifurcations in nonlinear opinion dynamics over k -regular graphs. Finally, in section VI we provide a brief discussion.

II. BACKGROUND

A. Notations and Lemmas

Let $\mathbb{R}$ denote the set of real numbers and $\mathbb{N}$ the set of positive integers. A mapping $f : U \rightarrow V$, where $U \subseteq \mathbb{R}^m$ and $V \subseteq \mathbb{R}^n$, is said to be of class C^ν for $\nu \in \mathbb{N}$ if it is ν -times continuously differentiable. For $f \in C^\nu$, we write

¹ P.G. is with the Centre for Systems and Control, Indian Institute of Technology Bombay, Mumbai - 400076, Maharashtra, India and with the Sibley School of Mechanical and Aerospace Engineering, Cornell University, Ithaca, NY, USA, 14850 guptapranav@iitb.ac.in

² R.B. is with the Centre for Systems and Control, Indian Institute of Technology Bombay, Mumbai - 400076, Maharashtra, India banavar@iitb.ac.in

³ A.B. is with the Sibley School of Mechanical and Aerospace Engineering, Cornell University, Ithaca, NY, USA, 14850, anastasiab@cornell.edu

$D_y f(y)$ for the Jacobian matrix of partial derivatives with respect to the variables y . For $y_0 \in \mathbb{R}^n$ and $r > 0$, the open ball of radius r centered at y_0 is denoted by $\mathcal{B}(y_0, r) := \{y \in \mathbb{R}^n : \|y - y_0\| < r\}$, where $\|\cdot\|$ is a vector norm on $\mathbb{R}^n$. For $A \in \mathbb{R}^{p \times q}$, we use $\|A\|$ to denote the spectral matrix norm induced by this vector norm. We write $\mathbf{0}_n, \mathbf{1}_n \in \mathbb{R}^n$ for the all-zero and all-one column vectors, respectively, and $\mathbf{0}_{p \times q}, \mathbf{1}_{p \times q} \in \mathbb{R}^{p \times q}$ for the corresponding matrices. For a matrix M , we denote by $\sigma_i(M) = \lambda_i(MM^T) = \lambda_i(M^T M)$ its i -th largest singular value and by $\sigma_{\min}(M)$ and $\sigma_{\max}(M)$ its smallest nonzero and largest singular values respectively. We let $\rho(M)$ denote the spectral radius of M .

Let $\mathcal{G} = (\mathcal{V}, \mathcal{E})$ denote a graph with vertex set $\mathcal{V}$ of cardinality $|\mathcal{V}| = n$ and edge set $\mathcal{E} \subseteq \mathcal{V} \times \mathcal{V}$. We associate with $\mathcal{G}$ the signed adjacency matrix $A \in \mathbb{R}^{n \times n}$, where the entry a_{ij} encodes the signed weight of the edge between nodes i and j , and $a_{ij} = 0$ if no such edge exists. A graph is said to be k -regular if every vertex has exactly k neighbors, i.e., the degree of each vertex is equal to k . Equivalently, the adjacency matrix A of a k -regular graph satisfies $A\mathbf{1}_n = k\mathbf{1}_n$.

Next we state two useful lemmas.

Lemma 2.1: Let $W \in \mathbb{R}^{n \times k}$ be a matrix with orthonormal columns, and let $P = WW^T$ denote the associated orthogonal projection onto the column space of W . Then, for any matrix $Y \in \mathbb{R}^{n \times m}$, $\|W^T Y\| = \|PY\|$.

Proof: By definition of spectral norm and properties of orthogonal projection matrices, $\|W^T Y\|^2 = \rho(Y^T WW^T Y) = \rho(Y^T PY) = \rho(Y^T P^2 Y) = \|PY\|^2$. ■

Lemma 2.2: Let $W \in \mathbb{R}^{n \times k}$ be a matrix with orthonormal columns, and let $P = WW^T$ denote the corresponding orthogonal projection. Then, for any matrix $M \in \mathbb{R}^{n \times n}$, $\sigma_i(W^T MW) = \sigma_i(PMP)$, $i = 1, \dots, k$.

Proof: By definition of singular value and orthogonal projections, $\sigma_i(W^T MW) = \lambda_i(W^T M^T WW^T MW) = \lambda_i(W^T M^T P MW) = \lambda_i(W^T M^T P P MW) = \sigma_i(PMW) = \lambda_i(PMWW^T M^T P) = \lambda_i(PMPM^T P) = \lambda_i(PMPPM^T P)V = \sigma_i(PMP)$. ■

B. Lyapunov-Schmidt Reduction

Lyapunov-Schmidt reduction is a method used to analyze parametrized nonlinear dynamical systems near their singular equilibria, leveraging the Implicit Function theorem. The method decomposes the state space of a dynamical system into the null space of its Jacobian and its orthogonal complement at the singular point. The Implicit Function theorem is then used to eliminate the component orthogonal to the null space in the local bifurcation diagram of the system, effectively projecting the high-dimensional equations describing the equilibria of the model onto the kernel [20, Ch. VII]. The resulting reduced equations capture the structure of the local bifurcation diagram of the original system. We now present a summary of the reduction procedure.

Let $X \subset \mathbb{R}^n$ and $\Lambda \subset \mathbb{R}^m$ be open sets. Consider the parametrized nonlinear dynamical system $\dot{x} = \Phi(x, p)$ where $x \in X$ is the state vector, $p \in \Lambda$ is the parameter vector, and $\Phi : X \times \Lambda \rightarrow \mathbb{R}^n$ is at least $\mathcal{C}^2$ and has a bifurcation at

$(x^*, p^*) \in X \times \Lambda$, i.e. $\Phi(x^*, p^*) = 0$ and $J := D_x \Phi(x^*, p^*)$ is singular (non-invertible) with $q := \dim(\ker J) > 0$.

Define orthonormal basis matrices $V \in \mathbb{R}^{n \times q}$ and $W \in \mathbb{R}^{n \times (n-q)}$ whose column vectors span $\ker(J)$ and $\ker(J)^\perp$ respectively. Explicitly,

$$V = [v_1 \ \dots \ v_q], \quad W = [w_{q+1} \ \dots \ w_n]. \quad (1)$$

By construction, $\forall i \in \{q+1, \dots, n\}$, $k \in \{1, \dots, q\}$ we have $w_i^\top v_k = 0$. Moreover, the projections onto $\ker(J)$ and $\ker(J)^\perp$ can be expressed as

$$P_\parallel = VV^\top, \quad P_\perp = WW^\top.$$

We denote the direct sum splitting of $\mathbb{R}^n$ into $\ker(J) \oplus \ker(J)^\perp$ by

$$x \mapsto (x^\parallel, x^\perp) := (P_\parallel x, P_\perp x) \in \ker(J) \times \ker(J)^\perp.$$

Define coordinate map $\mathbb{R}^q \times \mathbb{R}^{n-q} \ni (\alpha, \beta) \mapsto \Gamma(\alpha, \beta) \in X$:

$$x = \Gamma(\alpha, \beta) := \underbrace{V\alpha}_{x^\parallel} + \underbrace{W\beta}_{x^\perp} \quad \text{where} \quad \begin{cases} \alpha = V^\top x \in \mathbb{R}^q \\ \beta = W^\top x \in \mathbb{R}^{n-q} \end{cases} \quad (2)$$

where α and β represent the reduced coordinates of $x^\parallel$ and $x^\perp$, corresponding to components in $\ker(J)$ and its orthogonal complement $\ker(J)^\perp$, respectively. We employ this coordinate map as a shorthand throughout the analysis.

The Lyapunov-Schmidt reduction involves applying the Implicit Function theorem to obtain a locally valid implicit mapping $(\alpha, p) \mapsto \beta$. In particular, it guarantees the existence of open neighborhoods $\mathcal{N}((\alpha_0, p^*)) \subset \mathbb{R}^q \times \Lambda$ and $\mathcal{N}(\beta_0) \subset \mathbb{R}^{n-q}$ equipped with implicit map $\varphi : \mathcal{N}((\alpha_0, p^*)) \rightarrow \mathcal{N}(\beta_0)$ that expresses coordinates along the range of the Jacobian as a function of the coordinates along the null space and the bifurcation parameter, $\beta = \varphi(\alpha, p)$, satisfying

$$\Phi(V\alpha + W\varphi(\alpha, p), p) = 0 \quad (3)$$

for all $(\alpha, p) \in \mathcal{N}((\alpha_0, p^*))$.

C. Bounds of Validity for Lyapunov-Schmidt Reduction

A key limitation of Lyapunov-Schmidt reduction is that the equivalence between the reduced bifurcation equations and the equilibria of the original high-dimensional system holds only locally. Consequently, the analysis of bifurcations leveraging this reduction method is valid only in a small neighborhood of the singular point. Understanding the extent of these neighborhoods is crucial for evaluating the robustness of conclusions drawn from analysis of the reduced equations. To address this limitation, in previous work we derived a general bound of validity for Lyapunov-Schmidt reduction in systems of ordinary differential equations [18]. In this section we summarize our recent result.

In the context of Lyapunov-Schmidt reduction, we obtain computable lower bound approximations for the neighborhoods $\mathcal{N}((\alpha_0, p^*))$, $\mathcal{N}(\beta_0)$ as open balls $\mathcal{B}((\alpha_0, p^*), r_\parallel)$ and $\mathcal{B}(\beta_0, r_\perp)$. The radii $r_\parallel$ and $r_\perp$ bound the regions around the singular point $(\alpha, \beta, p) = (\alpha_0, \beta_0, p^*)$ where the reduced and full bifurcation diagrams coincide topologically.

In particular $r_{\parallel}$ defines the extent of equivalence along the critical subspace, i.e. the product between $\ker(J)$ and the parameter space Λ . Analogously, $r_{\perp}$ defines the extent of equivalence along $\ker(J)^{\perp}$. The algorithm to explicitly determine the bounding radii $r_{\parallel}$ and $r_{\perp}$ is described below.

Before we state the theorem on bounds, we define the following quantities. First, we define two scalar quantities that will appear in the statement of the bounds,

$$M_{\parallel} := \left\| \begin{bmatrix} \mathbf{0}_{(n-q) \times q} & W^{\top} D_p \Phi(x^*, p^*) \end{bmatrix} \right\|, \quad (4a)$$

$$M_{\perp} := \left\| (W^{\top} D_x \Phi(x^*, p^*) W)^{-1} \right\|. \quad (4b)$$

Next, we define the maps $\xi_{\parallel} : \mathbb{R}^q \times \Lambda \rightarrow \mathbb{R}^{(n-q) \times (q+m)}$ and $\xi_{\perp} : \mathbb{R}^q \times \mathbb{R}^{n-q} \times \Lambda \rightarrow \mathbb{R}^{(n-q) \times (n-q)}$ as

$$\xi_{\parallel}(\alpha, p) := W^{\top} \begin{bmatrix} \xi_{\parallel\alpha}(\alpha, p) & \xi_{\parallel p}(\alpha, p) \end{bmatrix}, \quad (5a)$$

$$\xi_{\perp}(\alpha, \beta, p) := W^{\top} \left(D_x \Phi(\Gamma(\alpha, \beta), p) - D_x \Phi(x^*, p^*) \right) W, \quad (5b)$$

where $\xi_{\parallel p}(\alpha, p) = D_p \Phi(\Gamma(\alpha, \beta_0), p) - D_p \Phi(x^*, p^*)$ and $\xi_{\parallel\alpha}(\alpha, p) = D_x \Phi(\Gamma(\alpha, \beta_0), p) V$. The maps (5) quantify first-order variations of the Jacobian $D_x \Phi$ and parameter derivative $D_p \Phi$ along the critical subspace and its orthogonal complement. We use $\mathcal{B}_{\parallel}(r_{\parallel}) := \mathcal{B}((\alpha_0, p^*), r_{\parallel}) \subset \mathbb{R}^q \times \Lambda$ and $\mathcal{B}_{\perp}(r_{\perp}) := \mathcal{B}(\beta_0, r_{\perp}) \subset \mathbb{R}^{n-q}$ as shorthand notation to denote open balls centered at the critical point. Finally, we define two scalar quantities as the maximum norm of $\xi_{\parallel}(\alpha, p), \xi_{\perp}(\alpha, \beta, p)$ over open balls centered around the critical point, and state the main result from [18]

$$L_{\parallel}(r_{\parallel}) := \sup \{ \|\xi_{\parallel}(\alpha, p)\| : (\alpha, p) \in \mathcal{B}_{\parallel}(r_{\parallel}) \}, \quad (6a)$$

$$L_{\perp}(r_{\parallel}, r_{\perp}) := \sup \{ \|\xi_{\perp}(\alpha, \beta, p)\| : (\alpha, p) \in \mathcal{B}_{\parallel}(r_{\parallel}) \text{ and } \beta \in \mathcal{B}_{\perp}(r_{\perp}) \}. \quad (6b)$$

Proposition 2.3: [18, Theorem 3.2] For all $r_{\parallel}, r_{\perp} > 0$ simultaneously satisfying

$$L_{\parallel}(r_{\parallel})r_{\parallel} + L_{\perp}(r_{\parallel}, r_{\perp})r_{\perp} < \frac{r_{\perp}}{M_{\perp}} - M_{\parallel}r_{\parallel} \quad (7a)$$

$$M_{\perp}L_{\perp}(r_{\parallel}, r_{\perp}) < 1 \quad (7b)$$

there exists a smooth implicit map $\varphi : \mathcal{B}_{\parallel}(r_{\parallel}) \rightarrow \mathcal{B}_{\perp}(r_{\perp})$ that satisfies (3), where the relevant quantities in (7) are defined by the equations (4a), (4b), (6a), (6b).

III. SIMPLIFIED GENERAL LS BOUNDS

Before stating our main results in the next section, we present some results simplifying the statement of the general bounds of Proposition 2.3. We first express the projections onto $\ker(J^*)$ and $\ker(J^*)^{\perp}$ as

$$P_{\parallel} = VV^{\top}, \quad P_{\perp} = WW^{\top} = I - P_{\parallel}.$$

Next we streamline the expressions (4a), (4b) and express these quantities in terms of projections of the Jacobian matrices of Φ at the bifurcation point (x^*, p^*) .

Lemma 3.1: $M_{\parallel} = \|P_{\perp} D_p \Phi(x^*, p^*)\|$

Proof: From (4a) and definition of the spectral norm,

$$\begin{aligned} M_{\parallel}^2 &= \rho \left(\begin{bmatrix} \mathbf{0}_{n-q} & W^{\top} D_p \Phi(x^*, p^*) \end{bmatrix}^{\top} \begin{bmatrix} \mathbf{0}_{n-q} & W^{\top} D_p \Phi(x^*, p^*) \end{bmatrix} \right) \\ &= \rho \left((D_p \Phi(x^*, p^*))^{\top} W W^{\top} D_p \Phi(x^*, p^*) \right) \\ &= \|W^{\top} D_p \Phi(x^*, p^*)\|^2. \end{aligned}$$

Taking the square root on both sides and applying Lemma 2.1 completes the proof. $\blacksquare$

Lemma 3.2: $M_{\perp} = 1/\sigma_{\min}(P_{\perp} D_x \Phi(x^*, p^*) P_{\perp})$.

Proof: From (4b) and definition of spectral norm,

$$\begin{aligned} M_{\perp} &= \sigma_{\max} \left((W^{\top} D_x \Phi(x^*, p^*) W)^{-1} \right) \\ &= \left(\sigma_{\min} (W^{\top} D_x \Phi(x^*, p^*) W) \right)^{-1} \end{aligned}$$

where the last step followed from [21, Theorem 3.3]. Then the lemma statement is a consequence of Lemma 2.2. $\blacksquare$

Next, we prove that the second inequality condition in the originally derived Proposition 2.3 is redundant, and state a simplified version of the bounds.

Theorem 3.3: For all $r_{\parallel}, r_{\perp} > 0$ satisfying

$$L_{\parallel}(r_{\parallel})r_{\parallel} + L_{\perp}(r_{\parallel}, r_{\perp})r_{\perp} < \frac{r_{\perp}}{M_{\perp}} - M_{\parallel}r_{\parallel} \quad (8)$$

there exists a smooth implicit map $\varphi : \mathcal{B}_{\parallel}(r_{\parallel}) \rightarrow \mathcal{B}_{\perp}(r_{\perp})$ that satisfies (3), where the quantities in (8) are defined by Lemma 3.1, Lemma 3.2, and equations (6a), (6b).

Proof: Suppose (7a) holds. Rearranging terms in this inequality, we can equivalently state that

$$M_{\perp}L_{\perp}(r_{\parallel}, r_{\perp}) + M_{\perp}(M_{\parallel} + L_{\parallel}(r_{\parallel})) \frac{r_{\parallel}}{r_{\perp}} < 1. \quad (9)$$

Observe that all the quantities on the left hand side of (9) are nonnegative scalars. Therefore by monotonicity of addition, it must necessarily hold that $M_{\perp}L_{\perp}(r_{\parallel}, r_{\perp}) < 1$, i.e. (7b) is satisfied. The Theorem statement then follows directly from Proposition 2.3. $\blacksquare$

IV. MAIN RESULTS: NETWORK BOUNDS

In this section we study the robustness of bifurcation diagrams for two general classes of networked nonlinear dynamical systems. First we consider systems of the form

$$\dot{x} = \Phi(x, p) := -Cx + pAS(x) + b \quad (10)$$

where $x \in \mathbb{R}^n$ is the state vector, $p \in \mathbb{R}$ is a parameter, $b \in \mathbb{R}^n$ is a constant input or bias, $A \in \mathbb{R}^{n \times n}$, and $C = \text{diag}(c_1, \dots, c_n)$ with $c_i \in \mathbb{R}^{>0}$. The function $S(z) = [S_1(z_1) \dots S_n(z_n)]^{\top}$ applies an element-wise smooth activation function to each input, and its derivative is denoted as $D_z S(z) = \text{diag}(S'_1(z_1), \dots, S'_n(z_n))$. We will refer to (10) as a *Hopfield* or *voltage* model as it follows the structure of classic and modern associative memory networks [1], [2]. Second we will consider systems of the form

$$\dot{x} = \Psi(x, p) := -Cx + S(pAx + b) \quad (11)$$

with all the quantities defined the same as for the Hopfield models. We will refer to (11) as a *firing rate* model [3], [4].

It has been shown that (10), (11) are related through changes of coordinates and time-varying input transformations [22].

In this section we establish computable bounds of validity for the equivalence of bifurcation diagrams of (10), (11) with their reduced bifurcation equations obtained via Lyapunov-Schmidt reduction at singular points, specializing Theorem 3.3. We seek to state these bounds in terms of interpretable quantities such as the network connectivity matrix A , the activation function S , and its derivatives.

A. Hopfield models

First we consider (10). Computing derivatives of the model with respect to x and p yields

$$D_x \Phi(x, p) = -C + pAD_z S(x), \quad (12a)$$

$$D_p \Phi(x, p) = AS(x). \quad (12b)$$

We define the Jacobian J_H^* at the singular point as

$$J_H^* := D_x \Phi(x^*, p^*) = -C + p^*AD_z S(x^*)$$

Next, we make the following assumptions:

Assumption 1: There exists $(x^*, p^*) \in \mathbb{R}^n \times \mathbb{R}^{>0}$ such that the following conditions hold:

- (i) The pair (x^*, p^*) is an equilibrium of (10), i.e.,

$$\Phi(x^*, p^*) = -Cx^* + p^*AS(x^*) + b = 0. \quad (13)$$

- (ii) The Jacobian J_H^* has a zero eigenvalue with algebraic and geometric multiplicity q and corresponding orthonormal right null eigenvectors $V = [v_1, \dots, v_q]$, orthonormal left null eigenvectors $W_0 = [w_1, \dots, w_q]$, and an orthonormal basis $W = [w_{q+1}, \dots, w_n]$ spanning $\ker(J_H^*)^\perp \cong \text{range}(J_H^*)$. Define the complementary orthogonal projections $P_\parallel = VV^\top$ and $P_\perp = WW^\top = I - W_0W_0^\top$. $\text{Re}(\lambda_j) \neq 0$ for all other eigenvalues.

To apply Theorem 3.3 to (10), we must compute the various quantities in the inequality condition (8). In the following sequence of Lemmas we derive expressions needed to define $L_\parallel(r_\parallel)$, $L_\perp(r_\parallel, r_\perp)$ in (8), exploiting the form of Hopfield dynamics to express these quantities in terms of the network matrix A , activation function S , and its first derivatives.

Lemma 4.1: $\|\xi_\parallel(\alpha, p)\| = \|P_\perp A [\Delta_\parallel(\alpha, p)V \quad \bar{S}_\parallel(\alpha)]\|$ where $\Delta_\parallel(\alpha, p)$ and $\bar{S}_\parallel(\alpha)$ are defined as

$$\Delta_\parallel(\alpha, p) := pD_z S(\Gamma(\alpha, \beta_0)) - p^*D_z S(x^*) \quad (14a)$$

$$\bar{S}_\parallel(\alpha) := S(\Gamma(\alpha, \beta_0)) - S(x^*) \quad (14b)$$

Proof: Following the definition (5a), we must compute $\xi_{\parallel\alpha}(\alpha, p)$ and $\xi_{\parallel p}(\alpha, p)$. Since $J_H^*V = 0$, we can write

$$(-C + p^*AD_z S(x^*))V = 0 \implies CV = p^*AD_z S(x^*)V,$$

which allows us to simplify $\xi_{\parallel\alpha}(\alpha, p)$ as

$$\begin{aligned} \xi_{\parallel\alpha}(\alpha, p) &= D_x \Phi(\Gamma(\alpha, \beta_0), p)V \\ &= [-C + pAD_z S(\Gamma(\alpha, \beta_0))]V \\ &= [-p^*AD_z S(x^*) + pAD_z S(\Gamma(\alpha, \beta_0))]V \\ &= A\Delta_\parallel(\alpha, p)V \end{aligned}$$

Next, we simplify $\xi_{\parallel p}(\alpha, p)$ as

$$\begin{aligned} \xi_{\parallel p}(\alpha, p) &:= D_p \Phi(\Gamma(\alpha, \beta_0), p) - D_p \Phi(x^*, p^*) \\ &= A(S(\Gamma(\alpha, \beta_0)) - S(x^*)) = A\bar{S}_\parallel(\alpha). \end{aligned}$$

Finally, combining our expressions for $\xi_{\parallel\alpha}$ and $\xi_{\parallel p}$ yields

$$\xi_\parallel(\alpha, p) = W^\top A [\Delta_\parallel(\alpha, p)V \quad \bar{S}_\parallel(\alpha)].$$

The desired expression for $\|\xi_\parallel(\alpha, p)\|$ follows directly by applying Lemma 2.1 with $Y = A [\Delta_\parallel(\alpha, p)V \quad \bar{S}_\parallel(\alpha)]$. ■
Lemma 4.2: $\|\xi_\perp(\alpha, \beta, p)\| = \|P_\perp A \Delta_\perp(\alpha, \beta, p)P_\perp\|$ where

$$\Delta_\perp(\alpha, \beta, p) := pD_z S(\Gamma(\alpha, \beta)) - p^*D_z S(x^*). \quad (15)$$

Proof: Building from its definition (5b), we have

$$\begin{aligned} \xi_\perp(\alpha, \beta, p) &= W^\top (D_x \Phi(\Gamma(\alpha, \beta), p) - D_x(x^*, p^*))W \\ &= W^\top (pAD_z S(\Gamma(\alpha, \beta)) - p^*AD_z S(x^*))W \\ &= W^\top A \Delta_\perp(\alpha, \beta, p)W. \end{aligned}$$

Then the lemma statement follows by the definition of spectral norm and Lemma 2.2. ■

With these lemmas established, we state the first result.

Theorem 4.3 (Bounds of bifurcation in Hopfield models):

Consider (10) under Assumption 1. For all $r_\parallel, r_\perp > 0$ satisfying

$$\begin{aligned} L_\parallel(r_\parallel)r_\parallel + L_\perp(r_\parallel, r_\perp)r_\perp \\ < \sigma_{\min}(P_\perp J_H^* P_\perp)r_\perp - \frac{1}{p^*}\|P_\perp(Cx^* - b)\|r_\parallel \end{aligned} \quad (16)$$

where the quantities $L_\parallel(r_\parallel)$, $L_\perp(r_\parallel, r_\perp)$ are defined as

$$\begin{aligned} L_\parallel(r_\parallel) &= \sup \left\{ \|P_\perp A [\Delta_\parallel(\alpha, p)V \quad \bar{S}_\parallel(\alpha)]\| \right. \\ &\quad \left. : (\alpha, p) \in \mathcal{B}((\alpha_0, p^*), r_\parallel) \right\}, \end{aligned}$$

$$\begin{aligned} L_\perp(r_\parallel, r_\perp) &= \sup \left\{ \|P_\perp A \Delta_\perp(\alpha, \beta, p)P_\perp\| \right. \\ &\quad \left. : (\alpha, p) \in \mathcal{B}((\alpha_0, p^*), r_\parallel), \beta \in \mathcal{B}(\beta_0, r_\perp) \right\}, \end{aligned}$$

with $\Delta_\parallel, \Delta_\perp, \bar{S}_\parallel$ defined in (14), (15), there exists a smooth implicit map $\varphi : \mathcal{B}((\alpha_0, p^*), r_\parallel) \rightarrow \mathcal{B}(\beta_0, r_\perp)$ that satisfies $\Phi(V\alpha + W\varphi(\alpha, p), p) = 0$.

Proof: From the equilibrium condition (13) we see that $AS(x^*) = \frac{1}{p^*}(Cx^* - b)$. The statement then immediately follows from applying Lemma 3.1 to obtain

$$M_\parallel = \|P_\perp AS(x^*)\| = \left\| \frac{1}{p^*}P_\perp(Cx^* - b) \right\|. \quad (18)$$

Substituting the values of (4b), (6a), (6b) from Lemma 3.2, Lemma 4.1, Lemma 4.2 into (8) from Theorem 3.3 yields the theorem statement. ■

B. Firing Rate models

Next, we repeat the presented analysis for (11). The derivatives of (11) with respect to x and p are

$$D_x \Psi(x, p) = -C + pD_z S(pAx + b)A \quad (19a)$$

$$D_p \Psi(x, p) = D_z S(pAx + b)Ax \quad (19b)$$

and the Jacobian J_F^* at the singular point evaluates to

$$J_F^* := D_x \Psi(x^*, p^*) = -C + p^* D_z S(p^* A x^* + b) A.$$

We make the following assumption:

Assumption 2: There exists $(x^*, p^*) \in \mathbb{R}^n \times \mathbb{R}^{>0}$ such that the following conditions hold:

(i) The pair (x^*, p^*) is an equilibrium of (11), i.e.,

$$\Psi(x^*, p^*) = -C x^* + S(p^* A x^* + b) = 0. \quad (20)$$

(ii) The Jacobian J_F^* has a zero eigenvalue with algebraic and geometric multiplicity q and corresponding orthonormal null eigenvectors $V = [v_1, \dots, v_q]$, orthonormal left null eigenvectors $W_0 = [w_1, \dots, w_q]$, and an orthonormal basis $W = [w_{q+1}, \dots, w_n]$ spanning $\ker(J_F^*)^\perp \cong \text{range}(J_F^{*\top})$. Define the complementary orthogonal projections $P_\parallel = VV^\top$ and $P_\perp = WW^\top = I - W_0W_0^\top$. $\text{Re}(\lambda_j) \neq 0$ for all other eigenvalues.

Analogously to the computation carried out for Hopfield networks, in the following sequence of Lemmas we derive expressions needed to define $L_\parallel(r_\parallel)$, $L_\perp(r_\parallel, r_\perp)$ in order to apply Theorem 3.3 to (11).

Lemma 4.4: $\|\xi_\parallel(\alpha, p)\| = \left\| P_\perp \begin{bmatrix} \Delta_x^\parallel(\alpha, p) AV & \Delta_p^\parallel(\alpha, p) \end{bmatrix} \right\|$ where

$$\Delta_x^\parallel(\alpha, p) := p D_z S(p A \Gamma(\alpha, \beta_0) + b) - p^* D_z S(p^* A x^* + b) \quad (21a)$$

$$\Delta_p^\parallel(\alpha, p) := D_z S(p A \Gamma(\alpha, \beta_0) + b) A \Gamma(\alpha, \beta_0) - D_z S(p^* A x^* + b) A x^* \quad (21b)$$

Proof: Following the definition (5a), we must compute $\xi_{\parallel\alpha}(\alpha, p)$ and $\xi_{\parallel p}(\alpha, p)$. Since $J_F^* V = 0$ we can write

$$C V = p^* D_z S(p^* A x^* + b) A V$$

which allows us to rewrite $\xi_{\parallel\alpha}(\alpha, p)$ as

$$\begin{aligned} \xi_{\parallel\alpha}(\alpha, p) &= D_x \Psi(\Gamma(\alpha, \beta_0), p) V \\ &= [-C + p D_z S(p A \Gamma(\alpha, \beta_0) + b) A] V \\ &= [p D_z S(p A \Gamma(\alpha, \beta_0) + b) - p^* D_z S(p^* A x^* + b)] A V =: \Delta_x^\parallel(\alpha, p) A V \end{aligned}$$

Next, we rewrite $\xi_{\parallel p}(\alpha, p)$ as

$$\begin{aligned} \xi_{\parallel p}(\alpha, p) &= D_p \Psi(V \alpha + W \beta_0, p) - D_p \Psi(x^*, p^*) \\ &= [D_z S(p A \Gamma(\alpha, \beta_0) + b) A \Gamma(\alpha, \beta_0) - D_z S(p^* A x^* + b) A x^*] =: \Delta_p^\parallel(\alpha, p) \end{aligned}$$

Finally, combining our expressions for $\xi_{\parallel\alpha}$ and $\xi_{\parallel p}$ yields

$$\xi_\parallel(\alpha, p) = W^\top \begin{bmatrix} \Delta_x^\parallel(\alpha, p) A V & \Delta_p^\parallel(\alpha, p) \end{bmatrix}.$$

The desired expression for $\|\xi_\parallel(\alpha, p)\|$ follows directly by applying Lemma 2.1 with $Y = A \begin{bmatrix} \Delta_x^\parallel(\alpha, p) V & \Delta_p^\parallel(\alpha, p) \end{bmatrix}$. ■

Lemma 4.5: $\|\xi_\perp(\alpha, \beta, p)\| = \|P_\perp \Delta_\perp(\alpha, \beta, p) A P_\perp\|$ where

$$\Delta_\perp(\alpha, \beta, p) := p D_z S(p A x + b) - p^* D_z S(p^* A x^* + b) \quad (22)$$

Proof: Building from its definition (5b), we have

$$\begin{aligned} \xi_\perp(\alpha, \beta, p) &= W^\top (D_x \Psi(\Gamma(\alpha, \beta), p) - D_x \Psi(x^*, p^*)) W \\ &= W^\top \left(-C + p D_z S(p A \Gamma(\alpha, \beta) + b) \right. \\ &\quad \left. + C - p^* D_z S(p^* A x^* + b) \right) A W \\ &= W^\top \Delta_\perp(\alpha, \beta, p) A W. \end{aligned}$$

Applying Lemma 2.2 to the above results in the desired expression for $\xi_\perp(\alpha, \beta, p)$. ■

Finally, we state an expression of the bounds for (11).

Theorem 4.6 (Bounds of bifurcation in Firing Rate models): Consider (11) under Assumption 2. For all $r_\parallel, r_\perp > 0$ satisfying

$$L_\parallel(r_\parallel) r_\parallel + L_\perp(r_\parallel, r_\perp) r_\perp < \sigma_{\min}(P_\perp J^* P_\perp) r_\perp - \|P_\perp D_z S(p^* A x^* + b) A x^*\| r_\parallel \quad (23)$$

where

$$L_\parallel(r_\parallel) = \sup \left\{ \left\| P_\perp \begin{bmatrix} \Delta_x^\parallel(\alpha, p) A V & \Delta_p^\parallel(\alpha, p) \end{bmatrix} \right\| : (\alpha, p) \in \mathcal{B}((\alpha_0, p^*), r_\parallel) \right\}$$

$$L_\perp(r_\parallel, r_\perp) = \sup \left\{ \|P_\perp \Delta_\perp(\alpha, \beta, p) A P_\perp\| : (\alpha, p) \in \mathcal{B}((\alpha_0, p^*), r_\parallel), \beta \in \mathcal{B}(\beta_0, r_\perp) \right\},$$

with $\Delta_x^\parallel$, $\Delta_p^\parallel$, $\Delta_\perp$ are defined in (21), (22), there exists a smooth implicit map $\varphi : \mathcal{B}((\alpha_0, p^*), r_\parallel) \rightarrow \mathcal{B}(\beta_0, r_\perp)$ that satisfies $\Psi(V \alpha + W \varphi(\alpha, p), p) = 0$.

Proof: Substituting the values of (4a), (4b), (6a), (6b) from Lemma 3.1, Lemma 3.2, Lemma 4.4, Lemma 4.5 into (8) from Theorem 3.3 yields the theorem statement. ■

V. APPLICATION TO CONSENSUS BIFURCATIONS

In recent years, models of the form (10),(11) have received attention in the literature on collective decision-making and opinion dynamics. In this context, components of the state vector x represent scalar opinions of individuals in a social network, which are updated in time according to a nonlinear saturating social rule. Of particular interest is the emergence of collective decision equilibria as bifurcations from the neutral $x = 0$ state. Such indecision-breaking bifurcations have been established in opinion dynamics models with both Hopfield-like structure [5], [7], [8] and firing rate-like structure [6], [10], [23]. In this section, we apply the bounds established in section IV to assess the validity region of the Lyapunov–Schmidt reduction analysis of indecision breaking supercritical pitchfork bifurcations in these nonlinear opinion dynamic models over regular balanced consensus graphs.

To specialize (10), (11) to nonlinear opinion dynamics (NOD) over regular graphs, we set $C = dI_n$ with $d > 0$, $p = u$, let A be the adjacency matrix of a k -regular, undirected and connected simple graph with unit edge weights, $S = \tanh$, and consider the unbiased case $b \equiv \mathbf{0}_n$. By the definition of a k -regular graph, the adjacency matrix A satisfies $A \mathbf{1}_n = k \mathbf{1}_n$. For both models, there is an indecision-breaking bifurcation point $(x^*, u^*) = (\mathbf{0}_n, \frac{d}{k})$ at which

consensus solutions emerge, see Fig. 1 for schematic. Note from (2) that $x^* = \mathbf{0}_n \implies (\alpha_0, \beta_0) = (0, \mathbf{0}_{n-1})$.




Fig. 1: Consensus pitchfork bifurcation in opinion dynamics over a regular graph at $(x^*, u^*) = (\mathbf{0}, \frac{d}{k})$; emerging branches of bistable equilibria lie on a manifold tangent to the consensus subspace $\{(x\mathbf{1}_n, u) : x, u \in \mathbb{R}\}$ at the bifurcation point.

A. Hopfield-structured NOD

Consider (10) for the n -node k -regular consensus graph:

$$\dot{x} = \Phi(x, u) = -dx + uA \tanh(x). \quad (25)$$

Jacobian $J^* := D_x \Phi(x^*, u^*)$ of system (25) simplifies to

$$J^* = -dI_n + u^* A \operatorname{diag}[\operatorname{sech}^2(x^*)] = \frac{d}{k}(A - kI_n). \quad (26)$$

We first characterize the orthonormal basis matrix V from (1). By the Perron–Frobenius theorem [24, Theorem 8.4.4], the adjacency matrix A of a k -regular, irreducible graph admits a simple largest eigenvalue k , with associated Perron eigenvector $\mathbf{1}_n$. Consequently, $\ker(J)$ is one-dimensional and spanned by the normalized vector $V = \frac{1}{\sqrt{n}}\mathbf{1}_n$.

We now evaluate the quantities defined in Theorem 4.3 in order to establish explicit bounds for the bifurcation. We begin with $M_{\parallel}$ and $L_{\parallel}(r_{\parallel})$, both of which evaluate to zero, as we justify below.

From its expression in (18), we find that

$$M_{\parallel} = \frac{1}{u^*} \|P_{\perp}(Cx^* - b)\| = \frac{k}{d} \|(I - VV^T)(dI_n \mathbf{0}_n)\| = 0.$$

To evaluate $L_{\parallel}(r_{\parallel})$, we first evaluate the terms $\Delta_{\parallel}(\alpha, u)$ and $\bar{S}_{\parallel}(\alpha)$ defined in (14) for $\xi_{\parallel}(\alpha, u)$ to obtain

$$\begin{aligned} \Delta_{\parallel}(\alpha, u) &= (us_{\alpha} - d/k)I_n, \quad \text{with } s_{\alpha} = \operatorname{sech}^2(\alpha/\sqrt{n}) \\ \bar{S}_{\parallel}(\alpha) &= t_{\alpha}\mathbf{1}_n, \quad \text{with } t_{\alpha} = \tanh(\alpha/\sqrt{n}). \end{aligned}$$

Substituting these expressions into Lemma 4.1 yields

$$\begin{aligned} \|\xi_{\parallel}(\alpha, u)\| &= \|P_{\perp}A[(us_{\alpha} - \frac{d}{k})V \quad t_{\alpha}\mathbf{1}_n]\| \\ &= \|P_{\perp}[(us_{\alpha} - \frac{d}{k})kV \quad t_{\alpha}k\mathbf{1}_n]\| \\ &= \|\mathbf{0}_{(n-1) \times 2}\| = 0 \end{aligned}$$

since both terms lie in the span of V , and are annihilated by $P_{\perp}$. Thus $\|\xi_{\parallel}(\alpha, u)\| = 0$, and from its definition in (6a) we conclude that $L_{\parallel}(r_{\parallel}) = 0$. Consequently, the inequality (8) reduces to the condition $M_{\perp}L_{\perp}(r_{\parallel}, r_{\perp}) < 1$. We now evaluate $M_{\perp}$ and $L_{\perp}(r_{\parallel}, r_{\perp})$ and state the bounds.

To evaluate $M_{\perp}$, observe that $\lambda_i(J^*) = \frac{d}{k}(\lambda_i(A) - k)$. For a k -regular, irreducible simple graph with adjacency matrix A , the Perron–Frobenius theorem guarantees that the largest eigenvalue of A is $\lambda_1(A) = k$, which is simple, and the subdominant eigenvalue satisfies $\lambda_2(A) < k$. Consequently,

$$\lambda_1(J^*) = 0, \quad \text{and} \quad \lambda_i(J^*) = \frac{d}{k}(\lambda_i(A) - k) < 0, \quad i > 1.$$

and hence, the smallest nonzero singular value of J^* is $\lambda_2(J^*) = \frac{d}{k}(\lambda_2(A) - k)$. Therefore,

$$M_{\perp} = \left(\frac{d}{k}|\lambda_2(A) - k|\right)^{-1} = \frac{k}{d(k - \lambda_2(A))}.$$

Note that the expression for $M_{\perp}$ depends inversely on the spectral gap $k - \lambda_2(A)$. Hence, larger spectral separation between the Perron root and the subdominant eigenvalue of A leads to smaller values of $M_{\perp}$.

Now, to evaluate $L_{\perp}(r_{\parallel}, r_{\perp})$, we first compute the expression for $\Delta_{\perp}(\alpha, \beta, u)$ defined in (14) for Lemma 4.2 as

$$\Delta_{\perp}(\alpha, \beta, u) = u \operatorname{diag}[\operatorname{sech}^2(\Gamma(\alpha, \beta))] - \frac{d}{k}I_n$$

To evaluate $\|\xi_{\perp}(\alpha, \beta, u)\|$, we first bound it from above using the submultiplicative property of the spectral norm

$$\begin{aligned} \|\xi_{\perp}(\alpha, \beta, u)\| &= \|P_{\perp}A\Delta_{\perp}(\alpha, \beta, u)P_{\perp}\| \\ &\leq \|P_{\perp}AP_{\perp}\| \|P_{\perp}\Delta_{\perp}(\alpha, \beta, u)P_{\perp}\| \end{aligned}$$

The spectral norm of the projection of A onto $\ker(J^*)^{\perp}$ simplifies to $\max_{i \neq 1} \{|\lambda_i(A)|\} = \max\{|\lambda_2(A)|, |\lambda_n(A)|\} =: |\lambda'(A)|$ where $\lambda_2(A)$ and $\lambda_n(A)$ denote the subdominant and smallest eigenvalues of A , respectively. For the projection of $\Delta_{\perp}(\alpha, \beta, u)$ onto $\ker(J^*)^{\perp}$, we find that its spectral norm is maximized at $(\alpha, \beta, u) = (0, \mathbf{0}_{n-1}, u^* \pm r_{\parallel})$. Since $\|\xi_{\parallel}(\alpha, u)\|$ obtains this upper bound at the same point, we conclude that $L_{\perp}(r_{\parallel}, r_{\perp}) = r_{\parallel} |\lambda'(A)|$.

Having computed all requisite quantities, we now state the bounds for the Hopfield regular balanced consensus network:

Theorem 5.1: The bounds of validity of the Lyapunov–Schmidt reduction from Proposition 2.3 around the pitchfork bifurcation at $(x^*, u^*) = (\mathbf{0}, \frac{d}{k})$ for the n -node k -regular balanced consensus opinion dynamic network (25) is

$$0 < r_{\parallel} < \frac{d(k - \lambda_2(A))}{k |\lambda'(A)|} \quad \text{and} \quad r_{\perp} > 0$$

along and normal to the consensus manifold $(\ker(J^*) \times \mathbb{R})$ (Figure 1) respectively.

Proof: Substituting the values of (4b) and (6a) into the reduced bounds inequality (7b) from Proposition 2.3 yields the desired bounds. ■

Corollary 1.1: For the case of the complete graph on n -nodes, i.e. $(n-1)$ -regular, the bounds from Theorem 5.1 simplify to $0 < r_{\parallel} < n$ and $r_{\perp} > 0$

Proof: For the complete graph, the adjacency matrix can be expressed as $A = \mathbf{1}_{n \times n} - I_n$ which has a null diagonal and unit non-diagonal entries, so each vertex has degree $k = n - 1$. The matrix $\mathbf{1}_{n \times n} =: C$ has $\lambda_1(C) = n$ with corresponding eigenvector $\mathbf{1}_n$, while the remaining eigenvalues $\lambda_{j \geq 2} = 0$




Fig. 2: Variation in bounds of validity for Lyapunov–Schmidt reduction for consensus bifurcation over balanced regular graphs. For each (n, k) pair, 100 independent random graphs were generated using the NetworkX package in Python. The reported mean values and corresponding error bars represent the average and standard deviation of the computed bounds on $r_{\parallel}$ across the sampled graphs. (a) illustrates the bounds with fixed degree $k \in \{12, 18, 24, 30, 36\}$, varying graph size n ; (b) illustrates the bounds with fixed graph size $n \in \{12, 18, 24, 30, 36\}$, varying degree k .

with the $n - 1$ eigenvectors orthogonal to $\mathbf{1}_n$. Consequently, $\lambda_1(A) = n - 1$ and $\lambda_{j \geq 2} = -1$. Substituting these into Theorem 5.1 yields $0 < r_{\parallel} \leq n$ and $r_{\perp} > 0$. ■

B. Firing-Rate-structured NOD

We now repeat this analysis for (11) over the same n –node k –regular consensus graph:

$$\dot{x} = \Psi(x, u) = -dx + \tanh(uAx). \quad (27)$$

We will now show that all the relevant quantities admit identical expressions between the Hopfield and Firing-Rate models. Firstly, Jacobian $J^* := D_x \Psi(x^*, u^*)$ of system (27) simplifies to $J^* = -dI_n + \frac{d}{k} \text{diag}[\text{sech}^2(uA\mathbf{0}_n)]A = \frac{d}{k}(A - kI_n)$ which is precisely the same expression obtained in the Hopfield case (26). Consequently, we have the same orthonormal basis $V = \frac{1}{\sqrt{n}}\mathbf{1}_n$ for $\ker(J^*)$.

To show that $M_{\parallel} = 0$, observe that

$$M_{\parallel} = \|P_{\perp} \text{diag}[\text{sech}^2(\frac{d}{k}A\mathbf{0}_n)]A\mathbf{0}_n\| = 0$$

and for $L_{\parallel}(r_{\parallel})$, we first compute the terms defined in (21) for $\xi_{\parallel}(\alpha, u)$ to obtain

$$\Delta_x(\alpha, u) = (us_{uk\alpha} - d/k)I_n, \quad \Delta_u(\alpha, u) = k\alpha s_{uk\alpha}\mathbf{1}_n,$$

where $s_{uk\alpha} = \text{sech}^2(uk\alpha/\sqrt{n})$. Substituting into Lemma 4.4

$$\begin{aligned} \|\xi_{\parallel}(\alpha, u)\| &= \|P_{\perp} [(us_{uk\alpha} - d/k)AV \quad k\alpha s_{uk\alpha}\mathbf{1}_n]\| \\ &= \|P_{\perp} [(uks_{uk\alpha} - d)V \quad k\alpha s_{uk\alpha}\mathbf{1}_n]\| = 0 \end{aligned}$$

since both terms both terms lie in the span of V , which is annihilated by $P_{\perp}$. From (6a) we conclude that $L_{\parallel}(r_{\parallel}) = 0$. Again, inequality (8) reduces to $M_{\perp}L_{\perp}(r_{\parallel}, r_{\perp}) < 1$. It remains to compute $M_{\perp}$ and $L_{\perp}(r_{\parallel}, r_{\perp})$.

Since J^* remains unchanged from the Hopfield model, we identically have $M_{\perp} = \frac{k}{d(k - \lambda_2(A))}$. To justify that $L_{\perp}(r_{\parallel}, r_{\perp})$ also remains unchanged, we compute the expression for defined in (22) for Lemma 4.5 to get $\Delta_{\perp}(\alpha, \beta, u) = u \text{diag}(\text{sech}^2(uA\Gamma(\alpha, \beta))) - \frac{d}{k} \text{diag}(\text{sech}^2(\frac{d}{k}A\mathbf{0}_n)) =$

$u \text{diag}(\text{sech}^2(uA\Gamma(\alpha, \beta))) - \frac{d}{k}I_n$. By the same reasoning as in the Hopfield case, we can say that that $\|\xi_{\parallel}(\alpha, u)\|$ attains its supremum at $(\alpha, \beta, u) = (0, \mathbf{0}_{n-1}, u^* \pm r_{\parallel})$. Consequently, identical to the Hopfield case, $L_{\perp}(r_{\parallel}, r_{\perp}) = r_{\parallel} |\lambda'(A)|$. Therefore, the bounds of validity for the reduced order model obtained via Lyapunov–Schmidt reduction at the indecision-breaking consensus bifurcation in the Firing Rate-form NOD model (27) coincide with those of the Hopfield-form NOD model (25), as stated in Theorem 5.1. We illustrate how these bounds depend on the network size n and node degree k in Figure 2.

VI. DISCUSSION

In this article we established explicit, computable bounds for the validity of Lyapunov–Schmidt reduction-based bifurcation analysis in representative classes of networked nonlinear systems, and demonstrated their utility by applying the bounds to an illustrative class of nonlinear consensus problems on regular graphs, for which the evaluation proves to be analytically computable. Notably, the derived expressions depend directly on interpretable features, such as the spectral properties of the underlying graph, providing insight into how network structure influences these bounds.

For future work, we aim to identify additional classes of systems with bifurcations analyzed with the Lyapunov–Schmidt reduction for which these bounds can be computed analytically.

REFERENCES

- [1] J. J. Hopfield, “Neurons with graded response have collective computational properties like those of two-state neurons,” *Proceedings of the national academy of sciences*, vol. 81, no. 10, pp. 3088–3092, 1984.
- [2] S. Betti, G. Baggio, F. Bullo, and S. Zampieri, “Input-driven dynamics for robust memory retrieval in hopfield networks,” *Science Advances*, vol. 11, no. 17, p. eadu6991, 2025.
- [3] H. R. Wilson and J. D. Cowan, “Excitatory and inhibitory interactions in localized populations of model neurons,” *Biophysical journal*, vol. 12, no. 1, pp. 1–24, 1972.
- [4] S. Betti, G. Baggio, F. Bullo, and S. Zampieri, “Firing rate models as associative memory: Synaptic design for robust retrieval,” *Neural Computation*, pp. 1–32, 2025.
- [5] A. Franci, V. Srivastava, and N. E. Leonard, “A realization theory for bio-inspired collective decision-making,” *arXiv preprint arXiv:1503.08526*, 2015.
- [6] A. Bizyaeva, A. Franci, and N. E. Leonard, “Nonlinear opinion dynamics with tunable sensitivity,” *IEEE Transactions on Automatic Control*, vol. 68, no. 3, pp. 1415–1430, 2023.
- [7] R. Gray, A. Franci, V. Srivastava, and N. E. Leonard, “Multiagent decision-making dynamics inspired by honeybees,” *IEEE Transactions on Control of Network Systems*, vol. 5, no. 2, pp. 793–806, 2018.
- [8] A. Fontan and C. Altafini, “Multiequilibria analysis for a class of collective decision-making networked systems,” *IEEE Transactions on Control of Network Systems*, vol. 5, no. 4, pp. 1931–1940, 2017.
- [9] P. U. Abara, F. Ticozzi, and C. Altafini, “Spectral conditions for stability and stabilization of positive equilibria for a class of nonlinear cooperative systems,” *IEEE Transactions on Automatic Control*, vol. 63, no. 2, pp. 402–417, 2017.
- [10] N. E. Leonard, A. Bizyaeva, and A. Franci, “Fast and flexible multiagent decision-making,” *Annual Review of Control, Robotics, and Autonomous Systems*, vol. 7, 2024.
- [11] Z. Tang, J. H. Park, and H. Shen, “Finite-time cluster synchronization of lur’e networks: A nonsmooth approach,” *IEEE Transactions on Systems, Man, and Cybernetics: Systems*, vol. 48, no. 8, pp. 1213–1224, 2017.
- [12] F. Liu, Q. Song, G. Wen, J. Lu, and J. Cao, “Bipartite synchronization of lur’e network under signed digraph,” *International Journal of Robust and Nonlinear Control*, vol. 28, no. 18, pp. 6087–6105, 2018.

- [13] F. A. Miranda-Villatoro, F. Forni, and R. J. Sepulchre, "Analysis of lur'e dominant systems in the frequency domain," *Automatica*, vol. 98, pp. 76–85, 2018.
- [14] B. C. Goodwin, "Oscillatory behavior in enzymatic control processes," *Advances in enzyme regulation*, vol. 3, pp. 425–437, 1965.
- [15] N. Rosenfeld, M. B. Elowitz, and U. Alon, "Negative autoregulation speeds the response times of transcription networks," *Journal of molecular biology*, vol. 323, no. 5, pp. 785–793, 2002.
- [16] T. S. Gardner, C. R. Cantor, and J. J. Collins, "Construction of a genetic toggle switch in *escherichia coli*," *Nature*, vol. 403, no. 6767, pp. 339–342, 2000.
- [17] M. Golubitsky and I. Stewart, *Dynamics and bifurcation in networks: theory and applications of coupled differential equations*. SIAM, 2023.
- [18] P. Gupta, A. Bizyaeva, and R. Banavar, "Estimates on the domain of validity for lyapunov-schmidt reduction," in *2024 IEEE 63rd Conference on Decision and Control (CDC)*, pp. 4393–4398, IEEE, 2024.
- [19] A. Jindal, D. Chatterjee, and R. Banavar, "Estimates of the size of the domain of the implicit function theorem: a mapping degree-based approach," *Mathematics of Control, Signals, and Systems*, vol. 36, no. 1, pp. 139–175, 2024.
- [20] M. Golubitsky and D. G. Schaeffer, *Singularities and Groups in Bifurcation Theory*, vol. 51 of *Applied Mathematical Sciences*. New York, NY: Springer-Verlag, 1985.
- [21] J. W. Demmel, *Applied numerical linear algebra*. SIAM, 1997.
- [22] K. D. Miller and F. Fumarola, "Mathematical equivalence of two common forms of firing rate models of neural networks," *Neural computation*, vol. 24, no. 1, pp. 25–31, 2012.
- [23] A. Bizyaeva, A. Matthews, A. Franci, and N. E. Leonard, "Patterns of nonlinear opinion formation on networks," in *2021 American Control Conference (ACC)*, pp. 2739–2744, IEEE, 2021.
- [24] R. A. Horn and C. R. Johnson, *Matrix analysis*. Cambridge university press, 2012.