

# INFECTION VS FATALITY OF COVID-19 IN NEW YORK STATE: EFFECTS OF DEMOGRAPHICS AND POOR AIR QUALITY

Department of Mathematics, Clarkson University

Presenter: Vijay Kumar

Date: 07/30/2020

Mentors: Dr. Sumona Mondal, Dr. Suresh Dhaniyala

Dr. Shantanu Sur, Dr. Supraja Gurajala


Collaborators: Chaya Chaipitakporn, Bridget Wangler



**AIR LAB**  
AEROSOL INSTRUMENTATION RESEARCH

# Background


- COVID-19 is ongoing pandemic, infected the whole world
- Caused more than 610K deaths
- New York State was hit badly and become an epicenter in US (March-June,2020)
- Exposure to poor air quality leads to the risk of COVID-19



# Objectives and Approach

## Objectives:

- Comparing the effect of COVID-19 within counties of New York State
- Estimating the different causes of variation
- Identifying the important risk factors causing infection, death and fatality rate




## Study Workflow

**Infection Group** is infection per 100K population  
**Death Group** is death per 100K population  
**Fatality Group** is death per 10k infected population

<https://www.census.gov/programs-surveys/acs>  
[https://aqs.epa.gov/aqsweb/documents/data\\_api.html](https://aqs.epa.gov/aqsweb/documents/data_api.html)  
<https://www.syracuse.com/data/>

# Clustering Analysis

- Classification of the counties, K-means clustering
- Infection and death rate cluster 1 and 2 are near the epicenter
- But Fatality rate counties cluster 1 is not near epicenter
- Highly infected cluster is the downstate NYS
- Lowly infected cluster are in the upstate and northern counties of NYS




Clusters of counties with infection, death and fatality rates

# Demographic factors

The demographic factors


1. Population Density (people per square miles)
  2. Population with age >55 years (%)
  3. Population of African American ethnicity (%)
  4. Population of Hispanic American ethnicity (%)
  5. Population below the poverty line(%)
  6. Distance from the epicenter-Manhattan (miles)
- Horizontal lines represent median with the actual values, while P-values were from non parametric Mann-Whitney-U tests after Bonferroni corrections
  - Age> 55 years is taking lead in 1<sup>st</sup> cluster of Fatality group
  - The Kruskal Wallis test demonstrated that all of our demographic parameters were significant at the 95% confidence level except poverty



Dot plots of Demographic variables in three groups with three clusters

# ARIMA models

- Line plots are observed AQI values
- 95% Confidence band AQI values using Autoregressive Integrated Moving Average models
- In dot plots horizontal lines represent median with the actual values
- P-values were from non parametric Mann-Whitney U tests
- Cluster 3 is significantly different from 1 & 2 in both plots
- ARIMA plot shows cluster 3 has consistently lower AQI values




Dot plot and ARIMA time series plots for the comparisons of Air Quality variables



# Stepwise Regression and significant factors

- Important factors are ranked by forward selection algorithm
- Infection and death groups have ethnicity as the most important predictor
- Fatality group was different than the other two groups
- Older age became the most prominent factor
- Bad air quality is 2<sup>nd</sup> or 3<sup>rd</sup> most important factor



Significant factors and relative contribution of different factors in three groups

# Conclusion

- Infection and death rates are high in counties located near the New York City
- In Fatality, several other counties take up the topmost positions even having a low infection rate
- Regression model shows **ethnicity (African-American and Hispanic)** and **poverty** are major risk factors for infection and death rate
- Fatality has a strong association with **age and  $PM_{2.5}$**
- Our results show distinct contributions by various risk factors to the COVID-19



# Acknowledgment

## Mentors:

- Prof. Sumona Mondal, Department of Mathematics, Clarkson University
- Prof. Suresh Dhaniyala, Department of Mechanical and Aeronautical Engineering, Clarkson University
- Prof. Supraja Gurajala, Department of Computer Science, SUNY Potsdam
- Prof. Shantanu Sur, Department of Biology, Clarkson University

## Colleagues:

- Chaya Chaipitakporn , David D. Reh School of Business, Clarkson University
- Bridget Wangler, Department of Mathematics, Clarkson University
- Morgan Busch, Department of Computer Science, Clarkson University

## Personal Contact

- Email: [vikumar@clarkson.edu](mailto:vikumar@clarkson.edu)

**AIR LAB**  
**AEROSOL INSTRUMENTATION RESEARCH**



# References

- [1] Guo, Y.-R. et al. The origin, transmission and clinical therapies on coronavirus disease 2019 (covid-19) outbreak—an update on the status. *Mil. Med. Res.* 7, 1–10 (2020)
- [2] Surveillances, V. The epidemiological characteristics of an outbreak of 2019 novel coronavirus diseases (covid-19)—china, 2020. *China CDC Wkly.* 2, 113–122 (2020)
- [3] Richardson, S. et al. Presenting characteristics, comorbidities, and outcomes among 5700 patients hospitalized with covid-19 in the new york city area. *Jama* (2020)
- [4] Dudley, J. P. & Lee, N. T. Disparities in age-specific morbidity and mortality from sars-cov-2 in china and the republic of korea. *Clin. Infect. Dis.* (2020).
- [5] Yang, Z. et al. Acute effects of air pollution on the incidence of hand, foot, and mouth disease in wuhan, china. *Atmospheric Environ.* 225, 117358 (2020)
- [6] <https://towardsdatascience.com/covid-19-infection-in-italy-mathematical-models-and-predictions-7784b4d7dd8d>



# Thank You!