

**Vijay Kumar<sup>1</sup>, Dinushani Senarathna<sup>1</sup>, Supraja Gurajala<sup>2</sup>, William Olsen<sup>5</sup>, Shantanu Sur<sup>4</sup>, Sumona Mondal<sup>1</sup>, Suresh Dhaniyala<sup>3</sup>**

<sup>1</sup>Department of Mathematics, <sup>2</sup>Department of Computer Science, <sup>3</sup>Department of Mechanical and Aerospace Engineering, <sup>4</sup>Department of Biology, <sup>5</sup>Department of Civil and Environmental Engineering, Clarkson University and Department of Computer Science, SUNY Potsdam

## Background

### Motivations:

- Low-cost air sensor data increasingly used in health studies
- Sensitivity, noise, and accuracy of data of sensors remain a concern
- Differences exist between data from EPA and low-cost sensors
- What is driving these differences, and can that be "calibrated" out?




Figure 1: Purple Air (PA) sensor


### Research objectives:

- Analyze the temporal differences in PM<sub>2.5</sub> data from EPA and purple air (PA) low-cost sensor network
- Characterize the influence of different aerosol sources on the EPA and PA data

## Methodology




**Data source:**  
EPA and Purple air (PA) data;  
Cook county, IL;  
October 2019 to September 2021



(A) Time series data of selected EPA & PA sites for the study duration (B) Scatter plot showing relation between selected EPA & PA sites in close vicinity

## Results

### Monitoring data:




Distribution of EPA and PA PM<sub>2.5</sub> over the analysis time period

- PA data on average higher median values than EPA for all sites
- PA data has a broader distribution

### Spectral analysis:


- Obtained power spectrum of the two data sources EPA and PA; For simplicity just averaged the data from the different sites for each of the sources



- Identified that there are selective frequencies (e.g., at 4, 8, 12, 24 hours); that influence both data sets possibly corresponding to traffic patterns, local industrial emission, etc.
- Power spectrum of the PA and EPA different at two ends of spectrum
  - PA has a higher relative power spectrum at low frequency (i.e., for time periods >1 day). Possibly suggesting an over contribution of regional aerosol to the signal
  - EPA has a higher power spectrum at short time periods – probably indicating its relatively better sensitivity to local sources e.g., traffic


## Results

### Relative contributions of temporal components of PM<sub>2.5</sub>



EPA E2 site PM<sub>2.5</sub> time series (A), decomposition of EPA site E2 PM<sub>2.5</sub> time series into: (B) Short-term component (C) Seasonal component (D) Long-term component

- Decomposed signal from the data sets into short term (< 24 hrs), seasonal (< 60 days), and long-term.
- EPA Short-term component contribution to total signal is 3 times that of PA – suggesting significant loss of sensitivity to fast sources such as traffic
- Long-term components have similar contribution



## Conclusion

- Lower accuracy of PA data relative to EPA data has been previously identified, but reasons are uncertain
- Most correction models based on regression; Models will not be valid if the distribution of sources change
- Our study shows PA data comparable in efficiency to EPA for long-term variations in the PM<sub>2.5</sub> data but less efficient in detecting short term variations (<12 hours) suggesting strong source dependence in sensitivity
- Models based on frequency-decomposition of data could enable corrections accounting for different sensitivity to different aerosol sources

## References

- Kuula, J., Mäkitalo, T., Aurela, M., Teirila, K., Varjonen, S., González, O., and Timonen, H.: Laboratory evaluation of particle-size selectivity of optical low-cost particulate matter sensors, *Atmospheric Measurement Techniques*, 13, 2413–2423, 2020, <https://doi.org/10.5194/amt-13-2413-2020>
- Tryner, I., Mehaffy, J., Miller-Lionberg, D., and Volkens, J.: Effects of aerosol type and simulated aging on performance of low-cost PM sensors, *Journal of Aerosol Science*, 150, 105 654, 2020, <https://doi.org/10.1016/j.jaerosci.2020.105654>
- Castell, N., Dauge, F. R., Schneider, P., Vogt, M., Lerner, U., Fishbain, B., Broday, D., and Bartonoval, A.: Can commercial low-cost environment international, 99, 293–302, 2017
- Tchepel, O. and Borrego, C.: Frequency analysis of air quality time series for traffic related pollutants, *Journal of Environmental Monitoring*, 12, 544–550, 2010, <https://doi.org/10.1039/B906717E>
- Zhang, Z., Kim, S.-J., and Ma, Z.: Significant decrease of PM<sub>2.5</sub> in Beijing based on long-term records and Kolmogorov-Zurbenko filter approach, 2018, <https://doi.org/10.4209/aaer.2017.01.0011>
- Bai, H., Gao, W., Zhang, Y., and Wang, L.: Assessment of health benefit of PM<sub>2.5</sub> reduction during COVID-19 lockdown in China Journal of Environmental Sciences, 115, 422–431, 2022, <https://doi.org/10.1016/j.jes.2021.01.022>

sensor, platforms contribute to air quality monitoring and exposure estimates? sensor, platforms contribute to air quality monitoring and exposure estimates?,  
and separating contributions from anthropogenic emissions and meteorology,  
<https://doi.org/10.1016/j.jes.2021.01.022>