

# Homework 02

## REAL VARIABLES I

### Fall 2026

**Due:** 2026-09-06 19:00

**Topics:** Lebesgue outer measure, outer measures generated by costs, interval coverings, and outer Jordan content

#### Instructions

Submission is through Gradescope, accessible through [Blackboard](#).

**Grading rubric:** Every problem/subproblem is graded according to this rubric (no intermediate scores):

- **0%:** No submission or no meaningful content.
- **30%:** Attempt at solution, outline of idea.
- **60%:** Serious attempt at solution. Solution fails to address important points.
- **80%:** Solutions with fixable mistakes or incompleteness.
- **100%:** Complete and correct solution.

You may receive up to a 10% penalty for very unclear or messy exposition.

Please start every solution on a separate page. Clearly mark the pages associated to every part. You may submit either computer typeset solutions or handwritten solutions. Handwritten solutions must be *very clearly* legible.

You are encouraged to work in groups. You may submit the entire homework assignment as a group (up to 4 people). If you only did some exercises as a group, instead of a solution you may clearly indicate whose submission I should look at for the solution to the problem.

Indicate any external help/notes/resources you used. You may not use generative AI in any way to help you with the HW.

#### 1. (Images of functions with bounded derivative)

##### IN CLASS

**1.a** (weight: 1.0) Let  $f: \mathbb{R} \rightarrow \mathbb{R}^2$  be differentiable and suppose that its derivative is bounded. Prove that the image  $f(\mathbb{R})$  has two-dimensional Lebesgue outer measure zero.

**Free hint.** First prove that  $\mathcal{L}_2^*(f([-N, N])) = 0$  for every  $N \in \mathbb{N}$ . Then use  $\sigma$ -additivity of outer measures

#### 2. (Equivalent interval definitions of Lebesgue outer measure)

**2.a** (weight: 1.0) A *dyadic interval* is a half-open interval of the form

$$[k2^{-m}, (k+1)2^{-m}), \quad k, m \in \mathbb{Z}.$$

On each of the following three collections, assign every interval its length:

1. bounded open intervals;
2. bounded closed intervals;
3. dyadic intervals.

Prove that all three associated generated outer measures are the same. In particular, each one equals the Lebesgue outer measure  $\mathcal{L}_1^*$ .

### 3. (Finite interval covers and outer Jordan content)

**3.a** (weight: 0.3) Prove that  $\mathcal{L}_1^*([0, 1] \setminus \mathbb{Q}) = 1$

The purpose of this exercise is to show that covering by a finite number of intervals will not suffice in the definition of Lebesgue outer measure.

**3.b** (weight: 0.7) The *outer Jordan content* of a set  $E \subseteq \mathbb{R}$  is defined by

$$J(E) = \inf \left\{ \sum_{I \in \mathcal{J}} \ell(I) : \mathcal{J} \text{ is a **finite** collection of bounded closed intervals and } E \subseteq \bigcup_{I \in \mathcal{J}} I \right\}.$$

As usual,  $\ell([a, b]) = b - a$ .

You do not need to provide a detailed proof of the fact that if  $E = [c, d]$  is a bounded closed interval and  $[c, d] \subseteq \bigcup_{n=1}^N [a_n, b_n]$  for some  $N \in \mathbb{N}$  then  $\sum_{n=1}^N (b_n - a_n) \geq d - c$ , but you should recall/provide a reference of a fact we proved in class that uses the argument that can be directly adapted to prove this claim.

Prove that  $J(E) = J(\bar{E})$ . Then exhibit a countable set  $E \subseteq \mathbb{R}$  such that  $J(E) = 1$ , whereas  $\mathcal{L}_1^*(E) = 0$ .