



## MA 106 Help Session-2

Guramrit Singh  
(210050061)

March 23, 2023

*These are some of the interesting questions discussed in the Help Session conducted on March 23, 2023.*

### Problem 1 (Tutorial 3: Q15)

Given  $n^2$  functions  $f_{ij}(x)$  each differentiable on the interval  $(a, b)$ , define  $f(x) = \det(f_{ij}(x))$  for each  $x \in (a, b)$ . Let  $A(x) = (f_{ij}(x))$ . Let  $A_i(x)$  be the matrix obtained from  $A(x)$  by differentiating the functions in the  $i^{\text{th}}$  row of  $A(x)$ . Prove that  $f'(x) = \sum_{i=1}^n \det A_i(x)$

**Solution:**

We prove it by induction on order of matrix i.e.  $n$ .

**Base case:**  $n = 1$

Trivial to see since  $f(x) = f_{11}(x)$  and so  $f'(x) = f'_{11}(x) = \det A_1(x) = \sum_{i=1}^1 \det A_i(x)$

**Induction hypothesis:** Assume the result  $f'(x) = \sum_{i=1}^{n-1} \det A_i(x)$  holds true for all  $(n-1) \times (n-1)$  matrices.

Now, we show that it is true for all  $n \times n$  matrices.

Let  $A_{jk}$  denote the sub-matrix of  $A$  obtained by removing the  $j^{\text{th}}$  row and the  $k^{\text{th}}$  column.

$$\det(A) = \sum_{i=1}^n (-1)^{i+1} f_{1i}(x) \det(A_{1i})$$

Differentiate both sides

$$f'(x) = \sum_{i=1}^n (-1)^{i+1} (f'_{1i}(x) \det(A_{1i}) + f_{1i}(x) (\det(A_{1i}))')$$

Note that  $(\det(A_{1i}))' = \sum_{j=1}^{n-1} \det(A_{1i_j}(x))$  by induction hypothesis as  $\forall i$   $A_{1i}$  is  $(n-1) \times (n-1)$  matrix .

$$\begin{aligned}
 f'(x) &= \sum_{i=1}^n (-1)^{i+1} (f'_{1i}(x) \det(A_{1i}) + f_{1i}(x) (\sum_{j=1}^{n-1} \det(A_{1i_j}(x)))) \\
 &= \sum_{i=1}^n (-1)^{i+1} f'_{1i}(x) \det(A_{1i}) + \sum_{j=1}^{n-1} \sum_{i=1}^n (-1)^{i+1} f_{1i}(x) \det(A_{1i_j}(x)) \\
 &= \det(A_1(x)) + \sum_{j=1}^{n-1} \det(A_{j+1}(x)) \\
 &= \sum_{i=1}^n \det(A_i(x))
 \end{aligned}$$

Hence proved

## Problem 2

Let  $\mathbf{A} \in \mathbb{R}^{9 \times 4}$  and  $\mathbf{B} \in \mathbb{R}^{7 \times 3}$ . Is there  $\mathbf{X} \in \mathbb{R}^{4 \times 7}$  such that  $\mathbf{X} \neq \mathbf{O}$  but  $\mathbf{AXB} = \mathbf{O}$ ?

**Hint:**  $\mathbf{AXB} = \mathbf{O}$  gives us  $9 \times 3 = 27$  equations to solve in  $4 \times 7 = 28$  variables.

Also note that the obtained system is homogeneous. The augmented matrix can have at most 27 pivots and therefore you can find at least one non-null solution to  $\mathbf{X}$

## Problem 3

Given a square matrix  $\mathbf{A}$  of order  $n \geq 2$ . Represent  $\mathbf{A}$  as

- (a) Sum of two invertible matrices
- (b) Sum of two non-invertible matrices

- (a)  $\mathbf{A} = \mathbf{L} + \mathbf{U}$  where  $\mathbf{L}$  and  $\mathbf{U}$  are lower and upper triangular matrices, select diagonal entries s.t. in both  $\mathbf{L}$  and  $\mathbf{U}$ , they are all non-zero. Then  $\mathbf{L}$  and  $\mathbf{U}$  are required invertible matrices. (Determinant of either lower or upper triangular matrix is just the product of diagonal entries)
- (b)  $\mathbf{A} = \mathbf{L} + \mathbf{U}$  where  $\mathbf{L}$  and  $\mathbf{U}$  are lower and upper triangular matrices respectively, select diagonal entries s.t. in both  $\mathbf{L}$  and  $\mathbf{U}$ , there is at least one zero entry. Then  $\mathbf{L}$  and  $\mathbf{U}$  are required non-invertible matrices. (Determinant of either lower or upper triangular matrix is just the product of diagonal entries)

## Problem 4

tut 1 q 10

## Problem 5

tut 2 q6... q12 build on this



## Problem 6

tut 3 q5