

Compounds of Transitional elements

321. The number of unpaired electrons in Zn^{2+} is
 (a) 2 (b) 3
 (c) 4 (d) 0
322. The trace metal present in insulin is
 (a) Iron (b) Cobalt
 (c) Zinc (d) Manganese
323. The chemical name of borax is
 (a) Sodium orthoborate
 (b) Sodium metaborate
 (c) Sodium tetraborate
 (d) Sodium tetraborate decahydrate
324. Hydrogen is not obtained when zinc reacts with
 (a) Cold water
 (b) Dilute H_2SO_4
 (c) Dilute HCl
 (d) Hot 20% $NaOH$
325. The metal which gives hydrogen on treatment with acid as well as sodium hydroxide is
 (a) Iron
 (b) Zinc
 (c) Copper
 (d) None of the above
326. To protect iron against corrosion, the most durable metal plating on it, is
 (a) Nickel plating
 (b) Tin plating
 (c) Copper plating
 (d) Zinc plating
327. The compound $ZnFe_2O_4$ is
 (a) A normal spinel compound
 (b) Interstitial compound
 (c) Covalent compound
 (d) Co-ordination compound
328. ZnO when heated with BaO at $1100^\circ C$ gives a compound. Identify the compound
 (a) $BaZnO_2$ (b) $BaO_2 + Zn$
 (c) $BaCdO_2$ (d) $Ba + ZnO_2$
329. Zn gives hydrogen gas with H_2SO_4 and HCl but not with HNO_3 because
 (a) NO_2 is reduced in preference to H_3O^+
 (b) HNO_3 is weaker acid than H_2SO_4 and HCl
 (c) Zn acts as oxidising agent when reacts with HNO_3
 (d) In electrochemical series Zn is placed above the hydrogen
330. The metal used for making radiation shield is
 (a) Aluminium (b) Iron
 (c) Zinc (d) Lead
331. Which of the following metal is obtained by leaching out process using a solution of $NaCN$ and then precipitating the metal by addition of zinc dust



- (a) Copper (b) Silver (a) Iron (b) Zinc
(c) Nickel (d) Iron (c) Silver (d) Lead
332. While extracting an element from its ore, the ore is ground and leached with dilute KCN solution to form the soluble product potassium argentocyanide. The element is
(a) Lead (b) Chromium
(c) Manganese (d) Silver
333. In Mc Arthur Forest method, silver is extracted from the solution of $Na[Ag(CN)_2]$ by the use of:
(b) Fe (b) Mg
(c) Cu (d) Zn
334. Iron obtained from blast furnace is known as
(a) Wrought iron (b) Cast iron
(c) Pig iron (d) Steel
335. Extraction of silver from commercial lead is possible by
(a) Mond's process
(b) Park's process
(c) Haber's process
(d) Clark's process
336. Impurities of lead in silver are removed by
(a) Park process
(b) Solvey process
(c) Cyanide process
(d) Amalgamation process
337. Park's process is used in the extraction of
(a) Iron (b) Zinc
(c) Silver (d) Lead
338. From argentite (Ag_2S) ore the method used in obtaining metallic silver is
(a) Fused mixture of Ag_2S and KCl is electrolysed
(b) Ag_2S is reduced with CO
(c) Ag_2S is roasted to Ag_2O which is reduced with carbon
(d) Treating argentite with $NaCN$ solution followed by metal displacement with zinc
339. In the extraction of zinc which gas is burnt in the jackets surrounding the retorts
(a) Water gas
(b) Producer gas
(c) Oil gas
(d) Coal gas
340. MacArther process is used for
(a) Hg (b) Fe
(c) Cl (d) O_2

