



Nature of Data

What is Data?

- Collection of *data objects (observations)* and their *attributes (features)*
- An *attribute* is a property or characteristic of an object
 - Examples: eye color of a person, temperature, etc.
 - Attribute is also known as feature, variable, field, characteristic.
- A collection of attributes describe an *object*
 - Object is also known as observation, record, point, case, sample, or instance

Features				
Tid	Refund	Marital Status	Taxable Income	Cheat
1	Yes	Single	125K	No
2	No	Married	100K	No
3	No	Single	70K	No
4	Yes	Married	120K	No
5	No	Divorced	95K	Yes
6	No	Married	60K	No
7	Yes	Divorced	220K	No
8	No	Single	85K	Yes
9	No	Married	75K	No
10	No	Single	90K	Yes

A More Complete View of Data


- Data may have parts
- The different parts of the data may have relationships
- More generally, data may have structure
- Data can be incomplete

Attribute Values

- **Attribute values** are numbers or symbols assigned to an attribute for a particular object
- Distinction between attributes and attribute values
 - Same attribute can be mapped to different attribute values
 - ◆ Example: height can be measured in feet or meters
 - Different attributes can be mapped to the same set of values
 - ◆ Example: Attribute values for ID and age are integers
 - ◆ But properties of attribute values can be different

Measurement of Length

- The way you measure an attribute may not match the attributes properties.



Types of Attributes

- There are different types of attributes
 - **Nominal**
 - ◆ Examples: ID numbers, eye color, zip codes
 - **Ordinal**
 - ◆ Examples: rankings (e.g., taste of potato chips on a scale from 1-10), grades, height {tall, medium, short}
 - **Interval**
 - ◆ Examples: calendar dates, temperatures in Celsius or Fahrenheit.
 - **Ratio**
 - ◆ Examples: length, time, counts

Properties of Attribute Values

- The type of an attribute depends on which of the following properties/operations it possesses:
 - Distinctness: $= \neq$
 - Order: $< >$
 - Differences are meaningful : $+ -$
 - Ratios are meaningful $* /$
 - Nominal attribute: distinctness
 - Ordinal attribute: distinctness & order
 - Interval attribute: distinctness, order & meaningful differences
 - Ratio attribute: all 4 properties/operations

		Attribute Type	Description	Examples	Operations
Categorical Qualitative		Nominal	Nominal attribute values only distinguish. ($=$, \neq)	zip codes, employee ID numbers, eye color, sex: <i>{male, female}</i>	mode, entropy, contingency correlation, χ^2 test
		Ordinal	Ordinal attribute values also order objects. ($<$, $>$)	hardness of minerals, <i>{good, better, best}</i> , grades, street numbers	median, percentiles, rank correlation, run tests, sign tests
Numeric Quantitative		Interval	For interval attributes, differences between values are meaningful. ($+$, $-$)	calendar dates, temperature in Celsius or Fahrenheit	mean, standard deviation, Pearson's correlation, t and F tests
		Ratio	For ratio variables, both differences and ratios are meaningful. ($*$, $/$)	temperature in Kelvin, monetary quantities, counts, age, mass, length, current	geometric mean, harmonic mean, percent variation

Categorical
QualitativeNumeric
Quantitative

Attribute Type	Transformation	Comments
Nominal	Any permutation of values	If all employee ID numbers were reassigned, would it make any difference?
Ordinal	An order preserving change of values, i.e., $new_value = f(old_value)$ where f is a monotonic function	An attribute encompassing the notion of good, better best can be represented equally well by the values {1, 2, 3} or by { 0.5, 1, 10}.
Interval	$new_value = a * old_value + b$ where a and b are constants	Thus, the Fahrenheit and Celsius temperature scales differ in terms of where their zero value is and the size of a unit (degree).
Ratio	$new_value = a * old_value$	Length can be measured in meters or feet.

Discrete and Continuous Attributes

● Discrete Attribute

- Has only a finite or countably infinite set of values
- Examples: zip codes, counts, or the set of words in a collection of documents
- Often represented as integer variables.
- Note: **binary attributes** are a special case of discrete attributes

● Continuous Attribute

- Has real numbers as attribute values
- Examples: temperature, height, or weight.
- Continuous attributes are typically represented as floating-point variables.

Key Messages for Attribute Types

- The types of operations you choose should be “meaningful” for the type of data you have
 - Distinctness, order, meaningful intervals, and meaningful ratios are only four properties of data
 - The data type you see – often numbers or strings – may not capture all the properties or may suggest properties that are not there
 - Analysis may depend on these other properties of the data
 - ◆ Many statistical analyses depend only on the distribution
 - Many times what is meaningful is measured by statistical significance
 - But in the end, what is meaningful is measured by the domain

Types of Data Sets

- Record
 - Data Matrix
 - Document Data
 - Transaction Data
- Graph
 - World Wide Web
 - Molecular Structures
- Ordered
 - Spatial Data
 - Temporal Data
 - Sequential Data
 - Genetic Sequence Data

Important Characteristics of Data

- Dimensionality (number of attributes)
 - ◆ High dimensional data brings a number of challenges
- Sparsity
 - ◆ Only presence counts
- Resolution
 - ◆ Patterns depend on the scale
- Size
 - ◆ Type of analysis may depend on size of data

Record Data

- Data that consists of a collection of records, each of which consists of a fixed set of attributes

<i>Tid</i>	Refund	Marital Status	Taxable Income	Cheat
1	Yes	Single	125K	No
2	No	Married	100K	No
3	No	Single	70K	No
4	Yes	Married	120K	No
5	No	Divorced	95K	Yes
6	No	Married	60K	No
7	Yes	Divorced	220K	No
8	No	Single	85K	Yes
9	No	Married	75K	No
10	No	Single	90K	Yes

Data Matrix

- If data objects have the same fixed set of numeric attributes, then the data objects can be thought of as points in a multi-dimensional space, where each dimension represents a distinct attribute
- Such data set can be represented by an m by n matrix, where there are m rows, one for each object, and n columns, one for each attribute

Projection of x Load	Projection of y load	Distance	Load	Thickness
10.23	5.27	15.22	2.7	1.2
12.65	6.25	16.22	2.2	1.1

Document Data

- Each document becomes a ‘term’ vector
 - Each term is a component (attribute) of the vector
 - The value of each component is the number of times the corresponding term occurs in the document.

	team	coach	play	ball	score	game	win	lost	timeout	season
Document 1	3	0	5	0	2	6	0	2	0	2
Document 2	0	7	0	2	1	0	0	3	0	0
Document 3	0	1	0	0	1	2	2	0	3	0


Transaction Data

- A special type of record data, where
 - Each record (transaction) involves a set of items.
 - For example, consider a grocery store. The set of products purchased by a customer during one shopping trip constitute a transaction, while the individual products that were purchased are the items.

<i>TID</i>	<i>Items</i>
1	Bread, Coke, Milk
2	Beer, Bread
3	Beer, Coke, Diaper, Milk
4	Beer, Bread, Diaper, Milk
5	Coke, Diaper, Milk

Graph Data

- Examples: Generic graph, a molecule, and webpages



Useful Links:

- [Bibliography](#)
- Other Useful Web sites
 - [ACM SIGKDD](#)
 - [KDnuggets](#)
 - [The Data Mine](#)

Knowledge Discovery and Data Mining Bibliography

(Gets updated frequently, so visit often!)

- [Books](#)
- [General Data Mining](#)

Book References in Data Mining and Knowledge Discovery


Usama Fayyad, Gregory Piatetsky-Shapiro, Padhraic Smyth, and Ramasamy uthurasamy, "Advances in Knowledge Discovery and Data Mining", AAAI Press/the MIT Press, 1996.

J. Ross Quinlan, "C4.5: Programs for Machine Learning", Morgan Kaufmann Publishers, 1993.
Michael Berry and Gordon Linoff, "Data Mining Techniques (For Marketing, Sales, and Customer Support)", John Wiley & Sons, 1997.

General Data Mining

Usama Fayyad, "Mining Databases: Towards Algorithms for Knowledge Discovery", Bulletin of the IEEE Computer Society Technical Committee on data Engineering, vol. 21, no. 1, March 1998.

Christopher Matheus, Philip Chan, and Gregory Piatetsky-Shapiro, "Systems for knowledge Discovery in databases", IEEE Transactions on Knowledge and Data Engineering, 5(6):903-913, December 1993.




Benzene Molecule: C₆H₆

Ordered Data

- Sequences of transactions

Items/Events



An element of the
sequence

Ordered Data

- Genomic sequence data


**GGTTCCGCCTTCAGCCCCGCGCC
CGCAGGGGCCCGCCCCGCGCCGTC
GAGAAGGGCCCGCCTGGCGGGCG
GGGGGAGGCGGGGCCCGCCGAGC
CCAACCGAGTCCGACCAGGTGCC
CCCTCTGCTCGGCCTAGACCTGA
GCTCATTAGGCGGCAGCGGACAG
GCCAAGTAGAACACGCGAAGCGC
TGGGCTGCCTGCTGCGACCAGGG**

Ordered Data

- Spatio-Temporal Data

**Average Monthly
Temperature of
land and ocean**

Jan



Data Quality

- Poor data quality negatively affects many data processing efforts
“The most important point is that poor data quality is an unfolding disaster.
 - *Poor data quality costs the typical company at least ten percent (10%) of revenue; twenty percent (20%) is probably a better estimate.”*

Thomas C. Redman, DM Review, August 2004

- Data mining example: a classification model for detecting people who are loan risks is built using poor data
 - Some credit-worthy candidates are denied loans
 - More loans are given to individuals that default


Data Quality ...

- What kinds of data quality problems?
- How can we detect problems with the data?
- What can we do about these problems?

- Examples of data quality problems:
 - Noise and outliers
 - Missing values
 - Duplicate data
 - Wrong data

Outliers

- **Outliers** are data objects with characteristics that are considerably different than most of the other data objects in the data set
 - **Case 1:** Outliers are noise that interferes with data analysis
 - **Case 2:** Outliers are the goal of our analysis
 - ◆ Credit card fraud
 - ◆ Intrusion detection



Missing Values

- Reasons for missing values
 - Information is not collected
(e.g., people decline to give their age and weight)
 - Attributes may not be applicable to all cases
(e.g., annual income is not applicable to children)
- Handling missing values
 - Eliminate data objects or variables
 - Estimate missing values
 - ◆ Example: time series of temperature
 - ◆ Example: census results
 - Ignore the missing value during analysis

Missing Values ...

- Missing completely at random (MCAR)
 - Missingness of a value is independent of attributes
 - Fill in values based on the attribute
 - Analysis may be unbiased overall
- Missing at Random (MAR)
 - Missingness is related to other variables
 - Fill in values based other values
 - Almost always produces a bias in the analysis
- Missing Not at Random (MNAR)
 - Missingness is related to unobserved measurements
 - Informative or non-ignorable missingness
- Not possible to know the situation from the data

Duplicate Data

- Data set may include data objects that are duplicates, or almost duplicates of one another
 - Major issue when merging data from heterogeneous sources
- Examples:
 - Same person with multiple email addresses
- Data cleaning
 - Process of dealing with duplicate data issues

Similarity and Dissimilarity Measures

- Similarity measure
 - Numerical measure of how alike two data objects are.
 - Is higher when objects are more alike.
 - Often falls in the range $[0,1]$
- Dissimilarity measure
 - Numerical measure of how different two data objects are
 - Lower when objects are more alike
 - Minimum dissimilarity is often 0
 - Upper limit varies
- Proximity refers to a similarity or dissimilarity

Similarity/Dissimilarity for Simple Attributes

The following table shows the similarity and dissimilarity between two objects, x and y , with respect to a single, simple attribute.

Attribute Type	Dissimilarity	Similarity
Nominal	$d = \begin{cases} 0 & \text{if } x = y \\ 1 & \text{if } x \neq y \end{cases}$	$s = \begin{cases} 1 & \text{if } x = y \\ 0 & \text{if } x \neq y \end{cases}$
Ordinal	$d = x - y / (n - 1)$ (values mapped to integers 0 to $n-1$, where n is the number of values)	$s = 1 - d$
Interval or Ratio	$d = x - y $	$s = -d, s = \frac{1}{1+d}, s = e^{-d},$ $s = 1 - \frac{d - \min_d}{\max_d - \min_d}$

Euclidean Distance


- Euclidean Distance

$$d(\mathbf{x}, \mathbf{y}) = \sqrt{\sum_{k=1}^n (x_k - y_k)^2}$$

where n is the number of dimensions (attributes) and x_k and y_k are, respectively, the k^{th} attributes (components) or data objects \mathbf{x} and \mathbf{y} .

- Standardization is necessary, if scales differ.

Euclidean Distance



point	x	y
p1	0	2
p2	2	0
p3	3	1
p4	5	1

	p1	p2	p3	p4
p1	0	2.828	3.162	5.099
p2	2.828	0	1.414	3.162
p3	3.162	1.414	0	2
p4	5.099	3.162	2	0

Distance Matrix

Minkowski Distance

- Minkowski Distance is a generalization of Euclidean Distance

$$d(\mathbf{x}, \mathbf{y}) = \left(\sum_{k=1}^n |x_k - y_k|^r \right)^{1/r}$$

Where r is a parameter, n is the number of dimensions (attributes) and x_k and y_k are, respectively, the k^{th} attributes (components) or data objects \mathbf{x} and \mathbf{y} .

Minkowski Distance: Examples

- $r = 1$. City block (Manhattan, taxicab, L_1 norm) distance.
 - A common example of this is the Hamming distance, which is just the number of bits that are different between two binary vectors
- $r = 2$. Euclidean distance
- $r \rightarrow \infty$. “supremum” (L_{\max} norm, L_{∞} norm) distance.
 - This is the maximum difference between any component of the vectors
- Do not confuse r with n , i.e., all these distances are defined for all numbers of dimensions.

Minkowski Distance

point	x	y
p1	0	2
p2	2	0
p3	3	1
p4	5	1

L1	p1	p2	p3	p4
p1	0	4	4	6
p2	4	0	2	4
p3	4	2	0	2
p4	6	4	2	0


L2	p1	p2	p3	p4
p1	0	2.828	3.162	5.099
p2	2.828	0	1.414	3.162
p3	3.162	1.414	0	2
p4	5.099	3.162	2	0

L_{∞}	p1	p2	p3	p4
p1	0	2	3	5
p2	2	0	1	3
p3	3	1	0	2
p4	5	3	2	0

Distance Matrix

Mahalanobis Distance


$$\text{mahalanobis}(\mathbf{x}, \mathbf{y}) = (\mathbf{x} - \mathbf{y})^T \Sigma^{-1} (\mathbf{x} - \mathbf{y})$$



Σ is the covariance matrix

For red points, the Euclidean distance is 14.7, Mahalanobis distance is 6.

Mahalanobis Distance



Covariance Matrix:

$$\Sigma = \begin{bmatrix} 0.3 & 0.2 \\ 0.2 & 0.3 \end{bmatrix}$$

A: (0.5, 0.5)

B: (0, 1)

C: (1.5, 1.5)

Mahal(A,B) = 5

Mahal(A,C) = 4

Common Properties of a Distance

- Distances, such as the Euclidean distance, have some well known properties.
 1. $d(\mathbf{x}, \mathbf{y}) \geq 0$ for all \mathbf{x} and \mathbf{y} and $d(\mathbf{x}, \mathbf{y}) = 0$ only if $\mathbf{x} = \mathbf{y}$. (Positive definiteness)
 2. $d(\mathbf{x}, \mathbf{y}) = d(\mathbf{y}, \mathbf{x})$ for all \mathbf{x} and \mathbf{y} . (Symmetry)
 3. $d(\mathbf{x}, \mathbf{z}) \leq d(\mathbf{x}, \mathbf{y}) + d(\mathbf{y}, \mathbf{z})$ for all points \mathbf{x} , \mathbf{y} , and \mathbf{z} . (Triangle Inequality)

where $d(\mathbf{x}, \mathbf{y})$ is the distance (dissimilarity) between points (data objects), \mathbf{x} and \mathbf{y} .

- A distance that satisfies these properties is a **metric**

Common Properties of a Similarity

- Similarities, also have some well known properties.
 1. $s(\mathbf{x}, \mathbf{y}) = 1$ (or maximum similarity) only if $\mathbf{x} = \mathbf{y}$.
 2. $s(\mathbf{x}, \mathbf{y}) = s(\mathbf{y}, \mathbf{x})$ for all \mathbf{x} and \mathbf{y} . (Symmetry)

where $s(\mathbf{x}, \mathbf{y})$ is the similarity between points (data objects), \mathbf{x} and \mathbf{y} .

Cosine Similarity

- If \mathbf{d}_1 and \mathbf{d}_2 are two vectors, then

$$\cos(\mathbf{d}_1, \mathbf{d}_2) = \langle \mathbf{d}_1, \mathbf{d}_2 \rangle / \|\mathbf{d}_1\| \|\mathbf{d}_2\| ,$$

where $\langle \mathbf{d}_1, \mathbf{d}_2 \rangle$ indicates inner product or vector **dot product** of vectors, \mathbf{d}_1 and \mathbf{d}_2 , and $\|\mathbf{d}\|$ is the length of vector \mathbf{d} .

- Example:

$$\mathbf{d}_1 = (3 \ 2 \ 0 \ 5 \ 0 \ 0 \ 0 \ 2 \ 0 \ 0)$$

$$\mathbf{d}_2 = (1 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 1 \ 0 \ 2)$$

$$\langle \mathbf{d}_1, \mathbf{d}_2 \rangle = 3*1 + 2*0 + 0*0 + 5*0 + 0*0 + 0*0 + 0*0 + 2*1 + 0*0 + 0*2 = 5$$

$$\|\mathbf{d}_1\| = (3*3 + 2*2 + 0*0 + 5*5 + 0*0 + 0*0 + 0*0 + 2*2 + 0*0 + 0*0)^{0.5} = (42)^{0.5} = 6.481$$

$$\|\mathbf{d}_2\| = (1*1 + 0*0 + 0*0 + 0*0 + 0*0 + 0*0 + 0*0 + 1*1 + 0*0 + 2*2)^{0.5} = (6)^{0.5} = 2.449$$

$$\cos(\mathbf{d}_1, \mathbf{d}_2) = 0.3150$$

Correlation measures linear relationship between objects

$$\text{corr}(\mathbf{x}, \mathbf{y}) = \frac{\text{covariance}(\mathbf{x}, \mathbf{y})}{\text{standard_deviation}(\mathbf{x}) * \text{standard_deviation}(\mathbf{y})} = \frac{s_{xy}}{s_x s_y},$$

where we are using the following standard statistical notation and definitions

$$\text{covariance}(\mathbf{x}, \mathbf{y}) = s_{xy} = \frac{1}{n-1} \sum_{k=1}^n (x_k - \bar{x})(y_k - \bar{y})$$


$$\text{standard_deviation}(\mathbf{x}) = s_x = \sqrt{\frac{1}{n-1} \sum_{k=1}^n (x_k - \bar{x})^2}$$

$$\text{standard_deviation}(\mathbf{y}) = s_y = \sqrt{\frac{1}{n-1} \sum_{k=1}^n (y_k - \bar{y})^2}$$

$$\bar{x} = \frac{1}{n} \sum_{k=1}^n x_k \text{ is the mean of } \mathbf{x}$$

$$\bar{y} = \frac{1}{n} \sum_{k=1}^n y_k \text{ is the mean of } \mathbf{y}$$

Visually Evaluating Correlation



Scatter plots showing the similarity from -1 to 1.

Drawback of Correlation

- $\mathbf{x} = (-3, -2, -1, 0, 1, 2, 3)$

- $\mathbf{y} = (9, 4, 1, 0, 1, 4, 9)$

$$y_i = x_i^2$$

- $\text{mean}(\mathbf{x}) = 0, \text{mean}(\mathbf{y}) = 4$

- $\text{std}(\mathbf{x}) = 2.16, \text{std}(\mathbf{y}) = 3.74$


- $\text{corr} = (-3)(5) + (-2)(0) + (-1)(-3) + (0)(-4) + (1)(-3) + (2)(0) + 3(5) / (6 * 2.16 * 3.74)$
 $= 0$

Density

- Measures the degree to which data objects are close to each other in a specified area
- The notion of density is closely related to that of proximity
- Concept of density is typically used for clustering and anomaly detection
- Examples:
 - Euclidean density
 - ◆ Euclidean density = number of points per unit volume
 - Probability density
 - ◆ Estimate what the distribution of the data looks like
 - Graph-based density
 - ◆ Connectivity

Euclidean Density: Grid-based Approach

- Simplest approach is to divide region into a number of rectangular cells of equal volume and define density as # of points the cell contains



0	0	0	0	0	0	0
0	0	0	0	0	0	0
4	17	18	6	0	0	0
14	14	13	13	0	18	27
11	18	10	21	0	24	31
3	20	14	4	0	0	0
0	0	0	0	0	0	0

Euclidean Density: Center-Based

- Euclidean density is the number of points within a specified radius of the point




Illustration of center-based density.