

Why Is the Sky Blue?

A Geometric Derivation of Rayleigh Scattering from Maxwell Theory

Gustaf Ullman

May 24, 2026

Abstract

The usual answer to the question why the sky is blue is that shorter wavelengths of sunlight are scattered more efficiently by the atmosphere than longer wavelengths. This answer is correct, but incomplete: it names the relevant effect without displaying the physical chain that produces it. This paper gives a conceptually explicit derivation of Rayleigh scattering from Maxwell theory, with particular attention to the geometrical structure of the electromagnetic field.

The aim is not pedagogical in the sense of simplification. It is structural: to make visible how a familiar phenomenon follows from the geometrical and dynamical structure of Maxwell theory. The intended reader is assumed to have some background in classical electrodynamics and relativity. The guiding idea is conceptual seeing: the Rayleigh law is not treated as a formula to be remembered, but as a consequence of the structure through which the phenomenon becomes physically intelligible.

The central chain is that an incident electromagnetic field polarizes a small molecule, the induced dipole radiates according to Maxwell's equations, the far-zone radiation field is proportional to the second time derivative of the dipole moment, and the observed intensity is obtained from the electromagnetic stress-energy structure, which is quadratic in the field. For harmonic radiation this gives a scattered field amplitude proportional to ω^2 , and hence an intensity proportional to ω^4 , or equivalently to λ^{-4} .

The derivation is then recast in spacetime algebra, where the electromagnetic field is represented as a bivector F , the electric and magnetic fields are observer-dependent projections of F , and the stress-energy tensor becomes the vector-valued linear map

$$T_F(a) = -\frac{\varepsilon_0}{2} F a F.$$

In this formulation Rayleigh scattering appears as a geometric process: an incident bivector field is projected in the molecular rest frame, induces a transverse oscillating dipole, generates an outgoing radiation bivector, and yields an observed energy flux through the stress-energy map.

1 Introduction

The blue colour of the sky is usually explained by saying that blue light is scattered more strongly than red light in the Earth's atmosphere. This is true, but it is not yet a full physical explanation. It gives the name and qualitative direction of the effect, but it does not yet explain why the scattering strength should scale as the inverse fourth power of wavelength.

The relevant mechanism is Rayleigh scattering. It applies when the scattering object is much smaller than the wavelength of the incident radiation. Air molecules have a typical

size of order 10^{-10} m, while visible light has wavelengths of order $4 \cdot 10^{-7}$ to $7 \cdot 10^{-7}$ m. Thus, for visible light in the atmosphere, the molecular scatterers are very small compared with the wavelength.

The purpose of this paper is not to give an elementary or popular account of the blue sky. The aim is instead structural clarification. By this I mean an explanation that does not merely attach a formula to a phenomenon, but makes visible the chain of physical and geometrical dependencies through which the phenomenon becomes intelligible. The intended reader is assumed to have some familiarity with classical electromagnetism, including Maxwell's equations, radiation from accelerated charges, and the Poynting vector.

In this respect the paper is also a small exercise in conceptual seeing. In a sense close to Hanson's account of theory-laden observation, a physical phenomenon becomes intelligible when it is seen under the right theoretical structure. The blue sky is not fully explained by attaching the phrase "short wavelengths scatter more" to it. It becomes intelligible when the Rayleigh law is seen as the joint consequence of induced molecular polarization, dipole radiation, and the quadratic stress-energy structure of the electromagnetic field.

The role of spacetime algebra is not to make the mathematics superficially easier. Rather, it displays more clearly the geometric unity of the electromagnetic field, observer-dependent field decompositions, and stress-energy flow. In STA the electromagnetic field is not fundamentally a pair $(\mathbf{E}, \mathbf{B})$, but a single spacetime bivector F . Electric and magnetic fields arise only after choosing a timelike direction u , that is, after specifying an observer or physical rest frame.

This also changes how the frequency dependence should be read. The frequency that enters the molecular response is not an absolute property of the wave alone. It is the frequency measured in the rest frame of the scatterer, given geometrically by the projection of the incident null wave vector k onto the molecular four-velocity u :

$$\omega = k \cdot u,$$

up to conventional factors of c . Thus the ω^4 dependence in the Rayleigh law is already tied to a physically realized observer projection.

In the Rayleigh regime the incident electromagnetic field is nearly uniform across each molecule. The field polarizes the molecule, displacing its electron cloud slightly relative to its nuclei. The molecule becomes a small induced electric dipole. Since the incident field oscillates, the induced dipole also oscillates. An oscillating dipole radiates. The scattered light is the radiation field of this driven dipole.

The familiar Rayleigh law is

$$I_{\text{sc}} \propto \frac{1}{\lambda^4}. \quad (1)$$

The derivation has two essential steps.

First, the far radiation field of an oscillating dipole is proportional to the second time derivative of the dipole moment. For harmonic radiation of angular frequency ω , this gives a factor ω^2 in the field amplitude.

Second, intensity is not proportional to field amplitude but to energy flux. Electromagnetic energy flux is quadratic in the field. In conventional vector notation this is expressed by the Poynting vector,

$$\mathbf{S} = \frac{1}{\mu_0} \mathbf{E} \times \mathbf{B}. \quad (2)$$

In relativistic form it is part of the electromagnetic stress-energy tensor. Thus an ω^2 dependence in the radiation field amplitude becomes an ω^4 dependence in intensity.

The blue sky is therefore not merely an elementary optical phenomenon. It is a macroscopic trace of the geometry and dynamics of Maxwell theory.

2 The Rayleigh Regime

Rayleigh scattering applies when the characteristic size a of the scatterer is much smaller than the wavelength λ of the incident radiation:

$$a \ll \lambda. \quad (3)$$

In this limit the electric field of the incident wave is approximately constant over the molecule. The molecule need not be treated as an extended object with appreciable phase differences across it. To leading order it may be treated as a pointlike polarizable system.

For an isotropic molecule, the induced electric dipole moment is proportional to the local electric field:

$$\mathbf{p}(t) = \alpha \mathbf{E}(t), \quad (4)$$

where α is the molecular polarizability. If the incident wave is monochromatic,

$$\mathbf{E}(t) = \mathbf{E}_0 \cos(\omega t), \quad (5)$$

then

$$\mathbf{p}(t) = \mathbf{p}_0 \cos(\omega t), \quad \mathbf{p}_0 = \alpha \mathbf{E}_0. \quad (6)$$

This is the elementary physical model behind Rayleigh scattering: a small polarizable molecule becomes a driven oscillating electric dipole. The derivation does not require the molecule to absorb and re-emit light as a sequence of separate events. In the classical description, the incident field drives a bound charge distribution, and the driven charge distribution radiates.

3 A Bound Electron Model

One can see the origin of the induced dipole more explicitly by modelling a bound electron as a driven oscillator. Let x be the displacement of an electron of charge $-e$ and mass m from its equilibrium position. In a simple classical model,

$$m\ddot{x} + m\gamma\dot{x} + m\omega_0^2 x = -eE_0 \cos(\omega t), \quad (7)$$

where ω_0 is the characteristic binding frequency and γ is a damping coefficient.

For frequencies far below resonance, $\omega \ll \omega_0$, and ignoring damping for simplicity, the displacement is approximately

$$x(t) \approx -\frac{e}{m\omega_0^2} E_0 \cos(\omega t). \quad (8)$$

The dipole moment is $p = -ex$, so

$$p(t) \approx \frac{e^2}{m\omega_0^2} E_0 \cos(\omega t). \quad (9)$$

Thus the polarizability is, in this simple model,

$$\alpha \approx \frac{e^2}{m\omega_0^2}. \quad (10)$$

Real molecular polarizabilities are more complicated, and for a full treatment one must take into account molecular structure, anisotropy, resonances, and quantum effects. None of these complications changes the basic point needed for the Rayleigh law: far from resonance and for scatterers much smaller than the wavelength, the leading response is dipolar and linear in the incident field.

4 Radiation from an Oscillating Dipole

Maxwell's equations imply that accelerated charges radiate. In the far zone, the electric field radiated by a localized time-dependent electric dipole $\mathbf{p}(t)$ is

$$\mathbf{E}_{\text{rad}}(\mathbf{r}, t) = \frac{1}{4\pi\epsilon_0 c^2 r} [\hat{\mathbf{n}} \times (\hat{\mathbf{n}} \times \ddot{\mathbf{p}}(t_r))], \quad (11)$$

where

$$t_r = t - \frac{r}{c} \quad (12)$$

is the retarded time, and $\hat{\mathbf{n}}$ is the unit vector from the dipole to the observer.

The double cross product projects the dipole acceleration onto the plane transverse to the observation direction. This expresses an important physical fact: radiation fields are transverse.

For a harmonic dipole,

$$\mathbf{p}(t) = \mathbf{p}_0 \cos(\omega t), \quad (13)$$

we have

$$\ddot{\mathbf{p}}(t) = -\omega^2 \mathbf{p}_0 \cos(\omega t). \quad (14)$$

Therefore the amplitude of the radiated electric field is proportional to

$$E_{\text{rad}} \propto \omega^2 p_0. \quad (15)$$

Since $p_0 = \alpha E_0$, the scattered field amplitude is proportional to

$$E_{\text{sc}} \propto \omega^2 \alpha E_0. \quad (16)$$

This is the first essential step in the Rayleigh law. The fourth power has not yet appeared. At this stage we have only obtained a second power in frequency at the level of field amplitude.

5 Energy Flux and the Poynting Vector

The observed brightness of light is not proportional to the electric field amplitude itself. It is proportional to electromagnetic energy flux.

In conventional vector notation the electromagnetic energy flux is given by the Poynting vector,

$$\mathbf{S} = \frac{1}{\mu_0} \mathbf{E} \times \mathbf{B}. \quad (17)$$

This expression is not an arbitrary definition of optical intensity. It follows from Maxwell's equations when they are written as a local energy conservation law. The electromagnetic energy density is

$$u_{\text{em}} = \frac{1}{2} \varepsilon_0 E^2 + \frac{1}{2\mu_0} B^2. \quad (18)$$

It is worth emphasizing what this quadratic dependence means physically. The electric field $\mathbf{E}$ is not itself an energy. It has the dimension of force per unit charge. The energy density arises only when the field strength is combined with the electromagnetic constants. In SI units,

$$\varepsilon_0 E^2 \quad (19)$$

has the dimension of energy per unit volume. Thus the statement that field energy is quadratic in the field should not be read as saying that force per charge is itself an energy. Rather, the electromagnetic field carries energy because work must be done to build up the field configuration, and this stored energy is quadratic in the field amplitude.

A simple way to see this is through a parallel-plate capacitor. Its stored energy is

$$U = \frac{1}{2} CV^2. \quad (20)$$

For plates of area A , separation d , and vacuum permittivity ε_0 , one has

$$C = \frac{\varepsilon_0 A}{d}, \quad E = \frac{V}{d}. \quad (21)$$

Hence

$$U = \frac{1}{2} \frac{\varepsilon_0 A}{d} (Ed)^2 = \frac{1}{2} \varepsilon_0 E^2 Ad. \quad (22)$$

Since Ad is the volume between the plates, the energy density is

$$u_E = \frac{U}{Ad} = \frac{1}{2} \varepsilon_0 E^2. \quad (23)$$

This is the same elementary structure as in Hooke's law. For a spring, $F = kx$, and the work required to stretch it from 0 to x is

$$W = \int_0^x kx' dx' = \frac{1}{2} kx^2. \quad (24)$$

Since $F = kx$, the same energy may also be written as

$$W = \frac{F^2}{2k}. \quad (25)$$

The force itself is not an energy, but in a linear system the work required to build up the deformation is quadratic in the amplitude, and may equivalently be written as quadratic in the corresponding force-like quantity. The electric field case has the same structural form: the field is not itself energy, but the work required to build up the field configuration is quadratic in the field strength.

This is the elementary electrostatic origin of the quadratic field-energy term. The magnetic term is analogous. The stress-energy formulation used below is the relativistic completion of this same fact: the electromagnetic field carries not only energy density, but also energy flux, momentum density, and stresses, all organized into a single quadratic object.

Together with the Poynting vector it satisfies Poynting's theorem,

$$\frac{\partial u_{\text{em}}}{\partial t} + \nabla \cdot \mathbf{S} = -\mathbf{J} \cdot \mathbf{E}. \quad (26)$$

The term $-\mathbf{J} \cdot \mathbf{E}$ is the rate at which the field does work on charges. Thus $\mathbf{S}$ is the local flux of electromagnetic energy.

For a plane electromagnetic wave in vacuum,

$$B = \frac{E}{c}, \quad (27)$$

and $\mathbf{E}$, $\mathbf{B}$, and the propagation direction are mutually orthogonal. Hence the magnitude of the Poynting vector is

$$S = \frac{1}{\mu_0} EB = \frac{1}{\mu_0 c} E^2 = \varepsilon_0 c E^2. \quad (28)$$

For a harmonic wave $E(t) = E_0 \cos(\omega t)$, the time-averaged intensity is

$$I = \langle S \rangle = \frac{1}{2} \varepsilon_0 c E_0^2. \quad (29)$$

Thus intensity is quadratic in field amplitude:

$$I \propto E_0^2. \quad (30)$$

Since the scattered field amplitude is proportional to ω^2 , the scattered intensity is proportional to

$$I_{\text{sc}} \propto \omega^4. \quad (31)$$

Using

$$\omega = \frac{2\pi c}{\lambda}, \quad (32)$$

we obtain

$$I_{\text{sc}} \propto \frac{1}{\lambda^4}. \quad (33)$$

The fourth power therefore comes from two distinct facts: the radiation field of a dipole contains a second time derivative, and energy flux is quadratic in the electromagnetic field.

6 Angular Dependence

The dipole radiation formula also gives the angular distribution of the scattered light. Suppose the induced dipole oscillates along a fixed direction. The radiated intensity in a direction making an angle θ with the dipole axis is proportional to

$$I(\theta) \propto \sin^2 \theta. \quad (34)$$

Thus an oscillating electric dipole radiates maximally perpendicular to its axis and does not radiate along its axis.

For unpolarized incident light, averaging over the two independent incident polarizations gives the familiar Rayleigh angular dependence

$$I(\theta) \propto 1 + \cos^2 \theta, \quad (35)$$

where θ is now the scattering angle between the incident and observed directions.

This angular dependence is also responsible for the partial polarization of the blue sky. Scattered skylight is most strongly polarized at approximately 90° from the Sun. This is not an additional phenomenon appended to Rayleigh scattering. It is already contained in the transverse structure of dipole radiation.

7 Rayleigh Scattering Cross Section

A more quantitative statement can be given in terms of the differential scattering cross section. For a small isotropic polarizable particle, the differential cross section for linearly polarized incident light has the form

$$\frac{d\sigma}{d\Omega} = \frac{k^4 |\alpha|^2}{16\pi^2 \varepsilon_0^2} |\mathbf{e}_{\text{sc}} \cdot \mathbf{e}_{\text{in}}|^2, \quad (36)$$

where $k = 2\pi/\lambda$, $\mathbf{e}_{\text{in}}$ is the incident polarization direction, and $\mathbf{e}_{\text{sc}}$ is the allowed transverse polarization direction of the scattered wave.

For unpolarized incident light one obtains

$$\frac{d\sigma}{d\Omega} = \frac{k^4 |\alpha|^2}{32\pi^2 \varepsilon_0^2} (1 + \cos^2 \theta). \quad (37)$$

The total cross section is proportional to

$$\sigma \propto k^4 |\alpha|^2 \propto \frac{1}{\lambda^4}. \quad (38)$$

The fourth power of the inverse wavelength is therefore not an empirical accident. It follows from the combination of dipole radiation and the quadratic energy structure of the electromagnetic field.

8 Spacetime Algebra Formulation

The preceding derivation is standard. It can, however, be formulated more transparently in spacetime algebra. In STA the electromagnetic field is a bivector field F . The conventional electric and magnetic fields are not fundamental independent fields. They arise from a split of F relative to a timelike direction u .

Let u be a future-directed unit timelike vector. In the present problem, u is the physically realized timelike direction along which the molecular response is defined: the four-velocity of the scattering molecule. This choice is not a mere coordinate convention. It specifies the rest frame in which the molecule is polarized and in which its induced dipole moment is defined.

The electric field measured in the frame u is

$$E_u = F \cdot u. \quad (39)$$

The magnetic field is obtained from the complementary projection, schematically

$$B_u \sim -I(F \wedge u), \quad (40)$$

where I is the spacetime pseudoscalar. Precise factors of c depend on the unit convention.

The important point is conceptual. The molecule is not driven by an absolute three-dimensional electric field. It is driven by the electric projection of the electromagnetic bivector in its own rest frame:

$$F_{\text{in}} \mapsto F_{\text{in}} \cdot u. \quad (41)$$

For an isotropic molecule the induced dipole is therefore

$$p = \alpha(F_{\text{in}} \cdot u), \quad (42)$$

with

$$p \cdot u = 0. \quad (43)$$

Thus p is spacelike relative to the molecular four-velocity u . It is the covariant expression of an electric dipole in the molecular rest frame.

For a monochromatic incident wave one may write schematically

$$F_{\text{in}}(x) = F_0 e^{-ik \cdot x}, \quad (44)$$

where k is a null wave vector,

$$k^2 = 0. \quad (45)$$

The frequency entering the molecular response is not an absolute scalar attached to the wave independently of a frame. It is the frequency measured in the molecular rest frame, given by the projection

$$\omega = k \cdot u, \quad (46)$$

up to conventional factors of c . Thus the ω^4 factor in the Rayleigh law is tied to a physically specified projection of the null wave vector onto the timelike direction defined by the scatterer.

The induced dipole then satisfies

$$\ddot{p} = -\omega^2 p, \quad (47)$$

where the dots denote differentiation with respect to the molecular proper time.

The far-zone scattered field is an outgoing radiation bivector. Schematically it has the form

$$F_{\text{sc}} \propto \frac{1}{r} k_{\text{sc}} \wedge \ddot{p}_{\perp}, \quad (48)$$

where k_{sc} is the outgoing null direction and $p_{\perp}$ is the part of the dipole transverse to the outgoing direction. This expression contains the geometrical content of dipole radiation: the outgoing electromagnetic field is a bivector formed from the propagation direction and the transverse dipole acceleration.

Using $\ddot{p} = -\omega^2 p$, we obtain

$$F_{\text{sc}} \propto \frac{\omega^2 \alpha}{r} k_{\text{sc}} \wedge (F_{\text{in}} \cdot u)_{\perp}. \quad (49)$$

This is the Rayleigh amplitude in STA form. It is intentionally written here as a structural expression rather than as a replacement for the full retarded solution. Its role is to display the geometrical origin of the scaling, transversality, and polarization.

9 Stress-Energy in Spacetime Algebra

In STA the electromagnetic stress-energy tensor is naturally written not as a matrix of components but as a vector-valued linear map. With signature $(+ - - -)$, one common convention is

$$T_F(a) = -\frac{\varepsilon_0}{2} F a F, \quad (50)$$

where a is a spacetime vector and the subscript emphasizes that the map is constructed from the field F . If $a = u$ is the observer's time direction, then $T_F(u)$ gives the electromagnetic energy-momentum density measured by that observer.

Relative to u , $T_F(u)$ contains both the energy density and the energy flux. In conventional language these correspond to

$$u_{\text{em}} = \frac{1}{2} \varepsilon_0 E^2 + \frac{1}{2\mu_0} B^2, \quad (51)$$

and

$$\mathbf{S} = \frac{1}{\mu_0} \mathbf{E} \times \mathbf{B}. \quad (52)$$

The Poynting vector is therefore not an additional optical concept. It is the spatial energy-flux part of the relativistic stress-energy structure of the field.

The expression $T_F(a) = -(\varepsilon_0/2) F a F$ is therefore not a remote formal construction. It is the spacetime-algebraic form of the same quadratic field-energy structure already visible in $u_E = (1/2)\varepsilon_0 E^2$, but now extended to the full relativistic energy-momentum content of the electromagnetic field.

For the scattered field, define

$$T_{F_{\text{sc}}}(a) = -\frac{\varepsilon_0}{2} F_{\text{sc}} a F_{\text{sc}}. \quad (53)$$

For an observer with four-velocity u , $T_{F_{\text{sc}}}(u)$ is the energy-momentum density of the scattered radiation measured by that observer. If n is a spatial unit direction in the observer's rest space, so that

$$n \cdot u = 0, \quad (54)$$

then the energy flux through a surface with spatial normal n is obtained, up to sign conventions, by the projection

$$\Phi_n \sim T_{F_{\text{sc}}}(u) \cdot n. \quad (55)$$

This is the spacetime-algebraic expression of the same content carried by the Poynting vector in the observer's three-space.

Since $T_{F_{\text{sc}}}$ is quadratic in F_{sc} , the flux is quadratic in the scattered field amplitude:

$$\Phi_n \propto F_{\text{sc}}^2. \quad (56)$$

But from the dipole radiation step,

$$F_{\text{sc}} \propto \omega^2. \quad (57)$$

Therefore

$$\Phi_n \propto \omega^4 \propto \lambda^{-4}. \quad (58)$$

In this formulation the Rayleigh law follows from a compact chain:

$$F_{\text{in}} \longrightarrow F_{\text{in}} \cdot u \longrightarrow p \longrightarrow \ddot{p} \longrightarrow F_{\text{sc}} \longrightarrow T_{F_{\text{sc}}}(u) \cdot n. \quad (59)$$

That is: incident bivector field, observer-dependent electric projection, induced dipole, dipole acceleration, outgoing radiation bivector, and stress-energy flux.

10 Why the Sky Is Blue

The Sun emits visible light over a broad range of wavelengths. When this light passes through the atmosphere, air molecules scatter it. Since Rayleigh scattering has the wavelength dependence

$$I_{\text{sc}} \propto \lambda^{-4}, \quad (60)$$

shorter visible wavelengths are scattered much more efficiently than longer ones. For example, comparing blue light at 450 nm with red light at 650 nm, the relative scattering strength is approximately

$$\left(\frac{650}{450}\right)^4 \approx 4.4. \quad (61)$$

Thus blue light is scattered several times more efficiently than red light.

When we look away from the Sun, we do not see direct sunlight. We see sunlight that has been scattered by the atmosphere into our line of sight. Since the shorter visible wavelengths are scattered more strongly, the scattered light is enriched in blue. This is why the sky appears blue.

The sky is not violet, even though violet light has a still shorter wavelength, for several reasons. The solar spectrum contains less violet than blue, some short-wavelength light is absorbed in the upper atmosphere, and the human eye is less sensitive to violet. Human colour perception also maps the mixture of short-wavelength scattered light into the perceptual category blue rather than pure violet.

At sunset the same physics explains the red colour of the Sun. When the Sun is near the horizon, its light travels a much longer path through the atmosphere. Along this long path, much of the blue light is scattered out of the direct beam. The light that remains in the direct solar direction is therefore enriched in longer wavelengths and appears yellow, orange, or red.

Atmospheric refraction affects the apparent position and shape of the setting Sun, and can produce subtle dispersive effects near the horizon. But the main reason for the red colour of the setting Sun is not refraction. It is the preferential Rayleigh scattering of shorter wavelengths out of the direct solar beam.

11 Discussion: Structural Clarification and Conceptual Seeing

This paper is not pedagogical by reducing the phenomenon to a simple image or an easily memorized rule. Rather, it is clarifying by refusing to stop at the rule. Its aim is to expose the chain of dependencies that connects a common optical phenomenon to the geometrical structure of electromagnetic theory.

The common explanation of the blue sky is correct but compressed. It says that blue light is scattered more strongly, but it often hides the physical chain behind this statement. A full explanation requires several layers of theory.

First, light is an electromagnetic field. Second, a small molecule in the field is polarized and becomes an induced dipole. Third, the oscillating dipole radiates because accelerated charges radiate according to Maxwell’s equations. Fourth, the far-zone radiation field is proportional to the second derivative of the dipole moment. Fifth, electromagnetic intensity is not linear in the field but quadratic, because it is part of the stress-energy structure of the field. These facts together imply the fourth-power wavelength dependence of Rayleigh scattering.

This is what is meant here by conceptual seeing. The point is not merely to know that $I_{\text{sc}} \propto \lambda^{-4}$, but to see the formula as an expression of a structure. In this respect the derivation is close in spirit to Hanson’s claim that seeing in science is theory-laden. The relevant theoretical structure does not merely decorate an otherwise neutral observation. It allows the phenomenon to be seen as what it is: not simply a blue colour overhead, but a macroscopic appearance of dipole radiation and electromagnetic stress-energy flow.

Spacetime algebra strengthens this point. The electromagnetic field is a single bivector F . Electric and magnetic fields are projections relative to a chosen timelike direction u . The molecule responds to $F \cdot u$, the electric field in its own rest frame. The frequency entering the response is $k \cdot u$, the projection of the incident null wave vector onto that same timelike direction. The scattered radiation is an outgoing bivector determined by the transverse dipole acceleration. The intensity is obtained from the stress-energy map

$$T_{F_{\text{sc}}}(a) = -\frac{\varepsilon_0}{2} F_{\text{sc}} a F_{\text{sc}}.$$

Thus Rayleigh scattering is not merely an optical formula, but a direct consequence of the geometry and dynamics of the electromagnetic field.

This kind of explanation is not elementary, but it is clarifying. It is meant for a reader who already has enough physics to recognize the ingredients, but who wants to see why they belong together. In that sense, the question “Why is the sky blue?” becomes a test case for a deeper conception of understanding in physics: not the ability to quote a formula, but the ability to see how the formula follows from the structure of the theory.

12 Conclusion

The blue colour of the sky is a visible consequence of Rayleigh scattering. But Rayleigh scattering itself is not a primitive fact. It follows from the interaction between an incident electromagnetic field and small polarizable molecules. In the Rayleigh regime, each molecule behaves as a driven electric dipole. The far radiation field of this dipole is proportional to ω^2 , while the observed intensity is quadratic in the field. The result is

$$I_{\text{sc}} \propto \omega^4 \propto \frac{1}{\lambda^4}. \quad (62)$$

In spacetime algebra this derivation takes a compact geometric form:

$$F_{\text{in}} \rightarrow F_{\text{in}} \cdot u \rightarrow p \rightarrow \ddot{p} \rightarrow F_{\text{sc}} \rightarrow T_{F_{\text{sc}}}(u) \cdot n. \quad (63)$$

The sky is blue because the electromagnetic field, when projected into the rest frames of atmospheric molecules, drives small dipoles whose radiation field and stress-energy structure favour shorter wavelengths. The everyday appearance of the sky is therefore a macroscopic trace of the geometric structure of Maxwell theory.

References

- [1] Lord Rayleigh, “On the light from the sky, its polarization and colour,” *Philosophical Magazine*, vol. 41, pp. 107–120, 1871.
- [2] J. D. Jackson, *Classical Electrodynamics*, 3rd ed., Wiley, 1999.
- [3] D. J. Griffiths, *Introduction to Electrodynamics*, 4th ed., Pearson, 2013.
- [4] C. F. Bohren and D. R. Huffman, *Absorption and Scattering of Light by Small Particles*, Wiley, 1983.
- [5] M. Born and E. Wolf, *Principles of Optics*, 7th ed., Cambridge University Press, 1999.
- [6] D. Hestenes, *Space-Time Algebra*, Gordon and Breach, 1966.
- [7] C. Doran and A. Lasenby, *Geometric Algebra for Physicists*, Cambridge University Press, 2003.
- [8] W. E. Baylis, *Electrodynamics: A Modern Geometric Approach*, Birkhäuser, 1999.
- [9] N. R. Hanson, *Patterns of Discovery: An Inquiry into the Conceptual Foundations of Science*, Cambridge University Press, 1958.
- [10] T. S. Kuhn, *The Structure of Scientific Revolutions*, University of Chicago Press, 1962.