

Theorems from Brezis' *Functional Analysis, Sobolev Spaces and Partial
Differential Equations* (first edition)

Gustaf Bjurstam

bjurstam@kth.se

1 The Hahn-Banach Theorems. Introduction to Conjugate Convex Functions

Theorem 1.1 (Hahn-Banach). *Let E be a vector space over \mathbb{R} , and let $p : E \rightarrow \mathbb{R}$ be a Minkowski functional. Let G be a linear subspace of E and let $g : G \rightarrow \mathbb{R}$ be a linear functional such that $g(x) \leq p(x)$ for all $x \in G$. There exists a linear functional $f : E \rightarrow \mathbb{R}$ such that*

$$f(x) = g(x), \forall x \in G,$$

and

$$f(x) \leq p(x), \forall x \in E.$$

Lemma 1.1 (Zorn). *Every nonempty ordered set that is inductive has a maximal element.*

Corollary 1.2. *Let G be a linear subspace of the real vector space E . If $g \in G^*$, then there exists $f \in E^*$ that extends g and such that $\|f\|_{E^*} = \|g\|_{G^*}$.*

Corollary 1.3. *Let E be a normed real vector space. For every $x \in E$ there is $f \in E^*$ such that $\|f\| = \|x\|$ and $\langle f, x \rangle = \|x\|^2$.*

Corollary 1.4. *For every x in the real normed vector space E we have*

$$\|x\| = \sup_{\substack{f \in E^* \\ \|f\| \leq 1}} |\langle f, x \rangle| = \max_{\substack{f \in E^* \\ \|f\| \leq 1}} |\langle f, x \rangle|.$$

Proposition 1.5. *Let E be a real normed vector space, and let $H = [f = \alpha] \subset E$ be an affine hyperplane. Then H is closed if and only if f is continuous.*