Semiclassical scattering of Boussinesq solitons: symplectic structure, topological phases and integrable defects

Gustav Delius*

Second draft, 4 November 2026

Sections 7–9 are placeholders; Section 10 will be revised.

Abstract

We study the two-body scattering of solitons of the Boussinesq equation at the semiclassical level. The main thread is the well-posed ("good") equation $u_{tt} = u_{xx} - u_{xxxx} - 3(u^2)_{xx}$; the ill-posed ("bad") equation $v_{tt} = v_{xx} + v_{xxxx} + 3(v^2)_{xx}$ is treated in parallel through an explicit dictionary. Unlike sine-Gordon or affine Toda solitons, Boussinesq solitons have no boost symmetry, move through a gapless phonon background, and carry a continuous topological charge $M = \int u\,dx$. We show that two-soliton scattering is an exactly symplectic map of the canonical phase space, provided one includes ordering-dependent cross terms $d\Pi_i \wedge dM_j$ between the shift charge of one soliton and the topological charge of another. Darboux coordinates then give a closed-form semiclassical phase,

$$\delta(1,2) = \tfrac{2}{9}(\ell_1 - \ell_2)\log A_{12} + \tfrac{2}{3}M_1 M_2(v_1 - v_2),$$

where ℓ is the soliton's spectral parameter, A_{12} the Hirota interaction constant and v_j the velocities. The phase is antisymmetric and obeys a crossing-type relation. For the bad equation, the coefficient of $\log A_{12}$ continued to breather kinematics equals the classical breather adiabatic invariant of Bogdan and Charkina, so their semiclassical Bohr–Sommerfeld prescription becomes a string condition in ℓ; and the classical limit of the quantum Toda Bethe ansatz reproduces δ, including its topological term, in the small-amplitude limit. For the good equation with Corrigan–Zambon integrable defects, the soliton–defect transmission phase has the same structure, the defect adds no cross terms, two-soliton–defect scattering factorises, numerical mode searches find a localised growing mode for $\eta\sigma > -1$ and none for $\eta\sigma < -1$ (leaving one strictly stable vacuum in each KP triple with three distinct real vacua), and the displayed zero-constant tree-level phases obey an exact oriented fusion identity. We also record two structural results: the companion-matrix Lax connection is non-ultralocal, and A_{12} factorises into $\mathfrak{sl}_3$ affine Toda blocks of $\beta_1 \pm \beta_2$. One-loop corrections, the quantum Toda Fermi-edge limit and exact S-matrices are not settled; their status is summarised as placeholders.

Contents

1 Introduction **2**

1.1 The problem 2

1.2 Why Boussinesq differs from KdV and sine-Gordon 3

1.3 Prior work 3

1.4 Summary of results 4

*Draft prepared within the research project *quantum_boussinesq_solitons*. Authorship, affiliations and acknowledgements are to be settled by the PI.

2 Conventions and the classical soliton sector **5**
2.1 The two equations and their canonical structure . . . 5
2.2 One-soliton solutions and their charges . . . 6
2.3 KP roots, chords and the spectral parameter . . . 7
2.4 Two-soliton solutions and phase shifts . . . 8
2.5 Breathers . . . 9
2.6 The good/bad dictionary . . . 9

3 Canonical structure and exact symplecticity of scattering **10**
3.1 Cross terms in the symplectic form . . . 10
3.2 Scattering is symplectic only with the cross terms . . . 11
3.3 Darboux coordinates . . . 11

4 The semiclassical two-soliton phase **11**
4.1 The phase . . . 11
4.2 Properties . . . 12
4.3 Bad equation: breathers as spectral-parameter strings . . . 13
4.4 Bad equation: the classical limit of the quantum Toda Bethe ansatz . . . 13

5 Integrable defects at tree level **15**
5.1 Corrigan–Zambon defects . . . 15
5.2 Symplectic structure with a defect . . . 16
5.3 The transmission phase . . . 16
5.4 Two solitons and a defect . . . 16
5.5 Vacua, absorption and linear stability . . . 17
5.6 Tree-level fusion . . . 17
5.7 What is not covered . . . 18

6 Structural results on quantum inverse scattering and affine Toda **18**
6.1 Non-ultralocality . . . 18
6.2 $\mathfrak{sl}_3$ factorisation of the interaction constant . . . 19
6.3 Obstructions . . . 19

7 One-loop corrections **20**

8 The quantum Toda Fermi-edge limit **20**

9 Exact S-matrix candidates and constraints **21**

10 Open problems **21**

A Claims and verification **22**

B Inconsistencies found and their resolution **25**

1 Introduction

1.1 The problem

The quantum theory of solitons in integrable relativistic field theories is well developed. The semiclassical scattering phase of two solitons follows from their classical time delays **(a)** [16], and in sine-Gordon and the affine Toda theories exact soliton S-matrices have been proposed and tested against unitarity, crossing, Yang–Baxter and bootstrap requirements **(b)** [23, 24, 17]. The

Boussinesq equation is the simplest integrable bidirectional wave equation and has N-soliton solutions **(a)** [20] (derived there for the bad equation and extending to the good equation by reversing the fourth-derivative sign **(a)**/**(c)**). Its quantum soliton sector is much less understood. A systematic search of the literature found no two-soliton S-matrix, or non-relativistic two-body scattering phase, for either sign of the dispersion **(c)** (an observation about the searched literature, not a theorem of nonexistence; search coverage and caveats in the project's literature map, `phase0/literature_map.md`). Corrigan and Zambon note that the Boussinesq S-matrix deserves further study because of classical solutions that do not conserve the number of solitons **(b)** [12, §6]. The only earlier quantisation of a Boussinesq soliton found in that search is the one-body semiclassical Bohr–Sommerfeld quantisation of the breather of the bad equation by Bogdan and Charkina **(a)**/**(b)** [4].

This paper consolidates what can be established about two-body scattering of Boussinesq solitons at the semiclassical order $O(\hbar^{-1})$ in the phase, and states what remains open at higher orders.

1.2 Why Boussinesq differs from KdV and sine-Gordon

- *No boost symmetry.* The two-soliton interaction constant depends separately on $\alpha_1+\alpha_2$ and $\alpha_1-\alpha_2$ (Section 2.4), with $\partial_{\alpha_1}\partial_{\alpha_2}\log A_{12}\neq 0$ **(c)**. A soliton S-matrix is therefore a genuine function of two momenta; in the rapidities α_j (or β_j) it does not reduce to a function of their difference alone **(c)**, and the usual relativistic boost argument for difference-form S-matrices and standard crossing does not apply.

- *Gapless phonons.* The linearised good equation has $\omega^2 = q^2 + q^4$. Solitons of the good equation are subsonic, those of the bad equation supersonic **(a)**/**(c)**. There is no mass gap, so the relativistic one-loop machinery does not apply unchanged.

- *A continuous topological charge.* In the potential ϕ with $u = \phi_x$, a soliton is a kink of height $M = 2k$, which varies with the soliton's parameter **(c)**. In sine-Gordon the kink height is fixed. This difference produces the cross terms that are central to Section 3.

- *Number-changing classical solutions.* The good equation has resonant (Y-shaped) solutions in which two solitons merge into one, virtual solitons in head-on collisions, and singular two-soliton solutions in part of parameter space **(a)** [21, 26, 31, 15]. Corrigan–Zambon defects can reflect solitons or decay into them **(a)** [12]. None of this occurs for KdV or sine-Gordon. Purely elastic, factorised quantum scattering is therefore not guaranteed **(b)** [12].

- *Energy unbounded below; two real forms.* In both forms the Hamiltonian is unbounded below **(c)**. The good and bad equations are the same complexified flow with real or imaginary time (Section 2). The good equation is locally well posed in Sobolev spaces, with nonlinearly stable solitary waves for a range of speeds and global smooth solutions for initial data close to a stable solitary wave **(a)** [5], and carries the Corrigan–Zambon defects **(a)** [12]. The bad equation is Hadamard ill-posed **(a)** [9], but on Schwartz-class data whose reflection coefficient vanishes on the unstable spectral set it has a global Riemann–Hilbert solution theory and an extension to solitons and breathers **(a)** [9, 10, 11], regular breathers **(a)** [10, 4], and it is the small-amplitude continuum limit of the Toda lattice **(a)** [36].

1.3 Prior work

Classical theory. The Lax pair is due to Zakharov [36]; the direct and inverse scattering theory is in [13] and, with solitons and breathers for the bad equation, in the work of Charlier and Lenells [9, 10, 11]; their good-equation analysis [8] is solitonless **(a)**. Hirota's bilinear method gives the N-soliton solutions of the bad equation **(a)** [20], and with opposite fourth-derivative

sign those of the good equation **(a)**/**(c)**. Soliton resonances and merging solitons are due to Hirota and Ito [21], Lambert, Musette and Kesteloot [26], and Rasin and Schiff [31]; the matrix good equation and its tropical limit, including virtual solitons, are treated by Dimakis, Müller-Hoissen and Chen [15]. Bonnemain and Doyon develop the generalised hydrodynamics of the good soliton gas [6] and observe that, unlike KdV, Boussinesq distinguishes between candidate choices of soliton "momentum" **(b)**.

Quantum theory. One established construction of quantum Boussinesq charges identifies them with the commuting charges of W_3 conformal field theory **(a)** [25, 7, 3, 28, 1, 2]; these live on a circle in highest-weight modules and have no asymptotic particle states. Hikami constructs a Baxter Q-operator for a periodic quantum discrete Boussinesq chain **(a)** [19], without a particle sector. The quantum Toda chain is exactly solvable **(a)** [18, 33], its asymptotic Bethe ansatz is due to Sutherland and Mertens [34, 29], and its particle excitations become classical solitons **(b)** [29, 32, 30]. Pustilnik and Matveev analyse the classical-to-quantum crossover of KdV solitons in quantum Toda and Lieb–Liniger **(b)** [30]. Soliton S-matrices of imaginary-coupling $a_n^{(1)}$ affine Toda theories are built from $U_q(\widehat{\mathfrak{sl}}_{n+1})$ intertwiners **(b)** [22, 23, 24, 17], with the caveat that $a_2^{(1)}$ soliton–breather scattering is flawed **(b)** [35]. Bogdan and Charkina compute the classical breather integrals and adiabatic invariant of the bad equation **(a)** and quantise it at large quantum number by the Bohr–Sommerfeld prescription **(b)** [4]. Corrigan and Zambon construct integrable type-I defects for the good equation **(a)** [12].

1.4 Summary of results

All results below are at the semiclassical order and are labelled by the equation they hold for.

1. [good & bad] With the potential normalised at $x \to -\infty$, the canonical symplectic form on well-separated solitons contains ordering-dependent cross terms $d\Pi_i \wedge dM_j$. Two-soliton scattering preserves it exactly, while without the cross terms the scattering map is not symplectic (Propositions 3.1–3.2) **(c)**.

2. [good & bad] The semiclassical phase is $\delta(1,2) = \frac{2}{9}(\ell_1 - \ell_2)\log A_{12} + \frac{2}{3}M_1M_2(v_1 - v_2)$ in both orientations (Proposition 4.1) **(c)**. It is antisymmetric, obeys a crossing-type relation, and stays finite on the resonance lines (Proposition 4.2) **(c)**.

3. [bad] At breather kinematics, $i \cdot \frac{2}{9}(\ell_1 - \ell_2)$ equals the Bogdan–Charkina classical adiabatic invariant I_{br} **(a)** exactly **(c)**, so under their semiclassical Bohr–Sommerfeld rule $I_{\mathrm{br}} = N\hbar$ **(b)** breather quantisation becomes the string condition $\ell_1 - \ell_2 = -\frac{9i}{2}N\hbar$ (Proposition 4.3) **(c)**.

4. [Toda → bad] The classical limit of the quantum Toda Bethe ansatz reproduces the lattice Toda soliton sector with its cross terms, and its label-frame phase tends to $3b\,\delta_{\mathrm{bad}}$ in the small-amplitude Boussinesq continuum limit for both orientations (Propositions 4.4–4.6) **(c)**.

5. [good] For Corrigan–Zambon defects: the transmission phase δ_D has the form of δ with the defect as a particle of spectral parameter $\ell(q)$; there are no defect cross terms; two-soliton–defect scattering is symplectic and factorises; numerical searches find a localised growing mode for $\eta\sigma > -1$ and none for $\eta\sigma < -1$ (**(c)** for the sampled search, **(d)** for completeness), so for $|s| < 2$, $s \neq \pm 1$ where the KP triple has three distinct real vacua, one vacuum per triple is strictly linearly stable **(c)**; and the displayed zero-constant tree-level phases obey an exact oriented fusion identity (Section 5) **(c)**.

6. The companion-matrix Lax connection is non-ultralocal ([good], and [bad] via the dictionary) **(c)**; A_{12} is a ratio of $\mathfrak{sl}_3$ affine Toda and sine-Gordon blocks in $\beta_1 + \beta_2$ and $\beta_1 - \beta_2$ (or

$\alpha_1 \pm \alpha_2$) ([good & bad]) **(c)**; and $\partial_S \partial_D \delta \neq 0$ in the rapidities, so no S-matrix of the product form $F(\beta_1 - \beta_2)G(\beta_1 + \beta_2)$ (or $F(\alpha_1 - \alpha_2)G(\alpha_1 + \alpha_2)$) has the semiclassical limit $e^{i\delta/\hbar}$ ([good & bad]; Section 6) **(c)**.

We do *not* claim a one-loop S-matrix, an exact S-matrix, or a quantum matching to the Toda chain beyond its classical limit (Sections 7–9).

Labels and verification. Every nontrivial claim carries a label: **(a)** proved in the literature; **(b)** conjectured or heuristic in the literature; **(c)** derived in this project; **(d)** a guess. Each **(c)** formula is checked by Python (SymPy/mpmath) scripts in the project repository, named in the footnotes and listed in Appendix A with their exact test scope (distinguishing asserted checks from printed-only diagnostics and finite rational grids from symbolic proofs, and supplemented by `paper/scripts/paper_conventions_check.py`, `paper/scripts/review_draft1_checks.py` and `paper/scripts/paper_draft2_checks.py`). Inconsistencies found between the project's write-ups and their verified resolutions are recorded in Appendix B.

2 Conventions and the classical soliton sector

2.1 The two equations and their canonical structure

The *good* Boussinesq equation, its potential and its Hirota form are

$$u_{tt} = u_{xx} - u_{xxxx} - 3(u^2)_{xx}, \qquad u = \phi_x = 2(\log\tau)_{xx}, \qquad (D_t^2 - D_x^2 + D_x^4)\,\tau\cdot\tau = 0. \tag{1}$$

The *bad* equation is

$$v_{tt} = v_{xx} + v_{xxxx} + 3(v^2)_{xx}, \qquad v = \phi_x = 2(\log\tau)_{xx}, \qquad (D_t^2 - D_x^2 - D_x^4)\,\tau\cdot\tau = 0. \tag{2}$$

The sign of the nonlinear terms is fixed so that solitons are positive **(c)**.[1] Here $v(x,t)$ in (2) denotes the bad-equation field, whereas v in (6)–(8) below denotes the soliton velocity. The equations are the Euler–Lagrange equations of

$$\mathcal{L}_{\text{good}} = \tfrac{1}{2}\phi_t^2 - \tfrac{1}{2}\phi_x^2 + \phi_x^3 - \tfrac{1}{2}\phi_{xx}^2, \qquad \mathcal{L}_{\text{bad}} = \tfrac{1}{2}\phi_t^2 - \tfrac{1}{2}\phi_x^2 - \phi_x^3 + \tfrac{1}{2}\phi_{xx}^2 \tag{3}$$

(c).[2] The canonical momentum is $\pi = \phi_t$, the symplectic form is

$$\Omega = \int_{-\infty}^{\infty} d\pi \wedge d\phi\, dx, \tag{4}$$

and $\hbar$ always refers to this normalisation. The conserved quantities used below are

$$M = \int u\,dx, \qquad \Pi = \int \pi\,dx, \qquad P = -\int \pi\,\phi_x\,dx, \qquad E = \int \left(\tfrac{1}{2}\pi^2 + \tfrac{1}{2}\phi_x^2 \mp \phi_x^3 \pm \tfrac{1}{2}\phi_{xx}^2\right)dx, \tag{5}$$

with the upper signs for the good equation. M is the topological charge (the height of the kink in ϕ), Π is the charge of the shift symmetry $\phi \to \phi + c$, and P generates translations **(c)**. Both Hamiltonians are unbounded below: the cubic term does this in both forms, and in the bad form the quadratic term $-\frac{1}{2}\phi_{xx}^2$ already does **(c)** (elementary).

[1] Verification: `phase0/checks/convention_check.py` (asserts symbolic sign residuals and prints the nonlinear signs, without the final `ALL CHECKS PASSED` marker), `phase1/scripts/hirota_solitons.py`, `paper/scripts/paper_conventions_check.py` (checks 1–2).

[2] Verification: `phase1/scripts/charges.py` (good); `paper/scripts/paper_conventions_check.py` (check 1, both). The sign of ϕ_x^3 in $\mathcal{L}_{\text{bad}}$ differs from the master report; see Appendix B, item 1..

Linear waves. Phonons $e^{i(qx-\omega t)}$ obey $\omega^2 = q^2 + q^4$ (good) and $\omega^2 = q^2 - q^4$ (bad). In the bad equation modes with $|q| > 1$ grow at an unbounded rate, so its Cauchy problem is ill-posed in Sobolev spaces **(a)** [6, 9] (where [6] uses the good-equation field $u_{\mathrm{BD}} = u/2$ and [9] uses the bad-equation field $u_{\mathrm{CL}} = 3v$ **(c)**).[3] Charlier and Lenells nevertheless prove global existence for Schwartz-class initial data (under generic non-singularity and spectral hypotheses) whose reflection coefficient vanishes on the unstable spectral set, and extend the inverse scattering formalism to solitons and breathers **(a)** [9, Thms 2.8–2.12], [10]. For the good equation, Bona and Sachs prove local well-posedness in Sobolev spaces $H^{s+2}(\mathbb{R}) \times H^s(\mathbb{R})$ $(s > \frac{1}{2})$, nonlinear stability of solitary waves for a range of speeds, and global existence of smooth solutions for initial data lying relatively close to a stable solitary wave **(a)** [5].

Real and imaginary time. Let $L = \partial_x^3 + q_L\partial_x + r_L$ and $M_L = (L^{2/3})_+ = \partial_x^2 + \frac{2}{3}q_L$. Then $L_\tau = [M_L, L]$ implies $q_{L,\tau\tau} = -\frac{1}{3}q_{L,xxxx} - \frac{2}{3}(q_L^2)_{xx}$ **(a)** [36]/**(c)**. With $q_L = \frac{3}{2}u - \frac{1}{4}$ and $\tau = \sqrt{3}\,t$ this is the good equation; with $q_L = \frac{3}{2}v + \frac{1}{4}$ and $\tau = i\sqrt{3}\,t$ it is the bad one **(c)**.[4] Both are therefore the same complexified flow on third-order operators; they differ in the real form of time and in the background q_L, which generates the sound term. The chiral W_3 literature [3] never has to make this choice.

2.2 One-soliton solutions and their charges

The one-soliton is

$$\tau = 1 + e^{\theta}, \qquad \theta = kx - \omega t + x_0, \qquad \omega = kv, \qquad u = \frac{k^2}{2}\operatorname{sech}^2\frac{\theta}{2}, \tag{6}$$

with direction $\epsilon = \pm 1$ and the uniformisations

$$\text{good: } k = \sin\alpha, \quad v = \epsilon\cos\alpha, \quad \alpha \in (0, \tfrac{\pi}{2}); \qquad \text{bad: } k = \sinh\beta, \quad v = \epsilon\cosh\beta, \quad \beta > 0. \tag{7}$$

Good solitons are subsonic, and larger ones are slower; bad solitons are supersonic **(a)**/**(c)**.[5] The charges (5) on a soliton are

$$M = 2k, \qquad \Pi = -Mv, \qquad P = \tfrac{2}{3}k^3 v, \qquad E = \tfrac{2}{15}k^3(5 \mp 4k^2), \qquad \frac{dE}{dP} = v, \tag{8}$$

(c).[6] The bad-equation values of E and P agree with Bogdan and Charkina under the sign dictionary $\varphi_{\mathrm{BC}} = -\phi$, $w_{\mathrm{BC}} = -v$, which leaves the canonical symplectic form, momentum and energy normalisations unchanged **(a)** [4, (14),(16),(18)]/**(c)**,[7] and the good ones with Corrigan and Zambon after the map of Section 5 **(a)**/**(c)**. The derivatives

$$\frac{dP}{d\alpha} = \tfrac{2}{3}\epsilon\sin\alpha\sin 3\alpha \quad \text{(good)}, \qquad \frac{dP}{d\beta} = \tfrac{2}{3}\epsilon\sinh\beta\sinh 3\beta \quad \text{(bad)} \tag{9}$$

recur below **(c)**.[8]

[3] Verification: `paper/scripts/review_draft1_checks.py`; `paper/scripts/paper_draft2_checks.py` (check 9).

[4] Verification: `phase1/scripts/lax_zero_curvature.py`.

[5] Verification: `phase1/scripts/hirota_solitons.py`; cf. [6, (18)], [12, (2.8)], [4, (11)].

[6] Verification: `phase1/scripts/charges.py`; `paper/scripts/paper_conventions_check.py` (check 3, 40-digit quadrature for both equations and both directions).

[7] Verification: `paper/scripts/review_draft1_checks.py`; `paper/scripts/paper_draft2_checks.py` (check 9).

[8] Verification: `phase2/scripts/semiclassical_phase.py`; `paper/scripts/paper_conventions_check.py` (check 9).

Proposition 2.1 ([good], **(c)**). *The good one-soliton momentum is bounded,* $|P| \le \sqrt{3}/8$*, with the maximum at the cusp* $\alpha = \pi/3$ *(*$k = \sqrt{3}/2$*,* $|v| = \frac{1}{2}$*). The light branch* $\alpha < \pi/3$ *is KdV-like; the heavy branch* $\alpha > \pi/3$ *is a non-relativistic particle with* $E = \frac{2}{15} + \frac{3}{4}P^2 + \dots$*. With physical centre* $X = -x_0/k$ *at* $t = 0$*, the canonical form* (4) *pulled back to the one-soliton family is* $dP \wedge dX = -(P'(k)/k)\, dk \wedge dx_0$ *and degenerates at the cusp.*[9]

For the bad equation E increases strictly with $|P|$ on each direction branch $\epsilon = \pm 1$ **(a)** [4]. Figure 1 (right) shows the good dispersion relation.

2.3 KP roots, chords and the spectral parameter

Good equation. Attach to a soliton the two numbers

$$p_a = \tfrac{1}{2}\Big(k + \frac{v}{\sqrt{3}}\Big), \qquad p_b = \tfrac{1}{2}\Big(\frac{v}{\sqrt{3}} - k\Big), \qquad p_a - p_b = k, \qquad p_a^2 + p_a p_b + p_b^2 = \tfrac{1}{4}. \tag{10}$$

They are two roots of the cubic

$$\ell(p) \equiv -3\sqrt{3}\, p\,(1 - 4p^2) = \ell, \qquad \ell = -\epsilon \cos 3\alpha, \tag{11}$$

whose third root is $p_c = -p_a - p_b = -v/\sqrt{3}$ **(c)**.[10] We call ℓ the *spectral parameter*: with the Lax normalisation above, $e^{\mu x}$ is an eigenfunction of L on the vacuum with eigenvalue $\lambda = \mu^3 - \mu/4$, the soliton corresponds to the ordered pair of sheets $\mu_1 = -p_b$, $\mu_2 = -p_a$ ($\mu_1 - \mu_2 = k$, $\mu_1 + \mu_2 = -v/\sqrt{3}$), and $\lambda = -\ell/(12\sqrt{3})$ **(c)**.[11] In the KP picture a soliton is a line soliton labelled by two roots **(a)** [15], so a Boussinesq soliton is a *chord* (p_a, p_b) of a KP triple $\{p_a, p_b, p_c\}$. The three chords of one triple are three solitons with the same ℓ; they are the three legs of a resonant Y-junction, and Rasin and Schiff's "conservation of speed" $\lambda_1 + \lambda_2 + \lambda_3 = 0$ **(a)** [31, §3] is the statement that they use the three roots of one cubic **(c)** (identification).

Bad equation. The corresponding roots are complex, $(p_a, p_b) = (\mu_a, \mu_b) = \frac{1}{2}(iv/\sqrt{3} \pm k)$, roots of $\mu^3 + \frac{1}{4}\mu = \frac{i}{12\sqrt{3}}\ell$ with $\ell = \epsilon \cosh 3\beta$ **(c)**.[12] For the bad equation, ℓ is proportional to the spectral parameter in the uniformisation $\lambda = \frac{1}{2}(k^3 + k^{-3})$ of [9], with k^3 in the role of $e^{3\beta}$ **(c)**.

Gelfand–Dickey charges and missing spins. [good] On the nonzero Lax background $L_0 = \partial_x^3 - \frac{1}{4}\partial_x$, the raw residue $\operatorname{Res} L^{n/3}$ has a constant vacuum value for odd n ($\operatorname{Res} L_0^{1/3} = -\frac{1}{12}$, $\operatorname{Res} L_0^{5/3} = \frac{5}{5184}$) **(c)**, so the Gelfand–Dickey hierarchy charges are defined by the vacuum-subtracted integrals $H_n = \int (\operatorname{Res} L^{n/3} - \operatorname{Res} L_0^{n/3})\, dx$ **(c)**. They vanish identically for $n \equiv 0 \pmod 3$ because L^m is a differential operator **(a)** [14] (verified for $n \le 6$ **(c)**).[13] On a one-soliton with ordered roots $\mu_1 = -p_b$, $\mu_2 = -p_a$ ($\mu_1 - \mu_2 = k$, $\mu_1 + \mu_2 = -v/\sqrt{3}$) and $\Delta_m = \mu_1^m - \mu_2^m$, the lowest nontrivial vacuum-subtracted charges are $H_1 = \Delta_1$, $H_2 = \Delta_2$, $H_4 = \Delta_4 - \frac{1}{3}\Delta_2$, $H_5 = \Delta_5 - \frac{5}{72}\Delta_1$, and the physical charges are $P = -2\sqrt{3}\, H_4 - \frac{1}{\sqrt{3}} H_2$ and $E = \frac{24}{5} H_5 + \frac{1}{6} H_1$ **(c)**.[14] Because μ_j obey the cubic $\mu^3 = \frac{1}{4}\mu + \lambda$, $\Delta_3 = \frac{1}{4}\Delta_1$ is not independent.

[9] Verification: `phase1/scripts/charges.py` (bounded P, cusp and heavy-branch expansion); `paper/scripts/review_draft1_checks.py` and `paper/scripts/paper_draft2_checks.py` (check 1; x_0-coordinate pullback by symbolic Jacobian and 50-digit quadrature).

[10] Verification: `corrigan-zambon/defect_transmission_phase.py` (check 2); `paper/scripts/paper_conventions_check.py` (check 4).

[11] Verification: `phase1/scripts/charges.py` (spectral picture); `paper/scripts/paper_conventions_check.py` (check 4) and `paper/scripts/paper_draft2_checks.py` (check 2c), including the ordered sign relation $\mu_{1,2} = -p_{b,a}$ between the Phase 1 and Phase 2 root variables.

[12] Verification: `phase2/scripts/smatrix_ansatz.py`; `phase2/scripts/consistency_checks.py` (check 5); `paper/scripts/paper_draft2_checks.py` (check 2g).

[13] Verification: `phase1/scripts/charges.py`; `paper/scripts/paper_draft2_checks.py` (check 2).

[14] Verification: `phase1/scripts/charges.py`; `paper/scripts/paper_draft2_checks.py` (check 2).

2.4 Two-soliton solutions and phase shifts

The N-soliton solution is

$$\tau = \sum_{\mu\in\{0,1\}^N} \exp\Big(\sum_i \mu_i\theta_i + \sum_{i<j}\mu_i\mu_j \log A_{ij}\Big), \qquad A_{ij} = -\frac{\mathcal{P}(k_i-k_j,\omega_i-\omega_j)}{\mathcal{P}(k_i+k_j,\omega_i+\omega_j)}, \tag{12}$$

with $\mathcal{P}(K,W) = W^2 - K^2 \pm K^4$ (upper sign good; lower sign bad, as derived in [20]) **(a)** [20]/**(c)**. The good-equation bilinear condition is verified in exact rational arithmetic at selected rational amplitudes in all direction patterns for $N \le 4$ **(c)**.[15] Position shifts are pairwise additive, so classical scattering of pure solitons is factorised **(a)**/**(c)**.

Proposition 2.2 ([good & bad], **(c)**). *Let $S = \alpha_1 + \alpha_2$, $D = \alpha_1 - \alpha_2$ (good) or $S = \beta_1 + \beta_2$, $D = \beta_1 - \beta_2$ (bad), with soliton 1 right-moving. Then*

$$\textit{good:}\quad A_{\rm ov} = \frac{1 - 3\cot^2\frac{S}{2}}{1 - 3\cot^2\frac{D}{2}}, \qquad A_{\rm ho} = \frac{1 - 3\tan^2\frac{D}{2}}{1 - 3\tan^2\frac{S}{2}}, \tag{13}$$

$$\textit{bad:}\quad A_{\rm ov} = \frac{1 + 3\coth^2\frac{S}{2}}{1 + 3\coth^2\frac{D}{2}}, \qquad A_{\rm ho} = \frac{1 + 3\tanh^2\frac{D}{2}}{1 + 3\tanh^2\frac{S}{2}}, \tag{14}$$

for overtaking ($\epsilon_1 = \epsilon_2$) and head-on ($\epsilon_1 = -\epsilon_2$) collisions. In both equations, away from coincident solitons and resonant denominators, A_{12} is the cross-ratio of the roots ((p_a, p_b) of (10) for the good equation and $(p_a, p_b) = (\mu_a, \mu_b)$ above for the bad equation),

$$A_{12} = \frac{(p_{1a} - p_{2a})(p_{1b} - p_{2b})}{(p_{1a} - p_{2b})(p_{1b} - p_{2a})}. \tag{15}$$

[16]

A_{12} agrees with Bonnemain–Doyon's eq. (16) **(a)** [6]/**(c)**. The bad constants lie in $(0, 1)$ **(c)**. The dependence on both S and D is the classical signature of the missing boost symmetry.

Proposition 2.3 ([good], **(c)**). *For the good equation (Fig. 1):*

1. *overtaking collisions are regular iff $\alpha_1 + \alpha_2 < \frac{2\pi}{3}$, and are then repulsive ($0 < A < 1$);*
2. *head-on collisions are regular and attractive ($A > 1$) for $\alpha_1 + \alpha_2 < \frac{\pi}{3}$, regular and repulsive for $|\alpha_1 - \alpha_2| > \frac{\pi}{3}$, and singular otherwise;*
3. *on each resonance line ($A = 0$ or ∞) the sum or difference of the two solitons' (k, ω) satisfies the soliton dispersion relation, and the degenerate τ-function describes a Y-junction in which two solitons merge or one splits.*

[17]

These regions agree with the conditions of Lambert, Musette and Kesteloot as quoted in [6, (29)–(33)] **(a)**/**(c)**. In root language a resonance line is where two roots of different solitons coincide, so that $\ell_1 = \ell_2$ **(c)**.[18]

[15] Verification: `phase1/scripts/hirota_solitons.py` (exact rational-amplitude checks for $N \le 4$; note that one separate asymptotic-profile comparison in that script is disabled by `or True`).

[16] Verification: `phase1/scripts/interaction_regimes.py`; `corrigan-zambon/defect_two_soliton_symplectic.py` (check 1); `phase2/scripts/consistency_checks.py` (check 5, bad); `paper/scripts/paper_conventions_check.py` (check 4, good); `paper/scripts/paper_draft2_checks.py` (check 2g, bad).

[17] Verification: `phase1/scripts/interaction_regimes.py` (400^2 grid).

[18] Verification: `corrigan-zambon/defect_two_soliton_symplectic.py` (check 5).




Figure 1: [good] *Left and centre:* regularity of the two-soliton solution in the (α_1, α_2) plane for overtaking and head-on collisions. Red: singular ($A < 0$). Blue: regular and repulsive ($0 < A < 1$). Green: regular and attractive ($A > 1$). The boundaries are resonance lines. *Right:* the one-soliton dispersion $E(P)$, with the light branch (solid) and heavy branch (dashed) meeting at the cusp $k = \sqrt{3}/2$. Figure from `phase1/scripts/figures.py`.

Position shifts. For a physical collision, label the solitons so that soliton 1 starts to the left of soliton 2 with $v_1 > v_2$. Then the asymptotic position shifts are

$$\Delta x_1 = -k_1^{-1} \log A_{12}, \qquad \Delta x_2 = +k_2^{-1} \log A_{12} \tag{16}$$

(c), checked against peak tracking of exact solutions in overtaking and attractive and repulsive head-on cases.[19] For small amplitudes in overtaking collisions, $\log A_{12}$ reduces to the KdV value $\log A_{\mathrm{KdV}} = 2 \log \left| \frac{k_1 - k_2}{k_1 + k_2} \right|$, so $\Delta x_1 \sim -2k_1^{-1} \log \left| \frac{k_1 - k_2}{k_1 + k_2} \right|$ and $\Delta x_2 \sim +2k_2^{-1} \log \left| \frac{k_1 - k_2}{k_1 + k_2} \right|$, while the head-on $\log A_{12} \to 0$ **(a)** [6]/**(c)**.[20]

2.5 Breathers

A breather is the real reduction $k_2 = \bar{k}_1$, $\omega_2 = \bar{\omega}_1$ of the two-soliton. It is regular iff $A(k, \bar{k}) > 1$.

Proposition 2.4 ([good & bad], **(c)**). *1.* [good] *For* $\alpha = a + ib$, $A(k, \bar{k}) = \dfrac{1 - 3\cot^2 a}{1 + 3\coth^2 b} < \frac{1}{4}$. *Every Hirota breather of the good equation on the zero background is singular.*

2. [bad] *For* $\beta_{1,2} = a \pm ib$ *(Bogdan–Charkina's parametrisation* $\kappa \pm ik = \sinh(a \pm ib)$*),* A_{12} *equals their* a_{12} *[4, (33)], and the breather is regular iff* $b > \pi/3$*, which is their existence boundary* $\pi/3 < b \leq \pi/2$ **(a)** *[4, (29)–(41)].*

[21]

Item 1 concerns the two-phase Hirota class on the zero background only. For the bad equation, breathers from the regular region of Charlier and Lenells are smooth for all t and those from the singular region are singular at every t **(a)** [10, Lemma A.4].

2.6 The good/bad dictionary

Table 1 collects the corresponding quantities. Results in this paper are stated for one equation or for both, as tagged.

[19] Verification: `phase1/scripts/interaction_regimes.py`.
[20] Verification: `paper/scripts/paper_draft2_checks.py` (check 3).
[21] Verification: `phase1/scripts/interaction_regimes.py` (good; grid and complex-k scan); `phase2/scripts/consistency_checks.py` (check 4, bad).

Table 1: Dictionary between the good and bad equations. Entries are **(c)** unless marked; the last three rows summarise Sections 2.5, 4.4 and 5.

	good	bad
Equation	$u_{tt} = u_{xx} - u_{xxxx} - 3(u^2)_{xx}$	$v_{tt} = v_{xx} + v_{xxxx} + 3(v^2)_{xx}$
Lagrangian	$\frac{1}{2}\phi_t^2 - \frac{1}{2}\phi_x^2 + \phi_x^3 - \frac{1}{2}\phi_{xx}^2$	$\frac{1}{2}\phi_t^2 - \frac{1}{2}\phi_x^2 - \phi_x^3 + \frac{1}{2}\phi_{xx}^2$
Phonons	$\omega^2 = q^2 + q^4$	$\omega^2 = q^2 - q^4$
Lax form	$q_L = \frac{3}{2}u - \frac{1}{4}$, $\tau = \sqrt{3}t$	$q_L = \frac{3}{2}v + \frac{1}{4}$, $\tau = i\sqrt{3}t$
Rapidity	$k = \sin\alpha$, $\alpha \in (0, \frac{\pi}{2})$	$k = \sinh\beta$, $\beta > 0$
Velocity	$v = \epsilon\cos\alpha$ (subsonic)	$v = \epsilon\cosh\beta$ (supersonic)
Spectral parameter	$\ell = -\epsilon\cos 3\alpha$	$\ell = \epsilon\cosh 3\beta$
M, Π	$2\sin\alpha$, $-\epsilon\sin 2\alpha$	$2\sinh\beta$, $-\epsilon\sinh 2\beta$
P	$\frac{2}{3}\epsilon\sin^3\alpha\cos\alpha$ (bounded)	$\frac{2}{3}\epsilon\sinh^3\beta\cosh\beta$
E	$\frac{2}{15}k^3(5 - 4k^2)$	$\frac{2}{15}k^3(5 + 4k^2)$
dP/d(rapidity)	$\frac{2}{3}\epsilon\sin\alpha\sin 3\alpha$	$\frac{2}{3}\epsilon\sinh\beta\sinh 3\beta$
A_{ov}; A_{ho}	(13)	(14)
Roots	real, $\frac{1}{2}(\pm k + v/\sqrt{3})$	complex, $\frac{1}{2}(iv/\sqrt{3} \pm k)$
Crossing map	$\alpha \to \alpha + \pi$	$\beta \to \beta + i\pi$
Two-soliton regularity	regions, resonances (Prop. 2.3)	always regular, $0 < A < 1$
Regular breathers	none (Hirota class)	for $b > \pi/3$ **(a)** [4]
Toda continuum limit	—	small amplitudes **(a)** [36]
CZ type-I defects	exist **(a)** [12]	no real type-I defect for $\lambda_{\mathrm{CZ}} \neq 0$

Remark 2.5 ([good & bad], **(c)**). The substitution $\alpha = i\beta$ maps the good-equation soliton data to the bad ones according to

$$(k,\, M,\, \Pi,\, P,\, E,\, \ell,\, v,\, A_{12})\big|_{\mathrm{good}} \;\longmapsto\; (ik,\, iM,\, i\Pi,\, -iP,\, -iE,\, -\ell,\, v,\, A_{12})\big|_{\mathrm{bad}},$$

in both orientations; consequently the semiclassical phase (21) below satisfies $\delta_{\mathrm{good}}(i\beta_1, i\beta_2) = -\delta_{\mathrm{bad}}(\beta_1, \beta_2)$.[22] This explains why a single formula covers both equations. It is a statement about formulas, not a map between real solutions.

3 Canonical structure and exact symplecticity of scattering

3.1 Cross terms in the symplectic form

Consider N well-separated solitons at positions X_j with parameters k_j, and normalise the potential by $\phi(-\infty) = c$. Then ϕ is a sum of kinks, ϕ_j rising from 0 to M_j across X_j, plus c.

Proposition 3.1 ([good & bad], **(c)**). *Up to exponentially small terms,*

$$\Omega = d\Pi_{\mathrm{tot}} \wedge dc + \sum_j dP_j \wedge dX_j + \sum_{i \neq j} \big[X_i > X_j\big]\, d\Pi_i \wedge dM_j. \tag{17}$$

The momentum density π_i of soliton i is localised at X_i, where ϕ_j equals 0 or M_j according to whether soliton i is left or right of soliton j. This gives the cross terms. The diagonal term

[22] Verification: `paper/scripts/paper_conventions_check.py` (check 5); the statement for A_{12} is in `phase1/scripts/interaction_regimes.py`.

is the one-soliton result of Proposition 2.1. In sine-Gordon the analogous cross term vanishes because the kink height $2\pi/\beta$ is fixed; here $M = 2k$ varies **(c)**.

3.2 Scattering is symplectic only with the cross terms

During a collision the parameters k_j and the constant c are conserved and $X_j \mapsto X_j + \Delta x_j$ with (16). Soliton 1 starts on the left and ends on the right, so the cross term changes from $d\Pi_2 \wedge dM_1$ to $d\Pi_1 \wedge dM_2$.

Proposition 3.2 ([good & bad], **(c)**). *The two-soliton scattering map preserves* (17)*:*

$$\sum_j dP_j \wedge d\Delta x_j = d\Pi_2 \wedge dM_1 - d\Pi_1 \wedge dM_2. \tag{18}$$

Without the cross terms the left side would have to vanish; it does not (for example it equals 19.79 *at* $\beta = (0.9, 0.3)$ *in the bad equation).*[23]

Equation (18) shows that the naive rule $\Delta x_j = -\partial\delta/\partial P_j$ cannot hold for any function δ: the shifts are not curl-free in the canonical momenta. The same identity holds on the Toda lattice in label coordinates (Proposition 4.5).

3.3 Darboux coordinates

In each ordering region define

$$X'_j = X_j + \gamma_j \sum_{i \text{ left of } j} M_i, \qquad \gamma_j = \frac{d\Pi_j/dk_j}{dP_j/dk_j} \tag{19}$$

(c). For the good equation $\gamma = -3\cos 2\alpha/(\sin\alpha \sin 3\alpha)$, which has a pole at the cusp **(c)**.[24] In these coordinates the cross terms are absorbed, and the shifts

$$\Delta X'_1 = -\frac{\log A_{12}}{k_1} + \gamma_1 M_2, \qquad \Delta X'_2 = +\frac{\log A_{12}}{k_2} - \gamma_2 M_1 \tag{20}$$

are curl-free in (P_1, P_2) **(c)**. Normalising ϕ at $+\infty$ instead gives the same in–out differences **(c)**.[25]

4 The semiclassical two-soliton phase

4.1 The phase

By stationary phase on two-particle wave packets, for a collision with incoming ordering $X_1 < X_2$ ($v_1 > v_2$) and away from $P'_j = 0$, a phase $\delta(1,2)$ satisfying $\Delta X'_j = -\partial\delta(1,2)/\partial P_j$ gives the leading semiclassical S-matrix $S \simeq e^{i\delta/\hbar}$ **(a)** [16]. We define $\delta(1,2)$ below as an antisymmetric function of the ordered pair of labels $(1,2)$, using the real principal logarithm on the regular region $A_{12} > 0$ (excluding coincident solitons $k_1 = k_2, \epsilon_1 = \epsilon_2$ and resonance lines $A_{12} \in \{0, \infty\}$).

Proposition 4.1 ([good & bad], **(c)**). *For overtaking and head-on collisions, in both equations,*

$$\boxed{\delta(1,2) = \tfrac{2}{9}(\ell_1 - \ell_2)\log A_{12} + \tfrac{2}{3} M_1 M_2 (v_1 - v_2)} \tag{21}$$

[23] Verification: Bad: `phase2/scripts/symplectic_scattering.py` (40 digits; overtaking and head-on, several points each). Good, regular region: `phase2/scripts/consistency_checks.py` (double precision) and `corrigan-zambon/defect_two_soliton_symplectic.py` (check 3, 10^{-40}, with a defect present).

[24] Verification: `paper/scripts/paper_conventions_check.py` (check 9).

[25] Verification: `phase2/scripts/semiclassical_phase.py`, `phase2/scripts/consistency_checks.py` (check 1).

with $\ell = -\epsilon\cos 3\alpha$ *(good) or* $\ell = \epsilon\cosh 3\beta$ *(bad). The second term can be written* $T_{12} \equiv \frac{2}{3}M_1M_2(v_1 - v_2) = -\frac{2}{3}(\Pi_1 M_2 - \Pi_2 M_1)$*. For the good equation the statement holds in the regular region* ($\alpha_j \neq \pi/3$, $A_{12} > 0$).[26]

Analytic proof of the phase-gradient identity. By (9) and (11), $dP_1 = \frac{2}{9}k_1\, d\ell_1$ in both equations **(c)**. Differentiating (21) with respect to the first soliton's parameter therefore gives $\partial_1\delta = (P_1'/k_1)\log A_{12} + \frac{2}{9}(\ell_1 - \ell_2)\partial_1 \log A_{12} - \frac{2}{3}(\Pi_1' M_2 - \Pi_2 M_1')$. In the root cross-ratio (15), each root satisfies $\ell(p) = -3\sqrt{3}p(1 - 4p^2)$, so for any pair of roots (p_{1r}, p_{2s}) of solitons 1 and 2,

$$\frac{\ell_1 - \ell_2}{p_{1r} - p_{2s}} = -3\sqrt{3}\big[1 - 4(p_{1r}^2 + p_{1r}p_{2s} + p_{2s}^2)\big],$$

cancelling every pole of $\partial_1 \log A_{12}$ identically **(c)**. Taking the alternating sum over $r, s \in \{a, b\}$ and using $p_{ja} - p_{jb} = M_j/2$ and $p_{ja}^2 - p_{jb}^2 = -\Pi_j/(2\sqrt{3})$ yields $\frac{2}{9}(\ell_1 - \ell_2)\partial_1 \log A_{12} = -\frac{1}{3}\Pi_1' M_2 - \frac{2}{3}\Pi_2 M_1'$ **(c)**. Adding $-\frac{2}{3}(\Pi_1' M_2 - \Pi_2 M_1')$ cancels the $\Pi_2 M_1'$ term and leaves $\partial_1\delta = (P_1'/k_1)\log A_{12} - \Pi_1' M_2 = -P_1'\Delta X_1'$ **(c)**.[27]

The remainder T_{12} was first found by symbolic integration and then identified by this root identity **(c)**. Three remarks:

- The dynamical part is (difference of spectral parameters)$\times \log A_{12}$. Because $dP = \frac{2}{9}k\, d\ell$ and $(\ell_1 - \ell_2)\partial_1 \log A_{12}$ reduces to a polynomial in the roots, integration by parts closes and the tree-level phase is elementary: no dilogarithm appears at $O(\hbar^{-1})$, in contrast to relativistic affine Toda theory **(c)** [23, 24].
- T_{12} is an Aharonov–Bohm-like "mutual statistics" phase between objects carrying shift charge Π and winding M **(d)** (interpretation).
- The same structure, (difference of spectral parameters)$\times \log A$ minus a multiple of $\Pi_1 M_2 - \Pi_2 M_1$, appears on the Toda lattice (Proposition 4.5). We conjecture that it is universal for integrable models with a continuous kink charge, KdV included **(d)**.

4.2 Properties

Proposition 4.2 ([good & bad], **(c)**). *1.* Unitarity: $\delta(1, 2) = -\delta(2, 1)$.

2. Crossing-type relation: $\delta_{\rm ov}(\beta_1, \beta_2 + i\pi) = -\delta_{\rm ho}(\beta_1, \beta_2)$ *(bad) and* $\delta_{\rm ov}(\alpha_1, \alpha_2 + \pi) = -\delta_{\rm ho}(\alpha_1, \alpha_2)$ *(good, regular head-on region). The shift maps a right-mover to a left-mover with* $(k, M, \ell) \to (-k, -M, -\ell)$ *and* $A_{\rm ov} \to 1/A_{\rm ho}$.

3. [good] Resonance lines: *where* $A_{12} \to 0$ *or* ∞ *one has* $\ell_1 = \ell_2$*, so* $\delta \to T_{12}$ *stays finite while the delays diverge logarithmically.*

4. [good] Cusp: *at* $\alpha_1 = \pi/3$*,* $dP_1/d\alpha_1 = 0$ *while* $\partial_{\alpha_1}\delta = -\Pi_1'(\pi/3)M_2 = -\epsilon_1 M_2 \neq 0$*; hence* δ *is smooth in* α_1 *but has a square-root branch point* $\delta(P_1, P_2) - \delta(P_{\rm cusp}, P_2) \sim \mp\sqrt{2/\sqrt{3}}\,\epsilon_1 M_2\sqrt{|P_{\rm cusp} - P_1|}$ *as a function of* P_1.[28]

[26] Verification: `phase2/scripts/semiclassical_phase.py` (symbolic integration of the Darboux shifts); `phase2/scripts/consistency_checks.py` (check 1, gradient test for both equations and orientations); `paper/scripts/paper_draft2_checks.py` (check 8, symbolic root-polynomial proof).

[27] Verification: `paper/scripts/paper_draft2_checks.py` (check 8).

[28] Verification: Items 1–2: `phase2/scripts/consistency_checks.py` (unitarity asserted; bad crossing printed) and `paper/scripts/paper_conventions_check.py` (check 6, both asserted at 40 digits). Item 3: `corrigan-zambon/defect_two_soliton_symplectic.py` (check 5, printed resonance-line limits). Item 4: `corrigan-zambon/defect_two_soliton_symplectic.py` (check 5, soliton–defect cusp asserted) and `paper/scripts/paper_draft2_checks.py` (check 8, two-body cusp square-root branch expansion asserted).

Near the cusp the wave-packet derivation needs a uniform (Airy-type) treatment, and α or the root, rather than P, is the natural label **(d)**. Past the resonance lines lie the Y-shaped solutions, which are outside the two-particle description.

Small amplitudes. [bad] For overtaking collisions with $\beta_1/\beta_2 = 1.7$ fixed and $\beta_j \to 0$, the ratio of the full phase derivative $\partial\delta/\partial\beta_1$ to the KdV-type term $2\beta_1 \log A_{12}$ tends to $0.564246498175\ldots = 1 + 2\beta_2/(\beta_1 \log A_{\mathrm{KdV}})$, with $A_{\mathrm{KdV}} = (\frac{\beta_1-\beta_2}{\beta_1+\beta_2})^2$, while the cross-term-only ratio tends to $-0.435753501825\ldots$ **(c)**. The cross-term contribution $-(d\Pi_1/d\beta_1)M_2 \to 4\beta_2$ to the phase derivative $\partial\delta/\partial\beta_1 = -P_1'\Delta X_1'$ is therefore of the same order as the KdV-type term **(c)**.[29] The heuristic of Bonnemain and Doyon identifies the quantum phase derivative with $P'_{\mathrm{GHD}}\Delta x$, with $P_{\mathrm{GHD}} = \eta^2/2$ **(b)** [6]. It reproduces the dynamical term but misses the cross-term contribution **(c)**; their classical hydrodynamics, which uses only Δx, is unaffected. [good & bad] For head-on collisions $\delta \to 4\beta_1\beta_2$, of which $\frac{16}{3}\beta_1\beta_2$ comes from T_{12} and $-\frac{4}{3}\beta_1\beta_2$ from the dynamical term (and likewise with α_j for the good equation) **(c)**.[30]

4.3 Bad equation: breathers as spectral-parameter strings

Bogdan and Charkina compute the classical integrals and adiabatic invariant I_{br} of the Tajiri–Murakami breather of the bad equation, verifying $dE = (\tilde{\omega}/2\pi)\, dI_{\mathrm{br}} + V\, dP$ **(a)**, and apply the semiclassical Bohr–Sommerfeld quantisation rule $I_{\mathrm{br}} = N\hbar$ at large quantum number $N \gg 1$ **(b)** [4, (64)–(71)].

Proposition 4.3 ([bad], **(c)**). *At $\beta_{1,2} = a \pm ib$,*

$$i \cdot \tfrac{2}{9}(\ell_1 - \ell_2) = -\tfrac{4}{9}\sinh 3a \,\sin 3b = I_{\mathrm{br}}, \tag{22}$$

the Bogdan–Charkina classical adiabatic invariant **(a)** *[4, (64)]. Hence the coefficient of* $\log A_{12}$ *in* (21) *is the breather action* **(c)**, *and under the Bohr–Sommerfeld prescription* $I_{\mathrm{br}} = N\hbar$ **(b)** *breather quantisation reads*

$$\ell_1 - \ell_2 = -\tfrac{9i}{2}\, N\hbar, \qquad N \in \mathbb{N}. \tag{23}$$

[31]

This ties the phase (21), built from Hirota shifts and the canonical symplectic form, to an independent classical action calculation in the literature **(c)**. Equation (23) is a string condition in the spectral parameter, the form in which bound states appear in Bethe-ansatz solvable models. Extrapolating this semiclassical condition to finite N suggests that an exact S-matrix has bound-state poles at $\ell_1 - \ell_2 \in -\frac{9i}{2}\hbar\mathbb{N}$, possibly with a renormalised $\hbar$ **(d)**. The good equation has no regular Hirota breathers on the zero background (Proposition 2.4), so there is no analogue of this check there.

4.4 Bad equation: the classical limit of the quantum Toda Bethe ansatz

Set-up. The Toda chain is $H = \sum_n (\frac{1}{2}p_n^2 + e^{-r_n})$ with $r_n = x_{n+1} - x_n$, mean spacing a and $b = e^{-a/2}$. Its small-amplitude long-wave limit is the bad equation **(a)** [36, 4]/**(c)**.[32] The one-soliton is $e^{-r_n} = b^2[1 + \sinh^2\kappa\, \mathrm{sech}^2(\kappa n - sbt\sinh\kappa)]$, with Toda direction sign $s = \pm 1$ (distinct from the defect parameter $s = \sigma_0$ of Section 5) and

$$M = \sum (r_n - a) = -2\kappa, \qquad \Pi = \sum p_n = 2sb\sinh\kappa, \qquad E = 2b^2(\sinh\kappa\cosh\kappa - \kappa),$$

[29] Verification: `phase2/scripts/consistency_checks.py` (check 6, reported as a finding); `paper/scripts/paper_conventions_check.py` (check 7) and `paper/scripts/paper_draft2_checks.py` (check 3), limit identified and asserted.

[30] Verification: `paper/scripts/paper_conventions_check.py` (check 7).

[31] Verification: `phase2/scripts/consistency_checks.py` (check 4, three parameter points, 10^{-16}); `paper/scripts/paper_conventions_check.py` (check 8, closed form).

[32] Verification: `phase1/scripts/toda_limit.py`.

where $E=\sum[\tfrac12 p_n^2+e^{-r_n}-b^2+b^2(r_n-a)]$. The canonical form $\sum dp_n\wedge dx_n$ pulled back to (κ,n_0) is $dP_n\wedge dn_0$ with label momentum $P_n=4sb(\kappa\cosh\kappa-\sinh\kappa)$, and $dE/dP_n=sb\sinh\kappa/\kappa$ is the signed label velocity **(c)**.[33] The two-soliton constant is $A_T=-\mathcal{P}_T(2\kappa_1-2\kappa_2,\omega_1-\omega_2)/\mathcal{P}_T(2\kappa_1+2\kappa_2,\omega_1+\omega_2)$ with $\mathcal{P}_T(K,W)=W^2-4b^2\sinh^2(K/2)$ and $\omega=2sb\sinh\kappa$ **(c)**.[34]

The relative two-body problem $(-\hbar^2\partial_x^2+e^{-x})\psi=q^2\psi$ with incoming wave $e^{-iqx/\hbar}$ has reflection amplitude $R=\Gamma(i\kappa)\hbar^{2i\kappa}/\Gamma(-i\kappa)$, $\kappa=2q/\hbar=\Delta p/\hbar$ **(a)**/**(c)**.[35] The asymptotic Bethe equations are $p_jL+\sum_l\Theta(p_j-p_l)=2\pi\hbar I_j$ with $\Theta(D)=2\hbar\,\mathrm{Im}\log\Gamma(1+iD/\hbar)+2D\log\hbar$ (using the unwrapped logarithm branch), whose classical limit is $2(D\log|D|-D)$; the method is that of Sutherland and Mertens **(b)** [34, 29].

Proposition 4.4 ([Toda → bad], **(c)**). *As $\hbar\to0$:*

1. *the ground-state quasi-momenta fill $[-2b,2b]$ with the arcsine density of total weight $1/a$, which is the Lax spectrum of the static lattice, and reproduce the ground-state energy per particle b^2;*
2. *a particle excitation at $p=2sb\cosh\kappa$, the soliton's Lax eigenvalue, has dressed momentum P_n/a and dressed energy, relative to the edge, equal to the soliton energy E;*
3. *the exact classical physical-frame shifts are $\Delta x_1=a\Delta n_1+M_2$ and $\Delta x_2=a\Delta n_2-M_1$, the extra terms being the displacement caused by the other soliton's compression; they are given by $\Delta x_2=-\partial\Phi(1|2)/\partial P_2$ and $\Delta x_1=+\partial\Phi(2|1)/\partial P_1$, where $\Phi(1|2)$ is the dressed Bethe phase of excitation 1 in the presence of 2;*
4. *$\Phi(1|2)+\Phi(2|1)$ is not separable; its mixed derivative with respect to (κ_1,κ_2) equals $M_1'(P_{n2}'/a-\Pi_2')+M_2'(P_{n1}'/a-\Pi_1')$ (primes denoting $d/d\kappa_j$), i.e. the Bethe ansatz carries precisely the cross terms of Proposition 3.1.*

[36]

Proposition 4.5 ([Toda → bad], **(c)**). *In the Toda label frame, two-soliton scattering satisfies $\sum_j dP_{n,j}\wedge d\Delta n_j=d\Pi_2\wedge dM_1-d\Pi_1\wedge dM_2$ (30 digits), the analogue of (18). The Darboux phase is*

$$\delta_T=(p_1-p_2)\log A_T+M_1M_2(v_1-v_2)=(p_1-p_2)\log A_T-(\Pi_1M_2-\Pi_2M_1),\tag{24}$$

with $p=2sb\cosh\kappa$ and $v=sb\sinh\kappa/\kappa$ the signed label velocity.[37]

Proposition 4.6 ([Toda → bad], **(c)**). *With $\kappa_j=\sqrt{3}\beta_j\to0$ and action scale $S_{\mathrm{Toda}}=3b\,S_{\mathrm{Bouss}}$,*

$$\delta_T/(3b\,\delta_{\mathrm{bad}})\to1$$

for overtaking and *head-on collisions; the error is $\approx6\times10^{-6}$ at $\beta\sim3\times10^{-3}$.*[38]

[33] Verification: `phase2/scripts/toda_soliton_classical.py`, `phase2/scripts/toda_label_symplectic.py`; signed velocity verified for both $s=\pm1$ in `paper/scripts/review_draft1_checks.py` and `paper/scripts/paper_draft2_checks.py` (check 4a).

[34] Verification: `phase1/scripts/toda_limit.py` (exact two-soliton in the lattice, 10^{-17}); `phase2/scripts/toda_label_symplectic.py`.

[35] Verification: Verified from the exact Bessel eigenfunction $\psi(x)=K_{i\kappa}(2e^{-x/2}/\hbar)$ and its $x\to+\infty$ asymptotics in `paper/scripts/review_draft1_checks.py` and `paper/scripts/paper_draft2_checks.py` (check 4b).

[36] Verification: `phase2/scripts/toda_bethe_classical.py` (10^{-8}, Chebyshev method); `phase2/scripts/toda_dressed_phase_check.py`; `phase2/scripts/toda_ba_vs_label.py`.

[37] Verification: `phase2/scripts/toda_label_symplectic.py` (symplectic identity); `phase2/scripts/toda_scaling_fast.py` (remainder fitted to $M_1M_2(v_1-v_2)$, ratio 1 to ten digits); `phase2/scripts/toda_ba_vs_label.py` ($\Phi(1|2)=\delta_T+M_1(P_{n2}/a-\Pi_2)+f(\kappa_1)$ and its mirror, to machine precision).

[38] Verification: `phase2/scripts/toda_scaling_fast.py`.

In head-on collisions the leading Boussinesq phase $4\beta_1\beta_2$ is dominated by the topological term ($\frac{16}{3}\beta_1\beta_2$ against $-\frac{4}{3}\beta_1\beta_2$). Agreement in that channel therefore confirms the topological term of (21) from an exactly solvable quantum model **(c)**. The split between the two terms of (24) is not preserved by the scaling limit; only their sum is **(c)**.

Limitation. The Toda continuum limit produces the bad equation only at leading order in amplitude, so Toda tests (21) only as $\beta \to 0$. Finite-β solitons have widths comparable to the dispersion length, where the bad equation is ill-posed and Toda and Boussinesq differ at $O(1)$ **(c)**. An exact S-matrix for finite-amplitude bad solitons would require a quantum regularisation that agrees with Boussinesq at the dispersion scale; we know of none **(d)**. The $O(\hbar)$ Toda corrections and the Fermi-edge double-scaling limit are discussed in Section 8.

5 Integrable defects at tree level

[good] Throughout this section the bulk theory is the good equation.

5.1 Corrigan–Zambon defects

Corrigan and Zambon construct integrable type-I defects for the good Boussinesq equation, with a defect Lagrangian $D = u_t F + v_t G + H$ at x_0 coupling the fields u ($x < x_0$) and v ($x > x_0$) **(a)** [12]. Within their ansatz, with the nonlinear term present, the defect coefficients are forced to be imaginary when the dispersion has the bad sign, so *no real type-I defect of this family exists for the bad equation*; for the linear equation real defects do exist **(c)**.[39] A type-II defect for the bad equation is not excluded **(d)**. This is why defects are a good-equation topic.

Dictionary. With $c = \lambda = \kappa = 1$ in [12], $u_{\mathrm{CZ}} = -3\phi$, $L_{\mathrm{CZ}} = 9\mathcal{L}$, $P_{\mathrm{CZ}} = 9P$, $\gamma = \operatorname{sech}\theta = k$ and $\nu = \tanh\theta = \epsilon\cos\alpha$ **(c)**.[40] The defect has parameter $s = \sigma_0$ and orientation $\eta = \pm 1$ (CZ's $\alpha_{\mathrm{CZ}} = \eta/\sqrt{3}$, $\beta_{\mathrm{CZ}} = 2\eta/\sqrt{3}$, obeying $\beta_{\mathrm{CZ}}^2 - \alpha_{\mathrm{CZ}}^2 = 1$ **(a)** [12, (3.22),(3.37)]). The sewing conditions were rederived from the action and checked on the exact solutions **(c)**.[41] One- and two-soliton solutions are purely transmitted, each e^{θ_j} being multiplied by

$$z = \frac{\sinh\theta + s\cosh\theta - \eta\sqrt{3}}{\sinh\theta + s\cosh\theta + \eta\sqrt{3}} \tag{25}$$

on the far side **(a)** [12, (5.5),(5.11)].

Proposition 5.1 ([good], **(c)**). *With the soliton roots* (10) *and the* defect root $q = -s/(2\sqrt{3})$,

$$z = \left(\frac{p_b - q}{p_a - q}\right)^{\eta}, \qquad \ell(q) = \tfrac{1}{2}(3s - s^3). \tag{26}$$

Formally $z^{-\eta}$ is the cross-ratio (15) *of the soliton with the pair (q, ∞): the defect behaves like half a soliton. The transmitted solution is regular iff $z \in (0, \infty)$, i.e. $q \notin [p_b, p_a]$; for $|s| > 2$ every soliton is regular, in agreement with [12]* **(a)**. *The delay is $\Delta x = -k^{-1}\log z^{\epsilon}$.*[42]

Corrigan and Zambon remark that no factor of the two-soliton constant seems closely related to z **(a)**; in root variables the relation is transparent.

[39] Verification: `corrigan-zambon/defect_check.py` (sewing conditions and momentum flux rederived from the action with the dispersion sign kept symbolic).

[40] Verification: `corrigan-zambon/defect_transmission_phase.py` (check 1).

[41] Verification: `corrigan-zambon/verify_defect_solutions.py` (10^{-28}–10^{-31} away from poles).

[42] Verification: `corrigan-zambon/defect_transmission_phase.py` (checks 2–3; delay from peak positions, 16 cases, 10^{-39}).

5.2 Symplectic structure with a defect

The defect adds $F\,\delta u + G\,\delta v$ at x_0 to the canonical one-form Θ. Asymptotically $F \to f(u,v)$ and $G \to g(u,v)$, so with $\Omega = d\Theta$ the defect contributes $\Omega_D = d(f\,du + g\,dv)|_{x_0} = (g_u - f_v)\,du \wedge dv|_{x_0}$ **(c)**. On every fixed vacuum sector $u - v$ at x_0 is a discrete label, so $du = dv$ and $\Omega_D = 0$ **(c)**.[43] Hence the type-I defect contributes no continuous canonical pair on a fixed vacuum sector and introduces no soliton–defect cross terms (this does not imply the absence of localised defect energy or momentum).[44] For one soliton, any map $X \mapsto X + \Delta x(P)$ is then symplectic, so the content lies in the phase and in the multi-soliton case.

5.3 The transmission phase

Proposition 5.2 ([good], **(c)**). *With $\Delta x = -\partial\delta_D/\partial P$, for all $\epsilon, \eta = \pm 1$ and all s in the regular region,*

$$\boxed{\delta_D = \tfrac{2}{9}\big(\ell - \ell_D\big)\log z^{\epsilon} - \tfrac{2}{3}\big(\Pi M_D - \Pi_D M\big)}, \qquad \ell_D = \ell(q), \quad M_D = \epsilon\eta\, q, \quad \Pi_D = \epsilon\eta\,\frac{s^2-2}{2\sqrt{3}}, \tag{27}$$

up to an additive s-dependent constant that the classical data do not fix. The two orientations are semiclassically inverse: $\delta_D(-\eta) = -\delta_D(\eta)$.[45]

The defect enters like a second particle with spectral parameter $\ell(q)$ and effective coefficients (M_D, Π_D) **(c)** (formula); the particle reading is **(d)**. In CZ's notation $\ell_D = -\sqrt{3}\,\eta\rho_0$, so the defect's spectral parameter is its constant ρ_0 **(c)**. Since M_D and Π_D depend on ϵ, they are effective coefficients of a mutual-statistics term rather than intrinsic defect charges.

5.4 Two solitons and a defect

Proposition 5.3 ([good], **(c)**). *For two solitons crossing a defect (overtaking and head-on, both orientations, s of both signs, regular region):*

1. *the position shifts are $\Delta x_j = \Delta x_j^{\mathrm{ss}} + \Delta x_j^{D}$, identical whether the soliton–soliton collision happens left or right of the defect;*
2. *the identity (18) holds with the soliton–soliton cross terms and no defect cross terms;*
3. *in the Darboux frame (19), $\Delta x_j' = -\partial_{P_j}(\delta_{12} + \delta_{1D} + \delta_{2D})$.*

[46]

Semiclassically the soliton–soliton–defect amplitude therefore factorises as $S_{12}T_1T_2$, and the compatibility relation $T_1T_2S_{12} = S_{12}T_2T_1$ is automatic for diagonal, purely transmitting scattering. The nontrivial content is that the defect shifts do not depend on the other soliton, and that the Darboux corrections are unaffected by the defect because $\Omega_D = 0$ **(c)**.

[43] Verification: `corrigan-zambon/defect_transmission_phase.py` (check 6); two-form sign checked in `paper/scripts/review_draft1_checks.py` and `paper/scripts/paper_draft2_checks.py` (check 5).

[44] We use the shift-invariant gauge $f = g = -\sigma^2/(4\mu)$ ($\mu = 1$, $\sigma = v - u + \eta\sqrt{3}\kappa$) with $g_u - f_v = \sigma$, in which the total Π is conserved through the defect. The example gauge printed in [12, (3.43),(3.47)] gives 2σ; this looks like a typo there **(c)** (`corrigan-zambon/README.md`).

[45] Verification: `corrigan-zambon/defect_transmission_phase.py` (check 4, 40 digits).

[46] Verification: `corrigan-zambon/defect_two_soliton_symplectic.py` (checks 2–4; shifts from peak positions at $t = \pm 1000$ to 10^{-27}; identities to 10^{-40}).

5.5 Vacua, absorption and linear stability

The static defect vacua are $\sigma \in \{s, \sigma_+, \sigma_-\}$ with $\sigma_\pm^2 + s\sigma_\pm + s^2 = 3$, equivalently the roots of $P_{\rm vac}(\sigma) \equiv \sigma^3 - 3\sigma + s(3 - s^2) = 0$ **(c)**. Writing $\sigma = -2\sqrt{3}\,q'$, this reads $q'^2 + qq' + q^2 = \frac{1}{4}$: the excited vacua $\sigma_\pm$ are the *other two roots of the KP triple containing* q **(c)**. Since $\sigma_\pm^2 + s\sigma_\pm + s^2 - 3$ has discriminant $\Delta = 3(4 - s^2)$ and $(s - \sigma_+)(s - \sigma_-) = 3(s^2 - 1)$, all three vacua of the triple are real and distinct iff $|s| < 2$ and $s \neq \pm 1$; at $|s| = 2$ or $s = \pm 1$ two roots coincide (at $\eta s = 2$ they coincide at the marginal point $\eta\sigma = -1$), while for $|s| > 2$ the excited roots $\sigma_\pm$ are nonreal complex conjugates, leaving $\sigma = s$ as the unique real vacuum **(c)**.[47] Absorption ($z \in \{0, \infty\}$) occurs iff $q = p_a$ or $q = p_b$ (which requires $|s| < 2$) **(a)** [12]/**(c)**. The defect with root q then absorbs the soliton (q, p) and becomes the defect with root p in the same triple, with exact energy–momentum balance between the soliton and the jump of the defect's (E_D, P_D) **(c)**.[48] On an absorption line $\ell = \ell(q)$, so δ_D stays finite while the delay diverges logarithmically **(c)**.[49]

Proposition 5.4 ([good], **(c)/(d)**). *Linearise the sewing conditions about a real static vacuum* $\sigma_{\rm vac}$*; the bulk equation becomes* $a_{tt} = a_{xx} - a_{xxxx}$.

1. *On every real vacuum phonons are purely transmitted,* $R = 0$, $|T| = 1$, *with* T *equal to CZ's linear-defect formula [12, (4.6)] at* $\sigma_0 \to \sigma_{\rm vac}$ **(c)**.

2. *Numerical searches over the spectral plane find a localised growing mode for* $\eta\,\sigma_{\rm vac} > -1$ *and find no growing mode for* $\eta\,\sigma_{\rm vac} < -1$ *(with* $\eta\,\sigma_{\rm vac} = -1$ *marginal)* **(c)***; that no growing mode exists anywhere in the complex plane for* $\eta\,\sigma_{\rm vac} < -1$ *is a completeness statement beyond the finite grid* **(d)**. *Every growing mode found has zero conserved energy; the defect's quadratic energy is indefinite* **(c)**.

3. *For* $|s| < 2$ *and* $s \neq \pm 1$, *the cubic* $P_{\rm vac}(\sigma) = \sigma^3 - 3\sigma + s(3 - s^2)$ *has derivative* $P'_{\rm vac}(\sigma) = 3(\sigma^2 - 1)$ *and satisfies* $P_{\rm vac}(-1) = (2 - s)(1 + s)^2 > 0$ *and* $P_{\rm vac}(+1) = -(2 + s)(1 - s)^2 < 0$*; by Rolle's theorem its three distinct real vacua lie in* $(-\infty, -1)$, $(-1, 1)$ *and* $(1, \infty)$, *so* exactly one vacuum per distinct real triple satisfies the strict stability condition $\eta\,\sigma_{\rm vac} < -1$ **(c)**.

[50]

Consequently, within the sampled mode search, a defect with ground vacuum $\sigma = s$ is strictly linearly stable for $\eta s < -1$ (marginal at $\eta s = -1$) and linearly unstable for $\eta s > -1$ **(c)/(d)**. Corrigan and Zambon's illustrative example ($s = 0.5$, $\eta = +1$) has a linearly unstable ground vacuum, with growth rate ≈ 0.4999; its exact soliton solutions are unaffected **(c)**. Soliton absorption by a strictly stable ground defect is possible only for $-2 < \eta s < -1$, where all three vacua of the triple are real and distinct; in that range both excited vacua $\sigma_\pm$ have $\eta\sigma_\pm > -1$ and are linearly unstable **(c)**, so the absorbed soliton leaves the defect in an unstable state, presumably a resonance that re-emits **(d)**. (For $\eta s < -2$ the ground vacuum is stable and the two excited roots are nonreal, so no real soliton is absorbed **(c)**.)

5.6 Tree-level fusion

Proposition 5.5 ([good], **(c)**). *Let* $|s| < 2$ *(*$s \neq \pm 1$*) and let a defect with root* q *absorb the soliton* $S = \{q, q'\}$ *(with* $k_S = |q - q'|$ *and velocity* $v_S = \sqrt{3}(q + q')$*) and become the defect with root*

[47] Verification: `paper/scripts/review_draft1_checks.py`; `paper/scripts/paper_draft2_checks.py` (check 6).

[48] Verification: `corrigan-zambon/defect_transmission_phase.py` (check 5, 10^{-40}).

[49] Verification: `corrigan-zambon/defect_two_soliton_symplectic.py` (check 5).

[50] Verification: `corrigan-zambon/defect_vacuum_stability.py` (fine scan of the imaginary axis plus Newton iteration from a complex grid); Rolle's-theorem separation and threshold cases checked symbolically in `paper/scripts/review_draft1_checks.py` and `paper/scripts/paper_draft2_checks.py` (check 6).

q'. *For the displayed zero-constant representatives* (27) *and every test soliton 3 in the regular region,*

$$\delta_D^{(q')}(3) - \delta_D^{(q)}(3) = \rho\,\big[\delta(3,S) + M_S\Pi_3\big], \qquad \rho = \epsilon_3\eta\,\mathrm{sgn}(q-q'). \tag{28}$$

The term $M_S\Pi_3$ is the cross term of soliton 3 with the ϕ-jump M_S stored in the excited defect. Shifting $\delta_D^{(q)} \mapsto \delta_D^{(q)} + C(q)$ by an s-dependent constant leaves the classical delay $\Delta x = -\partial_P\delta_D$ unchanged and adds $C(q') - C(q)$ to the left-hand side of (28).[51]

Identity (28) is a classical, leading-order soliton–soliton–defect bootstrap for the zero-constant representatives (27) **(c)**; fixing the absolute s-dependent phase constant requires quantum input **(d)**. All three solitons of one KP triple have the same ℓ. Promoting (28) to an exact relation between quantum amplitudes would require the excited defect vacua to be asymptotic states. For $-2 < \eta s < -1$, Proposition 5.4 shows that they are linearly unstable, so any quantum fusion argument built on (28) is heuristic **(d)**. Quantum fusion would also need the boundary renormalisation and localised charges of the defect, which are not available (Section 7).

5.7 What is not covered

For ordinary one- and two-soliton solutions the defect is purely transmitting **(a)** [12], so $R = 0$ at the order considered here **(c)**. Corrigan and Zambon's reflection occurs only for merging/splitting three-exponential solutions, and their defect can decay into two solitons **(a)** [12]. These processes change the soliton number and lie outside the fixed-number phase-space description used here; we have not derived a reflection amplitude. A guess is that semiclassical reflection is a resonance phenomenon, absorption into an excited vacuum followed by emission into another chord of the triple **(d)**.

6 Structural results on quantum inverse scattering and affine Toda

6.1 Non-ultralocality

An ordinary ultralocal Faddeev–Korepin–Sklyanin lattice quantisation requires an ultralocal r-matrix Poisson algebra for the Lax connection.

Proposition 6.1 ([good], **(c)**). *In canonical variables the Lax data of Section 2 are $q_L = \frac{3}{2}\phi_x - \frac{1}{4}$ and $r_L = \frac{1}{2}q_{L,x} + \frac{\sqrt{3}}{4}\pi$ (good real form; the bad one differs by factors of i). Their brackets are*

$$\{q_L(x), r_L(y)\} = \tfrac{3\sqrt{3}}{8}\,\delta'(x-y), \qquad \{q_L, q_L\} = \{r_L, r_L\} = 0,$$

so the companion-matrix Lax connection is non-ultralocal.[52]

This obstructs the standard ultralocal QISM construction, though not necessarily every possible lattice quantisation **(c)**/**(d)**. Two ways out are known in general, and neither yet gives a theory on the line. Maillet's r/s formalism with lattice regularisation [27] has not been constructed for Boussinesq. The Miura (free-field) route of Bazhanov, Hibberd and Khoroshkin [3] and Hikami [19] yields $U_q(\widehat{\mathfrak{sl}}_3)$ T- and Q-operators, but in the chiral setting on a circle, with no particle sector **(a)**. Since scattering is diagonal, the Yang–Baxter equation is automatic **(c)**; the "beyond Yang–Baxter" phenomena of [15] concern polarisations of the matrix equation and are vacuous for the scalar one **(c)**.

[51] Verification: `phase2/scripts/defect_smatrix_good_continuation.py` (check 4; 12 triangles, 10^{-40}); oriented sign $\rho = \epsilon_3\eta\,\mathrm{sgn}(q-q')$ and additive-constant shift verified in `paper/scripts/review_draft1_checks.py` and `paper/scripts/paper_draft2_checks.py` (check 7).

[52] Verification: `phase2/scripts/routes_BC_checks.py` (check B); smeared-bracket test in `paper/scripts/review_draft1_checks.py`.

6.2 $\mathfrak{sl}_3$ factorisation of the interaction constant

Proposition 6.2 ([good & bad], **(c)**). *Let* $A^{(12)}(\theta) = \dfrac{2\cosh\theta + 1}{2(1+\cosh\theta)}$ *be Hollowood's* $a_2^{(1)}$ *affine Toda interaction constant between the two fundamental species* **(a)** *[22, (1.19)]. For the bad equation, with* $S = \beta_1 + \beta_2$, $D = \beta_1 - \beta_2$,

$$A_{\rm ov} = \frac{A^{(12)}(S)}{A^{(12)}(D)} \cdot \frac{\tanh^2(D/2)}{\tanh^2(S/2)}, \qquad A_{\rm ho}(\beta_1,\beta_2) = \frac{1}{A_{\rm ov}(\beta_1,\beta_2+i\pi)}, \tag{29}$$

where $\tanh^2(\theta/2)$ *is the sine-Gordon two-soliton factor. Equivalently* $A_{\rm ov} = G(S/2)/G(D/2)$ *with*

$$G(y) = \frac{16\prod_{a=1,2}\sinh\frac{y+ia\pi/3}{2}\,\sinh\frac{y-ia\pi/3}{2}}{\sinh^2 y}.$$

For the good equation the corresponding form is

$$A_{\rm ov} = \frac{\sin(\frac{S}{2}+\frac{\pi}{3})\sin(\frac{S}{2}-\frac{\pi}{3})\,\sin^2\frac{D}{2}}{\sin(\frac{D}{2}+\frac{\pi}{3})\sin(\frac{D}{2}-\frac{\pi}{3})\,\sin^2\frac{S}{2}}.$$

[53]

The Boussinesq interaction is thus built from exactly the $\mathfrak{sl}_3$ blocks, but in the form $F(\beta_1-\beta_2)/F(\beta_1+\beta_2)$, the structure of scattering off a particle and its mirror image. This suggests a link with boundary (reflection) amplitudes and with the boundary affine Toda structure in [3] **(d)**.

6.3 Obstructions

- [good & bad] In the rapidities β_j (bad) or α_j (good), the semiclassical phase is not of product form: $\partial_S\partial_D\delta \neq 0$ (for example -0.55 at $(S,D) = (1.4, 0.6)$ for bad overtaking and -0.068 at $(S,D) = (0.7, 0.2)$ for good overtaking) **(c)**. Hence no S-matrix of the product form $F(\beta_1-\beta_2)G(\beta_1+\beta_2)$ (or $F(\alpha_1-\alpha_2)G(\alpha_1+\alpha_2)$) in the chosen rapidities, including pure $\mathfrak{sl}_3$ mirror forms and quantum dilogarithms of $\beta_1 \pm \beta_2$ separately, has the semiclassical limit $e^{i\delta/\hbar}$ **(c)**.[54] The weight $\frac{2}{9}(\ell_1-\ell_2)$ of $\log A_{12}$ is not a sum of functions of S and D, and T_{12} has no affine Toda analogue **(c)**.

- In the affine Toda constant the mass term $4m^2\sin^2(\pi a/3)$ appears where Boussinesq has K^4 **(c)**, so the discrete species label is replaced by the continuous wavenumber; no fixed-species affine Toda limit or dictionary producing Boussinesq solitons has been constructed here **(d)**.

- Soliton–breather scattering in $a_2^{(1)}$ Toda is flawed both classically and quantum mechanically **(b)** [35].

The structural similarity with affine Toda is therefore precise, but we have no dictionary that transfers its S-matrices.

[53] Verification: `phase2/scripts/routes_BC_checks.py` (check C, bad); `phase1/scripts/interaction_regimes.py` (good).

[54] Verification: `phase2/scripts/smatrix_rejected_forms.py` (bad); `paper/scripts/paper_draft2_checks.py` (check 10, both equations and orientations).

7 One-loop corrections

Pending

This section is pending the projected fluctuation determinant package and resolution of the open one-loop audits (`phase2/report.pdf`); no one-loop S-matrix is claimed here.

Phonon scattering on soliton and defect backgrounds. Linearising the τ-function of the good equation produces reflectionless phonon modes on N-soliton and Corrigan–Zambon defect backgrounds with factorised transmission amplitudes **(c)** (`phase2/oneloop_phase.pdf`, `phase2/oneloop_defect_phase.pdf`); for the bad equation, restricting the one-loop mode sum to the stable phonon band $|q| < 1$ (`phase2/oneloop.pdf`) is motivated by the Charlier–Lenells stable-subspace condition **(b)**, though identifying that linearised band truncation with the full nonlinear invariant manifold remains heuristic **(d)**. For the defect sewing determinant, independent checks correct a missing factor of 16 in the prefactor and an admissible-root branch change in the growing-mode pair **(c)** (`phase2/audit_frames_defects.pdf`).

Continuum integrals, charges and the projected determinant. The Faddeev–Korepin continuum integrals evaluate in closed form in terms of the Legendre function χ_2 and logarithms, and direct potential-field quadrature confirms the one-soliton spectral-density identity **(c)** (`phase2/audit_determinant_foundations.pdf`); the regulated co-moving mode sum reproduces the proposed one-loop charge Lagrangian when its cutoff endpoint is retained **(c)** (conditional, `phase2/audit_charges_action.pdf`); and potential-tail cross terms cancel from the finite collective symplectic density **(c)**, although the explicit projector does not preserve fixed potential endpoints **(c)** (`phase2/audit_projected_measure.pdf`). However, raw phase integrals are not exchange-antisymmetric and carry orientation-dependent discrepancies **(c)** (`phase2/independent_phase2h_audit.pdf`), and the full boundary-compatible fluctuation determinant, collective measure, defect UV counterterm and absolute phase normalisation remain unestablished **(d)**.

8 The quantum Toda Fermi-edge limit

Pending

This section is pending the audit of the Phase 2i Wiener–Hopf analysis (`phase2/toda_wh_and_kp_synthesis.pdf`) and the derivation of a common quantum momentum and frame dictionary (`phase2/report.pdf`). No Phase 2i claim is included in this draft.

Double-scaling limit and frame dictionary. In the Fermi-edge double-scaling limit $\beta \to 0$, $\hbar \propto \beta^2$ of the quantum Toda Bethe ansatz (where excitations lie within $O(\hbar)$ of the Fermi edge and $\delta/\hbar = O(1)$), the dressed two-particle phase probes small-amplitude quantum scattering of bad Boussinesq solitons, extending Pustilnik and Matveev's KdV analysis [30] to bidirectional collisions **(d)**. Within the implemented classical frame conversion, finite-$\hbar$ numerical data (`phase2/toda_route.pdf`, `phase2/toda_edge.pdf`, `phase2/matching.pdf`) approach the classical phase in the head-on channel and deviate from the Gamma candidate of Section 9 near threshold in the overtaking channel **(c)**. Independent Fredholm identities fix the dressed edge momentum $P_{\mathrm{dr}}(q) = \pi\hbar_T/a$ and an endpoint reciprocity bilinear for the symmetric dressed phase, revealing a nonseparable finite-$\hbar_T$ remainder after the classical frame subtraction **(c)** (`phase2/audit_toda_dictionary.pdf`); the physical quantum frame dictionary and its continuum matching at fixed quantum observables remain open **(d)**.

9 Exact S-matrix candidates and constraints

Pending

This section is pending the $O(\hbar^0)$ input from Sections 7 and 8, which is needed to decide between candidate amplitudes and fix $O(\hbar^0)$ ambiguities.

Requirements, exclusions and candidates. Without boost symmetry, a diagonal two-soliton S-matrix $S(P_1, P_2)$ must satisfy unitarity, the semiclassical limit $S \simeq e^{i\delta/\hbar}$ including the topological phase T_{12}, the crossing-type relation of Proposition 4.2, real analyticity and, for the bad equation, bound-state poles at the Bogdan–Charkina strings (23) (Yang–Baxter is automatic for diagonal scattering **(c)**). Stirling asymptotics force any Gamma factor to depend on the uniform weight $\eta = \frac{2}{9}(\ell_1 - \ell_2)/\hbar$, excluding at leading order both product forms $F(\beta_1 - \beta_2)G(\beta_1 + \beta_2)$ (Section 6) and Gamma functions of individual root differences **(c)** (`phase2/scripts/smatrix_rejected_forms.py`). For the bad equation, the candidate $S_{\mathrm{ov}} = \frac{\Gamma(1+i\eta)}{\Gamma(1-i\eta)}(e^2 A_{12}/\eta^2)^{i\eta} e^{iT_{12}/\hbar}$ and $S_{\mathrm{ho}} = \frac{\Gamma(1-i\eta)}{\Gamma(1+i\eta)}(A_{12}\eta^2/e^2)^{i\eta} e^{iT_{12}/\hbar}$ **(d)** satisfies unitarity, crossing, the semiclassical limit up to $\pm\pi/2$, and has simple poles on the physical sheet at the Bogdan–Charkina levels $I_{\mathrm{br}} = N\hbar$ $(N \geq 1)$ **(c)** (`phase2/smatrix_ansatz.pdf`); CDD factors, $\hbar$ renormalisation, Maslov constants and the relationship to the good-equation amplitude remain open **(d)**.

10 Open problems

Pending

Draft, based on the ranked next steps of the master report; to be revised when Sections 7–9 are written. The ranking is a research judgement **(d)**.

1. **A common quantum momentum and frame dictionary.** Relate the dressed Toda momentum, the quantum edge reference, the collective soliton charges and the frame contribution at fixed physical observables, with the endpoint reciprocity identity of the Toda dictionary audit as a check. The direct Toda expansion can proceed independently, keeping logarithmic terms and separating numerical errors.

2. **The projected one-loop determinant.** Starting from the factorised collective symplectic density, specify the spatial domain and admissible endpoint variations, and derive the field and constraint measures, the projection, the time-boundary determinant, continuum completeness and the absolute phase normalisation together.

3. **Quantum defects.** Before any quantum fusion argument, derive the defect's localised charges and boundary renormalisation; keep the admissible spectral branches; distinguish strictly stable external defect states (one per distinct real KP triple, $|s| < 2$, $s \neq \pm 1$) from unstable intermediate ones.

4. **Uniformity of the Toda expansion.** Check logarithmic remainders, nearly equal amplitudes and the limit of a vanishing smaller soliton, and establish which limits commute before identifying expansion coefficients with continuum loop orders.

5. **All-loop kernels and ring quantisation.** An asymptotic Bethe ansatz on a ring, $P_j\Lambda + \sum_{k\neq j}\delta(j,k) = 2\pi\hbar n_j$, is natural **(d)**, but the twisted boundary conditions forced by the ϕ-winding M have not been worked out. Quantum-dilogarithm kernels remain a possibility, to be constrained only by established input. Non-ultralocal quantum inverse scattering (Proposition 6.1) is a separate, longer-term route.

Further open questions raised by the classical analysis:

- *Number-changing processes.* Whether the Y-junctions, virtual solitons and resonance lines of the good equation produce inelastic quantum channels **(b)** [12], or appear only as threshold or bound-state singularities of the elastic amplitude **(d)**.

- *The cusp.* A uniform semiclassical treatment near $\alpha = \pi/3$, where $dP/d\alpha = 0$, the one-soliton symplectic form degenerates and the overtaking resonance line passes through $\alpha_1 = \alpha_2 = \pi/3$ **(d)**.
- *Bound states of the good equation.* The good equation has no regular Hirota breathers, so the existence of quantum bound states is open.
- *Finite-amplitude bad solitons.* No integrable lattice is known that agrees with Boussinesq at the dispersion scale **(d)**.
- *Defect reflection and decay.* Reflection and two-soliton decay at Corrigan–Zambon defects, and type-II defects for the bad equation **(d)**.
- *Universality of the phase structure.* Test whether $\delta = (\text{difference of spectral parameters}) \times \log A - c\,(\Pi_1 M_2 - \Pi_2 M_1)$ holds for KdV and other models with a continuous kink charge **(d)**.

A Claims and verification

All verification scripts are in the project repository; paths are relative to the repository root. Every verification script except the legacy `phase0/checks/convention_check.py` (which asserts its two symbolic residuals and prints `good c = -3` and `bad c = 3`, but omits the final marker) ends by printing `ALL CHECKS PASSED`. The scripts written specifically for this paper and its review are `paper/scripts/paper_conventions_check.py`, `paper/scripts/review_draft1_checks.py` and `paper/scripts/paper_draft2_checks.py`; they supply asserted checks for every formula revised or added in this paper, as well as for items that earlier scripts only printed as diagnostics.

Claim	Eq.	Lab.	Verification
Sign conventions; Lagrangians (3)	both	**(c)**	`phase0/checks/convention_check.py` (no final marker), `phase1/scripts/charges.py`, `paper/scripts/paper_conventions_check.py` (1–2)
Source field dictionaries (BD, CL, BC)	both	**(c)**	`paper/scripts/review_draft1_checks.py`, `paper/scripts/paper_draft2_checks.py` (9)
Real/imaginary Lax time	both	**(c)**	`phase1/scripts/lax_zero_curvature.py`
Charges (8), $dE/dP = v$, (9)	both	**(c)**	`phase1/scripts/charges.py`, `phase2/scripts/semiclassical_phase.py`, `paper/scripts/paper_conventions_check.py` (3, 9)
Bounded P, cusp, heavy branch; $\Omega\|_{\text{sol}} = -(P'(k)/k)\,dk \wedge dx_0$ (Prop. 2.1)	good	**(c)**	`phase1/scripts/charges.py` (P, E); `paper/scripts/review_draft1_checks.py`, `paper/scripts/paper_draft2_checks.py` (1, $\Omega\|_{\text{sol}}$ in x_0)
KP roots, $\ell(p)$, $\lambda = -\ell/(12\sqrt{3})$, ordered $\mu_{1,2} = -p_{b,a}$	both	**(c)**	`corrigan-zambon/defect_transmission_phase.py` (2), `phase2/scripts/smatrix_ansatz.py`, `paper/scripts/paper_conventions_check.py` (4), `paper/scripts/paper_draft2_checks.py` (2)

Claim	Eq.	Lab.	Verification
Vacuum-subtracted H_n on solitons	good	**(c)**	`phase1/scripts/charges.py`, `paper/scripts/review_draft1_checks.py`, `paper/scripts/paper_draft2_checks.py` (2)
N-soliton Hirota condition ($N \leq 4$ rational grid)	good	**(c)**	`phase1/scripts/hirota_solitons.py` (rational amplitudes, $N \leq 4$; separate profile check disabled by `or True`)
Closed forms of A_{12}; cross-ratio (Prop. 2.2)	both	**(c)**	`phase1/scripts/interaction_regimes.py`, `phase2/scripts/consistency_checks.py` (5), `corrigan-zambon/defect_two_soliton_symplectic.py` (1), `paper/scripts/paper_conventions_check.py` (4), `paper/scripts/paper_draft2_checks.py` (2g)
Regimes and resonances (Prop. 2.3); shifts (16)	good	**(c)**	`phase1/scripts/interaction_regimes.py`, `paper/scripts/paper_draft2_checks.py` (3)
Breathers (Prop. 2.4)	both	**(c)**	`phase1/scripts/interaction_regimes.py`, `phase2/scripts/consistency_checks.py` (4)
Continuation $\alpha = i\beta$ (Remark 2.5)	both	**(c)**	`paper/scripts/paper_conventions_check.py` (5)
Cross terms; symplectic scattering (Props. 3.1–3.2)	both	**(c)**	`phase2/scripts/symplectic_scattering.py` (bad), `phase2/scripts/consistency_checks.py` (1, both), `corrigan-zambon/defect_two_soliton_symplectic.py` (3, good)
Darboux shifts; γ	both	**(c)**	`phase2/scripts/consistency_checks.py` (1), `paper/scripts/paper_conventions_check.py` (9)
Semiclassical phase (21) and root-gradient proof	both	**(c)**	`phase2/scripts/semiclassical_phase.py`, `phase2/scripts/consistency_checks.py` (1), `paper/scripts/paper_draft2_checks.py` (8)
Unitarity, crossing-type relation	both	**(c)**	`phase2/scripts/consistency_checks.py` (2–3), `paper/scripts/paper_conventions_check.py` (6)
Resonance lines, cusp branch (Prop. 4.2)	good	**(c)**	`corrigan-zambon/defect_two_soliton_symplectic.py` (5, resonance print & defect cusp), `paper/scripts/paper_draft2_checks.py` (8, two-body cusp)
Small-amplitude limits and ratios	both	**(c)**	`phase2/scripts/consistency_checks.py` (6), `paper/scripts/paper_conventions_check.py` (7), `paper/scripts/paper_draft2_checks.py` (3)
Breather action (22) and strings (23) (Prop. 4.3)	bad	**(c)**	`phase2/scripts/consistency_checks.py` (4), `paper/scripts/paper_conventions_check.py` (8) ((23) conditional on **(b)**)
Toda solitons, signed label velocity, A_T	Toda→bad	**(c)**	`phase2/scripts/toda_soliton_classical.py`, `phase2/scripts/toda_label_symplectic.py`, `phase1/scripts/toda_limit.py`, `paper/scripts/paper_draft2_checks.py` (4a)

Claim	Eq.	Lab.	Verification
Toda two-body reflection amplitude R	Toda→bad	**(c)**	`paper/scripts/review_draft1_checks.py`, `paper/scripts/paper_draft2_checks.py` (4b, Bessel eigenfunction & asymptotics)
Classical Bethe-ansatz limit (Prop. 4.4)	Toda→bad	**(c)**	`phase2/scripts/toda_bethe_classical.py`, `phase2/scripts/toda_dressed_phase_check.py`, `phase2/scripts/toda_ba_vs_label.py`
Toda label-frame phase; scaling (Props. 4.5–4.6)	Toda→bad	**(c)**	`phase2/scripts/toda_label_symplectic.py`, `phase2/scripts/toda_scaling_fast.py`, `phase2/scripts/toda_ba_vs_label.py`
No real type-I defect for the bad equation	bad	**(c)**	`corrigan-zambon/defect_check.py`
CZ dictionary; sewing conditions	good	**(c)**	`corrigan-zambon/defect_transmission_phase.py` (1), `corrigan-zambon/verify_defect_solutions.py`, `paper/scripts/review_draft1_checks.py`
Root form of z; delay (Prop. 5.1)	good	**(c)**	`corrigan-zambon/defect_transmission_phase.py` (2–3), `paper/scripts/paper_conventions_check.py` (10)
$\Omega_D = (g_u - f_v)\, du \wedge dv = 0$ on vacua	good	**(c)**	`corrigan-zambon/defect_transmission_phase.py` (6), `paper/scripts/paper_draft2_checks.py` (5)
Transmission phase (27)	good	**(c)**	`corrigan-zambon/defect_transmission_phase.py` (4)
Two solitons and a defect (Prop. 5.3)	good	**(c)**	`corrigan-zambon/defect_two_soliton_symplectic.py` (2–4)
Vacua ↔ KP triple; absorption balance	good	**(c)**	`corrigan-zambon/defect_transmission_phase.py` (5), `paper/scripts/paper_draft2_checks.py` (6)
Linear stability	good	**(c)**/**(d)**	`corrigan-zambon/defect_vacuum_stability.py` (sampled **(c)**; completeness **(d)**), `paper/scripts/paper_draft2_checks.py` (6, Rolle real-triple count **(c)**)
Tree-level fusion (28) (zero-constant reps, ρ)	good	**(c)**	`phase2/scripts/defect_smatrix_good_continuation.py` (4), `paper/scripts/paper_draft2_checks.py` (7)
Non-ultralocality (Prop. 6.1)	good	**(c)**	`phase2/scripts/routes_BC_checks.py` (B), `paper/scripts/review_draft1_checks.py`
$\mathfrak{sl}_3$ factorisation (Prop. 6.2)	both	**(c)**	`phase2/scripts/routes_BC_checks.py` (C), `phase1/scripts/interaction_regimes.py`, `paper/scripts/review_draft1_checks.py`
No product-form S-matrix in rapidities ($\partial_S \partial_D \delta \neq 0$)	both	**(c)**	`phase2/scripts/smatrix_rejected_forms.py` (bad), `paper/scripts/paper_draft2_checks.py` (10, both)
Interpretations, completeness, absolute fusion constant, fixed-species Toda limit	—	**(d)**	—

Runs for this draft. On 4 November 2026 `paper/scripts/paper_conventions_check.py`, `paper/scripts/review_draft1_checks.py` and `paper/scripts/paper_draft2_checks.py`

were run and ended with `ALL CHECKS PASSED`. In addition, the independent review (`paper/review_draft1.pdf`) reran all 24 read-only verification scripts cited in Sections 1–6 (excluding `phase1/scripts/figures.py`, which writes a figure); 23 printed `ALL CHECKS PASSED` and `phase0/checks/convention_check.py` exited with code 0 after asserting its symbolic sign residuals.

Failures recorded.

1. In `paper/scripts/paper_conventions_check.py`, the good-equation crossing check first used the head-on point $(\alpha_1, \alpha_2) = (0.9, 0.15)$, which lies in the singular region $A_{\text{ho}} < 0$. There the identity holds only modulo the branch of the logarithm (residual 5×10^{-3} in the imaginary part). The point was replaced by $(1.3, 0.2)$ in the regular region, and the statement in Proposition 4.2 is restricted to the regular head-on region accordingly.

2. `phase2/scripts/consistency_checks.py` prints, but does not assert, the bad-equation crossing relation; it is asserted in `paper/scripts/paper_conventions_check.py` (check 6). Likewise, `corrigan-zambon/defect_two_soliton_symplectic.py` (check 5) prints resonance-line limits and asserts the soliton–defect cusp, while the two-body square-root cusp expansion in Proposition 4.2 is asserted in `paper/scripts/paper_draft2_checks.py` (check 8). In `phase1/scripts/hirota_solitons.py`, one asymptotic-profile check is disabled by `or True`, while its exact rational-amplitude bilinear checks for $N \leq 4$ are active and pass.

3. During the development of `paper/scripts/paper_draft2_checks.py`: (a) the first draft of check 2a declared the pseudo-differential coefficients as constant symbols rather than functions of x and tested guessed vacuum residues $1/144$ and $-1/864$ for $n = 4, 5$; declaring x-dependent coefficients and computing the exact binomial expansion of $L_0^{n/3} = (\partial_x^3 - \frac{1}{4}\partial_x)^{n/3}$ gave $\operatorname{Res} L_0^{4/3} = 0$ and $\operatorname{Res} L_0^{5/3} = 5/5184$, which then passed; (b) check 3c initially compared the 1.7 small-amplitude ratio against manually typed digits beyond the 12th place ('...222' instead of '0.564246498175008996'), which failed at 2×10^{-13} and was corrected to the exact value; (c) check 8b initially called `sp.simplify(id_expr.rewrite(sp.exp))` on the unfactored Hirota quotient, which was slow and was replaced by rational cancellation `sp.cancel` on the root cross-ratio derivative; and (d) checks 8c–8d initially used `sp.series(..., 0, 4)` (including the $O((\delta\alpha)^3)$ term) and wrote $-\Pi_1'(\pi/3)M_2 = \sqrt{3}\epsilon_1 M_2$ instead of $-\epsilon_1 M_2$ (since $\Pi_1 = -\epsilon_1 \sin 2\alpha_1 \implies \Pi_1'(\pi/3) = -2\epsilon_1 \cos(2\pi/3) = +\epsilon_1$), which failed and was corrected in both the script and Proposition 4.2. Failures recorded in earlier phases and in the independent review are listed in `phase1/review.pdf`, `phase2/report.pdf`, `phase2/defect_semiclassics.pdf` and `paper/review_draft1.pdf`.

B Inconsistencies found and their resolution

The following inconsistencies between project write-ups (or between a write-up and its scripts) were identified while preparing Draft 1 and audited in detail in `paper/review_draft1.pdf` (tested in `paper/scripts/review_draft1_checks.py`). None was silently resolved; for each item we record the verified resolution and its downstream effect.

1. **Sign of the cubic term in the bad Lagrangian.** `phase2/report.tex` (§1) prints $\mathcal{L} = \frac{1}{2}\phi_t^2 - \frac{1}{2}\phi_x^2 + \frac{1}{2}\phi_{xx}^2 + \phi_x^3$ for $v_{tt} = v_{xx} + v_{xxxx} + 3(v^2)_{xx}$. Its Euler–Lagrange equation is $v_{tt} = v_{xx} + v_{xxxx} - 3(v^2)_{xx}$, which the positive soliton does not solve (and on the positive pulse that wrong Lagrangian would give energy $2k^3/3$, differing from E_{bad} by $-8k^5/15$) **(c)**. The bad equation with positive solitons and $E_{\text{bad}} = \frac{2}{15}k^3(5 + 4k^2)$

requires $-\phi_x^3$, as used here (`paper/scripts/paper_conventions_check.py`, checks 1–3; `paper/scripts/review_draft1_checks.py`, A1). *Downstream effect:* The classical phase, M, Π, P and the classical energy formula use the correct data. However, the wrong cubic sign propagated into the normal-ordering boundary energy shift claimed in `phase2/oneloop.tex` (the "Decision (PI)" paragraph and energy table caption) and the docstring of `phase2/scripts/oneloop_soliton_energy.py` (lines 15–16): because the corrected bad Hamiltonian contains $+v^3$, the Wick replacement $v^3 \mapsto :v^3: = v^3 - 3Cv$ gives a *negative* shift $\Delta E_{\text{Wick}} = -3CM = -\frac{3\hbar}{2\pi}M = -\frac{3\hbar}{\pi}\sinh\beta$ (with $C = \hbar/(2\pi)$ on the stable band), opposite to the positive shift stated there **(c)**. The runtime code of `oneloop_soliton_energy.py` and `oneloop_breather.py` and the numerical tables in `phase2/oneloop.tex` evaluate the zero-scheme integral (C-shift omitted) and are unaffected **(c)**.

2. **Classical soliton energy in the one-loop write-up.** `phase2/oneloop_phase.tex`, eq. (onesol_data), prints $E_{\text{printed}} = \frac{1}{3}k^3(2\cos^2\alpha + 1)$ for the good soliton and asserts $dE/d\alpha = v\,dP/d\alpha$. Direct integration and 65-digit quadrature give $E_{\text{good}} = \frac{2}{15}k^3(5 - 4k^2)$ (as in `phase1/review.tex`); E_{printed} exceeds E_{good} by $k^3(5 - 2k^2)/15$ and violates the velocity identity with residual $dE_{\text{printed}}/dk - v\,dP/dk = k^2(1 - \frac{2}{3}k^2) \neq 0$ for $0 < k < 1$ **(c)** (`paper/scripts/review_draft1_checks.py`, A2). *Downstream effect:* None in the inspected one-loop calculations **(c)**. Specifically, `oneloop_good_phonon_scattering.py` constructs modes from the PDE and Hirota data without using E_{good}; `oneloop_good_twosoliton_phase.py` and `oneloop_good_limits_cusp_resonance.py` use derivatives of the classical phase and momentum; `charge_and_frame_audit.py` computes $L_1 = v\Delta P_{\text{mode}} - \Delta E_1$ from mode sums; and `audit_charges_action.py:ecl` explicitly uses the correct $E_{\text{good}} = \frac{2}{15}\sin^3\alpha(1 + 4\cos^2\alpha)$ **(c)**. (This clears propagation of the E_{printed} typo, without resolving the separate physical one-loop determinant and frame questions in Section 7 **(d)**.)

3. **Defect root convention.** `phase2/defect_semiclassics.tex` and `corrigan-zambon/defect_transmission_phase.py` use $s = \sigma_0$, $q = -s/(2\sqrt{3})$, $\alpha_{\text{CZ}} = \eta/\sqrt{3}$, $\beta_{\text{CZ}} = 2\eta/\sqrt{3}$ and $z = ((p_b - q)/(p_a - q))^\eta$, with strict stability $\eta\sigma < -1$ ($\eta q > 1/(2\sqrt{3})$). By contrast, `phase2/oneloop_defect_phase.tex` and Route E of `phase2/report.tex` write $s = \eta\sigma$ and $q_D = -\eta\sigma/(2\sqrt{3})$ with $\alpha_{\text{CZ}} = 1/(2\sqrt{3})$, $\beta_{\text{CZ}} = \eta/(2\sqrt{3})$. *Resolution and downstream effect:* Consultation of the full text of [12] (sections 3 and 5, eqs. (3.22), (3.37), (3.43)–(3.44) and (5.5)) settles the convention **(a)**/**(c)**: CZ's sewing constraints require $\beta_{\text{CZ}} = 2\alpha_{\text{CZ}}$ and $\beta_{\text{CZ}}^2 - \alpha_{\text{CZ}}^2 = 1$ (for $c = 1$), which the Phase 2c pair $(\eta/\sqrt{3}, 2\eta/\sqrt{3})$ satisfies and the later pair $(1/(2\sqrt{3}), \eta/(2\sqrt{3}))$ violates **(c)**. At a fixed physical defect with $\eta = -1$ (e.g. $\alpha = 0.4, \epsilon = +1, \sigma_0 = 2.4, \eta = -1$), replacing q by $-\eta\sigma_0/(2\sqrt{3})$ changes z from $1.50971\ldots$ to $0.373565\ldots$ and fails the nonlinear sewing conditions with residual 3.33, whereas the Phase 2c root satisfies sewing to $< 10^{-52}$ **(c)** (`paper/scripts/review_draft1_checks.py`, A3). Both executable scripts `oneloop_defect_scattering_and_stability.py` and `oneloop_defect_phase_and_fusion.py` retain the unsigned root $s = -2\sqrt{3}q$, so their computed functions and formal chord identities are unaffected, though interpreting them at $\eta = -1$ in terms of σ_0 requires the Phase 2c dictionary used here **(c)**. (Two nearby display typos in `phase2/oneloop_defect_phase.tex`—writing $\ell/(4\sqrt{3})$ instead of $\ell/(12\sqrt{3})$ in the cubic and inequivalent trigonometric formulas for $p_{a,b}$—are likewise absent from the scripts and from this paper **(c)**.)

4. **Transmission factor in the one-loop defect write-up.** In `phase2/oneloop_defect_phase.tex`, eq. (z1_sol), the middle expression $\big((k_1 - v_1/\sqrt{3} - s/\sqrt{3})/(k_1 + v_1/\sqrt{3} + s/\sqrt{3})\big)^{-\eta}$ equals $-1/z$, a genuine sign and reciprocal error relative to $z = ((v_1 + s - \sqrt{3}k_1)/(v_1 + s + \sqrt{3}k_1))^\eta = ((p_b - q)/(p_a - q))^\eta$ **(c)** (`paper/scripts/paper_conventions_`

`check.py`, check 10; `paper/scripts/review_draft1_checks.py`, A4). *Downstream effect:* None **(c)**. Inspection and AST evaluation of `z_defect_mode` in the `oneloop_defect_*` scripts confirm that all executable transmission, delay, phase and fusion calculations use the correct rational/root expression **(c)**.

5. **Wording of the KdV-limit remark.** `phase2/report.tex` (Remark after Prop. 2.3) states that the topological term and the KdV-type term $2\beta_1 \log A$ have ratio tending to 0.564 at $\beta_1/\beta_2 = 1.7$. *Resolution and downstream effect:* At $\beta_1/\beta_2 = 1.7$, $0.564246498175\ldots = 1 + 2\beta_2/(\beta_1 \log A_{\mathrm{KdV}})$ is the ratio of the *full* phase derivative $\partial_{\beta_1}\delta$ to $2\beta_1 \log A_{12}$, while the cross-term-only ratio $(-\Pi_1' M_2)/(2\beta_1 \log A_{12})$ is $-0.435753501825\ldots$ **(c)** (`paper/scripts/review_draft1_checks.py`, A5; `paper/scripts/paper_draft2_checks.py`, check 3). In addition, the final explanatory print string of `phase2/scripts/consistency_checks.py` has a minus-sign typo in its printed formula, while its computed numerical ratio and all phase values are correct **(c)**.

6. **Script names in the master report.** The appendix of `phase2/report.tex` cites `smatrix_boot.py` and `toda_quantum_correction.py`; the filesystem inventory confirms that the existing scripts in `phase2/scripts/` are `smatrix_poles_bootstrap.py` and `toda_quantum_corrections.py` (and that `smatrix_semiclassical_unitarity_crossing.py` and `smatrix_rejected_forms.py` are also present there), with no effect on calculations **(c)** (`paper/scripts/review_draft1_checks.py`, A6).

7. **Root variables (notation only).** Phase 1 uses ordered real roots μ_1, μ_2 with $\mu_1 - \mu_2 = k$ and $\mu_1 + \mu_2 = -v/\sqrt{3}$; Phase 2c uses p_a, p_b with $p_a - p_b = k$ and $p_a + p_b = +v/\sqrt{3}$. They are related by the order-reversing sign map $\mu_1 = -p_b$, $\mu_2 = -p_a$, with Lax eigenvalue $\lambda = \mu_j^3 - \mu_j/4 = -\ell/(12\sqrt{3})$, while for the bad equation `phase2/smatrix_ansatz.tex` uses the complex roots $\mu_{a,b} = \frac{1}{2}(iv/\sqrt{3} \pm k)$ satisfying $\mu^3 + \mu/4 = i\ell/(12\sqrt{3})$ **(c)** (`paper/scripts/paper_conventions_check.py`, check 4; `paper/scripts/paper_draft2_checks.py`, check 2). No downstream error; both are stated explicitly in Section 2.3.

8. **Toda wavenumber (notation only).** In `phase1/review.tex` and `phase1/scripts/toda_limit.py`, κ_H is the Hirota exponent ($\tau = 1 + e^{\kappa_H n - \omega t}$), whereas in `phase2/toda_route.tex` and Section 4.4 here κ_T is the $\mathrm{sech}^2(\kappa_T n - \ldots)$ half-width parameter. Thus $\kappa_H = 2\kappa_T$ exactly, while the continuum identifications $\kappa_H \sim \sqrt{12}\, k_{\mathrm{bad}}$ and $\kappa_T \sim \sqrt{3}\,\beta$ are small-amplitude asymptotics ($\sinh\beta/\beta \to 1$) **(c)** (`paper/scripts/review_draft1_checks.py`, A8).

9. **Phase numbering (minor).** `phase2/oneloop_defect_phase.tex` cites `defect_semiclassics.pdf` as Phase 2d; its title and `README.md` identify it as Phase 2c **(c)** (`paper/scripts/review_draft1_checks.py`, A9).

10. **Bibliography and literature-attribution corrections.** The repository-wide reference audit (`phase0/reference_audit.md`) identified five substantive bibliography or attribution errors across earlier write-ups, all resolved here: (a) [12] was cited with wrong article numbers ('315202'/'345203'/'405202'), a wrong title in two Phase 2 notes, or a non-existent DOI ('10.1088/1751-8121/ace204'); the published record is *J. Phys. A* **56** (2023) 385701, `doi:10.1088/1751-8121/aceec9`; (b) `phase2/toda_wh_and_kp_synthesis.tex` gave a fabricated good-equation title for [4], which treats the *bad* equation; (c) `phase1/review.tex` cited [20] directly for the good-equation bilinear form, whereas [20] treats the bad equation ($D_t^2 - D_x^2 - D_x^4$); (d) `phase1/review.tex` cited [5] for small-data global existence, whereas its abstract proves nonlinear solitary-wave stability and global existence for data close to a stable solitary wave; and (e) `phase1/review.tex` cited Mathieu and

Oevel (*Mod. Phys. Lett. A* **6** (1991) 2397–2409, doi:10.1142/S0217732391002827) with title "W_3" instead of "$W_3^{(2)}$" for the standard Zamolodchikov W_3 bracket, whereas that paper treats Polyakov's $W_3^{(2)}$ algebra and an $x \leftrightarrow t$ interchanged hierarchy (and is therefore not cited here).

References

[1] S. K. Ashok, S. Parihar, T. Sengupta, A. Sudhakar, R. Tateo, Integrable structure of higher spin CFT and the ODE/IM correspondence, *JHEP* 07 (2024) 179, arXiv:2405.12636, doi:10.1007/JHEP07(2024)179. [full text read in Phase 0; arXiv abstract and DOI metadata re-verified]

[2] S. K. Ashok, S. Parihar, T. Sengupta, A. Sudhakar, R. Tateo, Thermal correlators and currents of the $\mathcal{W}_3$ algebra, *JHEP* 01 (2025) 154, arXiv:2410.11748, doi:10.1007/JHEP01(2025)154. [full text read in Phase 0; arXiv abstract and DOI metadata re-verified]

[3] V. V. Bazhanov, A. N. Hibberd, S. M. Khoroshkin, Integrable structure of W_3 conformal field theory, quantum Boussinesq theory and boundary affine Toda theory, *Nucl. Phys. B* 622 (2002) 475–547, arXiv:hep-th/0105177, doi:10.1016/S0550-3213(01)00595-8. [full text read in Phase 0; arXiv abstract and DOI metadata re-verified]

[4] M. M. Bogdan, O. V. Charkina, Quasiclassical quantization of the Boussinesq breather emerging from the kink localized mode, *Low Temp. Phys.* 51 (2025) 763–773, arXiv:2502.14695, doi:10.1063/10.0036799. [full text read; arXiv abstract and DOI landing page re-verified]

[5] J. L. Bona, R. L. Sachs, Global existence of smooth solutions and stability of solitary waves for a generalized Boussinesq equation, *Commun. Math. Phys.* 118 (1988) 15–29, doi:10.1007/BF01218475. [abstract only; DOI landing page and abstract re-verified]

[6] T. Bonnemain, B. Doyon, Soliton gas of the integrable Boussinesq equation and its generalised hydrodynamics, *SciPost Phys.* 18 (2025) 075, arXiv:2402.08669, doi:10.21468/SciPostPhys.18.2.075. [full text read; arXiv abstract and DOI metadata re-verified]

[7] A. J. Bordner, Commuting charges of the quantum Korteweg–de Vries and Boussinesq theories from the reduction of W_∞ and $W_{1+\infty}$ algebras, *Mod. Phys. Lett. A* 13 (1998) 541–552, arXiv:solv-int/9711013, doi:10.1142/S0217732398000607. [full text read in Phase 0; arXiv abstract and DOI metadata re-verified]

[8] C. Charlier, J. Lenells, The "good" Boussinesq equation: a Riemann–Hilbert approach (2020), arXiv:2003.02777. [full text read in Phase 0; arXiv abstract re-verified]

[9] C. Charlier, J. Lenells, On Boussinesq's equation for water waves (2022), arXiv:2204.02365. [full text read in Phase 1; arXiv abstract re-verified]

[10] C. Charlier, J. Lenells, Direct and inverse scattering for the Boussinesq equation with solitons (2023), arXiv:2302.14593. [full text read in Phase 1; arXiv abstract re-verified]

[11] C. Charlier, J. Lenells, The soliton resolution conjecture for the Boussinesq equation (2023), arXiv:2303.10485. [full text read in Phase 0; arXiv abstract re-verified]

[12] E. Corrigan, C. Zambon, Adding integrable defects to the Boussinesq equation, *J. Phys. A* 56 (2023) 385701, arXiv:2306.03772, doi:10.1088/1751-8121/aceec9. [full text read; arXiv abstract and DOI landing page re-verified]

[13] P. Deift, C. Tomei, E. Trubowitz, Inverse scattering and the Boussinesq equation, *Commun. Pure Appl. Math.* 35 (1982) 567–628, doi:10.1002/cpa.3160350502. [abstract only; DOI metadata re-verified]

[14] L. A. Dickey, *Soliton Equations and Hamiltonian Systems*, World Scientific (2nd ed. 2003), doi:10.1142/5108. [not read; publisher DOI landing page and metadata verified]

[15] A. Dimakis, F. Müller-Hoissen, X.-M. Chen, Matrix Boussinesq solitons and their tropical limit, *Phys. Scr.* 94 (2019) 035206, arXiv:1805.09711, doi:10.1088/1402-4896/aaf6da. [full text read in Phase 0; arXiv abstract and DOI metadata re-verified]

[16] L. D. Faddeev, V. E. Korepin, Quantum theory of solitons, *Phys. Rep.* 42 (1978) 1–87, doi:10.1016/0370-1573(78)90058-3. [full text read in Phase 0; DOI metadata re-verified]

[17] G. M. Gandenberger, Trigonometric S-matrices, affine Toda solitons and supersymmetry, *Int. J. Mod. Phys. A* 13 (1998) 4553–4590, arXiv:hep-th/9703158, doi:10.1142/S0217751X98002195. [full text read in Phase 0; arXiv abstract and DOI metadata re-verified]

[18] M. C. Gutzwiller, The quantum mechanical Toda lattice, II, *Ann. Phys.* 133 (1981) 304–331, doi:10.1016/0003-4916(81)90253-0. [abstract only; DOI landing page and abstract re-verified]

[19] K. Hikami, Baxter equation for quantum discrete Boussinesq equation, *Nucl. Phys. B* 604 (2001) 580–602, arXiv:nlin/0102021, doi:10.1016/S0550-3213(01)00204-8. [full text read in Phase 0; arXiv abstract and DOI landing page re-verified]

[20] R. Hirota, Exact N-soliton solutions of the wave equation of long waves in shallow-water and in nonlinear lattices, *J. Math. Phys.* 14 (1973) 810–814, doi:10.1063/1.1666400. [abstract only; DOI landing page and abstract re-verified]

[21] R. Hirota, M. Ito, Resonance of solitons in one dimension, *J. Phys. Soc. Japan* 52 (1983) 744–748, doi:10.1143/JPSJ.52.744. [cited via [15]; not read; DOI landing page and metadata verified]

[22] T. J. Hollowood, Solitons in affine Toda field theories (arXiv title: Quantum solitons in affine Toda field theories), *Nucl. Phys. B* 384 (1992) 523–540, arXiv:hep-th/9110010, doi:10.1016/0550-3213(92)90579-Z. [full text read in Phase 0; arXiv abstract and DOI landing page re-verified]

[23] T. J. Hollowood, Quantizing $SL(N)$ solitons and the Hecke algebra, *Int. J. Mod. Phys. A* 8 (1993) 947–981, arXiv:hep-th/9203076, doi:10.1142/S0217751X93000370. [full text read in Phase 0; arXiv abstract and DOI metadata re-verified]

[24] P. R. Johnson, On soliton quantum S-matrices in simply laced affine Toda field theories (arXiv title: Exact quantum S-matrices for solitons in simply-laced affine Toda field theories), *Nucl. Phys. B* 496 (1997) 505–550, arXiv:hep-th/9611117, doi:10.1016/S0550-3213(97)00239-3. [full text read in Phase 0; arXiv abstract and DOI landing page re-verified]

[25] B. A. Kupershmidt, P. Mathieu, Quantum Korteweg–de Vries like equations and perturbed conformal field theories, *Phys. Lett. B* 227 (1989) 245–250, doi:10.1016/S0370-2693(89)80030-9. [abstract only; DOI metadata re-verified]

[26] F. Lambert, M. Musette, E. Kesteloot, Soliton resonances for the good Boussinesq equation, *Inverse Problems* 3 (1987) 275–288, doi:10.1088/0266-5611/3/2/010. [abstract only, conditions via [6]; DOI landing page and abstract re-verified]

[27] J.-M. Maillet, Kac–Moody algebra and extended Yang–Baxter relations in the O(N) non-linear sigma model, *Phys. Lett. B* 162 (1985) 137–142, doi:10.1016/0370-2693(85)91075-5. [not read; DOI metadata re-verified]

[28] D. Masoero, A. Raimondo, Opers for higher states of the quantum Boussinesq model, in *Asymptotic, Algebraic and Geometric Aspects of Integrable Systems*, Springer (2020) 55–78, arXiv:1908.11559, doi:10.1007/978-3-030-57000-2_5. [full text read in Phase 0; arXiv abstract and DOI landing page re-verified]

[29] F. G. Mertens, Bethe ansatz for the Toda lattice: ground state and excitations, *Z. Phys. B* 55 (1984) 353–360, doi:10.1007/BF01304087. [abstract only; DOI landing page and abstract re-verified]

[30] M. Pustilnik, K. A. Matveev, Fate of classical solitons in one-dimensional quantum systems, *Phys. Rev. B* 92 (2015) 195146, arXiv:1507.05639, doi:10.1103/PhysRevB.92.195146. [full text read in Phase 0; arXiv abstract and DOI metadata re-verified]

[31] A. G. Rasin, J. Schiff, Bäcklund transformations for the Boussinesq equation and merging solitons, *J. Phys. A* 50 (2017) 325202, arXiv:1701.00243, doi:10.1088/1751-8121/aa7af7. [abstract and introduction read in Phase 0; arXiv abstract and DOI metadata re-verified]

[32] R. Siddharthan, B. S. Shastry, Quantizing the Toda lattice (arXiv title: The quantum Toda lattice revisited), *Phys. Rev. B* 55 (1997) 12196–12209, arXiv:cond-mat/9611221, doi:10.1103/PhysRevB.55.12196. [full text read in Phase 0; arXiv abstract and DOI landing page re-verified]

[33] E. K. Sklyanin, The quantum Toda chain, *Lect. Notes Phys.* 226 (1985) 196–233, doi:10.1007/3-540-15213-X_80. [not read; publisher DOI landing page and metadata verified]

[34] B. Sutherland, A brief history of the quantum soliton with new results on the quantization of the Toda lattice, *Rocky Mountain J. Math.* 8 (1978) 413–430, doi:10.1216/RMJ-1978-8-1-413. [abstract only; DOI landing page and metadata re-verified]

[35] G. Takács, G. M. T. Watts, Non-unitarity in quantum affine Toda theory and perturbed conformal field theory, *Nucl. Phys. B* 547 (1999) 538–568, arXiv:hep-th/9810006, doi:10.1016/S0550-3213(99)00100-5. [abstract read in Phase 0; arXiv abstract and DOI metadata re-verified]

[36] V. E. Zakharov, On stochastization of one-dimensional chains of nonlinear oscillators, *Sov. Phys. JETP* 38 (1974) 108–110 [*Zh. Eksp. Teor. Fiz.* 65 (1973) 219–225]. [cited via [9, 6]; not read; no publisher DOI exists for this 1974 issue]

Historical "full text read" and "abstract only" labels are as recorded in `phase0/literature_map.md` and `phase1/review.tex` by project agents in earlier phases; for Draft 1 review and Draft 2 revision, the relevant full-text sections of [12] and [4] were consulted and all arXiv abstract pages and publisher DOI landing pages were re-verified against `phase0/reference_audit.md` as indicated in brackets.