

Lecture 3: Tasks Solutions

Gustave Kenedi


2026-02-06

Your Turn #1

```
library(tidyverse)

vote_share <- data.frame(prop_voters = c(0.25, 0.25, 0.35, 0.20, 0.20, 0.35,
                                          0.1, 0.35, 0.35, 0.2, 0.25, 0.25,
                                          0.2, 0.15, 0.25, 0.2, 0.15, 0.15, 0.2, 0.1))

vote_share |>
  ggplot(aes(x = prop_voters)) +
  geom_histogram()
```



Your Turn #2

Question 1

```
voter_population <- read.csv("https://www.dropbox.com/scl/fi/xdB5nzyoi7i5r9m80ae3r/population")
```

Question 2

```
library(moderndivide)

virtual_samples <- voter_population |>
  rep_sample_n(size = 50, reps = 1000)
```


Question 3

```
virtual_prop_pumpnickel <- virtual_samples |>
  group_by(replicate) |>
  summarise(prop_pumpnickel = mean(candidate == "Pumpnickel"))
```

Question 4

```
virtual_prop_pumpnickel |>  
  ggplot(aes(x = prop_pumpnickel)) +  
  geom_histogram()
```

``stat_bin()`` using ``bins = 30``. Pick better value ``binwidth``.



Question 5

Sampling distribution looks normally distributed. The most common voter shares are between 0.2 and 0.25. The shape of the histogram looks more bell-shaped than it did with 50 samples.

Question 6

It's rather unlikely that we sample 50 voters of which less than 15% are Pumpnickel voters.

Your Turn #3

Let $X_1, \dots, X_n$ be i.i.d. random variables with a finite expected value $E[X_i] = \mu$ and $\mathbb{V}(X_i) = \sigma^2 < \infty$.

Question 1

Show that $\mathbb{E}[S_n^2] = \mathbb{E}[X^2] - \mathbb{E}[\bar{X}_n^2]$.

Answer:

$$\mathbb{E}[S_n^2] = \mathbb{E} \left[\frac{1}{n} \sum_{i=1}^n (X_i - \bar{X}_n)^2 \right] \quad (1)$$

$$= \mathbb{E} \left[\frac{1}{n} \sum_{i=1}^n X_i^2 - \frac{1}{n} 2 \sum_{i=1}^n X_i \bar{X}_n + \frac{1}{n} \sum_{i=1}^n \bar{X}_n^2 \right] \quad (2)$$

$$= \frac{1}{n} \sum_{i=1}^n \mathbb{E}[X_i^2] - \mathbb{E} \left[\frac{1}{n} 2 \bar{X}_n \sum_{i=1}^n X_i \right] + \mathbb{E} [\bar{X}_n^2] \quad (3)$$

$$= \mathbb{E}[X^2] - \mathbb{E} \left[\frac{1}{n} 2 \bar{X}_n n \bar{X}_n \right] + \mathbb{E} [\bar{X}_n^2] \quad (4)$$

$$= \mathbb{E}[X^2] - 2\mathbb{E} [\bar{X}_n^2] + \mathbb{E} [\bar{X}_n^2] \quad (5)$$

$$= \mathbb{E}[X^2] - \mathbb{E} [\bar{X}_n^2] \quad (6)$$

Question 2

Show that $\mathbb{E}[X^2] = \sigma^2 + \mu^2$. Hint: recall the formula for the variance using expectations.

Answer:

Recall that $\mathbb{V}(X) = \mathbb{E}[X^2] - \mathbb{E}[X]^2$. We therefore have:

$$\mathbb{E}[X^2] = \mathbb{V}(X) + \mathbb{E}[X]^2 \quad (7)$$

$$= \sigma^2 + \mu^2 \quad (8)$$

Question 3

Show that $\mathbb{E}[\bar{X}_n^2] = \frac{\sigma^2}{n} + \mu^2$. *Hint: the formula for $\mathbb{V}(\bar{X}_n)$ will be useful.*

Answer:

Using the variance formula and replacing X by $\bar{X}_n$, we have:

$$\mathbb{E}[\bar{X}_n^2] = \mathbb{V}(\bar{X}_n) + \mathbb{E}[\bar{X}_n]^2 \quad (9)$$

$$= \frac{\sigma^2}{n} + \mu^2 \quad (10)$$

Question 4

Combine the above results to show that $\mathbb{E}[S_n^2] \neq \sigma^2$.

Answer:

We can use the results from questions 2 and 3:

$$\mathbb{E}[S_n^2] = \mathbb{E}[X^2] - \mathbb{E}[\bar{X}_n^2] \quad (11)$$

$$= \sigma^2 + \mu^2 - \left(\frac{\sigma^2}{n} + \mu^2\right) \quad (12)$$

$$= \sigma^2 - \frac{\sigma^2}{n} \quad (13)$$

$$= \frac{n-1}{n}\sigma^2 \quad (14)$$

$$\neq \sigma^2 \quad (15)$$

Question 5

Load this [voter data](#). Use **R** to compute the variance of Pumpernickel voters: (1) manually (i.e., taking the sum of deviations from the mean), and (2) using the **var** command. What do you notice?

Answer:

```
library(tidyverse)

voter_data <- read.csv("https://www.dropbox.com/scl/fi/lnz14t3a2cj6kw0l217yo/pumpnickel_voter_data.csv?dl=1")

voter_data |>
  mutate(deviation_from_mean = pumpnickel_voter - mean(pumpnickel_voter)) |>
  summarise(var_manual = 1/n() * sum(deviation_from_mean^2),
            var_command = var(pumpnickel_voter),
            var_manual_adjusted = n()/(n()-1) * var_manual)
```

```
  var_manual var_command var_manual_adjusted
1      0.1771    0.1772773          0.1772773
```

Question 6

Read the help `sd` (it is simpler than `var`) command's help file to see if you can understand what's going on.

```
?sd
```

Like var this uses denominator $n - 1$.

Your Turn #4

Question 1

Use `pnorm` to confirm the figure on slide 65, i.e., within ± 1 (2) standard deviation of the mean, the standard normal distribution contains 68% (95%) of observations.

Answer:

```
pnorm(q = 1, mean = 0, sd = 1) - pnorm(q = -1, mean = 0, sd = 1)
```

```
[1] 0.6826895
```

```
pnorm(q = 2, mean = 0, sd = 1) - pnorm(q = -2, mean = 0, sd = 1)
```

```
[1] 0.9544997
```

Question 2

A region has household incomes with mean $\mu = 58,000$ euros and standard deviation $\sigma = 15,000$ euros. An economist surveys 100 randomly selected households. What is the probability that the sample mean income is less than 55,000 euros?

Answer:

```
sample_mean = 55000
pop_mean = 58000
pop_sd = 15000
sample_size = 100
mean_se = pop_sd/sqrt(sample_size)

pnorm(sample_mean, mean = pop_mean, sd = mean_se)
```

```
[1] 0.02275013
```

Question 3

A country's true unemployment rate is 6.2%. A labor economist draws a random sample of 1,000 working-age adults. What is the probability that the sample unemployment rate falls between 5% and 7%?

Answer:

```
pop_prop = 0.062
pop_var = 0.062 * (1 - 0.062)
sample_size = 1000
se <- sqrt((pop_prop*(1-pop_prop)) / sample_size)

lower_bound = 0.05
upper_bound = 0.07

pnorm(upper_bound,
      mean = pop_prop, sd = se) -
  pnorm(lower_bound,
        mean = pop_prop, sd = se)
```

```
[1] 0.7951262
```

Question 4

Monthly consumer spending at retail stores has mean $\mu = 320$ euros and standard deviation $\sigma = 85$ euros. A retail analyst collects data from 50 randomly selected consumers. Below what value does the lowest 10% of sample mean spending fall?

Answer:

```
pop_mean = 320
pop_sd = 85
sample_size = 50
quantile = .1
qnorm(quantile, mean = pop_mean, sd = pop_sd/sqrt(sample_size))
```

```
[1] 304.5947
```