Assignment #5

Gustavo Estrela de Matos CSCE 433: Formal Languages and Automata

Monday 2nd May, 2016

Question 1.

- Eliminate the starting symbol from the right hand side
- Eliminate ϵ rules

$$S \to AB \mid aB \mid B$$

$$A \to aab$$

$$B \rightarrow bbA \mid bb$$

• Eliminate unity rules

$$S \rightarrow AB \mid aB \mid bbA \mid bb$$

$$A \to aab$$

$$B \rightarrow bbA \mid bb$$

• Eliminate rules with more than 2 symbols in the right hand side

$$S \rightarrow AB \mid aB \mid bY \mid bb$$

$$A \to aZ$$

$$B \rightarrow bY \mid bb$$

$$Y \to bA$$

$$Z \to ab$$

• Create unity rules for terminals

$$S \to AB \mid XB \mid VY \mid VV$$

$$A \to XZ$$

$$B \to VY \mid VV$$

$$Y \to VA$$

$$Z \to VX$$

$$V \to b$$

$$X \to a$$

Question 2.

- (a)
- (b)

```
q_100110 \vdash 0q_20110 \vdash 00q_2110 \vdash 000q_310 \vdash 0001q_30 \vdash 00010q_2 \vdash 0001q_40 \vdash 00010q_5 (c)
```

 $q_1111010 \vdash 1q_311010 \vdash 11q_31010 \vdash 111q_3010 \vdash 1110q_210 \vdash 11100q_30 \vdash 111000q_2 \vdash 11100q_40 \vdash 111000q_5$

Question 3.

(a)

The idea implemented in this TM is to match a's from left to right (replacing with A's) with the c's from right to left, and when this is done it should match b's from left to right with the A's from right to left.

(b)

We can split this machine in two other problems. First we divide the string in the middle, then we match every character of the two produced strings. To divide the string in two we are going to shift left the beggining of the string and the shift right the end of the string until the symbols we use for keeping track of the shifting meet in the middle of the first string. The second part is basically going to the beggining of the two strings and comparing the symbols.

Question 4.

This machine does the algorithm we normally do by hand, from right to left, bit by bit. The key idea is that we are summing N with 0 and starting the carry with 1. The carry is 1 when we are in the state q_1 and it is 0 when we are in the state q_2 .

Question 5.

As we saw in class, the symbol w_i is such that $1w_i = i_2$. Therefore, since $37 = (100101_2)$ we have that $1w_{37} = 100101$ and $w_{37} = 00101$.

Question 6.