

Molecular benchmark report: LiH, BeH₂, HF, and H₂O through `clifford_qc` A-CASE

September 18, 2026

Generated by `benchmarks/summarize_molecular.py`.

Source: `molecular/results/results_summary.json`. Do not edit by hand.

1 Results

Errors in mHa against the sector-exact ground energy of the reference’s own (N, S_z) sector; chemical accuracy is 1.5936 mHa.

Table 1: Adaptive A-CASE against classical baselines.

Molecule	n	N_e	Sector dim	M	A-CASE (mHa)	CCSD (mHa)	CISD (mHa)	SD (mHa)	$\kappa(S)$
BeH ₂	14	6	1225	11	+1.3411	+0.3934	+0.7526	+0.7526	1.00
BeH ₂ 2×	14	6	1225	31	+28.5144	+5.8613	+28.5110	—	1.00
H ₂ O	14	10	441	21	+1.4477	+0.1165	+0.7051	+0.7051	1.00
H ₂ O 2×	14	10	441	21	+55.4206	−9.7177	+54.5703	—	1.00
HF	12	10	36	5	+1.0725	+0.0000	+0.0000	+0.0000 [†]	1.00
LiH	12	4	225	6	+1.1264	+0.0105	+0.0133	+0.0133	1.00
LiH 3×	12	4	225	5	+1.5210	+0.1320	+0.1509	—	1.00

Reading the table. A-CASE reaches chemical accuracy on 5 of 7 geometries. CCSD has a smaller absolute error on 7 of 7 geometries. Negative CCSD errors are possible because coupled cluster is not variational. Some capped stretched runs miss chemical accuracy; inspect their errors and spin contamination alongside basis size.

The stopping rule is oracle-assisted. Where enabled, growth may halt when the error against the *exact* energy falls below threshold. Runs may also terminate at a budget or another stopping condition. The M column is the selected basis size, not a guarantee that the accuracy target was reached.

[†] The complete SD basis spans the entire sector for this molecule, so agreement with exact diagonalization is arithmetic rather than accuracy.

2 Observables

Read off the same solve that produced the adaptive energy, through the projected-observable route; no Ritz state is materialized.

Table 2: Double occupancy against its uncorrelated value, and the singlet check.

Molecule	$\langle d \rangle$	uncorrelated	difference	$\langle S^2 \rangle$
BeH ₂	0.426218803	0.428571429	$-2.35e-03$	$2.06e-04$
BeH ₂ 2 $\times$	0.395619366	0.428571429	$-3.30e-02$	$8.73e-18$
H ₂ O	0.711243361	0.714285714	$-3.04e-03$	$3.81e-04$
H ₂ O 2 $\times$	0.656131728	0.714285714	$-5.82e-02$	$2.66e-01$
HF	0.833034591	0.833333333	$-2.99e-04$	$3.01e-34$
LiH	0.332419497	0.333333333	$-9.14e-04$	$1.23e-32$
LiH 3 $\times$	0.325510212	0.333333333	$-7.82e-03$	$1.86e-02$

Conserving S_z does not enforce total spin. Nonzero $\langle S^2 \rangle$ indicates spin contamination of the selected state. A negligible projected S_z Lehmann response does not certify a zero physical susceptibility: relevant excited sectors and transition weights have not been validated.

3 Resource accounting

Table 3: Distinct Pauli words in H against the A-CASE element word universe $|\bigcup_{ij} \text{supp}(A_i^\dagger A_j) \cup \text{supp}(A_i^\dagger H A_j)|$. The two are different quantities.

Molecule	Words in H	Element universe	M	SD wall-clock (s)
BeH ₂	666	350,196	11	3018.3
BeH ₂ 2 $\times$	666	1,165,663	31	—
H ₂ O	1086	1,168,272	21	2461.3
H ₂ O 2 $\times$	1086	874,597	21	—
HF	631	21,156	5	54.2
LiH	631	38,250	6	440.3
LiH 3 $\times$	631	24,096	5	—