Publication list from all collaborations

- A.M. Sirunyan et al., **Inclusive search for a highly boosted Higgs boson decaying to a bottom quark-antiquark pair**, *Phys. Rev. Lett.* 120 (2018) 071802, doi:10.1103/PhysRevLett.120.071802, arXiv:1709.05543 [hep-ex]
- A.M. Sirunyan et al., Search for Higgs boson pair production in events with two bottom quarks and two tau leptons in proton-proton collisions at \sqrt{s} =13TeV, *Phys. Lett.* B778 (2018) 101–127, doi:10.1016/j.physletb.2018.01.001, arXiv:1707.02909 [hep-ex]
- A.M. Sirunyan et al., Search for supersymmetry with Higgs boson to diphoton decays using the razor variables at $\sqrt{s} = 13$ TeV, *Phys. Lett.* B779 (2018) 166–190, doi:10.1016/j.physletb.2017.12.069, arXiv:1709.00384 [hep-ex]
- A.M. Sirunyan et al., Search for resonant and nonresonant Higgs boson pair production in the $b\bar{b}$ final state in proton-proton collisions at $\sqrt{s}=13$ TeV, *JHEP*. 01 (2018) 054, doi:10.1007/JHEP01(2018)054, arXiv:1708.04188 [hep-ex]
- A.M. Sirunyan et al., Constraints on the double-parton scattering cross section from same-sign W boson pair production in proton-proton collisions at $\sqrt{s}=8$ TeV, JHEP.~02~(2018)~032,~doi:10.1007/JHEP02(2018)032,~arXiv:1712.02280~[hep-ex]
- A.M. Sirunyan et al., Search for supersymmetry in events with at least three electrons or muons, jets, and missing transverse momentum in proton-proton collisions at $\sqrt{s}=13$ TeV, $\it JHEP.~02~(2018)~067$, doi:10.1007/JHEP02(2018)067, arXiv:1710.09154 [hep-ex]
- A.M. Sirunyan et al., **Observation of the Higgs boson decay to a pair of** #tau leptons with the CMS detector, *Phys. Lett.* B779 (2018) 283–316, doi:10.1016/j.physletb.2018.02.004, arXiv:1708.00373 [hep-ex]
- A.M. Sirunyan et al., Search for the pair production of third-generation squarks with two-body decays to a bottom or charm quark and a neutralino in proton–proton collisions at $\sqrt{s} = 13$ TeV, *Phys. Lett.* B778 (2018) 263–291, doi:10.1016/j.physletb.2018.01.012, arXiv:1707.07274 [hep-ex]
- A.M. Sirunyan et al., Search for top squarks and dark matter particles in opposite-charge dilepton final states at $\sqrt{s} = 13$ TeV, *Phys. Rev.* D97 (2018) 032009, doi:10.1103/PhysRevD.97.032009, arXiv:1711.00752 [hep-ex]
- A.M. Sirunyan et al., Measurement of the associated production of a single top quark and a Z boson in pp collisions at $\sqrt{s} = \text{TeV}$, Phys. Lett. B779 (2018) 358–384, doi:10.1016/j.physletb.2018.02.025, arXiv:1712.02825 [hep-ex]
- A.M. Sirunyan et al., Search for low mass vector resonances decaying

- into quark-antiquark pairs in proton-proton collisions at $\sqrt{s} = 13$ TeV, JHEP.~01~(2018)~097,~doi:10.1007/JHEP01(2018)097,~arXiv:1710.00159~[hep-ex]
- A.M. Sirunyan et al., Search for Higgsino pair production in pp collisions at $\sqrt{s} = 13$ TeV in final states with large missing transverse momentum and two Higgs bosons decaying via $H \to b\bar{b}$, Phys. Rev. D97 (2018) 032007, doi:10.1103/PhysRevD.97.032007, arXiv:1709.04896 [hep-ex]
- A.M. Sirunyan et al., Azimuthal anisotropy of charged particles with transverse momentum up to 100 GeV/ c in PbPb collisions at $\sqrt{s_{NN}}$ =5.02 TeV, Phys. Lett. B776 (2018) 195–216, doi:10.1016/j.physletb.2017.11.041, arXiv:1702.00630 [hep-ex]
- A.M. Sirunyan et al., Search for supersymmetry in proton-proton collisions at 13 TeV using identified top quarks, *Phys. Rev.* D97 (2018) 012007, doi:10.1103/PhysRevD.97.012007, arXiv:1710.11188 [hep-ex]
- A.M. Sirunyan et al., Search for pair production of vector-like quarks in the bW \overline{b} W channel from proton-proton collisions at $\sqrt{s}=13$ TeV, *Phys. Lett.* B779 (2018) 82–106, doi:10.1016/j.physletb.2018.01.077, arXiv:1710.01539 [hep-ex]
- A.M. Sirunyan et al., Observation of electroweak production of samesign W boson pairs in the two jet and two same-sign lepton final state in proton-proton collisions at $\sqrt{s} = 13$ TeV, *Phys. Rev. Lett.* 120 (2018) 081801, doi:10.1103/PhysRevLett.120.081801, arXiv:1709.05822 [hep-ex]
- A.M. Sirunyan et al., Measurements of $t\bar{t}$ cross sections in association with b jets and inclusive jets and their ratio using dilepton final states in pp collisions at $\sqrt{s}=13$ TeV, Phys. Lett. B776 (2018) 355–378, doi:10.1016/j.physletb.2017.11.043, arXiv:1705.10141 [hep-ex]
- A.M. Sirunyan et al., Search for heavy resonances decaying to a top quark and a bottom quark in the lepton+jets final state in proton-proton collisions at 13 TeV, *Phys. Lett.* B777 (2018) 39–63, doi:10.1016/j.physletb.2017.12.006, arXiv:1708.08539 [hep-ex]
- A.M. Sirunyan et al., Pseudorapidity distributions of charged hadrons in proton-lead collisions at $\sqrt{s_{_{\rm NN}}} = 5.02$ and 8.16 TeV, *JHEP*. 01 (2018) 045, doi:10.1007/JHEP01(2018)045, arXiv:1710.09355 [hep-ex]
- A.M. Sirunyan et al., Search for pair production of excited top quarks in the lepton + jets final state, *Phys. Lett.* B778 (2018) 349, doi:10.1016/j.physletb.2018.01.049, arXiv:1711.10949 [hep-ex]
- A.M. Sirunyan et al., Measurement of the Λ_b polarization and angular parameters in $\Lambda_b \to J/\psi \Lambda$ decays from pp collisions at $\sqrt{s}=7$ and 8 TeV, (2018), arXiv:1802.04867 [hep-ex]
- A.M. Sirunyan et al., Search for single production of vector-like quarks

- decaying to a b quark and a Higgs boson, (2018), arXiv:1802.01486 [hep-ex]
- A.M. Sirunyan et al., Search for a heavy resonance decaying to a pair of vector bosons in the lepton plus merged jet final state at $\sqrt{s} = 13$ TeV, (2018), arXiv:1802.09407 [hep-ex]
- A.M. Sirunyan et al., Search for new physics in events with two soft oppositely charged leptons and missing transverse momentum in proton-proton collisions at $\sqrt{s} = 13$ TeV, (2018), arXiv:1801.01846 [hep-ex]
- A.M. Sirunyan et al., Comparing transverse momentum balance of b jet pairs in pp and PbPb collisions at $\sqrt{s_{\rm NN}}=$ 5.02 TeV, (2018), arXiv:1802.00707 [hep-ex]
- A.M. Sirunyan et al., Search for heavy neutral leptons in events with three charged leptons in proton-proton collisions at $\sqrt{s} = 13$ TeV, (2018), arXiv:1802.02965 [hep-ex]
- A.M. Sirunyan et al., Measurement of the inelastic proton-proton cross section at $\sqrt{s} = 13$ TeV, (2018), arXiv:1802.02613 [hep-ex]
- A.M. Sirunyan et al., Search for narrow resonances in the b-tagged dijet mass spectrum in proton-proton collisions at $\sqrt{s} = 8$ TeV, (2018), arXiv:1802.06149 [hep-ex]
- A.M. Sirunyan et al., **Observation of medium induced modifications of jet fragmentation in PbPb collisions using isolated-photon-tagged jets**, (2018), arXiv:1801.04895 [hep-ex]
- A.M. Sirunyan et al., Measurement of the $Z\gamma^* \to \tau\tau$ cross section in pp collisions at $\sqrt{s}=$ 13 TeV and validation of τ lepton analysis techniques, (2018), arXiv:1801.03535 [hep-ex]
- A.M. Sirunyan et al., Combined search for electroweak production of charginos and neutralinos in proton-proton collisions at $\sqrt{s} = 13$ TeV, (2018), arXiv:1801.03957 [hep-ex]
- A.M. Sirunyan et al., Search for natural and split supersymmetry in proton-proton collisions at $\sqrt{s} = 13$ TeV in final states with jets and missing transverse momentum, (2018), arXiv:1802.02110 [hep-ex]
- A.M. Sirunyan et al., Measurements of the pp \to ZZ production cross section and the Z \to 4 ℓ branching fraction, and constraints on anomalous triple gauge couplings at $\sqrt{s}=13 {\rm TeV},~Eur.~Phys.~J.~C78~(2018)~165,$ doi:10.1140/epjc/s10052-018-5567-9
- A.M. Sirunyan et al., Search for lepton-flavor violating decays of heavy resonances and quantum black holes to e μ final states in proton-

- proton collisions at $\sqrt{s} = 13$ TeV, (2018), arXiv:1802.01122 [hep-ex]
- S. Chatrchyan et al., Measurement of the mass difference between top quark and antiquark in pp collisions at $\sqrt{s} = 8$ TeV, *Phys. Lett.* B770 (2017) 50–71, doi:10.1016/j.physletb.2017.04.028, arXiv:1610.09551 [hep-ex]
- V. Khachatryan et al., Search for top quark decays via Higgs-boson-mediated flavor-changing neutral currents in pp collisions at $\sqrt{s}=8$ TeV, *JHEP*. 02 (2017) 079, doi:10.1007/JHEP02(2017)079, arXiv:1610.04857 [hep-ex]
- V. Khachatryan et al., **The CMS trigger system**, *JINST*. 12 (2017) P01020, doi:10.1088/1748-0221/12/01/P01020, arXiv:1609.02366 [physics.ins-det]
- V. Khachatryan et al., Search for heavy resonances decaying into a vector boson and a Higgs boson in final states with charged leptons, neutrinos, and b quarks, *Phys. Lett.* B768 (2017) 137–162, doi:10.1016/j.physletb.2017.02.040, arXiv:1610.08066 [hep-ex]
- V. Khachatryan et al., Measurements of differential production cross sections for a **Z** boson in association with jets in pp collisions at $\sqrt{s} = 8$ **TeV**, *JHEP*. 04 (2017) 022, doi:10.1007/JHEP04(2017)022, arXiv:1611.03844 [hep-ex]
- V. Khachatryan et al., Measurement of the total and differential inclusive B^+ hadron cross sections in pp collisions at $\sqrt{s}=$ 13 TeV, Phys. Lett. B771 (2017) 435–456, doi:10.1016/j.physletb.2017.05.074, arXiv:1609.00873 [hep-ex]
- V. Khachatryan et al., A search for new phenomena in pp collisions at $\sqrt{s}=13 {\rm TeV}$ in final states with missing transverse momentum and at least one jet using the $\alpha_{\rm T}$ variable, Eur. Phys. J. C77 (2017) 294, doi:10.1140/epjc/s10052-017-4787-8, arXiv:1611.00338 [hep-ex]
- V. Khachatryan et al., Inclusive search for supersymmetry using razor variables in pp collisions at $\sqrt{s} = 13$ TeV, Phys. Rev. D95 (2017) 012003, doi:10.1103/PhysRevD.95.012003, arXiv:1609.07658 [hep-ex]
- V. Khachatryan et al., Measurements of differential cross sections for associated production of a W boson and jets in proton-proton collisions at $\sqrt{s}=8$ TeV, Phys. Rev. D95 (2017) 052002, doi:10.1103/PhysRevD.95.052002, arXiv:1610.04222 [hep-ex]
- V. Khachatryan et al., Observation of charge-dependent azimuthal correlations in p-Pb collisions and its implication for the search for the chiral magnetic effect, Phys. Rev. Lett. 118 (2017) 122301, doi:10.1103/PhysRevLett.118.122301, arXiv:1610.00263 [nucl-ex]
- V. Khachatryan et al., Pseudorapidity dependence of long-range two-particle correlations in pPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, Phys. Rev.

- C96 (2017) 014915, doi:10.1103/PhysRevC.96.014915, arXiv:1604.05347 [nuclex]
- V. Khachatryan et al., Measurements of the associated production of a **Z** boson and b jets in pp collisions at $\sqrt{s} = 8$ TeV, Eur. Phys. J. C77 (2017) 751, doi:10.1140/epjc/s10052-017-5140-y, arXiv:1611.06507 [hep-ex]
- V. Khachatryan et al., Measurement of the production cross section of a W boson in association with two b jets in pp collisions at $\sqrt{s}=8{\rm TeV},\ Eur.\ Phys.\ J.\ C77\ (2017)\ 92,\ doi:10.1140/epjc/s10052-016-4573-z,\ arXiv:1608.07561\ [hep-ex]$
- V. Khachatryan et al., Measurement of inclusive jet cross sections in pp and PbPb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, Phys.~Rev. C96 (2017) 015202, doi:10.1103/PhysRevC.96.015202, arXiv:1609.05383 [nucl-ex]
- V. Khachatryan et al., Search for heavy neutrinos or third-generation leptoquarks in final states with two hadronically decaying τ leptons and two jets in proton-proton collisions at $\sqrt{s} = 13$ TeV, *JHEP*. 03 (2017) 077, doi:10.1007/JHEP03(2017)077, arXiv:1612.01190 [hep-ex]
- V. Khachatryan et al., Search for heavy gauge W' boson in events with an energetic lepton and large missing transverse momentum at $\sqrt{s} =$ 13 TeV, *Phys. Lett.* B770 (2017) 278–301, doi:10.1016/j.physletb.2017.04.043, arXiv:1612.09274 [hep-ex]
- V. Khachatryan et al., **Jet energy scale and resolution in the CMS experiment in pp collisions at 8 TeV**, *JINST*. 12 (2017) P02014, doi:10.1088/1748-0221/12/02/P02014, arXiv:1607.03663 [hep-ex]
- V. Khachatryan et al., Measurement of the $t\bar{t}$ production cross section using events in the e μ final state in pp collisions at $\sqrt{s}=$ 13 TeV, Eur. Phys. J. C77 (2017) 172, doi:10.1140/epjc/s10052-017-4718-8, arXiv:1611.04040 [hep-ex]
- V. Khachatryan et al., Search for Dark Matter and Supersymmetry with a Compressed Mass Spectrum in the Vector Boson Fusion Topology in Proton-Proton Collisions at $\sqrt{s} = 8$ TeV, Phys. Rev. Lett. 118 (2017) 021802, doi:10.1103/PhysRevLett.118.021802, arXiv:1605.09305 [hep-ex]
- V. Khachatryan et al., Measurement and QCD analysis of double-differential inclusive jet cross sections in pp collisions at $\sqrt{s}=8$ TeV and cross section ratios to 2.76 and 7 TeV, $\it{JHEP}.03$ (2017) 156, doi:10.1007/JHEP03(2017)156, arXiv:1609.05331 [hep-ex]
- V. Khachatryan et al., Measurement of differential cross sections for top quark pair production using the lepton+jets final state in proton-proton collisions at 13 TeV, *Phys. Rev.* D95 (2017) 092001, doi:10.1103/PhysRevD.95.092001, arXiv:1610.04191 [hep-ex]
- V. Khachatryan et al., Measurement of the transverse momentum spec-

- tra of weak vector bosons produced in proton-proton collisions at $\sqrt{s}=8$ TeV, JHEP.~02~(2017)~096,~doi:10.1007/JHEP02(2017)096,~arXiv:1606.05864 [hep-ex]
- V. Khachatryan et al., Search for CP violation in $t\bar{t}$ production and decay in proton-proton collisions at $\sqrt{s}=8$ TeV, *JHEP*. 03 (2017) 101, doi:10.1007/JHEP03(2017)101, arXiv:1611.08931 [hep-ex]
- V. Khachatryan et al., **Observation of the decay** $B^+ \to \psi(2S)\phi(1020)K^+$ in pp collisions at $\sqrt{s}=8$ TeV, Phys. Lett. B764 (2017) 66–86, doi:10.1016/j.physletb.2016.11.001, arXiv:1607.02638 [hep-ex]
- V. Khachatryan et al., Search for high-mass **Z** γ resonances in e⁺e⁻ γ and $\mu^+\mu^-\gamma$ final states in proton-proton collisions at $\sqrt{s}=8$ and 13 TeV, *JHEP*. 01 (2017) 076, doi:10.1007/JHEP01(2017)076, arXiv:1610.02960 [hep-ex]
- V. Khachatryan et al., Charged-particle nuclear modification factors in PbPb and pPb collisions at $\sqrt{s_{\rm NN}}=5.02$ TeV, *JHEP*. 04 (2017) 039, doi:10.1007/JHEP04(2017)039, arXiv:1611.01664 [nucl-ex]
- V. Khachatryan et al., **Search for supersymmetry in events with photons and missing transverse energy in pp collisions at 13 TeV**, *Phys. Lett.* B769 (2017) 391–412, doi:10.1016/j.physletb.2017.04.005, arXiv:1611.06604 [hep-ex]
- V. Khachatryan et al., Measurement of the WZ production cross section in pp collisions at $\sqrt{s}=7$ and 8 TeV and search for anomalous triple gauge couplings at $\sqrt{s}=8$ TeV, Eur. Phys. J. C77 (2017) 236, doi:10.1140/epjc/s10052-017-4730-z, arXiv:1609.05721 [hep-ex]
- V. Khachatryan et al., Search for top squark pair production in compressed-mass-spectrum scenarios in proton-proton collisions at $\sqrt{s}=8$ TeV using the α_T variable, *Phys. Lett.* B767 (2017) 403–430, doi:10.1016/j.physletb.2017.02.007, arXiv:1605.08993 [hep-ex]
- V. Khachatryan et al., Coherent J/ψ photoproduction in ultraperipheral PbPb collisions at $\sqrt{s_{NN}}=$ 2.76 TeV with the CMS experiment, *Phys. Lett.* B772 (2017) 489–511, doi:10.1016/j.physletb.2017.07.001, arXiv:1605.06966 [nucl-ex]
- V. Khachatryan et al., Search for heavy resonances decaying to tau lepton pairs in proton-proton collisions at $\sqrt{s}=13$ TeV, JHEP.~02~(2017)~048,~doi:10.1007/JHEP02(2017)048,~arXiv:1611.06594~[hep-ex]
- V. Khachatryan et al., Search for anomalous Wtb couplings and flavour-changing neutral currents in t-channel single top quark production in pp collisions at $\sqrt{s} = 7$ and 8 TeV, *JHEP*. 02 (2017) 028, doi:10.1007/JHEP02(2017)028, arXiv:1610.03545 [hep-ex]
- V. Khachatryan et al., Measurement of the WZ production cross section in pp collisions at $\sqrt(s) = 13$ TeV, Phys. Lett. B766 (2017) 268–290,

- doi:10.1016/j.physletb.2017.01.011, arXiv:1607.06943 [hep-ex]
- V. Khachatryan et al., Search for electroweak production of charginos in final states with two leptons in pp collisions at $\sqrt{s} = 8$ TeV, JHEP. 04 (2017) 018, doi:10.1007/JHEP04(2017)018, arXiv:1610.04870 [hep-ex]
- V. Khachatryan et al., Evidence for collectivity in pp collisions at the LHC, *Phys. Lett.* B765 (2017) 193–220, doi:10.1016/j.physletb.2016.12.009, arXiv:1606.06198 [nucl-ex]
- V. Khachatryan et al., Search for R-parity violating supersymmetry with displaced vertices in proton-proton collisions at $\sqrt{s}=8$ TeV, Phys. Rev. D95 (2017) 012009, doi:10.1103/PhysRevD.95.012009, arXiv:1610.05133 [hep-ex]
- V. Khachatryan et al., Measurement of electroweak-induced production of W γ with two jets in pp collisions at $\sqrt{s}=8$ TeV and constraints on anomalous quartic gauge couplings, *JHEP*. 06 (2017) 106, doi:10.1007/JHEP06(2017)106, arXiv:1612.09256 [hep-ex]
- V. Khachatryan et al., Measurement of the transverse momentum spectrum of the Higgs boson produced in pp collisions at $\sqrt{s}=8$ TeV using $H\to WW$ decays, JHEP. 03 (2017) 032, doi:10.1007/JHEP03(2017)032, arXiv:1606.01522 [hep-ex]
- V. Khachatryan et al., Search for single production of a heavy vector-like T quark decaying to a Higgs boson and a top quark with a lepton and jets in the final state, *Phys. Lett.* B771 (2017) 80–105, doi:10.1016/j.physletb.2017.05.019, arXiv:1612.00999 [hep-ex]
- V. Khachatryan et al., Searches for invisible decays of the Higgs boson in pp collisions at $\sqrt{s}=7$, 8, and 13 TeV, *JHEP*. 02 (2017) 135, doi:10.1007/JHEP02(2017)135, arXiv:1610.09218 [hep-ex]
- V. Khachatryan et al., Search for supersymmetry in events with one lepton and multiple jets in proton-proton collisions at $\sqrt{s} = 13$ TeV, Phys. Rev. D95 (2017) 012011, doi:10.1103/PhysRevD.95.012011, arXiv:1609.09386 [hep-ex]
- V. Khachatryan et al., Search for new phenomena in events with high jet multiplicity and low missing transverse momentum in proton–proton collisions at $\sqrt{s}=8$ TeV, *Phys. Lett.* B770 (2017) 257–267, doi:10.1016/j.physletb.2017.01.073, arXiv:1608.01224 [hep-ex]
- V. Khachatryan et al., Suppression of $\Upsilon(1S)$, $\Upsilon(2S)$ and $\Upsilon(3S)$ production in PbPb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, Phys. Lett. B770 (2017) 357–379, doi:10.1016/j.physletb.2017.04.031, arXiv:1611.01510 [nucl-ex]
- V. Khachatryan et al., Observation of $\Upsilon(1S)$ pair production in proton-proton collisions at $\sqrt{s} = 8$ TeV, *JHEP*. 05 (2017) 013,

- doi:10.1007/JHEP05(2017)013, arXiv:1610.07095 [hep-ex]
- V. Khachatryan et al., Search for high-mass diphoton resonances in proton–proton collisions at 13 TeV and combination with 8 TeV search, *Phys. Lett.* B767 (2017) 147–170, doi:10.1016/j.physletb.2017.01.027, arXiv:1609.02507 [hep-ex]
- V. Khachatryan et al., Suppression and azimuthal anisotropy of prompt and nonprompt J/ ψ production in PbPb collisions at $\sqrt{s_{_{\mathrm{NN}}}} = 2.76\mathrm{TeV}$, Eur. Phys. J. C77 (2017) 252, doi:10.1140/epjc/s10052-017-4781-1, arXiv:1610.00613 [nucl-ex]
- V. Khachatryan et al., **Multiplicity and rapidity dependence of strange hadron production in pp, pPb, and PbPb collisions at the LHC**, *Phys. Lett.* B768 (2017) 103–129, doi:10.1016/j.physletb.2017.01.075, arXiv:1605.06699 [nucl-ex]
- V. Khachatryan et al., Measurements of the $t\bar{t}$ production cross section in lepton+jets final states in pp collisions at 8 TeV and ratio of 8 to 7 TeV cross sections, Eur. Phys. J. C77 (2017) 15, doi:10.1140/epjc/s10052-016-4504-z, arXiv:1602.09024 [hep-ex]
- V. Khachatryan et al., Search for narrow resonances in dilepton mass spectra in proton-proton collisions at $\sqrt{s}=13$ TeV and combination with 8 TeV data, *Phys. Lett.* B768 (2017) 57–80, doi:10.1016/j.physletb.2017.02.010, arXiv:1609.05391 [hep-ex]
- V. Khachatryan et al., Measurement of the cross section for electroweak production of Z γ in association with two jets and constraints on anomalous quartic gauge couplings in proton–proton collisions at $\sqrt{s}=8$ TeV, *Phys. Lett.* B770 (2017) 380–402, doi:10.1016/j.physletb.2017.04.071, arXiv:1702.03025 [hep-ex]
- V. Khachatryan et al., Search for light bosons in decays of the 125 GeV Higgs boson in proton-proton collisions at $\sqrt{s} = 8$ TeV, JHEP. 10 (2017) 076, doi:10.1007/JHEP10(2017)076, arXiv:1701.02032 [hep-ex]
- V. Khachatryan et al., Search for new phenomena with multiple charged leptons in proton–proton collisions at $\sqrt{s} = 13$ TeV, Eur. Phys. J. C77 (2017) 635, doi:10.1140/epjc/s10052-017-5182-1, arXiv:1701.06940 [hep-ex]
- V. Khachatryan et al., Search for supersymmetry in the all-hadronic final state using top quark tagging in pp collisions at $\sqrt{s}=13$ TeV, Phys. Rev. D96 (2017) 012004, doi:10.1103/PhysRevD.96.012004, arXiv:1701.01954 [hep-ex]
- V. Khachatryan et al., Search for leptophobic Z bosons decaying into four-lepton final states in proton-proton collisions at \sqrt{s} =8TeV, *Phys. Lett.* B773 (2017) 563–584, doi:10.1016/j.physletb.2017.08.069,

- arXiv:1701.01345 [hep-ex]
- V. Khachatryan et al., Exclusive and semi-exclusive pi+pi- production in proton-proton collisions at sqrt(s) = 7 TeV, (2017), arXiv:1706.08310 [hep-ex]
- A.M. Sirunyan et al., Search for massive resonances decaying into WW, WZ or ZZ bosons in proton-proton collisions at $\sqrt{s} = 13$ TeV, JHEP. 03 (2017) 162, doi:10.1007/JHEP03(2017)162, arXiv:1612.09159 [hep-ex]
- A.M. Sirunyan et al., Cross section measurement of t-channel single top quark production in pp collisions at $\sqrt{s} = 13$ TeV, Phys. Lett. B772 (2017) 752–776, doi:10.1016/j.physletb.2017.07.047, arXiv:1610.00678 [hep-ex]
- A.M. Sirunyan et al., Measurements of the charm jet cross section and nuclear modification factor in pPb collisions at $\sqrt{s_{NN}}=5.02$ TeV, Phys. Lett. B772 (2017) 306–329, doi:10.1016/j.physletb.2017.06.053, arXiv:1612.08972 [nucl-ex]
- A.M. Sirunyan et al., Search for dijet resonances in proton–proton collisions at $\sqrt{s}=13$ TeV and constraints on dark matter and other models, *Phys. Lett.* B769 (2017) 520–542, doi:10.1016/j.physletb.2017.09.029, arXiv:1611.03568 [hep-ex]
- A.M. Sirunyan et al., Search for electroweak production of a vector-like quark decaying to a top quark and a Higgs boson using boosted topologies in fully hadronic final states, *JHEP*. 04 (2017) 136, doi:10.1007/JHEP04(2017)136, arXiv:1612.05336 [hep-ex]
- A.M. Sirunyan et al., Search for high-mass $Z\gamma$ resonances in proton-proton collisions at $\sqrt{s}=8$ and 13 TeV using jet substructure techniques, *Phys. Lett.* B772 (2017) 363–387, doi:10.1016/j.physletb.2017.06.062, arXiv:1612.09516 [hep-ex]
- A.M. Sirunyan et al., Relative Modification of Prompt (2S) and J/Yields from pp to PbPb Collisions at $\sqrt{s_{NN}} = 5.02 \,\text{TeV}$, Phys. Rev. Lett. 118 (2017) 162301, doi:10.1103/PhysRevLett.118.162301, arXiv:1611.01438 [nucl-ex]
- A.M. Sirunyan et al., Search for massive resonances decaying into WW, WZ, ZZ, qW, and qZ with dijet final states at sqrt(s) = 13 TeV, (2017), arXiv:1708.05379 [hep-ex]
- A.M. Sirunyan et al., Search for black holes in high-multiplicity final states in proton-proton collisions at $\sqrt{s} = 13$ TeV, *Phys. Lett.* B774 (2017) 279–307, doi:10.1016/j.physletb.2017.09.053, arXiv:1705.01403 [hep-ex]
- A.M. Sirunyan et al., Measurement of double-differential cross sections for top quark pair production in pp collisions at $\sqrt{s} = 8\text{TeV}$ and impact on parton distribution functions, Eur. Phys. J. C77 (2017) 459,

- doi:10.1140/epjc/s10052-017-4984-5, arXiv:1703.01630 [hep-ex]
- A.M. Sirunyan et al., Search for anomalous couplings in boosted WW/WZ $\rightarrow \ell\nu\bar{q}q$ production in proton-proton collisions at $\sqrt{s}=8$ TeV, Phys. Lett. B772 (2017) 21–42, doi:10.1016/j.physletb.2017.06.009, arXiv:1703.06095 [hep-ex]
- A.M. Sirunyan et al., Search for a heavy resonance decaying to a top quark and a vector-like top quark at $\sqrt{s} = 13$ TeV, *JHEP*. 09 (2017) 053, doi:10.1007/JHEP09(2017)053, arXiv:1703.06352 [hep-ex]
- A.M. Sirunyan et al., Measurement of the splitting function in pp and PbPb collisions at $\sqrt{s_{_{\rm NN}}} = 5.02$ TeV, (2017), arXiv:1708.09429 [nucl-ex]
- A.M. Sirunyan et al., Search for physics beyond the standard model in events with high-momentum Higgs bosons and missing transverse momentum in proton-proton collisions at 13 TeV, Submitted to: Phys. Rev. Lett. (2017), arXiv:1712.08501 [hep-ex]
- A.M. Sirunyan et al., Search for supersymmetry in multijet events with missing transverse momentum in proton-proton collisions at 13 TeV, *Phys. Rev.* D96 (2017) 032003, doi:10.1103/PhysRevD.96.032003, arXiv:1704.07781 [hep-ex]
- A.M. Sirunyan et al., Search for natural supersymmetry in events with top quark pairs and photons in pp collisions at sqrt(s) = 8 TeV, (2017), arXiv:1707.03325 [hep-ex]
- A.M. Sirunyan et al., Search for R-parity violating supersymmetry in pp collisions at $\sqrt{s} = 13$ TeV using b jets in a final state with a single lepton, many jets, and high sum of large-radius jet masses, Submitted to: Phys. Lett. B. (2017), arXiv:1712.08920 [hep-ex]
- A.M. Sirunyan et al., Measurement of angular parameters from the decay $B^0 \to K^{*0} \mu^+ \mu^-$ in proton-proton collisions at $\sqrt{s} = 8$ TeV, (2017), arXiv:1710.02846 [hep-ex]
- A.M. Sirunyan et al., Search for Low Mass Vector Resonances Decaying to Quark-Antiquark Pairs in Proton-Proton Collisions at $\sqrt{s} = 13$ TeV, Phys. Rev. Lett. 119 (2017) 111802, doi:10.1103/PhysRevLett.119.111802, arXiv:1705.10532 [hep-ex]
- A.M. Sirunyan et al., Evidence for the Higgs boson decay to a bottom quark-antiquark pair, (2017), arXiv:1709.07497 [hep-ex]
- A.M. Sirunyan et al., Search for new physics in the monophoton final state in proton-proton collisions at $\sqrt{s} = 13$ TeV, *JHEP*. 10 (2017) 073, doi:10.1007/JHEP10(2017)073, arXiv:1706.03794 [hep-ex]
- A.M. Sirunyan et al., Measurements of properties of the Higgs boson decaying into the four-lepton final state in pp collisions at $\sqrt{s} = 13$

- A.M. Sirunyan et al., **Identification of heavy-flavour jets with the CMS detector in pp collisions at 13 TeV**, *Submitted to: JINST*. (2017), arXiv:1712.07158 [physics.ins-det]
- A.M. Sirunyan et al., Search for single production of vector-like quarks decaying to a Z boson and a top or a bottom quark in proton-proton collisions at $\sqrt{s}=13$ TeV, *JHEP*. 05 (2017) 029, doi:10.1007/JHEP05(2017)029, arXiv:1701.07409 [hep-ex]
- A.M. Sirunyan et al., Search for excited states of light and heavy flavor quarks in the γ +jet final state in proton-proton collisions at $\sqrt{s} = 13$ TeV, (2017), arXiv:1711.04652 [hep-ex]
- A.M. Sirunyan et al., Search for Supersymmetry in pp Collisions at $\sqrt{s} = 13 \text{TeV}$ in the Single-Lepton Final State Using the Sum of Masses of Large-Radius Jets, Phys. Rev. Lett. 119 (2017) 151802, doi:10.1103/PhysRevLett.119.151802, arXiv:1705.04673 [hep-ex]
- A.M. Sirunyan et al., Non-Gaussian elliptic-flow fluctuations in PbPb collisions at $\sqrt{s_{_{\mathrm{NN}}}} = 5.02$ TeV, (2017), arXiv:1711.05594 [nucl-ex]
- A.M. Sirunyan et al., Measurement of vector boson scattering and constraints on anomalous quartic couplings from events with four leptons and two jets in proton–proton collisions at $\sqrt{s}=$ 13 TeV, *Phys. Lett.* B774 (2017) 682–705, doi:10.1016/j.physletb.2017.10.020, arXiv:1708.02812 [hep-ex]
- A.M. Sirunyan et al., **Principal-component analysis of two-particle azimuthal correlations in PbPb and** pPb **collisions at CMS**, Phys. Rev. C96 (2017) 064902, doi:10.1103/PhysRevC.96.064902, arXiv:1708.07113 [nuclex]
- A.M. Sirunyan et al., Search for dark matter produced with an energetic jet or a hadronically decaying W or Z boson at $\sqrt{s} = 13$ TeV, *JHEP*. 07 (2017) 014, doi:10.1007/JHEP07(2017)014, arXiv:1703.01651 [hep-ex]
- A.M. Sirunyan et al., Search for associated production of dark matter with a Higgs boson decaying to $b\bar{b}$ or $\gamma\gamma$ at $\sqrt{s}=13$ TeV, *JHEP*. 10 (2017) 180, doi:10.1007/JHEP10(2017)180, arXiv:1703.05236 [hep-ex]
- A.M. Sirunyan et al., Search for Z γ resonances using leptonic and hadronic final states in proton-proton collisions at $\sqrt{s} = 13$ TeV, (2017), arXiv:1712.03143 [hep-ex]
- A.M. Sirunyan et al., Measurement of the top quark mass using single top quark events in proton-proton collisions at $\sqrt{s} = 8$ TeV, Eur. Phys.

- J. C77 (2017) 354, doi:10.1140/epjc/s10052-017-4912-8, arXiv:1703.02530 [hepex]
- A.M. Sirunyan et al., Measurement of the top quark mass in the dileptonic $t\bar{t}$ decay channel using the mass observables $M_{b\ell}$, M_{T2} , and $M_{b\ell\nu}$ in pp collisions at $\sqrt{s}=8$ TeV, Phys. Rev. D96 (2017) 032002, doi:10.1103/PhysRevD.96.032002, arXiv:1704.06142 [hep-ex]
- A.M. Sirunyan et al., Study of Bose-Einstein correlations in pp, pPb, and PbPb collisions at the LHC, Submitted to: Phys. Rev. C. (2017), arXiv:1712.07198 [hep-ex]
- A.M. Sirunyan et al., Pseudorapidity and transverse momentum dependence of flow harmonics in pPb and PbPb collisions, (2017), arXiv:1710.07864 [nucl-ex]
- A.M. Sirunyan et al., Measurement of differential cross sections in the ϕ^* variable for inclusive Z boson production in pp collisions at $\sqrt{s} = 8$ TeV, (2017), arXiv:1710.07955 [hep-ex]
- A.M. Sirunyan et al., Search for a massive resonance decaying to a pair of Higgs bosons in the four b quark final state in proton-proton collisions at $\sqrt{s} = 13$ TeV, (2017), arXiv:1710.04960 [hep-ex]
- A.M. Sirunyan et al., Measurement of prompt and nonprompt charmonium suppression in PbPb collisions at 5.02 TeV, Submitted to: Eur. Phys. J. C. (2017), arXiv:1712.08959 [nucl-ex]
- A.M. Sirunyan et al., Measurement of the semileptonic $t\bar{t}$ + production cross section in pp collisions at $\sqrt{s} = 8$ TeV, *JHEP*. 10 (2017) 006, doi:10.1007/JHEP10(2017)006, arXiv:1706.08128 [hep-ex]
- A.M. Sirunyan et al., Search for new long-lived particles at $\sqrt{s} = 13$ TeV, (2017), arXiv:1711.09120 [hep-ex]
- A.M. Sirunyan et al., Electroweak production of two jets in association with a **Z** boson in proton-proton collisions at $\sqrt{s} = 13$ TeV, Submitted to: Eur. Phys. J. C. (2017), arXiv:1712.09814 [hep-ex]
- A.M. Sirunyan et al., Study of Jet Quenching with Z+jet Correlations in Pb-Pb and pp Collisions at $\sqrt{s}_{NN}=5.02 {\rm TeV},~Phys.~Rev.~Lett.~119~(2017)~082301,~doi:10.1103/PhysRevLett.119.082301,~arXiv:1702.01060~[nucl-ex]$
- A.M. Sirunyan et al., Search for top quark partners with charge 5/3 in proton-proton collisions at $\sqrt{s}=13$ TeV, *JHEP*. 08 (2017) 073, doi:10.1007/JHEP08(2017)073, arXiv:1705.10967 [hep-ex]
- A.M. Sirunyan et al., Search for new physics in final states with an energetic jet or a hadronically decaying W or Z boson and transverse momentum imbalance at $\sqrt{s} = 13$ TeV, (2017), arXiv:1712.02345 [hep-ex]
- A.M. Sirunyan et al., Azimuthal correlations for inclusive 2-jet, 3-jet,

- and 4-jet events in pp collisions at $\sqrt{s} = 13$ TeV, (2017), arXiv:1712.05471 [hep-ex]
- A.M. Sirunyan et al., Search for diboson resonances in the 2 \$ \$2 ν final state, (2017), arXiv:1711.04370 [hep-ex]
- A.M. Sirunyan et al., Search for associated production of a Z boson with a single top quark and for tZ flavour-changing interactions in pp collisions at $\sqrt{s} = 8$ TeV, JHEP. 07 (2017) 003, doi:10.1007/JHEP07(2017)003, arXiv:1702.01404 [hep-ex]
- A.M. Sirunyan et al., Search for electroweak production of charginos and neutralinos in multilepton final states in proton-proton collisions at $\sqrt{s} = 13$ TeV, (2017), arXiv:1709.05406 [hep-ex]
- M. Aaboud et al., Combination of inclusive and differential $t\bar{t}$ charge asymmetry measurements using ATLAS and CMS data at $\sqrt{s}=7$ and 8 TeV, Submitted to: JHEP. (2017), arXiv:1709.05327 [hep-ex]
- A.M. Sirunyan et al., Measurements of the pp $\to W\gamma\gamma$ and pp $\to Z\gamma\gamma$ cross sections and limits on anomalous quartic gauge couplings at $\sqrt{s}=8$ TeV, *JHEP*. 10 (2017) 072, doi:10.1007/JHEP10(2017)072, arXiv:1704.00366 [hep-ex]
- A.M. Sirunyan et al., Suppression of excited Upsilon states relative to the ground state in PbPb collisions at sqrt(sNN) = 5.02 TeV, (2017), arXiv:1706.05984 [hep-ex]
- A.M. Sirunyan et al., Search for vector-like light-flavor quark partners in proton-proton collisions at sqrt(s)=8 TeV, (2017), arXiv:1708.02510 [hep-ex]
- A.M. Sirunyan et al., Search for supersymmetry in events with one lepton and multiple jets exploiting the angular correlation between the lepton and the missing transverse momentum in proton-proton collisions at $\sqrt{s} = 13$ TeV, (2017), arXiv:1709.09814 [hep-ex]
- A.M. Sirunyan et al., Measurement of prompt and nonprompt J/ ψ production in pp and pPb collisions at $\sqrt{s_{\rm NN}} = 5.02 {\rm TeV}$, Eur. Phys. J. C77 (2017) 269, doi:10.1140/epjc/s10052-017-4828-3, arXiv:1702.01462 [nucl-ex]
- A.M. Sirunyan et al., Measurement of normalized differential t-tbar cross sections in the dilepton channel from pp collisions at sqrt(s) = 13 TeV, (2017), arXiv:1708.07638 [hep-ex]
- A.M. Sirunyan et al., Measurement of b hadron lifetimes in pp collisions at $\sqrt{s} = 8$ TeV, (2017), arXiv:1710.08949 [hep-ex]
- A.M. Sirunyan et al., Combination of searches for heavy resonances decaying to WW, WZ, ZZ, WH, and ZH boson pairs in proton-proton collisions at $\sqrt{s}=8$ and 13 TeV, Phys. Lett. B774 (2017) 533–558,

- doi:10.1016/j.physletb.2017.09.083, arXiv:1705.09171 [hep-ex]
- A.M. Sirunyan et al., Measurement of the inclusive energy spectrum in the very forward direction in proton-proton collisions at $\sqrt{s} = 13$ TeV, *JHEP*. 08 (2017) 046, doi:10.1007/JHEP08(2017)046, arXiv:1701.08695 [hep-ex]
- A.M. Sirunyan et al., Search for gauge-mediated supersymmetry in events with at least one photon and missing transverse momentum in pp collisions at $\sqrt{s} = 13$ TeV, (2017), arXiv:1711.08008 [hep-ex]
- A.M. Sirunyan et al., Search for the X(5568) state decaying into $B_s^0 \pi^{\pm}$ in proton-proton collisions at $\sqrt{s} = 8$ TeV, (2017), arXiv:1712.06144 [hep-ex]
- A.M. Sirunyan et al., Search for dark matter and unparticles in events with a Z boson and missing transverse momentum in proton-proton collisions at $\sqrt{s}=13$ TeV, $\it{JHEP}.~03~(2017)~061$, doi:10.1007/JHEP01(2018)056, arXiv:1701.02042 [hep-ex]
- A.M. Sirunyan et al., Measurement of the B^{\pm} Meson Nuclear Modification Factor in Pb-Pb Collisions at $\sqrt{s_{NN}}=5.02 {\rm TeV}$, Phys. Rev. Lett. 119 (2017) 152301, doi:10.1103/PhysRevLett.119.152301, arXiv:1705.04727 [hep-ex]
- A.M. Sirunyan et al., Measurement of prompt D^0 meson azimuthal anisotropy in PbPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, (2017), arXiv:1708.03497 [nucl-ex]
- A.M. Sirunyan et al., Measurement of associated **Z** + charm production in proton-proton collisions at $\sqrt{s} = 8$ TeV, (2017), arXiv:1711.02143 [hep-ex]
- A.M. Sirunyan et al., Measurement of quarkonium production cross sections in pp collisions at $\sqrt{s} = 13$ TeV, (2017), arXiv:1710.11002 [hep-ex]
- A.M. Sirunyan et al., Search for new physics in events with a leptonically decaying Z boson and a large transverse momentum imbalance in proton-proton collisions at $\sqrt{s}=$ 13 TeV, (2017), arXiv:1711.00431 [hep-ex]
- A.M. Sirunyan et al., Study of jet quenching with isolated-photon+jet correlations in PbPb and pp collisions at $\sqrt{s_{_{\mathrm{NN}}}} = 5.02$ TeV, (2017), arXiv:1711.09738 [nucl-ex]
- A.M. Sirunyan et al., Search for Evidence of the Type-III Seesaw Mechanism in Multilepton Final States in Proton-Proton Collisions at $\sqrt{s}=13\mathrm{TeV},\ Phys.\ Rev.\ Lett.\ 119\ (2017)\ 221802,$ doi:10.1103/PhysRevLett.119.221802, arXiv:1708.07962 [hep-ex]
- A.M. Sirunyan et al., Constraints on the chiral magnetic effect using charge-dependent azimuthal correlations in pPb and PbPb collisions at the LHC, (2017), arXiv:1708.01602 [nucl-ex]
- A.M. Sirunyan et al., Mechanical stability of the CMS strip tracker

- measured with a laser alignment system, JINST. 12 (2017) P04023, doi:10.1088/1748-0221/12/04/P04023, arXiv:1701.02022 [physics.ins-det]
- A.M. Sirunyan et al., Study of dijet events with a large rapidity gap between the two leading jets in pp collisions at $\sqrt{s}=7$ TeV, (2017), arXiv:1710.02586 [hep-ex]
- A.M. Sirunyan et al., Search for Charged Higgs Bosons Produced via Vector Boson Fusion and Decaying into a Pair of W and Z Bosons Using pp Collisions at $\sqrt{s} = 13 \text{TeV}$, Phys. Rev. Lett. 119 (2017) 141802, doi:10.1103/PhysRevLett.119.141802, arXiv:1705.02942 [hep-ex]
- A.M. Sirunyan et al., Search for decays of stopped exotic long-lived particles produced in proton-proton collisions at $\sqrt{s} = 13$ TeV, Submitted to: JHEP. (2017), arXiv:1801.00359 [hep-ex]
- A.M. Sirunyan et al., Measurement of the triple-differential dijet cross section in proton-proton collisions at $\sqrt{s}=8 {\rm TeV}$ and constraints on parton distribution functions, Eur. Phys. J. C77 (2017) 746, doi:10.1140/epjc/s10052-017-5286-7, arXiv:1705.02628 [hep-ex]
- A.M. Sirunyan et al., Challenges to the chiral magnetic wave using charge-dependent azimuthal anisotropies in pPb and PbPb collisions at $\sqrt{s_{_{\mathrm{NN}}}} = 5.02$ TeV, (2017), arXiv:1708.08901 [nucl-ex]
- A.M. Sirunyan et al., Search for standard model production of four top quarks in proton-proton collisions at $\sqrt{s} = 13$ TeV, *Phys. Lett.* B772 (2017) 336–358, doi:10.1016/j.physletb.2017.06.064, arXiv:1702.06164 [hep-ex]
- A.M. Sirunyan et al., Search for single production of vector-like quarks decaying into a b quark and a W boson in proton-proton collisions at $\sqrt{s} = 13$ TeV, Phys. Lett. B772 (2017) 634–656, doi:10.1016/j.physletb.2017.07.022, arXiv:1701.08328 [hep-ex]
- A.M. Sirunyan et al., Constraints on anomalous Higgs boson couplings using production and decay information in the four-lepton final state, *Phys. Lett.* B775 (2017) 1–24, doi:10.1016/j.physletb.2017.10.021, arXiv:1707.00541 [hep-ex]
- A.M. Sirunyan et al., Search for Higgs boson pair production in the $bb\tau\tau$ final state in proton-proton collisions at $\sqrt(s) = 8\text{TeV}$, Phys. Rev. D96 (2017) 072004, doi:10.1103/PhysRevD.96.072004, arXiv:1707.00350 [hep-ex]
- A.M. Sirunyan et al., **Measurements of jet charge with dijet events in pp collisions at** $\sqrt{s}=8$ **TeV**, *JHEP*. 10 (2017) 131, doi:10.1007/JHEP10(2017)131, arXiv:1706.05868 [hep-ex]
- A.M. Sirunyan et al., Search for the flavor-changing neutral current interactions of the top quark and the Higgs boson which decays into a pair of b quarks at $\sqrt{s} = 13$ TeV, (2017), arXiv:1712.02399 [hep-ex]
- A.M. Sirunyan et al., Search for tt resonances in highly boosted lep-

- ton+jets and fully hadronic final states in proton-proton collisions at $\sqrt{s}=13$ TeV, JHEP.~07~(2017)~001,~doi:10.1007/JHEP07(2017)001,~arXiv:1704.03366~[hep-ex]
- A.M. Sirunyan et al., Measurement of the $t\bar{t}$ production cross section using events with one lepton and at least one jet in pp collisions at $\sqrt{s}=13$ TeV, JHEP.~09~(2017)~051,~doi:10.1007/JHEP09(2017)051,~arXiv:1701.06228~[hep-ex]
- A.M. Sirunyan et al., Searches for W bosons decaying to a top quark and a bottom quark in proton-proton collisions at 13 TeV, *JHEP*. 08 (2017) 029, doi:10.1007/JHEP08(2017)029, arXiv:1706.04260 [hep-ex]
- A.M. Sirunyan et al., Measurement of the inclusive $t\bar{t}$ cross section in pp collisions at $\sqrt{s} = 5.02$ TeV using final states with at least one charged lepton, (2017), arXiv:1711.03143 [hep-ex]
- A.M. Sirunyan et al., **Particle-flow reconstruction and global event description with the CMS detector**, *JINST*. 12 (2017) P10003, doi:10.1088/1748-0221/12/10/P10003, arXiv:1706.04965 [physics.ins-det]
- A.M. Sirunyan et al., Search for pair production of vector-like T and B quarks in single-lepton final states using boosted jet substructure in proton-proton collisions at $\sqrt{s}=13$ TeV, *JHEP*. 11 (2017) 085, doi:10.1007/JHEP11(2017)085, arXiv:1706.03408 [hep-ex]
- A.M. Sirunyan et al., Search for a light pseudoscalar Higgs boson produced in association with bottom quarks in pp collisions at $\sqrt{s}=8$ TeV, JHEP. 11 (2017) 010, doi:10.1007/JHEP11(2017)010, arXiv:1707.07283 [hep-ex]
- A.M. Sirunyan et al., Observation of correlated azimuthal anisotropy Fourier harmonics in pp and pPb collisions at the LHC, (2017), arXiv:1709.09189 [nucl-ex]
- A.M. Sirunyan et al., Measurement of the cross section for top quark pair production in association with a W or Z boson in proton-proton collisions at $\sqrt{s}=$ 13 TeV, (2017), arXiv:1711.02547 [hep-ex]
- A.M. Sirunyan et al., Measurement of the underlying event activity in inclusive Z boson production in proton-proton collisions at $\sqrt{s} = 13$ TeV, (2017), arXiv:1711.04299 [hep-ex]
- A.M. Sirunyan et al., Measurement of the differential cross sections for the associated production of a W boson and jets in proton-proton collisions at $\sqrt{s}=13$ TeV, Phys. Rev. D96 (2017) 072005, doi:10.1103/PhysRevD.96.072005, arXiv:1707.05979 [hep-ex]
- A.M. Sirunyan et al., Search for direct production of supersymmetric partners of the top quark in the all-jets final state in

- **proton-proton collisions at** $\sqrt{s} = 13$ **TeV**, *JHEP*. 10 (2017) 005, doi:10.1007/JHEP10(2017)005, arXiv:1707.03316 [hep-ex]
- A.M. Sirunyan et al., Search for heavy resonances that decay into a vector boson and a Higgs boson in hadronic final states at $\sqrt{s}=13 \text{TeV}$, Eur. Phys. J. C77 (2017) 636, doi:10.1140/epjc/s10052-017-5192-z, arXiv:1707.01303 [hep-ex]
- A.M. Sirunyan et al., Search for dark matter produced in association with heavy-flavor quark pairs in proton-proton collisions at $\sqrt{s} = 13$ TeV, Eur. Phys. J. C77 (2017) 845, doi:10.1140/epjc/s10052-017-5317-4, arXiv:1706.02581 [hep-ex]
- A.M. Sirunyan et al., **Observation of top quark production in proton-nucleus collisions**, *Phys. Rev. Lett.* 119 (2017) 242001, doi:10.1103/PhysRevLett.119.242001, arXiv:1709.07411 [nucl-ex]
- A.M. Sirunyan et al., Search for a heavy composite Majorana neutrino in the final state with two leptons and two quarks at $\sqrt{s}=13$ TeV, *Phys. Lett.* B775 (2017) 315–337, doi:10.1016/j.physletb.2017.11.001, arXiv:1706.08578 [hep-ex]
- A.M. Sirunyan et al., Nuclear modification factor of D0 mesons in PbPb collisions at sqrt(s[NN]) = 5.02 TeV, (2017), arXiv:1708.04962 [nucl-ex]
- A.M. Sirunyan et al., Search for lepton flavour violating decays of the Higgs boson to $\mu\tau$ and e τ in proton-proton collisions at $\sqrt{s} = 13$ TeV, Submitted to: JHEP. (2017), arXiv:1712.07173 [hep-ex]
- A.M. Sirunyan et al., Measurement of the jet mass in highly boosted $t\bar{t}$ events from pp collisions at $\sqrt{s}=8 \text{TeV}$, Eur. Phys. J. C77 (2017) 467, doi:10.1140/epjc/s10052-017-5030-3, arXiv:1703.06330 [hep-ex]
- A.M. Sirunyan et al., Search for new physics with dijet angular distributions in proton-proton collisions at $\sqrt{s} = 13$ TeV, *JHEP*. 07 (2017) 013, doi:10.1007/JHEP07(2017)013, arXiv:1703.09986 [hep-ex]
- A.M. Sirunyan et al., Search for single production of a vector-like T quark decaying to a Z boson and a top quark in proton-proton collisions at sqrt(s) = 13 TeV, (2017), arXiv:1708.01062 [hep-ex]
- A.M. Sirunyan et al., Search for third-generation scalar leptoquarks and heavy right-handed neutrinos in final states with two tau leptons and two jets in proton-proton collisions at $\sqrt{s}=13$ TeV, JHEP.~07~(2017)~121,~doi:10.1007/JHEP07(2017)121,~arXiv:1703.03995~[hep-ex]
- A.M. Sirunyan et al., Search for supersymmetry in events with at least one photon, missing transverse momentum, and large transverse event activity in proton-proton collisions at $\sqrt{s} = 13$ TeV, *JHEP*. 12 (2017) 142, doi:10.1007/JHEP12(2017)142, arXiv:1707.06193 [hep-ex]
- A.M. Sirunyan et al., Measurements of the pp \rightarrow ZZ production cross sec-

- tion and the $Z \to 4\ell$ branching fraction, and constraints on anomalous triple gauge couplings at $\sqrt{s} = 13$ TeV, (2017), arXiv:1709.08601 [hep-ex]
- A.M. Sirunyan et al., Measurement of charged pion, kaon, and proton production in proton-proton collisions at $\sqrt{s} = 13$ TeV, *Phys. Rev.* D96 (2017) 112003, doi:10.1103/PhysRevD.96.112003, arXiv:1706.10194 [hep-ex]
- V. Khachatryan et al., Search for supersymmetry in events with a photon, a lepton, and missing transverse momentum in pp collisions at $\sqrt{s} = 8$ TeV, *Phys. Lett.* B757 (2016) 6–31, doi:10.1016/j.physletb.2016.03.039, arXiv:1508.01218 [hep-ex]
- V. Khachatryan et al., Search for neutral MSSM Higgs bosons decaying to $\mu^+\mu^-$ in pp collisions at $\sqrt{s}=7$ and 8 TeV, *Phys. Lett.* B752 (2016) 221–246, doi:10.1016/j.physletb.2015.11.042, arXiv:1508.01437 [hep-ex]
- V. Khachatryan et al., Search for new phenomena in monophoton final states in proton-proton collisions at $\sqrt{s} = 8$ TeV, *Phys. Lett.* B755 (2016) 102–124, doi:10.1016/j.physletb.2016.01.057, arXiv:1410.8812 [hep-ex]
- V. Khachatryan et al., Search for anomalous single top quark production in association with a photon in pp collisions at $\sqrt{s} = 8$ TeV, *JHEP*. 04 (2016) 035, doi:10.1007/JHEP04(2016)035, arXiv:1511.03951 [hep-ex]
- V. Khachatryan et al., Search for a low-mass pseudoscalar Higgs boson produced in association with a $b\bar{b}$ pair in pp collisions at $\sqrt{s}=8$ TeV, Phys. Lett. B758 (2016) 296–320, doi:10.1016/j.physletb.2016.05.003, arXiv:1511.03610 [hep-ex]
- V. Khachatryan et al., Search for dark matter and unparticles produced in association with a Z boson in proton-proton collisions at $\sqrt{s} =$ 8 TeV, *Phys. Rev.* D93 (2016) 052011, doi:10.1103/PhysRevD.93.052011, arXiv:1511.09375 [hep-ex]
- V. Khachatryan et al., Search for a massive resonance decaying into a Higgs boson and a W or Z boson in hadronic final states in proton-proton collisions at $\sqrt{s}=8$ TeV, $\it JHEP.~02~(2016)~145$, $\it doi:10.1007/JHEP02(2016)145$, $\it arXiv:1506.01443~[hep-ex]$
- V. Khachatryan et al., Search for narrow resonances decaying to dijets in proton-proton collisions at $\sqrt(s) = 13$ TeV, Phys. Rev. Lett. 116 (2016) 071801, doi:10.1103/PhysRevLett.116.071801, arXiv:1512.01224 [hep-ex]
- V. Khachatryan et al., Reconstruction and identification of lepton decays to hadrons and $_{\tau}$ at CMS, JINST. 11 (2016) P01019, doi:10.1088/1748-0221/11/01/P01019, arXiv:1510.07488 [physics.ins-det]
- V. Khachatryan et al., **Measurement of top quark polarisation** in t-channel single top quark production, *JHEP*. 04 (2016) 073, doi:10.1007/JHEP04(2016)073, arXiv:1511.02138 [hep-ex]
- V. Khachatryan et al., Search for $W' \to tb$ in proton-proton collisions

- at $\sqrt{s}=8$ TeV, JHEP. 02 (2016) 122, doi:10.1007/JHEP02(2016)122, arXiv:1509.06051 [hep-ex]
- V. Khachatryan et al., Measurement of the $t\bar{t}$ production cross section in the all-jets final state in pp collisions at $\sqrt{s}=8 \text{TeV}$, Eur. Phys. J. C76 (2016) 128, doi:10.1140/epjc/s10052-016-3956-5, arXiv:1509.06076 [hep-ex]
- V. Khachatryan et al., Search for pair-produced vectorlike B quarks in proton-proton collisions at \sqrt{s} =8 TeV, Phys. Rev. D93 (2016) 112009, doi:10.1103/PhysRevD.93.112009, arXiv:1507.07129 [hep-ex]
- V. Khachatryan et al., Measurement of the top quark mass using proton-proton data at $\sqrt{(s)}=7$ and 8 TeV, Phys. Rev. D93 (2016) 072004, doi:10.1103/PhysRevD.93.072004, arXiv:1509.04044 [hep-ex]
- V. Khachatryan et al., **Angular analysis of the decay** $B^0 \to K^{*0} \mu^+ \mu^-$ from pp collisions at $\sqrt{s} = 8$ TeV, *Phys. Lett.* B753 (2016) 424–448, doi:10.1016/j.physletb.2015.12.020, arXiv:1507.08126 [hep-ex]
- V. Khachatryan et al., Search for a Higgs boson decaying into $\gamma^* \gamma \to \ell \ell \gamma$ with low dilepton mass in pp collisions at $\sqrt{s} = 8$ TeV, *Phys. Lett.* B753 (2016) 341–362, doi:10.1016/j.physletb.2015.12.039, arXiv:1507.03031 [hep-ex]
- V. Khachatryan et al., Measurement of transverse momentum relative to dijet systems in PbPb and pp collisions at $\sqrt{s_{\rm NN}} = 2.76$ TeV, JHEP. 01 (2016) 006, doi:10.1007/JHEP01(2016)006, arXiv:1509.09029 [nucl-ex]
- V. Khachatryan et al., **Measurement of the ratio** $B(B_s^0 \to J/\psi f_0(980)) / B(B_s^0 \to J/\psi \phi(1020))$ in pp collisions at $\sqrt{s} = 7$ TeV, *Phys. Lett.* B756 (2016) 84–102, doi:10.1016/j.physletb.2016.02.047, arXiv:1501.06089 [hep-ex]
- V. Khachatryan et al., Measurement of the inclusive jet cross section in pp collisions at $\sqrt{s}=2.76 \,\mathrm{TeV},~Eur.~Phys.~J.~C76~(2016)~265,$ doi:10.1140/epjc/s10052-016-4083-z, arXiv:1512.06212 [hep-ex]
- V. Khachatryan et al., Measurement of long-range near-side two-particle angular correlations in pp collisions at $\sqrt{s} = 13$ TeV, *Phys. Rev. Lett.* 116 (2016) 172302, doi:10.1103/PhysRevLett.116.172302, arXiv:1510.03068 [nucl-ex]
- V. Khachatryan et al., **Measurement of the charge asymmetry in top quark pair production in pp collisions at** $\sqrt(s) = 8$ **TeV using a template method**, *Phys. Rev.* D93 (2016) 034014, doi:10.1103/PhysRevD.93.034014, arXiv:1508.03862 [hep-ex]
- V. Khachatryan et al., Search for the production of an excited bottom quark decaying to tW in proton-proton collisions at $\sqrt{s} = 8$ TeV, JHEP. 01 (2016) 166, doi:10.1007/JHEP01(2016)166, arXiv:1509.08141 [hep-ex]
- V. Khachatryan et al., Measurement of $t\bar{t}$ production with additional jet activity, including b quark jets, in the dilepton decay channel

- using pp collisions at $\sqrt{s}=8$ TeV, Eur. Phys. J. C76 (2016) 379, doi:10.1140/epjc/s10052-016-4105-x, arXiv:1510.03072 [hep-ex]
- V. Khachatryan et al., Search for a very light NMSSM Higgs boson produced in decays of the 125 GeV scalar boson and decaying into τ leptons in pp collisions at $\sqrt{s}=8$ TeV, *JHEP*. 01 (2016) 079, doi:10.1007/JHEP01(2016)079, arXiv:1510.06534 [hep-ex]
- V. Khachatryan et al., Measurement of the CP-violating weak phase ϕ_s and the decay width difference $\Delta\Gamma_s$ using the B $_s^0 \to J/\psi\phi$ (1020) decay channel in pp collisions at $\sqrt{s}=8$ TeV, Phys. Lett. B757 (2016) 97–120, doi:10.1016/j.physletb.2016.03.046, arXiv:1507.07527 [hep-ex]
- V. Khachatryan et al., Search for vector-like charge 2/3 T quarks in proton-proton collisions at sqrt(s) = 8 TeV, Phys. Rev. D93 (2016) 012003, doi:10.1103/PhysRevD.93.012003, arXiv:1509.04177 [hep-ex]
- V. Khachatryan et al., Inclusive and differential measurements of the $t\bar{t}$ charge asymmetry in pp collisions at $\sqrt{s}=8$ TeV, Phys. Lett. B757 (2016) 154–179, doi:10.1016/j.physletb.2016.03.060, arXiv:1507.03119 [hep-ex]
- V. Khachatryan et al., Search for the associated production of a Higgs boson with a single top quark in proton-proton collisions at $\sqrt{s}=8$ TeV, *JHEP*. 06 (2016) 177, doi:10.1007/JHEP06(2016)177, arXiv:1509.08159 [hep-ex]
- V. Khachatryan et al., Event generator tunes obtained from underlying event and multiparton scattering measurements, Eur. Phys. J. C76 (2016) 155, doi:10.1140/epjc/s10052-016-3988-x, arXiv:1512.00815 [hep-ex]
- V. Khachatryan et al., Search for supersymmetry in events with soft leptons, low jet multiplicity, and missing transverse energy in proton–proton collisions at \sqrt{s} =8 TeV, Phys. Lett. B759 (2016) 9–35, doi:10.1016/j.physletb.2016.05.033, arXiv:1512.08002 [hep-ex]
- V. Khachatryan et al., Search for W' decaying to tau lepton and neutrino in proton-proton collisions at $\sqrt(s) = 8$ TeV, Phys. Lett. B755 (2016) 196–216, doi:10.1016/j.physletb.2016.02.002, arXiv:1508.04308 [hep-ex]
- V. Khachatryan et al., Study of Z boson production in pPb collisions at $\sqrt{s_{NN}}=5.02$ TeV, Phys. Lett. B759 (2016) 36–57, doi:10.1016/j.physletb.2016.05.044, arXiv:1512.06461 [hep-ex]
- V. Khachatryan et al., Search for single production of scalar leptoquarks in proton-proton collisions at $\sqrt{s} = 8TeV$, Phys. Rev. D93 (2016) 032005, doi:10.1103/PhysRevD.95.039906, arXiv:1509.03750 [hep-ex]
- V. Khachatryan et al., Measurement of differential cross sections for Higgs boson production in the diphoton decay channel in pp collisions

- at $\sqrt{s} = 8$ TeV, Eur. Phys. J. C76 (2016) 13, doi:10.1140/epjc/s10052-015-3853-3, arXiv:1508.07819 [hep-ex]
- V. Khachatryan et al., Search for excited leptons in proton-proton collisions at sqrt(s) = 8 TeV, *JHEP*. 03 (2016) 125, doi:10.1007/JHEP03(2016)125, arXiv:1511.01407 [hep-ex]
- V. Khachatryan et al., Observation of top quark pairs produced in association with a vector boson in pp collisions at $\sqrt{s} = 8$ TeV, *JHEP*. 01 (2016) 096, doi:10.1007/JHEP01(2016)096, arXiv:1510.01131 [hep-ex]
- V. Khachatryan et al., Search for resonant $t\bar{t}$ production in proton-proton collisions at $\sqrt{s}=8$ TeV, Phys. Rev. D93 (2016) 012001, doi:10.1103/PhysRevD.93.012001, arXiv:1506.03062 [hep-ex]
- V. Khachatryan et al., Transverse momentum spectra of inclusive b jets in pPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, Phys. Lett. B754 (2016) 59, doi:10.1016/j.physletb.2016.01.010, arXiv:1510.03373 [nucl-ex]
- V. Khachatryan et al., Searches for a heavy scalar boson H decaying to a pair of 125 GeV Higgs bosons hh or for a heavy pseudoscalar boson A decaying to Zh, in the final states with $h \to \tau\tau$, Phys. Lett. B755 (2016) 217–244, doi:10.1016/j.physletb.2016.01.056, arXiv:1510.01181 [hep-ex]
- V. Khachatryan et al., Measurement of spin correlations in $t\bar{t}$ production using the matrix element method in the muon+jets final state in pp collisions at $\sqrt{s}=8$ TeV, *Phys. Lett.* B758 (2016) 321–346, doi:10.1016/j.physletb.2016.05.005, arXiv:1511.06170 [hep-ex]
- V. Khachatryan et al., Study of B Meson Production in p +Pb Collisions at $\sqrt{s_{NN}}=5.02$ TeV Using Exclusive Hadronic Decays, *Phys. Rev. Lett.* 116 (2016) 032301, doi:10.1103/PhysRevLett.116.032301, arXiv:1508.06678 [nucl-ex]
- V. Khachatryan et al., Measurement of the top quark pair production cross section in proton-proton collisions at $\sqrt(s) = 13$ TeV, *Phys. Rev. Lett.* 116 (2016) 052002, doi:10.1103/PhysRevLett.116.052002, arXiv:1510.05302 [hep-ex]
- V. Khachatryan et al., Search for pair production of first and second generation leptoquarks in proton-proton collisions at $\sqrt{s}=8$ TeV, Phys. Rev. D93 (2016) 032004, doi:10.1103/PhysRevD.93.032004, arXiv:1509.03744 [hep-ex]
- V. Khachatryan et al., Search for exotic decays of a Higgs boson into undetectable particles and one or more photons, *Phys. Lett.* B753 (2016) 363–388, doi:10.1016/j.physletb.2015.12.017, arXiv:1507.00359 [hep-ex]
- V. Khachatryan et al., A search for pair production of new light bosons decaying into muons, *Phys. Lett.* B752 (2016) 146–168,

- doi:10.1016/j.physletb.2015.10.067, arXiv:1506.00424 [hep-ex]
- V. Khachatryan et al., Measurement of differential and integrated fiducial cross sections for Higgs boson production in the four-lepton decay channel in pp collisions at $\sqrt{s}=7$ and 8 TeV, *JHEP*. 04 (2016) 005, doi:10.1007/JHEP04(2016)005, arXiv:1512.08377 [hep-ex]
- V. Khachatryan et al., Measurement of the inelastic cross section in proton–lead collisions at $\sqrt{s_{NN}}=5.02$ TeV, Phys. Lett. B759 (2016) 641–662, doi:10.1016/j.physletb.2016.06.027, arXiv:1509.03893 [hep-ex]
- V. Khachatryan et al., Search for neutral resonances decaying into a Z boson and a pair of b jets or τ leptons, *Phys. Lett.* B759 (2016) 369–394, doi:10.1016/j.physletb.2016.05.087, arXiv:1603.02991 [hep-ex]
- V. Khachatryan et al., **Search for heavy resonances decaying to two Higgs bosons in final states containing four b quarks**, *Eur. Phys. J.* C76 (2016) 371, doi:10.1140/epjc/s10052-016-4206-6, arXiv:1602.08762 [hep-ex]
- V. Khachatryan et al., Search for Higgs boson off-shell production in proton-proton collisions at 7 and 8 TeV and derivation of constraints on its total decay width, *JHEP*. 09 (2016) 051, doi:10.1007/JHEP09(2016)051, arXiv:1605.02329 [hep-ex]
- V. Khachatryan et al., Search for narrow resonances in dijet final states at $\sqrt(s) = 8$ TeV with the novel CMS technique of data scouting, *Phys. Rev. Lett.* 117 (2016) 031802, doi:10.1103/PhysRevLett.117.031802, arXiv:1604.08907 [hep-ex]
- V. Khachatryan et al., Correlations between jets and charged particles in PbPb and pp collisions at $\sqrt{s_{\rm NN}}=2.76$ TeV, *JHEP*. 02 (2016) 156, doi:10.1007/JHEP02(2016)156, arXiv:1601.00079 [nucl-ex]
- V. Khachatryan et al., Search for s channel single top quark production in pp collisions at $\sqrt{s} = 7$ and 8 TeV, *JHEP*. 09 (2016) 027, doi:10.1007/JHEP09(2016)027, arXiv:1603.02555 [hep-ex]
- V. Khachatryan et al., Measurement of the W boson helicity fractions in the decays of top quark pairs to lepton + jets final states produced in pp collisions at $\sqrt{s} = 8$ TeV, Phys. Lett. B762 (2016) 512–534, doi:10.1016/j.physletb.2016.10.007, arXiv:1605.09047 [hep-ex]
- V. Khachatryan et al., Measurement of the integrated and differential $t\bar{t}$ production cross sections for high- p_t top quarks in pp collisions at $\sqrt{s} = 8$ TeV, Phys. Rev. D94 (2016) 072002, doi:10.1103/PhysRevD.94.072002, arXiv:1605.00116 [hep-ex]
- V. Khachatryan et al., Search for Resonant Production of High-Mass Photon Pairs in Proton-Proton Collisions at \sqrt{s} =8 and 13 TeV, Phys. Rev. Lett. 117 (2016) 051802, doi:10.1103/PhysRevLett.117.051802,

- arXiv:1606.04093 [hep-ex]
- V. Khachatryan et al., Measurement of dijet azimuthal decorrelation in pp collisions at $\sqrt{s} = 8$ TeV, Eur. Phys. J. C76 (2016) 536, doi:10.1140/epjc/s10052-016-4346-8, arXiv:1602.04384 [hep-ex]
- V. Khachatryan et al., Search for supersymmetry in electroweak production with photons and large missing transverse energy in pp collisions at $\sqrt{s}=8$ TeV, Phys. Lett. B759 (2016) 479–500, doi:10.1016/j.physletb.2016.05.088, arXiv:1602.08772 [hep-ex]
- V. Khachatryan et al., Searches for R-parity-violating supersymmetry in pp collisions at $\sqrt(s) = 8$ TeV in final states with 0-4 leptons, Phys. Rev. D94 (2016) 112009, doi:10.1103/PhysRevD.94.112009, arXiv:1606.08076 [hep-ex]
- V. Khachatryan et al., Search for supersymmetry in the multijet and missing transverse momentum final state in pp collisions at 13 TeV, *Phys. Lett.* B758 (2016) 152–180, doi:10.1016/j.physletb.2016.05.002, arXiv:1602.06581 [hep-ex]
- V. Khachatryan et al., Search for dark matter in proton-proton collisions at 8 TeV with missing transverse momentum and vector boson tagged jets, *JHEP*. 12 (2016) 083, doi:10.1007/JHEP12(2016)083, arXiv:1607.05764 [hep-ex]
- V. Khachatryan et al., Measurement of the t-tbar production cross section in the e-mu channel in proton-proton collisions at sqrt(s) = 7 and 8 TeV, *JHEP*. 08 (2016) 029, doi:10.1007/JHEP08(2016)029, arXiv:1603.02303 [hep-ex]
- V. Khachatryan et al., Evidence for exclusive $\gamma\gamma \to W^+W^-$ production and constraints on anomalous quartic gauge couplings in pp collisions at $\sqrt{s}=7$ and 8 TeV, JHEP.~08~(2016)~119,~doi:10.1007/JHEP08(2016)119,~arXiv:1604.04464~[hep-ex]
- V. Khachatryan et al., Phenomenological MSSM interpretation of CMS searches in pp collisions at sqrt(s) = 7 and 8 TeV, *JHEP*. 10 (2016) 129, doi:10.1007/JHEP10(2016)129, arXiv:1606.03577 [hep-ex]
- V. Khachatryan et al., Search for direct pair production of supersymmetric top quarks decaying to all-hadronic final states in pp collisions at $\sqrt{s}=8{\rm TeV},~Eur.~Phys.~J.~C76~(2016)~460,~doi:10.1140/epjc/s10052-016-4292-5,~arXiv:1603.00765~[hep-ex]$
- V. Khachatryan et al., Search for heavy Majorana neutrinos in e \pm e \pm + jets and e \pm μ \pm + jets events in proton-proton collisions at $\sqrt{s} = 8$ TeV, *JHEP*. 04 (2016) 169, doi:10.1007/JHEP04(2016)169, arXiv:1603.02248 [hep-ex]
- V. Khachatryan et al., Measurement of the differential cross sections for top quark pair production as a function of kinematic event variables

- in pp collisions at \sqrt{s} =7 and 8 TeV, *Phys. Rev.* D94 (2016) 052006, doi:10.1103/PhysRevD.94.052006, arXiv:1607.00837 [hep-ex]
- V. Khachatryan et al., Search for lepton flavour violating decays of heavy resonances and quantum black holes to an e μ pair in proton-proton collisions at $\sqrt{s}=8$ TeV, Eur. Phys. J. C76 (2016) 317, doi:10.1140/epjc/s10052-016-4149-y, arXiv:1604.05239 [hep-ex]
- V. Khachatryan et al., Measurement of the mass of the top quark in decays with a J/ψ meson in pp collisions at 8 TeV, *JHEP*. 12 (2016) 123, doi:10.1007/JHEP12(2016)123, arXiv:1608.03560 [hep-ex]
- V. Khachatryan et al., Measurement of the differential cross section and charge asymmetry for inclusive pp \rightarrow W[±] + X production at \sqrt{s} = 8 TeV, Eur. Phys. J. C76 (2016) 469, doi:10.1140/epjc/s10052-016-4293-4, arXiv:1603.01803 [hep-ex]
- V. Khachatryan et al., Search for direct pair production of scalar top quarks in the single- and dilepton channels in proton-proton collisions at $\sqrt{s}=8$ TeV, JHEP.~07~(2016)~027,~doi:10.1007/JHEP07(2016)027,~arXiv:1602.03169~[hep-ex]
- V. Khachatryan et al., Measurement of electroweak production of a W boson and two forward jets in proton-proton collisions at $\sqrt{s} = 8$ TeV, *JHEP*. 11 (2016) 147, doi:10.1007/JHEP11(2016)147, arXiv:1607.06975 [hep-ex]
- V. Khachatryan et al., Search for dark matter particles in proton-proton collisions at $\sqrt{s} = 8$ TeV using the razor variables, *JHEP*. 12 (2016) 088, doi:10.1007/JHEP12(2016)088, arXiv:1603.08914 [hep-ex]
- V. Khachatryan et al., Studies of inclusive four-jet production with two b-tagged jets in proton-proton collisions at 7 TeV, Phys. Rev. D94 (2016) 112005, doi:10.1103/PhysRevD.94.112005, arXiv:1609.03489 [hep-ex]
- V. Khachatryan et al., Search for lepton flavour violating decays of the Higgs boson to $e\tau$ and $e\mu$ in proton–proton collisions at $\sqrt{s}=$ 8 TeV, Phys. Lett. B763 (2016) 472–500, doi:10.1016/j.physletb.2016.09.062, arXiv:1607.03561 [hep-ex]
- V. Khachatryan et al., Search for two Higgs bosons in final states containing two photons and two bottom quarks in proton-proton collisions at 8 TeV, *Phys. Rev.* D94 (2016) 052012, doi:10.1103/PhysRevD.94.052012, arXiv:1603.06896 [hep-ex]
- V. Khachatryan et al., Search for long-lived charged particles in proton-proton collisions at $\sqrt{s} = 13$ TeV, *Phys. Rev.* D94 (2016) 112004, doi:10.1103/PhysRevD.94.112004, arXiv:1609.08382 [hep-ex]
- V. Khachatryan et al., Decomposing transverse momentum balance contributions for quenched jets in PbPb collisions at $\sqrt{s_{\mathrm{NN}}} = 2.76$ TeV,

- JHEP.~11~(2016)~055,~doi:10.1007/JHEP11(2016)055,~arXiv:1609.02466~[nucl-ex]
- V. Khachatryan et al., Combined search for anomalous pseudoscalar HVV couplings in VH(H \rightarrow $b\bar{b}$) production and H \rightarrow VV decay, *Phys. Lett.* B759 (2016) 672–696, doi:10.1016/j.physletb.2016.06.004, arXiv:1602.04305 [hep-ex]
- V. Khachatryan et al., Measurement of the ZZ production cross section and $\mathbf{Z} \to \ell^+ \ell^- \ell'^+ \ell'^-$ branching fraction in pp collisions at \sqrt{s} =13 TeV, *Phys. Lett.* B763 (2016) 280–303, doi:10.1016/j.physletb.2017.09.030, arXiv:1607.08834 [hep-ex]
- V. Khachatryan et al., Azimuthal decorrelation of jets widely separated in rapidity in pp collisions at $\sqrt{s}=7$ TeV, *JHEP*. 08 (2016) 139, doi:10.1007/JHEP08(2016)139, arXiv:1601.06713 [hep-ex]
- V. Khachatryan et al., Search for supersymmetry in pp collisions at $\sqrt{s}=13$ TeV in the single-lepton final state using the sum of masses of large-radius jets, *JHEP*. 08 (2016) 122, doi:10.1007/JHEP08(2016)122, arXiv:1605.04608 [hep-ex]
- G. Aad et al., Measurements of the Higgs boson production and decay rates and constraints on its couplings from a combined ATLAS and CMS analysis of the LHC pp collision data at $\sqrt{s} = 7$ and 8 TeV, *JHEP*. 08 (2016) 045, doi:10.1007/JHEP08(2016)045, arXiv:1606.02266 [hep-ex]
- V. Khachatryan et al., $\Upsilon(nS)$ polarizations versus particle multiplicity in pp collisions at $\sqrt{s}=7$ TeV, *Phys. Lett.* B761 (2016) 31–52, doi:10.1016/j.physletb.2016.07.065, arXiv:1603.02913 [hep-ex]
- V. Khachatryan et al., Measurement of the double-differential inclusive jet cross section in proton–proton collisions at $\sqrt{s}=13 \text{TeV}$, Eur. Phys. J. C76 (2016) 451, doi:10.1140/epjc/s10052-016-4286-3, arXiv:1605.04436 [hep-ex]
- V. Khachatryan et al., Measurement of the top quark mass using charged particles in pp collisions at $\sqrt{s} = 8$ TeV, Phys. Rev. D93 (2016) 092006, doi:10.1103/PhysRevD.93.092006, arXiv:1603.06536 [hep-ex]
- V. Khachatryan et al., Measurement of inclusive jet production and nuclear modifications in pPb collisions at $\sqrt{s_{_{\mathrm{NN}}}} = 5.02$ TeV, Eur. Phys. J. C76 (2016) 372, doi:10.1140/epjc/s10052-016-4205-7, arXiv:1601.02001 [nucl-ex]
- V. Khachatryan et al., Search for R-parity violating decays of a top squark in proton-proton collisions at $\sqrt{s} = 8$ TeV, Phys. Lett. B760 (2016) 178–201, doi:10.1016/j.physletb.2016.06.039, arXiv:1602.04334 [hep-ex]
- V. Khachatryan et al., Search for massive WH resonances decaying into the $\ell\nu$ bb final state at $\sqrt{s}=8$ TeV, Eur. Phys. J. C76 (2016) 237,

- doi:10.1140/epjc/s10052-016-4067-z, arXiv:1601.06431 [hep-ex]
- V. Khachatryan et al., Measurement of the $Z\gamma \to \nu\bar{\nu}\gamma$ production cross section in pp collisions at $\sqrt{s}=8$ TeV and limits on anomalous $ZZ\gamma$ and $Z\gamma\gamma$ trilinear gauge boson couplings, *Phys. Lett.* B760 (2016) 448–468, doi:10.1016/j.physletb.2016.06.080, arXiv:1602.07152 [hep-ex]
- V. Khachatryan et al., Forward–backward asymmetry of Drell–Yan lepton pairs in pp collisions at $\sqrt{s}=8 \,\mathrm{TeV}$, Eur. Phys. J. C76 (2016) 325, doi:10.1140/epjc/s10052-016-4156-z, arXiv:1601.04768 [hep-ex]
- V. Khachatryan et al., Search for supersymmetry in pp collisions at sqrt(s) = 8 TeV in final states with boosted W bosons and b jets using razor variables, *Phys. Rev.* D93 (2016) 092009, doi:10.1103/PhysRevD.93.092009, arXiv:1602.02917 [hep-ex]
- G. Aad et al., Combined Measurement of the Higgs Boson Mass in pp Collisions at $\sqrt{s}=7$ and 8 TeV with the AT-LAS and CMS Experiments, *Phys. Rev. Lett.* 114 (2015) 191803, doi:10.1103/PhysRevLett.114.191803, arXiv:1503.07589 [hep-ex]
- S. Chatrchyan et al., Study of Z production in PbPb and pp collisions at $\sqrt{s_{\rm NN}} = 2.76$ TeV in the dimuon and dielectron decay channels, *JHEP*. 03 (2015) 022, doi:10.1007/JHEP03(2015)022, arXiv:1410.4825 [nucl-ex]
- S. Chatrchyan et al., Measurement of jet multiplicity distributions in $t\bar{t}$ production in pp collisions at $\sqrt{s}=7 \,\mathrm{TeV}$, Eur. Phys. J. C74 (2015) 3014, doi:10.1140/epjc/s10052-014-3014-0, arXiv:1404.3171 [hep-ex]
- V. Khachatryan et al., Searches for supersymmetry based on events with b jets and four W bosons in pp collisions at 8 TeV, *Phys. Lett.* B745 (2015) 5–28, doi:10.1016/j.physletb.2015.04.002, arXiv:1412.4109 [hep-ex]
- V. Khachatryan et al., Search for stealth supersymmetry in events with jets, either photons or leptons, and low missing transverse momentum in pp collisions at 8 TeV, *Phys. Lett.* B743 (2015) 503–525, doi:10.1016/j.physletb.2015.03.017, arXiv:1411.7255 [hep-ex]
- V. Khachatryan et al., Search for disappearing tracks in proton-proton collisions at $\sqrt{s} = 8$ TeV, *JHEP*. 01 (2015) 096, doi:10.1007/JHEP01(2015)096, arXiv:1411.6006 [hep-ex]
- V. Khachatryan et al., Search for long-lived particles that decay into final states containing two electrons or two muons in proton-proton collisions at $\sqrt{s}=8$ TeV, *Phys. Rev.* D91 (2015) 052012, doi:10.1103/PhysRevD.91.052012, arXiv:1411.6977 [hep-ex]
- V. Khachatryan et al., Measurements of differential and double-differential Drell-Yan cross sections in proton-proton collisions at 8 TeV, Eur. Phys. J. C75 (2015) 147, doi:10.1140/epjc/s10052-015-3364-2,

- arXiv:1412.1115 [hep-ex]
- V. Khachatryan et al., Measurement of the inclusive 3-jet production differential cross section in proton–proton collisions at 7 TeV and determination of the strong coupling constant in the TeV range, Eur. Phys. J. C75 (2015) 186, doi:10.1140/epjc/s10052-015-3376-y, arXiv:1412.1633 [hep-ex]
- V. Khachatryan et al., Search for Long-Lived Neutral Particles Decaying to Quark-Antiquark Pairs in Proton-Proton Collisions at $\sqrt{s} = 8$ TeV, Phys. Rev. D91 (2015) 012007, doi:10.1103/PhysRevD.91.012007, arXiv:1411.6530 [hep-ex]
- V. Khachatryan et al., Measurement of the $pp \to ZZ$ production cross section and constraints on anomalous triple gauge couplings in four-lepton final states at $\sqrt{s}=8$ TeV, Phys. Lett. B740 (2015) 250–272, doi:10.1016/j.physletb.2016.04.010, arXiv:1406.0113 [hep-ex]
- V. Khachatryan et al., Measurement of the cross section ratio $\sigma_{\bar{t}tb\bar{b}}/\sigma_{\bar{t}tj\bar{j}}$ in pp collisions at $\sqrt{s}=8$ TeV, Phys. Lett. B746 (2015) 132–153, doi:10.1016/j.physletb.2015.04.060, arXiv:1411.5621 [hep-ex]
- V. Khachatryan et al., Search for a standard model-like Higgs boson in the $\mu^+\mu^-$ and e^+e^- decay channels at the LHC, *Phys. Lett.* B744 (2015) 184–207, doi:10.1016/j.physletb.2015.03.048, arXiv:1410.6679 [hep-ex]
- V. Khachatryan et al., Search for quark contact interactions and extra spatial dimensions using dijet angular distributions in proton-proton collisions at $\sqrt{s}=8$ TeV, Phys. Lett. B746 (2015) 79–99, doi:10.1016/j.physletb.2015.04.042, arXiv:1411.2646 [hep-ex]
- V. Khachatryan et al., Measurement of electroweak production of two jets in association with a Z boson in proton-proton collisions at $\sqrt{s} = 8 \text{TeV}$, Eur. Phys. J. C75 (2015) 66, doi:10.1140/epjc/s10052-014-3232-5, arXiv:1410.3153 [hep-ex]
- V. Khachatryan et al., Search for physics beyond the standard model in dilepton mass spectra in proton-proton collisions at $\sqrt{s} = 8$ TeV, *JHEP*. 04 (2015) 025, doi:10.1007/JHEP04(2015)025, arXiv:1412.6302 [hep-ex]
- V. Khachatryan et al., Performance of the CMS missing transverse momentum reconstruction in pp data at $\sqrt{s}=8$ TeV, JINST. 10 (2015) P02006, doi:10.1088/1748-0221/10/02/P02006, arXiv:1411.0511 [physics.ins-det]
- V. Khachatryan et al., Precise determination of the mass of the Higgs boson and tests of compatibility of its couplings with the standard model predictions using proton collisions at 7 and 8 TeV, Eur. Phys. J. C75 (2015) 212, doi:10.1140/epjc/s10052-015-3351-7, arXiv:1412.8662 [hep-ex]
- V. Khachatryan et al., Long-range two-particle correlations of strange

- hadrons with charged particles in pPb and PbPb collisions at LHC energies, $Phys.\ Lett.\ B742\ (2015)\ 200-224,\ doi:10.1016/j.physletb.2015.01.034,\ arXiv:1409.3392\ [nucl-ex]$
- V. Khachatryan et al., Constraints on the spin-parity and anomalous HVV couplings of the Higgs boson in proton collisions at 7 and 8 TeV, *Phys. Rev.* D92 (2015) 012004, doi:10.1103/PhysRevD.92.012004, arXiv:1411.3441 [hep-ex]
- V. Khachatryan et al., Search for pair-produced resonances decaying to jet pairs in proton–proton collisions at $\sqrt{s} = 8$ TeV, *Phys. Lett.* B747 (2015) 98–119, doi:10.1016/j.physletb.2015.04.045, arXiv:1412.7706 [hep-ex]
- V. Khachatryan et al., Search for Displaced Supersymmetry in events with an electron and a muon with large impact parameters, *Phys. Rev. Lett.* 114 (2015) 061801, doi:10.1103/PhysRevLett.114.061801, arXiv:1409.4789 [hep-ex]
- V. Khachatryan et al., Measurement of the ratio of the production cross sections times branching fractions of $B_c^{\pm} \to J/\psi \pi^{\pm}$ and $B^{\pm} \to J/\psi K^{\pm}$ and $\mathcal{B}(B_c^{\pm} \to J/\psi \pi^{\pm} \pi^{\mp})/\mathcal{B}(B_c^{\pm} \to J/\psi \pi^{\pm})$ in pp collisions at $\sqrt{s}=$ 7 TeV, JHEP. 01 (2015) 063, doi:10.1007/JHEP01(2015)063, arXiv:1410.5729 [hep-ex]
- V. Khachatryan et al., Measurement of the production cross section ratio $\sigma(\mathbf{Xb2(1P)})$ / $\sigma(\mathbf{Xb1(1P)})$ in pp collisions at $\sqrt{s}=8$ TeV, Phys. Lett. B743 (2015) 383–402, doi:10.1016/j.physletb.2015.02.048, arXiv:1409.5761 [hep-ex]
- V. Khachatryan et al., Search for dark matter, extra dimensions, and unparticles in monojet events in proton–proton collisions at $\sqrt{s}=8$ TeV, Eur. Phys. J. C75 (2015) 235, doi:10.1140/epjc/s10052-015-3451-4, arXiv:1408.3583 [hep-ex]
- V. Khachatryan et al., Search for physics beyond the standard model in final states with a lepton and missing transverse energy in proton-proton collisions at sqrt(s) = 8 TeV, Phys. Rev. D91 (2015) 092005, doi:10.1103/PhysRevD.91.092005, arXiv:1408.2745 [hep-ex]
- V. Khachatryan et al., Search for Monotop Signatures in Proton-Proton Collisions at $\sqrt{s}=8$ TeV, Phys. Rev. Lett. 114 (2015) 101801, doi:10.1103/PhysRevLett.114.101801, arXiv:1410.1149 [hep-ex]
- V. Khachatryan et al., Differential cross section measurements for the production of a W boson in association with jets in proton–proton collisions at $\sqrt{s}=7$ TeV, *Phys. Lett.* B741 (2015) 12–37, doi:10.1016/j.physletb.2014.12.003, arXiv:1406.7533 [hep-ex]
- V. Khachatryan et al., Measurement of the W boson helicity in events with a single reconstructed top quark in pp collisions at $\sqrt{s} = 8$ TeV,

- JHEP. 01 (2015) 053, doi:10.1007/JHEP01(2015)053, arXiv:1410.1154 [hep-ex]
- V. Khachatryan et al., Constraints on parton distribution functions and extraction of the strong coupling constant from the inclusive jet cross section in pp collisions at $\sqrt{s} = 7 \text{TeV}$, Eur. Phys. J. C75 (2015) 288, doi:10.1140/epjc/s10052-015-3499-1, arXiv:1410.6765 [hep-ex]
- V. Khachatryan et al., Search for new resonances decaying via WZ to leptons in proton-proton collisions at $\sqrt{s} = 8$ TeV, *Phys. Lett.* B740 (2015) 83–104, doi:10.1016/j.physletb.2014.11.026, arXiv:1407.3476 [hep-ex]
- V. Khachatryan et al., Measurements of jet multiplicity and differential production cross sections of Z+ jets events in proton-proton collisions at $\sqrt{s}=7$ TeV, Phys. Rev. D91 (2015) 052008, doi:10.1103/PhysRevD.91.052008, arXiv:1408.3104 [hep-ex]
- V. Khachatryan et al., Search for vector-like T quarks decaying to top quarks and Higgs bosons in the all-hadronic channel using jet substructure, *JHEP*. 06 (2015) 080, doi:10.1007/JHEP06(2015)080, arXiv:1503.01952 [hep-ex]
- V. Khachatryan et al., Search for the standard model Higgs boson produced through vector boson fusion and decaying to $b\bar{b}$, Phys. Rev. D92 (2015) 032008, doi:10.1103/PhysRevD.92.032008, arXiv:1506.01010 [hep-ex]
- V. Khachatryan et al., Search for Third-Generation Scalar Leptoquarks in the t τ Channel in Proton-Proton Collisions at $\sqrt{s}=8$ TeV, JHEP. 07 (2015) 042, doi:10.1007/JHEP11(2016)056, arXiv:1503.09049 [hep-ex]
- V. Khachatryan et al., Search for a Higgs boson in the mass range from 145 to 1000 GeV decaying to a pair of W or Z bosons, *JHEP*. 10 (2015) 144, doi:10.1007/JHEP10(2015)144, arXiv:1504.00936 [hep-ex]
- V. Khachatryan et al., Search for supersymmetry with photons in pp collisions at \sqrt{s} =8 TeV, Phys. Rev. D92 (2015) 072006, doi:10.1103/PhysRevD.92.072006, arXiv:1507.02898 [hep-ex]
- V. Khachatryan et al., Production of leading charged particles and leading charged-particle jets at small transverse momenta in pp collisions at sqrt(s) = 8 TeV, Phys. Rev. D92 (2015) 112001, doi:10.1103/PhysRevD.92.112001, arXiv:1507.00233 [hep-ex]
- V. Khachatryan et al., Search for heavy Majorana neutrinos in $\mu^{\pm}\mu^{\pm}$ + jets events in proton-proton collisions at $\sqrt{s} = 8$ TeV, Phys. Lett. B748 (2015) 144–166, doi:10.1016/j.physletb.2015.06.070, arXiv:1501.05566 [hep-ex]
- V. Khachatryan et al., Measurement of diffraction dissociation cross sections in pp collisions at $\sqrt{s}=7$ TeV, Phys. Rev. D92 (2015) 012003, doi:10.1103/PhysRevD.92.012003, arXiv:1503.08689 [hep-ex]
- V. Khachatryan et al., Study of W boson production in pPb collisions at $\sqrt{s_{\mathrm{NN}}}$ = 5.02 TeV, Phys. Lett. B750 (2015) 565–586,

- doi:10.1016/j.physletb.2015.09.057, arXiv:1503.05825 [nucl-ex]
- V. Khachatryan et al., Performance of Electron Reconstruction and Selection with the CMS Detector in Proton-Proton Collisions at $\sqrt{s} = 8$ TeV, JINST. 10 (2015) P06005, doi:10.1088/1748-0221/10/06/P06005, arXiv:1502.02701 [physics.ins-det]
- V. Khachatryan et al., Search for a Standard Model Higgs Boson Produced in Association with a Top-Quark Pair and Decaying to Bottom Quarks Using a Matrix Element Method, Eur. Phys. J. C75 (2015) 251, doi:10.1140/epjc/s10052-015-3454-1, arXiv:1502.02485 [hep-ex]
- V. Khachatryan et al., Comparison of the Z/ γ^* + jets to γ + jets cross sections in pp collisions at $\sqrt{s}=8$ TeV, JHEP. 10 (2015) 128, doi:10.1007/JHEP04(2016)010, arXiv:1505.06520 [hep-ex]
- V. Khachatryan et al., Performance of Photon Reconstruction and Identification with the CMS Detector in Proton-Proton Collisions at sqrt(s) = 8 TeV, JINST. 10 (2015) P08010, doi:10.1088/1748-0221/10/08/P08010, arXiv:1502.02702 [physics.ins-det]
- V. Khachatryan et al., Search for neutral color-octet weak-triplet scalar particles in proton-proton collisions at $\sqrt{s} = 8$ TeV, *JHEP*. 09 (2015) 201, doi:10.1007/JHEP09(2015)201, arXiv:1505.08118 [hep-ex]
- V. Khachatryan et al., Search for Decays of Stopped Long-Lived Particles Produced in Proton-Proton Collisions at $\sqrt{s} = 8$ TeV, Eur. Phys. J. C75 (2015) 151, doi:10.1140/epjc/s10052-015-3367-z, arXiv:1501.05603 [hep-ex]
- V. Khachatryan et al., Pseudorapidity distribution of charged hadrons in proton-proton collisions at $\sqrt{s} = 13$ TeV, Phys. Lett. B751 (2015) 143–163, doi:10.1016/j.physletb.2015.10.004, arXiv:1507.05915 [hep-ex]
- V. Khachatryan et al., Measurement of the underlying event activity using charged-particle jets in proton-proton collisions at sqrt(s) = 2.76 TeV, *JHEP*. 09 (2015) 137, doi:10.1007/JHEP09(2015)137, arXiv:1507.07229 [hep-ex]
- V. Khachatryan et al., Measurement of the Z Production Cross Section in pp Collisions at 8 TeV and Search for Anomalous Triple Gauge Boson Couplings, *JHEP*. 04 (2015) 164, doi:10.1007/JHEP04(2015)164, arXiv:1502.05664 [hep-ex]
- V. Khachatryan et al., **Search for Lepton-Flavour-Violating Decays of the Higgs Boson**, *Phys. Lett.* B749 (2015) 337–362, doi:10.1016/j.physletb.2015.07.053, arXiv:1502.07400 [hep-ex]
- V. Khachatryan et al., Search for supersymmetry in the vector-boson fusion topology in proton-proton collisions at $\sqrt{s}=8$ TeV, *JHEP*. 11 (2015) 189, doi:10.1007/JHEP11(2015)189, arXiv:1508.07628 [hep-ex]
- V. Khachatryan et al., Search for a pseudoscalar boson decaying into a

- **Z** boson and the 125 GeV Higgs boson in $\ell^+\ell^-b\bar{b}$ final states, *Phys. Lett.* B748 (2015) 221–243, doi:10.1016/j.physletb.2015.07.010, arXiv:1504.04710 [hep-ex]
- V. Khachatryan et al., Constraints on the pMSSM, AMSB model and on other models from the search for long-lived charged particles in proton-proton collisions at sqrt(s) = 8 TeV, Eur. Phys. J. C75 (2015) 325, doi:10.1140/epjc/s10052-015-3533-3, arXiv:1502.02522 [hep-ex]
- V. Khachatryan et al., Limits on the Higgs boson lifetime and width from its decay to four charged leptons, *Phys. Rev.* D92 (2015) 072010, doi:10.1103/PhysRevD.92.072010, arXiv:1507.06656 [hep-ex]
- V. Khachatryan et al., Search for the production of dark matter in association with top-quark pairs in the single-lepton final state in proton-proton collisions at sqrt(s) = 8 TeV, JHEP. 06 (2015) 121, doi:10.1007/JHEP06(2015)121, arXiv:1504.03198 [hep-ex]
- V. Khachatryan et al., Measurement of the Z boson differential cross section in transverse momentum and rapidity in proton—proton collisions at 8 TeV, *Phys. Lett.* B749 (2015) 187–209, doi:10.1016/j.physletb.2015.07.065, arXiv:1504.03511 [hep-ex]
- V. Khachatryan et al., Evidence for transverse momentum and pseudorapidity dependent event plane fluctuations in PbPb and pPb collisions, *Phys. Rev.* C92 (2015) 034911, doi:10.1103/PhysRevC.92.034911, arXiv:1503.01692 [nucl-ex]
- V. Khachatryan et al., Measurement of the differential cross section for top quark pair production in pp collisions at $\sqrt{s} = 8\text{TeV}$, Eur. Phys. J. C75 (2015) 542, doi:10.1140/epjc/s10052-015-3709-x, arXiv:1505.04480 [hep-ex]
- V. Khachatryan et al., Angular coefficients of **Z** bosons produced in pp collisions at $\sqrt{s}=8$ TeV and decaying to $\mu^+\mu^-$ as a function of transverse momentum and rapidity, *Phys. Lett.* B750 (2015) 154–175, doi:10.1016/j.physletb.2015.08.061, arXiv:1504.03512 [hep-ex]
- V. Khachatryan et al., Measurements of the ZZ production cross sections in the $2l2\nu$ channel in proton–proton collisions at $\sqrt{s}=7$ and 8 TeV and combined constraints on triple gauge couplings, Eur. Phys. J. C75 (2015) 511, doi:10.1140/epjc/s10052-015-3706-0, arXiv:1503.05467 [hep-ex]
- V. Khachatryan et al., Search for Supersymmetry Using Razor Variables in Events with b-Tagged Jets in pp Collisions at $\sqrt{s}=8$ TeV, Phys. Rev. D91 (2015) 052018, doi:10.1103/PhysRevD.91.052018, arXiv:1502.00300 [hep-ex]
- V. Khachatryan et al., Search for diphoton resonances in the mass range from 150 to 850 GeV in pp collisions at $\sqrt{s} = 8$ TeV, Phys. Lett. B750

- (2015) 494–519, doi:10.1016/j.physletb.2015.09.062, arXiv:1506.02301 [hep-ex]
- V. Khachatryan et al., Measurements of the $\Upsilon(1\mathbf{S})$, $\Upsilon(2\mathbf{S})$, and $\Upsilon(3\mathbf{S})$ differential cross sections in pp collisions at $\sqrt{s}=7$ TeV, Phys. Lett. B749 (2015) 14–34, doi:10.1016/j.physletb.2015.07.037, arXiv:1501.07750 [hep-ex]
- V. Khachatryan et al., Search for a charged Higgs boson in pp collisions at $\sqrt{s}=8$ TeV, *JHEP*. 11 (2015) 018, doi:10.1007/JHEP11(2015)018, arXiv:1508.07774 [hep-ex]
- V. Khachatryan et al., Study of Final-State Radiation in Decays of Z Bosons Produced in pp Collisions at 7 TeV, Phys. Rev. D91 (2015) 092012, doi:10.1103/PhysRevD.91.092012, arXiv:1502.07940 [hep-ex]
- V. Khachatryan et al., Measurement of J/ and (2S) Prompt Double-Differential Cross Sections in pp Collisions at \sqrt{s} =7 TeV, Phys. Rev. Lett. 114 (2015) 191802, doi:10.1103/PhysRevLett.114.191802, arXiv:1502.04155 [hep-ex]
- V. Khachatryan et al., Search for resonances and quantum black holes using dijet mass spectra in proton-proton collisions at $\sqrt{s}=8$ TeV, Phys. Rev. D91 (2015) 052009, doi:10.1103/PhysRevD.91.052009, arXiv:1501.04198 [hep-ex]
- V. Khachatryan et al., **Search for neutral MSSM Higgs bosons decaying into a pair of bottom quarks**, *JHEP*. 11 (2015) 071, doi:10.1007/JHEP11(2015)071, arXiv:1506.08329 [hep-ex]
- V. Khachatryan et al., Search for a light charged Higgs boson decaying to $c\bar{s}$ in pp collisions at $\sqrt{s}=8$ TeV, *JHEP*. 12 (2015) 178, doi:10.1007/JHEP12(2015)178, arXiv:1510.04252 [hep-ex]
- V. Khachatryan et al., Searches for Supersymmetry using the M $_{T2}$ Variable in Hadronic Events Produced in pp Collisions at 8 TeV, JHEP.~05 (2015) 078, doi:10.1007/JHEP05(2015)078, arXiv:1502.04358 [hep-ex]
- V. Khachatryan et al., Evidence for Collective Multiparticle Correlations in p-Pb Collisions, *Phys. Rev. Lett.* 115 (2015) 012301, doi:10.1103/PhysRevLett.115.012301, arXiv:1502.05382 [nucl-ex]
- V. Khachatryan et al., Searches for third-generation squark production in fully hadronic final states in proton-proton collisions at $\sqrt{s}=8$ TeV, *JHEP*. 06 (2015) 116, doi:10.1007/JHEP06(2015)116, arXiv:1503.08037 [hep-ex]
- V. Khachatryan et al., Nuclear Effects on the Transverse Momentum Spectra of Charged Particles in pPb Collisions at $\sqrt{s_{_{\rm NN}}} = 5.02$ TeV, Eur. Phys. J. C75 (2015) 237, doi:10.1140/epjc/s10052-015-3435-4, arXiv:1502.05387 [nucl-ex]
- V. Khachatryan et al., Distributions of Topological Observables in Inclusive Three- and Four-Jet Events in pp Collisions at sqrt(s) =

- **7 TeV**, Eur. Phys. J. C75 (2015) 302, doi:10.1140/epjc/s10052-015-3491-9, arXiv:1502.04785 [hep-ex]
- V. Khachatryan et al., Search for resonant pair production of Higgs bosons decaying to two bottom quark-antiquark pairs in proton-proton collisions at 8 TeV, Phys. Lett. B749 (2015) 560–582, doi:10.1016/j.physletb.2015.08.047, arXiv:1503.04114 [hep-ex]
- V. Khachatryan et al., Search for Narrow High-Mass Resonances in Proton–Proton Collisions at $\sqrt{s}=8$ TeV Decaying to a Z and a Higgs Boson, *Phys. Lett.* B748 (2015) 255–277, doi:10.1016/j.physletb.2015.07.011, arXiv:1502.04994 [hep-ex]
- S. Chatrchyan et al., **Search for baryon number violation in top-quark decays**, *Phys. Lett.* B731 (2014) 173–196, doi:10.1016/j.physletb.2014.02.033, arXiv:1310.1618 [hep-ex]
- S. Chatrchyan et al., **Observation of a peaking structure in the** $J/\psi\phi$ mass spectrum from $B^{\pm} \rightarrow J/\psi\phi K^{\pm}$ decays, *Phys. Lett.* B734 (2014) 261–281, doi:10.1016/j.physletb.2014.05.055, arXiv:1309.6920 [hep-ex]
- S. Chatrchyan et al., Evidence of b-Jet Quenching in PbPb Collisions at $\sqrt{s_{NN}}=2.76$ TeV, Phys. Rev. Lett. 113 (2014) 132301, doi:10.1103/PhysRevLett.115.029903, arXiv:1312.4198 [nucl-ex]
- S. Chatrchyan et al., Study of the production of charged pions, kaons, and protons in pPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, Eur. Phys. J. C74 (2014) 2847, doi:10.1140/epjc/s10052-014-2847-x, arXiv:1307.3442 [hep-ex]
- S. Chatrchyan et al., Measurement of the $t\bar{t}$ production cross section in the dilepton channel in pp collisions at $\sqrt{s}=8$ TeV, JHEP.~02~(2014)~024,~doi:10.1007/JHEP02(2014)024,~arXiv:1312.7582~[hep-ex]
- S. Chatrchyan et al., Probing color coherence effects in pp collisions at $\sqrt{s}=7 \, \mathrm{TeV}$, Eur. Phys. J. C74 (2014) 2901, doi:10.1140/epjc/s10052-014-2901-8, arXiv:1311.5815 [hep-ex]
- S. Chatrchyan et al., Measurement of the $W\gamma$ and $Z\gamma$ inclusive cross sections in pp collisions at $\sqrt{s}=7$ TeV and limits on anomalous triple gauge boson couplings, Phys. Rev. D89 (2014) 092005, doi:10.1103/PhysRevD.89.092005, arXiv:1308.6832 [hep-ex]
- S. Chatrchyan et al., Searches for light- and heavy-flavour three-jet resonances in pp collisions at $\sqrt{s}=8$ TeV, Phys. Lett. B730 (2014) 193–214, doi:10.1016/j.physletb.2014.01.049, arXiv:1311.1799 [hep-ex]
- S. Chatrchyan et al., **Determination of the top-quark pole mass and strong coupling constant from the t t-bar production cross section in pp collisions at \sqrt{s}=7 TeV,** *Phys. Lett.* **B728 (2014) 496–517, doi:10.1016/j.physletb.2014.08.040, arXiv:1307.1907 [hep-ex]**
- S. Chatrchyan et al., Search for supersymmetry in pp collisions at \sqrt{s} =8

- TeV in events with a single lepton, large jet multiplicity, and multiple b jets, *Phys. Lett.* B733 (2014) 328–353, doi:10.1016/j.physletb.2014.04.023, arXiv:1311.4937 [hep-ex]
- S. Chatrchyan et al., Measurement of higher-order harmonic azimuthal anisotropy in PbPb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, Phys. Rev. C89 (2014) 044906, doi:10.1103/PhysRevC.89.044906, arXiv:1310.8651 [nucl-ex]
- S. Chatrchyan et al., Modification of jet shapes in PbPb collisions at $\sqrt{s_{NN}}=2.76$ TeV, Phys. Lett. B730 (2014) 243–263, doi:10.1016/j.physletb.2014.01.042, arXiv:1310.0878 [nucl-ex]
- S. Chatrchyan et al., Measurement of the triple-differential cross section for photon+jets production in proton-proton collisions at \sqrt{s} =7 TeV, *JHEP*. 06 (2014) 009, doi:10.1007/JHEP06(2014)009, arXiv:1311.6141 [hep-ex]
- S. Chatrchyan et al., Measurement of the properties of a Higgs boson in the four-lepton final state, *Phys. Rev.* D89 (2014) 092007, doi:10.1103/PhysRevD.89.092007, arXiv:1312.5353 [hep-ex]
- S. Chatrchyan et al., **Search for top squark and higgsino production using diphoton Higgs boson decays**, *Phys. Rev. Lett.* 112 (2014) 161802, doi:10.1103/PhysRevLett.112.161802, arXiv:1312.3310 [hep-ex]
- S. Chatrchyan et al., Measurement of the muon charge asymmetry in inclusive $pp \to W + X$ production at $\sqrt{s} = 7$ TeV and an improved determination of light parton distribution functions, *Phys. Rev.* D90 (2014) 032004, doi:10.1103/PhysRevD.90.032004, arXiv:1312.6283 [hep-ex]
- S. Chatrchyan et al., Search for Flavor-Changing Neutral Currents in Top-Quark Decays $t \to Zq$ in pp Collisions at $\sqrt{s} = 8$ TeV, Phys. Rev. Lett. 112 (2014) 171802, doi:10.1103/PhysRevLett.112.171802, arXiv:1312.4194 [hep-ex]
- S. Chatrchyan et al., Event activity dependence of Y(nS) production in $\sqrt{s_{NN}}$ =5.02 TeV pPb and \sqrt{s} =2.76 TeV pp collisions, *JHEP*. 04 (2014) 103, doi:10.1007/JHEP04(2014)103, arXiv:1312.6300 [nucl-ex]
- S. Chatrchyan et al., Search for pair production of excited top quarks in the lepton + jets final state, *JHEP*. 06 (2014) 125, doi:10.1007/JHEP06(2014)125, arXiv:1311.5357 [hep-ex]
- S. Chatrchyan et al., Measurement of four-jet production in proton-proton collisions at $\sqrt{s}=7$ TeV, Phys. Rev. D89 (2014) 092010, doi:10.1103/PhysRevD.89.092010, arXiv:1312.6440 [hep-ex]
- S. Chatrchyan et al., Measurement of associated W + charm production in pp collisions at $\sqrt{s} = 7$ TeV, *JHEP*. 02 (2014) 013, doi:10.1007/JHEP02(2014)013, arXiv:1310.1138 [hep-ex]
- S. Chatrchyan et al., Inclusive search for a vector-like T quark with charge $\frac{2}{3}$ in pp collisions at $\sqrt{s} = 8$ TeV, Phys. Lett. B729 (2014) 149–171,

- doi:10.1016/j.physletb.2014.01.006, arXiv:1311.7667 [hep-ex]
- S. Chatrchyan et al., Measurement of the production cross section for a W boson and two b jets in pp collisions at \sqrt{s} =7 TeV, *Phys. Lett.* B735 (2014) 204–225, doi:10.1016/j.physletb.2014.06.041, arXiv:1312.6608 [hep-ex]
- S. Chatrchyan et al., Search for top-quark partners with charge 5/3 in the same-sign dilepton final state, *Phys. Rev. Lett.* 112 (2014) 171801, doi:10.1103/PhysRevLett.112.171801, arXiv:1312.2391 [hep-ex]
- S. Chatrchyan et al., Study of double parton scattering using W + 2-jet events in proton-proton collisions at $\sqrt{s} = 7$ TeV, *JHEP*. 03 (2014) 032, doi:10.1007/JHEP03(2014)032, arXiv:1312.5729 [hep-ex]
- S. Chatrchyan et al., Measurement of the top-quark mass in all-jets $t\bar{t}$ events in pp collisions at \sqrt{s} =7 TeV, Eur. Phys. J. C74 (2014) 2758, doi:10.1140/epjc/s10052-014-2758-x, arXiv:1307.4617 [hep-ex]
- S. Chatrchyan et al., Search for the standard model Higgs boson produced in association with a W or a Z boson and decaying to bottom quarks, *Phys. Rev.* D89 (2014) 012003, doi:10.1103/PhysRevD.89.012003, arXiv:1310.3687 [hep-ex]
- S. Chatrchyan et al., Search for anomalous production of events with three or more leptons in pp collisions at $\sqrt(s) = 8$ TeV, Phys. Rev. D90 (2014) 032006, doi:10.1103/PhysRevD.90.032006, arXiv:1404.5801 [hep-ex]
- S. Chatrchyan et al., Measurement of WZ and ZZ production in pp collisions at $\sqrt{s} = 8$ TeV in final states with b-tagged jets, Eur. Phys. J. C74 (2014) 2973, doi:10.1140/epjc/s10052-014-2973-5, arXiv:1403.3047 [hep-ex]
- S. Chatrchyan et al., Measurement of jet fragmentation in PbPb and pp collisions at $\sqrt{s_{NN}}=2.76$ TeV, Phys. Rev. C90 (2014) 024908, doi:10.1103/PhysRevC.90.024908, arXiv:1406.0932 [nucl-ex]
- S. Chatrchyan et al., Search for $WW\gamma$ and $WZ\gamma$ production and constraints on anomalous quartic gauge couplings in pp collisions at $\sqrt{s} =$ 8 TeV, Phys. Rev. D90 (2014) 032008, doi:10.1103/PhysRevD.90.032008, arXiv:1404.4619 [hep-ex]
- S. Chatrchyan et al., Measurement of the production cross sections for a Z boson and one or more b jets in pp collisions at sqrt(s) = 7 TeV, *JHEP*. 06 (2014) 120, doi:10.1007/JHEP06(2014)120, arXiv:1402.1521 [hep-ex]
- S. Chatrchyan et al., **Description and performance of track and primary-vertex reconstruction with the CMS tracker**, *JINST.* 9 (2014) P10009, doi:10.1088/1748-0221/9/10/P10009, arXiv:1405.6569 [physics.ins-det]
- S. Chatrchyan et al., Measurement of differential cross sections for the production of a pair of isolated photons in pp collisions at \sqrt{s}

- 7TeV, Eur. Phys. J. C74 (2014) 3129, doi:10.1140/epjc/s10052-014-3129-3, arXiv:1405.7225 [hep-ex]
- S. Chatrchyan et al., Measurement of the ratio of inclusive jet cross sections using the anti- k_T algorithm with radius parameters R=0.5 and 0.7 in pp collisions at $\sqrt{s} = 7$ TeV, Phys. Rev. D90 (2014) 072006, doi:10.1103/PhysRevD.90.072006, arXiv:1406.0324 [hep-ex]
- S. Chatrchyan et al., Search for supersymmetry with razor variables in pp collisions at \sqrt{s} =7 TeV, Phys. Rev. D90 (2014) 112001, doi:10.1103/PhysRevD.90.112001, arXiv:1405.3961 [hep-ex]
- S. Chatrchyan et al., Studies of dijet transverse momentum balance and pseudorapidity distributions in pPb collisions at $\sqrt{s_{\rm NN}}=5.02{\rm TeV}$, Eur. Phys. J. C74 (2014) 2951, doi:10.1140/epjc/s10052-014-2951-y, arXiv:1401.4433 [nucl-ex]
- S. Chatrchyan et al., Search for W' \rightarrow tb decays in the lepton + jets final state in pp collisions at $\sqrt{s} = 8$ TeV, JHEP. 05 (2014) 108, doi:10.1007/JHEP05(2014)108, arXiv:1402.2176 [hep-ex]
- S. Chatrchyan et al., Search for new physics in the multijet and missing transverse momentum final state in proton-proton collisions at \sqrt{s} = 8 TeV, *JHEP*. 06 (2014) 055, doi:10.1007/JHEP06(2014)055, arXiv:1402.4770 [hep-ex]
- S. Chatrchyan et al., Measurement of inclusive W and Z boson production cross sections in pp collisions at $\sqrt{s} = 8$ TeV, *Phys. Rev. Lett.* 112 (2014) 191802, doi:10.1103/PhysRevLett.112.191802, arXiv:1402.0923 [hep-ex]
- S. Chatrchyan et al., Evidence for the 125 GeV Higgs boson decaying to a pair of τ leptons, *JHEP*. 05 (2014) 104, doi:10.1007/JHEP05(2014)104, arXiv:1401.5041 [hep-ex]
- S. Chatrchyan et al., Measurement of pseudorapidity distributions of charged particles in proton-proton collisions at $\sqrt{s} = 8$ TeV by the CMS and TOTEM experiments, Eur. Phys. J. C74 (2014) 3053, doi:10.1140/epjc/s10052-014-3053-6, arXiv:1405.0722 [hep-ex]
- S. Chatrchyan et al., Search for invisible decays of Higgs bosons in the vector boson fusion and associated ZH production modes, Eur. Phys. J. C74 (2014) 2980, doi:10.1140/epjc/s10052-014-2980-6, arXiv:1404.1344 [hep-ex]
- S. Chatrchyan et al., **Observation of the associated production of a single top quark and a** W **boson in** pp **collisions at** $\sqrt{s} = 8$ **TeV**, Phys. Rev. Lett. 112 (2014) 231802, doi:10.1103/PhysRevLett.112.231802, arXiv:1401.2942 [hep-ex]
- S. Chatrchyan et al., Evidence for the direct decay of the 125 GeV Higgs boson to fermions, *Nature Phys.* 10 (2014) 557–560, doi:10.1038/nphys3005,

- arXiv:1401.6527 [hep-ex]
- S. Chatrchyan et al., **Alignment of the CMS tracker with LHC and cosmic ray data**, *JINST*. 9 (2014) P06009, doi:10.1088/1748-0221/9/06/P06009, arXiv:1403.2286 [physics.ins-det]
- S. Chatrchyan et al., Studies of azimuthal dihadron correlations in ultra-central PbPb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, *JHEP*. 02 (2014) 088, doi:10.1007/JHEP02(2014)088, arXiv:1312.1845 [nucl-ex]
- V. Khachatryan et al., Search for excited quarks in the γ +jet final state in proton–proton collisions at $\sqrt{s}=8$ TeV, Phys. Lett. B738 (2014) 274–293, doi:10.1016/j.physletb.2014.09.048, arXiv:1406.5171 [hep-ex]
- V. Khachatryan et al., **Measurement of Prompt** $\psi(2S) \to J/\psi$ **Yield Ratios in Pb-Pb and** p-p **Collisions at** $\sqrt{s_{NN}} = \mathbf{2.76}$ **TeV**, *Phys. Rev. Lett.* 113 (2014) 262301, doi:10.1103/PhysRevLett.113.262301, arXiv:1410.1804 [nucl-ex]
- V. Khachatryan et al., Search for heavy neutrinos and W bosons with right-handed couplings in proton-proton collisions at $\sqrt{s} = 8 \text{TeV}$, Eur. Phys. J. C74 (2014) 3149, doi:10.1140/epjc/s10052-014-3149-z, arXiv:1407.3683 [hep-ex]
- V. Khachatryan et al., Search for massive resonances decaying into pairs of boosted bosons in semi-leptonic final states at $\sqrt{s} = 8$ TeV, JHEP. 08 (2014) 174, doi:10.1007/JHEP08(2014)174, arXiv:1405.3447 [hep-ex]
- V. Khachatryan et al., Search for massive resonances in dijet systems containing jets tagged as W or Z boson decays in pp collisions at \sqrt{s} = 8 TeV, *JHEP*. 08 (2014) 173, doi:10.1007/JHEP08(2014)173, arXiv:1405.1994 [hep-ex]
- V. Khachatryan et al., Constraints on the Higgs boson width from off-shell production and decay to Z-boson pairs, *Phys. Lett.* B736 (2014) 64–85, doi:10.1016/j.physletb.2014.06.077, arXiv:1405.3455 [hep-ex]
- V. Khachatryan et al., **Measurement of prompt** J/ψ **pair production in pp collisions at** $\sqrt{s} = 7$ **Tev**, JHEP. 09 (2014) 094, doi:10.1007/JHEP09(2014)094, arXiv:1406.0484 [hep-ex]
- V. Khachatryan et al., Study of hadronic event-shape variables in multijet final states in pp collisions at sqrt(s) = 7 TeV, *JHEP*. 10 (2014) 87, doi:10.1007/JHEP10(2014)087, arXiv:1407.2856 [hep-ex]
- V. Khachatryan et al., **Observation of the diphoton decay of the Higgs boson and measurement of its properties**, Eur. Phys. J. C74 (2014) 3076, doi:10.1140/epjc/s10052-014-3076-z, arXiv:1407.0558 [hep-ex]
- V. Khachatryan et al., Measurement of the t-channel single-topquark production cross section and of the $\mid V_{tb} \mid$ CKM matrix element in pp collisions at $\sqrt{s}=$ 8 TeV, JHEP. 06 (2014) 090,

- doi:10.1007/JHEP06(2014)090, arXiv:1403.7366 [hep-ex]
- V. Khachatryan et al., Searches for heavy Higgs bosons in two-Higgs-doublet models and for $t \to ch$ decay using multilepton and diphoton final states in pp collisions at 8 TeV, Phys. Rev. D90 (2014) 112013, doi:10.1103/PhysRevD.90.112013, arXiv:1410.2751 [hep-ex]
- V. Khachatryan et al., Measurement of the $t\bar{t}$ production cross section in pp collisions at $\sqrt{s}=8$ TeV in dilepton final states containing one τ lepton, Phys. Lett. B739 (2014) 23–43, doi:10.1016/j.physletb.2014.10.032, arXiv:1407.6643 [hep-ex]
- V. Khachatryan et al., **Measurement of the ratio** $\mathcal{B}(t \to Wb)/\mathcal{B}(t \to Wq)$ in pp collisions at $\sqrt{s} = 8$ TeV, Phys. Lett. B736 (2014) 33–57, doi:10.1016/j.physletb.2014.06.076, arXiv:1404.2292 [hep-ex]
- V. Khachatryan et al., **Search for the associated production of the Higgs boson with a top-quark pair**, *JHEP*. 09 (2014) 087, doi:10.1007/JHEP09(2014)087, arXiv:1408.1682 [hep-ex]
- V. Khachatryan et al., Search for Standard Model Production of Four Top Quarks in the Lepton + Jets Channel in pp Collisions at \sqrt{s} = 8 TeV, *JHEP*. 11 (2014) 154, doi:10.1007/JHEP11(2014)154, arXiv:1409.7339 [hep-ex]
- V. Khachatryan et al., Search for pair production of third-generation scalar leptoquarks and top squarks in proton–proton collisions at \sqrt{s} =8 TeV, *Phys. Lett.* B739 (2014) 229–249, doi:10.1016/j.physletb.2014.10.063, arXiv:1408.0806 [hep-ex]
- V. Khachatryan et al., Search for jet extinction in the inclusive jet- p_t spectrum from proton-proton collisions at $\sqrt{s}=8$ TeV, Phys. Rev. D90 (2014) 032005, doi:10.1103/PhysRevD.90.032005, arXiv:1405.7653 [hep-ex]
- V. Khachatryan et al., **Identification techniques for highly boosted W** bosons that decay into hadrons, *JHEP*. 12 (2014) 017, doi:10.1007/JHEP12(2014)017, arXiv:1410.4227 [hep-ex]
- V. Khachatryan et al., Search for neutral MSSM Higgs bosons decaying to a pair of tau leptons in pp collisions, *JHEP*. 10 (2014) 160, doi:10.1007/JHEP10(2014)160, arXiv:1408.3316 [hep-ex]
- S. Chatrchyan et al., Search for heavy quarks decaying into a top quark and a W or Z boson using lepton + jets events in pp collisions at \sqrt{s} = 7 TeV, JHEP. 01 (2013) 154, doi:10.1007/JHEP01(2013)154, arXiv:1210.7471 [hep-ex]
- S. Chatrchyan et al., Search for a narrow spin-2 resonance decaying to a pair of Z vector bosons in the semileptonic final state, *Phys. Lett.* B718 (2013) 1208–1228, doi:10.1016/j.physletb.2012.11.063, arXiv:1209.3807 [hep-ex]
- S. Chatrchyan et al., Search for new physics in events with photons, jets,

- and missing transverse energy in pp collisions at $\sqrt{s} = 7$ TeV, JHEP. 03 (2013) 111, doi:10.1007/JHEP03(2013)111, arXiv:1211.4784 [hep-ex]
- S. Chatrchyan et al., Measurement of the sum of WW and WZ production with W+dijet events in pp collisions at $\sqrt{s}=7$ TeV, Eur. Phys. J. C73 (2013) 2283, doi:10.1140/epjc/s10052-013-2283-3, arXiv:1210.7544 [hep-ex]
- S. Chatrchyan et al., Measurements of differential jet cross sections in proton-proton collisions at $\sqrt{s}=7$ TeV with the CMS detector, *Phys. Rev.* D87 (2013) 112002, doi:10.1103/PhysRevD.87.112002, arXiv:1212.6660 [hep-ex]
- S. Chatrchyan et al., Search for a non-standard-model Higgs boson decaying to a pair of new light bosons in four-muon final states, *Phys. Lett.* B726 (2013) 564–586, doi:10.1016/j.physletb.2013.09.009, arXiv:1210.7619 [hep-ex]
- S. Chatrchyan et al., Forward-backward asymmetry of Drell-Yan lepton pairs in pp collisions at $\sqrt{s} = 7$ TeV, Phys. Lett. B718 (2013) 752–772, doi:10.1016/j.physletb.2012.10.082, arXiv:1207.3973 [hep-ex]
- S. Chatrchyan et al., Search for a W 'boson decaying to a bottom quark and a top quark in pp collisions at $\sqrt{s}=7$ TeV, $Phys.\ Lett.\ B718\ (2013)$ 1229–1251, doi:10.1016/j.physletb.2012.12.008, arXiv:1208.0956 [hep-ex]
- S. Chatrchyan et al., Studies of jet quenching using isolated-photon+jet correlations in PbPb and pp collisions at $\sqrt{s_{NN}}=2.76$ TeV, Phys. Lett. B718 (2013) 773–794, doi:10.1016/j.physletb.2012.11.003, arXiv:1205.0206 [nuclex]
- S. Chatrchyan et al., Search for contact interactions in $\mu^+\mu^-$ events in pp collisions at $\sqrt{s}=7$ TeV, Phys. Rev. D87 (2013) 032001, doi:10.1103/PhysRevD.87.032001, arXiv:1212.4563 [hep-ex]
- S. Chatrchyan et al., **Search for flavor changing neutral currents in top quark decays in pp collisions at 7 TeV**, *Phys. Lett.* B718 (2013) 1252–1272, doi:10.1016/j.physletb.2012.12.045, arXiv:1208.0957 [hep-ex]
- S. Chatrchyan et al., Study of the Mass and Spin-Parity of the Higgs Boson Candidate Via Its Decays to Z Boson Pairs, *Phys. Rev. Lett.* 110 (2013) 081803, doi:10.1103/PhysRevLett.110.081803, arXiv:1212.6639 [hep-ex]
- S. Chatrchyan et al., **Search in leptonic channels for heavy resonances decaying to long-lived neutral particles**, *JHEP*. 02 (2013) 085, doi:10.1007/JHEP02(2013)085, arXiv:1211.2472 [hep-ex]
- S. Chatrchyan et al., **Identification of b-quark jets with the CMS experiment**, *JINST*. 8 (2013) P04013, doi:10.1088/1748-0221/8/04/P04013, arXiv:1211.4462 [hep-ex]
- S. Chatrchyan et al., Search for long-lived particles decaying to photons and missing energy in proton-proton collisions at $\sqrt{s} = 7$ TeV, Phys.

- $Lett. \ B722 \ (2013) \ 273-294, \ doi:10.1016/j.physletb.2013.04.027, \ arXiv:1212.1838 \\ [hep-ex]$
- S. Chatrchyan et al., **Measurement of the inelastic proton-proton cross section at** $\sqrt{s}=7$ **TeV**, *Phys. Lett.* B722 (2013) 5–27, doi:10.1016/j.physletb.2013.03.024, arXiv:1210.6718 [hep-ex]
- S. Chatrchyan et al., Search for heavy narrow dilepton resonances in pp collisions at $\sqrt{s} = 7$ TeV and $\sqrt{s} = 8$ TeV, $Phys.\ Lett.\ B720\ (2013)\ 63-82$, doi:10.1016/j.physletb.2013.02.003, arXiv:1212.6175 [hep-ex]
- S. Chatrchyan et al., Measurement of the $t\bar{t}$ production cross section in pp collisions at $\sqrt{s}=7$ TeV with lepton + jets final states, Phys. Lett. B720 (2013) 83–104, doi:10.1016/j.physletb.2013.02.021, arXiv:1212.6682 [hep-ex]
- S. Chatrchyan et al., Search for exotic resonances decaying into WZ/ZZ in pp collisions at $\sqrt{s}=7$ TeV, JHEP. 02 (2013) 036, doi:10.1007/JHEP02(2013)036, arXiv:1211.5779 [hep-ex]
- S. Chatrchyan et al., Measurement of differential top-quark pair production cross sections in pp colisions at $\sqrt{s} = 7$ TeV, Eur. Phys. J. C73 (2013) 2339, doi:10.1140/epjc/s10052-013-2339-4, arXiv:1211.2220 [hep-ex]
- S. Chatrchyan et al., Search for supersymmetry in final states with a single lepton, b-quark jets, and missing transverse energy in proton-proton collisions at $\sqrt{s}=7$ TeV, Phys. Rev. D87 (2013) 052006, doi:10.1103/PhysRevD.87.052006, arXiv:1211.3143 [hep-ex]
- S. Chatrchyan et al., Measurement of the ZZ production cross section and search for anomalous couplings in 2 l2l ' final states in pp collisions at $\sqrt{s}=7$ TeV, JHEP. 01 (2013) 063, doi:10.1007/JHEP01(2013)063, arXiv:1211.4890 [hep-ex]
- S. Chatrchyan et al., Search for pair production of third-generation leptoquarks and top squarks in pp collisions at $\sqrt{s}=7$ TeV, Phys. Rev. Lett. 110 (2013) 081801, doi:10.1103/PhysRevLett.110.081801, arXiv:1210.5629 [hep-ex]
- S. Chatrchyan et al., Measurement of the elliptic anisotropy of charged particles produced in PbPb collisions at $\sqrt{s_{NN}}$ =2.76 TeV, Phys. Rev. C87 (2013) 014902, doi:10.1103/PhysRevC.87.014902, arXiv:1204.1409 [nucl-ex]
- S. Chatrchyan et al., Search for anomalous production of highly boosted Z bosons decaying to $\mu^+\mu^-$ in proton-proton collisions at $\sqrt{s}=7$ TeV, Phys. Lett. B722 (2013) 28–47, doi:10.1016/j.physletb.2013.03.037, arXiv:1210.0867 [hep-ex]
- S. Chatrchyan et al., Inclusive search for supersymmetry using the razor variables in pp collisions at $\sqrt{s} = 7$ TeV, Phys. Rev. Lett. 111 (2013) 081802,

- doi:10.1103/PhysRevLett.111.081802, arXiv:1212.6961 [hep-ex]
- S. Chatrchyan et al., **Observation of a diffractive contribution to dijet** production in proton-proton collisions at $\sqrt{s} = 7$ TeV, *Phys. Rev.* D87 (2013) 012006, doi:10.1103/PhysRevD.87.012006, arXiv:1209.1805 [hep-ex]
- S. Chatrchyan et al., Measurement of the azimuthal anisotropy of neutral pions in PbPb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, Phys. Rev. Lett. 110 (2013) 042301, doi:10.1103/PhysRevLett.110.042301, arXiv:1208.2470 [nucl-ex]
- S. Chatrchyan et al., Search for supersymmetry in final states with missing transverse energy and 0, 1, 2, or at least 3 b-quark jets in 7 TeV pp collisions using the variable alphaT, *JHEP*. 01 (2013) 077, doi:10.1007/JHEP01(2013)077, arXiv:1210.8115 [hep-ex]
- S. Chatrchyan et al., Measurement of the Y(1S), Y(2S) and Y(3S) polarizations in pp collisions at $\sqrt{s}=7$ TeV, Phys. Rev. Lett. 110 (2013) 081802, doi:10.1103/PhysRevLett.110.081802, arXiv:1209.2922 [hep-ex]
- S. Chatrchyan et al., Search for Z 'resonances decaying to $t\bar{t}$ in dilepton+jets final states in pp collisions at $\sqrt{s}=7$ TeV, Phys. Rev. D87 (2013) 072002, doi:10.1103/PhysRevD.87.072002, arXiv:1211.3338 [hep-ex]
- S. Chatrchyan et al., Search for heavy resonances in the W/Z-tagged dijet mass spectrum in pp collisions at 7 TeV, *Phys. Lett.* B723 (2013) 280–301, doi:10.1016/j.physletb.2013.05.040, arXiv:1212.1910 [hep-ex]
- S. Chatrchyan et al., Search for supersymmetry in pp collisions at $\sqrt{s} = 7$ TeV in events with a single lepton, jets, and missing transverse momentum, Eur. Phys. J. C73 (2013) 2404, doi:10.1140/epjc/s10052-013-2404-z, arXiv:1212.6428 [hep-ex]
- S. Chatrchyan et al., Evidence for associated production of a single top quark and W boson in pp collisions at $\sqrt{s} = 7$ TeV, Phys. Rev. Lett. 110 (2013) 022003, doi:10.1103/PhysRevLett.110.022003, arXiv:1209.3489 [hep-ex]
- S. Chatrchyan et al., Study of exclusive two-photon production of W^+W^- in pp collisions at $\sqrt{s}=7$ TeV and constraints on anomalous quartic gauge couplings, JHEP.~07~(2013)~116,~doi:10.1007/JHEP07(2013)116,~arXiv:1305.5596~[hep-ex]
- S. Chatrchyan et al., Jet and underlying event properties as a function of charged-particle multiplicity in proton–proton collisions at $\sqrt{s}=$ 7 TeV, Eur. Phys. J. C73 (2013) 2674, doi:10.1140/epjc/s10052-013-2674-5, arXiv:1310.4554 [hep-ex]
- S. Chatrchyan et al., Measurement of the B(s) to mu+ mu- branching fraction and search for B0 to mu+ mu- with the CMS Experiment, *Phys. Rev. Lett.* 111 (2013) 101804, doi:10.1103/PhysRevLett.111.101804, arXiv:1307.5025 [hep-ex]
- S. Chatrchyan et al., Measurement of masses in the $t\bar{t}$ system by kine-

- matic endpoints in pp collisions at $\sqrt{s} = 7$ TeV, Eur. Phys. J. C73 (2013) 2494, doi:10.1140/epjc/s10052-013-2494-7, arXiv:1304.5783 [hep-ex]
- S. Chatrchyan et al., Angular analysis and branching fraction measurement of the decay $B^0 \to K^{*0} \mu^+ \mu^-$, Phys. Lett. B727 (2013) 77–100, doi:10.1016/j.physletb.2013.10.017, arXiv:1308.3409 [hep-ex]
- S. Chatrchyan et al., Measurement of the prompt J/ψ and $\psi(2S)$ polarizations in pp collisions at $\sqrt{s} = 7$ TeV, Phys. Lett. B727 (2013) 381–402, doi:10.1016/j.physletb.2013.10.055, arXiv:1307.6070 [hep-ex]
- S. Chatrchyan et al., Measurement of the X(3872) production cross section via decays to J/psi pi pi in pp collisions at sqrt(s) = 7 TeV, JHEP. 04 (2013) 154, doi:10.1007/JHEP04(2013)154, arXiv:1302.3968 [hep-ex]
- S. Chatrchyan et al., Energy Calibration and Resolution of the CMS Electromagnetic Calorimeter in pp Collisions at $\sqrt{s} = 7$ TeV, JINST. 8 (2013) P09009, doi:10.1088/1748-0221/8/09/P09009, arXiv:1306.2016 [hep-ex]
- S. Chatrchyan et al., Search for physics beyond the standard model in events with τ leptons, jets, and large transverse momentum imbalance in pp collisions at $\sqrt{s}=7$ TeV, Eur. Phys. J. C73 (2013) 2493, doi:10.1140/epjc/s10052-013-2493-8, arXiv:1301.3792 [hep-ex]
- S. Chatrchyan et al., Study of the underlying event at forward rapidity in pp collisions at $\sqrt{s} = 0.9$, 2.76, and 7 TeV, *JHEP*. 04 (2013) 072, doi:10.1007/JHEP04(2013)072, arXiv:1302.2394 [hep-ex]
- S. Chatrchyan et al., Search for pair-produced dijet resonances in four-jet final states in pp collisions at \sqrt{s} =7 TeV, Phys. Rev. Lett. 110 (2013) 141802, doi:10.1103/PhysRevLett.110.141802, arXiv:1302.0531 [hep-ex]
- S. Chatrchyan et al., Measurement of the W-boson helicity in topquark decays from $t\bar{t}$ production in lepton+jets events in pp collisions at $\sqrt{s}=7$ TeV, *JHEP*. 10 (2013) 167, doi:10.1007/JHEP10(2013)167, arXiv:1308.3879 [hep-ex]
- S. Chatrchyan et al., Measurement of the hadronic activity in events with a Z and two jets and extraction of the cross section for the electroweak production of a Z with two jets in pp collisions at \sqrt{s} = 7 TeV, *JHEP*. 10 (2013) 062, doi:10.1007/JHEP10(2013)062, arXiv:1305.7389 [hep-ex]
- S. Chatrchyan et al., Measurement of the top-antitop production cross section in the tau+jets channel in pp collisions at sqrt(s) = 7 TeV, Eur. Phys. J. C73 (2013) 2386, doi:10.1140/epjc/s10052-013-2386-x, arXiv:1301.5755 [hep-ex]
- S. Chatrchyan et al., Search for supersymmetry in events with oppositesign dileptons and missing transverse energy using an artificial neural

- $\bf network,\ Phys.\ Rev.\ D87\ (2013)\ 072001,\ doi:10.1103/PhysRevD.87.072001,\ arXiv:1301.0916\ [hep-ex]$
- S. Chatrchyan et al., Observation of a new boson with mass near 125 GeV in pp collisions at $\sqrt{s} = 7$ and 8 TeV, *JHEP*. 06 (2013) 081, doi:10.1007/JHEP06(2013)081, arXiv:1303.4571 [hep-ex]
- S. Chatrchyan et al., Searches for new physics using the $t\bar{t}$ invariant mass distribution in pp collisions at \sqrt{s} =8 TeV, Phys. Rev. Lett. 111 (2013) 211804, doi:10.1103/PhysRevLett.111.211804, arXiv:1309.2030 [hep-ex]
- S. Chatrchyan et al., **Search for new physics in final states with a lepton** and missing transverse energy in pp collisions at the LHC, *Phys. Rev.* D87 (2013) 072005, doi:10.1103/PhysRevD.87.072005, arXiv:1302.2812 [hep-ex]
- S. Chatrchyan et al., Search for a new bottomonium state decaying to $\Upsilon(1S)\pi^+\pi^-$ in pp collisions at $\sqrt{s}=8$ TeV, Phys. Lett. B727 (2013) 57–76, doi:10.1016/j.physletb.2013.10.016, arXiv:1309.0250 [hep-ex]
- S. Chatrchyan et al., Search for contact interactions using the inclusive jet p_T spectrum in pp collisions at $\sqrt{s} = 7$ TeV, Phys. Rev. D87 (2013) 052017, doi:10.1103/PhysRevD.87.052017, arXiv:1301.5023 [hep-ex]
- S. Chatrchyan et al., Search for supersymmetry in hadronic final states with missing transverse energy using the variables α_T and b-quark multiplicity in pp collisions at $\sqrt{s}=8$ TeV, Eur. Phys. J. C73 (2013) 2568, doi:10.1140/epjc/s10052-013-2568-6, arXiv:1303.2985 [hep-ex]
- S. Chatrchyan et al., Measurement of the production cross section for $Z\gamma \to \nu\bar{\nu}\gamma$ in pp collisions at $\sqrt{s}=7$ TeV and limits on $ZZ\gamma$ and $Z\gamma\gamma$ triple gauge boson couplings, *JHEP*. 10 (2013) 164, doi:10.1007/JHEP10(2013)164, arXiv:1309.1117 [hep-ex]
- S. Chatrchyan et al., **Multiplicity and transverse momentum dependence** of two- and four-particle correlations in pPb and PbPb collisions, *Phys. Lett.* B724 (2013) 213–240, doi:10.1016/j.physletb.2013.06.028, arXiv:1305.0609 [nucl-ex]
- S. Chatrchyan et al., Searches for long-lived charged particles in pp collisions at \sqrt{s} =7 and 8 TeV, *JHEP*. 07 (2013) 122, doi:10.1007/JHEP07(2013)122, arXiv:1305.0491 [hep-ex]
- S. Chatrchyan et al., Rapidity distributions in exclusive **Z** + jet and γ + jet events in pp collisions at \sqrt{s} = 7 TeV, Phys. Rev. D88 (2013) 112009, doi:10.1103/PhysRevD.88.112009, arXiv:1310.3082 [hep-ex]
- S. Chatrchyan et al., Measurement of associated production of vector bosons and top quark-antiquark pairs at sqrt(s) = 7 TeV, *Phys. Rev. Lett.* 110 (2013) 172002, doi:10.1103/PhysRevLett.110.172002, arXiv:1303.3239 [hep-ex]
- S. Chatrchyan et al., Search for a Higgs boson decaying into a

- b-quark pair and produced in association with b quarks in proton–proton collisions at 7 TeV, *Phys. Lett.* B722 (2013) 207–232, doi:10.1016/j.physletb.2013.04.017, arXiv:1302.2892 [hep-ex]
- S. Chatrchyan et al., Search for narrow resonances using the dijet mass spectrum in pp collisions at \sqrt{s} =8 TeV, Phys. Rev. D87 (2013) 114015, doi:10.1103/PhysRevD.87.114015, arXiv:1302.4794 [hep-ex]
- S. Chatrchyan et al., Measurement of neutral strange particle production in the underlying event in proton-proton collisions at sqrt(s) = 7 TeV, Phys. Rev. D88 (2013) 052001, doi:10.1103/PhysRevD.88.052001, arXiv:1305.6016 [hep-ex]
- S. Chatrchyan et al., The performance of the CMS muon detector in proton-proton collisions at sqrt(s) = 7 TeV at the LHC, JINST. 8 (2013) P11002, doi:10.1088/1748-0221/8/11/P11002, arXiv:1306.6905 [physics.ins-det]
- S. Chatrchyan et al., Searches for Higgs bosons in pp collisions at sqrt(s) = 7 and 8 TeV in the context of four-generation and fermiophobic models, *Phys. Lett.* B725 (2013) 36–59, doi:10.1016/j.physletb.2013.06.043, arXiv:1302.1764 [hep-ex]
- S. Chatrchyan et al., Measurement of the Λ_b^0 lifetime in pp collisions at $\sqrt{s}=7$ TeV, JHEP. 07 (2013) 163, doi:10.1007/JHEP07(2013)163, arXiv:1304.7495 [hep-ex]
- S. Chatrchyan et al., **Interpretation of Searches for Supersymmetry with simplified Models**, *Phys. Rev.* D88 (2013) 052017, doi:10.1103/PhysRevD.88.052017, arXiv:1301.2175 [hep-ex]
- S. Chatrchyan et al., Measurement of the differential and double-differential Drell-Yan cross sections in proton-proton collisions at $\sqrt{s}=7$ TeV, $\it JHEP.~12~(2013)~030,~doi:10.1007/JHEP12(2013)030,~arXiv:1310.7291~[hep-ex]$
- S. Chatrchyan et al., Event shapes and azimuthal correlations in Z + jets events in pp collisions at $\sqrt{s} = 7$ TeV, Phys. Lett. B722 (2013) 238–261, doi:10.1016/j.physletb.2013.04.025, arXiv:1301.1646 [hep-ex]
- S. Chatrchyan et al., Measurement of the ratio of the inclusive 3-jet cross section to the inclusive 2-jet cross section in pp collisions at $\sqrt{s}=7$ TeV and first determination of the strong coupling constant in the TeV range, Eur. Phys. J. C73 (2013) 2604, doi:10.1140/epjc/s10052-013-2604-6, arXiv:1304.7498 [hep-ex]
- S. Chatrchyan et al., Measurement of the $t\bar{t}$ production cross section in the all-jet final state in pp collisions at $\sqrt{s}=7$ TeV, *JHEP*. 05 (2013) 065, doi:10.1007/JHEP05(2013)065, arXiv:1302.0508 [hep-ex]
- S. Chatrchyan et al., Search for a Higgs boson decaying into a Z and a photon in pp collisions at sqrt(s) = 7 and 8 TeV, Phys. Lett. B726

- (2013) 587–609, doi:10.1016/j.physletb.2013.09.057, arXiv:1307.5515 [hep-ex]
- S. Chatrchyan et al., Studies of jet mass in dijet and W/Z + jet events, *JHEP*. 05 (2013) 090, doi:10.1007/JHEP05(2013)090, arXiv:1303.4811 [hep-ex]
- S. Chatrchyan et al., **Search for gluino mediated bottom- and top-squark production in multijet final states in pp collisions at 8 TeV**, *Phys. Lett.* B725 (2013) 243–270, doi:10.1016/j.physletb.2013.06.058, arXiv:1305.2390 [hep-ex]
- S. Chatrchyan et al., Search for top squarks in *R*-parity-violating supersymmetry using three or more leptons and b-tagged jets, *Phys. Rev. Lett.* 111 (2013) 221801, doi:10.1103/PhysRevLett.111.221801, arXiv:1306.6643 [hep-ex]
- S. Chatrchyan et al., Search for microscopic black holes in pp collisions at sqrt(s) = 8 TeV, JHEP. 07 (2013) 178, doi:10.1007/JHEP07(2013)178, arXiv:1303.5338 [hep-ex]
- S. Chatrchyan et al., Search for the standard model Higgs boson produced in association with a top-quark pair in pp collisions at the LHC, *JHEP*. 05 (2013) 145, doi:10.1007/JHEP05(2013)145, arXiv:1303.0763 [hep-ex]
- S. Chatrchyan et al., Measurement of the $\Upsilon(1S), \Upsilon(2S)$, and $\Upsilon(3S)$ cross sections in pp collisions at $\sqrt{s} = 7$ TeV, Phys. Lett. B727 (2013) 101–125, doi:10.1016/j.physletb.2013.10.033, arXiv:1303.5900 [hep-ex]
- S. Chatrchyan et al., Search for a standard-model-like Higgs boson with a mass in the range 145 to 1000 GeV at the LHC, Eur. Phys. J. C73 (2013) 2469, doi:10.1140/epjc/s10052-013-2469-8, arXiv:1304.0213 [hep-ex]
- S. Chatrchyan et al., Measurement of the cross section and angular correlations for associated production of a Z boson with b hadrons in pp collisions at $\sqrt{s}=7$ TeV, $\it JHEP.~12~(2013)~039$, doi:10.1007/JHEP12(2013)039, arXiv:1310.1349 [hep-ex]
- S. Chatrchyan et al., Search for excited leptons in pp collisions at $\sqrt{s}=7$ TeV, Phys.~Lett.~B720~(2013)~309–329,~doi:10.1016/j.physletb.2013.02.031,~arXiv:1210.2422~[hep-ex]
- S. Chatrchyan et al., **Observation of long-range near-side angular correlations in proton-lead collisions at the LHC**, *Phys. Lett.* B718 (2013) 795–814, doi:10.1016/j.physletb.2012.11.025, arXiv:1210.5482 [nucl-ex]
- S. Chatrchyan et al., Search for supersymmetry in events with photons and low missing transverse energy in pp collisions at $\sqrt{s}=7$ TeV, Phys. Lett. B719 (2013) 42–61, doi:10.1016/j.physletb.2012.12.055, arXiv:1210.2052 [hep-ex]
- S. Chatrchyan et al., Search for fractionally charged particles in pp collisions at $\sqrt{s} = 7$ TeV, Phys. Rev. D87 (2013) 092008,

- doi:10.1103/PhysRevD.87.092008, arXiv:1210.2311 [hep-ex]
- S. Chatrchyan et al., Search for narrow resonances and quantum black holes in inclusive and b-tagged dijet mass spectra from pp collisions at $\sqrt{s}=7$ TeV, JHEP. 01 (2013) 013, doi:10.1007/JHEP01(2013)013, arXiv:1210.2387 [hep-ex]
- S. Chatrchyan et al., Exclusive photon-photon production of muon pairs in proton-proton collisions at $\sqrt{s} = 7$ TeV, *JHEP*. 01 (2012) 052, doi:10.1007/JHEP01(2012)052, arXiv:1111.5536 [hep-ex]
- S. Chatrchyan et al., Inclusive search for squarks and gluinos in pp collisions at $\sqrt{s}=7$ TeV, Phys. Rev. D85 (2012) 012004, doi:10.1103/PhysRevD.85.012004, arXiv:1107.1279 [hep-ex]
- S. Chatrchyan et al., Search for signatures of extra dimensions in the diphoton mass spectrum at the Large Hadron Collider, *Phys. Rev. Lett.* 108 (2012) 111801, doi:10.1103/PhysRevLett.108.111801, arXiv:1112.0688 [hep-ex]
- S. Chatrchyan et al., Measurement of the charge asymmetry in top-quark pair production in proton-proton collisions at $\sqrt{s}=7$ TeV, Phys. Lett. B709 (2012) 28–49, doi:10.1016/j.physletb.2012.01.078, arXiv:1112.5100 [hep-ex]
- S. Chatrchyan et al., J/ψ and ψ_{2S} production in pp collisions at $\sqrt{s}=7$ TeV, JHEP. 02 (2012) 011, doi:10.1007/JHEP02(2012)011, arXiv:1111.1557 [hep-ex]
- S. Chatrchyan et al., **Jet Production Rates in Association with** W and Z Bosons in pp Collisions at $\sqrt{s}=7$ TeV, JHEP. 01 (2012) 010, doi:10.1007/JHEP01(2012)010, arXiv:1110.3226 [hep-ex]
- S. Chatrchyan et al., Measurement of the Production Cross Section for Pairs of Isolated Photons in pp collisions at $\sqrt{s} = 7$ TeV, JHEP. 01 (2012) 133, doi:10.1007/JHEP01(2012)133, arXiv:1110.6461 [hep-ex]
- S. Chatrchyan et al., Forward Energy Flow, Central Charged-Particle Multiplicities, and Pseudorapidity Gaps in W and Z Boson Events from pp Collisions at $\sqrt{s} = 7$ TeV, Eur. Phys. J. C72 (2012) 1839, doi:10.1140/epjc/s10052-011-1839-3, arXiv:1110.0181 [hep-ex]
- S. Chatrchyan et al., Measurement of the Rapidity and Transverse Momentum Distributions of Z Bosons in pp Collisions at $\sqrt{s}=7$ TeV, Phys. Rev. D85 (2012) 032002, doi:10.1103/PhysRevD.85.032002, arXiv:1110.4973 [hep-ex]
- S. Chatrchyan et al., Search for a light pseudoscalar Higgs boson in the dimuon decay channel in pp collisions at $\sqrt{s} = 7$ TeV, Phys. Rev. Lett. 109 (2012) 121801, doi:10.1103/PhysRevLett.109.121801, arXiv:1206.6326 [hep-ex]
- S. Chatrchyan et al., Search for quark compositeness in dijet angular

- distributions from pp collisions at $\sqrt{s} = 7$ TeV, JHEP. 05 (2012) 055, doi:10.1007/JHEP05(2012)055, arXiv:1202.5535 [hep-ex]
- S. Chatrchyan et al., Measurement of the $t\bar{t}$ production cross section in the dilepton channel in pp collisions at $\sqrt{s} = 7$ TeV, JHEP. 11 (2012) 067, doi:10.1007/JHEP11(2012)067, arXiv:1208.2671 [hep-ex]
- S. Chatrchyan et al., Search for resonant $t\bar{t}$ production in lepton+jets events in pp collisions at $\sqrt{s}=7$ TeV, JHEP. 12 (2012) 015, doi:10.1007/JHEP12(2012)015, arXiv:1209.4397 [hep-ex]
- S. Chatrchyan et al., Inclusive and differential measurements of the $t\bar{t}$ charge asymmetry in proton-proton collisions at $\sqrt{s}=7$ TeV, *Phys. Lett.* B717 (2012) 129–150, doi:10.1016/j.physletb.2012.09.028, arXiv:1207.0065 [hep-ex]
- S. Chatrchyan et al., Measurement of the top-quark mass in $t\bar{t}$ events with lepton+jets final states in pp collisions at $\sqrt{s}=7$ TeV, JHEP. 12 (2012) 105, doi:10.1007/JHEP12(2012)105, arXiv:1209.2319 [hep-ex]
- S. Chatrchyan et al., Search for the standard model Higgs boson in the decay channel H to ZZ to 4 leptons in pp collisions at $\sqrt{s}=7$ TeV, Phys. Rev. Lett. 108 (2012) 111804, doi:10.1103/PhysRevLett.108.111804, arXiv:1202.1997 [hep-ex]
- S. Chatrchyan et al., Inclusive *b*-jet production in *pp* collisions at $\sqrt{s}=7$ TeV, *JHEP*. 04 (2012) 084, doi:10.1007/JHEP04(2012)084, arXiv:1202.4617 [hep-ex]
- S. Chatrchyan et al., Search for stopped long-lived particles produced in pp collisions at $\sqrt{s}=7$ TeV, JHEP. 08 (2012) 026, doi:10.1007/JHEP08(2012)026, arXiv:1207.0106 [hep-ex]
- S. Chatrchyan et al., Measurement of the top-quark mass in $t\bar{t}$ events with dilepton final states in pp collisions at $\sqrt{s}=7$ TeV, Eur. Phys. J. C72 (2012) 2202, doi:10.1140/epjc/s10052-012-2202-z, arXiv:1209.2393 [hep-ex]
- S. Chatrchyan et al., Measurement of the single-top-quark t-channel cross section in pp collisions at $\sqrt{s} = 7$ TeV, JHEP. 12 (2012) 035, doi:10.1007/JHEP12(2012)035, arXiv:1209.4533 [hep-ex]
- S. Chatrchyan et al., Search for heavy Majorana neutrinos in $\mu^{\pm}\mu^{\pm}$ + jets and $e^{\pm}e^{\pm}$ + jets events in pp collisions at $\sqrt{s}=7$ TeV, *Phys. Lett.* B717 (2012) 109–128, doi:10.1016/j.physletb.2012.09.012, arXiv:1207.6079 [hep-ex]
- S. Chatrchyan et al., **Combined search for the quarks of a sequential fourth generation**, *Phys. Rev.* D86 (2012) 112003, doi:10.1103/PhysRevD.86.112003, arXiv:1209.1062 [hep-ex]
- S. Chatrchyan et al., Measurement of jet fragmentation into charged particles in pp and PbPb collisions at $\sqrt{s_{NN}}=2.76$ TeV, $\it{JHEP}.~10$

- (2012) 087, doi:10.1007/JHEP10(2012)087, arXiv:1205.5872 [nucl-ex]
- S. Chatrchyan et al., Measurement of the inclusive production cross sections for forward jets and for dijet events with one forward and one central jet in pp collisions at $\sqrt{s} = 7$ TeV, JHEP. 06 (2012) 036, doi:10.1007/JHEP06(2012)036, arXiv:1202.0704 [hep-ex]
- S. Chatrchyan et al., Search for high-mass resonances decaying into -lepton pairs in pp collisions at $\sqrt{s} = 7$ TeV, *Phys. Lett.* B716 (2012) 82–102, doi:10.1016/j.physletb.2012.07.062, arXiv:1206.1725 [hep-ex]
- S. Chatrchyan et al., Search for the standard model Higgs boson in the H to ZZ to $\ell\ell\tau\tau$ decay channel in pp collisions at $\sqrt{s}=7$ TeV, JHEP.~03 (2012) 081, doi:10.1007/JHEP03(2012)081, arXiv:1202.3617 [hep-ex]
- S. Chatrchyan et al., Measurement of the cross section for production of bb^- bar X, decaying to muons in pp collisions at $\sqrt{s} = 7$ TeV, JHEP. 06 (2012) 110, doi:10.1007/JHEP06(2012)110, arXiv:1203.3458 [hep-ex]
- S. Chatrchyan et al., Search for new physics with long-lived particles decaying to photons and missing energy in pp collisions at $\sqrt{s} = 7$ TeV, JHEP. 11 (2012) 172, doi:10.1007/JHEP11(2012)172, arXiv:1207.0627 [hep-ex]
- S. Chatrchyan et al., Search for narrow resonances in dilepton mass spectra in pp collisions at $\sqrt{s} = 7$ TeV, Phys. Lett. B714 (2012) 158–179, doi:10.1016/j.physletb.2012.06.051, arXiv:1206.1849 [hep-ex]
- S. Chatrchyan et al., Study of the dijet mass spectrum in $pp \rightarrow W+$ jets events at $\sqrt{s}=7$ TeV, Phys. Rev. Lett. 109 (2012) 251801, doi:10.1103/PhysRevLett.109.251801, arXiv:1208.3477 [hep-ex]
- S. Chatrchyan et al., Search for supersymmetry in hadronic final states using MT2 in pp collisions at $\sqrt{s}=7$ TeV, JHEP. 10 (2012) 018, doi:10.1007/JHEP10(2012)018, arXiv:1207.1798 [hep-ex]
- S. Chatrchyan et al., Search for large extra dimensions in dimuon and dielectron events in pp collisions at $\sqrt{s} = 7$ TeV, *Phys. Lett.* B711 (2012) 15–34, doi:10.1016/j.physletb.2012.03.029, arXiv:1202.3827 [hep-ex]
- S. Chatrchyan et al., Search for a W' or Techni- ρ Decaying into WZ in pp Collisions at $\sqrt{s}=7$ TeV, Phys. Rev. Lett. 109 (2012) 141801, doi:10.1103/PhysRevLett.109.141801, arXiv:1206.0433 [hep-ex]
- S. Chatrchyan et al., Search for Anomalous $t\bar{t}$ Production in the Highly-Boosted All-Hadronic Final State, *JHEP*. 09 (2012) 029, doi:10.1007/JHEP09(2012)029, arXiv:1204.2488 [hep-ex]
- S. Chatrchyan et al., Search for new physics in the multijet and missing transverse momentum final state in proton-proton collisions at $\sqrt{s} = 7$ TeV, *Phys. Rev. Lett.* 109 (2012) 171803, doi:10.1103/PhysRevLett.109.171803,

- arXiv:1207.1898 [hep-ex]
- S. Chatrchyan et al., **Observation of sequential Upsilon suppression in PbPb collisions**, *Phys. Rev. Lett.* 109 (2012) 222301, doi:10.1103/PhysRevLett.109.222301, arXiv:1208.2826 [nucl-ex]
- S. Chatrchyan et al., Measurement of the pseudorapidity and centrality dependence of the transverse energy density in PbPb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, Phys. Rev. Lett. 109 (2012) 152303, doi:10.1103/PhysRevLett.109.152303, arXiv:1205.2488 [nucl-ex]
- S. Chatrchyan et al., Search for dark matter and large extra dimensions in monojet events in pp collisions at $\sqrt{s} = 7$ TeV, JHEP. 09 (2012) 094, doi:10.1007/JHEP09(2012)094, arXiv:1206.5663 [hep-ex]
- S. Chatrchyan et al., Search for anomalous production of multilepton events in pp collisions at $\sqrt{s}=7$ TeV, JHEP. 06 (2012) 169, doi:10.1007/JHEP06(2012)169, arXiv:1204.5341 [hep-ex]
- S. Chatrchyan et al., Search for leptonic decays of W ' bosons in pp collisions at $\sqrt{s}=7$ TeV, JHEP. 08 (2012) 023, doi:10.1007/JHEP08(2012)023, arXiv:1204.4764 [hep-ex]
- S. Chatrchyan et al., **Shape, Transverse Size, and Charged Hadron Multiplicity of Jets in pp Collisions at 7 TeV**, *JHEP*. 06 (2012) 160, doi:10.1007/JHEP06(2012)160, arXiv:1204.3170 [hep-ex]
- S. Chatrchyan et al., **Observation of a new Xi(b) baryon**, *Phys. Rev. Lett.* 108 (2012) 252002, doi:10.1103/PhysRevLett.108.252002, arXiv:1204.5955 [hep-ex]
- S. Chatrchyan et al., **Jet momentum dependence of jet quenching in PbPb collisions at** $\sqrt{s_{NN}}=2.76$ **TeV**, *Phys. Lett.* B712 (2012) 176–197, doi:10.1016/j.physletb.2012.04.058, arXiv:1202.5022 [nucl-ex]
- S. Chatrchyan et al., Suppression of non-prompt J/ψ , prompt J/ψ , and Y(1S) in PbPb collisions at $\sqrt{s_{NN}}=2.76$ TeV, JHEP. 05 (2012) 063, doi:10.1007/JHEP05(2012)063, arXiv:1201.5069 [nucl-ex]
- S. Chatrchyan et al., Study of W boson production in PbPb and pp collisions at $\sqrt{s_{NN}}=2.76$ TeV, Phys. Lett. B715 (2012) 66–87, doi:10.1016/j.physletb.2012.07.025, arXiv:1205.6334 [nucl-ex]
- S. Chatrchyan et al., Ratios of dijet production cross sections as a function of the absolute difference in rapidity between jets in proton-proton collisions at $\sqrt{s}=7$ TeV, Eur. Phys. J. C72 (2012) 2216, doi:10.1140/epjc/s10052-012-2216-6, arXiv:1204.0696 [hep-ex]
- S. Chatrchyan et al., Study of the inclusive production of charged pions, kaons, and protons in pp collisions at $\sqrt{s} = 0.9$, 2.76, and 7 TeV, Eur.

- Phys. J. C72 (2012) 2164, doi:10.1140/epjc/s10052-012-2164-1, arXiv:1207.4724 [hep-ex]
- S. Chatrchyan et al., Search for the standard model Higgs boson produced in association with W and Z bosons in pp collisions at $\sqrt{s}=7$ TeV, JHEP. 11 (2012) 088, doi:10.1007/JHEP11(2012)088, arXiv:1209.3937 [hep-ex]
- S. Chatrchyan et al., Search for supersymmetry in events with b-quark jets and missing transverse energy in pp collisions at 7 TeV, *Phys. Rev.* D86 (2012) 072010, doi:10.1103/PhysRevD.86.072010, arXiv:1208.4859 [hep-ex]
- S. Chatrchyan et al., **Search for** $B_s^0 \to \mu^+ \mu^-$ **and** $B^0 \to \mu^+ \mu^-$ **decays**, *JHEP*. 04 (2012) 033, doi:10.1007/JHEP04(2012)033, arXiv:1203.3976 [hep-ex]
- S. Chatrchyan et al., Search for a Higgs boson in the decay channel H to $\mathbf{ZZ}(*)$ to q qbar ℓ^- l+ in pp collisions at $\sqrt{s}=7$ TeV, JHEP.~04~(2012)~036,~doi:10.1007/JHEP04(2012)036,~arXiv:1202.1416~[hep-ex]
- S. Chatrchyan et al., Search for heavy long-lived charged particles in pp collisions at $\sqrt{s}=7$ TeV, Phys. Lett. B713 (2012) 408–433, doi:10.1016/j.physletb.2012.06.023, arXiv:1205.0272 [hep-ex]
- S. Chatrchyan et al., Search for third-generation leptoquarks and scalar bottom quarks in pp collisions at $\sqrt{s}=7$ TeV, JHEP. 12 (2012) 055, doi:10.1007/JHEP12(2012)055, arXiv:1210.5627 [hep-ex]
- S. Chatrchyan et al., Search for charge-asymmetric production of W bosons in $t\bar{t}+$ jet events from pp collisions at $\sqrt{s}=$ 7 TeV, Phys. Lett. B717 (2012) 351–370, doi:10.1016/j.physletb.2012.09.048, arXiv:1206.3921 [hep-ex]
- S. Chatrchyan et al., Search for microscopic black holes in pp collisions at $\sqrt{s}=7$ TeV, JHEP. 04 (2012) 061, doi:10.1007/JHEP04(2012)061, arXiv:1202.6396 [hep-ex]
- S. Chatrchyan et al., Measurement of the underlying event in the Drell-Yan process in proton-proton collisions at $\sqrt{s} = 7$ TeV, Eur. Phys. J. C72 (2012) 2080, doi:10.1140/epjc/s10052-012-2080-4, arXiv:1204.1411 [hep-ex]
- S. Chatrchyan et al., Search for Dark Matter and Large Extra Dimensions in pp Collisions Yielding a Photon and Missing Transverse Energy, *Phys. Rev. Lett.* 108 (2012) 261803, doi:10.1103/PhysRevLett.108.261803, arXiv:1204.0821 [hep-ex]
- S. Chatrchyan et al., Measurement of the underlying event activity in pp collisions at $\sqrt{s}=0.9$ and 7 TeV with the novel jet-area/median approach, JHEP. 08 (2012) 130, doi:10.1007/JHEP08(2012)130, arXiv:1207.2392 [hep-ex]
- S. Chatrchyan et al., Search for exclusive or semi-exclusive photon pair production and observation of exclusive and semi-exclusive electron

- pair production in pp collisions at $\sqrt{s} = 7$ TeV, JHEP. 11 (2012) 080, doi:10.1007/JHEP11(2012)080, arXiv:1209.1666 [hep-ex]
- S. Chatrchyan et al., Search for the standard model Higgs boson decaying into two photons in pp collisions at $\sqrt{s} = 7$ TeV, Phys. Lett. B710 (2012) 403–425, doi:10.1016/j.physletb.2012.03.003, arXiv:1202.1487 [hep-ex]
- S. Chatrchyan et al., Combined results of searches for the standard model Higgs boson in pp collisions at $\sqrt{s} = 7$ TeV, Phys. Lett. B710 (2012) 26–48, doi:10.1016/j.physletb.2012.02.064, arXiv:1202.1488 [hep-ex]
- S. Chatrchyan et al., Measurement of the relative prompt production rate of chi(c2) and chi(c1) in pp collisions at $\sqrt{s} = 7$ TeV, Eur. Phys. J. C72 (2012) 2251, doi:10.1140/epjc/s10052-012-2251-3, arXiv:1210.0875 [hep-ex]
- S. Chatrchyan et al., Measurement of the mass difference between top and antitop quarks, *JHEP*. 06 (2012) 109, doi:10.1007/JHEP06(2012)109, arXiv:1204.2807 [hep-ex]
- S. Chatrchyan et al., Search for three-jet resonances in pp collisions at $\sqrt{s} = 7$ TeV, Phys. Lett. B718 (2012) 329–347, doi:10.1016/j.physletb.2012.10.048, arXiv:1208.2931 [hep-ex]
- S. Chatrchyan et al., Search for a light charged Higgs boson in top quark decays in pp collisions at $\sqrt{s} = 7$ TeV, JHEP. 07 (2012) 143, doi:10.1007/JHEP07(2012)143, arXiv:1205.5736 [hep-ex]
- S. Chatrchyan et al., Search for neutral Higgs bosons decaying to tau pairs in pp collisions at $\sqrt{s} = 7$ TeV, Phys. Lett. B713 (2012) 68–90, doi:10.1016/j.physletb.2012.05.028, arXiv:1202.4083 [hep-ex]
- S. Chatrchyan et al., Measurement of isolated photon production in pp and PbPb collisions at $\sqrt{s_{NN}}=2.76$ TeV, $Phys.\ Lett.\ B710\ (2012)\ 256–277,$ doi:10.1016/j.physletb.2012.02.077, arXiv:1201.3093 [nucl-ex]
- S. Chatrchyan et al., **Measurement of the Z**/ γ *+b-jet cross section in pp collisions at $\sqrt{s} = 7$ TeV, *JHEP*. 06 (2012) 126, doi:10.1007/JHEP06(2012)126, arXiv:1204.1643 [hep-ex]
- S. Chatrchyan et al., Measurement of the top quark pair production cross section in pp collisions at $\sqrt{s}=7$ TeV in dilepton final states containing a τ , Phys.~Rev. D85 (2012) 112007, doi:10.1103/PhysRevD.85.112007, arXiv:1203.6810 [hep-ex]
- S. Chatrchyan et al., Search for pair produced fourth-generation uptype quarks in pp collisions at $\sqrt{s}=7$ TeV with a lepton in the final state, $Phys.\ Lett.\ B718\ (2012)\ 307–328,\ doi:10.1016/j.physletb.2012.10.038,\ arXiv:1209.0471\ [hep-ex]$
- S. Chatrchyan et al., Search for a fermiophobic Higgs boson in pp collisions at $\sqrt{s} = 7$ TeV, JHEP. 09 (2012) 111, doi:10.1007/JHEP09(2012)111,

- arXiv:1207.1130 [hep-ex]
- S. Chatrchyan et al., Search for pair production of first- and second-generation scalar leptoquarks in pp collisions at $\sqrt{s} = 7$ TeV, Phys. Rev. D86 (2012) 052013, doi:10.1103/PhysRevD.86.052013, arXiv:1207.5406 [hep-ex]
- S. Chatrchyan et al., Centrality dependence of dihadron correlations and azimuthal anisotropy harmonics in PbPb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, Eur. Phys. J. C72 (2012) 2012, doi:10.1140/epjc/s10052-012-2012-3, arXiv:1201.3158 [nucl-ex]
- S. Chatrchyan et al., Search for the standard model Higgs boson decaying to bottom quarks in pp collisions at $\sqrt{s} = 7$ TeV, Phys. Lett. B710 (2012) 284–306, doi:10.1016/j.physletb.2012.02.085, arXiv:1202.4195 [hep-ex]
- S. Chatrchyan et al., Measurement of the Λ_b cross section and the $\bar{\Lambda}_b$ to Λ_b ratio with $J/\Psi\Lambda$ decays in pp collisions at $\sqrt{s}=7$ TeV, Phys. Lett. B714 (2012) 136–157, doi:10.1016/j.physletb.2012.05.063, arXiv:1205.0594 [hep-ex]
- S. Chatrchyan et al., **Performance of CMS muon reconstruction in** pp **collision events at** $\sqrt{s} = 7$ **TeV**, JINST. 7 (2012) P10002, doi:10.1088/1748-0221/7/10/P10002, arXiv:1206.4071 [physics.ins-det]
- S. Chatrchyan et al., Azimuthal anisotropy of charged particles at high transverse momenta in PbPb collisions at $\sqrt{s_{NN}}=2.76$ TeV, Phys. Rev. Lett. 109 (2012) 022301, doi:10.1103/PhysRevLett.109.022301, arXiv:1204.1850 [nucl-ex]
- S. Chatrchyan et al., Measurement of the electron charge asymmetry in inclusive W production in pp collisions at $\sqrt{s}=7$ TeV, Phys. Rev. Lett. 109 (2012) 111806, doi:10.1103/PhysRevLett.109.111806, arXiv:1206.2598 [hep-ex]
- S. Chatrchyan et al., **A search for a doubly-charged Higgs boson** in pp collisions at $\sqrt{s}=7$ TeV, Eur. Phys. J. C72 (2012) 2189, doi:10.1140/epjc/s10052-012-2189-5, arXiv:1207.2666 [hep-ex]
- S. Chatrchyan et al., Search for heavy bottom-like quarks in 4.9 inverse femtobarns of pp collisions at $\sqrt{s}=7$ TeV, $\it{JHEP}.$ 05 (2012) 123, doi:10.1007/JHEP05(2012)123, arXiv:1204.1088 [hep-ex]
- S. Chatrchyan et al., **Performance of tau-lepton reconstruction and identification in CMS**, *JINST*. 7 (2012) P01001, doi:10.1088/1748-0221/7/01/P01001, arXiv:1109.6034 [physics.ins-det]
- S. Chatrchyan et al., Study of high-pT charged particle suppression in PbPb compared to pp collisions at $\sqrt{s_{NN}} = 2.76$ TeV, Eur. Phys. J. C72 (2012) 1945, doi:10.1140/epjc/s10052-012-1945-x, arXiv:1202.2554 [nucl-ex]
- S. Chatrchyan et al., Observation of Z decays to four leptons with the

- **CMS** detector at the LHC, *JHEP*. 12 (2012) 034, doi:10.1007/JHEP12(2012)034, arXiv:1210.3844 [hep-ex]
- S. Chatrchyan et al., A New Boson with a Mass of 125 GeV Observed with the CMS Experiment at the Large Hadron Collider, *Science*. 338 (2012) 1569–1575, doi:10.1126/science.1230816
- S. Chatrchyan et al., Search for heavy lepton partners of neutrinos in proton-proton collisions in the context of the type III seesaw mechanism, *Phys. Lett.* B718 (2012) 348–368, doi:10.1016/j.physletb.2012.10.070, arXiv:1210.1797 [hep-ex]
- S. Chatrchyan et al., Search for heavy neutrinos and W[R] bosons with right-handed couplings in a left-right symmetric model in pp collisions at sqrt(s) = 7 TeV, Phys. Rev. Lett. 109 (2012) 261802, doi:10.1103/PhysRevLett.109.261802, arXiv:1210.2402 [hep-ex]
- S. Chatrchyan et al., Search for supersymmetry in events with a lepton, a photon, and large missing transverse energy in pp collisions at $\sqrt{s} = 7$ TeV, JHEP. 06 (2011) 093, doi:10.1007/JHEP06(2011)093, arXiv:1105.3152 [hep-ex]
- S. Chatrchyan et al., Search for First Generation Scalar Leptoquarks in the e ν jj channel in pp collisions at $\sqrt{s} = 7$ TeV, Phys. Lett. B703 (2011) 246–266, doi:10.1016/j.physletb.2011.07.089, arXiv:1105.5237 [hep-ex]
- S. Chatrchyan et al., Charged particle transverse momentum spectra in pp collisions at $\sqrt{s} = 0.9$ and 7 TeV, JHEP. 08 (2011) 086, doi:10.1007/JHEP08(2011)086, arXiv:1104.3547 [hep-ex]
- S. Chatrchyan et al., Search for a Vector-like Quark with Charge 2/3 in t + Z Events from pp Collisions at $\sqrt{s} = 7$ TeV, Phys. Rev. Lett. 107 (2011) 271802, doi:10.1103/PhysRevLett.107.271802, arXiv:1109.4985 [hep-ex]
- S. Chatrchyan et al., Search for Supersymmetry in Events with b Jets and Missing Transverse Momentum at the LHC, *JHEP*. 07 (2011) 113, doi:10.1007/JHEP07(2011)113, arXiv:1106.3272 [hep-ex]
- S. Chatrchyan et al., Search for Three-Jet Resonances in pp Collisions at $\sqrt{s}=7$ TeV, Phys. Rev. Lett. 107 (2011) 101801, doi:10.1103/PhysRevLett.107.101801, arXiv:1107.3084 [hep-ex]
- S. Chatrchyan et al., Measurement of the Drell-Yan Cross Section in pp Collisions at $\sqrt{s}=7$ TeV, JHEP.~10~(2011)~007,~doi:10.1007/JHEP10(2011)007,~arXiv:1108.0566~[hep-ex]
- S. Chatrchyan et al., **Determination of Jet Energy Calibration and Transverse Momentum Resolution in CMS**, *JINST*. 6 (2011) P11002, doi:10.1088/1748-0221/6/11/P11002, arXiv:1107.4277 [physics.ins-det]
- S. Chatrchyan et al., Search for a W' boson decaying to a muon and a neutrino in pp collisions at $\sqrt{s} = 7$ TeV, Phys. Lett. B701 (2011) 160–179,

- doi:10.1016/j.physletb.2011.05.048, arXiv:1103.0030 [hep-ex]
- S. Chatrchyan et al., Long-range and short-range dihadron angular correlations in central PbPb collisions at a nucleon-nucleon center of mass energy of 2.76 TeV, *JHEP*. 07 (2011) 076, doi:10.1007/JHEP07(2011)076, arXiv:1105.2438 [nucl-ex]
- S. Chatrchyan et al., **Search for a Heavy Bottom-like Quark in** pp **Collisions at** $\sqrt{s}=7$ **TeV**, Phys. Lett. B701 (2011) 204–223, doi:10.1016/j.physletb.2011.05.074, arXiv:1102.4746 [hep-ex]
- S. Chatrchyan et al., Measurement of the Top-antitop Production Cross Section in pp Collisions at $\sqrt{s}=7$ TeV using the Kinematic Properties of Events with Leptons and Jets, Eur. Phys. J. C71 (2011) 1721, doi:10.1140/epjc/s10052-011-1721-3, arXiv:1106.0902 [hep-ex]
- S. Chatrchyan et al., Search for physics beyond the standard model using multilepton signatures in pp collisions at $\sqrt{s} = 7$ TeV, Phys. Lett. B704 (2011) 411–433, doi:10.1016/j.physletb.2011.09.047, arXiv:1106.0933 [hep-ex]
- S. Chatrchyan et al., Search for Light Resonances Decaying into Pairs of Muons as a Signal of New Physics, *JHEP*. 07 (2011) 098, doi:10.1007/JHEP07(2011)098, arXiv:1106.2375 [hep-ex]
- S. Chatrchyan et al., Measurement of the Underlying Event Activity at the LHC with $\sqrt{s} = 7$ TeV and Comparison with $\sqrt{s} = 0.9$ TeV, JHEP. 09 (2011) 109, doi:10.1007/JHEP09(2011)109, arXiv:1107.0330 [hep-ex]
- S. Chatrchyan et al., Measurement of the Polarization of W Bosons with Large Transverse Momenta in W+Jets Events at the LHC, *Phys. Rev. Lett.* 107 (2011) 021802, doi:10.1103/PhysRevLett.107.021802, arXiv:1104.3829 [hep-ex]
- S. Chatrchyan et al., Search for Large Extra Dimensions in the Diphoton Final State at the Large Hadron Collider, *JHEP*. 05 (2011) 085, doi:10.1007/JHEP05(2011)085, arXiv:1103.4279 [hep-ex]
- S. Chatrchyan et al., Measurement of the lepton charge asymmetry in inclusive W production in pp collisions at $\sqrt{s} = 7$ TeV, JHEP. 04 (2011) 050, doi:10.1007/JHEP04(2011)050, arXiv:1103.3470 [hep-ex]
- S. Chatrchyan et al., Search for B(s) and B to dimuon decays in pp collisions at 7 TeV, Phys. Rev. Lett. 107 (2011) 191802, doi:10.1103/PhysRevLett.107.191802, arXiv:1107.5834 [hep-ex]
- S. Chatrchyan et al., Search for New Physics with a Mono-Jet and Missing Transverse Energy in pp Collisions at $\sqrt{s}=7$ TeV, Phys. Rev. Lett. 107 (2011) 201804, doi:10.1103/PhysRevLett.107.201804, arXiv:1106.4775 [hep-ex]
- S. Chatrchyan et al., Search for Resonances in the Dijet Mass Spectrum from 7 TeV pp Collisions at CMS, Phys. Lett. B704 (2011) 123–142,

- doi:10.1016/j.physletb.2011.09.015, arXiv:1107.4771 [hep-ex]
- S. Chatrchyan et al., Measurement of the Inclusive Z Cross Section via Decays to Tau Pairs in pp Collisions at $\sqrt{s} = 7$ TeV, JHEP. 08 (2011) 117, doi:10.1007/JHEP08(2011)117, arXiv:1104.1617 [hep-ex]
- S. Chatrchyan et al., Search for Neutral MSSM Higgs Bosons Decaying to Tau Pairs in pp Collisions at $\sqrt{s} = 7$ TeV, Phys. Rev. Lett. 106 (2011) 231801, doi:10.1103/PhysRevLett.106.231801, arXiv:1104.1619 [hep-ex]
- S. Chatrchyan et al., **Dependence on pseudorapidity and centrality** of charged hadron production in PbPb collisions at a nucleon-nucleon centre-of-mass energy of **2.76 TeV**, *JHEP*. 08 (2011) 141, doi:10.1007/JHEP08(2011)141, arXiv:1107.4800 [nucl-ex]
- S. Chatrchyan et al., Indications of suppression of excited Υ states in PbPb collisions at $\sqrt{S_{NN}}=$ 2.76 TeV, Phys. Rev. Lett. 107 (2011) 052302, doi:10.1103/PhysRevLett.107.052302, arXiv:1105.4894 [nucl-ex]
- S. Chatrchyan et al., **A search for excited leptons in** pp **Collisions at** $\sqrt{s} =$ **7 TeV**, Phys. Lett. B704 (2011) 143–162, doi:10.1016/j.physletb.2011.09.021, arXiv:1107.1773 [hep-ex]
- S. Chatrchyan et al., Measurement of the B^0 production cross section in pp Collisions at $\sqrt{s}=7$ TeV, Phys. Rev. Lett. 106 (2011) 252001, doi:10.1103/PhysRevLett.106.252001, arXiv:1104.2892 [hep-ex]
- S. Chatrchyan et al., Search for supersymmetry in pp collisions at $\sqrt{s}=7$ TeV in events with a single lepton, jets, and missing transverse momentum, JHEP. 08 (2011) 156, doi:10.1007/JHEP08(2011)156, arXiv:1107.1870 [hep-ex]
- S. Chatrchyan et al., Measurement of the differential dijet production cross section in proton-proton collisions at $\sqrt{s}=7$ TeV, *Phys. Lett.* B700 (2011) 187–206, doi:10.1016/j.physletb.2011.05.027, arXiv:1104.1693 [hep-ex]
- S. Chatrchyan et al., **Measurement of** $W\gamma$ and $Z\gamma$ production in pp collisions at $\sqrt{s}=7$ TeV, Phys. Lett. B701 (2011) 535–555, doi:10.1016/j.physletb.2011.06.034, arXiv:1105.2758 [hep-ex]
- S. Chatrchyan et al., **Observation and studies of jet quenching in PbPb collisions at nucleon-nucleon center-of-mass energy = \mathbf{2.76} TeV,** *Phys. Rev.* **C84 (2011) 024906, doi:10.1103/PhysRevC.84.024906, arXiv:1102.1957 [nucl-ex]**
- S. Chatrchyan et al., Missing transverse energy performance of the CMS detector, JINST. 6 (2011) P09001, doi:10.1088/1748-0221/6/09/P09001, arXiv:1106.5048 [physics.ins-det]
- S. Chatrchyan et al., Study of Z boson production in PbPb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, Phys. Rev. Lett. 106 (2011) 212301,

- doi:10.1103/PhysRevLett.106.212301, arXiv:1102.5435 [nucl-ex]
- S. Chatrchyan et al., Measurement of the Differential Cross Section for Isolated Prompt Photon Production in pp Collisions at 7 TeV, *Phys. Rev.* D84 (2011) 052011, doi:10.1103/PhysRevD.84.052011, arXiv:1108.2044 [hep-ex]
- S. Chatrchyan et al., Measurement of the Strange B Meson Production Cross Section with J/Psi ϕ Decays in pp Collisions at $\sqrt{s}=7$ TeV, Phys. Rev. D84 (2011) 052008, doi:10.1103/PhysRevD.84.052008, arXiv:1106.4048 [hep-ex]
- S. Chatrchyan et al., Measurement of the t-channel single top quark production cross section in pp collisions at $\sqrt{s} = 7$ TeV, Phys. Rev. Lett. 107 (2011) 091802, doi:10.1103/PhysRevLett.107.091802, arXiv:1106.3052 [hep-ex]
- S. Chatrchyan et al., Search for Supersymmetry in pp Collisions at $\sqrt{s}=7$ TeV in Events with Two Photons and Missing Transverse Energy, Phys. Rev. Lett. 106 (2011) 211802, doi:10.1103/PhysRevLett.106.211802, arXiv:1103.0953 [hep-ex]
- S. Chatrchyan et al., Measurement of energy flow at large pseudorapidities in pp collisions at $\sqrt{s} = 0.9$ and 7 TeV, JHEP. 11 (2011) 148, doi:10.1007/JHEP11(2011)148, arXiv:1110.0211 [hep-ex]
- S. Chatrchyan et al., Measurement of the ratio of the 3-jet to 2-jet cross sections in pp collisions at $\sqrt{s} = 7$ TeV, Phys. Lett. B702 (2011) 336–354, doi:10.1016/j.physletb.2011.07.067, arXiv:1106.0647 [hep-ex]
- S. Chatrchyan et al., Search for Resonances in the Dilepton Mass Distribution in pp Collisions at $\sqrt(s) = 7$ TeV, JHEP.~05 (2011) 093, doi:10.1007/JHEP05(2011)093, arXiv:1103.0981 [hep-ex]
- S. Chatrchyan et al., Measurement of the weak mixing angle with the Drell-Yan process in proton-proton collisions at the LHC, *Phys. Rev.* D84 (2011) 112002, doi:10.1103/PhysRevD.84.112002, arXiv:1110.2682 [hep-ex]
- S. Chatrchyan et al., Measurement of the $t\bar{t}$ Production Cross Section in pp Collisions at 7 TeV in Lepton + Jets Events Using b-quark Jet Identification, Phys. Rev. D84 (2011) 092004, doi:10.1103/PhysRevD.84.092004, arXiv:1108.3773 [hep-ex]
- S. Chatrchyan et al., Search for Supersymmetry at the LHC in Events with Jets and Missing Transverse Energy, *Phys. Rev. Lett.* 107 (2011) 221804, doi:10.1103/PhysRevLett.107.221804, arXiv:1109.2352 [hep-ex]
- S. Chatrchyan et al., **Measurement of the Inclusive** W and Z **Production Cross Sections in** pp **Collisions at** $\sqrt{s} = 7$ **TeV**, JHEP. 10 (2011) 132, doi:10.1007/JHEP10(2011)132, arXiv:1107.4789 [hep-ex]
- S. Chatrchyan et al., Measurement of the Inclusive Jet Cross Section

- in pp Collisions at $\sqrt{s} = 7$ TeV, Phys. Rev. Lett. 107 (2011) 132001, doi:10.1103/PhysRevLett.107.132001, arXiv:1106.0208 [hep-ex]
- S. Chatrchyan et al., Search for New Physics with Jets and Missing Transverse Momentum in pp collisions at $\sqrt{s} = 7$ TeV, JHEP. 08 (2011) 155, doi:10.1007/JHEP08(2011)155, arXiv:1106.4503 [hep-ex]
- V. Khachatryan et al., Search for a heavy gauge boson W ' in the final state with an electron and large missing transverse energy in pp collisions at $\sqrt{s}=7$ TeV, Phys. Lett. B698 (2011) 21–39, doi:10.1016/j.physletb.2011.02.048, arXiv:1012.5945 [hep-ex]
- V. Khachatryan et al., Measurement of the Isolated Prompt Photon Production Cross Section in pp Collisions at $\sqrt{s} = 7$ TeV, Phys. Rev. Lett. 106 (2011) 082001, doi:10.1103/PhysRevLett.106.082001, arXiv:1012.0799 [hep-ex]
- V. Khachatryan et al., Search for Pair Production of First-Generation Scalar Leptoquarks in pp Collisions at $\sqrt{s} = 7$ TeV, Phys. Rev. Lett. 106 (2011) 201802, doi:10.1103/PhysRevLett.106.201802, arXiv:1012.4031 [hep-ex]
- V. Khachatryan et al., Search for Pair Production of Second-Generation Scalar Leptoquarks in pp Collisions at $\sqrt{s} = 7$ TeV, Phys. Rev. Lett. 106 (2011) 201803, doi:10.1103/PhysRevLett.106.201803, arXiv:1012.4033 [hep-ex]
- V. Khachatryan et al., Charged particle multiplicities in pp interactions at $\sqrt{s} = 0.9$, 2.36, and 7 TeV, JHEP. 01 (2011) 079, doi:10.1007/JHEP01(2011)079, arXiv:1011.5531 [hep-ex]
- V. Khachatryan et al., Search for Stopped Gluinos in pp collisions at $\sqrt{s}=7$ TeV, Phys. Rev. Lett. 106 (2011) 011801, doi:10.1103/PhysRevLett.106.011801, arXiv:1011.5861 [hep-ex]
- V. Khachatryan et al., **Search for Microscopic Black Hole Signatures at the Large Hadron Collider**, *Phys. Lett.* B697 (2011) 434–453, doi:10.1016/j.physletb.2011.02.032, arXiv:1012.3375 [hep-ex]
- V. Khachatryan et al., Measurements of Inclusive W and Z Cross Sections in pp Collisions at $\sqrt{s} = 7$ TeV, JHEP. 01 (2011) 080, doi:10.1007/JHEP01(2011)080, arXiv:1012.2466 [hep-ex]
- V. Khachatryan et al., **Prompt and non-prompt** J/ψ **production in** pp **collisions at** $\sqrt{s}=7$ **TeV**, Eur. Phys. J. C71 (2011) 1575, doi:10.1140/epjc/s10052-011-1575-8, arXiv:1011.4193 [hep-ex]
- V. Khachatryan et al., **Upsilon Production Cross-Section in pp Collisions at** sqrts = 7 **TeV**, Phys. Rev. D83 (2011) 112004, doi:10.1103/PhysRevD.83.112004, arXiv:1012.5545 [hep-ex]
- V. Khachatryan et al., Measurement of Dijet Angular Distributions and Search for Quark Compositeness in pp Collisions at sqrts = 7 TeV,

- $Phys. \quad Rev. \ Lett. \quad 106 \ (2011) \ \ 201804, \ \ doi:10.1103/PhysRevLett.106.201804, \\ arXiv:1102.2020 \ [hep-ex]$
- V. Khachatryan et al., First Measurement of Hadronic Event Shapes in pp Collisions at $\sqrt{s}=7$ TeV, Phys. Lett. B699 (2011) 48–67, doi:10.1016/j.physletb.2011.03.060, arXiv:1102.0068 [hep-ex]
- V. Khachatryan et al., Inclusive b-hadron production cross section with muons in pp collisions at $\sqrt{s} = 7$ TeV, JHEP. 03 (2011) 090, doi:10.1007/JHEP03(2011)090, arXiv:1101.3512 [hep-ex]
- V. Khachatryan et al., Measurement of Bose-Einstein Correlations in pp Collisions at $\sqrt{s} = 0.9$ and 7 TeV, JHEP. 05 (2011) 029, doi:10.1007/JHEP05(2011)029, arXiv:1101.3518 [hep-ex]
- V. Khachatryan et al., Measurement of the B^+ Production Cross Section in pp Collisions at $\sqrt{s}=7$ TeV, Phys. Rev. Lett. 106 (2011) 112001, doi:10.1103/PhysRevLett.106.112001, arXiv:1101.0131 [hep-ex]
- V. Khachatryan et al., **Search for Supersymmetry in pp Collisions at 7 TeV in Events with Jets and Missing Transverse Energy**, *Phys. Lett.* B698 (2011) 196–218, doi:10.1016/j.physletb.2011.03.021, arXiv:1101.1628 [hep-ex]
- V. Khachatryan et al., Strange Particle Production in pp Collisions at $\sqrt{s}=0.9$ and 7 TeV, JHEP.~05~(2011)~064,~doi:10.1007/JHEP05(2011)064,~arXiv:1102.4282~[hep-ex]
- V. Khachatryan et al., Search for Heavy Stable Charged Particles in pp collisions at $\sqrt{s}=7$ TeV, JHEP.~03~(2011)~024,~doi:10.1007/JHEP03(2011)024,~arXiv:1101.1645~[hep-ex]
- V. Khachatryan et al., Measurement of $B\bar{B}$ Angular Correlations based on Secondary Vertex Reconstruction at $\sqrt{s}=7$ TeV, JHEP. 03 (2011) 136, doi:10.1007/JHEP03(2011)136, arXiv:1102.3194 [hep-ex]
- V. Khachatryan et al., **Dijet Azimuthal Decorrelations in** pp **Collisions at** $\sqrt{s} = 7$ **TeV**, Phys. Rev. Lett. 106 (2011) 122003, doi:10.1103/PhysRevLett.106.122003, arXiv:1101.5029 [hep-ex]
- S. Chatrchyan et al., Commissioning and Performance of the CMS Pixel Tracker with Cosmic Ray Muons, *JINST*. 5 (2010) T03007, doi:10.1088/1748-0221/5/03/T03007, arXiv:0911.5434 [physics.ins-det]
- S. Chatrchyan et al., **Performance of the CMS Level-1 Trigger during Commissioning with Cosmic Ray Muons**, *JINST*. 5 (2010) T03002, doi:10.1088/1748-0221/5/03/T03002, arXiv:0911.5422 [physics.ins-det]
- S. Chatrchyan et al., Measurement of the Muon Stopping Power in Lead Tungstate, JINST. 5 (2010) P03007, doi:10.1088/1748-0221/5/03/P03007,

- arXiv:0911.5397 [physics.ins-det]
- S. Chatrchyan et al., Commissioning and Performance of the CMS Silicon Strip Tracker with Cosmic Ray Muons, *JINST*. 5 (2010) T03008, doi:10.1088/1748-0221/5/03/T03008, arXiv:0911.4996 [physics.ins-det]
- S. Chatrchyan et al., **Performance of CMS Muon Reconstruction** in **Cosmic-Ray Events**, *JINST*. 5 (2010) T03022, doi:10.1088/1748-0221/5/03/T03022, arXiv:0911.4994 [physics.ins-det]
- S. Chatrchyan et al., **Performance of the CMS Cathode Strip Chambers with Cosmic Rays**, *JINST*. 5 (2010) T03018, doi:10.1088/1748-0221/5/03/T03018, arXiv:0911.4992 [physics.ins-det]
- S. Chatrchyan et al., **Performance of the CMS Hadron Calorimeter with Cosmic Ray Muons and LHC Beam Data**, *JINST*. 5 (2010) T03012, doi:10.1088/1748-0221/5/03/T03012, arXiv:0911.4991 [physics.ins-det]
- S. Chatrchyan et al., CMS Data Processing Workflows during an Extended Cosmic Ray Run, *JINST*. 5 (2010) T03006, doi:10.1088/1748-0221/5/03/T03006, arXiv:0911.4842 [physics.ins-det]
- S. Chatrchyan et al., **Aligning the CMS Muon Chambers with the Muon Alignment System during an Extended Cosmic Ray Run**, *JINST*. 5 (2010) T03019, doi:10.1088/1748-0221/5/03/T03019, arXiv:0911.4770 [physics.ins-det]
- S. Chatrchyan et al., **Time Reconstruction and Performance of the CMS Electromagnetic Calorimeter**, *JINST*. 5 (2010) T03011, doi:10.1088/1748-0221/5/03/T03011, arXiv:0911.4044 [physics.ins-det]
- S. Chatrchyan et al., Commissioning of the CMS Experiment and the Cosmic Run at Four Tesla, *JINST*. 5 (2010) T03001, doi:10.1088/1748-0221/5/03/T03001, arXiv:0911.4845 [physics.ins-det]
- S. Chatrchyan et al., **Performance of the CMS Drift Tube Chambers with Cosmic Rays**, *JINST*. 5 (2010) T03015, doi:10.1088/1748-0221/5/03/T03015, arXiv:0911.4855 [physics.ins-det]
- S. Chatrchyan et al., **Performance of CMS Hadron Calorimeter Timing and Synchronization using Test Beam, Cosmic Ray, and LHC Beam Data**, *JINST*. 5 (2010) T03013, doi:10.1088/1748-0221/5/03/T03013, arXiv:0911.4877 [physics.ins-det]
- S. Chatrchyan et al., **Identification and Filtering of Uncharacteristic Noise in the CMS Hadron Calorimeter**, *JINST*. 5 (2010) T03014, doi:10.1088/1748-0221/5/03/T03014, arXiv:0911.4881 [physics.ins-det]
- S. Chatrchyan et al., Commissioning of the CMS High-Level Trigger with Cosmic Rays, JINST. 5 (2010) T03005, doi:10.1088/1748-

- 0221/5/03/T03005, arXiv:0911.4889 [physics.ins-det]
- S. Chatrchyan et al., **Performance of the CMS Drift-Tube Local Trigger with Cosmic Rays**, *JINST*. 5 (2010) T03003, doi:10.1088/1748-0221/5/03/T03003, arXiv:0911.4893 [physics.ins-det]
- S. Chatrchyan et al., Calibration of the CMS Drift Tube Chambers and Measurement of the Drift Velocity with Cosmic Rays, *JINST.* 5 (2010) T03016, doi:10.1088/1748-0221/5/03/T03016, arXiv:0911.4895 [physics.ins-det]
- S. Chatrchyan et al., **Fine Synchronization of the CMS Muon Drift-Tube Local Trigger using Cosmic Rays**, *JINST*. 5 (2010) T03004, doi:10.1088/1748-0221/5/03/T03004, arXiv:0911.4904 [physics.ins-det]
- S. Chatrchyan et al., **Alignment of the CMS Muon System with Cosmic-Ray and Beam-Halo Muons**, *JINST*. 5 (2010) T03020, doi:10.1088/1748-0221/5/03/T03020, arXiv:0911.4022 [physics.ins-det]
- S. Chatrchyan et al., **Performance Study of the CMS Barrel Resistive Plate Chambers with Cosmic Rays**, *JINST*. 5 (2010) T03017, doi:10.1088/1748-0221/5/03/T03017, arXiv:0911.4045 [physics.ins-det]
- S. Chatrchyan et al., **Performance and Operation of the CMS Electromagnetic Calorimeter**, *JINST*. 5 (2010) T03010, doi:10.1088/1748-0221/5/03/T03010, arXiv:0910.3423 [physics.ins-det]
- S. Chatrchyan et al., Precise Mapping of the Magnetic Field in the CMS Barrel Yoke using Cosmic Rays, *JINST*. 5 (2010) T03021, doi:10.1088/1748-0221/5/03/T03021, arXiv:0910.5530 [physics.ins-det]
- S. Chatrchyan et al., Alignment of the CMS Silicon Tracker during Commissioning with Cosmic Rays, *JINST*. 5 (2010) T03009, doi:10.1088/1748-0221/5/03/T03009, arXiv:0910.2505 [physics.ins-det]
- V. Khachatryan et al., **Observation of Long-Range Near-Side Angular Correlations in Proton-Proton Collisions at the LHC**, *JHEP*. 09 (2010) 091, doi:10.1007/JHEP09(2010)091, arXiv:1009.4122 [hep-ex]
- V. Khachatryan et al., **Search for Dijet Resonances in 7 TeV pp Collisions at CMS**, *Phys. Rev. Lett.* 105 (2010) 211801, doi:10.1103/PhysRevLett.105.211801, arXiv:1010.0203 [hep-ex]
- V. Khachatryan et al., **Measurement of the charge ratio of atmospheric muons with the CMS detector**, *Phys. Lett.* B692 (2010) 83–104, doi:10.1016/j.physletb.2010.07.033, arXiv:1005.5332 [hep-ex]
- V. Khachatryan et al., First Measurement of the Underlying Event Activity at the LHC with $\sqrt{s} = 0.9$ TeV, Eur. Phys. J. C70 (2010) 555–572, doi:10.1140/epjc/s10052-010-1453-9, arXiv:1006.2083 [hep-ex]
- V. Khachatryan et al., CMS Tracking Performance Results from early

- **LHC Operation**, Eur. Phys. J. C70 (2010) 1165–1192, doi:10.1140/epjc/s10052-010-1491-3, arXiv:1007.1988 [physics.ins-det]
- V. Khachatryan et al., Search for Quark Compositeness with the Dijet Centrality Ratio in pp Collisions at $\sqrt{s} = 7$ TeV, Phys. Rev. Lett. 105 (2010) 262001, doi:10.1103/PhysRevLett.105.262001, arXiv:1010.4439 [hep-ex]
- V. Khachatryan et al., First Measurement of Bose-Einstein Correlations in proton-proton Collisions at \sqrt{s} =0.9 and 2.36 TeV at the LHC, *Phys. Rev. Lett.* 105 (2010) 032001, doi:10.1103/PhysRevLett.105.032001, arXiv:1005.3294 [hep-ex]
- V. Khachatryan et al., Transverse-momentum and pseudorapidity distributions of charged hadrons in pp collisions at $\sqrt{s}=7$ TeV, Phys. Rev. Lett. 105 (2010) 022002, doi:10.1103/PhysRevLett.105.022002, arXiv:1005.3299 [hep-ex]
- V. Khachatryan et al., Transverse momentum and pseudorapidity distributions of charged hadrons in pp collisions at $\sqrt{s} = 0.9$ and 2.36 TeV, *JHEP*. 02 (2010) 041, doi:10.1007/JHEP02(2010)041, arXiv:1002.0621 [hep-ex]
- W. Adam et al., Alignment of the CMS Silicon Strip Tracker during stand-alone Commissioning, *JINST*. 4 (2009) T07001, doi:10.1088/1748-0221/4/07/T07001, arXiv:0904.1220 [physics.ins-det]
- W. Adam et al., Performance studies of the CMS Strip Tracker before installation, JINST.~4~(2009)~P06009, doi:10.1088/1748-0221/4/06/P06009, arXiv:0901.4316 [physics.ins-det]
- W. Adam et al., The CMS tracker operation and performance at the Magnet Test and Cosmic Challenge, *JINST*. 3 (2008) P07006, doi:10.1088/1748-0221/3/07/P07006
- S. Chatrchyan et al., **The CMS Experiment at the CERN LHC**, *JINST*. 3 (2008) S08004, doi:10.1088/1748-0221/3/08/S08004
- G.L. Bayatian et al., **CMS technical design report**, **volume II: Physics performance**, *J. Phys.* G34 (2007) 995–1579, doi:10.1088/0954-3899/34/6/S01
- S. Chekanov et al., **Measurement of neutral current cross sections at high Bjorken-x with the ZEUS detector at HERA**, Eur. Phys. J. C49 (2007) 523–544, doi:10.1140/epjc/s10052-006-0164-8, arXiv:hep-ex/0608014 [hep-ex]
- S. Chekanov et al., Measurement of azimuthal asymmetries in neutral current deep inelastic scattering at HERA, Eur. Phys. J. C51 (2007) 289–299, doi:10.1140/epjc/s10052-007-0310-y, arXiv:hep-ex/0608053 [hep-ex]
- S. Chekanov et al., **Event shapes in deep inelastic scattering at HERA**, *Nucl. Phys.* B767 (2007) 1–28, doi:10.1016/j.nuclphysb.2006.05.016, arXiv:hep-

- ex/0604032 [hep-ex]
- S. Chekanov et al., Measurement of prompt photons with associated jets in photoproduction at HERA, Eur. Phys. J. C49 (2007) 511–522, doi:10.1140/epjc/s10052-006-0134-1, arXiv:hep-ex/0608028 [hep-ex]
- S. Chekanov et al., Inclusive-jet and dijet cross-sections in deep inelastic scattering at HERA, Nucl. Phys. B765 (2007) 1–30, doi:10.1016/j.nuclphysb.2006.09.018, arXiv:hep-ex/0608048 [hep-ex]
- D.G. d'Enterria et al., CMS physics technical design report: Addendum on high density QCD with heavy ions, J. Phys. G34 (2007) 2307–2455, doi:10.1088/0954-3899/34/11/008
- J. Nash et al., CMS expression of interest in the SLHC, (2007)
- G.L. Bayatian et al., CMS Physics, (2006)
- S. Chekanov et al., Forward jet production in deep inelastic ep scattering and low-x parton dynamics at HERA, *Phys. Lett.* B632 (2006) 13–26, doi:10.1016/j.physletb.2005.09.066, arXiv:hep-ex/0502029 [hep-ex]
- S. Chekanov et al., Measurement of high-Q**2 deep inelastic scattering cross sections with a longitudinally polarised positron beam at HERA, *Phys. Lett.* B637 (2006) 210–222, doi:10.1016/j.physletb.2006.04.047, arXiv:hep-ex/0602026 [hep-ex]
- S. Chekanov et al., Study of the pion trajectory in the photoproduction of leading neutrons at HERA, *Phys. Lett.* B610 (2005) 199–211, doi:10.1016/j.physletb.2005.01.101, arXiv:hep-ex/0404002 [hep-ex]
- S. Chekanov et al., **Search for pentaquarks decaying to Xi-pi in deep inelastic scattering at HERA**, *Phys. Lett.* B610 (2005) 212–224, doi:10.1016/j.physletb.2005.02.016, arXiv:hep-ex/0501069 [hep-ex]
- S. Chekanov et al., Search for lepton-flavor violation at HERA, Eur. Phys. J. C44 (2005) 463–479, doi:10.1140/epjc/s2005-02399-1, arXiv:hep-ex/0501070 [hep-ex]
- S. Chekanov et al., Measurement of inelastic J/ψ production in deep inelastic scattering at HERA, Eur. Phys. J. C44 (2005) 13–25, doi:10.1140/epjc/s2005-02346-2, arXiv:hep-ex/0505008 [hep-ex]
- S. Chekanov et al., Exclusive electroproduction of phi mesons at HERA, Nucl.~Phys.~B718~(2005)~3-31,~doi:10.1016/j.nuclphysb.2005.04.009,~arXiv:hep-ex/0504010~[hep-ex]
- S. Chekanov et al., **Measurement of charm fragmentation ratios and fractions in photoproduction at HERA**, Eur. Phys. J. C44 (2005) 351–366, doi:10.1140/epjc/s2005-02397-3, arXiv:hep-ex/0508019 [hep-ex]
- S. Chekanov et al., An NLO QCD analysis of inclusive cross-section and jet-production data from the zeus experiment, Eur. Phys. J. C42 (2005)

- 1-16, doi:10.1140/epjc/s2005-02293-x, arXiv:hep-ph/0503274 [hep-ph]
- S. Chekanov et al., Multijet production in neutral current deep inelastic scattering at HERA and determination of alpha(s), Eur. Phys. J. C44 (2005) 183–193, doi:10.1140/epjc/s2005-02347-1, arXiv:hep-ex/0502007 [hep-ex]
- S. Chekanov et al., **Study of deep inelastic inclusive and diffractive scattering with the ZEUS forward plug calorimeter**, *Nucl. Phys.* B713 (2005) 3–80, doi:10.1016/j.nuclphysb.2005.02.001, arXiv:hep-ex/0501060 [hep-ex]
- S. Chekanov et al., Inclusive jet cross sections and dijet correlations in D*+- photoproduction at HERA, Nucl. Phys. B729 (2005) 492–525, doi:10.1016/j.nuclphysb.2005.09.021, arXiv:hep-ex/0507089 [hep-ex]
- S. Chekanov et al., **Isolated tau leptons in events with large missing transverse momentum at HERA**, *Phys. Lett.* B583 (2004) 41–58, doi:10.1016/j.physletb.2003.12.054, arXiv:hep-ex/0311028 [hep-ex]
- S. Chekanov et al., **Bose-Einstein correlations in one and two-dimensions in deep inelastic scattering**, *Phys. Lett.* B583 (2004) 231–246, doi:10.1016/j.physletb.2003.12.068, arXiv:hep-ex/0311030 [hep-ex]
- S. Chekanov et al., **Search for contact interactions, large extra dimensions and finite quark radius in e p collisions at HERA**, *Phys. Lett.* B591 (2004) 23–41, doi:10.1016/j.physletb.2004.03.081, arXiv:hep-ex/0401009 [hep-ex]
- S. Chekanov et al., Measurement of D*+- production in deep inelastic e+- p scattering at HERA, Phys. Rev. D69 (2004) 012004, doi:10.1103/PhysRevD.69.012004, arXiv:hep-ex/0308068 [hep-ex]
- S. Chekanov et al., Beauty photoproduction measured using decays into muons in dijet events in e p collisions at s**(1/2) = 318-GeV, *Phys. Rev.* D70 (2004) 012008, doi:10.1103/PhysRevD.74.059906, arXiv:hep-ex/0312057 [hep-ex]
- S. Chekanov et al., **Observation of K0(s) K0(s) resonances in deep inelastic scattering at HERA**, *Phys. Lett.* B578 (2004) 33–44, doi:10.1016/j.physletb.2003.10.049, arXiv:hep-ex/0308006 [hep-ex]
- S. Chekanov et al., Search for QCD instanton induced events in deep inelastic ep scattering at HERA, Eur. Phys. J. C34 (2004) 255–265, doi:10.1140/epjc/s2004-01735-3, arXiv:hep-ex/0312048 [hep-ex]
- S. Chekanov et al., **High Q**2 neutral current cross-sections in e+ p** deep inelastic scattering at s**(1/2) = 318-GeV, *Phys. Rev.* D70 (2004) 052001, doi:10.1103/PhysRevD.70.052001, arXiv:hep-ex/0401003 [hep-ex]
- S. Chekanov et al., **Photoproduction of D*+- mesons associated with a leading neutron**, *Phys. Lett.* B590 (2004) 143–160,

- doi:10.1016/j.physletb.2004.03.076, arXiv:hep-ex/0401017 [hep-ex]
- S. Chekanov et al., **Dissociation of virtual photons in events with a leading proton at HERA**, Eur. Phys. J. C38 (2004) 43–67, doi:10.1140/epjc/s2004-02047-4, arXiv:hep-ex/0408009 [hep-ex]
- S. Chekanov et al., **The Dependence of dijet production on photon virtuality in ep collisions at HERA**, Eur. Phys. J. C35 (2004) 487–500, doi:10.1140/epjc/s2004-01885-2, arXiv:hep-ex/0404033 [hep-ex]
- S. Chekanov et al., **Evidence for a narrow baryonic state decaying to K0(S) p and K0(S) anti-p in deep inelastic scattering at HERA**, *Phys. Lett.* B591 (2004) 7–22, doi:10.1016/j.physletb.2004.04.024, arXiv:hep-ex/0403051 [hep-ex]
- S. Chekanov et al., **Substructure dependence of jet cross sections at HERA and determination of alpha(s)**, *Nucl. Phys.* B700 (2004) 3–50, doi:10.1016/j.nuclphysb.2004.08.049, arXiv:hep-ex/0405065 [hep-ex]
- S. Chekanov et al., **Exclusive electroproduction of J/psi mesons at HERA**, *Nucl. Phys.* B695 (2004) 3–37, doi:10.1016/j.nuclphysb.2004.06.034, arXiv:hep-ex/0404008 [hep-ex]
- S. Chekanov et al., **Search for a narrow charmed baryonic state decaying** to D*+- p-+ in ep collisions at HERA, Eur. Phys. J. C38 (2004) 29–41, doi:10.1140/epjc/s2004-02042-9, arXiv:hep-ex/0409033 [hep-ex]
- S. Chekanov et al., **Measurement of beauty production in deep inelastic scattering at HERA**, *Phys. Lett.* B599 (2004) 173–189, doi:10.1016/j.physletb.2004.08.048, arXiv:hep-ex/0405069 [hep-ex]
- S. Chekanov et al., **Observation of isolated high E(T) photons** in deep inelastic scattering, *Phys. Lett.* B595 (2004) 86–100, doi:10.1016/j.physletb.2004.05.033, arXiv:hep-ex/0402019 [hep-ex]
- S. Chekanov et al., Measurements of inelastic J / psi and psiprime photoproduction at HERA, Eur. Phys. J. C27 (2003) 173–188, doi:10.1140/epjc/s2002-01130-2, arXiv:hep-ex/0211011 [hep-ex]
- S. Chekanov et al., Measurement of high Q**2 e- p neutral current cross-sections at HERA and the extraction of xF(3), Eur. Phys. J. C28 (2003) 175–201, doi:10.1140/epjc/s2003-01163-y, arXiv:hep-ex/0208040 [hep-ex]
- S. Chekanov et al., **Observation of the strange sea in the proton via inclusive phi meson production in neutral current deep inelastic scattering at HERA**, *Phys. Lett.* B553 (2003) 141–158, doi:10.1016/S0370-2693(02)03206-9, arXiv:hep-ex/0211025 [hep-ex]
- S. Chekanov et al., A ZEUS next-to-leading-order QCD analysis of data on deep inelastic scattering, *Phys. Rev.* D67 (2003) 012007,

- doi:10.1103/PhysRevD.67.012007, arXiv:hep-ex/0208023 [hep-ex]
- S. Chekanov et al., Measurement of proton dissociative diffractive photoproduction of vector mesons at large momentum transfer at HERA, Eur. Phys. J. C26 (2003) 389–409, doi:10.1140/epjc/s2002-01079-0, arXiv:hep-ex/0205081 [hep-ex]
- S. Chekanov et al., Scaling violations and determination of alpha(s) from jet production in gamma p interactions at HERA, *Phys. Lett.* B560 (2003) 7–23, doi:10.1016/S0370-2693(03)00380-0, arXiv:hep-ex/0212064 [hep-ex]
- S. Chekanov et al., Study of the azimuthal asymmetry of jets in neutral current deep inelastic scattering at HERA, *Phys. Lett.* B551 (2003) 226–240, doi:10.1016/S0370-2693(02)03072-1, arXiv:hep-ex/0210064 [hep-ex]
- S. Chekanov et al., Measurement of subjet multiplicities in neutral current deep inelastic scattering at HERA and determination of alpha(s), *Phys. Lett.* B558 (2003) 41–58, doi:10.1016/S0370-2693(03)00216-8, arXiv:hep-ex/0212030 [hep-ex]
- S. Chekanov et al., Measurement of event shapes in deep inelastic scattering at HERA, Eur. Phys. J. C27 (2003) 531–545, doi:10.1140/epjc/s2003-01148-x, arXiv:hep-ex/0211040 [hep-ex]
- S. Chekanov et al., Leading proton production in e+p collisions at HERA, Nucl. Phys. B658 (2003) 3–46, doi:10.1016/S0550-3213(03)00152-4, arXiv:hep-ex/0210029 [hep-ex]
- S. Chekanov et al., A Search for resonance decays to lepton + jet at HERA and limits on leptoquarks, *Phys. Rev.* D68 (2003) 052004, doi:10.1103/PhysRevD.68.052004, arXiv:hep-ex/0304008 [hep-ex]
- S. Chekanov et al., **Dijet angular distributions in photoproduction of charm at HERA**, *Phys. Lett.* B565 (2003) 87–101, doi:10.1016/S0370-2693(03)00752-4, arXiv:hep-ex/0302025 [hep-ex]
- S. Chekanov et al., Measurement of the open charm contribution to the diffractive proton structure function, *Nucl. Phys.* B672 (2003) 3–35, doi:10.1016/j.nuclphysb.2003.09.001, arXiv:hep-ex/0307068 [hep-ex]
- S. Chekanov et al., Jet production in charged current deep inelastic e+ p scattering at HERA, Eur. Phys. J. C31 (2003) 149–164, doi:10.1140/epjc/s2003-01358-2, arXiv:hep-ex/0306018 [hep-ex]
- S. Chekanov et al., Measurement of high Q**2 charged current crosssections in e+ p deep inelastic scattering at HERA, Eur. Phys. J. C32 (2003) 1–16, doi:10.1140/epjc/s2003-01363-5, arXiv:hep-ex/0307043 [hep-ex]
- S. Chekanov et al., Measurement of deeply virtual Compton scattering at HERA, *Phys. Lett.* B573 (2003) 46–62, doi:10.1016/j.physletb.2003.08.048,

- arXiv:hep-ex/0305028 [hep-ex]
- S. Chekanov et al., Search for single top production in ep collisions at HERA, Phys. Lett. B559 (2003) 153–170, doi:10.1016/S0370-2693(03)00333-2, arXiv:hep-ex/0302010 [hep-ex]
- S. Chekanov et al., Inclusive jet cross-sections in the Breit frame in neutral current deep inelastic scattering at HERA and determination of alpha(s), *Phys. Lett.* B547 (2002) 164–180, doi:10.1016/S0370-2693(02)02763-6, arXiv:hep-ex/0208037 [hep-ex]
- S. Chekanov et al., Measurement of the Q**2 and energy dependence of diffractive interactions at HERA, Eur. Phys. J. C25 (2002) 169–187, doi:10.1007/s10052-002-1003-1, arXiv:hep-ex/0203039 [hep-ex]
- S. Chekanov et al., Leading neutron production in e+p collisions at HERA, Nucl. Phys. B637 (2002) 3–56, doi:10.1016/S0550-3213(02)00439-X, arXiv:hep-ex/0205076 [hep-ex]
- S. Chekanov et al., **Measurement of diffractive production of D*+-(2010) mesons in deep inelastic scattering at HERA**, *Phys. Lett.* B545 (2002) 244–260, doi:10.1016/S0370-2693(02)02595-9, arXiv:hep-ex/0206020 [hep-ex]
- S. Chekanov et al., Measurement of high Q 2 charged current cross-sections in e $^-$ p deep inelastic scattering at HERA, *Phys. Lett.* B539 (2002) 197–217, doi:10.1016/S0370-2693(02)03097-6, arXiv:hep-ex/0205091 [hep-ex]