

Trigonometric Formulas and the Congruence Theorems

Euclidean Geometry Label triangle in the usual way: sides a, b, c , angles α, β, γ .

SAS Given a, γ, b :

1. Cosine rule $c^2 = a^2 + b^2 - 2ab \cos \gamma$ gives c .
2. Sine rule $\sin \alpha = \frac{a}{c} \sin \gamma$ gives α .
3. $\Sigma_{\Delta} = 180^\circ$ gives β (or sine rule again).

SSS Given a, b, c :

1. Cosine rule $\cos \alpha = \frac{b^2 + c^2 - a^2}{2bc}$ gives α .
2. Cos rule or sine rule for β .
3. $\Sigma_{\Delta} = 180^\circ$ gives γ .

SAA Given b, α, β :

1. $\Sigma_{\Delta} = 180^\circ$ gives γ .
2. Sine rule $a = b \frac{\sin \alpha}{\sin \beta}$ and $c = b \frac{\sin \gamma}{\sin \beta}$.

ASA Given α, c, β :

1. $\Sigma_{\Delta} = 180^\circ$ gives γ .
2. Sine rule $a = c \frac{\sin \alpha}{\sin \gamma}$ and $b = c \frac{\sin \beta}{\sin \gamma}$.

SSA One/two solutions (not a congruence theorem!) Given a, b, α :

1. Sine rule $\sin \beta = \frac{b}{a} \sin \alpha$ gives two values for β (one acute, one obtuse). If $\alpha \geq 90^\circ$, then only acute angle is valid.
2. Could instead use cosine rule $a^2 = c^2 + b^2 - 2bc \cos \alpha \implies c = b \cos \alpha \pm \sqrt{a^2 - b^2 \sin^2 \alpha}$ gives one/two values for c .

Hyperbolic Geometry Same notation where sides are *hyperbolic* lengths.

Pythagoras/trig for right-triangles

$$\cosh h = \cosh x \cosh y \quad \sin \theta = \frac{\sinh y}{\sinh h} \quad \cos \theta = \frac{\tanh x}{\tanh h} \quad \tan \theta = \frac{\tanh y}{\sinh x}$$

Cos rules $\cosh c = \cosh a \cosh b - \sinh a \sinh b \cos \gamma$ and³⁴ $\cos \gamma = -\cos \alpha \cos \beta + \sin \alpha \sin \beta \cosh c$

Sine rule $\frac{\sin \alpha}{\sinh a} = \frac{\sin \beta}{\sinh b} = \frac{\sin \gamma}{\sinh c}$

SAS Given a, γ, b :

1. Cosine rule gives (cosh) c .
2. Sine rule $\sin \alpha = \frac{\sinh a}{\sinh c} \sin \gamma$ gives α .
3. Sine rule again $\sin \beta = \frac{\sinh b}{\sinh c} \sin \gamma$ gives β (can't use Σ_{Δ} !).

SSS Given a, b, c :

1. Cosine rule $\cos \alpha = \frac{\cosh b \cosh c - \cosh a}{\sinh a \sinh b}$ gives α .
2. Repeat twice more, or use sine rule for β, γ .

SAA Given b, α, β :

1. Sine rule $\sinh a = \frac{\sin \alpha}{\sin \beta} \sinh b$ gives a .
2. Cosine formula by dropping perpendicular to compute

$$c = \tanh^{-1}(\cos \alpha \tanh b) + \tanh^{-1}(\cos \beta \tanh a)$$

3. Sine rule $\sin \gamma = \sin \alpha \frac{\sinh c}{\sinh a}$.

ASA (Two methods) Given α, c, β :

1. Drop perp from angle at $\beta = \beta_1 + \beta_2$ to get two right triangles: compute $\beta_1, \beta_2 = \beta - \beta_1$, basic trig and Pythag (mess).
2. Use second cosine rule to compute γ , now have AAA data...

AAA Given angles α, β, γ .

1. Use second cosine rule three times to find sides.

SSA Two solutions (not a congruence theorem!) Given a, b, α :

1. Sine rule $\sin \beta = \frac{\sinh b}{\sinh a} \sin \alpha$ gives two values for β (one acute, one obtuse). Can use cosine rule as in Euclidean geometry, but result ugly since not a quadratic for c ...
2. Now in one of above situations.

³⁴In Euclidean limit, second version has $\gamma = \pi - \alpha - \beta$, which reduces to $\cosh c = 1$ ($c = 0$).