

# Gene Expression Characteristics of Tumor and Adjacent Non-Tumor Tissues of Pancreatic Ductal Adenocarcinoma in-silico

Emine Güven<sup>1\*</sup>,

<sup>1</sup> Department of Biomedical Engineering, Engineering Faculty, Düzce University, Yörük/Düzce Merkez/Düzce, 81620, Turkey

*\*Corresponding author:* Emine Güven. Department of Biomedical Engineering, Faculty of Engineering, Duzce University, Duzce, Turkey. Tel: +9 0538-8733459, E-mail: emine.guven@duzce.edu.tr

**Background:** One of the mortal and the commonest malignant tumor is pancreatic ductal adenocarcinoma (PDAC). Microarray has become an important tool in the research of PDAC genes and target therapeutic drugs.

**Objectives:** This study intended to clarify the promising prognostic and biomarker targets in PDAC using GSE78229 and GSE62452 datasets publicly accessible at NCBI Gene Expression Omnibus database.

**Materials and Methods:** Employing GEOquery, Biobase, gplots, and ggplot2 packages in R software 3.6 that is based on expression analysis, we detect 221 differentially expressed genes (DEGs) of down-regulation, whereas we find 207 up-regulated genes. The gene ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) of pathway enrichments investigation of DEGs were studied.

**Results:** 28 KEGG pathways linked to pancreatic ductal adenocarcinoma (PDAC) were detected, in which the endoplasmic reticulum protein processing pathway was noted to be significant. The following 21 hub genes were detected through NetworkAnalyst founded on protein-protein interaction (PPI) network by the STRING tool: UBC, RACK1, RPL18A, RPL8, RPS23, RPS3A, RPS6, RPL10, RPL7A, RPLP1, ACTB, RPL39, P4HB, PABPC1, HSP90B1, HSPA8, GAPDH, EXOC4, and JAK1. In the TCGA database, the quantification of the expression of levels and survival probabilities were examined down and up-regulated DEGs and overall survival (OS) DEGs were investigated by Kaplan–Meier (KM) plotter (KM plotter).

**Conclusions:** The constructed study of protein-protein interactions and DAVID KEGG pathway enrichment study indicated as such ‘protein processing in endoplasmic reticulum’, ‘epstein-barr virus infection’, ‘platelet activation’, ‘ribosome’, ‘leukocyte’, ‘transendothelial migration’, and ‘protein digestion’ which had a close association with PDAC. Other hub genes discussed in this study, can be noted as promising targets for PDAC and related diseases diagnosis and treatment.

**Keywords:** biomarker; differentially expressed genes; gene ontology pathway enrichment; pancreatic ductal adenocarcinoma; cancer; endoplasmic reticulum

## 1. Introduction

One of the most deadly cancer is pancreatic ductal adenocarcinoma (PDAC) which has a 5-year overall viability rate of 3% due to the diagnosis at a distant stage (1). There have been significant improvements in terms of treatments such as pancreatectomy, radiotherapy, adjuvant, and neoadjuvant chemotherapies, and palliative care in the previous decades (2, 3). However, pancreatectomy still stays the most efficient treatment, specifically for the initial phase of pancreatic cancer (4). Thus, updated knowledge of the simple and basic mechanism of pancreatic

cancer is necessary for more useful and curable therapies and the advancement of patient survival.

Microarray has become an important tool in the research of PDAC genes and target therapeutic drugs. Recent studies suggest an extensive gene expression analysis of PDAC and related diseases by reviewing expressed gene datasets through a comparison between tumor and normal tissues in the framework of PDAC (5,6). Moreover, relative investigation of the different levels of expressed genes stays moderately constrained, and a dependable biomarker profile would be a need to develop new gene targets (7).

The protein expression alterations in the advancement and growth of PDAC and related diseases require comprehensive analysis. Furthermore, the relations among the detected DEGs, specifically protein-protein interaction (PPI) networks and underlying signaling pathways should be clarified. Prospective molecular markers from such investigations can later be checked utilizing other methods to be used in the initial stage of PDAC.

Wang et al. and Yang et al., 2016 recently performed experiments from pancreatic tissue from patients with pancreatic ductal adenocarcinoma of microarray gene-expression profiles (8,9) "(data obtainable at NCBI GEO database accession GSE78229 and GSE62452)." By studying their hub nodes globally and between tumor and normal samples constructing PPI networks, the goal of this project is to study the pathway and genetic mechanisms of PDAC and related diseases growth and to come up with candidate biomarkers for diagnosis, therapeutic targets, and predictions.

Earlier studies tackling pancreatic cancer and related diseases underlying biomarkers, due to the implicit evaluation of source and progenitor populations, need to support experimental studies with numerical analysis and statistical methods in addition to previous experiments on mice (10). Traditional therapeutic alternatives, particularly chemotherapy, are not efficient enough fighting PDAC, and notwithstanding advancements over the last 15 years, the rate of survival has not increased and has become one of the most lethal cancer types (11). Thus, constant efforts of the advancement of novel therapeutic alternatives are a need.

The generation of countless gene expression profiles of pathological samples has been directed by the advancement of high throughput sequencing that is publicly reachable via the Gene Expression Omnibus (GEO) database (12). Whereas only a small part of these datasets has been studied, the different facets of the machinery of pancreatic tumor fast growth and resilience to therapies should be on focus. The deposited datasets are re-analyzed and used to offer beneficial outcomes for further examination in-silico.

Throughout the re-examination procedure, differentially expressed genes (DEGs) are first

detected with the underlying methods, and following the pathways, molecular and biological functions of the genes concerned are examined. Previously, various experimental studies are designed to understand pancreatic cancer (13,14). The studies have concentrated only on the detection of the significant genes so far. This study suggests by pairwise comparison of the tumor and normal samples in-silico suggests novel gene targets. Thereby, this study focused on two GEO datasets which contained paired samples such that tumor and adjacent non-tumor samples, and the microarray expression dataset was analyzed. The investigation provided the detection of the DEGs, and Gene Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway enrichment analysis were subsequently studied to examine the biological process, cellular component, and molecular function, of the pathways and genes and proteins. Moreover, a protein-protein interaction (PPI) network was created and associated subnetwork was analyzed to study the hub genes of PDAC. This study may offer fresh knowledge into the machinery of pancreatic tumor expansion involving its subsequent core genes. Further, the pathways concerned in this study may assist as promising targets for the therapies of PDAC.

## **2. Objectives**

In this study, two GEO datasets were chosen, which included tumor and adjacent non-tumor tissues of the microarray expression datasets. The analysis of the datasets conducted for the detection of the DEGs, and Gene Ontology (GO) and KEGG pathway enrichment analysis were thereafter investigated in terms of the biological process, cellular component, and molecular function of the pathways were highlighted. Moreover, a PPI network was created and a subnetwork analysis was done to examine the core genes in PDAC tissues. This study aims to contribute understanding into the machinery of PDAC development and its subsequent core genes. The significant pathways that are revealed via GO and KEGG enrichments may assist as promising targets for the therapies of PDAC.

## **3. Materials and Methods**

### 3.1. The gene expression datasets

The publicly accessible dataset of gene expression from pancreatic tumor and normal samples pulled out from the GEO database with

GSE62452 and GSE78229 which the affymetrix gene-expression data of these 50 samples were also included in the previous submission as GEO




Figure 1. The raw GSE62452 and GSE78229 microarray datasets subject to quality control (QC) to examine high-dimensional structure of datasets using “uniform manifold approximate and projection” (UMAP).

accession number GSE62452 (8,9). Genomic information ranging from gene sequences to protein structure predictions was acquired. **Fig.1** displays “uniform manifold approximate and projection” (UMAP) GSE62452 and GSE78229 microarray datasets for the quality control (QC) (15).

The combined dataset contains a total of expression of 33,297 probes of in total 111 samples i.e., 50 tumor and 61 adjacent non-tumor tissues. Utilizing Bioconductor’s GEOquery package in subsequent conventional techniques, the gene expression datasets are studied (16). The list of other packages employed in this study is Biobase, biomaRT, gplots, and ggplot2 packages in R version 3.6.3 (17–19). To rectify multiple testing, calculate the adjusted *p-value*, and avoid Type I errors, we used Benjamini-Hochberg Procedure. To adjust the statistical tests locally, a hypergeometric model was performed for both

the down-regulated and up-regulated DEGs, and false discovery rates (FDR) were estimated (20).

### 3.2. Gene expression data analysis codes

Analysis was performed in the R programming language. Scripts in R programming and data analysis of the GSE datasets can be reached at [https://github.com/guven-code/bioinformaticsAnalysis\\_PDAC](https://github.com/guven-code/bioinformaticsAnalysis_PDAC) repository.

Before conducting the analyses, the datasets with the low-quality reads and low number of reads were filtered out and the rest of expression values transformed to a base-2 logarithmic scale. The study then compared specimens into two groups provided that PDAC tumor and adjacent non-tumor tissues. The expression values were normalized by taking the averages of the samples. The analysis of paired samples was conducted as calculating fold-change difference between the averages of the categories. The vast majority of

these approaches are not suitable to study microarray gene expression datasets. Although statistical methods to correct for multiple comparisons are relevant for a great deal of time namely Bonferroni correction (21).

This study used a widely utilized t-distribution and its versions as a statistical model. The variance in averages between samples divided by differences of the mean expression criterion. To detect down and up-regulated DEGs between each group via t-test, statistical significance threshold is fixed at *p-value* cutoff 0.05 and  $\log_2[\text{fold cut-off}] > 5$ .

### 3.3. Clustering analysis and Validation of Differentially Expressed Genes

Prior to transformation of gene expression values to base-2 logarithmic scale, datasets were pulled out by utilizing GEOquery package in Bioconductor (18). The study utilized gplots using heatmap.2 function and ggplot function in ggplots2 packages of R to construct heatmaps of DEGs and bar plots of GO pathways respectively (17,19). The gene expression pattern of DEGs in PDAC tumor and adjacent non-tumor tissues is validated via clustering analysis of DEGs.

### 3.4. GO and KEGG Enrichment Analysis of the Pathways

Prior to gene expression measurements of annotations for down-regulated and up-regulated and DEGs, probe IDs are mapped to the official gene symbols using Biomart package in R language.

Subsequently, GO annotations of biological processes, molecular functions, and cellular components via DAVID 6.8 ([www.david.ncifcrf.gov](http://www.david.ncifcrf.gov)) enrichments which stands for the Database for Annotation, Visualization and Integrated Discovery were studied (22). Each annotation type was retrieved utilizing DAVID and KEGG Kyoto Encyclopedia Genes and Genomes (22,23).

All annotated pathways were carefully reviewed and further partitioned according to the characteristics of their biological and molecular meanings.

the standard deviation that is generally defined as the noisiness of the samples, t-test compares the differences of the mean expression levels between the two samples. Biomart package is utilized to annotate probes to official gene symbols.

To detect DEGs, the converted gene symbols are filtered according to *p-value* and fold change

### 3.5. The protein-protein interaction (PPI) network

NetworkAnalyst publicly reachable at <https://www.networkanalyst.ca/>, offers the study of the PPI networks for single gene lists and expression values utilizing STRING Interactome (24).

For broad examination of the regulatory mechanisms in PDAC and related diseases, down and up-regulated DEGs of pancreatic tumor and normal tissues are employed to construct a human PPI network. The core genes of the network were detected with previously reported GO classification and enrichment.

The results of DAVID are then compared with NetworkAnalyst enrichments performed with KEGG (25).

### 3.6. Survival analysis

In reference to the database of The Cancer Genome Atlas Program (TCGA) (26) was used to perform survival analysis via Kaplan–Meier (KM) plotter. The analysis was carried out using down and up DEGs relying on the gene expression values in PDAC. In contrast to non-tumor tissues, gene expression levels present important individual differences in tumor tissues.

Low expression level shows the transcripts per million value (TPM) is equal to or below the upper quartile whereas high expression level shows the TPM is above the upper quartile.


## 4. Results

### 4.1. Experimental data analysis

With the gene expression result of the microarray expression datasets, we detect DEGs in a total of 428 genes from pancreatic tumor and non-tumor tissues which was shown with a volcano plot (Fig. 2). We find the down-regulated and up-regulated DEGs of pancreatic tumor and normal tissues comparison. The expression values pulled

out, and a heatmap was created to show the tumor and normal tissues discrepancy (**Fig. 3**). DEGs were selected with common *t-test*, and labelled with *p-value* < 0.05 and  $\log_2|\text{fold change}| > 5$ .

Here, the examination detected 207 differentially expressed genes of up-regulation, whereas it found 221 down-regulated genes.



**Figure 2.** (A) The boxplot shows gene expression values of each patient of the raw data without normalization. (B) Plots displaying the expression discrepancy in PDAC tumor and normal tissues comparison. Black illustrates no change (NO), red illustrates down-regulated (Down), and blue illustrates up-regulated (Up) DEGs, FC, fold change.

#### 4.2 GO enrichment and KEGG pathways

**Table 1** shows the significant enrichment of DEGs using biological processes (BP) translational initiation (GO:0006413), nuclear-transcribed mRNA catabolic process (GO:0000184), SRP-dependent cotranslational protein targeting to membrane (GO:0006614), negative regulation of apoptotic process (GO:0043066), and cell-cell adhesion (GO:0098609). The significant enrichment of GO terms in molecular function (MF) comprises protein binding (GO:0005515), cadherin binding involved in cell-cell adhesion (GO:0098641), poly(A) RNA binding (GO:0044822), structural constituent of ribosome (GO:0003735), and extracellular matrix structural constituent (GO:0005201). Lastly, GO term significant

enrichments in cellular component (CC) is resulted extracellular exosome (GO:0070062), focal adhesion (GO:0005925), extracellular matrix (GO:0031012), extracellular space (GO:0005615), and membrane (GO:0016020).

In **Table 2**, the top 11 GO terms of the down-regulated DEGs and the top 15 GO terms of the up-regulated DEGs and were ranked according to the gene counts and p-value criterion. The up-regulated DEGs mainly enriched were associated with nuclear-transcribed mRNA catabolic process, translational initiation, SRP-dependent cotranslational protein targeting to membrane, viral transcription, and translation in the ‘biological process’ group, while concerning the ‘molecular function’ group poly(A) RNA



(hsa05169), platelet activation (hsa04611), ribosome (hsa03010), leukocyte transendothelial migration (hsa03010). Among these pathways, endoplasmic reticulum protein processing

pathway might have a vital influence on multiple protein process which also has a role was aberrant in this disease.

**Table 1.** Gene expression dataset retrieved with top significant pathways GO annotations analysis of the DEGs in PDAC.


---



Category	Term	Count	p-value	Genes
GOTERM_BP	GO:0006413~translational initiation	12	3.05E-06	RPL7A, EIF2S3, RPL10, RPL18A, RPLP1, RPS6, RPS3A, PABPC1, RPL8, RPL39, EIF1, RPS23
GOTERM_BP	GO:0000184~nuclear-transcribed mRNA catabolic process, nonsense-mediated decay	11	5.74E-06	RPL7A, RPL10, RPL18A, PPP2R1A, RPLP1, RPS6, RPS3A, PABPC1, RPL8, RPL39, RPS23
GOTERM_BP	GO:0006614~SRP-dependent cotranslational protein targeting to membrane	9	4.58E-05	RPL7A, RPL10, RPL18A, RPLP1, RPS6, RPS3A, RPL8, RPL39, RPS23
GOTERM_BP	GO:0043066~negative regulation of apoptotic process	19	6.12E-05	CD74, ANXA1, DUSP1, PRKDC, ANXA4, RPS6, ANXA5, RPS3A, SOD2, MT3, HSP90B1, NME1-NME2, GREM1, GOLPH3, UBC, MDM2, PDCC4, PDE3A, ANGPTL4
GOTERM_BP	GO:0098609~cell-cell adhesion	14	1.01E-04	YWHAЕ, RAB1A, HSPA8, SND1, RAB10, LIMA1, RPL7A, MYO1B, EIF2S3, SLK, KIF5B, PRDX1, RACK1, SPTBN1
GOTERM_MF	GO:0005515~protein binding	151	1.11E-06	
GOTERM_MF	GO:0098641~cadherin binding involved in cell-cell adhesion	16	6.76E-06	YWHAЕ, RAB1A, HSPA8, ANXA1, SND1, RAB10, LIMA1, RPL7A, MYO1B, EIF2S3, SLK, KIF5B, PRDX1, RACK1, CTNNA2, SPTBN1
GOTERM_MF	GO:0044822~poly(A) RNA binding	33	2.92E-05	YWHAЕ, RPL10, PRKDC, RPL8, RPL7A, PPP1CC, RPL18A, PRDX1, UBC, RACK1, DHX37, TNPO1, SPTBN1, HSPA8, DDX18, SSB, RPS6, RPS3A, HLA-A, DDX50, QKI, SND1, EIF1, CNOT1, HNRNPH1, SUB1, HNRNPA2B1, LUC7L3, SRSF5, PABPC1, P4HB, SLC25A5, RPS23
GOTERM_MF	GO:0003735~structural constituent of ribosome	12	1.70E-04	RPL7A, RPLP0P6, RPL10, RPL18A, RPLP1, RPS6, RPS3A, SLC25A5, RPL8, RPL39, RPS23, SLC25A6
GOTERM_MF	GO:0005201~extracellular matrix structural constituent	7	2.41E-04	COL1A1, COL3A1, VCAN, COL1A2, BGN, LAMB1, FBN1
GOTERM_CC	GO:0070062~extracellular exosome	86	3.56E-15	
GOTERM_CC	GO:0005925~focal adhesion	26	4.79E-11	YWHAЕ, RPLP1, RPL8, ACTB, ACTG1, HSP90B1, CORO1C, LIMA1, RPL7A, PPP1CC, B2M, JAK1, ACTR3, HSPA8, ANXA1, ANXA5, ADAM10, MSN, RPS3A, RAB10, MMP14, ARPC2, VIM, PABPC1, P4HB, BCAR1
GOTERM_CC	GO:0031012~extracellular matrix	22	2.79E-10	HSPA8, MMP7, PRKDC, BGN, LAMB1, RPS3A, ACTG1, HSP90B1, COL1A1, COL3A1, MMP14, VCAN, SFRP2, COL1A2, PRDX1, VIM, P4HB, SLC25A5, GAPDH, PRSS2, SLC25A6, FBN1
GOTERM_CC	GO:0005615~extracellular space	41	5.77E-07	
GOTERM_CC	GO:0016020~membrane	56	8.46E-07	
KEGG_PATHWAY	hsa04141:Protein processing in endoplasmic reticulum	12	3.08E-04	EDEM3, HSPA8, LMAN1, TRAM1, EIF2AK1, SEL1L, DNAJA2, DNAJC10, UBQLN1, SSR1, P4HB, HSP90B1
KEGG_PATHWAY	hsa05169:Epstein-Barr virus infection	9	0.00198029	USP7, MDM2, PLCG2, HLA-DRA, TNFAIP3, HLA-A, VIM, HLA-G, JAK1
KEGG_PATHWAY	hsa04611:Platelet activation	9	0.00294902	COL1A1, PPP1CC, COL3A1, COL1A2, ROCK1, GNAQ, PLCG2, ACTB, ACTG1
KEGG_PATHWAY	hsa03010:Ribosome	9	0.00389219	RPL7A, RPL10, RPL18A, RPLP1, RPS6, RPS3A, RPL8, RPL39, RPS23
KEGG_PATHWAY	hsa04670:Leukocyte transendothelial migration	8	0.00567818	ROCK1, PECAM1, PLCG2, MSN, CTNNA2, ACTB, BCAR1, ACTG1

Abbreviations- gene ontology: GO; biological process: BP; cell component: CC; Kyoto Encyclopedia of Genes and Genomes: KEGG ( by the p-value)





**Figure 4.** A bar plot of the DEGs the 30 top GO annotations regarding biological processes (BP), cellular component (CC), molecular function (MF) is shown by colors of red, green, and blue respectively. The top 10 KEGG pathway enrichments of the DEGs is shown by the purple bars. The bars on the x-axis represented gene counts.


#### 4.3. The protein-protein interaction network and KEGG pathway enrichment

**Fig. 5** shows the PPI map between the set of input DEGs. The expressions of the nodes and their degree of connection were symbolized by green to purple and fields, respectively in the visualized networks. The genes with the best 19 scores according to the *p-value* are identified; these proteins also determine the functionality of the

PPI network. This network was constructed to detect the hub proteins using the betweenness centrality matrix. UBC, RACK1, RPL18A, RPL8, RPS23, RPS3A, RPS6, RPL10, RPL7A, RPLP1, ACTB, RPL39, P4HB, PABPC1, HSP90B1, HSPA8, GAPDH, EXOC4, and JAK1 were detected as the most connected hub proteins (**Fig. 5, Tables 1 and 2**). Best scoring genes comprise PPI network KEGG enrichment

analysis represents involvement ribosome (hsa03010), Ubiquitin mediated proteolysis (hsa04120), protein processing in endoplasmic reticulum (hsa04141), pathways in cancer (hsa05200) (**Table 3**). The analysis observed that DAVID and PPI network KEGG enrichment analysis revealed protein processing in endoplasmic reticulum (hsa04141), epstein-Barr virus infection (hsa05169), platelet activation

(hsa04611), ribosome (hsa03010), leukocyte transendothelial migration (hsa04670) pathways in common. The PPI network of DEGs in PDAC tumor and adjacent non-tumor tissues. Hub genes can be listed as UBC and ribosomal protein (RP) gene family sub-setting both small (RPS) and large (RPL) subunits.



**Figure 5.** The human PPI network of DEGs was created via NetworkAnalyst. Hub genes are shown in the network displayed is (A) Subnetwork 1 and (B) Subnetwork 2 human PPI networks. The “green” colored nodes present up-regulated DEGs whereas the “purple” indicates the down-regulated DEGs. The gradual color alteration presents the gene expression levels. The number of the edges where the nodes linked each other presents the “node degree”. The node sizes show a hierarchy of the significant genes in-degree centrality, the greater quantity of neighbors a node has.

Table 2. GO annotation results of the DEGs from 50 tumor and 61 adjacent non-tumor tissues of PDAC

patients. A, Down-regulated

Category	Term/gene function	count	%	P-value	Genes
GOTERM_BP	GO:0021942~radial glia guided migration of Purkinje cell	2	1.0929	0.02125823	DAB1, CTNNA2
GOTERM_BP	GO:0021756~striatum development	2	1.0929	0.0461805	CNTNAP2, SLITRK5
GOTERM_BP	GO:0016311~dephosphorylation	3	1.6393	0.05261072	ALPP, PON1, LPIN1
GOTERM_BP	GO:0060134~prepulse inhibition	2	1.0929	0.05434804	FABP7, CTNNA2
GOTERM_BP	GO:0007409~axonogenesis	3	1.6393	0.06623069	SLITRK5, KERA, CTNNA2
GOTERM_MF	GO:0005261~cation channel activity	2	1.09290	0.09604912	CATSPER3, TRPM6
GOTERM_CC	GO:0014069~postsynaptic density	4	2.18579	0.04441086	DAB1, MAP1B, CTNNA2, MT3
GOTERM_CC	GO:0043025~neuronal cell body	5	2.73224	0.04615778	CNTNAP2, DAB1, KLHL14, FABP7, RACK1
GOTERM_CC	GO:0005615~extracellular space	11	6.01093	0.06080918	OLFM3, PON1, IFNK, KRT78, SOSTDC1, GAST, KERA, PXDNL, ANGPTL4, MT3, CPA4
GOTERM_CC	GO:0043197~dendritic spine	3	1.63934	0.06842851	TENM2, MAP1B, MT3
GOTERM_CC	GO:0043005~neuron projection	4	2.18579	0.08135679	TENM2, DAB1, KLHL14, STON2
KEGG_PATHWAY	hsa04670:Leukocyte transendothelial migration	3	1.63934	0.08903048	PLCG2, CTNNA2, BCAR1

B, Up-regulated

Category	Term/gene function	count	%	P-value	Genes
GOTERM_BP	GO:0000184~nuclear-transcribed mRNA catabolic process, nonsense-mediated decay	16	6.80851	5.02E-12	RPL4, RPL30, RPL10, RPLP1, RPS6, RPL8, PNRC2, RPL7A, RPS25, RPL18A, PPP2R1A, PABPC1, RPL39, RPS24, RPS23, RPL19
GOTERM_BP	GO:0006413~translational initiation	16	6.80851	3.95E-11	RPL4, RPL30, RPL10, RPLP1, RPS6, RPL8, EIF1, RPL7A, RPS25, EIF2S3, RPL18A, PABPC1, RPL39, RPS24, RPS23, RPL19
GOTERM_BP	GO:0006614~SRP-dependent cotranslational protein targeting to membrane	13	5.53191	6.22E-10	RPL4, RPL30, RPL10, RPLP1, RPS6, RPL8, RPL7A, RPS25, RPL18A, RPL39, RPS24, RPS23, RPL19
GOTERM_BP	GO:0019083~viral transcription	13	5.53191	4.82E-09	RPL4, RPL30, RPL10, RPLP1, RPS6, RPL8, RPL7A, RPS25, RPL18A, RPL39, RPS24, RPS23, RPL19
GOTERM_BP	GO:0006412~translation	15	6.38298	1.08E-06	RPL4, RPL30, RPL10, RPLP1, RPS6, RPL8, RPL7A, RPS25, RPL18A, SLC25A5, RPL39, RPS24, SLC25A6, RPS23, RPL19

GOTERM_MF	GO:0044822~poly(A) RNA binding	40	17.0213	7.38E-11	
GOTERM_MF	GO:0005515~protein binding	138	58.7234	1.59E-10	
GOTERM_MF	GO:0003735~structural constituent of ribosome	15	6.3830	1.41E-07	RPL4, RPL30, RPLP0P6, RPL10, RPLP1, RPS6, RPL8, RPL7A, RPL18A, SLC25A5, RPL39, RPS24, SLC25A6, RPS23, RPL19
GOTERM_MF	GO:0098641~cadherin binding involved in cell-cell adhesion	16	6.8085	6.44E-07	YWHAE, RAB1A, HSPA8, ANXA1, SND1, RAB10, LIMA1, RPL7A, MYO1B, EIF2S3, SLK, EPCAM, KIF5B, PRDX1, RACK1, SPTBN1
GOTERM_MF	GO:0003723~RNA binding	19	8.0851	2.89E-05	RPL4, DDX18, RPL30, SSB, RPL8, DDX50, QKI, HSP90B1, RPL7A, RPS25, RPL18A, HNRNPH1, HNRNPA2B1, HNRNPD, PDCD4, PABPC1, SRSF5, RPL39, RPL19
GOTERM_CC	GO:0070062~extracellular exosome	90	38.2979	7.45E-25	
GOTERM_CC	GO:0031012~extracellular matrix	25	10.6383	6.89E-15	RPL30, DDX5, PRKDC, ACTG1, HSP90B1, PRDX1, PRSS2, HSPA8, MMP7, BGN, LAMB1, COL1A1, RPS25, COL3A1, MMP14, VCAN, SFRP2, COL1A2, CANX, VIM, P4HB, SLC25A5, GAPDH, FBN1, SLC25A6
GOTERM_CC	GO:0005925~focal adhesion	27	11.4894	5.36E-14	YWHAE, RPL4, RPL30, RPLP1, RPL8, ACTB, ACTG1, HSP90B1, CORO1C, LIMA1, RPL7A, PPP1CC, B2M, JAK1, RPL19, ACTR3, HSPA8, ANXA1, ANXA5, ADAM10, MSN, RAB10, MMP14, ARPC2, VIM, PABPC1, P4HB
GOTERM_CC	GO:0016020~membrane	60	25.5319	2.68E-12	
GOTERM_CC	GO:0022625~cytosolic large ribosomal subunit	10	4.2553	3.49E-08	RPL4, RPL7A, RPL30, RPLP0P6, RPL10, RPL18A, RPLP1, RPL8, RPL39, RPL19
KEGG_PATHWAY	hsa03010:Ribosome	13	5.5319	2.38E-06	RPL4, RPL30, RPL10, RPLP1, RPS6, RPL8, RPL7A, RPS25, RPL18A, RPL39, RPS24, RPS23, RPL19
KEGG_PATHWAY	hsa04141:Protein processing in endoplasmic reticulum	13	5.5319	2.25E-05	EDEM3, HSPA8, TRAM1, EIF2AK1, SEL1L, HSP90B1, LMAN1, CANX, DNAJA2, DNAJC10, UBQLN1, SSR1, P4HB
KEGG_PATHWAY	hsa05169:Epstein-Barr virus infection	10	4.2553	1.87E-04	USP7, MDM2, HLA-B, PLCG2, HLA-DRA, TNFAIP3, HLA-A, VIM, HLA-G, JAK1
KEGG_PATHWAY	hsa04612:Antigen processing and presentation	8	3.4042	2.56E-04	HSPA8, CD74, CANX, HLA-B, HLA-DRA, HLA-A, B2M, HLA-G
KEGG_PATHWAY	hsa05168:Herpes simplex infection	11	4.6808	9.22E-04	CD74, PPP1CC, USP7, EIF2AK1, CSNK2B, HLA-B, HLA-DRA, HLA-A, SRSF5, HLA-G, JAK1

---

### 3.4. The role of the endoplasmic reticulum protein processing pathway


One of the goals of this study because of its strong connection with pancreatic cancer and other related diseases through the significantly enriched pathways of DEGs in the analysis of the “endoplasmic reticulum protein processing” pathway. The endoplasmic reticulum (ER) is one of cytoplasmic organelles whereupon excretory or membrane proteins are compounded. Shortly, ER stress is a disparity within the protein folding capacity of ER and its protein pack that results from the collection of cranking proteins (27,28). ER stress has been considered to be engaged in the Parkinson's, Alzheimer's, and other deformational diseases. These diseases are caused by a few of the particular morbidic unfolding proteins (29,30).

We have performed the primary DEGs associated with ER protein processing pathway in **Fig. 6**. There were 12 DEGs specifically engaged in ER protein processing pathway, containing EDEM3, HSPA8, LMAN1, TRAM1, EIF2AK1, SEL1L, DNAJA2, DNAJC10, UBQLN1, SSR1, P4HB, HSP90B1 (see **Table 3**). We hypothesize that by virtue of over-expression of ER-associated proteins, the entire endoplasmic reticulum protein processing pathway might be unsettled in PDAC according to **Fig. 6**. We observed “endoplasmic reticulum protein processing pathway” genes in the PPI network of the entire DEGs. Related DEGs of the dataset are enriched with the ER protein processing pathway that was deciphered as new hub genes.

In **Fig. 6A**, hub genes of this pathway are selected based on expression values. HSPA8, HSP90B1, P4HB, LMAN1, UBQLN1, and DNAJA2 are the most significant genes regarding gene expression values. The most expressed HSP gene family also performs a key position which is demonstrated in subnetwork 2 (**Fig. 6B**) of the ER protein processing pathway. Heat Shock Protein 90 Beta Family Member 1 (HSP90B1) might be a gene that is associated with this pathway, folding and transforming molecular chaperones with key roles in organizing other proteins. The HSP90B1 protein is contained in the endoplasmic reticulum. HSP90B1 expression is associated with several pathogenic conditions, including tumor formation. Other DEGs in subnetwork 2 are listed as HSPA8, P4HB, and DNAJA2 in pancreatic tumor and normal tissues. These findings verify the vital duty of the ER protein processing pathway engaged in PDAC and related diseases treatment, proposing updated molecular targets to the basic drug agents.

### 3.5 Survival Analysis

To anticipate the prognostic values of the 6 hub genes of down and up-regulated DEGs, KM plotter was employed (**Fig. 7**). Among the DEGs examined, the results displayed that the low expression of OLFM3, DHX37, and RLPL23AP64 was related to worse overall survival (OS) for PDAC patients (**Fig. 7A-C**). Additionally, over-expression levels of PRDX1, JAK1, and GNG12 were related to poor OS for PDAC patients (**Fig. 7D-F**).




**Figure 6.** The PPI network of DEGs in PDAC tissues identified by NetworkAnalyst emphasizing the endoplasmic reticulum protein processing pathway associated genes are shown with “blue” nodes. (A) The numbers (bigger to slower) represent gene expression values of HSPA8, HSP90B1, P4HB, LMAN1, UBQLN1, and DNAJA2 respectively of subnetwork 1. (B) HPS gene family has the key genes HSP90B1 and HSPA8 of the subnetwork 2.



**Table 3.** Top 10 most excessive KEGG pathway enrichment analysis of global DEGs in PDAC tumor and non-tumor tissues micro-array gene expression dataset.

Term	Count	p-value	Genes
hsa04141:Protein processing in endoplasmic reticulum	12	3.08E-04	EDEM3, HSPA8, LMAN1, TRAM1, EIF2AK1, SEL1L, DNAJA2, DNAJC10, UBQLN1, SSR1, P4HB, HSP90B1
hsa05169:Epstein-Barr virus infection	9	0.00198029	USP7, MDM2, PLCG2, HLA-DRA, TNFAIP3, HLA-A, VIM, HLA-G, JAK1
hsa04611:Platelet activation	9	0.00294902	COL1A1, PPP1CC, COL3A1, COL1A2, ROCK1, GNAQ, PLCG2, ACTB, ACTG1
hsa03010:Ribosome	9	0.0038922	RPL7A, RPL10, RPL18A, RPLP1, RPS6, RPS3A, RPL8, RPL39, RPS23
hsa04670:Leukocyte transendothelial migration	8	0.00567818	ROCK1, PECAM1, PLCG2, MSN, CTNNA2, ACTB, BCAR1, ACTG1
hsa04974:Protein digestion and absorption	7	0.00597994	COL1A1, CPA2, COL3A1, CPA1, COL1A2, SLC38A2, PRSS2
hsa05168:Herpes simplex infection	10	0.00724955	CD74, PPP1CC, USP7, EIF2AK1, CSNK2B, HLA-DRA, HLA-A, SRSF5, HLA-G, JAK1
hsa04612:Antigen processing and presentation	6	0.01371483	HSPA8, CD74, HLA-DRA, HLA-A, B2M, HLA-G
hsa04510:Focal adhesion	10	0.01503298	COL1A1, PDGFRA, PPP1CC, COL3A1, COL1A2, ROCK1, LAMB1, ACTB, BCAR1, ACTG1
hsa05164:Influenza A	9	0.01624626	IVNS1ABP, HSPA8, EIF2AK1, HLA-DRA, NXT2, PRSS2, ACTB, JAK1, ACTG1



**Figure 7.** Prognostic values of six DEGs in PDAC patients. Effect of expression levels on PDAC patients of survival. Down-regulated DEGs **(A)** OLFM3 ( $p = 0.075$ ), **(B)** DHX37 ( $p = 0.076$ ), **(C)** RLPL23AP64 ( $p = 0.93$ ) and up-regulated DEGs **(D)** PRDX1 ( $p = 0.29$ ), **(E)** JAK1 ( $p = 0.59$ ), and **(F)** GNG12 ( $p = 0.044$ ).

**Table 4.** The top 15 genes of PPI network of DEGs in PDAC tumor and non-tumor tissues gene expression data.

Gene ID	Genes	Node Degree	Betweenness centrality	Expression
7316	UBC	72	3370.87	6.89073
10399	RACK1	11	58.71	6.71587613
6142	RPL18A	10	56.87	6.71812216
6132	RPL8	10	56.87	6.14703757
6228	RPS23	9	38.72	5.46938802
6189	RPS3A	9	38.72	5.67086
6194	RPS6	9	38.72	5.43906883
6134	RPL10	9	0.78	7.22927072
6130	RPL7A	9	0.78	5.87863243
6176	RPLP1	8	0.5	4.95154703
60	ACTB	7	9.5	7.05993541
6170	RPL39	7	3.36	5.51250622
5034	P4HB	4	164	6.27703351
26986	PABPC1	4	20.22	5.54081207

#### 4. Discussion

The prevalence of pancreatic ductal adenocarcinoma and the related survival rates have demonstrated a decrease in tendency in the past years (1). One study showed that PDAC patients survive for only 4 months typically without therapies. Moreover, for patients who undergo surgery and take required therapies the survival is not significantly increased. Thus, precise quick identification of PDAC and the advancement of powerful specific remedies is of fundamental significance. Recent research detected hub genes in PDAC that were stated to be of diagnostic significance (13). In this project, the combination of GSE78229 and GSE62452 datasets from patients with pancreatic ductal adenocarcinoma of microarray gene expression profiles were comprehensively studied, holding the expressions of 50 tumor and 61 non-tumor tissues. 221 DEGs of down-regulation were identified, whereas we found 207 up-regulated DEGs using R language (**Figs. 2 and 3**). And further, GO functional enrichments and KEGG pathway analyses were performed which showed locational and functional information of these DEGs. The outcomes of the GO enrichments show that most of the DEGs were significantly enriched in processes of vital cell organizations and functions, including extracellular matrix-associated proteins, extracellular exosome formation, extracellular matrix organization, extracellular space, and extracellular region (**Tables 1 and 2**). In addition, KEGG pathway study resulted mostly in the upregulated DEGs were entailed in ribosome (hsa03010), protein processing in endoplasmic reticulum (hsa04141), Epstein-Barr virus infection (hsa05169), antigen processing and presentation (hsa04612), Herpes simplex infection (hsa05168) (**Table 3**). The DEGs are involved in other pathways such as protein, digestion and absorption, PI3K/Akt signaling pathway, focal adhesion, pancreatic secretion, oocyte meiosis, bacterial invasion of epithelial cells, and hippo signaling pathway might be of importance (**Fig. 4**). Recent studies have discovered an important role for 'ribosome' pathway genes involved in ribosome biogenesis in early pancreatic development (31). Another recent review proposed the interaction of focal adhesions with the extracellular matrix might

advance epithelial-mesenchymal transition (EMT), therefore foster cell carcinogenesis (32). Moreover, the PI3K/Akt signaling pathway (hsa04151) is the other significant pathway in the understanding of the PDAC (33). To obtain an in-depth understanding of these DEGs, this study analyzed the constructed PPI network and found that UBC, RACK1, RPL18A, RPL8, RPS23, RPS3A, RPS6, RPL10, RPL7A, RPLP1, ACTB, RPL39, P4HB, PABPC1, HSP90B1, HSPA8, GAPDH, EXOC4, and JAK1 (**Fig. 5**) were the hub genes, which may be critical to the molecular and biological mechanisms underlying the development of pancreatic tumor and may thus serve as promising therapeutic targets. Furthermore, mutations employ in ER protein processing pathway genes and many related pathway genes provide pancreatic ductal adenocarcinoma carcinogenesis. Moreover, recent research found that metabolic alteration is classified among features of cancer, particularly the malfunction of pancreatic secretion. Metabolic and functional changes in ribosomal and ER protein processing pathways are evident in pancreatic cancer (34,35). In addition to their standard roles, ER protein processing pathway further rules metabolism characteristics of aggregation of misfolded proteins in the endoplasmic reticulum which drives ER stress and then triggers the unfolded protein response (UPR) signaling pathway (**Fig. 6**). Ubiquitin-C (UBC) was detected as one of the core genes with the biggest degree of connectivity (Table 4). One study revealed that UBC expression associates with increased patient survival in PDAC (36). UBC is a key gene that directly engages with other genes such as RACK1 and ribosomal protein gene family (RP), proposes that it might be a central component that leads to a bad prognosis of PDAC regulated by perineural invasion (37). UBC has a vital task in diseases comprising renal cancer and lung cancer. RACK1 is a scaffold protein that is a receptor at the same time that initiates signal transduction of protein kinase C that is participated in most of the intracellular signal pathways (38). Earlier research has uncovered that RACK1 is closely linked to the development and progression of several cancer types, along with gastric cancer and hepatocellular carcinoma (39). Although the

exact involvement of RACK1 in human PDAC growth still stays unidentified, it was established that RACK1 was significantly up-regulated in human PDAC samples and cell lines (40). Taken together, the results of the bioinformatics analysis of four GEO microarray datasets of PDAC indicated that ribosome (hsa03010) and protein processing in endoplasmic reticulum (hsa04141), participate in the onset and development of PDAC. The low expression of OLFM3, DHX37, and RLPL23AP64, as well as the overexpression of PRDX1, JAK1, and GNG12, were observably related to unsatisfactory survival effects in patients with PDAC (Fig. 7). However, further studies need to be implemented to discover the molecular mechanisms and biological functions of the DEGs, biological processes, cellular components, molecular function, and KEGG pathways to estimate whether they can serve as novel potential biomarkers or therapeutic targets in PDAC patients. Additional studies are required for clinical lab confirmation of predicted proteins that are expressed in PDAC tumor and non-tumor datasets and to express at the developmental stage of pancreatic ductal adenocarcinoma. More research is needed in the field of cancer biology to detect pancreatic cancer and subset diseases at their early stage. This paper also emphasizes the importance of microarray experiments in comprehending pancreatic cancer and related diseases and approach to study several results of gene expression data, like differentially expressed genes analysis, pathway and process identification, and protein-protein interaction network study.

## References

1. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2018. *CA Cancer J Clin* [Internet]. 2018 [cited 2020 Oct 22];68(1):7–30. Available from: <https://acsjournals.onlinelibrary.wiley.com/doi/abs/10.3322/caac.21442>
2. Gillen S, Schuster T, Büschenfelde CM, Friess H, Kleeff J. Preoperative/Neoadjuvant Therapy in Pancreatic Cancer: A Systematic Review and Meta-analysis of Response and Resection Percentages. *PLOS Med* [Internet]. 2010 Nis [cited 2020 Oct 22];7(4):e1000267. Available from:

## 5. Conclusion

An accurate and faster identification of PDAC and the advancement of strong particular remedies is of importance. It is identified 221 differentially expressed genes (DEGs) of down-regulation, whereas we find 207 up-regulated genes. 28 KEGG pathways linked to pancreatic PDAC were detected, in which the endoplasmic reticulum protein processing pathway was noted to be significant. 21 hub genes were identified regarding PPI network: UBC, RACK1, RPL18A, RPL8, RPS23, RPS3A, RPS6, RPL10, RPL7A, RPLP1, ACTB, RPL39, P4HB, PABPC1, HSP90B1, HSPA8, GAPDH, EXOC4, and JAK1. The constructed study of PPIs, DAVID, and KEGG pathway enrichment studies agreed that ‘protein processing in endoplasmic reticulum’, ‘epstein-barr virus infection’, ‘platelet activation’, ‘ribosome’, ‘leukocyte’, ‘transendothelial migration’, and ‘protein digestion’ which may relieve a close association with PDAC.

## Abbreviations

DEGs: Differentially Expressed Genes  
 PDAC: Pancreatic Adenocarcinoma  
 DAVID: The Database for Annotation, Visualization and Integrated Discovery  
 KEGG: Kyoto Encyclopedia of Genes and Genomes  
 GEO: Gene Expression Omnibus  
 GO: Gene Ontology  
 ER: Endoplasmic Reticulum  
 PPI: Protein-protein Interaction  
 BP: Biological Process  
 MF: Molecular Function  
 CC: Cellular Component  
<https://journals.plos.org/plosmedicine/article?id=10.1371/journal.pmed.1000267>

3. Tempero MA, Malafa MP, Behrman SW, Benson AB, Casper ES, Chiorean EG, et al. Pancreatic adenocarcinoma, version 2.2014. *J Natl Compr Canc Netw*. 2014;12(8):1083–93.
4. Lambert A, Schwarz L, Borbath I, Henry A, Van Laethem J-L, Malka D, et al. An update on treatment options for pancreatic adenocarcinoma. *Ther Adv Med Oncol* [Internet]. 2019 Jan [cited 2020 Oct 22];11:175883591987556. Available from: <http://journals.sagepub.com/doi/10.1177/1758835919875568>

5. Alldinger I, Dittert D, Peiper M, Fusco A, Chiappetta G, Staub E, et al. Gene expression analysis of pancreatic cell lines reveals genes overexpressed in pancreatic cancer. *Pancreatolgy*. 2005;5(4–5):370–9.
6. Jones S, Zhang X, Parsons DW, Lin JC-H, Leary RJ, Angenendt P, et al. Core signaling pathways in human pancreatic cancers revealed by global genomic analyses. *science*. 2008;321(5897):1801–6.
7. Grützmann R, Boriss H, Ammerpohl O, Lüttges J, Kalthoff H, Schackert HK, et al. Meta-analysis of microarray data on pancreatic cancer defines a set of commonly dysregulated genes. *Oncogene*. 2005;24(32):5079–88.
8. Wang J, Yang S, He P, Schetter AJ, Gaedcke J, Ghadimi BM, et al. Endothelial Nitric Oxide Synthase Traffic Inducer (NOSTRIN) is a Negative Regulator of Disease Aggressiveness in Pancreatic Cancer. *Clin Cancer Res*. 2016 Dec 15;22(24):5992–6001.
9. Yang S, He P, Wang J, Schetter A, Tang W, Funamizu N, et al. A Novel MIF Signaling Pathway Drives the Malignant Character of Pancreatic Cancer by Targeting NR3C2. *Cancer Res*. 2016 Jul 1;76(13):3838–50.
10. Dineen SP, Roland CL, Greer R, Carbon JG, Toombs JE, Gupta P, et al. Smac mimetic increases chemotherapy response and improves survival in mice with pancreatic cancer. *Cancer Res*. 2010;70(7):2852–61.
11. Rahib L, Smith BD, Aizenberg R, Rosenzweig AB, Fleshman JM, Matrisian LM. Projecting Cancer Incidence and Deaths to 2030: The Unexpected Burden of Thyroid, Liver, and Pancreas Cancers in the United States. *Cancer Res* [Internet]. 2014 Jun 1 [cited 2020 Oct 28];74(11):2913–21. Available from: <http://cancerres.aacrjournals.org/cgi/doi/10.1158/0008-5472.CAN-14-0155>
12. Barrett T, Troup DB, Wilhite SE, Ledoux P, Rudnev D, Evangelista C, et al. NCBI GEO: mining tens of millions of expression profiles—database and tools update. *Nucleic Acids Res* [Internet]. 2007 Jan 1 [cited 2021 Jun 30];35(suppl\_1):D760–5. Available from: <https://doi.org/10.1093/nar/gkl887>
13. Lv K, Yang J, Sun J, Guan J. Identification of key candidate genes for pancreatic cancer by bioinformatics analysis. *Exp Ther Med* [Internet]. 2019 Jul 1 [cited 2020 Nov 12];18(1):451–8. Available from: <http://www.spandidos-publications.com/10.3892/etm.2019.7619/abstract>
14. Kwon M-S, Kim Y, Lee S, Namkung J, Yun T, Yi SG, et al. Integrative analysis of multi-omics data for identifying multi-markers for diagnosing pancreatic cancer. *BMC Genomics* [Internet]. 2015 Aug 17 [cited 2021 Jun 28];16(9):S4. Available from: <https://doi.org/10.1186/1471-2164-16-S9-S4>
15. Diaz-Papkovich A, Anderson-Trocmé L, Gravel S. A review of UMAP in population genetics. *J Hum Genet* [Internet]. 2021 Jan [cited 2021 Apr 2];66(1):85–91. Available from: <https://www.nature.com/articles/s10038-020-00851-4>
16. Davis S, Meltzer P. GEOquery: A bridge between the Gene Expression Omnibus (GEO) and BioConductor. *Bioinforma Oxf Engl*. 2007 Aug 1;23:1846–7.
17. Warnes GR, Bolker B, Bonebakker L, Gentleman R, Huber W, Liaw A, et al. gplots: Various R programming tools for plotting data. *R Package Version*. 2009;2(4):1.
18. Durinck S, Moreau Y, Kasprzyk A, Davis S, De Moor B, Brazma A, et al. BioMart and Bioconductor: a powerful link between biological databases and microarray data analysis. *Bioinformatics*. 2005;21(16):3439–40.
19. Wickham H. ggplot2. *WIREs Comput Stat* [Internet]. 2011 [cited 2021 Jul 6];3(2):180–5. Available from: <https://wires.onlinelibrary.wiley.com/doi/abs/10.1002/wics.147>
20. Dudoit S, Shaffer JP, Boldrick JC. Multiple hypothesis testing in microarray experiments. *Stat Sci*. 2003;71–103.
21. Tarca AL, Romero R, Draghici S. Analysis of microarray experiments of gene expression profiling. *Am J Obstet Gynecol*. 2006 Aug;195(2):373–88.
22. Sherman BT, Lempicki RA. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc*. 2009;4(1):44.
23. Kanehisa M, Sato Y, Kawashima M, Furumichi M, Tanabe M. KEGG as a reference resource for gene and protein annotation. *Nucleic Acids Res*. 2016;44(D1):D457–62.
24. Szklarczyk D, Franceschini A, Wyder S, Forslund K, Heller D, Huerta-Cepas J, et al. STRING v10: protein–protein interaction networks, integrated over the tree of life. *Nucleic Acids Res*. 2015;43(D1):D447–52.

25. Zhou G, Soufan O, Ewald J, Hancock REW, Basu N, Xia J. NetworkAnalyst 3.0: a visual analytics platform for comprehensive gene expression profiling and meta-analysis. *Nucleic Acids Res.* 2019;47(W1):W234–41. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6260831/>
26. Chandrashekar DS, Bashel B, Balasubramanya SAH, Creighton CJ, Ponce-Rodriguez I, Chakravarthi BV, et al. UALCAN: a portal for facilitating tumor subgroup gene expression and survival analyses. *Neoplasia.* 2017;19(8):649–58.
27. Kelly E, Greene CM, Carroll TP, McElvaney NG, O'Neill SJ. Selenoprotein S/SEPS1 modifies endoplasmic reticulum stress in Z variant alpha1-antitrypsin deficiency. *J Biol Chem.* 2009 Jun 19;284(25):16891–7.
28. Hidvegi T, Schmidt BZ, Hale P, Perlmutter DH. Accumulation of mutant alpha1-antitrypsin Z in the endoplasmic reticulum activates caspases-4 and -12, NFkappaB, and BAP31 but not the unfolded protein response. *J Biol Chem.* 2005 Nov 25;280(47):39002–15.
29. Forman MS, Lee VM-Y, Trojanowski JQ. “Unfolding” pathways in neurodegenerative disease. *Trends Neurosci.* 2003 Aug;26(8):407–10.
30. Gow A, Sharma R. The unfolded protein response in protein aggregating diseases. *Neuromolecular Med.* 2003;4(1–2):73–94.
31. Provost E, Weier CA, Leach SD. Multiple Ribosomal Proteins Are Expressed at High Levels in Developing Zebrafish Endoderm and Are Required for Normal Exocrine Pancreas Development. *Zebrafish* [Internet]. 2013 Jun [cited 2021 Jul 5];10(2):161–9. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3673614/>
32. Burridge K. Focal adhesions: a personal perspective on a half century of progress. *FEBS J* [Internet]. 2017 [cited 2021 Jul 5];284(20):3355–61. Available from: <https://febs.onlinelibrary.wiley.com/doi/abs/10.1111/febs.14195>
33. Jiang N, Dai Q, Su X, Fu J, Feng X, Peng J. Role of PI3K/AKT pathway in cancer: the framework of malignant behavior. *Mol Biol Rep* [Internet]. 2020 [cited 2020 Nov 2];47(6):4587–629. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7295848/>
34. Grant TJ, Hua K, Singh A. Molecular Pathogenesis of Pancreatic Cancer. *Prog Mol Biol Transl Sci* [Internet]. 2016 [cited 2021 Jul 5];144:241–75. Available from:
35. Wang W, Nag S, Zhang X, Wang M-H, Wang H, Zhou J, et al. Ribosomal Proteins and Human Diseases: Pathogenesis, Molecular Mechanisms, and Therapeutic Implications. *Med Res Rev* [Internet]. 2015 Mar [cited 2021 Jul 5];35(2):225–85. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4710177/>
36. Chu KY, Li H, Wada K, Johnson JD. Ubiquitin C-terminal hydrolase L1 is required for pancreatic beta cell survival and function in lipotoxic conditions. *Diabetologia* [Internet]. 2012 Jan 1 [cited 2021 Jul 5];55(1):128–40. Available from: <https://doi.org/10.1007/s00125-011-2323-1>
37. Yang Y-H, Zhang Y-X, Gui Y, Liu J-B, Sun J-J, Fan H. Analysis of the autophagy gene expression profile of pancreatic cancer based on autophagy-related protein microtubule-associated protein 1A/1B-light chain 3. *World J Gastroenterol* [Internet]. 2019 May 7 [cited 2020 Nov 3];25(17):2086–98. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6506580/>
38. Adams DR, Ron D, Kiely PA. RACK1, A multifaceted scaffolding protein: Structure and function. *Cell Commun Signal.* 2011;9(1):1–24.
39. Li J, Xie D. RACK1, a versatile hub in cancer. *Oncogene.* 2015;34(15):1890–8.
40. Wang L, Liu H-L, Li Y, Yuan P. Proteomic analysis of pancreatic intraepithelial neoplasia and pancreatic carcinoma in rat models. *World J Gastroenterol WJG.* 2011;17(11):1434.



