

CLASS QUIZ SOLUTIONS: JANUARY 7: INVERSE TRIGONOMETRIC FUNCTIONS

MATH 153, SECTION 55 (VIPUL NAIK)

1. PERFORMANCE REVIEW

28 people took this 4-question quiz. The scores are as follows:

- Score of 0: 1 person.
- Score of 1: 2 people.
- Score of 2: 7 people.
- Score of 3: 7 people.
- Score of 4: 11 people.

The mean score was 2.89, the median score was 3, and the modal score was 4.

Here is the question-wise performance (full solutions in the next section):

- (1) Option (B): 18 people got this correct.
- (2) Option (C): 22 people got this correct.
- (3) Option (B): 20 people got this correct.
- (4) Option (A): 21 people got this correct.

2. SOLUTIONS

- (1) What is the domain of $\arcsin \circ \arcsin$? Here, *domain* refers to the maximal possible subset of \mathbb{R} on which the function is defined.
 - (A) $[-1, 1]$
 - (B) $[-\sin 1, \sin 1]$
 - (C) $[-\arcsin 1, \arcsin 1]$
 - (D) $[-\sin(2/\pi), \sin(2/\pi)]$
 - (E) $[-\arcsin(2/\pi), \arcsin(2/\pi)]$

Answer: Option (B)

Explanation: For x to be in the domain of this function, we need to be able to apply \arcsin twice to x . Thus, the result obtained by applying \arcsin once to x must be in the domain of \arcsin , which is $[-1, 1]$. Thus, x must be in the inverse image of $[-1, 1]$ under \arcsin , which is $[\sin(-1), \sin 1]$. Since \sin is odd, this can be rewritten as $[-\sin 1, \sin 1]$.

Post-performance review: 18 people got this correct. 5 people chose (A), 3 people chose (C), 1 person chose (D), and 1 person chose (E).

- (2) Suppose f is a polynomial with degree at least one and positive leading coefficient. Consider the function $g(x) := \arctan(f(x))$. What can we say about the horizontal asymptotes of the graph $y = g(x)$?
 - (A) The horizontal asymptote is $y = \pi/2$ both for $x \rightarrow +\infty$ and for $x \rightarrow -\infty$, regardless of f .
 - (B) The horizontal asymptote is $y = \pi/2$ for $x \rightarrow +\infty$ and $-\pi/2$ for $x \rightarrow -\infty$, regardless of f .
 - (C) The horizontal asymptote is $y = \pi/2$ for $x \rightarrow +\infty$, and as $x \rightarrow -\infty$, it is $y = \pi/2$ if f has even degree and $y = -\pi/2$ if f has odd degree.
 - (D) The horizontal asymptote is $y = f(\pi/2)$ both for $x \rightarrow +\infty$ and for $x \rightarrow -\infty$.
 - (E) The horizontal asymptote is $y = f(\pi/2)$ for $x \rightarrow +\infty$ and as $x \rightarrow -\infty$, it is $y = f(\pi/2)$ if f has even degree and $y = f(-\pi/2)$ if f has odd degree.

Answer: Option (C)

Explanation: The key observation is that as $u \rightarrow +\infty$, $\arctan u \rightarrow \pi/2$, and as $u \rightarrow -\infty$, $\arctan u \rightarrow -\pi/2$. The question now is what happens to $f(x)$ as x approaches $\pm\infty$.

Since f has positive degree and positive leading coefficient, $f(x) \rightarrow +\infty$ as $x \rightarrow +\infty$. The behavior as $x \rightarrow -\infty$ depends on whether the degree of f is even or odd. If the even, then it goes to $+\infty$, and if the latter, then it goes to $-\infty$.

Combining these observations gives us (C).

Post-performance review: 22 people got this correct. 2 people chose (A), 2 people chose (B), 2 people chose (E).

- (3) Consider the function $f(x) := \arcsin(\sin x)$ on the domain $[\pi/2, 3\pi/2]$. Which of the following is $f(x)$ equal to on that domain?
- (A) $\pi + x$
 - (B) $\pi - x$
 - (C) $x - \pi$
 - (D) $(3\pi/2) - x$
 - (E) $x - (\pi/2)$

Answer: Option (B)

Explanation: Note that $\sin(f(x)) = \sin(\arcsin(\sin x)) = \sin x$. We note that of the options given here, $\pi - x$ is the only option satisfying this constraint. Additionally, we need to check that $x \mapsto \pi - x$, sends the interval $[\pi/2, 3\pi/2]$ to the interval $[-\pi/2, \pi/2]$, which it does.

Post-performance review: 20 people got this correct. 5 people chose (C), 1 person chose (A), 1 person chose (D), 1 person chose (E).

- (4) Consider the function $f(x) := \arccos(\sin x)$ on all of \mathbb{R} . What can we say about the function f ?
- (A) f is periodic, continuous, and piecewise linear.
 - (B) f is periodic and continuous but is not piecewise linear.
 - (C) f is continuous and piecewise linear but not periodic.
 - (D) f is periodic but not continuous.
 - (E) f is continuous but not periodic or piecewise linear.

Answer: Option (A)

Explanation: First, note that the function *does* make sense on all of \mathbb{R} : \sin is defined everywhere and has range $[-1, 1]$, and this is precisely the domain of \arccos . Since both \sin and \arccos are continuous on their respective domains, the composite function is also continuous.

Since \sin has a period of 2π , the composite is also periodic. (In this case, the period turns out to be exactly 2π , though in general the period of a composite function could be smaller than that of the inner function).

It remains to justify piecewise linearity. Working out the definition reveals this. On the interval $[2n\pi - \pi/2, 2n\pi + \pi/2]$, the correct definition is $2n\pi + \pi/2 - x$. On an interval $[2n\pi + \pi/2, 2n\pi + 3\pi/2]$, the correct definition is $x - 2n\pi - \pi/2$. Thus, the definition is piecewise linear on intervals of length π , and each of the pieces has slope 1. In fact, the graph of the function has a sawtooth-like shape.

Post-performance review: 21 people got this correct. 5 people chose (B), 1 person chose (C), and 1 person chose (D).