

Fast and Small




The Elementary World

Elements

Then




Elements




John Dalton

Reihen	Gruppe I. — R'O	Gruppe II. — RO	Gruppe III. — R'O ³	Gruppe IV. RH ⁴ RO ⁴	Gruppe V. RH ⁵ R'O ⁵	Gruppe VI. RH ⁶ RO ⁶	Gruppe VII. RH R'O ⁷	Gruppe VIII. — RO ⁴
1	H=1							
2	Li=7	Be=9,4	B=11	C=12	N=14	O=16	F=19	
3	Na=23	Mg=24	Al=27,3	Si=28	P=31	S=32	Cl=35,5	
4	K=39	Ca=40	—=44	Ti=48	V=51	Cr=52	Mn=55	Fe=56, Co=59, Ni=59, Cu=63.
5	(Cu=63)	Zn=65	—=68	—=72	As=75	Se=78	Br=80	
6	Rb=86	Sr=87	?Yt=88	Zr=90	Nb=94	Mo=96	—=100	Ru=104, Rh=104, Pd=106, Ag=108.
7	(Ag=108)	Cd=112	In=113	Sn=118	Sb=122	Te=125	J=127	
8	Cs=133	Ba=137	?Di=138	?Ce=140	—	—	—	— — — —
9	(—)	—	—	—	—	—	—	
10	—	—	?Er=178	?La=180	Ta=182	W=184	—	Os=195, Ir=197, Pt=198, Au=199.
11	(Au=199)	Hg=200	Tl=204	Pb=207	Bi=208	—	—	
12	—	—	—	Th=231	—	U=240	—	— — — —




Dmitri Mendeleev

Elementary, ... Watson!




Modern View



If the protons and neutrons in this picture were 10 cm across, then the quarks and electrons would be less than 0.1 mm in size and the entire atom would be about 10 km across.

The Standard Model



$$\begin{aligned}\mathcal{L} = & -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} \\ & + i\bar{\psi} \not{D} \psi + \text{h.c.} \\ & + \chi_i^\dagger y_{ij} \chi_j \phi + \text{h.c.} \\ & + |D_\mu \phi|^2 - V(\phi)\end{aligned}$$

Modern View

PDF | Page 1 / 5

Details

Science
Volume 360, Issue 6385
Apr 2018

ARTICLE
Measurement of the fine-structure constant as a test of the Standard Model

View article page

Richard H. Parker, Chenghui Yu, Weicheng Zhong, Brian Estey and Holger Müller

Copyright © 2018 The Authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original U.S. Government Works
<http://www.sciencemag.org/about/science-licenses-journal-article-reuse>
This is an article distributed under the terms of the [Science Journals Default License](https://doi.org/10.1126/science.aap7706).
<https://doi.org/10.1126/science.aap7706>

Publisher American Association for the Advancement of Science
ISSN 0036-8075
eISSN 1095-9203
Print April 13, 2018

RESEARCH

METROLOGY

Measurement of the fine-structure constant as a test of the Standard Model

Richard H. Parker,^{1*} Chenghui Yu,^{1*} Weicheng Zhong,¹ Brian Estey,¹ Holger Müller^{1,2†}



Measurements of the fine-structure constant α require methods from across subfields and are thus powerful tests of the consistency of theory and experiment in physics. Using the recoil frequency of cesium-133 atoms in a matter-wave interferometer, we recorded the most accurate measurement of the fine-structure constant to date: $\alpha = 1/137.035999046(27)$ at 2.0×10^{-10} accuracy. Using multiphoton interactions (Bragg diffraction and Bloch oscillations), we demonstrate the largest phase effects (12 million radians) of any Ramsey-Bordé interferometer and control systematic of the electron gyromagnetic anomaly $g_e - 2$ via the Standard Model of particle physics is now limited by the uncertainty in $g_e - 2$; a 2.5σ tension rejects dark photons as the reason for the unexplained part of the muon's magnetic moment at a 99% confidence level. Implications for dark-sector candidates and electron substructure may be a sign of physics beyond the Standard Model that warrants further investigation.

The fine-structure constant α characterizes the strength of the electromagnetic interaction between elementary charged particles. It has been measured by various methods



ronic physics (1–3). This path leads to an accuracy of 0.24 part per billion (ppb) (4–6) and was until now the most accurate measurement of α .

d


The Language of Nature



Quantum Field Theory



A Feynman diagram





Gauge theory




Emmy Noether

Quantum Field Theory

Fields




A Feynman




Emmy Noether

Quantum Field Theory


Fields




Standard Model Interactions (Forces Mediated by Gauge Bosons)




X is any fermion in the Standard Model.




X is electrically charged.





X is any quark.




U is a up-type quark;
D is a down-type quark.




L is a lepton and ν is the corresponding neutrino.



X is a photon or Z-boson.





X and Y are any two electroweak bosons such that charge is conserved.




Emmy Noether



Quantum Field Theory



A Feynman diagram



Gauge theory



Emmy Noether



A little
problem with
infinities



The Standard Model

Standard Model of FUNDAMENTAL PARTICLES AND INTERACTIONS

The Standard Model summarizes the current knowledge in Particle Physics. It is the quantum theory that includes the theory of strong interactions (quantum chromodynamics or QCD) and the unified theory of weak and electromagnetic interactions (electroweak). Gravity is included on this chart because it is one of the fundamental interactions even though not part of the "Standard Model."

FERMIONS


matter constituents
spin = 1/2, 3/2, 5/2, ...

Leptons spin = 1/2			Quarks spin = 1/2		
Flavor	Mass GeV/c ²	Electric charge	Flavor	Approx. Mass GeV/c ²	Electric charge
ν_e electron neutrino	$<1 \times 10^{-8}$	0	u up	0.003	2/3
e electron	0.000511	-1	d down	0.006	-1/3
ν_μ muon neutrino	<0.0002	0	c charm	1.3	2/3
μ muon	0.106	-1	s strange	0.1	-1/3
ν_τ tau neutrino	<0.02	0	t top	175	2/3
τ tau	1.7771	-1	b bottom	4.3	-1/3

Spin is the intrinsic angular momentum of particles. Spin is given in units of $\hbar$, which is the quantum unit of angular momentum, where $\hbar = h/2\pi = 6.58 \times 10^{-25} \text{ GeV s} = 1.05 \times 10^{-34} \text{ J s}$.

Electric charges are given in units of the proton's charge. In SI units the electric charge of the proton is $1.60 \times 10^{-19} \text{ coulombs}$.

The **energy** unit of particle physics is the electronvolt (eV), the energy gained by one electron in crossing a potential difference of one volt. **Masses** are given in GeV/c^2 (remember $E = mc^2$), where $1 \text{ GeV} = 10^9 \text{ eV} = 1.60 \times 10^{-10} \text{ joule}$. The mass of the proton is $0.938 \text{ GeV}/c^2 = 1.67 \times 10^{-27} \text{ kg}$.



BOSONS

force carriers
spin = 0, 1, 2, ...

Unified Electroweak spin = 1			Strong (color) spin = 1		
Name	Mass GeV/c ²	Electric charge	Name	Mass GeV/c ²	Electric charge
γ photon	0	0	g gluon	0	0
W^-	80.4	-1	Color Charge Each quark carries one of three types of "strong charge," also called "color charge." These charges have nothing to do with the colors of visible light. There are eight possible types of color charge for gluons. Just as electrically-charged particles interact by exchanging photons, in strong interactions color-charged particles interact by exchanging gluons. Leptons, photons, and W and Z bosons have no strong interactions and hence no color charge.		
W^+	80.4	+1			
Z^0	91.187	0			

Quarks Confined in Mesons and Baryons

One cannot isolate quarks and gluons; they are confined in color-neutral particles called **hadrons**. This confinement (binding) results from multiple exchanges of gluons among the color-charged constituents. As color-charged particles (quarks and gluons) move apart, the energy in the color-force field between them increases. This energy eventually is converted into additional quark-antiquark pairs (see figure below). The quarks and antiquarks then combine into hadrons; these are the particles seen to emerge. Two types of hadrons have been observed in nature: **mesons** $q\bar{q}$ and **baryons** qqq .

Residual Strong Interaction

The strong binding of color-neutral protons and neutrons to form nuclei is due to residual strong interactions between their color-charged constituents. It is similar to the residual electrical interaction that binds electrically neutral atoms to form molecules. It can also be viewed as the exchange of mesons between the hadrons.

PROPERTIES OF THE INTERACTIONS

Baryons qqq and Antibaryons $\bar{q}\bar{q}\bar{q}$					
Baryons are fermionic hadrons. There are about 120 types of baryons.					
Symbol	Name	Quark content	Electric charge	Mass GeV/c ²	Spin
p	proton	uud	1	0.938	1/2
$\bar{p}$	anti-proton	$\bar{u}\bar{u}\bar{d}$	-1	0.938	1/2
n	neutron	udd	0	0.940	1/2
Λ	lambda	uds	0	1.116	1/2
Ω^-	omega	sss	-1	1.672	3/2


Property \ Interaction	Gravitational	Weak	Electromagnetic	Strong	
		(Electroweak)		Fundamental	Residual
Acts on:	Mass – Energy	Flavor		Color Charge	See Residual Strong Interaction Note
Particles experiencing:	All	Quarks, Leptons		Quarks, Gluons	Hadrons
Particles mediating:	Graviton (not yet observed)	W^+ W^- Z^0		Gluons	Mesons
Strength relative to electromag for two u quarks at:	10^{-41}	0.8	1	25	Not applicable to quarks
for two protons in nucleus	10^{-41}	10^{-4}	1	60	
	10^{-36}	10^{-7}	1	Not applicable to hadrons	20

Mesons $q\bar{q}$					
Mesons are bosonic hadrons. There are about 140 types of mesons.					
Symbol	Name	Quark content	Electric charge	Mass GeV/c ²	Spin
π^+	pion	$u\bar{d}$	+1	0.140	0
K^-	kaon	$s\bar{u}$	-1	0.494	0
ρ^+	rho	$u\bar{d}$	+1	0.770	1
B^0	B-zero	$d\bar{b}$	0	5.279	0
η_c	eta-c	$c\bar{c}$	0	2.980	0


The Missing Piece



The Standard Model



The Standard Model



FERMIONS

Leptons spin = 1/2		
Flavor	Mass GeV/c ²	Electric charge
ν_e electron neutrino	$<1 \times 10^{-8}$	0
e electron	0.000511	-1
ν_μ muon neutrino	<0.0002	0
μ muon	0.106	-1
ν_τ tau neutrino	<0.02	0
τ tau	1.7771	-1

Spin is the intrinsic angular momentum of a particle, measured in units of the quantum unit of angular momentum, $\hbar/2$.

Electric charges are given in units of the proton charge, which is 1.60×10^{-19} coulombs.

The **energy** unit of particle physics is the electron volt (eV). The energy of a proton in crossing a potential difference of 1 volt is $E = mc^2$, where $1 \text{ GeV} = 10^9 \text{ eV} = 1.6 \times 10^{-10} \text{ J}$.

Baryons qqq and Antibaryons $\bar{q}\bar{q}\bar{q}$

Baryons are fermionic hadrons. There are about 120 types of baryons.

Symbol	Name	Quark content	Electric charge
p	proton	uud	1
$\bar{p}$	anti-proton	$\bar{u}\bar{u}\bar{d}$	-1
n	neutron	udd	0
Λ	lambda	uds	0
Ω^-	omega	sss	-1

S

Force carriers spin = 0, 1, 2, ...

Strong (color) spin = 1		
Name	Mass GeV/c ²	Electric charge
g gluon	0	0

Color Charge

Each quark carries one of three types of "strong charge," also called "color charge." These charges have nothing to do with the colors of visible light. There are eight possible types of color charge for gluons. Just as electrons, in strong interactions color-charged particles, and W and Z bosons have no strong

are confined in color-neutral particles called hadrons. Multiple exchanges of gluons among the constituents (quarks and gluons) move apart, the energy eventually is converted into additional quarks and antiquarks then combine into hadrons. Two types of hadrons have been observed in




neutrons to form nuclei is due to residual strong interactions. It is similar to the residual electromagnetic interactions that hold atoms together to form molecules. It can also be observed in the decay of hadrons.

Mesons $q\bar{q}$

Mesons are bosonic hadrons. There are about 140 types of mesons.

Name	Quark content	Electric charge	Mass GeV/c ²	Spin
pion	$u\bar{d}$	+1	0.140	0
kaon	$s\bar{u}$	-1	0.494	0
rho	$u\bar{d}$	+1	0.770	1
B-zero	$d\bar{b}$	0	5.279	0
eta-c	$c\bar{c}$	0	2.980	0


Bound states



How much do we know?

Then

Every object of creation is made of atoms (paramāṇu) which in turn connect with each other to form molecules (aṇu). Atoms are eternal, and their combinations constitute the empirical material world.



Now



Particle Data Group <https://pdg.lbl.gov/>

A good set of units

SI units are NOT useful.
Too big!

Make life simple by setting

$$c = \hbar = 1$$

$$1 \text{ eV} = 1.6 \times 10^{-19} \text{ J}$$

Actually, particle theorists are lazy (or clever, depending on your point of view)—they seldom include the c 's and $\hbar$'s ($\hbar \equiv h/2\pi$) in their formulas. You're just supposed to fit them in for yourself at the end, to make the dimensions come out right. As they say in the business, “set $c = \hbar = 1$.” This amounts to working in units such that time is measured in centimeters and mass and energy in inverse centimeters; the unit of time is the time it takes light to travel 1 centimeter, and the unit of energy is the energy of a photon whose wavelength is 2π centimeters.

Conversion Factor	$\hbar = c = 1$ Units	Actual Dimension
$1 \text{ kg} = 5.61 \times 10^{26} \text{ GeV}$	GeV	$\frac{\text{GeV}}{c^2}$
$1 \text{ m} = 5.07 \times 10^{15} \text{ GeV}^{-1}$	GeV^{-1}	$\frac{\hbar c}{\text{GeV}}$
$1 \text{ sec} = 1.52 \times 10^{24} \text{ GeV}^{-1}$	GeV^{-1}	$\frac{\hbar}{\text{GeV}}$
$e = \sqrt{4\pi\alpha}$	—	$(\hbar c)^{1/2}$

By choosing units with $\hbar = c = 1$, it becomes unnecessary to write $\hbar$ and c explicitly in the formulas, thus saving a lot of time and trouble. We can always use dimensional analysis to work out unambiguously where the $\hbar$'s and c 's enter any formula. Hence, with a slight but permissible laziness, it is customary to speak of mass (m), momentum (mc), and energy (mc^2) all in terms of GeV, and to measure length ($\hbar/mc$) and time ($\hbar/mc^2$) in units of GeV^{-1} .

$$1 \text{ TeV} = 10^3 \text{ GeV} = 10^6 \text{ MeV} = 10^9 \text{ KeV} = 10^{12} \text{ eV}$$

$$1 \text{ fermi} \equiv 1 \text{ F} = 10^{-13} \text{ cm} = 5.07 \text{ GeV}^{-1}$$

$$(1 \text{ F})^2 = 10 \text{ mb} = 10^4 \mu\text{b} = 10^7 \text{ nb} = 10^{10} \text{ pb}$$

$$(1 \text{ GeV})^{-2} = 0.389 \text{ mb}$$



The charge of an electron

So far, we have not considered the elementary charge e , which measures how strongly electrons, say, interact electromagnetically with each other. To obtain a dimensionless measure of the strength of this interaction, we compare the electrostatic energy of repulsion between two electrons one natural unit of length apart with the rest mass energy of an electron:

$$\alpha = \frac{1}{4\pi} \frac{e^2}{(\hbar/mc)} / mc^2 = \frac{e^2}{4\pi\hbar c} \approx \frac{1}{137}.$$


we have adopted the rationalized Heaviside–Lorentz system of electromagnetic units. That is, the 4π factors appear in the force equations rather than in the Maxwell equations, and ϵ_0 is set equal to unity. This choice, which is conventional in particle physics, reduces Maxwell's equations to their simplest possible form. The value of α is, of course, the same in all systems of units, but the numerical value of e is different.

For historical reasons, α is known as the fine structure constant. Unfortunately, this name conveys a false impression. We have seen that the charge of an electron is not strictly constant but varies with distance because of quantum effects; hence α must be regarded as a variable, too. The value $\frac{1}{137}$ is the asymptotic value of α



Mysterious Neutrinos

β decay




Conservation of charge requires that B carry one more unit of positive charge than A . (We now realize that the underlying process here is the conversion of a neutron, in A , into a proton, in B ; but remember that in 1930 the neutron had not yet been discovered.) Thus the ‘daughter’ nucleus (B) lies one position farther along on the periodic table. There are many examples of beta decay: potassium goes to calcium (${}^{40}_{19}\text{K} \rightarrow {}^{40}_{20}\text{Ca}$), copper goes to zinc (${}^{64}_{29}\text{Cu} \rightarrow {}^{64}_{30}\text{Zn}$), tritium goes to helium (${}^3_1\text{H} \rightarrow {}^3_2\text{He}$), and so on.*

Now, it is a characteristic of two-body decays ($A \rightarrow B + C$) that the outgoing energies are kinematically determined, in the center-of-mass frame. Specifically, if the ‘parent’ nucleus (A) is at rest, so that B and e come out back-to-back with equal and opposite momenta, then conservation of energy dictates that the electron energy is

$$E = \left(\frac{m_A^2 - m_B^2 + m_e^2}{2m_A} \right) c^2$$

In modern terminology, the fundamental beta decay process is



(neutron goes to proton plus electron plus antineutrino).




Table 1.1 The lepton family, 1962–1976

	Lepton number	Electron number	Muon number
Leptons			
e^-	1	1	0
ν_e	1	1	0
μ^-	1	0	1
ν_μ	1	0	1
Antileptons			
e^+	-1	-1	0
$\bar{\nu}_e$	-1	-1	0
μ^+	-1	0	-1
$\bar{\nu}_\mu$	-1	0	-1

Strange Particles

For a brief period in 1947, it was possible to believe that the major problems of elementary particle physics were solved.

In December 1947, Rochester and Butler discovered a neutral particle decaying to two pions

$$K^0 \rightarrow \pi^+ + \pi^-$$

In 1949, another Kaon was discovered


$$K^+ \rightarrow \pi^+ + \pi^+ + \pi^-$$

In due course, many more mesons were discovered – the η , the ϕ , the ω , the ρ 's, and so on.

In 1950, another Baryon was discovered

$$\Lambda \rightarrow p^+ + \pi^-$$

Not only were the new particles unexpected; there is a more technical sense in which they seemed 'strange': they are *produced* copiously (on a time scale of about 10^{-23} seconds), but they *decay* relatively slowly (typically about 10^{-10} seconds). This suggested that the mechanism involved in their production is entirely different from that which governs their disintegration. In modern language, the strange particles are *produced* by the *strong* force (the same one that holds the nucleus together), but they *decay* by the *weak* force (the one that accounts for beta decay and all other neutrino processes).




'Why is the proton stable?' Why, for example, doesn't it decay into a positron and a photon:



$$p^+ \rightarrow e^+ + \gamma$$

Baryon Number

There are patterns



Baryons




Mesons


The Quark Model

$$\begin{aligned}p &= uud \\n &= udd \\ \Delta^{++} &= uuu.\end{aligned}$$

1. Every baryon is composed of three quarks (and every *antibaryon* is composed of three *antiquarks*).
2. Every meson is composed of a quark and an antiquark.





The quarks



The antiquarks

The Quark Model

The quarks have colour
but the "world" is colourless



1. Every baryon is composed of three quarks (and every *antibaryon* is composed of three *antiquarks*).
2. Every meson is composed of a quark and an antiquark.

So, what do we have?

Charge: All three interactions, of course, conserve electric charge. In the case of the weak interactions, the lepton (or quark) that comes out may not have the same charge as the one that went in, but if so, the difference is carried away by the W .

Color: The electromagnetic and weak interactions do not affect color. At a strong vertex the quark color does change, but the difference is carried off by the gluon. (The direct gluon–gluon couplings also conserve color.) However, since naturally occurring particles are always colorless, the observable manifestation of color conservation is pretty trivial: zero in, zero out.

Baryon number: In all the primitive vertices, if a quark goes *in*, a quark comes *out*, so the total number of quarks present is a constant. In this arithmetic we count *antiquarks* as *negative*, so that, for example, at the vertex $q + \bar{q} \rightarrow g$ the quark number is zero before and zero after. Of course, we never see individual quarks, only baryons (with quark number 3), antibaryons (quark number -3), and mesons (quark number zero). So, in practice, it is more convenient to speak

of the conservation of baryon number (1 for baryons, -1 for antibaryons, and 0 for everything else). The baryon number is just $\frac{1}{3}$ the quark number. Notice that there is no analogous conservation of *meson number*; since mesons carry zero quark number, a given collision or decay can produce as many mesons as it likes, consistent with conservation of energy.

Lepton number: The strong forces do not touch leptons at all; in an electromagnetic interaction the same particle comes out (accompanied by a photon) as went in; and in the weak interactions if a lepton goes in, a lepton comes out (not necessarily the *same* one, this time). So, lepton number is absolutely conserved. Until recently there appeared to be no cross-generation mixing among the leptons, so electron number, muon number, and tau number were all separately conserved. This remains true in most cases, but neutrino oscillations indicate that it is not absolute.*

So, what do we have?

Flavor: What about quark *flavor*? Flavor is conserved at a strong or electromagnetic vertex, but *not* at a weak vertex, where an up-quark may turn into a down quark or a strange quark, with nothing at all picking up the lost upness or supplying the ‘gained’ downness or strangeness. Because the weak forces *are* so weak, we say that the various flavors are *approximately* conserved. In fact, as you may remember, it was precisely this approximate conservation that led Gell-Mann to introduce the notion of strangeness in the first place. He ‘explained’ the fact that strange particles are always produced in pairs:

$$\pi^-(d\bar{u}) + p^+(uud) \rightarrow K^+(u\bar{s}) + \Sigma^-(dds)$$

for instance, but

$$\pi^-(d\bar{u}) + p^+(uud) \not\rightarrow \pi^+(u\bar{d}) + \Sigma^-(dds)$$

by arguing that the latter violates conservation of strangeness. (Actually, this *is* a possible *weak* interaction, but it will never be seen in the laboratory, because it must compete

against enormously more probable strong processes that do conserve strangeness.) In *decays*, however, the nonconservation of strangeness is very conspicuous, because for many particles this is the only way they *can* decay; there *is* no competition from strong or electromagnetic processes. The Λ , for instance, is the lightest strange baryon; if it is to decay, it must go to n (or p) plus something. But the lightest strange meson is the K , and n (or p) plus K weighs substantially more than the Λ . If the Λ decays at all (and it *does* as we know: $\Lambda \rightarrow p^+ + \pi^-$ 64% of the time, and $\Lambda \rightarrow n + \pi^0$ 36% of the time), then strangeness cannot be conserved, and the reaction must proceed by the weak interaction. By contrast, the Δ^0 (with a strangeness of zero) can go to $p^+ + \pi^-$ or $n + \pi^0$ by the *strong* interaction, and its lifetime is accordingly much shorter.

So, what do we have?

Flavor: What about quark *flavor*? Flavor is conserved at a strong or electromagnetic vertex, but *not* at a weak vertex, where an up-quark may turn into a down quark or a strange quark, with nothing at all picking up the lost upness or supplying the ‘gained’ downness or strangeness. Because the weak forces *are* so weak, we say that the various flavors are *approximately* conserved. In fact, as you may remember, it was precisely this approximate conservation that led Gell-Mann to introduce the notion of strangeness in the first place. He ‘explained’ the fact that strange particles are always produced in pairs:

$$\pi^{-}(d\bar{u}) + p^{+}(uud) \rightarrow K^{+}(u\bar{s}) + \Sigma^{-}(dds)$$

for instance, but

$$\pi^{-}(d\bar{u}) + p^{+}(uud) \not\rightarrow \pi^{+}(u\bar{d}) + \Sigma^{-}(dds)$$

by arguing that the latter violates conservation of strangeness. (Actually, this *is* a possible *weak* interaction, but it will never be seen in the laboratory, because it must compete

against enormously more probable strong processes that do conserve strangeness.) In *decays*, however, the nonconservation of strangeness is very conspicuous, because for many particles this is the only way they *can* decay; there *is* no competition from strong or electromagnetic processes. The Λ , for instance, is the lightest strange baryon; if it is to decay, it must go to n (or p) plus something. But the lightest strange meson is the K , and n (or p) plus K weighs substantially more than the Λ . If the Λ decays at all (and it *does* as we know: $\Lambda \rightarrow p^{+} + \pi^{-}$ 64% of the time, and $\Lambda \rightarrow n + \pi^{0}$ 36% of the time), then strangeness cannot be conserved, and the reaction must proceed by the weak interaction. By contrast, the Δ^{0} (with a strangeness of zero) can go to $p^{+} + \pi^{-}$ or $n + \pi^{0}$ by the *strong* interaction, and its lifetime is accordingly much shorter.

Which are allowed?

Home work

(a) $p + \bar{p} \rightarrow \pi^+ + \pi^0$

(c) $\Sigma^0 \rightarrow \Lambda + \pi^0$

(e) $e^+ + e^- \rightarrow \mu^+ + \mu^-$

(g) $\Delta^+ \rightarrow p + \pi^0$

(i) $e + p \rightarrow \nu_e + \pi^0$

(k) $p \rightarrow e^+ + \gamma$

(m) $n + \bar{n} \rightarrow \pi^+ + \pi^- + \pi^0$

(o) $K^- \rightarrow \pi^- + \pi^0$

(q) $\Sigma^0 \rightarrow \Lambda + \gamma$

(s) $\Xi^0 \rightarrow p + \pi^-$

(u) $\pi^0 \rightarrow \gamma + \gamma$

(b) $\eta \rightarrow \gamma + \gamma$

(d) $\Sigma^- \rightarrow n + \pi^-$

(f) $\mu^- \rightarrow e^- + \bar{\nu}_e$

(h) $\bar{\nu}_e + p \rightarrow n + e^+$

(j) $p + p \rightarrow \Sigma^+ + n + K^0 + \pi^+ + \pi^0$

(l) $p + p \rightarrow p + p + p + \bar{p}$

(n) $\pi^+ + n \rightarrow \pi^- + p$


(p) $\Sigma^+ + n \rightarrow \Sigma^- + p$

(r) $\Xi^- \rightarrow \Lambda + \pi^-$

(t) $\pi^- + p \rightarrow \Lambda + K^0$

(v) $\Sigma^- \rightarrow n + e + \bar{\nu}_e$


Is that all?



Is that all?


Fifth dimension

Space is warped by energy throughout five-dimensional space-time. As a result, gravity is much weaker on our brane.




Is that all?


SUPERSYMMETRY AND PARTICLE PAIRS



Standard Model Particles



Hypothetical SUSY Particles



Supersymmetry remains just an interesting, if mathematically elegant, hypothesis right now. It would explain why particles have the masses they do and shed light on dark matter. Many versions of the theory suggest the LHC's current experiments will be energetic enough to produce the heavier supersymmetric particles — if they exist.

Where to test?




- ▶ Approximately 27km in circumference. Proton beams travel at close to the speed of light to collide.
- ▶ The beams travel in opposite directions in separate beam pipes kept at ultrahigh vacuum. They are guided around the accelerator ring by a strong magnetic field maintained by superconducting electromagnets chilled to -271.3°C (1.9K) – a temperature colder than outer space.

Proton + proton collider



The Large Hadron Collider at CERN

Where to test?




guided around the curve of light to collide.
in vacuum. They are
superconducting
face.

Proton + proton collider



The Large Hadron Collider at CERN

Where to test?



ABOUT NEWS SCIENCE RESOURCES Q SEARCH | EN -

A three-in-one vacuum system

The LHC is unusual in that it has three separate vacuum systems: one for the beam pipes, one for insulating the cryogenically cooled magnets and one for insulating the helium distribution line.

To avoid colliding with gas molecules inside the accelerator, the beams of particles in the LHC must travel in a vacuum as empty as interstellar space. In the cryomagnets and the helium distribution line, the vacuum serves a different purpose. Here, it acts as a thermal insulator, to reduce the amount of heat that seeps from the surrounding room-temperature environment into the [cryogenic parts](#) which are kept at at 1.9 K (-271.3°C).

The largest vacuum system in the world

With a total of 104 kilometres of piping under vacuum, the vacuum system of the LHC is among the largest in the world. The insulating vacuum, equivalent to some 10^{-6} mbar, is made up of an impressive 50 km of piping, with a combined volume of 15,000 cubic metres, more than enough to fill the nave of a cathedral. Building this vacuum system required more than 250,000 welded joints and 18,000 vacuum seals. The remaining 54 km of pipes under vacuum are the beam pipes, through which the LHC's two beams travel. The pressure in these pipes is in the order of 10^{-10} to 10^{-11} mbar, a vacuum almost as rarefied as that found on the surface of the Moon. The LHC's vacuum systems are fitted with 170 Bayard-Alpert ionisation gauges and 1084 Pirani and Penning gauges to monitor the vacuum pressure.

A vacuum thinner than the interstellar void

Ultra-high vacuum is needed for the pipes in which particle beams travel. This includes 48 km of arc sections, kept at 1.9 K, and 6 km of straight sections, kept at room temperature, where beam-control systems and the insertion regions for the experiments are located.


In the arcs, the ultra-high vacuum is maintained by cryogenic pumping of 9000 cubic metres of gas. As the beam pipes are cooled to extremely low temperatures, the gases condense and adhere to the walls of the beam pipe by adsorption. Just under two weeks of pumping are required to bring the pressures down below 1.013×10^{-10} mbar (or 10^{-13} atmospheres).

Two important design features maintain the ultra-high vacuum in the room-temperature sections. Firstly, these sections make widespread use of a non-evaporable "getter coating" – developed and industrialized at CERN – that absorbs residual molecules when heated. The coating consists of a thin liner of titanium-zirconium-vanadium alloy deposited inside the beam pipes. It acts as a distributed pumping system, effective for removing all gases except methane and the noble gases.



The Large Hadron Collider at CERN

Where to test?



CMS

SEARCH | EN

ne for

In the

t that seeps

nt to some

Building

through

of the

at

ures,

the pressures down below

are sections. Firstly, these sections make widespread use of a non-evaporable


absorbs residual molecules when heated. The coating consists of a thin liner of titanium-zirconium-

pipes. It acts as a distributed pumping system, effective for removing all gases except methane and the noble gases.



The Large Hadron Collider at CERN

Where to test?



Gold + polarised proton collider

Where to test?




Proton + anti-proton collider




Tevatron

Where to test?



Neutrino Experiments


Inside a collider



Inside a collider




CMS Experiment at LHC, CERN
Data recorded: Wed Nov 25 12:21:51 2015 CET
Run/Event: 262548 / 14582169
Lumi section: 309




Inside a collider



Run: 313100
Event: 196478531
2016-11-18 23:23:28 CEST




How to see?




How to see?

proton



proton



How to see?

